Bmp8a Is a Novel Player in Regulation of Antiviral Immunity

Shenjie Zhong, Haoyi Li, Yun-Sheng Wang, Ying Wang, Guangdong Ji, Hong-Yan Li, Shicui Zhang*, Zhenhui Liu**

1 College of Marine Life Science and Institute of Evolution & Marine Biodiversity, Ocean University of China, Qingdao

266003, China

2 Laboratory for Marine Biology and Biotechnology, Pilot National Laboratory for Marine Science and Technology

(Qingdao), Qingdao 266003, China

26   *Correspondence author: e-mail: sczhang@ouc.edu.cn

27   **Correspondence author: e-mail: zhenhuiliu@ouc.edu.cn

Abstract

Bone morphogenetic protein (BMP) is a kind of classical multi-functional growth factor that plays a vital role in the formation and maintenance of bone, cartilage, muscle, blood vessels, and the regulation of energy balance. Whether BMP plays a role in antiviral immunity is unknown. Here we demonstrate that Bmp8a is a newly-identified positive regulator for antiviral immune responses. The *bmp8a*−/− zebrafish, when infected with the viruses of GCRV, SVCV or TSVDV, show significantly reduced antiviral immunity, increased viral load and morbidity. We also show for the first time that Bmp8a interacts with Alk6a, which promotes the phosphorylation of Tbk1 and Irf3 through p38 MAPK pathway, and induces the production of type I IFNs in response to virus infection. Upon virus infection, *bmp8a* expression is activated through the binding of Stat1a/Stat1b to the GAS motifs in *bmp8a* promoter region, enlarging the antiviral innate immune signal. Our study uncovers a previously unrecognized role of Bmp8a in regulation of antiviral immune responses and provides a new target for controlling viral infection.

Introduction

Viruses infect all groups of living things and produce a variety of diseases. Host cell detection of viral infection depends on the recognition of pathogen-associated molecular patterns (PAMPs) from viral proteins or nucleic acids by the pattern recognition receptors (PRRs) such as Toll-like receptors (TLRs), RIG-I-like receptors (RLRs), NOD-like receptors (NLRs) and cytoplasmic DNA or RNA sensors[1-6]. The PRRs, once stimulated by their appropriate ligands, trigger distinct intracellular signaling pathways that converge on the activation of IFN regulatory factor 3 (IRF3) and/or IFN regulatory factor 7 (IRF7), which then translocate into the nucleus and activate the production of interferons (IFNs) that play important roles in inhibiting virus replication[7,8].

IFNs are classified into type I IFNs that include IFNα and IFNβ; type II IFN which has only a single member IFN-γ; and type III IFNs that consist of IFN-λs[9]. All these three types of IFNs trigger intracellular signaling cascades generally via the Janus kinase signal transducer and activator of transcription (JAK-STAT) pathway[10]. Canonically, type I IFNs initiate the signaling via binding to a heterodimeric receptor complex IFN-α/β receptor (IFNAR), inducing phosphorylation of transcription factors STAT1 and STAT2. The phosphorylated STATs form a trimolecular complex with IRF9, which translocates to the nucleus and binds to IFN-stimulated response elements (ISREs) of IFN-stimulated genes (ISGs), resulting in transcriptional activation of the target ISGs, including those encoding numerous cytokines and antiviral proteins[11-16]. In contrast, type II IFN binds to its unique receptor, IFN-γ receptor (IFNGR), resulting in the phosphorylation of STAT1 and the formation of STAT1 homodimers that recognize gamma-activated sequences (GASs) present in the promoter regions of IFN-γ-regulated genes[9].

Bone morphogenetic proteins (BMPs) are potent growth factors belonging to the transforming growth factor beta (TGFβ) superfamily, which are intercellular signaling molecules with multiple functions in development and differentiation, such as skeletal formation and hematopoiesis, as well as lipid oxidation[17-21]. However, accumulating data suggests that BMPs also play important roles in the regulation of immune responses[22-27]. For example, BMP7 has been shown to drive Langerhans cell differentiation[28]. Interestingly, BMP signaling appears to have different responses in T cells: BMP4 and BMP6 were shown to promote T cell proliferation, whereas BMP2 was seen to inhibit T cell proliferation[29,30]. In addition, although BMPs were found to promote tumor growth by inhibiting the functions of cytotoxic T lymphocytes (CTLs) and dendritic cells (DCs), they were also shown to inhibit cancer cell proliferation and promote the activity of NK cells that typically display antitumour activity[31]. Therefore, it remains controversial regarding the functions of BMPs in immune responses.

In humans and mice, two closely related BMP8 genes, *BMP8A* and *BMP8B*, have been identified[32,33]. BMP8B has been found to play roles in the regulation of thermogenesis in mature brown adipose tissue and spermatogenesis in male germ cells, while BMP8A shown to closely correlate with the progression of spermatogenesis[34,35]. In contrast, only one

bmp8 gene, *bmp8a*, has been isolated in zebrafish, and shown to be involved in the regulation of lipid metabolism[36]. To

date, whether Bmp8a plays a role in immune responses is completely unknown. The present study was thus performed

to answer this question. We demonstrated that Bmp8a was a previously unrecognized regulator functioning in antiviral

immune responses of zebrafish *Danio rerio*, providing a new target for control of viral infection.

Results

Bmp8a inhibited RNA virus replication *in vitro*. To explore the function of zebrafish Bmp8a in the antiviral immune

response, we infected both the wild-type zebrafish liver (ZFL) cells and ZFL cells over-expressing bmp8a as well as the

epithelioma papulosum cyprini (EPC) cells and EPC cells over-expressing bmp8a with Grass carp reovirus (GCRV), a

dsRNA virus, and then monitored the cytopathic effect（CPE）and the viral titers of the supernatants. In response to GCRV

challenge, an apparent CPE was observed in the control cells, while the bmp8a-overexpressing ZFL cells had much less

CPE (Fig. 1a). In accordance with the CPE results, a significant decrease of the viral titers was observed in the

supernatants of bmp8a-overexpressing ZFL cells (Fig. 1b). Similarly, the GCRV-induced CPE and GCRV yields of the

supernatants were also markedly reduced in the bmp8a-overexpressing EPC cells (Fig. 1c, d). These suggested that

Bmp8a suppressed the replication of the RNA virus in the cells.

Bmp8a was involved in antiviral responses *in vivo*. To test the function of Bmp8a in host defense against virus infection

in vivo, we generated *bmp8a* deficient (*bmp8a$^{-/-}$*) zebrafish using the TALEN approach (Supplementary Fig. 1a), which

caused 7 nucleotides deletion in the exon 4 (Supplementary Fig. 1b, c). We challenged both the wild-type and *bmp8a$^{-/-}$*

mutant zebrafish by intraperitoneal injection with GCRV. Compared with wild-type zebrafish, the levels of GCRV RNA in

the liver, kidney, intestine, and spleen of *bmp8a$^{-/-}$* zebrafish were considerably increased (Fig. 1e). Accordingly, *bmp8a$^{-/-}$*

zebrafish exhibited significantly reduced survival rate than wild-type zebrafish upon GCRV infection (Fig. 1f). When the

wild-type and *bmp8a$^{-/-}$* zebrafish were similarly challenged with spring viremia of carp virus (SVCV), a negative ssRNA

virus, the survival rate of *bmp8a$^{-/-}$* zebrafish was also remarkably lower than that of wild-type zebrafish (Fig. 1g). Moreover,

when the wild-type and *bmp8a$^{-/-}$* zebrafish were infected with turbot skin verruca disease virus (TSVDV) isolated from

skin (Supplementary Fig. 2), mortality occurred in *bmp8a$^{-/-}$* zebrafish at 108 h postinfection, and all the fish died at 156 h

post infection, while none of the wild type zebrafish were dead during the period (Fig. 1h). These data indicated that

bmp8a$^{-/-}$ zebrafish were more susceptible to virus infection than wild-type zebrafish, suggesting involvement of Bmp8a in

antiviral immune responses *in vivo*.

Fig.1 Bmp8a inhibits RNA virus replication both *in vitro* and *in vivo*. a, b ZFL cells were transfected with bmp8a (1μg) or empty vector (1μg), respectively. The cells were infected with GCRV (5×10^4TCID$_{50}$ per ml) at 24 h post-transfection, and the culture supernatants were collected at 72 h post-infection. The cell monolayers were fixed with 4% paraformaldehyde for 1 h and stained with 0.5% crystal violet for 2 h (**a**), and the viral titers of the supernatants were determined by TCID$_{50}$ assays (**b**). **c, d** Similar as (**a**, **b**) but in EPC cells. **e** The expression of GCRV RNA in the liver, kidney, intestine, and spleen from wild-type (WT) or *bmp8a*$^{-/-}$ zebrafish injected i.p. with 50 μl of GCRV (10^8TCID$_{50}$ per ml) for 72 h. **f, g** Kaplan–Meier analysis of the overall survival of WT (*n* = 20) or *bmp8a*$^{-/-}$ zebrafish (*n* = 20) which were injected i.p. with 50 μl of GCRV (10^8TCID$_{50}$ per ml) or SVCV (10^8TCID$_{50}$ per ml) and monitored every 12 h after infection. **h** Kaplan–Meier analysis of the overall survival of WT (*n* = 8) or *bmp8a*$^{-/-}$ zebrafish (*n* = 8) which were injected i.p. with 50 μl of TSVDV (crude virus extracts filtered by a 0.45 μm microporous membrane) and monitored every 12 h after infection. The expression of zebrafish *actb1* was used as an internal control for the qRT-PCR. Data were from three independent experiments (**a–e**) or two independent experiments (**f-h**). Data were analyzed by Student's *t*-test (two-tailed) or log-rank

(Mantel–Cox) test and were presented as mean ± SD (*$p < 0.05$, ***$p < 0.001$). Source data are provided as a Source

Data file.

Bmp8a promoted expression of antiviral genes. In zebrafish, type I IFNs contained 4 members *ifnφ1*, *ifnφ2*, *ifnφ3*, and

ifnφ4, among which *ifnφ1* and *ifnφ3* were shown to be involved in the antiviral response. In contrast, in carp, only one type

I IFN was identified to fulfill antiviral role in EPC cell[37,38]. To pinpoint the role of Bmp8a in the antiviral immune response,

both the wild-type and *bmp8a*$^{-/-}$ zebrafish were injected intraperitoneally with GCRV, and the expression of type I IFN

genes and the antiviral protein gene *mx* were detected using quantitative reverse transcription PCR (qRT-PCR). We found

that the expression of both *ifnφ1* and *ifnφ3* as well as *mx* was all significantly down-regulated in the liver, kidney, intestine,

and spleen of *bmp8a*$^{-/-}$ zebrafish, compared with those of wild-type fish (Fig. 2a-d). We then over-expressed bmp8a in

ZFL cells and found that ZFL cells with over-expressed Bmp8a showed markedly higher expression of *ifnφ1*, *ifnφ3*, and

mx than that of control cells, infected with or without GCRV (Fig. 2e-h). In addition, *bmp8a* knockdown caused remarkably

decreased expression of *ifnφ1*, *ifnφ3*, and *mx* in ZFL cells (Fig. 2i-l). Similarly, the expression of *ifn* and *mx* in EPC cells

with over-expressed Bmp8a was also remarkably up-regulated than that in control cells (Fig. 2m-p). Moreover, EPC cells

cotransfected with bmp8a expressing plasmid and IFNφ1, IFNφ3, or EPC IFN promoter-driven luciferase plasmid, followed

by infection with or without GCRV, had markedly increased intracellular IFNφ1, IFNφ3, or EPC IFN promoter-driven

luciferase activities (Fig. 2q-s). All these data denoted that Bmp8a promoted the expression of both type I IFN and *mx*

genes.

Fig.2 Bmp8a promotes antiviral innate immune responses. a-d Expression of *ifnφ1*, *ifnφ3*, and *mx* mRNA in the liver,

kidney, intestine, and spleen from WT or *bmp8a^-/-* zebrafish injected i.p. with 50 μl of GCRV (10^8TCID$_{50}$ per ml) for 72 h.

e, **f** Expression of *ifnφ1*, *ifnφ3*, and *mx* mRNA after transfected with bmp8a (2 μg) or empty vector (2 μg) in ZFL cells. The

cells were collected at 36 h (**e**) or 48 h (**f**) post transfection. **g, h** Expression of *ifnφ1*, *ifnφ3*, and *mx* mRNA after transfected

with bmp8a (2 μg) or empty vector (2 μg) in ZFL cells for 24 h, followed by infection with GCRV for another 24 h (**g**) or 36

h (**h**). **i**, **l** Expression of *ifnφ1*, *ifnφ3*, and *mx* mRNA after *bmp8a* knockdown in ZFL cells. The cells were collected at 36 h

(**i**) or 48 h (**l**) post knockdown. **k**, **l** Expression of *ifnφ1*, *ifnφ3*, and *mx* mRNA after knockdown *bmp8a* in ZFL cells for 24

h, followed by infection with GCRV for another 24h (**k**) or 36 h (**l**). **m**, **n** Expression of EPC *ifn* and EPC *mx* mRNA after

transfected with bmp8a (2 μg) or empty vector (2 μg) in EPC cells. The cells were collected at 36 h (**m**) or 48 h (**n**) post

transfection. **o**, **p** Expression of EPC *ifn* and EPC *mx* mRNA after transfected with bmp8a (2 μg) or empty vector (2 μg) in

EPC cells for 24 h, followed by infection with GCRV for another 24 h (**o**) or 36 h (**p**). The expression of zebrafish *actb1* or

EPC *actin* was used as an internal control for the qRT-PCR. **q-s** EPC cells were transfected with IFNφ1pro-luc (200 ng,

q), IFNφ3pro-luc (200 ng, **r**) or EPC IFN pro-luc (200 ng, **s**) respectively, with or without bmp8a (200 ng), followed by

infection with GCRV. At the indicated time points, cells were collected for luciferase assays. *Renilla* luciferase was used

as the internal control. Data were from three independent experiments and were analyzed by Student's *t*-test (two-tailed)

and were presented as mean ± SD (**$p < 0.01$, ***$p < 0.001$). Source data are provided as a Source Data file.

Bmp8a activated Tbk1-Irf3-Ifn antiviral signaling via p38 MAPK pathway.   To putatively examine the molecular mechanism

by which Bmp8a stimulates *ifn* and *mx* expression, we carried out transcriptome analysis of the livers of *bmp8a$^{-/-}$* mutant and

wild-type zebrafish, that both had been intraperitoneally injected with GCRV. Kyoto Encyclopedia of Genes and Genomes

(KEGG) analyses revealed that, compared with wild-type zebrafish, the down-regulated genes in the mutant were remarkably

enriched in the RIG-I-like receptor signaling pathway, NF-kappa B signaling pathway, TNF signaling pathway, and protein

digestion and absorption that function in the antiviral immune process (Supplementary Fig. 3). In agreement with transcriptome

analysis, we found that the expressions of *tbk1*, *irf3*, and *irf7* were significantly upregulated in bmp8a-overexpressing ZFL cells

than control cells, infected with or without GCRV (Fig. 3a-d). Similar expression patterns of *tbk1*, *irf3*, and *irf7* were also observed

in EPC cells, infected with or without GCRV (Fig. 3e-h). In addition, *bmp8a* knockdown resulted in considerably reduced

expression of *tbk1*, *irf3*, and *irf7* in ZFL cells than control cells, infected with or without GCRV (Fig. 3i-l). We then evaluated the

activation of Tbk1 and Irf3 by immunoblot assays in both bmp8a-overexpressing and wild-type ZFL cells as well as EPC cells,

infected with or without GCRV. It was found that phosphorylation levels of Tbk1 and Irf3 were significantly increased in both

types of the cells with overexpressed Bmp8a (Fig. 3m-p). Moreover, *bmp8a* knockdown markedly reduced the phosphorylation

levels of Tbk1 and Irf3 in ZFL cells, infected with or without GCRV (Fig. 3q, r). Furthermore, in luciferase activity assays,

transfection of dominant negative mutations of *tbk1* (tbk1-K38M), *irf3* (irf3DN), or *irf7* (irf7DN) in EPC cells induced a significant

loss in the ability of Bmp8a to activate the IFN promoter (Fig. 3s-u). This was further supported by the observations *in vivo* that

the expression of *tbk1*, *irf3*, and *irf7* in the liver, kidney, intestine, and spleen was significantly downregulated in *bmp8a$^{-/-}$* mutant

than that in wild-type zebrafish (Fig. 3v-y). These data together revealed that Bmp8a induced *ifn* and *mx* expression via

promoting Tbk1-Irf3-Ifn antiviral signaling.

BMPs transmit signals through smad1/5/8, smad2/3, ERK, JNK, or p38 MAPK pathways[39-41]. Thus, the effects of p38

MAPK inhibitor SB203580, JNK inhibitor SP600125, MEK1/2 inhibitor U0126, SMAD1/5/8 inhibitor DMH1, and smad2/3

inhibitor TP0427736 HCl on the expression of *ifn* was tested in bmp8a-overexpressing ZFL cells and EPC cells. We have

shown above that Bmp8a overexpression increased the expression of *ifn* (Fig. 2). Here we found that only p38 MAPK

inhibitor SB203580 significantly reduced the expression of *ifnφ1* and *ifnφ3* in ZFL cells and the expression of *ifn* in EPC

cells (Fig. 4a-c). Furthermore, the IFN promoter-driven luciferase assays revealed that the luciferase activities were

markedly reduced in EPC cells upon treatment with p38 MAPK inhibitor SB203580 (Fig. 4d-f), consistent with the

observation that Bmp8a induced the expression of *ifn* through p38 MAPK pathway. Taken the previous report that the

MAPK signaling increased the phosphorylation of TBK1 and IRF3 upon viral infection into consideration[42], we suggested

that Bmp8a activated Tbk1-Irf3-Ifn antiviral signaling via p38 MAPK pathway.

Fig.3 Bmp8a increases Tbk1-Irf3-Ifn antiviral signaling. a, **b**, **e**, **f** Expression of *irf3*, *irf7*, and *tbk1* mRNA after

transfected with 2 μg bmp8a or empty vector in ZFL (**a**, **b**) or EPC (**e**, **f**) cells. The cells were collected at 36 h (**a**, **e**) or 48

h (**b**, **f**) post transfection. **c**, **d**, **g**, **h** Expression of *irf3*, *irf7*, and *tbk1* mRNA after transfected with 2 μg bmp8a or empty

vector in ZFL (**c**, **d**) or EPC (**g**, **h**) cells for 24 h, followed by infection with GCRV for another 24 h (**c**, **g**) or 36 h (**d**, **h**). **i, j,**

k, l Expression of *irf3*, *irf7*, and *tbk1* mRNA after *bmp8a* knockdown in ZFL cells. The cells were collected at 36 h (**i**) and

48 h (**j**) post knockdown or at 24 h (**k**) and 36 h (**l**) post infected with GCRV. **m**, **o** Immunoblot analysis of phosphorylated

(p-) Tbk1 and Irf3 after transfected with 2 μg bmp8a or empty vector in ZFL (**m**) or EPC (**o**) cells. The cells were collected

at 36 h or 48 h post transfection for Immunoblot analysis. **n**, **p** Immunoblot analysis of phosphorylated (p-) Tbk1 and Irf3

after transfected with 2 μg bmp8a or empty vector in ZFL (**n**) or EPC (**p**) cells for 24 h, followed by infection with GCRV

for another 24 h or 36 h. **q**, **r** Immunoblot analysis of phosphorylated (p-) TBK1 and IRF3 after *bmp8a* knockdown in ZFL

cells. The cells were collected at 36 h and 48 h post knockdown or at 24 h and 36 h post infected with GCRV. **s-u** EPC

cells were cotransfected with IFN-φ1pro-luc (200 ng, **s**), IFN-φ3pro-luc (200 ng, **t**) or EPC IFNpro-luc (200ng, **u**), and

bmp8a (100 ng) together with each of the dominant negative plasmids including tbk1-K38M (100 ng), irf3DN (100 ng) and

irf7DN (100 ng). At 48 h post transfection, the cells were collected for luciferase assays. *Renilla* luciferase was used as

the internal control. **v-y** Expression of *irf3*, *irf7*, and *tbk1* mRNA in the liver, kidney, intestine, and spleen from WT or *bmp8a*

$^{-/-}$ zebrafish injected i.p. with 50 μl of GCRV (10^8TCID$_{50}$ per ml). The expression of zebrafish *actb1* or EPC *actin* was used

as an internal control for the qRT-PCR. Data were from three independent experiments and were analyzed by Student's

t-test (two-tailed) and were presented as mean ± SD (**$p < 0.01$, ***$p < 0.001$). Source data are provided as a Source

Data file.

Fig.4 Bmp8a promotes the IFN expression via p38 MAPK pathway. a, **b** Expression of *ifnφ1* (**a**) and *ifnφ3* (**b**) mRNA

after transfected with bmp8a (2 µg) in ZFL cells for 24 h, followed by treatment with SB203580, SP600125, U0126, DMH1

and TP0427736 HCl for another 24 h. **c** similar as (**a**, **b**) but in EPC cells. The expression of zebrafish *actb1* or EPC *actin*

was used as an internal control for the qRT-PCR. **d-f** EPC cells were cotransfected with IFNφ1pro-luc (200 ng, **d**),

IFNφ3pro-luc (200 ng, **e**) or EPC IFNpro-luc (200 ng, **f**), pRL-TK (20 ng) together with bmp8a (200 ng) or empty vector

(200 ng), respectively. At 24 h post transfection, cells were treated with or without SB203580 for another 24 h and then

harvested for detection of luciferase activity. *Renilla* luciferase was used as the internal control. Data were from three

independent experiments and were analyzed by Student's *t*-test (two-tailed) and were presented as mean ± SD (**$p <$

0.01, ***$p < 0.001$). Source data are provided as a Source Data file.

BMP type I receptor Alk6a interacted with Bmp8a and participated in antiviral immunity. BMPs exert their biological

effects through the sequential activation of two types of transmembrane receptors, namely BMP receptor type I and BMP

receptor type II, that both possess intrinsic serine/threonine kinase activity[43,44]. To detect which receptor is involved in the

regulation of the antiviral immune responses, the expression of both type I receptor genes (*alk2*, *alk3*, *alk6a*) and type II

receptor genes (*bmpr2a*, *bmpr2b*, *actr2a*, *actr2b*) in ZFL cells transfected with poly(I:C) or infected with GCRV were

measured. Among these receptor genes, *alk6a* showed the highest expression upon poly(I:C) or virus challenge (Fig. 5a,

b). We also overexpressed all these receptors in ZFL and EPC cells, and examined the expression of *ifn* in these cells.

The upregulation of *ifnφ1* was only found in the alk6a-overexpressing ZFL cells (Fig. 5c). Moreover, the overexpression

of *alk6a* or *alk2* resulted in significantly increased expression of *ifnφ3* in ZFL cells and *ifn* in EPC cells, with overexpressed

alk6a being more effective (Fig. 5d, e). Based on these findings, the function of Alk6a was selected for further test.

Compared to wild-type ZFL cells, the expressions of antiviral protein genes *ifnφ1, ifnφ3* and *mx* as well as the antiviral

signaling genes *tbk1*, *irf3*, and *irf7* were all markedly upregulated in alk6a-overexpressed ZFL cells with or without GCRV

infection (Fig. 5f, g). The similar results were also observed in the alk6a-overexpressed EPC cells (Fig. 5h, i). Because

the phosphorylation of Gly-Ser (GS) domain of the type I receptor is required for its activation, we thus constructed

dominant-negative mutant of alk6a (alk6a-ΔGS) plasmid (Fig. 5j), and then transferred it into both ZFL and EPC cells. It

revealed that the dominant-negative mutation of *alk6a* significantly reduced the expression of *mx, ifn* (*ifnφ1* and *ifnφ3* in

ZFL cells), *tbk1*, *irf3*, and *irf7* in ZFL and EPC cells, with or without GCRV infection (Fig. 5k-n). The data suggested that

Alk6a was apparently involved in the Tbk1-Irf3/7-Ifn antiviral signaling.

Considering we have revealed that Bmp8a activated Tbk1-Irf3-Ifn antiviral signaling via p38 MAPK pathway, we then

examined whether the the levels of phosphorylated p38 MAPK protein were regulated by Alk6a. Clearly, compared to wild-

type ZFL or EPC cells, phosphorylation levels of p38 MAPK were significantly increased in alk6a-overexpressed both

types of the cells with or without GCRV infection (Fig. 6a-d). Moreover, dominant-negative mutation of *alk6a* significantly

reduced the phosphorylation levels of p38 MAPK in ZFL or EPC cells, with or without GCRV infection (Fig. 6e-h). Thus,

Alk6a was shown to be involved in the phosphorylation of p38 MAPK. Next, we investigated whether the antiviral immune

responses of Bmp8a were mediated by Alk6a. In the IFN promoter-driven luciferase assays, it was found that the ability

of Bmp8a to activate the IFN promoter was markedly blocked after the dominant-negative mutation of *alk6a* in EPC cells

(Fig. 6i-k). Importantly, co-immunoprecipitation (Co-IP) experiments confirmed that Bmp8a interacted with Alk6a (Fig. 6l).

Collectively, our data demonstrated that BMP type I receptor Alk6a interacted with Bmp8a, and participated in the antiviral

immunity. To our best knowledge, Alk6a is the first BMP receptor identified thus far which is being directly required for

antiviral immune responses.

Fig. 5 Alk6a is involved in the antiviral innate immune responses. a, **b** Expression of *alk2*, *alk3*, *alk6a*, *bmpr2a*,

bmpr2b, *actr2a*, *actr2b* mRNA in ZFL cells stimulated with GCRV (5 × 10⁴TCID₅₀ per ml, **a**) or poly(I:C) (2 μg/ml, **b**) for 48

h. **c-e** Expression of *ifnφ1* (**c**) and *ifnφ3* (**d**) mRNA in ZFL cells or EPC *ifn* (**e**) in EPC cells which were transfected with 2

μg of alk2, alk3, alk6a, bmpr2a, bmpr2b, actr2a, actr2b or empty vector for 48 h. **f, h** Expression of *irf3*, *irf7*, *tbk1*, *ifn* (or

ifnφ1 and *ifnφ3),* and *mx* mRNA after transfected with 2 μg of pcDNA3.1-alk6a or empty vector in ZFL (**f**) or EPC (**h**) cells

for 48 h. **g, i** Expression of *irf3*, *irf7*, *tbk1*, *ifn* (or *ifnφ1* and *ifnφ3),* and *mx* mRNA after transfected with 2 μg of of pcDNA3.1-

alk6a or empty vector in ZFL (**g**) or EPC (**i**) cells for 24 h, followed by infection with GCRV for another 36 h. **j** Schematic

drawing of the alk6a-ΔGS mutation that the GS domain of Alk6a was deleted. **k, m** Expression of *irf3*, *irf7*, *tbk1*, *ifn* (or

ifnφ1 and *ifnφ3),* and *mx* mRNA after transfected with 2 μg pcDNA3.1-alk6a-ΔGS or empty vector in ZFL (**k**) or EPC (**m**)

cells for 48 h. **l, n** Expression of *irf3*, *irf7*, *tbk1*, *ifn* (or *ifnφ1* and *ifnφ3),* and *mx* mRNA after transfected with 2 μg pcDNA3.1-

272   alk6a-ΔGS or empty vector in ZFL (**l**) or EPC (**n**) cells for 24 h, followed by infection with GCRV for another 36 h. The

273   expression of zebrafish *actb1* or EPC *actin* was used as an internal control for the qRT-PCR. Data were from three

274   independent experiments and were analyzed by Student's *t*-test (two-tailed) and were presented as mean ± SD (*$p < 0.05$,

275   **$p < 0.01$, *** $p < 0.001$, ns means no significant difference). Source data are provided as a Source Data file.

Fig. 6 Bmp8a activates the IFN expression through Alk6a. a, **c** Immunoblot analysis of phosphorylated (p-) p38 MAPK

after transfected with 2 μg alk6a or empty vector in ZFL (**a**) or EPC (**c**) cells. The cells were collected at 36 h or 48 h post

transfection for Immunoblot analysis. **b**, **d** Immunoblot analysis of phosphorylated (p-) p38 MAPK after transfected with 2

μg alk6a or empty vector in ZFL (**b**) or EPC (**d**) cells for 24 h, followed by infection with GCRV (5 × 10^4TCID$_{50}$ per ml) for

another 24 h or 36 h. **e, g** Immunoblot analysis of phosphorylated (p-) p38 MAPK after transfected with 2 µg alk6a-ΔGS

or empty vector in ZFL (**e**) or EPC (**g**) cells. The cells were collected at 36 h or 48 h post transfection for Immunoblot

analysis. **f**, **h** Immunoblot analysis of phosphorylated (p-) p38 MAPK after transfected with 2 µg alk6a or empty vector in

ZFL (**f**) or EPC (**h**) cells for 24 h, followed by infection with GCRV (5 × 10^{4}TCID$_{50}$ per ml) for another 24 h or 36 h. **i-k** EPC

cells were cotransfected with IFN-φ1pro-luc (200 ng, **i**), IFN-φ3pro-luc (200 ng, **j**) or EPC IFNpro-luc (200ng, **k**), and

bmp8a (100 ng) together with or without the dominant negative plasmids alk6a-ΔGS (100 ng). At 48 h post transfection,

the cells were collected for luciferase assays. *Renilla* luciferase was used as the internal control. **l** Co-immunoprecipitation

and immunoblot analysis of EPC cells cotransfected with alk6a-Flag (1 µg) and bmp8a-HA (1µg). Data were from three

independent experiments and were analyzed by Student's *t*-test (two-tailed) and were presented as mean ± SD (*** *p*<

0.001). Source data are provided as a Source Data file.

Virus induced expression of *bmp8a*. Next, we measured the expression of *bmp8a* in ZFL cells upon virus infection.

GCRV significantly induced the expression of *bmp8a* at 12 h post-stimulation, and reached a peak at 48 h (Fig. 7a). We

also detected the expression of *bmp8a* upon infection with GCRV or poly(I:C) *in vivo*. The expression of *bmp8a* in the liver,

kidney, intestine, and spleen of zebrafish infected with GCRV or poly(I:C) was all significantly elevated (Fig. 7b-i). This

indicated that the expression of *bmp8a* was inducible by infection with virus or its mimic poly(I:C).

Fig. 7 GCRV or poly(I:C) increases the expression of *bmp8a*. a Expression of *bmp8a* mRNA in ZFL cells after infected

with GCRV (5×10^4 TCID$_{50}$ per ml) for 48 h. **b-e** Expression of *bmp8a* mRNA in the liver (**b**), spleen (**c**), intestine (**d**), and

kidney (**e**) from zebrafish injected i.p. with poly(I:C) (10 μg/fish). **f-i** Expression of *bmp8a* mRNA in the liver (**f**), spleen (**g**),

intestine (**h**), and kidney (**i**) from zebrafish injected i.p. with 50μl of GCRV (10^8 TCID$_{50}$ per ml). Zebrafish injected i.p. with

PBS or MEM were used as the control. The expression of *actb1* served as a control for the qRT-PCR. Data were from

three independent experiments and were analyzed by Student's *t*-test (two-tailed) and were presented as mean ± SD (*p

< 0.05, **p < 0.01, ***p < 0.001, ns means no significant difference). Source data are provided as a Source Data file.

Binding of Stat1 to GAS motifs activated *bmp8a* expression. To explore how virus induces expression of *bmp8a*, we

searched for the transcription factor binding sites in *bmp8a* promoter region at the web http://jaspar.genereg.net/. Two

gamma-activated sites (GAS), 5'-ATTCCGGGAAA-3' (P1) and 5'-TTTACTAGAAC-3' (P2), in *bmp8a* promoter region were

identified (Supplementary Fig. 4). In mammals, the transcription factor STAT1 homodimer was known to bind to the GAS

motif in the promoter region of IFN-γ triggered downstream genes[11,16]. We thus wonder if Stat1 can bind to the *bmp8a*

promoter region to activate the expression of *bmp8a*. In zebrafish, there exist two Stat1 factors, Stat1a (GenBank

accession number NM_131480.1) and Stat1b (GenBank accession number NM_200091.2). Thus we constructed

dominant-negative mutant plasmids of stat1a-ΔC and stat1b-ΔC, that were transferred into ZFL cells, respectively. It was

found that the expression of *bmp8a* was significantly blocked in both cases (Fig. 8a, b). By contrast, the overexpression

of either *stat1a* or *stat1b* markedly promoted *bmp8a* promoter-driven luciferase activities (Fig. 8c). In addition, when one

of the GAS motifs in *bmp8a* promoter region was deleted, *bmp8a* promoter-driven luciferase activities were significantly

reduced; when both of the GAS motifs were deleted, *bmp8a* promoter-driven luciferase activities were reduced even more

significantly (Fig. 8d, e). These suggested that both Stat1a and Stat1b may interact with the GAS motifs.

To verity the binding of Stat1a/Stat1b to the GAS motifs in *bmp8a* promoter region, the recombinant proteins of both

Stat1a and Stat1b were expressed and purified (Fig. 8f, g), and used for electrophoretic mobility shift assay (EMSA). As

shown in Fig. 8h-k, a remarkable band of protein-DNA complex was observed (lane 2). The protein-DNA complex band

was obviously weaker in the presence of an excess unlabeled competitor probe, compared to that of biotin labeled probe

group (lane 3), but the intensity of the band showed little change in the presence of GAS mutant probes (lane 4). To detect

the specific binding, a supershift experiment with the antibody against His tag was performed. It revealed that His tag

antibody caused a specific supershift of slower migrating protein-DNA-antibody complex (Fig. 8h-k, lane 5). There was no

protein-DNA complex observed in the negative control group (Fig. 8h-k, lane 1). These demonstrated that both Stat1a and

Stat1b could directly bind to the GAS motifs in *bmp8a* promoter region, suggesting that *bmp8a* expression was subjected

to the regulation by binding of Stat1a/Stat1b to the GAS motifs.

Fig. 8 Stat1 binds the GAS sites and activates the *bmp8a* transcriptions upon virus stimulation. a, b Expression of

bmp8a mRNA in ZFL cells after transfected with stat1a-ΔC or stat1b-ΔC for 36 h (**a**) or 48 h (**b**). The expression of *actb1*

served as a control for the qRT-PCR. **c** Dual luciferase report assay was used to analyze the transcription abilities of

Stat1a and Stat1b in activation of *bmp8a* in EPC cells. pGl3-bmp8a (200 ng) was transfected into EPC cells together with stat1a (200 ng), stat1b (200 ng) or empty vector (200 ng). After 48 h, the transfected cells were collected for luciferase assays. **d** Schematic drawing of wild-type and GAS motif mutation Luc-report plasmids. **e** The 200 ng of pGL3-bmp8a-promoter, pGL3-bmp8a-promoter-ΔP1, pGL3-bmp8a-promoter-ΔP2 or pGL3-bmp8a-promoter-ΔP1 and ΔP2 was transfected into EPC cells along with stat1a (200 ng), stat1b (200 ng) or empty vector (200 ng), respectively. After 48 h, the transfected cells were collected for luciferase assays. *Renilla* luciferase was used as the internal control. **f, g** SDS-PAGE and Western-blotting analysis of rStat1a (**f**) and rStat1b (**g**). Lane M: protein molecular standard; Lane 1: negative control for IPTG induced *E. coli* (without rStat1a or rStat1b); Lane2: induced rStat1a or rStat1b (the whole cell lysate); Lane 3: induced rStat1a or rStat1b (the supernatant); Lane 4: purified rStat1a or rStat1b; Lane 5: western blot analysis of the sample in Lane 4. **h-k** EMSA was performed to validate the interaction of rStat1a or rStat1b with the GAS motif (P1 or P2) in the bmp8a promoter region. Lane 1: negative group; Lane 2: positive group; Lane 3: an excess unlabeled competitor probe; Lane 4: an excess unlabeled competitor probe containing a mutated runt binding site; Lane 5: Super-shift assays were performed by adding antibody against His tag. Data were from three independent experiments (**a–c, e**) or two independent experiments (**f-k**). Data were analyzed by Student's *t*-test (two-tailed) and were presented as mean ± SD (**p < 0.01,***p < 0.001). Source data are provided as a Source Data file.

Discussion

BMPs, canonical multi-functional growth factors, are suggested to play roles in regulation of immune responses, but it remains controversial over their functions in immunity. We show here for the first time that Bmp8a is a previously unrecognized factor involved in the regulation of antiviral immune responses in zebrafish. Evidences supporting this nature of Bmp8a include: Bmp8a inhibits RNA viruses replication *in vitro* and *in vivo*; Bmp8a promotes the expression of antiviral protein genes including type I IFNs and *mx*; the expression of *bmp8a* is induced by CGRV or poly(I:C), and Bmp8a activates TBK1-IRF3-IFN signaling. Bmp8a is a member of the TGF-β superfamily, which includes more than 30 genes encoding TGFβ, BMPs and activins. TGFβ has been shown to have both pro- and anti-inflammatory effects depending on the context it is acting in the immune system[45]. Like TGFβ, BMP members have also been shown to play different immunoregulatory roles[46]. BMP2, 4 and 7 deficiency or knockdown results in partial loss of the thymic capsule, reduced size of thymus and increased *Helicobacter pylori*-induced inflammation, whereas BMP6 inhibits macrophage growth by inducing cell cycle arrest *in vitro*[27,47-50]. The finding that Bmp8a is a positive regulator in antiviral immune responses provides a new angle for our understanding of the functions of TGF-β superfamily members including BMPs.

On one hand, it is generally regarded that the recognition of virus by PRRs elicits the activation of IRF3 and NF-κB, and induces the production of IFNα/β, that then activate JAK-STAT pathway, eventually leading to the production of ISGs

such as cytokines and antiviral proteins. On the other hand, BMPs are known to transmit signals through both Smad-

dependent pathways (smad1/5/8 and smad2/3 pathways) and Smad-independent pathways (ERK, JNK, and p38 MAPK

pathways). Here we clearly demonstrate that Bmp8a promotes the expression of endogenous type I IFNs independent of

Smad signaling pathways. Moreover, it is p38 MAPK pathway, rather than ERK and JNK pathways, that is involved in the

antiviral responses induced by Bmp8a. Furthermore, we show that Alk6a participates in Tbk1-Irf3/7-Ifn antiviral signaling

via interaction with Bmp8a. In addition, Alk6a was found to be required to induce the phosphorylation of p38 MAPK, which

is consistent with the findings in mouse that Alk6 knockdown suppresses p38 MAPK phosphorylation[51]. Previous studies

have shown that MAPK signaling pathway is crucial in the phosphorylation of TBK1 and IRF3 upon viral infection[42]. Thus,

we probably discover a new pathway of Bmp8a signaling in antiviral immune responses that Bmp8a acts as a positive

regulator through the promotion of phosphorylation of Tbk1 via p38 MAPK pathway, i.e., Bmp8a binds to Alk6a, which

induces p38 MAPK phosphorylation, and in turn enhances Tbk1 and Irf3 phosphorylation, eventually resulting in increased

synthesis of type I IFN (Fig. 9).

Our study also show that the challenge with GCRV and poly(I:C) both significantly promotes the expression of

zebrafish *bmp8a*. Searching for the binding sites of transcript factors in *bmp8a* promoter region reveals that two IFN-γ

activation sites (GAS) that may interact with STAT1 are identified. Functionally, we demonstrate that Stat1a and Stat1b,

orthologues of human STAT1, can directly bind the GAS sites in *bmp8a* promoter, resulting in activation of *bmp8a*

expression. It has been reported that in black carp, Stat1a and Stat1b can both form homodimer and heterodimer *in vivo*

like their mammalian counterpart[52]. It is thus highly likely that in zebrafish, Stat1a and Stat1b can form homodimer or

heterodimer, which then binds to the GAS sites in *bmp8a* promoter region. In zebrafish, IFN-γ has been found to be able

to induce the expression of both *stat1a* and *stat1b*, and the GAS motif is required for Ifn-γ activation[53,54]. It is widely known

that the JAK-STAT pathway is one of the most important candidate pathways through which IFN-γ works. Activation of

JAK-STAT signaling leads to transcription of various downstream ISGs for antiviral activity[55]. We thus think that Bmp8a is

another important antiviral factor which can be activated through Ifn-γ-Jak-Stat1 pathway (Fig. 9).

In summary, we demonstrate for the first time that Bmp8a is a positive regulator in antiviral immune responses, which

functions through promotion of phosphorylation of Tbk1 via p38 MAPK pathway. Upon virus infection, the expression of

bmp8a is activated by Stat1a/Stat1b directly. Bmp8a knockdown leads to significantly decreased phosphorylation of Tbk1

and Irf3. As a consequence, *bmp8a*-deficient zebrafish produces significantly lowered type I IFNs in response to RNA

virus infection, exhibits significantly reduced immune responses, and increases significantly higher viral load and morbidity

in vivo. Our study is significant to understand the regulatory mechanism of BMP involving in antiviral immune response

and provides a new target for controlling viral infection.

Fig. 9 Schematic illustration of Bmp8a serving as positive regulator in the antiviral immune. Upon virus infection,

the transcriptions of *bmp8a* are activated through the Jak-Stat1 pathway. The Bmp8a binds to BMP type I receptor Alk6a,

promoting phosphorylation of Tbk1 and Irf3 to induce the expression of *Ifn* through p38 MAPK pathway.

Methods

Cells and viruses. The epithelioma papulosum cyprini cells (EPC), zebrafish liver cells (ZFL) were acquired from the

China Zebrafish Resource Center (CZRC) and cultured according to the instructions from the CZRC. EPC and ZFL cells

were grown in a 28 °C incubator supplied with 5% CO_2. The flounder gill cells (FG) were grown in a 22 °C incubator

supplied. The EPC or FG cells were cultured in MEM media supplemented with 10% fetal bovine serum (FBS) (Gibco),

100 U/ml penicillin and 100 μg/ml streptomycin. The ZFL cells were maintained in DMEM/F-12 media supplemented with

10% FBS, 100 U/ml penicillin and 100 μg/ml streptomycin. Grass carp reovirus (GCRV), a dsRNA virus, and spring viremia

of carp virus (SVCV), a negative ssRNA virus, were kindly provided by Yibing Zhang (Institute of Hydrobiology, Chinese

Academy of Sciences), and tittered according to the method of Reed and Muench[56], by a 50% tissue culture infective dose

($TCID_{50}$) assay on EPC cells. The turbot skin verruca disease virus (TSVDV) was isolated from the focal site of turbot skin

in our lab.

Zebrafish. The animals used in the experiment followed the ethical guidelines established by the Institutional Animal Care

and Use Committee of the Ocean University of China (permit number, SD2007695). The zebrafish *bmp8a* knockout mutant

lines (*bmp8a*[-/-]) were established from the zebrafish AB line using TALENs technology. The TALENs for zebrafish *bmp8a*

were assembled using the Unit Assembly (UA) method as previously described[57]. All the UA method starting vectors were

the gifts from laboratories of Bo Zhang at Peking University and of Shuo Lin at University of California Los Angeles.

TALENs mRNAs were transcribed using the mMESSAGE mMACHINE SP6 kit (Invitrogen, #AM1340) and purified using

the RNeasy Mini kit (QIAGEN, #74106). To generate zebrafish mutant lines, TALEN mRNAs (50–200 pg) were

microinjected into 1-cell stage zebrafish embryos. Two days after injection, genomic DNA was isolated from 8 to 10 pooled

larvae. The target genomic regions were amplified by nested PCR and subcloned into the pGEM-T vector. Single colonies

were genotyped by sequencing. To obtain germline mutations, the TALEN-injected embryos were raised to adulthood and

outcrossed with wild-type (WT) fish. The F1 progeny were genotyped by sequencing. To obtain homozygous mutants,

heterozygous mutant of the same mutation were obtained and self-crossed. The primers used in this study were listed in

Supplemental Table 1.

Plasmid construction. The open reading frames (ORF) of zebrafish bmp8a, stat1a, stat1b, alk2, alk3, alk6a, actr2a,

actr2b, bmpr2a and bmpr2b were cloned into the pcDNA3.1 expression vector for the eukaryotic expression. The ORF of

zebrafish stat1a and stat1b were cloned into the pET-28a expression vector for the expression of recombinant proteins.

The promoter regions of bmp8a, bmp8a-ΔP1 (delete the GAS motif of 5´-attccgggaaa-3´ in the bmp8a promoter), bmp8a-

ΔP2 (delete another GAS motif of 5´-tttactagaac-3´ in the bmp8a promoter), bmp8a-ΔP1and ΔP2 (delete both the two

GAS motifs), IFNφ1, IFNφ3, and EPC IFN were cloned into the pGL3-basic vector. Dominant negative mutant plasmids

including tbk1-K38M, stat1a-ΔC, stat1b-ΔC, irf3DN, and irf7DN were described previously[58,59]. Dominant negative mutant

plasmids alk6a-ΔGS (delete the sequence of 5´-agctccggttctggctcagga-3´ encoding the GS domain) was cloned into the

pcDNA3.1 expression vector. The deletions and mutations were created using Mut Express II Fast Mutagenesis Kit V2

(Vazyme, #C214-01). The primers used in this study were listed in Supplemental Table 2.

Transcriptome Sequencing. The zebrafish were fasted for 24 h prior to tissue collection. Three mutant (*bmp8a*$^{-/-}$) and

three wild type (WT) adult zebrafish (0.4 g ± 0.05 g) were randomly selected from each group and were injected

intraperitoneally with 50 µl of GCRV (10^8TCID$_{50}$ per ml) per fish. Seventy-two hours after the injection, the livers from three

fish each group were pooled together for total RNA extraction. Three independent samples from each group were prepared

for transcriptome sequencing. mRNAs were enriched by magnetic beads combining the oligo (dT). cDNA fragments were

subjected to the end repair process by the addition of a single 'A' base and the ligation of Illumina sequencing adapters.

The ligation products were selected by agarose gel electrophoresis, amplified by PCR, and sequenced using Illumina

HiSeq 4000 by Annoroad Technology (Beijing, China).

Identification of DEGs. The RPKM (reads per kb per million reads) method was used to calculate and normalize the

abundances of genes[60]. DEGs across groups were identified by the edgeR package (http://www.r-project.org/). Genes

with a fold change ≥ 2 and a false discovery rate (FDR) < 0.05 in a comparison were regarded as significantly expressed.

DEGs were then subjected to enrichment analysis of KEGG pathways.

Viral infection *in vitro*. EPC, FG, or ZFL cells were plated 24 h before infection. Cells were infected with GCRV (5 ×

$10^4 TCID_{50}$ per ml) or TSVDV (crude virus extracts filtered by a 0.45 μm microporous membrane) for 1 h in medium without

FBS. After infection, cells were washed with PBS and then medium was added with FBS. The cells and cell-free

supernatants were harvested at the indicated times.

Viral infection *in vivo*. For *in vivo* viral infection, the adult WT and *bmp8a*[-/-] zebrafish were infected via intraperitoneal

injection using 50 μL of SVCV ($10^8 TCID_{50}$ per ml), GCRV ($10^8 TCID_{50}$ per ml), or TSVDV (crude virus extracts filtered by a

0.45 μm microporous membrane) per fish. Total RNA was extracted and viral RNA expression in the liver, kidney, intestine,

and spleen were examined by qRT-PCR. For the survival experiments, zebrafish were monitored for survival after SVCV,

GCRV, or TSVDV infection.

Crystal violet staining. EPC or ZFL cells were transfected with bmp8a or empty vector plasmid. At 24 h post transfection,

cells were treated with GCRV (5 × $10^4 TCID_{50}$ per ml). At 72 h after virus challenge, cells were fixed with 4% formaldehyde

for 1 h and stained with 0.5% crystal violet for 2 h. After washing with tap water, visible plaques were photographed.

Luciferase assays. EPC cells were seeded in 24-well plates overnight and cotransfected with various plasmids using

FuGENE HD Transfection Reagent (Promega, #E2311) following the manufacturer's instruction. If necessary, the cells

were infected with GCRV (5 × $10^4 TCID_{50}$ per ml). At the indicated time points, luciferase activities were measured with the

Luc-Pair Duo-Luciferase Assay Kits 2.0 (iGene Biotechnology, #LF002). Data were normalized by calculating the ratio

between firefly luciferase activity and *Renilla* luciferase activity. The primers used for cloning the promoter are shown in

Supplemental Table 2.

RNA interference. Small interfering RNA (siRNA) of bmp8a and si-NC (negative control) were purchased from RiboBio.

The oligonucleotide sequences (siRNA) specific targeting *bmp8a* mRNA was as follows: 5′-GTTCTTCAGAGCTAGTCAA-

3′. Transfection was performed with Lipofectamine RNAiMAX (Invitrogen, #13778075) according to the manufacturer's

protocols. Total RNA was extracted for qRT-PCR analysis.

Drug treatment and analysis. SMAD1/5/8 inhibitor DMH1(Selleck, #S7146), smad2/3 inhibitor TP0427736 HCl (Selleck,

#S8700), p38 MAPK inhibitor SB203580 (Beyotime, #S1863), JNK inhibitor SP600125 (Beyotime, #S1876), and MEK1/2

inhibitor U0126 (Beyotime, #S1901) were dissolved in dimethyl ulfoxide (DMSO). ZFL or EPC cells were treated with each

inhibitor, which were diluted with culture medium at concentrations of 10 μM for 24 h. Total cellular RNA was extracted for

qRT-PCR analysis.

RNA quantification. Total RNA was isolated using the miRNeasy Mini Kit (QIAGEN, #217004) according to the

manufacturer's instructions. RNA treated with DNase and the cDNAs were synthesized with PrimeScript™ RT reagent Kit

with gDNA Eraser (TaKaRa, #RP047A). Samples without reverse transcriptase were also added as control. Gene

expression was determined by amplifying the cDNA with ChamQ SYBR Color qPCR Master Mix(Vazyme, #Q431-02) by

an ABI 7500 Fast Real-Time PCR System (Applied Biosystems). The expression of zebrafish *actb1* or EPC *actin* was

used as an internal control. The $2^{-\Delta\Delta Ct}$ method was used to calculate relative expression changes. All qRT-PCR

experiments were performed in triplicate and repeated three times. Primers used are listed in Supplementary Table 1.

Preparation of recombinant protein. The recombinant protein of rStat1a or rStat1b was expressed and purified as

previous description[61]. Briefly, the recombinant plasmid (pET-28a-stat1a or pET-28a-stat1b) was transformed into *Trans*B

(DE3) chemically competent cell. The positive transformants were incubated in liquid LB medium containing 50 μg/mL

kanamycin at 37 °C by shaking at 160 rpm. When the cells grow to OD600 = 0.5, isopropyl β-D-thiogalactoside (IPTG)

was added to a final concentration of 1 mM, and incubated at 19 °C with shaking at 120 rpm for 14 h. After incubation, the

cells were harvested by centrifugation, re-suspended in PBS, and broken by ultrasound. The supernatant was centrifuged

at 12,000 g for 20 min, and the protein was purified by Ni-NTA Sepharose column. The concentration of purified protein

was determined by BCA (bicinchonininc acid) method. The purified recombinant proteins were stored at -80 °C for

subsequent experiment.

Electrophoretic mobility shift assay. Electrophoretic mobility shift assay (EMSA) was performed according to the

instruction of Chemiluminescent EMSA Kit (Beyotime, #GS009). Oligonucleotides (Supplementary Table 1) were

synthesized and biotinylated by Sangon Biotech Company as required and the complementary oligonucleotides were

annealed to form the double DNA probe. The recombinant protein was pre-incubated in EMSA/Gel-Shift binding buffer

with poly (dI-dC) at 25 °C for 25 min in the presence or absence of Non-Biotin probe (0.5 μM). Then, all samples were

mixed with biotinylated probes (0.5 μM) and incubated at 25°C for 30 min. The binding reaction without recombinant

protein was set as the negative control. For supershift analysis, anti-His tag antibody was incubated with the reaction

mixture for another 30 min after the labeled probe was added. After separation in a native 6% polyacrylamide gel, the free

probes and DNA-protein compound were transferred to a Hybond-N⁺ nylon membrane, followed by crosslinking under UV

light. After immersion in the blocking buffer, membranes were incubated with streptavidin conjugated HRP for 20 min with

shaking and then washed three times with washing buffer. Membranes were exposed briefly in Western Lightning-ECL

and then exposure to Automatic X-ray Film Processor (Smpicgg).

Immunoblot analysis and co-immunoprecipitation. For Immunoblot analysis, cells were lysed in NP-40 buffer (150

mM NaCl, 0.5% EDTA, 50 mM Tris, 1% NP40, proteinase inhibitors, and protein phosphatase inhibitors) for 30 min on ice.

For co-immunoprecipitation (co-IP), whole-cell extracts were collected 48 h after transfection and lysed in NP-40 buffer on

ice for 30 min. After centrifugation for 15 min at 12,000 × g, 4 °C, supernatants were collected and incubated with Protein

A/G PLUS-Agarose (Santa Cruz, #sc-2003) coupled to specific antibodies for overnight with rotation at 4°C. Beads were

washed 3 × times with NP-40 buffer. Bound proteins were eluted by boiling 7min with 2 x SDS sample buffer. For

immunoblot analysis, immunoprecipitates or whole-cell lysates were separated by 12.5% SDS–PAGE gels, electro-

transferred to PVDF membranes and blocked for 4 h with 5% no-fat milk solution, followed by blotting with the appropriate

antibodies and detection by enhanced chemiluminiscence (ECL). Antibodies from Cell Signaling Technology: TBK1

(1:1000, #3504T), p-TBK1(Ser172) (1:1000, #5483T). Antibodies from Bioss: IRF3 (1:1000, #bs-2993R), p-IRF3 (Ser386)

(1:1000, #bsm-52170R), p38MAPK (1:1000, #bs-0637R), p-p38MAPK (Thr180 + Tyr182) (1:1000, #bs-2210R), Actin

(1:2000, #bs-0061R). Antibodies from Beyotime: HA tag (1:1000, #AH158) and Flag tag (1:1000, #AF519). Antibodies

from CWBIO: His tag (1:5000, #CW0285) and goat anti-rabbit IgG HRP secondary antibody (1:8000, #CW0103S).

Antibodies from Abcam: VeriBlot for IP Detection (1:5000, #ab131366).

Statistical analysis and reproducibility. Representative experiments have been repeated at least two to three times.

All statistical analysis was performed using Graphpad Prism 8.0.1. Data are presented as mean ± SEM. Statistical

significance was assessed using the unpaired, two-tailed Student's *t*-test. Survival analyses were performed using the

Kaplan-Meier method and assessed using the log-rank (Mantel-Cox) test. *$p < 0.05$, **$p < 0.01$, *** $p < 0.001$; ns, not

significant, $p > 0.05$.

Data availability. All data are available from the corresponding author upon reasonable request and that a Source Data

file is available for this article.

References

1. Thaiss, C. A., Levy, M., Itav, S. & Elinav, E. Integration of Innate Immune Signaling. *Trends Immunol*. **37**, 84-101

(2016).

2. Akira, S. & Hemmi, H. Recognition of pathogen-associated molecular patterns by TLR family. *Immunol. Lett*. **85**,

85-95 (2003).

3. Chen, S. N., Zou, P. F. & Nie, P. Retinoic acid-inducible gene I (RIG-I)-like receptors (RLRs) in fish: current

knowledge and future perspectives. *Immunology* **151**, 16-25 (2017).

4. Keating, S. E., Baran, M. & Bowie, A. G. Cytosolic DNA sensors regulating type I interferon induction. *Trends*

Immunol. **32**, 574-581 (2011).

5. Parvatiyar, K. et al. The helicase DDX41 recognizes the bacterial secondary messengers cyclic di-GMP and cyclic

di-AMP to activate a type I interferon immune response. *Nat. Immunol*. **13**, 1155-1161 (2012).

6. Zhou, P. et al. MLL5 suppresses antiviral innate immune response by facilitating STUB1-mediated RIG-I

degradation. *Nat. Commun*. **9**, 1243 (2018).

7. Honda, K. et al. IRF-7 is the master regulator of type-I interferon-dependent immune responses. *Nature* **434**, 772-

777 (2005).

8. Tamura, T., Yanai, H., Savitsky, D. & Taniguchi, T. The IRF family transcription factors in immunity and oncogenesis.

Annu. Rev. Immunol. **26**, 535-584 (2008).

9. Li, W., Hofer, M. J., Jung, S. R., Lim, S. L. & Campbell, I. L. IRF7-dependent type I interferon production induces

lethal immune-mediated disease in STAT1 knockout mice infected with lymphocytic choriomeningitis virus. *J. Virol*.

88, 7578-7588 (2014).

10. Schindler, C. & Plumlee, C. Inteferons pen the JAK-STAT pathway. *Semin. Cell Dev. Biol*. **19**, 311-318 (2008).

11. Decker, T., Kovarik, P. & Meinke, A. GAS elements: a few nucleotides with a major impact on cytokine-induced

gene expression. *J. Interferon Cytokine Res*. **17**, 121-134 (1997).

12. Fu, X. Y., Kessler, D. S., Veals, S. A., Levy, D. E. & Darnell, J. E. Jr. ISGF3, the transcriptional activator induced

by interferon alpha, consists of multiple interacting polypeptide chains. *Proc. Natl Acad. Sci. USA* **87**, 8555-8559

(1990).

13. Levy, D. E., Kessler, D. S., Pine, R. & Darnell, J. E. Jr. Cytoplasmic activation of ISGF3, the positive regulator of

interferon-alpha-stimulated transcription, reconstituted in vitro. *Genes Dev*. **3**, 1362-1371 (1989).

14. Levy, D. E., Kessler, D. S., Pine, R., Reich, N. & Darnell, J. E. Jr. Interferon-induced nuclear factors that bind a

shared promoter element correlate with positive and negative transcriptional control. *Genes Dev*. **2**, 383-393 (1988).

15. Ren, Y. et al. Deubiquitinase USP2a Sustains Interferons Antiviral Activity by Restricting Ubiquitination of Activated

STAT1 in the Nucleus. *PLoS Pathog.* **12**, e1005764 (2016).

16.   Shuai, K., Schindler, C., Prezioso, V. R. & Darnell, J. E. Jr. Activation of transcription by IFN-gamma: tyrosine

phosphorylation of a 91-kD DNA binding protein. *Science* **258**, 1808-1812 (1992).

17.   Bragdon, B., Moseychuk, O., Saldanha, S., King, D., Julian, J. & Nohe, A. Bone morphogenetic proteins: a critical

review. *Cell Signal* **23**, 609-620 (2011).

18.   Crisan, M. et al. BMP signalling differentially regulates distinct haematopoietic stem cell types. *Nat. Commun.* **6**,

8040 (2015).

19.   Wu, M., Chen, G. & Li, Y. P. TGF-beta and BMP signaling in osteoblast, skeletal development, and bone formation,

homeostasis and disease. *Bone Res.* **4**, 16009 (2016).

20.   Xue, Y. et al. Organizer-derived Bmp2 is required for the formation of a correct Bmp activity gradient during

embryonic development. *Nat. Commun.* **5**, 3766 (2014).

21.   Yu, Y., Mutlu, A. S., Liu, H. & Wang, M. C. High-throughput screens using photo-highlighting discover BMP signaling

in mitochondrial lipid oxidation. *Nat. Commun.* **8**, 865 (2017).

22.   Huber, S. & Schramm, C. Role of activin A in the induction of Foxp3+ and Foxp3- CD4+ regulatory T cells. *Crit.*

Rev. Immunol. **31**, 53-60 (2011).

23.   Martínez, V. G. et al. The canonical BMP signaling pathway is involved in human monocyte-derived dendritic cell

maturation. *Immunol. Cell Biol.* **89**, 610-618 (2011).

24.   Martínez, V. G. et al. The BMP Pathway Participates in Human Naive CD4+ T Cell Activation and Homeostasis.

PloS One **10**, e0131453 (2015).

25.   Phillips, D. J., de Kretser, D. M. & Hedger, M. P. Activin and related proteins in inflammation: not just interested

bystanders. *Cytokine Growth Factor Rev.* **20**, 153-164 (2009).

26.   Seeger, P., Musso, T. & Sozzani, S. The TGF-β superfamily in dendritic cell biology. *Cytokine Growth Factor Rev.*

26, 647-657 (2015).

27.   Takabayashi, H. et al. Anti-inflammatory activity of bone morphogenetic protein signaling pathways in stomachs of

mice. *Gastroenterology* **147**, 396-406 (2014).

28.   Yasmin, N. et al. Identification of bone morphogenetic protein 7 (BMP7) as an instructive factor for human

epidermal Langerhans cell differentiation. *J. Exp. Med.* **210**, 2597-2610 (2013).

29.   Sivertsen, E. A., Huse, K., Hystad, M. E., Kersten, C., Smeland, E. B. & Myklebust, J. H. Inhibitory effects and

target genes of bone morphogenetic protein 6 in Jurkat TAg cells. *Eur. J. Immunol.* **37**, 2937-2948 (2007).

30.   Varas, A. et al. Interplay between BMP4 and IL-7 in human intrathymic precursor cells. *Cell cycle* **8**, 4119-4126

(2009).

31.   Chen, W. & Ten Dijke, P. Immunoregulation by members of the TGFβ superfamily. *Nat. Rev. Immunol.* **16**, 723-740

600   (2016).

601   32.   Zhao, G. Q., Deng, K., Labosky, P. A., Liaw, L. & Hogan, B. L. The gene encoding bone morphogenetic protein 8B

602   is required for the initiation and maintenance of spermatogenesis in the mouse. *Genes Dev.* **10**, 1657-1669 (1996).

603   33.   Zhao, G. Q. & Hogan, B. L. Evidence that mouse Bmp8a (Op2) and Bmp8b are duplicated genes that play a role

604   in spermatogenesis and placental development. *Mech. Dev.* **57**, 159-168 (1996).

605   34.   Walton, K. D. et al. Villification in the mouse: Bmp signals control intestinal villus patterning. *Development* **143**,

606   427-436 (2016).

607   35.   Ying, Y., Qi, X. & Zhao, G. Q. Induction of primordial germ cells from murine epiblasts by synergistic action of BMP4

608   and BMP8B signaling pathways. *Proc. Natl Acad. Sci. USA* **98**, 7858-7862 (2001).

609   36.   Zhong, S. j. et al. Spatial and temporal expression of *bmp8a* and its role in regulation of lipid metabolism in

610   zebrafish *Danio rerio. Gene Rep.* **10**, 33-41 (2018).

611   37.   Robertsen, B. The interferon system of teleost fish. *Fish Shellfish Immunol.* **20**, 172-191 (2006).

612   38.   Zhang, Y. B. et al. The innate immune response to grass carp hemorrhagic virus (GCHV) in cultured Carassius

613   auratus blastulae (CAB) cells. *Dev. Comp. Immunol.* **31**, 232-243 (2007).

614   39.   Derynck, R., Akhurst, R. J. & Balmain, A. TGF-beta signaling in tumor suppression and cancer progression. *Nat.*

615   *Genet.* **29**, 117-129 (2001).

616   40.   Drummond, A. E. TGFbeta signalling in the development of ovarian function. *Cell Tissue Res.* **322**, 107-115 (2005).

617   41.   Oh, S. P., Yeo, C. Y., Lee, Y., Schrewe, H., Whitman, M. & Li, E. Activin type IIA and IIB receptors mediate Gdf11

618   signaling in axial vertebral patterning. *Genes Dev.* **16**, 2749-2754 (2002).

619   42.   Shi, Y. et al. Exosomal Interferon-Induced Transmembrane Protein 2 Transmitted to Dendritic Cells Inhibits

620   Interferon Alpha Pathway Activation and Blocks Anti-Hepatitis B Virus Efficacy of Exogenous Interferon Alpha.

621   *Hepatology* **69**, 2396-2413 (2019).

622   43.   Heldin, C. H., Miyazono, K. & ten Dijke, P. TGF-beta signalling from cell membrane to nucleus through SMAD

623   proteins. *Nature* **390**, 465-471 (1997).

624   44.   Shi, Y. & Massagué, J. Mechanisms of TGF-beta signaling from cell membrane to the nucleus. *Cell* **113**, 685-700

625   (2003).

626   45.   Worthington, J. J., Fenton, T. M., Czajkowska, B. I., Klementowicz, J. E. & Travis, M. A. Regulation of TGFβ in the

627   immune system: an emerging role for integrins and dendritic cells. *Immunobiology* **217**, 1259-1265 (2012).

628   46.   Tu, E., Chia, P. Z. & Chen, W. TGFβ in T cell biology and tumor immunity: Angel or devil? *Cytokine Growth Factor*

629   *Rev.* **25**, 423-435 (2014).

47.   Aleman-Muench, G. R. & Soldevila, G. When versatility matters: activins/inhibins as key regulators of immunity.

Immunol. Cell Biol. **90**, 137-148 (2012).

48.   Bleul, C. C. & Boehm, T. BMP signaling is required for normal thymus development. *J. Immunol*. **175**, 5213-5221

(2005).

49.   Gordon, J., Patel, S. R., Mishina, Y. & Manley, N. R. Evidence for an early role for BMP4 signaling in thymus and

parathyroid morphogenesis. *Dev. Biol*. **339**, 141-154 (2010).

50.   Lee, M. K. et al. TGF-beta activates Erk MAP kinase signalling through direct phosphorylation of ShcA. *EMBO J*.

26, 3957-3967 (2007).

51.   Itoh, S. et al. Trps1 plays a pivotal role downstream of Gdf5 signaling in promoting chondrogenesis and apoptosis

of ATDC5 cells. *Genes Cells* **13**, 355-363 (2008).

52.   Wu, H., Zhang, Y., Lu, X., Xiao, J., Feng, P. & Feng, H. STAT1a and STAT1b of black carp play important roles in

the innate immune defense against GCRV. *Fish Shellfish Immunol*. **87**, 386-394 (2019).

53.   Zou, J. & Secombes, C. J. Teleost fish interferons and their role in immunity. *Dev. Comp. Immunol*. **35**, 1376-1387

(2011).

54.   Ruan, B. Y. et al. Two type II IFN members, IFN-γ and IFN-γ related (rel), regulate differentially IRF1 and IRF11 in

zebrafish. *Fish Shellfish Immunol*. **65**, 103-110 (2017).

55.   Boehm, U., Klamp, T., Groot, M. & Howard, J. C. Cellular responses to interferon-gamma. *Annu. Rev. Immunol*.

15, 749-795 (1997).

56.   Reed, L. J. & Muench, H. A Simple Method of Estimating Fifty Percent Endo-Points. *Am. J. Hyg*. **27**, 493-797

(1938).

57.   Huang, P., Xiao, A., Tong, X., Zu, Y., Wang, Z. & Zhang, B. TALEN construction via "Unit Assembly" method and

targeted genome modifications in zebrafish. *Methods* **69**, 67-75 (2014).

58.   Feng, H., Zhang, Y. B., Zhang, Q. M., Li, Z., Zhang, Q. Y. & Gui, J. F. Zebrafish IRF1 regulates IFN antiviral

response through binding to IFNφ1 and IFNφ3 promoters downstream of MyD88 signaling. *J. Immunol*. **194**, 1225-

1238 (2015).

59.   Sun, F., Zhang, Y. B., Liu, T. K., Shi, J., Wang, B. & Gui, J. F. Fish MITA serves as a mediator for distinct fish IFN

gene activation dependent on IRF3 or IRF7. *J. Immunol*. **187**, 2531-2539 (2011).

60.   Mortazavi, A., Williams, B. A., McCue, K., Schaeffer, L. & Wold, B. Mapping and quantifying mammalian

transcriptomes by RNA-Seq. *Nat. Methods* **5**, 621-628 (2008).

61.   Sun, C., Hu, L., Liu, S., Gao, Z. & Zhang, S. Functional analysis of domain of unknown function (DUF) 1943,

DUF1944 and von Willebrand factor type D domain (VWD) in vitellogenin2 in zebrafish. *Dev. Comp. Immunol*. **41**,

661   469-476 (2013).

662 62. Qin, L., Wang. Y. G. & Zhang, Z. Multiple verrucous protrusions on skin in turbot Scophthalmus maximus. *J. Dalian*

663   *Ocean Univ*. **23**, 479-483 (2008) (in Chinese with English abstract).

664
665 **Author contributions**

666  Z.L.and S.-C.Z. conceived and coordinated the project. Z.L., G.J., and S.Z. designed the experiments. S.Z., H.L. and

667 Y.-S.W. performed experiments. H.-Y.L. and Y.W. contributed to establish the *bmp8a* knockout mutant lines (*bmp8a*$^{-/-}$)

668 using TALENs technology. S.Z. performed the statistical analysis. Z.L., S.-C.Z. and S.Z. wrote the manuscript, with input

669 from the other authors. All authors reviewed the manuscript and approved the final version.

671 **Acknowledgements**

672  This work was supported by the grants of National Natural Science Foundation of China (31572259) and National

673 Key Research and Development Project of the Ministry of Science and Technology（2018YFD0900505）.

674
675 **Conflict of Interest**

676  The work is under a patent in China "A method to improve the antiviral immunity of fish (Application No.

677 201910454499.6)".