Bmp8a Is a Novel Player in Regulation of Antiviral Immunity

Shenjie Zhong[1,2], Haoyi Li[1,2], Yun-Sheng Wang[1,2], Ying Wang[1,2], Guangdong Ji[1,2], Hong-Yan Li[1,2], Shicui

Zhang[1,2]*, Zhenhui Liu[1,2]**

[1]College of Marine Life Science and Institute of Evolution & Marine Biodiversity, Ocean University of China,

Qingdao 266003, China

[2]Laboratory for Marine Biology and Biotechnology, Pilot National Laboratory for Marine Science and Technology

(Qingdao), Qingdao 266003, China

24 *Correspondence author: e-mail: sczhang@ouc.edu.cn

25 **Correspondence author: e-mail: zhenhuiliu@ouc.edu.cn

Abstract

Bone morphogenetic protein (BMP) is a kind of classical multi-functional growth factor that plays a vital role in the

formation and maintenance of bone, cartilage, muscle, blood vessels, and the regulation of energy balance.

However, whether BMP plays a role in antiviral immunity is unknown. Here we demonstrate that Bmp8a is a newly-

identified positive regulator for antiviral immune responses. The *bmp8a*$^{-/-}$ zebrafish, when infected with the viruses

of GCRV, SVCV or TSVDV, show significantly reduced antiviral immunity, increased viral load and morbidity. We

also show for the first time that Bmp8a interacts with Alk6a, which promotes the phosphorylation of Tbk1 and Irf3

through p38 MAPK pathway, and induces the production of type I IFNs in response to virus infection. Upon virus

infection, *bmp8a* expression is activated through the binding of Stat1a/Stat1b to the GAS motifs in *bmp8a* promoter

region, enlarging the antiviral innate immune signal. Our study uncovers a previously unrecognized role of Bmp8a

in regulation of antiviral immune responses and provides a new target for controlling viral infection.

Introduction

Viruses infect all groups of living things and produce a variety of diseases. Host cell detection of viral infection depends on the recognition of pathogen-associated molecular patterns (PAMPs) from viral proteins or nucleic acids by the pattern recognition receptors (PRRs) such as Toll-like receptors (TLRs), RIG-I-like receptors (RLRs), NOD-like receptors (NLRs) and cytoplasmic DNA or RNA sensors (*Thaiss et al., 2016; Akira et al., 2003; Chen et al., 2017; Keating et al., 2011; Parvatiyar et al., 2012*). The PRRs, once stimulated by their appropriate ligands, trigger distinct intracellular signaling pathways that converge on the activation of IFN regulatory factor 3 (IRF3) and/or IFN regulatory factor 7 (IRF7), which then translocate into the nucleus and activate the production of interferons (IFNs) that play important roles in inhibiting virus replication (*Honda et al., 2005; Tamura et al., 2008*).

IFNs are classified into type I IFNs that include IFNα and IFNβ; type II IFN which has only a single member IFN-γ; and type III IFNs that consist of IFN-λs (*Li et al., 2014*). All these three types of IFNs trigger intracellular signaling cascades generally via the Janus kinase signal transducer and activator of transcription (JAK-STAT) pathway (*Schindler et al., 2008*). Canonically, type I IFNs initiate the signaling via binding to a heterodimeric receptor complex IFN-α/β receptor (IFNAR), inducing phosphorylation of transcription factors STAT1 and STAT2. The phosphorylated STATs form a trimolecular complex with IRF9, which translocates to the nucleus and binds to IFN-stimulated response elements (ISREs) of IFN-stimulated genes (ISGs), resulting in transcriptional activation of the target ISGs, including those encoding numerous cytokines and antiviral proteins (*Decker et al., 1997; Fu et al., 1990; Levy et al., 1989; Levy et al., 1988; Ren et al., 2016; Shuai et al., 1992*). In contrast, type II IFN binds to its unique receptor, IFN-γ receptor (IFNGR), resulting in the phosphorylation of STAT1 and the formation of STAT1 homodimers that recognize gamma-activated sequences (GASs) present in the promoter regions of IFN-γ-regulated genes (*Li et al., 2014*).

Bone morphogenetic proteins (BMPs) are potent growth factors belonging to the transforming growth factor beta (TGFβ) superfamily, which are intercellular signaling molecules with multiple functions in development and differentiation, such as skeletal formation and hematopoiesis, as well as lipid oxidation (*Bragdon et al., 2011; Wu et al., 2016*). However, accumulating data suggests that BMPs also play important roles in the regulation of immune responses (*Huber et al., 2011; Martínez et al., 2011; Phillips et al., 2009; Seeger et al., 2015; Takabayashi et al., 2014*). For example, BMP7 has been shown to drive Langerhans cell differentiation (*Yasmin et al., 2013*). Interestingly, BMP signaling appears to have different responses in T cells: BMP4 and BMP6 were shown to promote T cell proliferation, whereas BMP2 was seen to inhibit T cell proliferation (*Sivertsen et al., 2007; Varas et al., 2009*). In addition, although BMPs were found to promote tumor growth by inhibiting the functions of

cytotoxic T lymphocytes (CTLs) and dendritic cells (DCs), they were also shown to inhibit cancer cell proliferation

and promote the activity of NK cells that typically display antitumour activity (**Chen et al., 2016**). Therefore, it

remains controversial regarding the functions of BMPs in immune responses.

In humans and mice, two closely related BMP8 genes, *BMP8A* and *BMP8B*, have been identified (**Zhao et**

al., 1996). BMP8B has been found to play roles in the regulation of thermogenesis in mature brown adipose tissue

and spermatogenesis in male germ cells, while BMP8A shown to closely correlate with the progression of

spermatogenesis (**Walton et al., 2016; Ying et al., 2001**). In contrast, only one *bmp8* gene, *bmp8a*, has been

isolated in zebrafish, and shown to be involved in the regulation of lipid metabolism (**Zhong et al., 2018**). To date,

whether Bmp8a plays a role in immune responses is completely unknown. The present study was thus performed

to answer this question. We demonstrated that Bmp8a was a previously unrecognized regulator functioning in

antiviral immune responses of zebrafish *Danio rerio*, providing a new target for control of viral infection.

Results

Bmp8a inhibited RNA virus replication *in vitro*.

To explore the function of zebrafish Bmp8a in the antiviral immune response, we infected both the wild-type

zebrafish liver (ZFL) cells and ZFL cells over-expressing bmp8a as well as the epithelioma papulosum cyprini

(EPC) cells and EPC cells over-expressing bmp8a with Grass carp reovirus (GCRV), a dsRNA virus, and then

monitored the cytopathic effect（CPE）and the viral titers of the supernatants. In response to GCRV challenge,

an apparent CPE was observed in the control cells, while the bmp8a-overexpressing ZFL cells had much less CPE

(**Figure 1A**). In accordance with the CPE results, a significant decrease of the viral titers was observed in the

supernatants of bmp8a-overexpressing ZFL cells (**Figure 1B**). Similarly, the GCRV-induced CPE and GCRV yields

of the supernatants were also markedly reduced in the bmp8a-overexpressing EPC cells (**Figure 1C-D**). These

suggested that Bmp8a suppressed the replication of the RNA virus in the cells.

Bmp8a was involved in antiviral responses *in vivo*.

To test the function of Bmp8a in host defense against virus infection *in vivo*, we generated *bmp8a* deficient

(*bmp8a$^{-/-}$*) zebrafish using the TALEN approach (**Figure 1 – Figure supplement 1A**), which caused 7 nucleotides

deletion in the exon 4 (**Figure 1 – Figure supplement 1B-C**). We challenged both the wild-type and *bmp8a$^{-/-}$*

mutant zebrafish by intraperitoneal injection with GCRV. Compared with wild-type zebrafish, the levels of GCRV

RNA in the liver, kidney, intestine, and spleen of *bmp8a$^{-/-}$* zebrafish were considerably increased (**Figure 1E**).

Accordingly, *bmp8a$^{-/-}$* zebrafish exhibited significantly reduced survival rate than wild-type zebrafish upon GCRV

infection (**Figure 1F**). When the wild-type and *bmp8a$^{-/-}$* zebrafish were similarly challenged with spring viremia of

carp virus (SVCV), a negative ssRNA virus, the survival rate of *bmp8a*^{−/−} zebrafish was also remarkably lower than

that of wild-type zebrafish (***Figure 1G***). Moreover, when the wild-type and *bmp8a*^{−/−} zebrafish were infected with

turbot skin verruca disease virus (TSVDV) isolated from skin (***Figure 1 – Figure supplement 2***), mortality occurred

in *bmp8a*^{−/−} zebrafish at 108 h postinfection, and all the fish died at 156 h post infection, while none of the wild type

zebrafish were dead during the period (***Figure 1H***). These data indicated that *bmp8a*^{−/−} zebrafish were more

susceptible to virus infection than wild-type zebrafish, suggesting involvement of Bmp8a in antiviral immune

responses *in vivo*.

Bmp8a promoted expression of antiviral genes.

In zebrafish, type I IFNs contained 4 members *ifnφ1*, *ifnφ2*, *ifnφ3*, and *ifnφ4*, among which *ifnφ1* and *ifnφ3* were

shown to be involved in the antiviral response. In contrast, in carp, only one type I IFN was identified to fulfill

antiviral role in EPC cell (***Robertsen, 2006; Zhang et al., 2007***). To pinpoint the role of Bmp8a in the antiviral

immune response, both the wild-type and *bmp8a*^{−/−} zebrafish were injected intraperitoneally with GCRV, and the

expression of type I IFN genes and the antiviral protein gene *mx* were detected using quantitative reverse

transcription PCR (qRT-PCR). We found that the expression of both *ifnφ1* and *ifnφ3* as well as *mx* was all

significantly down-regulated in the liver, kidney, intestine, and spleen of *bmp8a*^{−/−} zebrafish, compared with those

of wild-type fish (***Figure 2A-D***). We then over-expressed bmp8a in ZFL cells and found that ZFL cells with over-

expressed Bmp8a showed markedly higher expression of *ifnφ1*, *ifnφ3*, and *mx* than that of control cells, infected

with or without GCRV (***Figure 2E-H***). In addition, *bmp8a* knockdown caused remarkably decreased expression of

ifnφ1, *ifnφ3*, and *mx* in ZFL cells (***Figure 2I-L***). Similarly, the expression of *ifn* and *mx* in EPC cells with over-

expressed Bmp8a was also remarkably up-regulated than that in control cells (***Figure 2M-P***). Moreover, EPC cells

cotransfected with bmp8a expressing plasmid and IFNφ1, IFNφ3, or EPC IFN promoter-driven luciferase plasmid,

followed by infection with or without GCRV, had markedly increased intracellular IFNφ1, IFNφ3, or EPC IFN

promoter-driven luciferase activities (***Figure 2Q-S***). All these data denoted that Bmp8a promoted the expression

of both type I IFN and *mx* genes.

Bmp8a activated Tbk1-Irf3-Ifn antiviral signaling via p38 MAPK pathway.

To putatively examine the molecular mechanism by which Bmp8a stimulates *ifn* and *mx* expression, we carried

out transcriptome analysis of the livers of *bmp8a*^{−/−} mutant and wild-type zebrafish, that both had been

intraperitoneally injected with GCRV. Kyoto Encyclopedia of Genes and Genomes (KEGG) analyses revealed that,

compared with wild-type zebrafish, the down-regulated genes in the mutant were remarkably enriched in the RIG-

I-like receptor signaling pathway, NF-kappa B signaling pathway, TNF signaling pathway, and protein digestion

and absorption that function in the antiviral immune process (***Figure 3 – Figure supplement 1***). In agreement

with transcriptome analysis, we found that the expressions of *tbk1*, *irf3*, and *irf7* were significantly upregulated in

bmp8a-overexpressing ZFL cells than control cells, infected with or without GCRV (***Figure 3A-D***). Similar

expression patterns of *tbk1*, *irf3*, and *irf7* were also observed in EPC cells, infected with or without GCRV (***Figure***

3E-H). In addition, *bmp8a* knockdown resulted in considerably reduced expression of *tbk1*, *irf3*, and *irf7* in ZFL

cells than control cells, infected with or without GCRV (***Figure 3I-L***). We then evaluated the activation of Tbk1 and

Irf3 by immunoblot assays in both bmp8a-overexpressing and wild-type ZFL cells as well as EPC cells, infected

with or without GCRV. It was found that phosphorylation levels of Tbk1 and Irf3 were significantly increased in both

types of the cells with overexpressed Bmp8a (***Figure 3M-P***). Moreover, *bmp8a* knockdown markedly reduced the

phosphorylation levels of Tbk1 and Irf3 in ZFL cells, infected with or without GCRV (***Figure 3Q-R***). Furthermore,

in luciferase activity assays, transfection of dominant negative mutations of *tbk1* (tbk1-K38M), *irf3* (irf3DN), or *irf7*

(irf7DN) in EPC cells induced a significant loss in the ability of Bmp8a to activate the IFN promoter (***Figure 3S-U***).

This was further supported by the observations *in vivo* that the expression of *tbk1*, *irf3*, and *irf7* in the liver, kidney,

intestine, and spleen was significantly downregulated in *bmp8a$^{-/-}$* mutant than that in wild-type zebrafish (***Figure***

3V-Y). These data together revealed that Bmp8a induced *ifn* and *mx* expression via promoting Tbk1-Irf3-Ifn

antiviral signaling.

BMPs transmit signals through smad1/5/8, smad2/3, ERK, JNK, or p38 MAPK pathways (***Derynck et al.,***

2001; Drummond, 2005; Oh et al.,2002). Thus, the effects of p38 MAPK inhibitor SB203580, JNK inhibitor

SP600125, MEK1/2 inhibitor U0126, SMAD1/5/8 inhibitor DMH1, and smad2/3 inhibitor TP0427736 HCl on the

expression of *ifn* was tested in bmp8a-overexpressing ZFL cells and EPC cells. We have shown above that Bmp8a

overexpression increased the expression of *ifn* (***Figure 2***). Here we found that only p38 MAPK inhibitor SB203580

significantly reduced the expression of *ifnφ1* and *ifnφ3* in ZFL cells and the expression of *ifn* in EPC cells (***Figure***

4A-C). Furthermore, the IFN promoter-driven luciferase assays revealed that the luciferase activities were

markedly reduced in EPC cells upon treatment with p38 MAPK inhibitor SB203580 (***Figure 4D-F***), consistent with

the observation that Bmp8a induced the expression of *ifn* through p38 MAPK pathway. Taken the previous report

that the MAPK signaling increased the phosphorylation of TBK1 and IRF3 upon viral infection into consideration

(***Shi et al., 2019***), we suggested that Bmp8a activated Tbk1-Irf3-Ifn antiviral signaling via p38 MAPK pathway.

BMP type I receptor Alk6a interacted with Bmp8a and participated in antiviral immunity.

BMPs exert their biological effects through the sequential activation of two types of transmembrane receptors,

namely BMP receptor type I and BMP receptor type II, that both possess intrinsic serine/threonine kinase activity

(**Heldin et al., 1997; Shi et al., 2003**). To detect which receptor is involved in the regulation of the antiviral immune

responses, the expression of both type I receptor genes (*alk2*, *alk3*, *alk6a*) and type II receptor genes (*bmpr2a*,

bmpr2b, *actr2a*, *actr2b*) in ZFL cells transfected with poly(I:C) or infected with GCRV were measured. Among these

receptor genes, *alk6a* showed the highest expression upon poly(I:C) or virus challenge (**Figure 5A-B**). We also

overexpressed all these receptors in ZFL and EPC cells, and examined the expression of *ifn* in these cells. The

upregulation of *ifnφ1* was only found in the alk6a-overexpressing ZFL cells (**Figure 5C**). Moreover, the

overexpression of *alk6a* or *alk2* resulted in significantly increased expression of *ifnφ3* in ZFL cells and *ifn* in EPC

cells, with overexpressed *alk6a* being more effective (**Figure 5D-E**). Based on these findings, the function of Alk6a

was selected for further test. Compared to wild-type ZFL cells, the expressions of antiviral protein genes *ifnφ1,*

ifnφ3 and *mx* as well as the antiviral signaling genes *tbk1*, *irf3*, and *irf7* were all markedly upregulated in alk6a-

overexpressed ZFL cells with or without GCRV infection (**Figure 5F-G**). The similar results were also observed in

the alk6a-overexpressed EPC cells (**Figure 5H-I**). Because the phosphorylation of Gly-Ser (GS) domain of the

type I receptor is required for its activation, we thus constructed dominant-negative mutant of alk6a (alk6a-ΔGS)

plasmid (**Figure 5J**), and then transferred it into both ZFL and EPC cells. It revealed that the dominant-negative

mutation of *alk6a* significantly reduced the expression of *mx, ifn* (*ifnφ1* and *ifnφ3* in ZFL cells), *tbk1*, *irf3*, and *irf7*

in ZFL and EPC cells, with or without GCRV infection (**Figure 5K-N**). The data suggested that Alk6a was apparently

involved in the Tbk1-Irf3/7-Ifn antiviral signaling.

Considering we have revealed that Bmp8a activated Tbk1-Irf3-Ifn antiviral signaling via p38 MAPK pathway,

we then examined whether the the levels of phosphorylated p38 MAPK protein were regulated by Alk6a. Clearly,

compared to wild-type ZFL or EPC cells, phosphorylation levels of p38 MAPK were significantly increased in alk6a-

overexpressed both types of the cells with or without GCRV infection (**Figure 6A-D**). Moreover, dominant-negative

mutation of *alk6a* significantly reduced the phosphorylation levels of p38 MAPK in ZFL or EPC cells, with or without

GCRV infection (**Figure 6E-H**). Thus, Alk6a was shown to be involved in the phosphorylation of p38 MAPK. Next,

we investigated whether the antiviral immune responses of Bmp8a were mediated by Alk6a. In the IFN promoter-

driven luciferase assays, it was found that the ability of Bmp8a to activate the IFN promoter was markedly blocked

after the dominant-negative mutation of *alk6a* in EPC cells (**Figure 6I-K**). Importantly, co-immunoprecipitation (Co-

IP) experiments confirmed that Bmp8a interacted with Alk6a (**Figure 6L**). Collectively, our data demonstrated that

BMP type I receptor Alk6a interacted with Bmp8a, and participated in the antiviral immunity. To our best knowledge,

Alk6a is the first BMP receptor identified thus far which is being directly required for antiviral immune responses.

Virus induced expression of *bmp8a*.

Next, we measured the expression of *bmp8a* in ZFL cells upon virus infection. GCRV significantly induced the

expression of *bmp8a* at 12 h post-stimulation, and reached a peak at 48 h (***Figure 7A***). We also detected the

expression of *bmp8a* upon infection with GCRV or poly(I:C) *in vivo*. The expression of *bmp8a* in the liver, kidney,

intestine, and spleen of zebrafish infected with GCRV or poly(I:C) was all significantly elevated (***Figure 7B-I***). This

indicated that the expression of *bmp8a* was inducible by infection with virus or its mimic poly(I:C).

Binding of Stat1 to GAS motifs activated *bmp8a* expression.

To explore how virus induces expression of *bmp8a*, we searched for the transcription factor binding sites in *bmp8a*

promoter region at the web http://jaspar.genereg.net/. Two gamma-activated sites (GAS), 5'-ATTCCGGGAAA-3'

(P1) and 5'-TTTACTAGAAC-3' (P2), in *bmp8a* promoter region were identified (***Figure 8 – Figure supplement 1***).

In mammals, the transcription factor STAT1 homodimer was known to bind to the GAS motif in the promoter region

of IFN-γ triggered downstream genes (***Decker et al., 1997; Shuai et al., 1992***). We thus wonder if Stat1 can bind

to the *bmp8a* promoter region to activate the expression of *bmp8a*. In zebrafish, there exist two Stat1 factors,

Stat1a (GenBank accession number NM_131480.1) and Stat1b (GenBank accession number NM_200091.2).

Thus we constructed dominant-negative mutant plasmids of stat1a-ΔC and stat1b-ΔC, that were transferred into

ZFL cells, respectively. It was found that the expression of *bmp8a* was significantly blocked in both cases (***Figure***

8A-B). By contrast, the overexpression of either *stat1a* or *stat1b* markedly promoted *bmp8a* promoter-driven

luciferase activities (***Figure 8C***). In addition, when one of the GAS motifs in *bmp8a* promoter region was deleted,

bmp8a promoter-driven luciferase activities were significantly reduced; when both of the GAS motifs were deleted,

bmp8a promoter-driven luciferase activities were reduced even more significantly (***Figure 8D-E***). These suggested

that both Stat1a and Stat1b may interact with the GAS motifs.

To verity the binding of Stat1a/Stat1b to the GAS motifs in *bmp8a* promoter region, the recombinant proteins

of both Stat1a and Stat1b were expressed and purified (***Figure 8F-G***), and used for electrophoretic mobility shift

assay (EMSA). As shown in ***Figure 8H-K***, a remarkable band of protein-DNA complex was observed (lane 2). The

protein-DNA complex band was obviously weaker in the presence of an excess unlabeled competitor probe,

compared to that of biotin labeled probe group (lane 3), but the intensity of the band showed little change in the

presence of GAS mutant probes (lane 4). To detect the specific binding, a supershift experiment with the antibody

against His tag was performed. It revealed that His tag antibody caused a specific supershift of slower migrating

protein-DNA-antibody complex (***Figure 8H-K***, lane 5). There was no protein-DNA complex observed in the

negative control group (***Figure 8H-K***, lane 1). These demonstrated that both Stat1a and Stat1b could directly bind

to the GAS motifs in *bmp8a* promoter region, suggesting that *bmp8a* expression was subjected to the regulation

by binding of Stat1a/Stat1b to the GAS motifs.

Discussion

BMPs, canonical multi-functional growth factors, are suggested to play roles in regulation of immune

responses, but it remains controversial over their functions in immunity. We show here for the first time that Bmp8a

is a previously unrecognized factor involved in the regulation of antiviral immune responses in zebrafish. Evidences

supporting this nature of Bmp8a include: Bmp8a inhibits RNA viruses replication *in vitro* and *in vivo*; Bmp8a promotes

the expression of antiviral protein genes including type I IFNs and *mx*; the expression of *bmp8a* is induced by CGRV

or poly(I:C), and Bmp8a activates TBK1-IRF3-IFN signaling. Bmp8a is a member of the TGF-β superfamily, which

includes more than 30 genes encoding TGFβ, BMPs and activins. TGFβ has been shown to have both pro- and

anti-inflammatory effects depending on the context it is acting in the immune system (***Worthington et al., 2012***).

Like TGFβ, BMP members have also been shown to play different immunoregulatory roles (***Tu et al., 2014***). BMP2,

4 and 7 deficiency or knockdown results in partial loss of the thymic capsule, reduced size of thymus and increased

Helicobacter pylori-induced inflammation, whereas BMP6 inhibits macrophage growth by inducing cell cycle arrest

in vitro (***Aleman-Muench and Soldevila, 2012; Takabayashi et al., 2014; Bleul et al., 2005; Gordon et al., 2010;***

Lee et al., 2007). The finding that Bmp8a is a positive regulator in antiviral immune responses provides a new

angle for our understanding of the functions of TGF-β superfamily members including BMPs.

On one hand, it is generally regarded that the recognition of virus by PRRs elicits the activation of IRF3 and

NF-κB, and induces the production of IFNα/β, that then activate JAK-STAT pathway, eventually leading to the

production of ISGs such as cytokines and antiviral proteins. On the other hand, BMPs are known to transmit signals

through both Smad-dependent pathways (smad1/5/8 and smad2/3 pathways) and Smad-independent pathways

(ERK, JNK, and p38 MAPK pathways). Here we clearly demonstrate that Bmp8a promotes the expression of

endogenous type I IFNs independent of Smad signaling pathways. Moreover, it is p38 MAPK pathway, rather than

ERK and JNK pathways, that is involved in the antiviral responses induced by Bmp8a. Furthermore, we show that

Alk6a participates in Tbk1-Irf3/7-Ifn antiviral signaling via interaction with Bmp8a. In addition, Alk6a was found to

be required to induce the phosphorylation of p38 MAPK, which is consistent with the findings in mouse that Alk6

knockdown suppresses p38 MAPK phosphorylation (***Itoh et al., 2008***). Previous studies have shown that MAPK

signaling pathway is crucial in the phosphorylation of TBK1 and IRF3 upon viral infection (***Shi et al., 2019***). Thus,

we probably discover a new pathway of Bmp8a signaling in antiviral immune responses that Bmp8a acts as a

positive regulator through the promotion of phosphorylation of Tbk1 via p38 MAPK pathway, i.e., Bmp8a binds to

Alk6a, which induces p38 MAPK phosphorylation, and in turn enhances Tbk1 and Irf3 phosphorylation, eventually

resulting in increased synthesis of type I IFN (**Figure 9**).

Our study also show that the challenge with GCRV and poly(I:C) both significantly promotes the expression

of zebrafish *bmp8a*. Searching for the binding sites of transcript factors in *bmp8a* promoter region reveals that two

IFN-γ activation sites (GAS) that may interact with STAT1 are identified. Functionally, we demonstrate that Stat1a

and Stat1b, orthologues of human STAT1, can directly bind the GAS sites in *bmp8a* promoter, resulting in activation

of *bmp8a* expression. It has been reported that in black carp, Stat1a and Stat1b can both form homodimer and

heterodimer *in vivo* like their mammalian counterpart (**Wu et al., 2019**). It is thus highly likely that in zebrafish,

Stat1a and Stat1b can form homodimer or heterodimer, which then binds to the GAS sites in *bmp8a* promoter

region. In zebrafish, IFN-γ has been found to be able to induce the expression of both *stat1a* and *stat1b*, and the

GAS motif is required for Ifn-γ activation (**Zou and Secombes, 2011; Ruan et al., 2017**). It is widely known that

the JAK-STAT pathway is one of the most important candidate pathways through which IFN-γ works. Activation of

JAK-STAT signaling leads to transcription of various downstream ISGs for antiviral activity (**Boehm et al., 1997**).

We thus think that Bmp8a is another important antiviral factor which can be activated through Ifn-γ-Jak-Stat1

pathway (**Figure 9**).

In summary, we demonstrate for the first time that Bmp8a is a positive regulator in antiviral immune responses,

which functions through promotion of phosphorylation of Tbk1 via p38 MAPK pathway. Upon virus infection, the

expression of *bmp8a* is activated by Stat1a/Stat1b directly. Bmp8a knockdown leads to significantly decreased

phosphorylation of Tbk1 and Irf3. As a consequence, *bmp8a*-deficient zebrafish produces significantly lowered

type I IFNs in response to RNA virus infection, exhibits significantly reduced immune responses, and increases

significantly higher viral load and morbidity *in vivo*. Our study is significant to understand the regulatory mechanism

of BMP involving in antiviral immune response and provides a new target for controlling viral infection.

300 **Materials and Methods**

301 **Key Resources Table**

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
antibody	anti-TBK1 (rabbit, monoclonal)	Cell Signaling Technology	3504	WB(1:1000)
antibody	anti-TBK1 pSer172 (rabbit, monoclonal)	Cell Signaling Technology	5483	WB(1:1000)
antibody	anti-IRF3 (rabbit, polyclonal)	Bioss	2993R	WB(1:1000)
antibody	anti-IRF3 pSer386 (rabbit, monoclonal)	Bioss	52170R	WB(1:1000)
antibody	anti-p38MAPK (rabbit, polyclonal)	Bioss	0637R	WB(1:1000)
antibody	anti-p38 MAPK pThr180/pTyr182 (rabbit, polyclonal)	Bioss	2210R	WB(1:1000)
antibody	anti-β-Actin (rabbit, polyclonal)	Bioss	0061R	WB(1:2000)
antibody	anti-HA (mouse, monoclonal)	Beyotime	AH158	WB(1:1000)
antibody	anti-Falg (mouse, monoclonal)	Beyotime	AF519	WB(1:1000)
antibody	anti-His (mouse, monoclonal)	CWBIO	CW0285	WB(1:5000)
antibody	goat anti-rabbit IgG HRP	CWBIO	CW0103S	WB(1:8000)
antibody	VeriBlot for IP Detection	Abcam	ab131366	WB(1:5000)
other	Protein A/G PLUS-Agarose	Santa Cruz	sc-2003	
kit	Chemiluminescent EMSA Kit	Beyotime	GS009	
kit	PrimeScript™ RT reagent Kit with gDNA Eraser	TaKaRa	RP047A	
kit	miRNeasy Mini Kit	QIAGEN	217004	
chemical compound	DMH1	Selleck	S7146	10 µM
chemical compound	TP0427736 HCl	Selleck	S8700	10 µM
chemical compound	SB203580	Beyotime	S1863	10 µM
chemical compound	SP600125	Beyotime	S1876	10 µM
chemical compound	U0126	Beyotime	S1901	10 µM
kit	Lipofectamine RNAiMAX	Invitrogen	13778075	
kit	Luc-Pair Duo-Luciferase Assay Kits	iGene Biotechnology	LF002	
kit	FuGENE HD Transfection Reagent	Promega	E2311	
kit	Mut Express II Fast Mutagenesis Kit	Vazyme	C214-01	
kit	mMESSAGE mMACHINE SP6 kit	Invitrogen	AM1340	

302 **Cells and viruses.** The epithelioma papulosum cyprini cells (EPC), zebrafish liver cells (ZFL) were acquired from

303 the China Zebrafish Resource Center (CZRC) and cultured according to the instructions from the CZRC. EPC and

304 ZFL cells were grown in a 28 °C incubator supplied with 5% CO_2. The flounder gill cells (FG) were grown in a 22 °C

305 incubator supplied. The EPC or FG cells were cultured in MEM media supplemented with 10% fetal bovine serum

306 (FBS), 100 U/ml penicillin and 100 µg/ml streptomycin. The ZFL cells were maintained in DMEM/F-12 media

307 supplemented with 10% FBS, 100 U/ml penicillin and 100 µg/ml streptomycin. Grass carp reovirus (GCRV), a

308 dsRNA virus, and spring viremia of carp virus (SVCV), a negative ssRNA virus, were kindly provided by Yibing

309 Zhang (Institute of Hydrobiology, Chinese Academy of Sciences), and tittered according to the method of Reed

310 and Muench (**Reed and Muench, 1938**), by a 50% tissue culture infective dose ($TCID_{50}$) assay on EPC cells. The

311 turbot skin verruca disease virus (TSVDV) was isolated from the focal site of turbot skin in our lab.

312 **Zebrafish**. The zebrafish *bmp8a* knockout mutant lines ($bmp8a^{-/-}$) were established from the zebrafish AB line

313 using TALENs technology. The TALENs for zebrafish *bmp8a* were assembled using the Unit Assembly (UA)

314 method as previously described (**Huang et al., 2014**). All the UA method starting vectors were the gifts from

315 laboratories of Bo Zhang at Peking University and of Shuo Lin at University of California Los Angeles. TALENs

316 mRNAs were transcribed using the mMESSAGE mMACHINE SP6 kit and purifified using the RNeasy Mini kit. To

317 generate zebrafish mutant lines, TALEN mRNAs (50–200 pg) were microinjected into 1-cell stage zebrafish

318 embryos. Two days after injection, genomic DNA was isolated from 8 to 10 pooled larvae. The target genomic

319 regions were amplified by nested PCR and subcloned into the pGEM-T vector. Single colonies were genotyped by

320 sequencing. To obtain germline mutations, the TALEN-injected embryos were raised to adulthood and outcrossed

321 with wild-type (WT) fish. The F1 progeny were genotyped by sequencing. To obtain homozygous mutants,

322 heterozygous mutant of the same mutation were obtained and self-crossed. The primers used in this study were

323 listed in Supplemental Table 1.

324 **Plasmid construction.** The open reading frames (ORF) of zebrafish bmp8a, stat1a, stat1b, alk2, alk3, alk6a,

325 actr2a, actr2b, bmpr2a and bmpr2b were cloned into the pcDNA3.1 expression vector for the eukaryotic expression.

326 The ORF of zebrafish stat1a and stat1b were cloned into the pET-28a expression vector for the expression of

327 recombinant proteins. The promoter regions of bmp8a, bmp8a-ΔP1 (delete the GAS motif of 5´-attccgggaaa-3´ in

328 the bmp8a promoter), bmp8a-ΔP2 (delete another GAS motif of 5´-tttactagaac-3´ in the bmp8a promoter), bmp8a-

329 ΔP1and ΔP2 (delete both the two GAS motifs), IFNφ1, IFNφ3, and EPC IFN were cloned into the pGL3-basic

330 vector. Dominant negative mutant plasmids including tbk1-K38M, stat1a-ΔC, stat1b-ΔC, irf3DN, and irf7DN were

331 described previously (**Feng et al., 2015; Sun et al., 2011**). Dominant negative mutant plasmids alk6a-ΔGS (delete

332 the sequence of 5´-agctccggttctggctcagga-3´ encoding the GS domain) was cloned into the pcDNA3.1 expression

333 vector. The deletions and mutations were created using Mut Express II Fast Mutagenesis Kit. The primers used in

334 this study were listed in Supplemental Table 2.

335 **Transcriptome Sequencing.** The zebrafish were fasted for 24 h prior to tissue collection. Three mutant (*bmp8a*⁻

$^{/-}$) and three wild type (WT) adult zebrafish (0.4 g ± 0.05 g) were randomly selected from each group and were

injected intraperitoneally with 50 μl of GCRV (10^8TCID$_{50}$ per ml) per fish. Seventy-two hours after the injection, the

livers from three fish each group were pooled together for total RNA extraction. Three independent samples from

each group were prepared for transcriptome sequencing. mRNAs were enriched by magnetic beads combining

the oligo (dT). cDNA fragments were subjected to the end repair process by the addition of a single 'A' base and

the ligation of Illumina sequencing adapters. The ligation products were selected by agarose gel electrophoresis,

amplified by PCR, and sequenced using Illumina HiSeq 4000 by Annoroad Technology (Beijing, China).

Identification of DEGs. The RPKM (reads per kb per million reads) method was used to calculate and normalize

the abundances of genes (*Mortazavi et al., 2008*). DEGs across groups were identified by the edgeR package

(http://www.r-project.org/). Genes with a fold change ≥ 2 and a false discovery rate (FDR) < 0.05 in a comparison

were regarded as significantly expressed. DEGs were then subjected to enrichment analysis of KEGG pathways.

Viral infection *in vitro*. EPC, FG, or ZFL cells were plated 24 h before infection. Cells were infected with GCRV

(5×10^4TCID$_{50}$ per ml) or TSVDV (crude virus extracts filtered by a 0.45 μm microporous membrane) for 1 h in

medium without FBS. After infection, cells were washed with PBS and then medium was added with FBS. The

cells and cell-free supernatants were harvested at the indicated times.

Viral infection *in vivo*. For *in vivo* viral infection, the adult WT and *bmp8a$^{-/-}$* zebrafish were infected via

intraperitoneal injection using 50 μL of SVCV (10^8TCID$_{50}$ per ml), GCRV (10^8TCID$_{50}$ per ml), or TSVDV (crude

virus extracts filtered by a 0.45 μm microporous membrane) per fish. Total RNA was extracted and viral RNA

expression in the liver, kidney, intestine, and spleen were examined by qRT-PCR. For the survival experiments,

zebrafish were monitored for survival after SVCV, GCRV, or TSVDV infection.

Crystal violet staining. EPC or ZFL cells were transfected with bmp8a or empty vector plasmid. At 24 h post

transfection, cells were treated with GCRV (5×10^4TCID$_{50}$ per ml). At 72 h after virus challenge, cells were fixed

with 4% formaldehyde for 1 h and stained with 0.5% crystal violet for 2 h. After washing with tap water, visible

plaques were photographed.

Luciferase assays. EPC cells were seeded in 24-well plates overnight and cotransfected with various plasmids

using FuGENE HD Transfection Reagent following the manufacturer's instruction. If necessary, the cells were

infected with GCRV (5×10^4TCID$_{50}$ per ml). At the indicated time points, luciferase activities were measured with

the Luc-Pair Duo-Luciferase Assay Kits. Data were normalized by calculating the ratio between firefly luciferase

activity and *Renilla* luciferase activity. The primers used for cloning the promoter are shown in

Supplemental Table 2.

RNA interference. Small interfering RNA (siRNA) of bmp8a and si-NC (negative control) were purchased from

RiboBio. The oligonucleotide sequences (siRNA) specific targeting *bmp8a* mRNA was as follows: 5′-

GTTCTTCAGAGCTAGTCAA-3′. Transfection was performed with Lipofectamine RNAiMAX according to the

manufacturer's protocols. Total RNA was extracted for qRT-PCR analysis.

Drug treatment and analysis. SMAD1/5/8 inhibitor DMH1, smad2/3 inhibitor TP0427736 HCl, p38 MAPK inhibitor

SB203580, JNK inhibitor SP600125, and MEK1/2 inhibitor U0126 were dissolved in dimethyl ulfoxide (DMSO).

ZFL or EPC cells were treated with each inhibitor, which were diluted with culture medium at concentrations of 10

µM for 24 h. Total cellular RNA was extracted for qRT-PCR analysis.

RNA quantification. Total RNA was isolated using the miRNeasy Mini Kit according to the manufacturer's

instructions. RNA treated with DNase and the cDNAs were synthesized with PrimeScript™ RT reagent Kit with

gDNA Eraser. Samples without reverse transcriptase were also added as control. Gene expression was

determined by amplifying the cDNA with ChamQ SYBR Color qPCR Master Mix by an ABI 7500 Fast Real-Time

PCR System. The expression of zebrafish *actb1* or EPC *actin* was used as an internal control. The $2^{-\Delta\Delta Ct}$ method

was used to calculate relative expression changes. All qRT-PCR experiments were performed in triplicate and

repeated three times. Primers used are listed in Supplementary Table 1.

Preparation of recombinant protein. The recombinant protein of rStat1a or rStat1b was expressed and purified

as previous description (***Sun et al., 2013***). Briefly, the recombinant plasmid (pET-28a-stat1a or pET-28a-stat1b)

was transformed into *Trans*B (DE3) chemically competent cell. The positive transformants were incubated in liquid

LB medium containing 50 µg/mL kanamycin at 37 °C by shaking at 160 rpm. When the cells grow to OD600 = 0.5,

isopropyl β-D-thiogalactoside (IPTG) was added to a final concentration of 1 mM, and incubated at 19 °C with

shaking at 120 rpm for 14 h. After incubation, the cells were harvested by centrifugation, re-suspended in PBS,

and broken by ultrasound. The supernatant was centrifuged at 12,000 g for 20 min, and the protein was purified

by Ni-NTA Sepharose column. The concentration of purified protein was determined by BCA (bicinchonininc acid)

method. The purified recombinant proteins were stored at -80 °C for subsequent experiment.

Electrophoretic mobility shift assay. Electrophoretic mobility shift assay (EMSA) was performed according to

the instruction of Chemiluminescent EMSA Kit. Oligonucleotides (Supplementary Table 1) were synthesized and

biotinylated by Sangon Biotech Company as required and the complementary oligonucleotides were annealed to

form the double DNA probe. The recombinant protein was pre-incubated in EMSA/Gel-Shift binding buffer with

poly (dI-dC) at 25 °C for 25 min in the presence or absence of Non-Biotin probe (0.5 µM). Then, all samples were

mixed with biotinylated probes (0.5 µM) and incubated at 25°C for 30 min. The binding reaction without

recombinant protein was set as the negative control. For supershift analysis, anti-His tag antibody was incubated

with the reaction mixture for another 30 min after the labeled probe was added. After separation in a native 6%

polyacrylamide gel, the free probes and DNA-protein compound were transferred to a Hybond-N$^+$ nylon membrane,

followed by crosslinking under UV light. After immersion in the blocking buffer, membranes were incubated with

streptavidin conjugated HRP for 20 min with shaking and then washed three times with washing buffer. Membranes

were exposed briefly in Western Lightning-ECL and then exposure to Automatic X-ray Film Processor.

Immunoblot analysis and co-immunoprecipitation. For Immunoblot analysis, cells were lysed in NP-40 buffer

(150 mM NaCl, 0.5% EDTA, 50 mM Tris, 1% NP40, proteinase inhibitors, and protein phosphatase inhibitors) for

30 min on ice. For co-immunoprecipitation (co-IP), whole-cell extracts were collected 48 h after transfection and

lysed in NP-40 buffer on ice for 30 min. After centrifugation for 15 min at 12,000 × g, 4 °C, supernatants were

collected and incubated with Protein A/G PLUS-Agarose coupled to specific antibodies for overnight with rotation

at 4°C. Beads were washed 3 × times with NP-40 buffer. Bound proteins were eluted by boiling 7min with 2 x SDS

sample buffer. For immunoblot analysis, immunoprecipitates or whole-cell lysates were separated by 12.5% SDS–

PAGE gels, electro-transferred to PVDF membranes and blocked for 4 h with 5% no-fat milk solution, followed by

blotting with the appropriate antibodies and detection by enhanced chemiluminiscence (ECL). Antibodies from Cell

Signaling Technology: anti-TBK1 and anti-TBK1 pSer172. Antibodies from Bioss: anti-IRF3, anti-IRF3 pSer386,

anti-p38MAPK, anti-p38 MAPK pThr180/pTyr182, anti-β-Actin. Antibodies from Beyotime: anti-HA and anti-Falg.

Antibodies from CWBIO: anti-His and goat anti-rabbit IgG HRP secondary antibody. Antibodies from Abcam:

VeriBlot for IP Detection.

Statistical analysis. Representative experiments have been repeated at least two to three times. All statistical

analysis was performed using Graphpad Prism 8.0.1. Data are presented as mean ± SEM. Statistical significance

was assessed using the unpaired, two-tailed Student's t-test. Survival analyses were performed using the Kaplan-

Meier method and assessed using the log-rank (Mantel-Cox) test. *$p < 0.05$, **$p < 0.01$, *** $p < 0.001$; ns, non-

significant, $p > 0.05$.

Data availability. All data are available from the corresponding author upon reasonable request and that a Source

Data file is available for this article.

Conflict of Interest

The work is under a patent in China "A method to improve the antiviral immunity of fish (Application No.

201910454499.6)".

Author contributions

Z.L.and S.-C.Z. conceived and coordinated the project. Z.L., G.J., and S.Z. designed the experiments. S.Z.,

H.L. and Y.-S.W. performed experiments. H.-Y.L. and Y.W. contributed to establish the *bmp8a* knockout mutant

lines (*bmp8a$^{-/-}$*) using TALENs technology. S.Z. performed the statistical analysis. Z.L., S.-C.Z. and S.Z. wrote the

manuscript, with input from the other authors. All authors reviewed the manuscript and approved the final version.

Acknowledgements

This work was supported by the grants of National Natural Science Foundation of China (31572259) and

National Key Research and Development Project of the Ministry of Science and Technology (2018YFD0900505).

Ethics

Animal experimentation: The animals used in the experiment followed the ethical guidelines established by the

Institutional Animal Care and Use Committee of the Ocean University of China (permit number, SD2007695).

 Figure legends

Figure 1. Bmp8a inhibits RNA virus replication both *in vitro* and *in vivo*. (**A**, **B**) ZFL cells were transfected

with bmp8a or empty vector, respectively. The cells were infected with GCRV (5×10^4TCID$_{50}$ per ml) at 24 h post-

transfection, and the culture supernatants were collected at 72 h post-infection. The cell monolayers were fixed

with 4% paraformaldehyde for 1 h and stained with 0.5% crystal violet for 2 h (**A**), and the viral titers of the

supernatants were determined by TCID$_{50}$ assays (**B**). (**C**, **D**) Similar as (**A**, **B**) but in EPC cells. (**E**) The expression

of GCRV RNA in the liver, kidney, intestine, and spleen from wild-type (WT) or *bmp8a$^{-/-}$* zebrafish injected i.p. with

50 µl of GCRV (10^8TCID$_{50}$ per ml) for 72 h. (**B, D, E**) Data were from three independent experiments. Values are

expressed as mean ± SEM and analyzed by Student's *t*-test (***p < 0.001). (**F**, **G**) Survival curve of WT and *bmp8a$^-$*

$^{/-}$ zebrafish i.p. injected with 50 µl of GCRV (10^8TCID$_{50}$ per ml) or SVCV (10^8TCID$_{50}$ per ml). Values are expressed

as percent survival of 20 zebrafish for each genotype. *p<0.05 by log-rank test. (**H**) Survival curve of WT and

bmp8a$^{-/-}$ zebrafish i.p. injectedwith 50 µl of TSVDV (crude virus extracts filtered by a 0.45 µm microporous

membrane). Values are expressed as percent survival of 8 zebrafish for each genotype. **p<0.01 by log-rank test.

Source data are provided as a Source Data file.

Figure 2. Bmp8a promotes antiviral innate immune responses. (**A**–**D**) Expression of *ifnφ1*, *ifnφ3*, and *mx*

mRNA in the liver (**A**), kidney (**B**), intestine (**C**), and spleen (**D**) from WT or *bmp8a$^{-/-}$* zebrafish injected i.p. with 50

µl of GCRV (10^8TCID$_{50}$ per ml) for 72 h. (**E**, **F**) Expression of *ifnφ1*, *ifnφ3*, and *mx* mRNA after transfected with

bmp8a or empty vector in ZFL cells. The cells were collected at 36 h (**E**) or 48 h (**F**) post transfection. (**G**, **H**)

Expression of *ifnφ1*, *ifnφ3*, and *mx* mRNA after transfected with bmp8a or empty vector in ZFL cells for 24 h,

followed by infection with GCRV (5×10^4TCID$_{50}$ per ml) for another 24 h (**G**) or 36 h (**H**). (**I**, **L**) Expression of *ifnφ1*,

ifnφ3, and *mx* mRNA after *bmp8a* knockdown in ZFL cells. The cells were collected at 36 h (**I**) or 48 h (**L**) post

knockdown. (**K**, **L**) Expression of *ifnφ1*, *ifnφ3*, and *mx* mRNA after knockdown *bmp8a* in ZFL cells for 24 h, followed

by infection with GCRV (5×10^4TCID$_{50}$ per ml) for another 24h (**K**) or 36 h (**L**). (**M**, **N**) Expression of EPC *ifn* and

EPC *mx* mRNA after transfected with bmp8a or empty vector in EPC cells. The cells were collected at 36 h (**M**) or

48 h (**N**) post transfection. (**O**, **P**) Expression of EPC *ifn* and EPC *mx* mRNA after transfected with bmp8a or empty

vector in EPC cells for 24 h, followed by infection with GCRV (5×10^4TCID$_{50}$ per ml) for another 24 h (**O**) or 36 h

(**P**). (**Q**–**S**) EPC cells were transfected with pGL3-IFNφ1-promoter-luc (**Q**), pGL3-IFNφ3-promoter-luc (**R**) or pGL3-

EPC IFN-promoter-luc (**S**) respectively, with or without bmp8a, followed by infection with GCRV (5×10^4TCID$_{50}$ per

ml). *Renilla* luciferase was used as the internal control. Data were from three independent experiments. Values are

expressed as mean ± SEM and analyzed by Student's *t*-test (*p < 0.05, **p < 0.01, ***p < 0.001). Source data are

provided as a Source Data file.

Figure 3. Bmp8a increases Tbk1-Irf3-Ifn antiviral signaling. (**A**, **B**, **E**, **F**) Expression of *irf3*, *irf7*, and *tbk1* mRNA

after transfected with bmp8a or empty vector in ZFL (**A**, **B**) or EPC (**E**, **F**) cells. The cells were collected at 36 h

(**A**, **E**) or 48 h (**B**, **F**) post transfection. (**C**, **D**, **G**, **H**) Expression of *irf3*, *irf7*, and *tbk1* mRNA after transfected with

bmp8a or empty vector in ZFL (**C**, **D**) or EPC (**G**, **H**) cells for 24 h, followed by infection with GCRV (5×10^4TCID$_{50}$

per ml) for another 24 h (**C**, **G**) or 36 h (**D**, **H**). (**I, J, K, L**) Expression of *irf3*, *irf7*, and *tbk1* mRNA after *bmp8a*

knockdown in ZFL cells. The cells were collected at 36 h (**I**) and 48 h (**J**) post knockdown or at 24 h (**K**) and 36 h

(**L**) post infected with GCRV (5×10^4TCID$_{50}$ per ml). (**M**, **O**) Immunoblot analysis of phosphorylated (p-) Tbk1 and

Irf3 after transfected with bmp8a or empty vector in ZFL (**M**) or EPC (**O**) cells. The cells were collected at 36 h or

48 h post transfection for Immunoblot analysis. (**N**, **P**) Immunoblot analysis of phosphorylated (p-) Tbk1 and Irf3

after transfected with bmp8a or empty vector in ZFL (**N**) or EPC (**P**) cells for 24 h, followed by infection with GCRV

(5×10^4TCID$_{50}$ per ml) for another 24 h or 36 h. (**Q**, **R**) Immunoblot analysis of phosphorylated (p-) TBK1 and IRF3

after *bmp8a* knockdown in ZFL cells. The cells were collected at 36 h and 48 h post knockdown or at 24 h and 36

h post infected with GCRV (5×10^4TCID$_{50}$ per ml). (**S**–**U**) EPC cells were cotransfected with pGL3-IFNφ1-promoter-

luc (**S**), pGL3-IFNφ3-promoter-luc (**T**) or pGL3-EPC IFN-promoter-luc (**U**), and bmp8a together with each of the

dominant negative plasmids including tbk1-K38M, irf3DN and irf7DN. At 48 h post transfection, the cells were

collected for luciferase assays. *Renilla* luciferase was used as the internal control. (**V**–**Y**) Expression of *irf3*, *irf7*,

and *tbk1* mRNA in the liver (**V**), kidney (**W**), intestine (**X**), and spleen (**Y**) from WT or *bmp8a*$^{-/-}$ zebrafish injected

i.p. with 50 μl of GCRV (10^8TCID$_{50}$ per ml). Data were from three independent experiments. Values are expressed as

mean ± SEM and analyzed by Student's *t*-test (**p < 0.01, ***p < 0.001). Source data are provided as a Source

Data file.

Figure 4. Bmp8a promotes the IFN expression via p38 MAPK pathway. (**A**, **B**) Expression of *ifnφ1* (**A**) and

ifnφ3 (**B**) mRNA after transfected with bmp8a in ZFL cells for 24 h, followed by treatment with SB203580,

SP600125, U0126, DMH1 and TP0427736 HCl for another 24 h. **C** similar as (**A**, **B**) but in EPC cells. (**D**–**F**) EPC

cells were cotransfected with pGL3-IFNφ1-promoter-luc (**D**), pGL3-IFNφ1-promoter-luc (**E**) or pGL3-EPC IFN-

promoter-luc (**F**), pRL-TK together with bmp8a or empty vector, respectively. At 24 h post transfection, cells were

treated with or without SB203580 for another 24 h and then harvested for detection of luciferase activity. *Renilla*

luciferase was used as the internal control. Data were from three independent experiments. Values are expressed

as mean ± SEM and analyzed by Student's *t*-test (**p < 0.01, ***p < 0.001). Source data are provided as a Source

Data file.

Figure 5. Alk6a is involved in the antiviral innate immune responses. (A, B) Expression of *alk2*, *alk3*, *alk6a*,

bmpr2a, *bmpr2b*, *actr2a*, *actr2b* mRNA in ZFL cells stimulated with GCRV (5 × 10^4TCID$_{50}$ per ml, **A**) or poly(I:C)

(2 μg/ml, **B**) for 48 h. (**C–E**) Expression of *ifnφ1* (**C**) and *ifnφ3* (**D**) mRNA in ZFL cells or EPC *ifn* (**E**) in EPC cells

which were transfected with alk2, alk3, alk6a, bmpr2a, bmpr2b, actr2a, actr2b or empty vector for 48 h. (**F, H**)

Expression of *irf3*, *irf7*, *tbk1*, *ifn* (or *ifnφ1* and *ifnφ3*), and *mx* mRNA after transfected with alk6a or empty vector in

ZFL (**F**) or EPC (**H**) cells for 48 h. (**G, I**) Expression of *irf3*, *irf7*, *tbk1*, *ifn* (or *ifnφ1* and *ifnφ3*), and *mx* mRNA after

transfected with alk6a or empty vector in ZFL (**G**) or EPC (**I**) cells for 24 h, followed by infection with GCRV (5 ×

10^4TCID$_{50}$ per ml) for another 36 h. **J** Schematic drawing of the alk6a-ΔGS mutation that the GS domain of Alk6a

was deleted. (**K, M**) Expression of *irf3*, *irf7*, *tbk1*, *ifn* (or *ifnφ1* and *ifnφ3*), and *mx* mRNA after transfected with

alk6a-ΔGS or empty vector in ZFL (**K**) or EPC (**M**) cells for 48 h. (**L, N**) Expression of *irf3*, *irf7*, *tbk1*, *ifn* (or *ifnφ1*

and *ifnφ3*), and *mx* mRNA after transfected with alk6a-ΔGS or empty vector in ZFL (**L**) or EPC (**N**) cells for 24 h,

followed by infection with GCRV (5 × 10^4TCID$_{50}$ per ml) for another 36 h. Data were from three independent

experiments. Values are expressed as mean ± SEM and analyzed by Student's *t*-test (*p < 0.05, **p < 0.01, ***p<

0.001, ns non-significant). Source data are provided as a Source Data file.

Figure 6. Bmp8a activates the IFN expression through Alk6a. (A, C) Immunoblot analysis of phosphorylated

(p-) p38 MAPK after transfected with alk6a or empty vector in ZFL (**A**) or EPC (**C**) cells. The cells were collected

at 36 h or 48 h post transfection for Immunoblot analysis. (**B, D**) Immunoblot analysis of phosphorylated (p-) p38

MAPK after transfected with alk6a or empty vector in ZFL (**B**) or EPC (**D**) cells for 24 h, followed by infection with

GCRV (5 × 10^4TCID$_{50}$ per ml) for another 24 h or 36 h. (**E, G**) Immunoblot analysis of phosphorylated (p-) p38

MAPK after transfected with alk6a-ΔGS or empty vector in ZFL (**E**) or EPC (**G**) cells. The cells were collected at

36 h or 48 h post transfection for Immunoblot analysis. (**F, G**) Immunoblot analysis of phosphorylated (p-) p38

MAPK after transfected with alk6a or empty vector in ZFL (**F**) or EPC (**G**) cells for 24 h, followed by infection with

GCRV (5 × 10^4TCID$_{50}$ per ml) for another 24 h or 36 h. (**I–K**) EPC cells were cotransfected with pGL3-IFNφ1-

promoter-luc (**I**), pGL3-IFNφ3-promoter-luc (**J**) or pGL3-EPC IFN-promoter-luc (**K**), and bmp8a together with or

without the dominant negative plasmids alk6a-ΔGS. At 48 h post transfection, the cells were collected for luciferase

assays. *Renilla* luciferase was used as the internal control. **L** Co-immunoprecipitation and immunoblot analysis of

EPC cells cotransfected with alk6a-Flag and bmp8a-HA. Data were from three independent experiments. Values

are expressed as mean ± SEM and analyzed by Student's *t*-test (***p<0.001). Source data are provided as a

Source Data file.

Figure 7. GCRV or poly(I:C) increases the expression of *bmp8a*. (A) Expression of *bmp8a* mRNA in ZFL cells

after infected with GCRV (5×10^4TCID$_{50}$ per ml) for 48 h. (**B**–**E**) Expression of *bmp8a* mRNA in the liver (**B**), spleen (**C**), intestine (**D**), and kidney (**E**) from zebrafish injected i.p. with poly(I:C) (10 μg/fish). (**F**–**I**) Expression of *bmp8a* mRNA in the liver (**F**), spleen (**G**), intestine (**H**), and kidney (**I**) from zebrafish injected i.p. with 50μl of GCRV (10^8TCID$_{50}$ per ml). Zebrafish injected i.p. with PBS or MEM were used as the control. The expression of *actb1* served as a control for the qRT-PCR. Data were from three independent experiments. Values are expressed as mean ± SEM and analyzed by Student's *t*-test (*p < 0.05, **p < 0.01, ***p < 0.001, ns non-significant). Source data are provided as a Source Data file.

Figure 8. Stat1 binds the GAS sites and activates the *bmp8a* transcriptions upon virus stimulation. (**A, B**) Expression of *bmp8a* mRNA in ZFL cells after transfected with stat1a-ΔC or stat1b-ΔC for 36 h (**A**) or 48 h (**B**). (**C**) Dual luciferase report assay was used to analyze the transcription abilities of Stat1a and Stat1b in activation of *bmp8a* in EPC cells. pGl3-bmp8a-promoter-luc was transfected into EPC cells together with stat1a, stat1b or empty vector. After 48 h, the transfected cells were collected for luciferase assays. (**D**) Schematic drawing of wild-type and GAS motif mutation Luc-report plasmids. (**E**) The pGL3-bmp8a-promoter, pGL3-bmp8a-promoter-ΔP1, pGL3-bmp8a-promoter-ΔP2 or pGL3-bmp8a-promoter-ΔP1 and ΔP2 was transfected into EPC cells along with stat1a, stat1b or empty vector, respectively. After 48 h, the transfected cells were collected for luciferase assays. *Renilla* luciferase was used as the internal control. Data were from three independent experiments. Values are expressed as mean ± SEM and analyzed by Student's *t*-test (**p < 0.01, ***p < 0.001). (**F, G**) SDS-PAGE and Western-blotting analysis of rStat1a (**F**) and rStat1b (**G**). Lane M: protein molecular standard; Lane 1: negative control for IPTG induced *E. coli* (without rStat1a or rStat1b); Lane2: induced rStat1a or rStat1b (the whole cell lysate); Lane 3: induced rStat1a or rStat1b (the supernatant); Lane 4: purified rStat1a or rStat1b; Lane 5: western blot analysis of the sample in Lane 4. (**H**–**K**) EMSA was performed to validate the interaction of rStat1a or rStat1b with the GAS motif (P1 or P2) in the bmp8a promoter region. Lane 1: negative group; Lane 2: positive group; Lane 3: an excess unlabeled competitor probe; Lane 4: an excess unlabeled competitor probe containing a mutated runt binding site; Lane 5: Super-shift assays were performed by adding antibody against His tag. Source data are provided as a Source Data file.

Figure 9. Schematic illustration of Bmp8a serving as positive regulator in the antiviral immune. Upon virus infection, the transcriptions of *bmp8a* are activated through the Jak-Stat1 pathway. The Bmp8a binds to BMP type I receptor Alk6a, promoting phosphorylation of Tbk1 and Irf3 to induce the expression of *Ifn* through p38 MAPK pathway.

Figure 1 – Figure supplement 1. Generation of zebrafish *bmp8a* mutant alleles using TALENs technology.

(**A**) Partial structure and sequence of the *bmp8a* gene, showing the binding sites (underlined) of the left and right

TALENs. (**B**) Sequences of *bmp8a* homozygous mutant. Deletions of 7 bp are indicated by dashes letters. The

sequences of *bmp8a* in wild type (WT) zebrafish were used as a control. (**C**) Bmp8a in WT, 434 amino acids (aa)

long, contains a TGFβ-propeptide domain (yellow) and a TGFβ domain (green). The *bmp8a*$^{-/-}$ encodes a truncated

281-amino-acid-long polypeptide that contains a partial TGFβ-propeptide domains (yellow) followed by a

frameshift leading to a premature stop codon.

Figure 1 – Figure supplement 2. A new virus (TSVDV) was isolated from the focal site of turbot skin. (**A**)

Skin ulceration appeared on the skin surface (green arrow) of turbot infected with TSVDV (***Qin et al., 2008***). (**B**)

The FG cells had obvious cytopathic effect (CPE) after TSVDV infection.

Figure 3 – Figure supplement 1. Kyoto Encyclopedia of Genes and Genomes (KEGG) analysis of the top

10 pathways by enrichment of differential expressed genes (DEGs) between *bmp8a*$^{-/-}$ and WT liver tissue

of zebrafish challenged by GCRV. Rich ratio is the ratio of differentially expressed gene numbers annotated in

this pathway terms to all gene numbers annotated in this pathway term. The *q*-value means corrected p-value, q

< 0.05 as significantly enriched.

Figure 8 – Figure supplement 1. Sequence analysis of *bmp8a* promoter region. Two GAS motifs (P1 and P2)

have been found in the *bmp8a* promoter region, which were underlined. The transcriptional start site was

designated by +1.

References

Thaiss, C. A., Levy, M., Itav, S. & Elinav, E. Integration of Innate Immune Signaling. *Trends Immunol*. **37**, 84-101

(2016).

Akira, S. & Hemmi, H. Recognition of pathogen-associated molecular patterns by TLR family. *Immunol. Lett*. **85**,

85-95 (2003).

Chen, S. N., Zou, P. F. & Nie, P. Retinoic acid-inducible gene I (RIG-I)-like receptors (RLRs) in fish: current

knowledge and future perspectives. *Immunology* **151**, 16-25 (2017).

Keating, S. E., Baran, M. & Bowie, A. G. Cytosolic DNA sensors regulating type I interferon induction. *Trends*

Immunol. **32**, 574-581 (2011).

Parvatiyar, K. et al. The helicase DDX41 recognizes the bacterial secondary messengers cyclic di-GMP and cyclic

di-AMP to activate a type I interferon immune response. *Nat. Immunol*. **13**, 1155-1161 (2012).

Honda, K. et al. IRF-7 is the master regulator of type-I interferon-dependent immune responses. *Nature* **434**, 772-

777 (2005).

Tamura, T., Yanai, H., Savitsky, D. & Taniguchi, T. The IRF family transcription factors in immunity and oncogenesis.

Annu. Rev. Immunol. **26**, 535-584 (2008).

Li, W., Hofer, M. J., Jung, S. R., Lim, S. L. & Campbell, I. L. IRF7-dependent type I interferon production induces

lethal immune-mediated disease in STAT1 knockout mice infected with lymphocytic choriomeningitis virus. *J. Virol*.

88, 7578-7588 (2014).

Schindler, C. & Plumlee, C. Inteferons pen the JAK-STAT pathway. *Semin. Cell Dev. Biol*. **19**, 311-318 (2008).

Decker, T., Kovarik, P. & Meinke, A. GAS elements: a few nucleotides with a major impact on cytokine-induced

gene expression. *J. Interferon Cytokine Res*. **17**, 121-134 (1997).

Fu, X. Y., Kessler, D. S., Veals, S. A., Levy, D. E. & Darnell, J. E. Jr. ISGF3, the transcriptional activator induced

by interferon alpha, consists of multiple interacting polypeptide chains. *Proc. Natl Acad. Sci. USA* **87**, 8555-8559

(1990).

Levy, D. E., Kessler, D. S., Pine, R. & Darnell, J. E. Jr. Cytoplasmic activation of ISGF3, the positive regulator of

interferon-alpha-stimulated transcription, reconstituted in vitro. *Genes Dev*. **3**, 1362-1371 (1989).

Levy, D. E., Kessler, D. S., Pine, R., Reich, N. & Darnell, J. E. Jr. Interferon-induced nuclear factors that bind a

shared promoter element correlate with positive and negative transcriptional control. *Genes Dev*. **2**, 383-393

(1988).

Ren, Y. et al. Deubiquitinase USP2a Sustains Interferons Antiviral Activity by Restricting Ubiquitination of Activated

STAT1 in the Nucleus. *PLoS Pathog*. **12**, e1005764 (2016).

Shuai, K., Schindler, C., Prezioso, V. R. & Darnell, J. E. Jr. Activation of transcription by IFN-gamma: tyrosine

phosphorylation of a 91-kD DNA binding protein. *Science* **258**, 1808-1812 (1992).

Bragdon, B., Moseychuk, O., Saldanha, S., King, D., Julian, J. & Nohe, A. Bone morphogenetic proteins: a critical

review. *Cell Signal* **23**, 609-620 (2011).

Wu, M., Chen, G. & Li, Y. P. TGF-beta and BMP signaling in osteoblast, skeletal development, and bone formation,

homeostasis and disease. *Bone Res*. **4**, 16009 (2016).

Huber, S. & Schramm, C. Role of activin A in the induction of Foxp3+ and Foxp3- CD4+ regulatory T cells. *Crit*.

Rev. Immunol. **31**, 53-60 (2011).

Martínez, V. G. et al. The canonical BMP signaling pathway is involved in human monocyte-derived dendritic cell

maturation. *Immunol. Cell Biol*. **89**, 610-618 (2011).

Martínez, V. G. et al. The BMP Pathway Participates in Human Naive CD4+ T Cell Activation and Homeostasis.

PloS One **10**, e0131453 (2015).

Phillips, D. J., de Kretser, D. M. & Hedger, M. P. Activin and related proteins in inflammation: not just interested

bystanders. *Cytokine Growth Factor Rev*. **20**, 153-164 (2009).

Seeger, P., Musso, T. & Sozzani, S. The TGF-β superfamily in dendritic cell biology. *Cytokine Growth Factor Rev*.

26, 647-657 (2015).

Takabayashi, H. et al. Anti-inflammatory activity of bone morphogenetic protein signaling pathways in stomachs of

mice. *Gastroenterology* **147**, 396-406 (2014).

Yasmin, N. et al. Identification of bone morphogenetic protein 7 (BMP7) as an instructive factor for human

epidermal Langerhans cell differentiation. *J. Exp. Med*. **210**, 2597-2610 (2013).

Sivertsen, E. A., Huse, K., Hystad, M. E., Kersten, C., Smeland, E. B. & Myklebust, J. H. Inhibitory effects and

target genes of bone morphogenetic protein 6 in Jurkat TAg cells. *Eur. J. Immunol*. **37**, 2937-2948 (2007).

Varas, A. et al. Interplay between BMP4 and IL-7 in human intrathymic precursor cells. *Cell cycle* **8**, 4119-4126

(2009).

Chen, W. & Ten Dijke, P. Immunoregulation by members of the TGFβ superfamily. *Nat. Rev. Immunol*. **16**, 723-

740 (2016).

Zhao, G. Q. & Hogan, B. L. Evidence that mouse Bmp8a (Op2) and Bmp8b are duplicated genes that play a role

in spermatogenesis and placental development. *Mech. Dev*. **57**, 159-168 (1996).

Walton, K. D. et al. Villification in the mouse: Bmp signals control intestinal villus patterning. *Development* **143**,

427-436 (2016).

Ying, Y., Qi, X. & Zhao, G. Q. Induction of primordial germ cells from murine epiblasts by synergistic action of

BMP4 and BMP8B signaling pathways. *Proc. Natl Acad. Sci. USA* **98**, 7858-7862 (2001).

Zhong, S. j. et al. Spatial and temporal expression of *bmp8a* and its role in regulation of lipid metabolism in

zebrafish *Danio rerio. Gene Rep.* **10**, 33-41 (2018).

Robertsen, B. The interferon system of teleost fish. *Fish Shellfish Immunol.* **20**, 172-191 (2006).

Zhang, Y. B. et al. The innate immune response to grass carp hemorrhagic virus (GCHV) in cultured Carassius

auratus blastulae (CAB) cells. *Dev. Comp. Immunol.* **31**, 232-243 (2007).

Derynck, R., Akhurst, R. J. & Balmain, A. TGF-beta signaling in tumor suppression and cancer progression. *Nat.*

Genet. **29**, 117-129 (2001).

Drummond, A. E. TGFbeta signalling in the development of ovarian function. *Cell Tissue Res.* **322**, 107-115 (2005).

Oh, S. P., Yeo, C. Y., Lee, Y., Schrewe, H., Whitman, M. & Li, E. Activin type IIA and IIB receptors mediate Gdf11

signaling in axial vertebral patterning. *Genes Dev.* **16**, 2749-2754 (2002).

Shi, Y. et al. Exosomal Interferon-Induced Transmembrane Protein 2 Transmitted to Dendritic Cells Inhibits

Interferon Alpha Pathway Activation and Blocks Anti-Hepatitis B Virus Efficacy of Exogenous Interferon Alpha.

Hepatology **69**, 2396-2413 (2019).

Heldin, C. H., Miyazono, K. & ten Dijke, P. TGF-beta signalling from cell membrane to nucleus through SMAD

proteins. *Nature* **390**, 465-471 (1997).

Shi, Y. & Massagué, J. Mechanisms of TGF-beta signaling from cell membrane to the nucleus. *Cell* **113**, 685-700

(2003).

Worthington, J. J., Fenton, T. M., Czajkowska, B. I., Klementowicz, J. E. & Travis, M. A. Regulation of TGFβ in the

immune system: an emerging role for integrins and dendritic cells. *Immunobiology* **217**, 1259-1265 (2012).

Tu, E., Chia, P. Z. & Chen, W. TGFβ in T cell biology and tumor immunity: Angel or devil? *Cytokine Growth Factor*

Rev. **25**, 423-435 (2014).

Aleman-Muench, G. R. & Soldevila, G. When versatility matters: activins/inhibins as key regulators of immunity.

Immunol. Cell Biol. **90**, 137-148 (2012).

Bleul, C. C. & Boehm, T. BMP signaling is required for normal thymus development. *J. Immunol.* **175**, 5213-5221

(2005).

Gordon, J., Patel, S. R., Mishina, Y. & Manley, N. R. Evidence for an early role for BMP4 signaling in thymus and

parathyroid morphogenesis. *Dev. Biol.* **339**, 141-154 (2010).

Lee, M. K. et al. TGF-beta activates Erk MAP kinase signalling through direct phosphorylation of ShcA. *EMBO J.*

26, 3957-3967 (2007).

Itoh, S. et al. Trps1 plays a pivotal role downstream of Gdf5 signaling in promoting chondrogenesis and apoptosis

of ATDC5 cells. *Genes Cells* **13**, 355-363 (2008).

Wu, H., Zhang, Y., Lu, X., Xiao, J., Feng, P. & Feng, H. STAT1a and STAT1b of black carp play important roles in

the innate immune defense against GCRV. *Fish Shellfish Immunol.* **87**, 386-394 (2019).

Zou, J. & Secombes, C. J. Teleost fish interferons and their role in immunity. *Dev. Comp. Immunol.* **35**, 1376-1387

(2011).

Ruan, B. Y. et al. Two type II IFN members, IFN-γ and IFN-γ related (rel), regulate differentially IRF1 and IRF11 in

zebrafish. *Fish Shellfish Immunol.* **65**, 103-110 (2017).

Boehm, U., Klamp, T., Groot, M. & Howard, J. C. Cellular responses to interferon-gamma. *Annu. Rev. Immunol.*

15, 749-795 (1997).

Reed, L. J. & Muench, H. A Simple Method of Estimating Fifty Percent Endo-Points. *Am. J. Hyg.* **27**, 493-797

(1938).

Huang, P., Xiao, A., Tong, X., Zu, Y., Wang, Z. & Zhang, B. TALEN construction via "Unit Assembly" method and

targeted genome modifications in zebrafish. *Methods* **69**, 67-75 (2014).

Feng, H., Zhang, Y. B., Zhang, Q. M., Li, Z., Zhang, Q. Y. & Gui, J. F. Zebrafish IRF1 regulates IFN antiviral

response through binding to IFNφ1 and IFNφ3 promoters downstream of MyD88 signaling. *J. Immunol.* **194**, 1225-

1238 (2015).

Sun, F., Zhang, Y. B., Liu, T. K., Shi, J., Wang, B. & Gui, J. F. Fish MITA serves as a mediator for distinct fish IFN

gene activation dependent on IRF3 or IRF7. *J. Immunol.* **187**, 2531-2539 (2011).

Mortazavi, A., Williams, B. A., McCue, K., Schaeffer, L. & Wold, B. Mapping and quantifying mammalian

transcriptomes by RNA-Seq. *Nat. Methods* **5**, 621-628 (2008).

Sun, C., Hu, L., Liu, S., Gao, Z. & Zhang, S. Functional analysis of domain of unknown function (DUF) 1943,

DUF1944 and von Willebrand factor type D domain (VWD) in vitellogenin2 in zebrafish. *Dev. Comp. Immunol.* **41**,

469-476 (2013).

Qin, L., Wang. Y. G. & Zhang, Z. Multiple verrucous protrusions on skin in turbot Scophthalmus maximus. *J. Dalian*

Ocean Univ. **23**, 479-483 (2008) (in Chinese with English abstract).

725
726
727
728

729 **Figure 1**

Figure 2

Figure 3

752 **Figure 4**

771   **Figure 5**

Figure 6

Figure 7

Figure 8

805   **Figure 9**

824 **Figure 1 – Figure supplement 1**

827 **Figure 1 – Figure supplement 2**

830 **Figure 3 – Figure supplement 1**

Figure 8 – Figure supplement 1

```
-1439  TCACTTTCGTCAATTGGTGAAGTTTGTTCCTCATGACTGGCTTGCTTGGTTTGGGACTTGTGGAGCT
-1372  GCGCATCGATGGATTTGTTCTTCAGTGTTTGGACTTTCAGTAGGGAAAATTAAACC
```
                                                                           GAS motif
```
-1316  ACACTGAACTGAACTTCAACTCTGAAAACTGGTCTGACTCCGTTTCAATTTACTAGAACTT
```
                                                                      P2
```
-1255  CTCTGTTAAGCTGATCTGACACAATCTACATTGTAAAAGTGCTATAGAAATAAAGATGAAATAAATT
-1188  TTTATAATATTTACCTAAGTACACTGCTCCAAAAAATAAAGGGAACACTTAAACAACACAATATAA
-1122  CCATGTGTTGCCTTTATCTTTTTAGCAGTATGTTTATGCATTTGGTGCTGTAATGAACAACTACTAAT
-1054  ATAACAATCTGAATATTCCTAACAGCCACTCTAAAAGTTTAAATAATATATTTTCAAGTGCACACGA
 -987  TATCAGTATTGTCACTATCACAATACATTGAATTGTTGAGTATCAACATACGTCTCATACACAGTGA
 -920  TCTCTCACTGCTTGGCTACCGCTGAACCCACAGAATACAAATTTAGTTTTCATATTACTGTAGCCAC
 -853  ACAAACTCCGTCCCATTCTGTGCAAAGACTGGATCACATCTCCACTCCAAACCACATGATAATTGT
 -787  CAGCAATTTGTGAGGTATGTGTAGAAGATGAGACGGTTTGATAGGGCAGATGGAGACTGAGGTGA
 -722  GAAGGGTTCAGGATAGAGAAAAGGGCGGAGGTTTGAGATGATGGAAACCCTCCCTCTGCGTAAG
 -658  CCTGTTTCTTGGAGGGAAGGAGCTCGCCTGTGTTCAGAGTAACCCTCCAGCGCAAGCAGCACACA
 -593  AGCTAAGAATACCACAGCTCAACAAAGTTCAGCGGTGAACCAATACAATTCCTTCTCAGAGTTTTC
 -527  AGGAGCCTTGGATTTTCCATCCTTTGGATTATTTGTTTTTTTTTTTAAAGCTCAACTTTTCATGGATTA
 -458  CCTCAAGACAGTTTTGTTTTTCAATTAAACAGAGGATTTCACCAGTGAAGAGTTCTCAAAAGAAGC
 -392  AGGTGTGCTTTTTAAAGGTTGACCGCTGTTTTTAGTTACTCTACCCAAAGACTTTCCATTCAACATT
 -325  GTGCTTTGTCCTTCCAACATCCAAAATCCAGTTTCCAGGATATATTGTCATCCATCTTTGTAAGTGC
 -258  ACTATAAATTATGAGATTGCTTTGGTAAATAAAGTGACGGCATGAGGAACGACTGTCAAGAGGCG
 -193  ATAAACTGAGCAAGTAAACATTGCACTCTTCCAAATAACGATGACTGCTGTGGAATCTGAGTGGCT
```
                                                                        GAS motif
```
-127  TAGACGTTCGAGTGTTGTTCGGGGTTGAACTTCGAGACGGTAGTCAGACCATTCCGGGAAACCA
```
                                                                    P1
```
 -62  ACGTCGGGTGAACAGCTAAAAGGACAAATATCTTTACAGACGGCTGGGCTCTTCGCTACACCA
```
       +1
```
   1  ATGGACAGACACGAGGTT
```

846 **Supplementary Table 1**

Primer names	Sequences (5' to 3')	Applications
irf3-F	GCTGTTCGTGCCCTGGTTGTATG	
irf3-R	GCACATCGTCGCTGTTGGAGTC	
irf7-F	GCTGGAATTGCTCAACTTGGTGG	
irf7-R	CAGTCTGTGGTCGGGAAGCGTAT	
mx-F	CCTGGCAGAATCTATGCTGAAAAC	
mx-R	AAGGCAGTTTTATTTTCGGTTGTG	
ifnφ1-F	GTGGAGGACCAGGTGAAGTT	
ifnφ1-R	GATTGACCCTTGCGTTGC	
ifnφ3-F	TCTATACCATTCCCGCAAAA	
ifnφ3-R	GGATCACCGAAGTTCTCAAAC	
tbk1-F	TGATTGGAGACGACGGATTC	
tbk1-R	CCACTCGATCTTCCCATTTTC	
alk2-F	TGGGTTGGTGCTGTGGGAGA	
alk2-R	CGGGTGAGCGGTCAGTAGTT	
alk3-F	GCATCACGCTTAACCATCC	qRT-PCR for
alk3-R	TCCCAAATCCACCAGAACA	zebrafish
alk6a-F	AGTGTCTCCGACAGATGGGG	
alk6a-R	TGTGAGTTCAGGGTTTGGGT	
bmpr2a-F	GAGATACCAAAACTCGACCCG	
bmpr2a-R	TCCTTGTGACCCCGAAACC	
bmpr2b-F	GTGATGATGGGCGAGTGTAA	
bmpr2b-R	CGATAAAGCGAGCAATGTTT	
actr2a-F	TTGATGGCGGGTTCTATGC	
actr2a-R	TCCTCGGATTCAGCGAGTG	
actr2b-F	TGCTATTTTACCACGACCCT	
actr2b-R	CACAACACCAGATAAACCTTGA	
bmp8a-F	TCGCTGGCTTCTCCATCCT	
bmp8a-R	ATCACATTCCCCGTCACAA	
actb1-F	CCGTGACATCAAGGAGAAGC	
actb1-R	TACCGCAAGATTCCATACCC	
EPC *actin*-F	TGTTCCAGCCATCCTTCTTG	
EPC *actin*-R	TGATTTTCATTGTGCTGGGG	
EPC *ifn*-F	ATGAAAACTCAAATGTGGACGTA	
EPC *ifn*-R	GATAGTTTCCACCCATTTCCTTAA	
EPC *irf3*-F	AACAAGAATGACACTGCGGA	
EPC *irf3*-R	AACTCGGGAGGGACTTTCAT	qRT-PCR for EPC
EPC *irf7*-F	AAAGTCTTCGTCAGCACCAGCG	cells
EPC *irf7*-R	CTCTCCGAAGCACAGGTAGATGGT	
EPC *tbk1*-F	TCAGAAGTTTGAGAACGGGAAGA	
EPC *tbk1*-R	CGTAGACCACGATGCGGTGTAAG	
EPC *mx*-F	GGCTGGAGCAGGTGTTGGTATC	
EPC *mx*-R	TCCACCAGGTCCGGCTTTGTTAA	
GCRV VP5-F	CTCCCCGTGAGCGTATTT	qRT-PCR for
GCRV VP5-R	GTTAGCAGCGGTAGTGACTTG	GCRV
P1 biotin probe-F	biotin-AGACCATTCCGGGAAAACCAA	
P1 biotin probe-R	biotin-TTGGTTTTCCCGGAATGGTCT	
P1 non-biotin probe-F	AGACCATTCCGGGAAAACCAA	
P1 non-biotin probe-R	TTGGTTTTCCCGGAATGGTCT	
P1 mutant non-biotin probe-F	AGACCAGCCCGGGGCAACCAA	
P1 mutant non-biotin probe-R	TTGGTTGCCCCGGGCTGGTCT	
P2 biotin probe-F	biotin-TTCAATTTACTAGAACTTCTC	EMSA
P2 biotin probe-R	Biotin-GAGAAGTTCTAGTAAATTGAA	
P2 non-biotin probe-F	TTCAATTTACTAGAACTTCTC	
P2 non-biotin probe-R	GAGAAGTTCTAGTAAATTGAA	
P2 mutant non-biotin probe-F	TTCAATGCACTAGGCCTTCTC	
P2 mutant non-biotin probe-R	GAGAAGGCCTAGTGCATTGAA	
bmp8a talen-F	TGGGCCTCGCTCTAAGC	
bmp8a talen-R	ACTGATCGAGACTTCTT	
bmp8a F1-F	GGAACTCTGAATCTGCGTCT	
bmp8a R1-R	TCACAGGAGGGCGAATAG	TALEN
bmp8a F2-F	GACAGACATTCAGGCACGTA	
bmp8a R2-R	GCCGTCCACTGCTATGAT	

847 Supplementary Table 1. Oligonucleotides used in this study.

848 **Supplementary Table 2**

Primer names	Sequences (5' to 3')	Applications
pcDNA3.1-bmp8a-F	CGGGATCCGCCACCATGGACAGACACGAGGT	Eukaryotic expression
pcDNA3.1-bmp8a-R	GGAATTCCTATAAACAGCCACAATTC	
pcDNA3.1-stat1a-F	GGGGTACCGCCACCATGACTCAGTGGTTGGAGC	
pcDNA3.1-stat1a-R	GCTCTAGACTATGTCATTGTCATCAAG	
pcDNA3.1-stat1b-F	GGGGTACCGCCACCATGACGCTCTGGAACCA	
pcDNA3.1-stat1b-R	GCTCTAGACTAAAACCTTGCAACGGGT	
pcDNA3.1-alk2-F	CCCAAGCTTGCCACCATGGGGCATTGCAGCAC	
pcDNA3.1-alk2-R	CGGGATCCTCAGCAGTCGGTTTTGCCCTT	
pcDNA3.1-alk3-F	CCCAAGCTTGCCACCATGCGTCAGCTTTTGT	
pcDNA3.1-alk3-R	CGGGATCCTCAGATTTTAATGTCTTGAGAT	
pcDNA3.1-alk6a-F	CGGGATCCGCCACCATGGATCTCTATAGGA	
pcDNA3.1-alk6a-R	GCTCTAGATCAGACTTTAATGTCCTGGGAT	
pcDNA3.1-actr2a-F	CCCAAGCTTGCCACCATGGGACCTGCAACAAA	
pcDNA3.1-actr2a-R	GGAATTCTCATAGACTAGACTCCTTTGGG	
pcDNA3.1-actr2b-F	CCCAAGCTTGCCACCATGTTCGCTTCTCTGCTC	
pcDNA3.1-actr2b-R	GGAATTCTCAGATGCTGGACTCTTTGGGC	
pcDNA3.1-bmpr2a-F	GGAATTCGCCACCATGGCAGTTGAAGGCCGAATA	
pcDNA3.1-bmpr2a-R	GCTCTAGATCAAAAGTCACAGTCATTAGGATCTG	
pcDNA3.1-bmpr2b-F	GGAATTCGCCACCATGAGAGTGCGGATCCTAAAC	
pcDNA3.1-bmpr2b-R	GCTCTAGATCACAGGCAGGTCATATCTGG	
pet-28a-stat1a-F	GGAATTCATGACTCAGTGGTTGGAGCTTC	Prokaryotic expression
pet-28a-stat1a-R	CCCAAGCTTTGTCATTGTCATCAAGTCTGC	
pet-28a-stat1b-F	CGGGATCCATGACGCTCTGGAACCAGCTG	
pet-28a-stat1b-R	CCGCTCGAGAAACCTTGCAACGGGTCTTG	
pGL3-bmp8a-promoter-F	GGGGTACCTCACTTTCGTCAATTGGTGAA	Luciferase assay
pGL3-bmp8a-promoter-R	CCGCTCGAGTGGTGTAGCGAAGAGCCCAGC	
pGL3-bmp8a-promoter-ΔP1-F	GGTCTGACTCCGTTTCAATTCTCTGTTAAGCTGATCTGACACAA	
pGL3-bmp8a-promoter-ΔP1-R	TTGAAACGGAGTCAGACCAGTTTTCAGAGTTGA	
pGL3-bmp8a-promoter-ΔP2-F	TAGTCAGACCACCAACGTCGGGTGAACAGCTAA	
pGL3-bmp8a-promoter-ΔP2-R	ACGTTGGTGGTCTGACTACCGTCTCGAAGTTCA	
pGL3-IFNφ1-promoter-F	TCTCGAGAAGTGAAGTGAAAGTGCA	
pGL3-IFNφ1-promoter-R	CAAGCTTGTTCTCATCTTTGCGTGT	
pGL3-IFNφ3-promoter-F	GGGTACCGTTCACTGAGCCTGCATG	
pGL3-IFNφ3-promoter-R	TCTCGAGGATGCCTATACGGAATCA	
pGL3-EPC IFN-promoter-F	GGGGTACCGACCTTGAAATACTTTGGAATCAGGT	
pGL3-EPC IFN-promoter-R	CCCAAGCTTGCACAAACATATACGTCCACATT	
pcmv-bmp8a-HA-F	AGCCCGGGCGGATCCAAGCTTGCCACCATGGACAGACACGA	CO-IP
pcmv-bmp8a-HA-R	ATCGAATTCCTGCAGAAGCTTTAAACAGCCACAATTCTTGACCA	
pcmv-alk6a-Flag-F	CGCTCTAGCCCGGGCGGATCCGCCACCATGGATCTCTATAGGAGC	
pcmv-alk6a-Flag-R	TTCCTGCAGAAGCTTGGATCCGACTTTAATGTCCTGGGATTCAGAC	
pcDNA3.1-alk6a-ΔGS-F	ATAGAGCAATCCCAGCTCCCTCTACTGGTGCAGCG	Dominant negative mutant
pcDNA3.1-alk6a-ΔGS-R	GAGCTGGGATTGCTCTATGAGATCCCGCAGAGA	
pcDNA3.1-irf3DN-F	GGGGTACCGCCACCATGGGATCTGAGGGCTCTCAA	
pcDNA3.1-irf3DN-R	GGAATTCTTAGCAGAGCTCCATCATTTG	
pcDNA3.1-irf7DN-F	GGGGTACCGCCACCATGAAGAGTGCGATCCAGAGTG	
pcDNA3.1-irf7DN-R	GGAATTCTTATTCCACTGAAGGCAGACC	
pcDNA3.1-stat1a-ΔC-F	TCCGAGCTCGGTACCAAGCTTATGACTCAGTGGTTGGAGCTTCA	
pcDNA3.1-stat1a-ΔC-R	CAGCGGTTTAAACTTAAGCTTTCAAATGATGTCAGGAAGTGAAACA	
pcDNA3.1-stat1b-ΔC-F	TCCGAGCTCGGTACCAAGCTTATGACGCTCTGGAACCAGCTG	
pcDNA3.1-stat1b-ΔC-R	CAGCGGTTTAAACTTAAGCTTTCACATGGAGATGGTTGCCAG	
pcDNA3.1-tbk1 K38M-F	CCGTCATGGTGTTCAACAACCTGGTGTTCAACAACCTGAGTTTCCTG	
pcDNA3.1-tbk1 K38M-R	TGTTGAACACCATGACGGCGTACAGATTGACGGCGTACAGATCGCC	

849   Supplementary Table 2. Oligonucleotides used in this study.