The lncRNA *MARS* modulates the epigenetic reprogramming of the marneral cluster in response to ABA

Thomas Roulé[1,2], Federico Ariel[3], Caroline Hartmann[1,2], Jose Gutierrez-Marcos[4], Nosheen Hussain[4], Martin Crespi[1,2]* and Thomas Blein[1,2]

[1] Institute of Plant Sciences Paris-Saclay, Centre Nationale de la Recherche, Institut National de la Recherche Agronomique, Université Evry, Université Paris-Saclay, 91405 Orsay, France

[2] Institute of Plant Sciences Paris-Saclay, Université de Paris, 91405 Orsay, France

[3] Instituto de Agrobiotecnología del Litoral, CONICET, FBCB, Universidad Nacional del Litoral, Colectora Ruta Nacional 168 km 0, 3000 Santa Fe, Argentina

[4] School of Life Sciences, University of Warwick, Coventry, UK

*Correspondence to: MC (martin.crespi@universite-paris-saclay.fr)

ABSTRACT

Clustered organization of biosynthetic non-homologous genes is emerging as a characteristic feature of plant genomes. The co-regulation of clustered genes seems to largely depend on epigenetic reprogramming and three-dimensional chromatin conformation. Here we identified the long noncoding RNA (lncRNA) *MARneral Silencing* (*MARS*), localized inside the Arabidopsis marneral cluster, and which controls the local epigenetic activation of its surrounding region in response to ABA. *MARS* modulates the POLYCOMB REPRESSIVE COMPLEX 1 (PRC1) component LIKE-HETEROCHROMATIN PROTEIN 1 (LHP1) binding throughout the cluster in a dose-dependent manner, determining H3K27me3 deposition and chromatin condensation. In response to ABA, *MARS* decoys LHP1 away from the cluster and promotes the formation of a chromatin loop bringing together the *MARNERAL SYNTHASE 1* (*MRN1*) locus and a distal ABA-responsive enhancer. The enrichment of co-regulated lncRNAs in clustered metabolic genes suggests that the acquisition of noncoding transcriptional units constitute an additional regulatory layer driving the evolution of biosynthetic pathways.

KEYWORDS: lncRNA, enhancer, cluster, chromatin conformation, LHP1, ABA, seed germination, epigenetics, marneral

INTRODUCTION

In eukaryotes, functionally related genes are usually scattered across the genome. However, a growing number of operon-like clustered organization of non-homologous genes participating in common metabolic pathways point at an emerging feature of animal, fungi and plant genomes[1].

In plants, synthesis of numerous secondary metabolic compounds is important for the dynamic interaction with their environment, affecting their life and survival[2]. Terpenoids are bioactive molecules of diverse chemical structure[3]. In *Arabidopsis thaliana*, the biosynthesis of four triterpenes, namely thalianol[4], tirucalla-7,24-dien-3b-ol[5], arabidiol[6] and marneral[7], is governed by enzymes encoded by genes organized in clusters[8]. The thalianol and marneral related genes are located in the smallest metabolic clusters identified in plants to date, each being less than 40kb in size[8]. Both compounds are derived from 2,3-oxidosqualene and the corresponding gene clusters contain the oxidosqualene cyclases (OSCs), thalianol synthase (THAS) and marneral synthase (MRN1), respectively. The marneral cluster includes two additional protein-coding genes, *CYP705A12* and *CYP71A16*, participating in marneral oxidation[7].

Growing evidence indicates that the co-regulation of clustered genes relies on epigenetic mechanisms. It has been shown that the deposition of the histone variant H2A.Z positively correlates with transcriptionally active clusters. Accordingly, nucleosome stability precluding gene expression is dependent on ARP6, a component of the SWR1 chromatin remodeling complex required for the deposition of H2A.Z into nucleosomes[9]. Additionally, it was shown that the thalianol and marneral clusters exhibit increased expression in the Polycomb mutant *curly leaf* (*clf*) with compromised H3K27me3 deposition, and reduced expression in the trithorax-group protein mutant *pickle* (*pkl*), a positive regulator that counteracts H3K27me3 silencing[10]. Strikingly, it has been recently shown that biosynthetic gene clusters are embedded in local hot spots of three-dimensional (3D) contacts that segregate cluster regions from the surrounding chromosome environment in a tissue-dependent manner. Notably, H3K27me3 appeared as a central feature of the 3D domains at silenced clusters[11].

Long noncoding RNAs (lncRNAs) have emerged as important regulators of eukaryotic gene expression at different levels[12]. In plants, several lncRNAs have been shown to interact with the Polycomb Repressive Complex 1 and 2 components LIKE HETEROCHROMATIN PROTEIN 1 (LHP1) and CLF, respectively, which are related to H3K27me3 distribution[13,14]. Furthermore, it has been proposed that lncRNAs can modulate the transcriptional activity of

neighboring genes by shaping local 3D chromatin conformation[15–17]. Here we show that the
marneral cluster in Arabidopsis includes three noncoding transcriptional units. Among them,
the lncRNA *MARS* influences the expression of marneral cluster genes in response to ABA
through modification of the epigenetic landscape. *MARS* deregulation affects H3K27me3
distribution, LHP1 deposition and chromatin condensation throughout the cluster.
Furthermore, an ABA responsive chromatin loop dynamically regulates *MRN1* transcriptional
activation by bringing together the *MRN1* proximal promoter and an enhancer element
enriched in ABA-related transcription factors (TF) binding sites. *MARS*-mediated control of
the marneral cluster affects seed germination in response to ABA. The general co-regulation
of genes located within lncRNA-containing clusters in Arabidopsis points to noncoding
transcription as an important feature in coordinated transcriptional activity of clustered loci.

RESULTS

The marneral gene cluster includes three noncoding transcriptional
units

The small marneral cluster includes three genes: marneral synthase (*MRN1*),
CYP705A12 and *CYP71A16* that *are* two P450 cytochrome-encoding genes (**Fig. 1a**), all
participating in the biosynthesis and metabolism of the triterpene marneral[7].

The advent of novel sequencing technologies has allowed the identification of an
increasing number of lncRNAs throughout the *Arabidopsis* genome. According to the latest
annotation (Araport 11[18]), three additional transcriptional units are located within the marneral
cluster, between the *CYP71A16* and the *MRN1* loci. The *AT5G00580* and the pair of
antisense genes *AT5G06325* and *AT5G06335* are located upstream of the *MRN1* gene at
6kpb and 3kbp, respectively (**Fig. 1a**). The 1,941bp-long *AT5G00580* locus generates four
transcript isoforms ranging from 636 nt to 1,877 nt in length (**Fig. 1b and 1c**). In contrast,
each of the antisense genes *AT5G06325* and *AT5G06335* are transcribed into only one RNA
molecule of 509 nt and 367 nt, respectively (**Fig. 1a**). All these transcripts were classified as
lncRNAs when using two coding prediction tools, CPC[19] and CPC2[20], because of their low
coding potential and their length (over 200 nt), similarly to previously characterized lncRNAs
(*COLDAIR*[21]; *APOLO*[15]; and *ASCO*[22]) (**Fig. 1d**).

According to available transcriptomic datasets (Araport11), *AT5G00580*
transcriptional accumulation positively correlates with that of marneral genes, whereas
AT5G06325 and *AT5G06335* RNAs do not (**Supplementary Fig. 1**). Notably, our analysis of

the transcriptional behavior of the noncoding gene *AT5G00580* and the marneral cluster
protein-coding genes revealed a correlated expression in response to phosphate and nitrate
starvation, heat stress, as well as to exogenous auxin and ABA (**Fig. 1e**). Interestingly, the
AT5G00580 lncRNA exhibited the strongest induction in response to heat stress and
exogenous ABA, in comparison with *MRN1* and the two *CYP* genes (**Fig. 1e**). Altogether, our
observations uncovered that the marneral cluster includes three noncoding transcriptional
units, one of which is actively transcribed and co-regulated with its neighboring protein-
coding genes.

The lncRNA *MARS* shapes the transcriptional response of the marneral gene cluster to ABA

It has been shown that lncRNAs can regulate the expression of their neighboring
genes through epigenetic mechanisms[14,23]. Thus, we wondered if the lncRNA derived from
the *AT5G00580* locus may regulate the transcriptional activity of the protein-coding genes
included in the marneral cluster. To this end, we modified the lncRNA expression without
affecting the cluster DNA region using an RNAi construct targeting the first exon of
AT5G00580, and isolated two independent lines. The RNAi line 1 impaired the transcriptional
accumulation of AT5G00580 without affecting the rest of the cluster. Interestingly, the RNAi
line 2 exhibited a strong down-regulation of *AT5G00580* together with a significant basal
repression of the two *CYP* genes (**Fig. 2a**). Strikingly, the response of the three protein-
coding genes of the marneral cluster to exogenous ABA was significantly deregulated in both
independent RNAi lines. Transcriptional levels of *MRN1* and the two *CYP* genes increased
earlier in RNAi seedlings (15 min) than in the wild-type (Col-0, 30 min) (**Fig. 2b bottom**
panel). In addition, the transcriptional accumulation of these genes later reached three-fold
higher levels in the RNAi lines compared to Col-0 (**Fig. 2b top panel**). To further support our
observations, we isolated the insertional mutant *SALK_133089* located 200 bp upstream the
transcription start site (TSS) of *AT5G00580* gene. Although this insertion did not affect
AT5G00580 basal levels (**Fig. 2a**), it partially impaired its induction in response to ABA. In
agreement with both RNAi lines, *MRN1* and *CYP705A12* genes were strongly induced by
exogenous ABA (**Supplementary Fig. 2**), in contrast to *CYP71A16,* whose promoter region
may be locally affected by the T-DNA insertion. Notably, *MRN1*, which encodes the marneral
synthase, is the gene in the cluster most strongly affected by *AT5G00580* down-regulation.
Collectively, our results indicate that the noncoding transcriptional activity derived from the
AT5G00580 locus represses the dynamic expression of the marneral cluster genes, mainly
MRN1, in response to ABA. Therefore, we named *the AT5G00580*-derived noncoding
transcript *MARneral Silencing* (*MARS*) lncRNA.

MARS affects seed germination likely through its impact on *MRN1* expression

The phytohormone ABA has been implicated in the perception and transduction of environmental signals participating also in a wide range of growth and developmental events such as seed maturation, dormancy and germination[24]. Considering that the marneral cluster exhibited a strong *MARS*-dependent response to ABA, we wondered what the physiological impact of *MARS* deregulation was during seed germination. To this end, we assessed seed germination in Col-0 and *MARS* down-regulated lines (RNAi lines 1/2 and *SALK_133089*) with or without exogenous ABA. Notably, *MARS* silencing resulted in a delayed germination compared to Col-0 (**Supplementary Fig. 3a**), as revealed by an increase in T50 (time for 50% of germination) of nearly 5 hours (**Supplementary Fig. 3b**). Interestingly, in response to 0.5µM ABA, the germination of RNAi-*MARS* was further delayed than Col-0 with an increase of T50 of nearly 10 hours (**Supplementary Fig. 3c and 3d**). Accordingly, transgenic plants over-expressing *MRN1* also exhibited a delayed germination phenotype regardless of the treatment with ABA (**Supplementary Fig. 3**). The behavior of *35S:MRN1* seedlings suggests that *MRN1* up-regulation in RNAi-*MARS* lines could be linked to the increased sensitivity to ABA (**Fig. 2b**). Altogether, our results indicate that *MARS* can modulate seed germination through the regulation of the expression of *MRN1*.

MARS controls the epigenetic status of the marneral locus

It has been shown that gene clusters in plants are tightly regulated by epigenetic modifications, including the repressive mark H3K27me3[10]. According to publicly available ChIP-Seq datasets[25], the marneral cluster region is highly enriched in H3K27me3 deposition in shoots, extensively coinciding with LHP1 recognition (**Supplementary Fig. 4**). Consistently, an ATAC-Seq available dataset[26] revealed that the marneral cluster exhibits a high chromatin condensation in shoots (**Supplementary Fig. 4**). Thus, the marneral cluster is characterized by an epigenetically silent status in aerial organs coinciding with its low expression level[10].

We wondered if the transcriptional activation of the marneral cluster in response to exogenous ABA was associated with a dynamic epigenetic reprogramming. We first assessed H3K27me3 deposition across the marneral cluster, including the gene body of *MRN1*, *MARS* and the two *CYP* loci (**Fig. 3a and Supplementary Fig. 5**). Interestingly,

exogenous ABA triggered a strong reduction of H3K27me3 deposition throughout the
marneral cluster (**Fig. 3a and Supplementary Fig. 5**). Strikingly, H3K27me3 basal levels
were also significantly lower in RNAi-*MARS* seedlings. Remarkably, H3K27me3 deposition
was even lower across the body of all genes of the cluster in response to ABA in the RNAi-
MARS lines when compared with Col-0, in agreement with the stronger induction by ABA of
this subset of genes upon *MARS* silencing (**Fig. 3a, Supplementary Fig. 5 and Fig. 2b**).
Furthermore, we assessed LHP1 recognition of the marneral cluster. In agreement with
previous observations (**Supplementary Fig. 4**[25]), LHP1 distribution was high in *MRN1*
promoter and more weakly across *MARS* gene body and the intergenic region between
CYP71A16 and *MARS* (**Fig. 3b and Supplementary Fig. 6**). Remarkably, LHP1 recognition
was strongly impaired in response to ABA as well as in RNAi-*MARS* seedlings (**Fig. 3b and**
Supplementary Fig. 6). Therefore, our results indicate that ABA triggers an epigenetic
reprogramming of the marneral cluster, likely in a process involving the lncRNA *MARS*.

MARS is directly recognized by LHP1 and modulates local chromatin condensation

It has been shown that the deposition of the repressive mark H3K27me3 and the
concomitant recognition of the plant PRC1 component LHP1 are correlated with high
chromatin condensation[27]. Therefore, we determined by Formaldehyde-Assisted Isolation of
Regulatory Elements (FAIRE) the chromatin condensation of the whole marneral cluster. In
contrast to Col-0 showing a highly condensed chromatin, RNAi-*MARS* seedlings exhibit a
lower chromatin condensation in control conditions. Notably, the global chromatin status of
the cluster was even less condensed in RNAi-*MARS* seedlings in response to ABA (**Fig. 4a**
and Supplementary Fig. 7), in agreement with a decrease of both H3K27me3 deposition
and LHP1 binding (**Fig. 3b and Supplementary Fig. 6**) and the concomitant transcriptional
activation of the clustered genes (**Fig. 2b**). Consistently with our observations, *lhp1* mutant
seedlings also showed a global chromatin decondensation in control conditions, comparable
to Col-0 in response to ABA. Notably, further chromatin decondensation triggered by ABA
was completely impaired (**Fig. 4b and Supplementary Fig. 8**), supporting the role of LHP1
in the dynamic epigenetic silencing of the marneral cluster.

It has been shown that LHP1 can recognize RNAs *in vitro*[13] and the lncRNA *APOLO*
in vivo[15]. Moreover, it has been proposed that *APOLO* over-accumulation can decoy LHP1
away from target chromatin[28]. Therefore, we wondered whether *MARS* lncRNA was able to
interact with the chromatin-related protein LHP1 participating in the modulation of the local
epigenetic environment. Thus, we first determined that *MARS* was enriched in the nucleus,

compared with total RNA, as the previously characterized lncRNAs *APOLO* and *ASCO* that
interact respectively with nuclear epigenetic and splicing machineries, and the spliceosome
structural ncRNA *U6* (**Supplementary Fig. 9a**). Then, we confirmed by RNA
immunoprecipitation (RIP) that LHP1 can interact with *MARS in vivo,* in contrast to the *MRN1*
mRNA or housekeeping RNAs taken as negative controls (**Fig. 4c**).

LHP1 binding to the marneral cluster was impaired both in response to exogenous
ABA (inducing *MARS*, **Fig. 1e and Supplementary Fig. 3b**) and in RNAi-*MARS* seedlings,
hinting at a stoichiometry-dependent action of *MARS* on LHP1 recognition. Therefore, we
used chromatin extracts from RNAi-*MARS* line 1 seedling, with very low *MARS* transcript
levels (**Fig. 2a**) to assess LHP1 recognition of the marneral cluster upon the addition of
increasing concentrations of *in vitro*-transcribed *MARS* RNA. Strikingly, we found that low
MARS RNA concentrations (between 0.01 and 0.1 µg of RNA; **Fig. 4d and Supplementary
Fig. 9b**) successfully promoted LHP1 binding to the cluster, in contrast to higher
concentrations (between 1 and 10 µg of RNA), supporting the relevance of *MARS*-LHP1
stoichiometry for LHP1-target recognition (**Fig. 4d and Supplementary Fig. 9b**). Altogether,
our results suggest that the physical interaction of the nuclear-enriched lncRNA *MARS* to
LHP1 modulates its binding to proximal chromatin likely participating in the modulation of the
dynamic chromatin condensation of the marneral cluster.

215   *MARS* expression modulates an LHP1-dependent chromatin loop
216   bringing together the *MRN1* locus and an ABA enhancer element

It was recently observed that the spatial conformation of cluster-associated domains
differs between transcriptionally active and silenced clusters. In Arabidopsis, segregating 3D
contacts are distinguished among organs, in agreement with the corresponding
transcriptional activity of clustered genes[11]. Therefore, we explored whether *MARS* could
participate in the dynamic regulation of the local 3D chromatin conformation modulating the
transcription of the marneral cluster. According to available HiC datasets[25,29] there is a
significant interaction linking the intergenic region between *CYP71A16* and *MARS* and the
MRN1 locus (indicated as "Chromatin loop" in **Fig. 5a**). By using Chromatin Conformation
Capture (3C), we determined that the formation of this chromatin loop drastically increased
over 30 min with exogenous ABA and remained formed for at least 4 hours (**Fig 5b**). Thus,
the formation of this chromatin loop positively correlates with the transcriptional accumulation
of the marneral cluster genes in response to ABA (**Fig. 2b**).

The *MARS* locus is encompassed in the ABA-dependent chromatin loop (**Fig. 5a**). In
order to determine the role of *MARS* in the modulation of local 3D chromatin conformation,
we assessed the formation of the chromatin loop in RNAi-*MARS* lines. Notably, RNAi-*MARS*
seedlings exhibit enhanced chromatin loop formation, which remained unchanged in
response to exogenous ABA (**Fig. 5b**). Interestingly, LHP1 has been implicated in shaping
local 3D conformation of target regions[25], suggesting that the LHP1-*MARS* module may
dynamically switch the epigenetic status of the marneral cluster from a condensed-linear to a
decondensed-3D structured chromatin conformation. Supporting this hypothesis, *lhp1* mutant
seedlings exhibited enhanced chromatin loop formation compared to Col-0 (**Fig. 5c**). Overall,
our results demonstrate that a chromatin loop within the marneral cluster is regulated by
LHP1 and the interacting lncRNA *MARS*, encoded in the region encompassed by the loop.

To better understand the role of the *MARS*-dependent chromatin loop in response to
ABA we looked for ABA-related *cis* regulatory sequences throughout the marneral cluster.
We extracted from ChIP-seq the binding distribution of 13 ABA-related transcription factors
(TFs)[30]. Interestingly, a high enrichment in ABA TF binding sites was found at the *MARS*
locus, as well as in the intergenic region between the *CYP71A16* and *MARS* loci, notably
surrounding the contact point brought into close spatial proximity with the *MRN1* locus by the
ABA-dependent 3D chromatin loop (**Fig. 5a**). We thus assessed the activating capacity of
this region, potentially acting as a distant enhancer element of the *MRN1* proximal promoter.
To this end, we made use of a *GUS*-based reporter system (as described in [31]), fusing two
regions of interest to a minimal *35S* promoter. Two additional DNA regions nearby the
putative enhancers were used as negative controls: one between *CYP705A12* and
CYP71A16 and the other at the 3' end of *AT5G42620* locus (**Fig. 5a** indicated in red). Among
the two putative enhancers regions tested, one was able to activate *GUS* expression
(Intergenic region 2, **Fig. 5a and 5d**), coinciding with the region showing a high enrichment
of ABA-related TF binding sites close to the chromatin loop anchor point (**Fig. 5a**).
Collectively, our results indicate that an ABA-driven chromatin loop brings into close spatial
proximity the *MRN1* locus and a transcriptional activation site likely acting as an ABA
enhancer element. Notably, this chromatin reorganization process depends on the LHP1-
MARS module.

Long noncoding RNAs as emerging regulators of gene clusters

Physically linked genes organized in clusters are generally coregulated[8]. Considering
that the lncRNA *MARS* is implicated in the regulation of the marneral cluster, we wondered
whether the presence of noncoding transcriptional units may constitute a relevant feature of

gene cluster organization. Therefore, we collected two lists of Arabidopsis clustered genes,
i.e. one of co-expressed neighboring genes[10] and one of predicted metabolic gene clusters
(by PlantiSMASH[32]). According to the latest Arabidopsis genome annotation (Araport11),
among the 390 clusters of co-expressed neighboring genes), 189 (48%) contain at least one
embedded lncRNA inside the cluster. Most importantly, among the 45 metabolic clusters, 28
(62%) include lncRNAs (**Fig. 6a**). Furthermore, among the clusters containing a lncRNA, a
correlation analysis based on the maximum strength of co-expression between a lncRNA
and any clustered gene revealed that the metabolic clusters exhibit a significantly higher
correlation than co-expressed clusters (**Fig. 6b**). Altogether, our analyses suggest that
lncRNA-mediated local epigenetic remodeling may constitute an emerging feature of non-
homologous genes metabolic clusters in plants.

DISCUSSION

The cell nucleus is a dynamic arrangement of DNA, RNAs and proteins[33,34]. Genome
topology has emerged as an important feature in the complex network of mechanisms
regulating gene activity and genome connectivity, leading to regionalized chromosomal
spatial distribution and the clustering of diverse genomic regions with similar expression
patterns[35].

In the last few years, noncoding transcription has been implicated in shaping 3D
nuclear organization[36]. Notably, RNase-A micro-injection into the nucleus revealed that long
nuclear-retained RNAs maintained euchromatin in a biologically active decondensed state,
whereas heterochromatin domains exhibited an RNA-independent structure[37,38]. More
recently, HiC analyses were performed in mammalian cells exposed or not to RNase, before
and after crosslinking, or upon transcriptional inhibition[39]. As a result, it was observed that
topologically associated domains (TAD) boundaries remained mostly unaffected by RNase
treatment, whereas compartmental interactions suffered a subtle disruption. In contrast,
transcriptional inhibition led to weaker TAD boundaries, hinting at different roles of steady-
state RNA vs. active transcription in nuclear organization[39].

In plants, several lncRNAs have been implicated in local chromatin conformation
dynamics affecting the transcriptional activity of neighboring genes[14,40]. Notably, the lncRNA
COLDWRAP participates in the formation of an intragenic chromatin loop blocking the
transcription of the flowering time regulator *FLOWERING LOCUS C* (*FLC*[16]) in response to
cold, in a process involving the recruitment of PRC2 by direct interaction with the component
CLF. The lncRNA *APOLO* also controls the transcriptional activity of its neighboring gene
PINOID (*PID*) by dynamically modulating the formation of an intergenic chromatin loop

encompassing the divergent promoter of *PID* and *APOLO*[15], in a process involving the PRC1
component LHP1. More recently, it was proposed that high levels of *APOLO* can decoy
LHP1 away from multiple loci in *trans*, modulating the 3D conformation of target genes[28]. In
rice, the expression of the leucine-rich repeat receptor kinase clustered genes *RLKs* is
modulated by the locally-encoded lncRNA *LRK ANTISENSE INTERGENIC RNA* (*LAIR*). It
was proposed that *LAIR* may directly recruit OsMOF (MALES ABSENT ON THE FIRST) and
OsWDR5 (WD REPEAT DOMAIN 5), involved in H4K16 acetylation and chromatin
remodeling[41]. Here we showed that the lncRNA *MARS* contributes to the co-regulation of a
set of physically linked genes in *cis* in Arabidopsis. We demonstrated that the relative
abundance of *in vitro*-transcribed *MARS* fine-tunes LHP1 binding to the cluster region in a
stoichiometry-dependent manner, thus explaining how *MARS* affects H3K27me3 deposition
and chromatin condensation. It has been shown in yeast that histone depletion boosts
chromatin flexibility and facilitates chromatin loop formation on the kilobase pair scale[42]. In
agreement thereof, we uncovered here the dynamic role of the LHP1-*MARS* module
affecting nucleosome distribution across the marneral cluster in response to ABA, thus
promoting the formation of an intra-cluster chromatin loop.

It has been recently observed that biosynthetic gene clusters are embedded in local
three-dimensionally organized hot spots that segregate the region from the surrounding
chromosome environment[11]. Here we showed that active noncoding transcriptional units
within the cluster may contribute to 3D conformation dynamics switching from silent to active
states. Our results indicated that the *MARS*-dependent chromatin loop may bring the *MRN1*
locus and a distal ABA-responsive element into close spatial proximity, likely acting as an
enhancer. Notably, *MARS*-dependent LHP1 and H3K27me3 removal in Col-0, RNAi-*MARS*
and the *lhp1* mutant correlated with chromatin decondensation, loop formation and increased
marneral genes transcriptional activity in response to ABA. According to this model,
chromatin loop conformation is related to LHP1 binding and is modulated by *MARS*. LHP1
recognition at basal *MARS* levels maintains a possibly linear conformation of the region,
precluding the enhancer-*MRN1* locus interaction, whereas the positively activating chromatin
loop is formed in the absence of LHP1. *MARS* transcriptional accumulation directly
modulates LHP1 binding to the marneral cluster and high levels of *MARS* then titrate LHP1
away from the cluster (**Fig. 6c**; in response to ABA). Interestingly, when *MARS* levels are too
low compared to basal levels (as in the RNAi lines), recruitment of LHP1 to the cluster is also
impaired (**Fig. 6c;** *MARS* repression).

In mammals, growing evidence supports the role of lncRNAs in chromatin
conformation determination[43] and enhancer activity (e.g. *PVT1*[44] and *CCAT1-L*[45]). Here, we

showed that the nuclear-enriched lncRNA *MARS* brings together the *MRN1* proximal
promoter and a putative enhancer element enriched in ABA-responsive TF binding sites.
Interestingly, it has been shown that human lncRNAs can modulate the binding of TFs to
their target chromatin (*DHFR*[46]) and *PANDA*[47], whereas TFs have been implicated in
chromatin loop formation in plants[35]. Furthermore, it was shown that in addition to the TF NF-
YA, the lncRNA *PANDA* interacts with the scaffold-attachment-factor A (SAFA) as well as
with PRC1 and PRC2 to modulate cell senescence[48]. Therefore, further research will be
needed to determine what ABA-responsive TFs are in control of the marneral cluster and to
elucidate how they participate in chromatin loop formation along the area, in relation with the
PRC1-interacting lncRNA *MARS*.

Plants are a tremendous source of diverse chemicals which are important for their life
and survival[10]. Marneral biosynthesis has been linked to root and leaf development, flowering
time and embryogenesis[2]. Here we found that the Arabidopsis marneral cluster is activated
by the phytohormone ABA, in a lncRNA-dependent epigenetic reprogramming. *MARS*
deregulation affects the cluster response to ABA, impacting seed germination. Interestingly,
noncoding transcription had already been associated with seed germination. The *DELAY OF*
GERMINATION 1 (*DOG1*) locus is a major actor regulating seed dormancy strength in
Arabidopsis. Indeed, an antisense lncRNA (*asDOG1*) is able to repress *DOG1* transcription
in mature plants. Notably, it was observed that *DOG1* suppression is released by shutting
down antisense transcription, which is induced by ABA and drought[49].

It was proposed that the marneral cluster was founded by the duplication of ancestral
genes, independent events of gene rearrangement and the recruitment of additional genes[7].
The exploration of the noncoding transcriptome in Arabidopsis recently served to identify
ecotype-specific lncRNA-mediated responses to the environment[50]. It was suggested that the
noncoding genome may participate in multiple mechanisms involved in ecotype adaptation.
Collectively, our results indicate that the acquisition of novel noncoding transcriptional units
within biosynthetic gene clusters may constitute an additional regulatory layer behind their
natural variation in plant evolution.

METHODS

Lead contact and materials availability

362 Further information and requests for resources and reagents should be directed to
363 and will be fulfilled by the Lead Contact, Martin Crespi (martin.crespi@ips2.universite-paris-
364 saclay.fr).

365 Plant lines generated in this study are available from the Lead Contact with a
366 completed Materials Transfer Agreement.

Lines selection and generation

368 All plants used in this study are in Columbia-0 background. RNAi-*MARS* were
369 obtained using the pFRN binary vector[51] bearing 250bp of the first exon of *MARS* gene (see
370 primers in **Supplementary Table 1**), previously sub-cloned into the pENTR/D-TOPO vector.
371 Arabidopsis plants were transformed using *Agrobacterium tumefaciens* Agl-0[52]. The T-DNA
372 inserted line *SALK_133089* was ordered to NASC (N633089).

Growth conditions

374 Seeds were sown in plates vertically disposed in a growing chamber in long day
375 conditions (16 h in light 150uE; 8 h in dark; 21°C) for all the experiments. Plants were grown
376 on solid half-strength MS medium (MS/2) supplemented with 0.7% sucrose, and without
377 sucrose for the germination assay. For nitrate starvation assay, KNO_3 and $Ca(NO_3)_2$ were
378 replaced from MS/2 by a corresponding amount of KCl and $CaCl_2$ respectively, 2.25 mM
379 NH_4HCO_3 was added for nitrate-containing medium. For the phosphate starvation assay,
380 growth medium contained 0.15 mM $MgSO_4$, 2.1 mM NH_4NO_3, 1.9 mM KNO_3, 0.34 mM $CaCl_2$,
381 0.5 µM KI, 10 µM $FeCl_2$, 10 µM H_3BO_3, 10 µM $MnSO_4$, 3 µM $ZnSO_4$, 0.1 µM $CuSO_4$, 0.1 µM
382 $CoCl_2$, 0.1 µM Na_2MoO_4, 0.5 g.L^{-1} sucrose supplemented with 500uM Pi for Pi containing
383 medium versus 10uM for Pi free medium. All media were supplemented with 0.8g/L agar
384 (Sigma-Aldrich, A1296 #BCBL6182V) and buffered at pH 5.6 with 3.4mM 2-(N-morpholino)
385 ethane sulfonic acid. For the treatment with exogenous ABA or auxin, seedlings were
386 sprayed with 10uM ABA and 10uM 1-Naphthaleneacetic acid (NAA), respectively. For heat
387 stress, plates were transferred to a growth chamber at 37°C under the same lightning
388 conditions. For nitrate and phosphate starvation assays, seedlings have been transferred at
389 day 12 after sowing (DAS) from respectively nitrate and phosphate containing medium to
390 nitrate and phosphate free medium. Finally, for seed germination assay, 0.5uM ABA was
391 supplemented or not to the medium. Germination rate was evaluated twice a day. Seeds
392 were considered germinated when the seed coats were perforated by elongating radicle. For
393 all the experiments, samples were taken from 12 DAS starting two hours after light
394 illumination, at different time-points, after cross-linking or not, depending on the experiment.

RT-qPCR

Total RNA was extracted from whole seedlings using TRI Reagent (Sigam-Aldrich) and treated with DNase (Fermentas) as indicated by the manufacturers. Reverse transcription was realized on 1ug total RNA using the Maxima Reverse Transcriptase (Thermo Scientific). qPCR was performed on a Light Cycler 480 with SYBR Green master I (Roche) in standard protocol (40 cycles, 60°C annealing). Primers used in this study are listed in **Supplementary Table 1**. Data were analyzed using the delta delta Ct method using *PROTEIN PHOSPHATASE 2A SUBUNIT A3* (*AT1G13320*) for gene normalization[53] and time 0 for time-course experiment.

Chromatin Immunoprecipitation (ChIP)

ChIP was performed using anti-IgG (Millipore,Cat#12-370), anti-H3K27me3 (Millipore, Cat#07-449) and anti-LHP1 (Covalab, Pab0923-P), as previously described[15], starting from two grams of seedlings crosslinked in 1% (v/v) formaldehyde. Chromatin was sonicated in a water bath Bioruptor Plus (Diagenode; 60 cycles of 30s ON and 30s OFF pulses at high intensity). ChIP was performed in an SX-8G IP-Star Compact Automated System (Diagenode). Antibody coated into Protein A Dynabeads (Invitrogen) were incubated 12 hours at 4 °C with the samples. Recovering of immunoprecipitated DNA was realized using Phenol:Chloroform:Isoamilic Acid (25:24:1, Sigma) followed by ethanol precipitation and analyzed by qPCR. For input samples, non-immunoprecipitated sonicated chromatin was processed in parallel.

In-vitro transcribed *MARS* RNA was obtained from a PCR product amplified from wild-type genomic DNA using the T7 promoter on the 5' PCR primer (**Supplementary Table 1**). PCR products were verified using agarose electrophoresis, and purified using NucleoSpin kit (Macherey-Nagel). 1µg of purified DNA was used for *in-vitro* transcription following the manufacturer instructions (HiScribe T7 High Yield RNA Synthesis Kit, NEB). Purified non-crosslinked chromatin obtained from five grams of *MARS* RNAi line 1 seedlings were resuspending in 1 mL of nuclei lysis buffer and split into five tubes. An increasing quantity of *MARS* RNA was added to each tube from 0 to 10 µg RNA and incubated under soft rotation during 3 h at 4 °C. Chromatin samples were then cross-linked using 1% (v/v) of formaldehyde for five minutes. Sonication and the following ChIP steps were performed as above.

Formaldehyde-Assisted Isolation of Regulatory Elements (FAIRE)

427   FAIRE was performed as described by [54]. After chromatin purification as for ChIP,
428   only 50 µl from the 500 µl of purified chromatin were used (diluted to 500 µl with 10 mM Tris-
429   HCl pH 8). For the qPCR, the same set of primers as for ChIP were used.

Nuclear purification

Non-cross-linked seedlings were used to assess the sub-cellular localization of RNAs.
To obtain the nuclear fraction, chromatin was purified as for ChIP and resuspended, after the
sucrose gradient, into 1mL of TRI Reagent (Sigam-Aldrich). For the total fraction, 200 µL of
cell suspension in cell resuspension solution, were collected and completed with 800 µL of
TRI Reagent to follow with the RNA extraction. RNA's samples were treated by DNase, and
RT was performed using random hexamers prior to qPCR analysis.

RNA immunoprecipitation (RIP)

For RIP, the *lhp1* mutant complemented with the *ProLHP1:LHP1:GFP*[55] were used
after 4 hours of treatment with ABA. After crosslinking and chromatin extraction as for ChIP,
ten percent of resuspended chromatin was conserved at -20 °C as the input. Chromatin was
sonicated in a water bath Bioruptor Plus (Diagenode; 5 cycles of 30 s ON and 30 s OFF
pulses at high intensity). Anti-LHP1 RIP was performed using the anti-GFP antibody (Abcam
ab290), as previously described[15]. Results were expressed as the percentage of cDNA
detected after IP taking the input value as 100%.

Chromosome conformation capture (3C)

3C was performed as described by [56]. Briefly, chromatin was extracted from two
grams of cross-linked seedlings as for ChIP. Overnight digestion at 37 °C were performed
using 400U of Hind III enzyme (NEB). Digested DNA was ligated during 5 h incubation at 16
°C with 100 U of T4 DNA ligase (NEB). DNA was recovered after reverse crosslinking and
Proteinase K treatment (Invitrogen) by Phenol:Chloroform:Isoamyl Acid (25:24:1; Sigma)
extraction and ethanol precipitation. Interaction frequency was calculated by qPCR using a
DNA region uncut by Hind III to normalize the amount of DNA.

Transcriptional activation assay in tobacco leaves

The *GUS* reporter system for validating the activity of the putative enhancer element
was adapted from [31]. Different DNA fragments were cloned in the GreenGate system[57] fused
to a minimal 35S promoter element from CAMV (synthesized by Eurofins Genomics). The

sub-unit B3 from 35S promoter element from CAMV[58] was synthesized and used as a

positive control. All primers used for cloning are indicated in **Supplementary Table 1**.

A. tumefaciens-mediated transient transformation was performed on 5-week-old

tobacco plants using a needle-less syringe. Together with enhancer constructs, another

vector containing mCherry driven by 35S promoter was co-transfected to control the

transformation efficiency. Two leaf discs were collected near the infiltration site. One, to

determine the transfection efficiency by mCherry fluorescence observation under

epifluorescent microscope. The second was used for GUS staining, as previously

described[59]. Samples were incubated 4 h in the dark at 37 °C before observation.

Identification of lncRNA loci in Arabidopsis gene clusters

The genes of co-expressed clusters were retrieved from [10]. The boundaries of the

gene clusters were extracted using Araport11 annotations. The boundaries of the metabolic

clusters were extracted from the plantiSMASH predicted clusters on Arabidopsis[32]. Using

Araport11 GFF, the lncRNAs (genes with a locus type annotated as "long_noncoding_rna",

"novel_transcribed_region" or "other_rna") present within the boundaries of the cluster were

retrieved.

Gene expression correlation analysis

To compute the correlation of expression in different organ of Arabidopsis we used

the 113 RNA-seq datasets that were used for the Araport11 annotations (10.1111/tpj.13415).

These datasets were generated from untreated or mock-treated wild-type Col-0 plants. After

removing the adaptors with Trim Galore with default parameters, the reads were mapped on

TAIR10 with STAR v2.7.2a (10.1093/bioinformatics/bts635) and the parameters '--

alignIntronMin 20 --alignIntronMax 3000'. Gene expression was then quantified with

featureCounts v2.0.0 (10.1093/bioinformatics/btt656) with the parameters "-B -C -p -s 0"

using the GFF of Araport11. Raw counts were then normalized by median of ratios using the

DESeq2 R package[60].

For the correlation of expression inside the marneral cluster, the expression of the

genes of the clusters and 25kb around it (four genes upstream and two downstream) were

collected for the correlation analysis. Pearson's correlations for each pair of genes were

computed after log 2 transformation of the normalized counts. The correlation value and

associated p-value were plotted with the corrplot R package[61].

Inside each co-expressed and metabolic clusters of genes, Pearson's correlation was computed between every possible pairs of lncRNA and coding gene as for the genes inside the marneral cluster. The maximum correlation value was kept as an indication of lncRNAs correlation with the genes of the cluster.

Quantification and statistical analyses

For all the experiments, at least two independent biological samples were considered. For RT-qPCR, each sample was prepared from a pool of 5 to 10 individual seedlings. For biochemistry assays (ChIP, FAIRE, nuclear purification, RIP and 3C) two to five grams of seedlings were prepared for each independent biological sample. For validation of enhancer function, the four leaf discs were taken from four independent tobacco plants. The tests used for statistical analyses are indicated in the respective figure legends. Statistical test and associated plots have been generated using R software (v3.6.3[62]) with the help of the tidyverse package[63].

ACKNOWLEDGMENTS

IPS2 benefits from the support of Saclay Plant Sciences-SPS (ANR-17-EUR-0007). We thank Jeremie Bazin and Aurélie Christ from IPS2 for the helpful discussion about results interpretation and design of the experiments. We also thank Moussa Benhamed (IPS2) for helpful advice on epigenetic regulation. We thank Olivier Martin for critical reading of the manuscript.

AUTHORS' CONTRIBUTIONS

TR, FA, TB and MC conceived and designed the experiments. TR performed the experiments. TR and TB analyzed the data. All authors discussed the results and edited the manuscript.

COMPETING FINANCIAL INTERESTS

The authors declare no competing financial interests.

REFERENCES

1.   Nützmann, H.-W., Scazzocchio, C. & Osbourn, A. Metabolic Gene Clusters in Eukaryotes. *Annu.*

515   *Rev. Genet.* **52**, 159–183 (2018).

516   2.   Go, Y. S. *et al.* Identification of marneral synthase, which is critical for growth and
517   development in Arabidopsis. *Plant J.* **72**, 791–804 (2012).

518   3.   Yasumoto, S., Fukushima, E. O., Seki, H. & Muranaka, T. Novel triterpene oxidizing activity of
519   Arabidopsis thaliana CYP716A subfamily enzymes. *FEBS Lett.* **590**, 533–540 (2016).

520   4.   Field, B. & Osbourn, A. E. Clusters in Different Plants. *Science (80-.).* **194**, 543–547 (2008).

521   5.   Boutanaev, A. M. *et al.* Investigation of terpene diversification across multiple sequenced
522   plant genomes. *Proc. Natl. Acad. Sci. U. S. A.* **112**, E81–E88 (2015).

523   6.   Castillo, D. A., Kolesnikova, M. D. & Matsuda, S. P. T. An effective strategy for exploring
524   unknown metabolic pathways by genome mining. *J. Am. Chem. Soc.* **135**, 5885–5894 (2013).

525   7.   Field, B. *et al.* Formation of plant metabolic gene clusters within dynamic chromosomal
526   regions. *Proc. Natl. Acad. Sci.* **108**, 16116–16121 (2011).

527   8.   Nützmann, H. W., Huang, A. & Osbourn, A. Plant metabolic clusters – from genetics to
528   genomics. *New Phytol.* **211**, 771–789 (2016).

529   9.   Nützmann, H. W. & Osbourn, A. Regulation of metabolic gene clusters in Arabidopsis thaliana.
530   *New Phytol.* **205**, 503–510 (2015).

531   10.   Yu, N. *et al.* Delineation of metabolic gene clusters in plant genomes by chromatin signatures.
532   *Nucleic Acids Res.* **44**, 2255–2265 (2016).

533   11.   Nützmann, H. *et al.* Active and repressed biosynthetic gene clusters have spatially distinct
534   chromosome states. 1–10 (2020) doi:10.1073/pnas.1920474117.

535   12.   Rinn, J. L. & Chang, H. Y. Long Noncoding RNAs: Molecular Modalities to Organismal
536   Functions. *Annu. Rev. Biochem.* **89**, 283–308 (2020).

537   13.   Berry, S., Rosa, S., Howard, M., Bühler, M. & Dean, C. Disruption of an RNA-binding hinge
538   region abolishes LHP1-mediated epigenetic repression. *Genes Dev.* **31**, 2115–2120 (2017).

539   14.   Lucero, L., Fonouni-Farde, C., Crespi, M. & Ariel, F. Long noncoding RNAs shape transcription
540   in plants. *Transcription* **00**, 1–12 (2020).

541   15.   Ariel, F. *et al.* Noncoding transcription by alternative rna polymerases dynamically regulates
542   an auxin-driven chromatin loop. *Mol. Cell* **55**, 383–396 (2014).

16.   Kim, D.-H. & Sung, S. Vernalization-triggered intragenic chromatin-loop formation by long

noncoding RNAs. *Dev. Cell* **176**, 100–106 (2017).

17.   Gagliardi, D. *et al.* Dynamic regulation of chromatin topology and transcription by inverted
repeat-derived small RNAs in sunflower. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 17578–17583
(2019).

18.   Cheng, C. Y. *et al.* Araport11: a complete reannotation of the Arabidopsis thaliana reference
genome. *Plant J.* **89**, 789–804 (2017).

19.   Kong, L. *et al.* CPC: Assess the protein-coding potential of transcripts using sequence features
and support vector machine. *Nucleic Acids Res.* **35**, 345–349 (2007).

20.   Kang, Y. J. *et al.* CPC2: A fast and accurate coding potential calculator based on sequence
intrinsic features. *Nucleic Acids Res.* **45**, W12–W16 (2017).

21.   Heo, J. B. & Sung, S. Vernalization-mediated epigenetic silencing by a long intronic noncoding
RNA. *Science (80-.).* **331**, 76–79 (2011).

22.   Bardou, F. *et al.* Long Noncoding RNA Modulates Alternative Splicing Regulators in
Arabidopsis. *Dev. Cell* **30**, 166–176 (2014).

23.   Jarroux, J., Morillon, A. & Pinskaya, M. Long Non Coding RNA Biology. *Adv. Exp. Med. Biol.*
1008, 1–46 (2017).

24.   Vishwakarma, K. *et al.* Abscisic acid signaling and abiotic stress tolerance in plants: A review
on current knowledge and future prospects. *Front. Plant Sci.* **8**, 1–12 (2017).

25.   Veluchamy, A. *et al.* LHP1 Regulates H3K27me3 Spreading and Shapes the Three-Dimensional
Conformation of the Arabidopsis Genome. *PLoS One* **11**, 1–25 (2016).

26.   Sijacic, P., Bajic, M., McKinney, E. C., Meagher, R. B. & Deal, R. B. Changes in chromatin
accessibility between Arabidopsis stem cells and mesophyll cells illuminate cell type-specific
transcription factor networks. *Plant J.* **94**, 215–231 (2018).

27.   Yang, X., Tong, A., Yan, B. & Wang, X. Governing the silencing state of chromatin: The roles of
polycomb repressive complex 1 in arabidopsis. *Plant Cell Physiol.* **58**, 198–206 (2017).

28.   Ariel, F. *et al.* R-Loop Mediated trans Action of the APOLO Long Noncoding RNA. *Mol. Cell* **77**,
1–11 (2020).

29.   Liu, C. *et al.* Genome-wide analysis of chromatin packing in Arabidopsis thaliana at single-gene
resolution. *Genome Res.* **26**, 1057–1068 (2016).

30. Song, L. *et al.* A transcription factor hierarchy defines an environmental stress response
network. *Science (80-.).* **354**, 598 (2016).

31. Yan, W. *et al.* Dynamic control of enhancer activity drives stage-specific gene expression
during flower morphogenesis. *Nat. Commun.* **10**, 1–16 (2019).

32. Kautsar, S. A., Suarez Duran, H. G., Blin, K., Osbourn, A. & Medema, M. H. PlantiSMASH:
Automated identification, annotation and expression analysis of plant biosynthetic gene
clusters. *Nucleic Acids Res.* **45**, W55–W63 (2017).

33. Cavalli, G. & Misteli, T. Functional implications of genome topology. *Nat. Struct. Mol. Biol.* **20**,
290–299 (2013).

34. Gibcus, J. H. & Dekker, J. The Hierarchy of the 3D Genome. *Mol. Cell* **49**, 773–782 (2013).

35. Rodriguez-Granados, N. Y. *et al.* Put your 3D glasses on: Plant chromatin is on show. *J. Exp.*
Bot. **67**, 3205–3221 (2016).

36. Quinodoz, S. & Guttman, M. Long non-coding RNAs: An emerging link between gene
regulation and nuclear organization. *Trends Cell Biol.* **24**, 651–663 (2014).

37. Caudron-Herger, M. *et al.* Coding RNAs with a non-coding function: Maintenance of open
chromatin structure. *Nucleus* **2**, (2011).

38. Caudron-Herger, M. & Rippe, K. Nuclear architecture by RNA. *Curr. Opin. Genet. Dev.* **22**, 179–
187 (2012).

39. Barutcu, A. R., Blencowe, B. J. & Rinn, J. L. Differential contribution of steady-state RNA and
active transcription in chromatin organization . *EMBO Rep.* **20**, 1–13 (2019).

40. Gagliardi, D. & Manavella, P. A. Short-range regulatory chromatin loops in plants. *New Phytol.*
1–6 (2020) doi:10.1111/nph.16632.

41. Wang, Y. *et al.* Overexpressing lncRNA LAIR increases grain yield and regulates neighbouring
gene cluster expression in rice. *Nat. Commun.* **9**, 1–9 (2018).

42. Diesinger, P. M., Kunkel, S., Langowski, J. & Heermann, D. W. Histone depletion facilitates
chromatin loops on the kilobasepair scale. *Biophys. J.* **99**, 2995–3001 (2010).

43. Gil, N. & Ulitsky, I. Regulation of gene expression by cis-acting long non-coding RNAs. *Nat.*
Rev. Genet. **21**, 102–117 (2020).

44. Cho, S. W. *et al.* Promoter of lncRNA Gene PVT1 Is a Tumor-Suppressor DNA Boundary

602   Element. *Cell* **173**, 1398-1412.e22 (2018).

603 45. Xiang, J. F. *et al.* Human colorectal cancer-specific CCAT1-L lncRNA regulates long-range
604   chromatin interactions at the MYC locus. *Cell Res.* **24**, 513–531 (2014).

605 46. Martianov, I., Ramadass, A., Serra Barros, A., Chow, N. & Akoulitchev, A. Repression of the
606   human dihydrofolate reductase gene by a non-coding interfering transcript. *Nature* **445**, 666–
607   670 (2007).

608 47. Hung, T. *et al.* Extensive and coordinated transcription of noncoding RNAs within cell-cycle
609   promoters. *Nat. Genet.* **43**, 621–629 (2011).

610 48. Puvvula, P. K. *et al.* Long noncoding RNA PANDA and scaffold-attachment-factor SAFA control
611   senescence entry and exit. *Nat. Commun.* **5**, (2014).

612 49. Yatusevich, R. *et al.* Antisense transcription represses Arabidopsis seed dormancy QTL DOG 1
613   to regulate drought tolerance . *EMBO Rep.* **18**, 2186–2196 (2017).

614 50. Blein, T. *et al.* Landscape of the non-coding transcriptome response of two Arabidopsis
615   ecotypes to phosphate starvation. *Plant Physiol.* **183**, pp.00446.2020 (2020).

616 51. Ariel, F. *et al.* Two direct targets of cytokinin signaling regulate symbiotic nodulation in
617   medicago truncatula. *Plant Cell* **24**, 3838–3852 (2012).

618 52. Clough, S. J. & Bent, A. F. Floral dip: A simplified method for Agrobacterium-mediated
619   transformation of Arabidopsis thaliana. *Plant J.* **16**, 735–743 (1998).

620 53. Czechowski, T., Stitt, M., Altmann, T., Udvardi, M. K. & Scheible, W. Genome-Wide
621   Identification and Testing of Superior Reference Genes for Transcript Normalization in
622   Arabidopsis. *Plant Physiol.* **139**, 5–17 (2005).

623 54. Simon, J. M., Giresi, P. G., Davis, I. J. & Lieb, J. D. Using formaldehyde-assisted isolation of
624   regulatory elements (FAIRE) to isolate active regulatory DNA. *Nat. Protoc.* **7**, 256–267 (2012).

625 55. Nakahigashi, K., Jasencakova, Z., Schubert, I. & Goto, K. The Arabidopsis HETEROCHROMATIN
626   PROTEIN1 homolog (TERMINAL FLOWER2) silences genes within the euchromatic region but
627   not genes positioned in heterochromatin. *Plant Cell Physiol.* **46**, 1747–1756 (2005).

628 56. Louwers, M., Splinter, E., van Driel, R., de Laat, W. & Stam, M. Studying physical chromatin
629   interactions in plants using Chromosome Conformation Capture (3C). *Nat. Protoc.* **4**, 1216–
630   1229 (2009).

57.   Lampropoulos, A. *et al.* GreenGate - A novel, versatile, and efficient cloning system for plant

transgenesis. *PLoS One* **8**, (2013).

58.   Moreno-risueno, M. A. *et al.* NIH Public Access. **329**, 1306–1311 (2010).

59.   Jefferson, R. A., Kavanagh, T. A. & Bevan, M. W. GUS fusions: ,B-glucuronidase as a sensitive

and versatile gene fusion marker in higher plants. *EMBO J.* **6**, 3901–3907 (1987).

60.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biol.* **15**, 1–21 (2014).

61.   Taiyun Wei and Viliam Simko (2017). R package "corrplot": Visualization of a Correlation

Matrix (Version 0.84). Available online at https://github.com/taiyun/corrplot

62.   R Core Team (2018). R: A language and environment for statistical computing. R Foundation

for Statistical Computing, Vienna, Austria. Available online at https://www.R-project.org/.

63.   Wickham, H. *et al.* Welcome to the Tidyverse. *J. Open Source Softw.* **4**, 1686 (2019).

Fig. 1, *AT5G00580* **is a lncRNA transcribed from the marneral cluster locus and its expression correlates with its neighboring genes**

a, Schematic illustration of the marneral cluster. Genes are indicated with plain rectangles and white arrows indicate the sense of transcription. The square indicates the region displayed in (B).

b, Schematic illustration of the different isoforms of *AT5G00580* transcripts. First line corresponds to *AT5G00580* genomic region whereas the other lines present the various isoforms. For each isoform, exons are indicated with rectangles and introns with solid lines. Black arrows indicate the primers used for the cDNA amplification in (C).

c, cDNA amplification of *AT5G00580* isoforms. The position of the primers used for the amplification are displayed on B. MWM stands for Molecular Weight Marker (GeneRuler 1 kb Plus DNA Ladder, Thermo Scientific).

d, Coding potential of the transcripts located in the marneral cluster genomic region. Scores were calculated using CPC1 (left) and CPC2 (right) programs[19,20]. For each algorithm, the threshold between coding and non-coding genes is displayed with a horizontal solid black line. Coding genes are situated above the threshold, whereas non coding genes are situated below. *COLDAIR*, *APOLO* and *ASCO* are used as positive controls of non-coding transcripts.

e, Dynamic transcriptional levels of co-regulated genes of the marneral cluster in response to phosphate and nitrate starvation, heat stress, and exogenous ABA and auxin. Gene expression data are represented as the mean ± standard error (n ≥ 3) of the log2 fold change compared to time 0h.

a Transcript abundance

b Transcript abundance

Fig. 2, *MARS* transcriptional activity modulates the response to ABA of the marneral cluster

a, Transcript levels of the marneral cluster genes in control conditions in RNAi lines targeting *AT5G00580*/*MARS* and *SALK_133089* line. Transcriptional levels are represented as the mean ± standard error (n = 3) of the log2 fold change compared to Col-0. Letters indicate statistic group determined by one-way analysis of variance (ANOVA) followed by Tukey's post-hoc test. For each gene, each letter indicates statistical difference between genotypes (p ≤ 0.05).

b, Transcript levels of the genes of the marneral cluster in response to ABA treatment in RNAi lines targeting *AT5G00580/MARS*. Gene expression data are represented as the mean ± standard error (n = 3) of the log2 fold change compared to time 0h.

Fig. 3, *MARS* modulates the epigenetic landscape of *MRN1* locus

a, H3K27me3 deposition over the *MRN1* gene in Col-0 and RNAi-*MARS* seedlings under control conditions and in response to ABA. Higher values of ChIP-qPCR indicate more H3K27me3.

b, LHP1 binding to the *MRN1* gene in Col-0 and RNAi-*MARS* seedlings in the same conditions as in (A). Higher values of ChIP-qPCR indicate more LHP1 deposition.In (A) and (B), values under the dotted line are considered as not enriched. Results are represented as the mean ± standard error (n = 2) of the H3K27me3/IgG or LHP1/IgG ratio. Numbers are p-value of the difference between the two genotypes determined by Student t-test.

Fig. 4, *MARS* **influences chromatin condensation of** *MRN1* **gene through its interaction with LHP1 protein**

a, Chromatin condensation in *MRN1* gene of Col-0 and RNAi-*MARS* seedlings in control conditions and in response to ABA, determined by Formaldehyde Assisted Isolation of Regulatory Element (FAIRE)-qPCR.

b, Evolution of the chromatin condensation in *MRN1* gene of Col-0 and *lhp1* mutant subjected to ABA treatment determined by Formaldehyde Assisted Isolation of Regulatory Element (FAIRE) qPCR.

In **a** and **b**, results are expressed as the mean ± standard error (n = 3) of the percentage of input (signal measured before isolation of decondensed region of chromatin, free of nucleosomes). Lower value indicates more condensed chromatin. Numbers are p-value of the difference between the two genotypes determined by Student t-test.

c, LHP1-*MARS* interaction was assessed by RNA immunoprecipitation (RIP) using *LHP1-GFP* seedlings. Negative controls include a housekeeping gene (*PP2A*) and *MRN1* mRNA. The interaction between *APOLO* and LHP1 was taken as a positive control[15]. Results are expressed as the mean ± standard error (n = 4) of the percentage of input (signal measured before immunoprecipitation).

d, LHP1 binding to the *MRN1* promoter region in chromatin from RNAi-*MARS* seedlings upon increasing amounts of *in-vitro* transcribed *MARS* RNA. After incubation (see Methods), the samples were crosslinked for LHP1 ChIP-qPCR. Higher values indicate more LHP1-DNA interaction. Results are expressed as the mean ± standard error (n = 2) of the LHP1/Igg ratio.

In **c** and **d** numbers are p-value of the difference between the different corresponding genes determined by Student t-test.

Fig. 5, An LHP1-dependent chromatin loop brings together the *MRN1* locus and a putative enhancer element in response to ABA

a, Schematic illustration of the loop linking the *MRN1* locus with the intergenic region between *CYP71A16* and *MARS*. Forward (F) and Reverse (R) oligonucleotides used for 3C-qPCR (in B–C) are indicated with arrows. The orange track shows the number of different ABA-related transcription factor binding sites (HB6, HB7, GBF2, GBF3, MYB3, MYB44, NF-YC2, NF-YB2, ANAC102, ANAC032, ABF1, ABF3, ABF4, RD26, ZAT6, FBH3, DREBA2A, AT5G04760, HAT22 and HSFA6A) found along the marneral cluster[30]. Green and red rectangles indicate the putative enhancer region and the negative controls, respectively, tested for the GUS-based reporter system in **d**.

b, Relative chromatin loop formation in response to ABA in Col-0 and RNAi-*MARS* seedlings. Results are expressed as the mean ± standard error (n = 2) from 3C-qPCR using primer F and R shown on **a**, compared to time 0h.

c, Relative chromatin loop formation in response to ABA treatment in Col-0 and *lhp1* mutant. Data are represented as the mean ± standard error (n = 3) from 3C-qPCR using primer F and R shown on **a**, compared to time 0h.

d, Constructs used for GUS-based reporter system are illustrated on the left. Corresponding transformed tobacco leaf discs are on the right (n = 4). First line represents the positive control in which 35S sub-unit controls *GUS* expression. The second and third lines show two independent negative controls in which *GUS* gene is driven by a genomic region that does not contain ABA-related binding sites as indicated in **a**. In the remaining lines, the transcriptional activity is assessed for the two intergenic regions indicated in **a**.

Fig. 6, Regulation of metabolic clusters in plants by lncRNA

a, The proportion of metabolic clusters including lncRNA loci is higher than for other clusters. The co-expressed clusters were predicted in [10] and correspond to co-expressed neighboring genes. The metabolic clusters are co-expressed neighboring genes involved in the biosynthesis of a particular secondary metabolite predicted by plantiSMASH[32].

b, Maximum level of correlation between a lncRNA and any clustered gene calculated in each cluster. The number shown above indicates the p-value of the difference between the two type of clusters determined by Student t-test.

c, The lncRNA *MARS* regulates the expression of the marneral cluster genes through epigenetic reprogramming and chromatin conformation. In control conditions (upper panel) the chromatin of the marneral cluster is enriched in H3K27me3 and LHP1, which results in a condensed and possibly linear chromatin conformation. In response to ABA (bottom left panel) *MARS* over-accumulated transcripts titrate LHP1 away from the cluster. The decrease of LHP1 deposition diminishes H3K27me3 distribution, relaxes the chromatin and as a consequence allows the formation of a chromatin loop that brings together the enhancer element and *MRN1* proximal promoter, leading to transcriptional activation. When *MARS* is repressed, LHP1 recruitment to the cluster is impaired, thus leading to a similar chromatin state: decrease in H3K27me3 mark, chromatin decondensation and increase in chromatin loop conformation. Under this chromatin state, the clustered genes become highly responsive to the ABA treatment.