Rapid endothelial infection, endothelialitis and vascular damage characterise

SARS-CoV-2 infection in a human lung-on-chip model

Vivek V Thacker *[1], Kunal Sharma^[1], Neeraj Dhar ^[1], Gian-Filippo Mancini[2], Jessica

Sordet-Dessimoz[2], John D McKinney[1]

[1]Global Health Institute, Ecole Polytechnique Federale de Lausanne, Switzerland

[2] Histology Core Facility, Ecole Polytechnique Federale de Lausanne, Switzerland

^ equal contribution

*Corresponding author: Vivek V Thacker, vivekvthacker@gmail.com

10
Abstract

Severe cases of COVID-19 present with hypercoagulopathies and systemic

endothelialitis of the lung microvasculature. The dynamics of vascular damage, and

whether it is a direct consequence of endothelial infection or an indirect consequence

of immune cell mediated cytokine storms is unknown. This is in part because *in vitro*

models are typically epithelial cell monocultures or fail to recapitulate vascular

physiology. We use a vascularised lung-on-chip model where, consistent with

monoculture reports, low numbers of SARS-CoV-2 virions are released apically from

alveolar epithelial cells. However, rapid infection of the underlying endothelial layer

leads to the generation of clusters of endothelial cells with low or no CD31 expression,

a progressive loss of endothelial barrier integrity, and a pro-coagulatory

microenvironment. These morphological changes do not occur if these cells are

exposed to the virus apically. Viral RNA persists in individual cells, which generates a

response that is skewed towards NF-KB mediated inflammation, is typified by IL-6

secretion even in the absence of immune cells, and is transient in epithelial cells but

persistent in endothelial cells. Perfusion with Tocilizumab, an inhibitor of trans IL-6

signalling slows the loss of barrier integrity but does not prevent the formation of

endothelial cell clusters with reduced CD31 expression. SARS-CoV-2 mediated

endothelial cell damage occurs despite a lack of rapid viral replication, in a cell-type

specific manner and independently of immune-cell mediated cytokine storms, whose

effect would only exacerbate the damage.

Introduction

Organ-on-chip technologies recreate key aspects of human physiology in a bottom-up and modular manner[1]. In the context of infectious diseases, this allows for studies of cell dynamics[2,3], infection tropism[4], and the role of physiological factors in disease pathogenesis in more native settings[5]. This is particularly relevant for the study of respiratory infectious diseases[6], where the vast surface area of the alveoli poses a challenge to direct experimental observation.

COVID-19, caused by the novel *betacoronavirus* SARS-CoV-2, first manifests as an infection of the upper airways. Severe cases are marked by progression into the lower airways and alveoli. Here, it manifests as an atypical form of acute respiratory distress syndrome (ARDS) characterized by good lung compliance measurements[7,8], and elevated levels of coagulation markers such as D-dimers[9], and pro-inflammatory markers in the blood[10]. Autopsy reports show numerous microvascular thrombi in the lungs of deceased patients together with evidence of the intracellular presence of the virus in vascular cells[11,12]. These reports suggest that infection of and alterations to lung microvasculature plays a key role in COVID-19 pathogenesis[13,14], which is further reinforced by the early observation of endothelial cell damage, platelet aggregation and the formation of microthrombi in an ante-mortem study[15]. Yet most *in vitro* studies have focused on monocultures of upper airway respiratory cells. In studies with alveolar epithelial cells, SARS-CoV-2 has been shown to replicate poorly both in the A549 lung adenocarcinoma cell line[16] and in primary alveolar epithelial cells *ex vivo*[17] and has been reported to be unable to infect primary lung microvascular endothelial cells[18], which are at odds with the reported medical literature. There is therefore an urgent need for a better understanding of the pathogenesis of SARS-CoV-2 in alveolar epithelial cells, especially in a more realistic model of the alveolar space that is vascularized. The

lung-on-chip model is well-suited to this purpose because it includes a vascular compartment maintained under flow, and infection can occur at the air-liquid interface[19], two key physiological features that are lacking in organoid models[20–22]. We therefore establish a human lung-on-chip model for SARS-CoV-2 infections, and probe the viral growth kinetics, cellular localization and responses to a low dose infection using qRT-PCR, ELISA, RNAScope, immunofluorescence and confocal imaging (Fig. S1A).

Results

SARS-CoV-2 infection of a human lung-on-chip alters expression of viral entry factors

A human lung-on-chip (LoC) model for SARS-CoV-2 pathogenesis (Fig. 1A) mimics the alveolar space using primary human alveolar epithelial cells ('epithelial') and lung microvascular endothelial cells ('endothelial'), which form confluent monolayers on the apical and vascular sides of the porous membrane in the chip (Fig. 1B, C). The modular nature of the technology allowed us to recreate otherwise identical LoCs either without ('w/o') (Fig. 1B) or with the addition of CD14+ macrophages (representative image in Fig. S1B) to the epithelial layer on the apical side of the chip. The latter configuration was used for a subset of experiments as indicated throughout the text. We first characterized the epithelial and endothelial cells used, both in monoculture and on-chip at the air-liquid interface to verify that the cells mimicked human alveolar physiology. In epithelial cells, *ACE2* expression was low both in monoculture (Fig S2A) and on-chip (Fig. 1E) consistent with transcriptomic[18,23] and proteomic analyses[24] of human tissue. Expression on-chip was five-fold lower than in monoculture (Fig. S2B). Low *ACE2* expression was also observed from RNAscope assays on uninfected

LoCs (Fig. 1D). Although *ACE2* expression was particularly low in endothelial cells in

monoculture (Fig. S2A, expression increased 10-fold on-chip (Fig. S2B) so that levels

in epithelial and endothelial cells were comparable (Fig. 1E). This upregulation in

ACE2 expression could possibly be due to the effects of shear stress and flow in the

vascular channel[25]. Neuropilin-1, an integrin-binding protein[26] has recently been

reported to be an alternative receptor for SARS-CoV-2 entry[27,28] that is highly

expressed in pulmonary microvascular cells. Consistent with these reports, *NRP1*

expression was between one and four orders of magnitude higher than *ACE2* expression

in both cell types in monoculture (Fig. S2A) and on-chip (Fig. 1E). On-chip, *NRP1*

expression was 6-fold higher in endothelial cells compared to epithelial cells (Fig S2B).

Lastly, mature alveolar epithelial cells retained differentiation in type I and type II cells,

which we verified by immunostaining for type II (pro-SPC) and type I (Podoplanin)

markers in monocultures (Fig. S3A-C) as well as through qRT-PCR measurements for

type II (*ABCA3*, *SFTPC*) and type I (*AQP5*, *PDPN*, *CAV1*) markers in cells from the

epithelial layer of uninfected LoCs (Fig. S3D). Endothelial cells showed characteristic

expression of Platelet Endothelial Cell Adhesion Molecule (PECAM-1 or CD31) at cell

junctions (Fig. 1C, additional images in Fig. 4C, D).

A low dose infection (200-300 plaque forming units (PFUs), multiplicity of infection

(MOI)=0.003) of the apical surface of the LoC resulted in decreases in *ACE2* expression

in epithelial and *NRP1* expression in both epithelial and endothelial cells at 1 dpi (Fig.

1F, I). Expression of both cell surface receptors remained downregulated at 3 dpi (Fig.

1J). These results are consistent with observations using RNAscope assays, where

ACE2 levels in the epithelial layer declined over subsequent days (Fig. 1D, G, H,

p=0.014). On the other hand, infection resulted in increased expression of the

TMPRSS2 protease[29] required to cleave the spike protein for viral entry at 1 dpi in both

cell types relative to uninfected controls (Fig. 1 F, I), with variable expression levels

across chips. This trend was also maintained at 3 dpi (Fig. 1J). Overall, these results

show that both ACE2 and NRP1 likely play a role in pathogenesis in the alveolar space,

aided by increased TMPRSS2 levels. In comparison, infection did not alter levels of

type II or type I alveolar epithelial cell markers significantly (Fig. S3E, F).

Infection of the alveolar space is characterized by a lack of productive infection,

slow intracellular replication, and transmission to the endothelial layer

We first characterized the progression of infection by measuring the release of viral

genomes via qRT-PCR and the number of infectious virions released via measurement

of plaque forming units (PFUs). Infected LoCs were monitored daily for the release of

infected viral progeny (1) *apically* – on the epithelial layer and (2) in the cell culture

media flowed through the vascular channel ('apical wash' and 'vascular effluent' in

Fig. 1A). A low number of viral genomes were released apically from the epithelial

layer, and the number of genomes detected decreased over 1-3 dpi (Fig. 2A). Genome

copy numbers were 100-fold lower than the starting inoculum (between 200-300 PFU).

Low numbers of infectious virions were also detected via measurements of PFU from

eight LoCs w/o macrophages at 1 dpi and two LoCs w/o macrophages at 2 dpi (Fig.

2B). These results showed that, on the whole, productive infection of the epithelial cell

layer did not occur on-chip, although instances of small numbers of epithelial cells with

high levels of spike protein (Fig. S5A) or viral genomic RNA (Fig. S9) were detected

via microscopy and likely contribute to the low levels of infectious virions released.

No viral genomes were detected in the vascular effluent (Fig. 2B), and the lack of

infectious particles in the effluent was confirmed for two LoCs with and w/o

macrophages by plaque forming unit assays (data not shown). This ruled out the

possibility of direct release of virions from the epithelial cell layer into the vascular

channel and suggested that either dissemination of virions to the endothelial cell layer

did not occur, or that infection of the endothelial layer could occur but no viral genomes

or infectious virions were released *apically* into the vascular channel by endothelial

cells.

To address this, we further quantified intracellular viral RNA loads from the total RNA

extracted from cells of the apical and vascular channels of three infected LoCs w/o

macrophages at 1 dpi. This revealed a significant number of viral transcripts in cells

from both the epithelial and endothelial layers (Fig. 2C, Fig. S4C). Additional

quantification of intracellular viral RNA from an infected LoC w/o macrophages using

a SARS-CoV-2 specific one-step qRT-PCR kit revealed >10^4 genomes in both

epithelial and endothelial cells (Fig. S4A); genome copy numbers exceeded those for

cellular housekeeping gene *RNAseP* (Fig. S4B). Similar numbers of intracellular viral

genome copy numbers have been reported for infections of alveolar epithelial cells in

monoculture[17] and these numbers are modest in comparison to cells such as airway

epithelial cells and Vero E6 cells where SARS-CoV-2 replicates productively [30]. The

difference in replication kinetics can also be appreciated from the fact that whereas the

N gene is most abundant during productive infection in Vero E6 cells [31], we obtained a

ratio of SARS-CoV-2 *N* to SARS-CoV-2 *E* <5 (Fig. 2D, Fig. S4D), consistent with the

observation of low numbers of extracellular infection virions (Fig. 2B).

The detection of SARS-CoV-2 genomes in the endothelial layer within 1 dpi suggests

rapid dissemination through *basolateral transfer* from the epithelial to the endothelial

layer can occur. Although basolateral transmission has not been reported previously to

be the significant mode of transmission for both SARS-CoV-2[32] and SARS-CoV-1[33]

infections of monocultures of upper airway cells at the air-liquid interface, this could

reflect aspects of infection unique to alveolar epithelial cells, or more generally, of

infection of cells where productive infection is rare.

SARS-CoV-2 infection of endothelial cells on-chip but not in monoculture alters

cell morphologies

Recently published autopsy reports of COVID-19 patients revealed features of

exudative alveolar damage[12], and a rapid loss of alveolar epithelial cell barrier integrity

could account for the basolateral transfer to endothelial cells. The LoC model is

compatible with microscopy-based assays, which allowed us to assess changes in

cellular physiology with high spatial resolution using confocal microscopy. Even at 3

dpi, the epithelial layer maintained a high degree of confluency (Fig, 3D, Fig. S6F),

with only a few signs of cell death (Fig. 3D, white arrows). In stark contrast, staining

for F-actin localisation revealed striking changes to the morphology of cells in the

vascular channel. In comparison to the confluent layer of cells in an uninfected control

LoC (3D view in Fig 3A, Fig. S6A) infected LoCs showed profound alterations in

endothelial layer architecture (Fig 3B, C). At 2 dpi, cellular clusters characterised by

an increased cell density and stronger nucleic acid staining (Fig 3B, yellow arrows)

coexisted with areas with normal nucleic acid staining levels but reduced cell-to-cell

contact (Fig 3B, white arrows). By 3 dpi, a significant loss of cell confluency was

observed (Fig 3C, Fig S6B-D), with large cell clusters no longer visible. To quantify

these changes across multiple areas of the chip, we enumerated the number of nuclei

per unit area of the membrane surface for infected LoCs at 1 and 3 dpi and uninfected

controls. On the epithelial side, the density of nuclei declines progressively due to

infection irrespective of the presence (p=0.010, p=9E-4) or absence (p=0.054, p=0.029)

of macrophages compared to uninfected controls (Fig. 3E). In contrast, the density of

cellular nuclei increases in the endothelial layer at 1 dpi before decreasing back to levels

in the control samples at 3 dpi (Fig. 3F), consistent with our observation of the transient

formation of endothelial cell clusters. As a further measure of quantification, we

measured the proportion of the surface area of the endothelial cell layer with low or no

actin staining using a range of intensity thresholds (Fig. S2E). At 3 dpi, a much larger

proportion of the surface area of the endothelial layer of infected LoCs had low or no

actin staining compared to an uninfected control for all intensity thresholds (Fig. S2E).

Immunostaining for the viral S protein at 3 dpi (Fig 3G for endothelial and Fig 3H for

epithelial layers, additional examples in Fig. S6B-D, S6F and controls with secondary

antibody only in Fig. S8D-G) show clear evidence of individual infected epithelial (Fig.

3I) and endothelial cells (Fig. 3J). In some cases, S proteins appear to co-localize with

the periphery of the nucleus (Fig. 3I, 3J, white arrows). The transfer of viral proteins

from the apical to vascular channel and difference in confluency between the epithelial

and endothelial layers at 3 dpi can be better appreciated from a 3D view of the two

channels together with the connecting pores in the PDMS membrane; S protein can be

clearly localised to the interface of the two cell types in the pores of the membrane

(yellow arrows, Fig, 3K). Epithelial and endothelial cells exhibit very different

physiological responses to low levels of infection, and vascular damage by 3 dpi was a

consistent observation across all infected LoCs (n=12) maintained until 3 dpi in this

study

We next sought to understand if these responses were specific to the lung-on-chip

physiology and co-culture, and so examined infections of epithelial and endothelial cell

monoculture. At 1 dpi, heavily infected epithelial cells in monoculture with 1000 PFU

showed similar levels and intracellular distributions of spike protein (Fig. S5B). In stark

contrast, monoculture infections of endothelial cells with viral innocula up to 100-fold

greater than the apical inoculum used in the LoC experiments (Fig. 2A) showed few

foci of viral S proteins. Endothelial cells retained their morphology and confluency

(Fig. S7) up to 2 dpi, in agreement with recent reports of the inability of apical

inoculations of SARS-CoV-2 to infect lung microvascular endothelial cells [18]. This

suggests that the site of contact of SARS-CoV-2 with endothelial cells (apical vs

basolateral) may play an important role – a physiological factor that can be recapitulated

in the LoC but not in simpler models.

SARS-CoV-2 infected endothelial cells lose expression of tight junction markers

and adopt a pro-coagulatory phenotype

Actin staining definitively demonstrated the vascular damage caused as a result of loss

of cell confluence, we therefore probed cells from both layers in infected and uninfected

LoCs for changes in expression of tight junction proteins that might have caused these

changes Maximum intensity projections from the epithelial cell layer of an uninfected

control showed strong expression of both actin (Fig. 4A) and TJP1 at the cell junctions

(Fig. 4B). Similarly, images from a confluent endothelial cell layer in an uninfected

control LoC showed actin staining correlated with high expression of Zonula Occludens

- 1 (ZO-1 or TJP1) (Fig. 4C) as well as high expression of the endothelial cell specific

marker CD31 at cell junctions (Fig. 4D). These results were corroborated by qRT-PCR

measurement of high levels of expression of *TJP1* and *CD31* as well as that of VE-

cadherin (*CDH5*) in uninfected controls (Fig. S8A).

Consistent with Fig. 3, changes to the expression pattern of TJP1 in the epithelial layer

at 1 dpi were subtle, TJP1 expression at cell junctions continued to be observed across

much of the epithelial cell layer (Fig. 4E, additional images in Fig. S8B, C). SARS-CoV-2 infection, on the other hand, profoundly altered CD31 expression and distribution in the endothelial layer. 3D views of regions of interest in the endothelial layer of an infected LoC showed this to be a dynamic process that resulted in two significant changes: first, an increase in vascular permeability from the loss of tight junctions between CD31+ cells (white arrows in Fig. 4F, G), and second, the presence of cells that showed little or no CD31 expression (Fig. 4F-H). Cells displaying the 'CD31- phenotype' typically had higher levels of spike protein (shown in amber in Fig. 4F-G), and an increased local cell density and nuclear labelling consistent with the endothelial cell clusters identified in Figure 3. This link is strengthened by examination of zooms of two endothelial cell clusters (Fig. 4I, J, and Fig. 4K, L) from another LoC at 1 dpi. This confirmed that cells in both clusters were CD31- (Fig. 4I, Fig. 4K) but nevertheless express TJP1 and showed strong and altered actin distributions (Fig. 4J, Fig. 4L), morphological features consistent with the clusters identified in independent experiments in Fig. 3. These changes are further corroborated by qRT-PCR measurements which show a progressive and sharp reduction in overall levels of *CD31* and *CDH5* expression in the endothelial layer at 1 dpi and 3 dpi respectively compared to uninfected controls (Fig. 4M). In contrast, the reduction in *TJP1* expression at 1 dpi is less pronounced and expression levels do not decline further at 3 dpi.

These dramatic changes in expression of tight junction proteins also resulted in series of complex alterations to the expression of endothelial cell proteins with direct roles in coagulatory pathways (Fig. 4N). For example, infection did not alter expression of *pro-coagulatory* Tissue Factor (*F3*) released by endothelial cells as part of the extrinsic coagulation pathway[34], but induced a progressive decrease in the expression of *anti-coagulatory* Tissue Factor Pathway Inhibitor (*TFPI*) [35]. A more modest but similarly

consistent decrease was also observed for the expression of Thrombomodulin (*THBD*)
that also has an anti-coagulatory function. Although infection also decreased the
expression of von Willebrand Factor (*VWF*), elevated levels of VWF protein were
observed on the surface of the CD31- endothelial cells at 1 dpi via immunofluorescence
(bright pink, Fig. 4F-G). In contrast, expression of plasminogen inhibitor activation
(PAI-1 expressed by *SERPINE1*) decreased only by 3 dpi. Taken together with the loss
of CD31 expression, these results point to the generation of a pro-coagulatory
environment in the vascular channel of infected chips.

SARS-COV-2 persists in individual epithelial and endothelial cells in the alveolar
space

Our results thus far showed that infection on-chip itself lead to heterogenous outcomes
that were strongly cell-type specific. We therefore performed a series of RNAscope
assays with confocal microscopy to probe the spatial distribution and cell-to-cell
variability in gene expression at the single-cell level to complement observations from
qRT-PCR measurements. We began by examining levels of viral RNA and antisense
RNA generated during viral replication *in situ* using probes for *S* RNA and *ORF1AB*
antisense RNA respectively. Representative images for 232 x 232 μm^2 fields of view
from the apical and vascular channels of chips at 1 and 3 dpi are shown in Fig 5A-H.
At 1 dpi, a few examples of heavily infected cells are visible on the epithelial layer (Fig.
S9A-C) and viral genomic RNA can be detected throughout the cytoplasm. Consistent
with the observations in Fig. S5, such cells are likely to be the source of infectious
virions released in the apical wash in Fig. 2A, B. However, in the vast majority of cells
on the epithelial layer, the virus is detected only at low levels at 1 dpi (Fig. 5A), and
infection does not appear to be limited by (Fig. 5C, white arrows) or even correlated to

cellular *ACE2* expression at the time of fixation (Fig. 5D, yellow arrows). Both genomic

and antisense RNA are detected in endothelial cells at 1 dpi (Fig. 5B), indicating that

intracellular viral replication can also occur in these cells. Heavily infected endothelial

cells were not observed, in agreement with the lack of infectious virions in the vascular

effluent (Fig. 2C). Endothelial cell clusters are clearly identifiable by their

morphological features, and zoomed-in images (Fig 5E, F) show that viral infection of

endothelial cells with normal and clustered morphology. As with the epithelial layer,

instances of localization of viral RNA with the nucleus is also observed (Fig. 5F, white

arrowheads). At 3 dpi, infection in both layers (Fig 5G, 5H) and the loss of endothelial

layer integrity is evident (compare Fig. 5G vs. Fig. 5H). We did not detect heavily

infected epithelial cells at 3 dpi, in agreement with the diminishing viral titre (Fig. 2A).

We normalized the number of RNA dots detected either on a per-cell basis (Fig. 5I-K)

or per-field-of-view basis (Fig. S10), to enable quantification of differences between

the two layers while accounting for differences in cell numbers and cell death. The

median values of intensities were similar across all conditions (Fig. S11). In the

epithelial layer, viral infection and replication per cell were either similar (Fig. 5J, L)

or lower by 3 dpi (Fig. 5I, p=0.018; Fig. 5K, p=0.006). In contrast, viral RNA

accumulates in individual endothelial cells (Fig. 5K, Fig. 5L, p=0.004). The presence

of macrophages generally leads to lower levels of viral RNA replication (Fig. 5I, J) but

does not prevent the accumulation of viral genomes intracellularly in endothelial cells

(Fig. 5L). At 3 dpi, *ACE2* levels are higher on a per-cell basis in endothelial vs epithelial

cells (Fig. 5M, p=0.028), due to a reduction in *ACE2* levels on the epithelial layer.

Normalization on a per-field-of-view basis shows predominantly a *declining* trend for

levels of replication (Fig. S10A, B) and intracellular viral RNA (Fig. S10C, D) in both

cell types, as well as a *decline* in *ACE2* expression (Fig. S10E), consistent with qRT-

PCR measurements (Fig. 1J, Fig. S10F). These results highlight the cumulative effect

of differential cell numbers and cell death in these two layers (Fig. 3E, F). Overall, virus

levels are low and decline in both layers by 3 dpi, yet the virus persists and continues

to replicate within individual infected cells, in agreement with reports of clinical

infection.

Persistent NF-KB inflammatory responses to SARS-CoV-2 infection in

endothelial but not epithelial cells

Concurrently, we investigated host responses to the infection. Expression levels of NF-

KB related pro-inflammatory genes (*TNFA*, *IL6*, *IL1B*) at 1 dpi was an order of

magnitude higher than that of interferon genes in both cell types, with little to no

interferon stimulatory gene expression detected in both cell types (Fig. 6A). Responses

were also cell-type specific; endothelial cells from infected LoCs showed upregulation

of *TNFA*, *IL6* and *IFNB*, *IFNL1* and *IFNL3* genes whereas only expression of *IFNB*

and *IFNL1* in epithelial cells were higher than uninfected controls (Fig. 6B). In both

cell types, expression of *IP10* and *ISG15* were lower than uninfected controls,

consistent with the action of a number of SARS-COV-2 proteins in inhibiting type I

interferon signalling [36,37].

The LoC platform also allows the simultaneous detection of secreted cytokines in the

vascular effluent from the same chips. ELISA assays for IL-1B and IP10 did not detect

these cytokines in the effluent (data not shown), the lack of IL-1B secretion despite

high expression (Fig. 6B), suggests that it is not exported[38]. IL-6, on the other hand,

was consistently detected up to 3 dpi in the vascular effluent from LoCs both in the

presence or absence of macrophages (Fig. 6C). No IL-6 was detected in the vascular

effluent from uninfected controls. IL-6 levels were not significantly different between these two categories of LoCs, which strongly suggested a non-immune cell source for this cytokine (Fig. 6C).

Given the pleiotropic nature of IL-6[39], we also examined expression of the IL6R receptor and the metallopeptidase ADAM17 which sheds the TNF-alpha receptor and the IL-6R receptor[40] in both the epithelial and endothelial layer at 1 dpi (Fig. 6D). Expression of *ADAM17*, which has been shown to enhance vascular permeability[41], was on average 7-fold higher in the endothelial layer at 1 dpi (Fig. 6E), suggestive of the activation of the IL-6 signalling pathway.

COVID-19 patients with severe infection have elevated IL-6 levels, and we therefore sought to examine the cell-type specific *IL6* expression at the single cell resolution. We therefore used RNAscope to probe the spatial distribution of *IL6* expression in LoCs reconstituted with macrophages. Consistent with the qRT-PCR data (Fig. S12), *IL6* expression in the control LoC is higher in the epithelial layer (Fig. S13A) than the endothelial layer (Fig. S13B), and immunostaining with an anti-CD45 antibody confirmed that the *IL6* expression in the epithelial layer of the chip did not co-localise with macrophages (Fig. S13A, B). At 1 dpi, individual cells with higher levels of *IL6* expression were observed in both epithelial and endothelial layers (Fig. 6F, G), but only differences in expression in the endothelia layer were statistically significant (Fig. 6J). Cells with high *IL6* expression were also unlikely to be macrophages (Fig. S13 C-E).

By 3 dpi *IL6* expression in the epithelial layer had returned to the control levels (Fig. 6H, J, S14A) whereas expression increased dramatically and was more widespread in the endothelial layer (Fig. 6I, J, S14A, S14B). Expression did not appear to correlate with level of infection – zooms of cells with comparable levels of infection in Fig. 6I with low (Fig. 6K), medium (Fig. 6L) and high (Fig. 6 M, N) *IL6* expression are shown.

Expression in the endothelial layer at 3 dpi was higher than that in the endothelial layer at 1 dpi, the epithelial layer at 3 dpi, and the uninfected controls, both on a per-cell and per-field-of-view basis (Fig. 6J, Fig. S14A, p=0.004, p=0.014, p=0.008 respectively). Unlike viral RNA levels (Fig. S10B, D) *IL6* expression in the endothelial layer *does not diminish* over 1-3 dpi and would therefore appear to be the major contributor to IL-6 secretion in the vascular effluent as infection progresses.

Inhibition of trans IL-6 signalling reduces but does not eliminate endothelial cell inflammation

The activation of IL-6 signalling observed suggested *trans* IL-6 signalling as a potential target to ameliorate the vascular inflammation observed. A similar rationale has led to numerous clinical trials of the anti-IL-6R monoclonal antibody Tocilizumab[42] as a repurposed therapeutic for COVID-19. Tocilizumab administration at 56 μg/mL via continuous perfusion over 2 days post infection *per se* did not abrogate IL-6 secretion, consistent with its mode of action (Fig. 6C). We therefore undertook an investigation of cell morphologies in the endothelial layer of a treated and untreated LoC via actin staining. This revealed that whereas endothelial cell confluency was, on the whole, better retained in the treated chip (Fig. 7A, zooms in Fig. 7B, untreated control in Fig. 7D, zoom in Fig. 7E) Tocilizumab administration *did not prevent* the occurrence of endothelial cell clusters (Fig. 7A, zooms in Fig. 7C, untreated control in Fig. 7D, zoom in Fig. 7F). Consistent with these results, immunostaining for CD31 in another treated LoC showed an improvement in the integrity of the endothelial cell layer with tight junctions observable at 2 dpi (3D views in Fig. 7G, H) yet endothelial cell clusters with reduced CD31 expression and strong actin staining were nevertheless formed (3D views in Fig. 7G, H, indicated by yellow arrows).

To quantify the improvement in permeability, we compared regions of interest (ROIs)

that excluded areas of cellular hyperplasia across at least six fields of view from the

endothelial layer of LoCs with and without Tocilizumab perfusion, and identified areas

with low or no F-actin staining as those with reduced confluence, as in Fig. S6E. A plot

of the proportion of pixels with intensities below a defined cut-off threshold (Fig. 7I)

showed that the untreated LoC did have a significantly higher proportion of pixels with

intensities lower than 5% (p=0.032) of the maximum intensity (p=0.022 for 3% and

p=0.038 for 2%). Inhibition of IL-6 signalling through Tocilizumab is therefore able to

ameliorate *some* but *not all* of the vascular damage observed.

Discussion

The alveolar space has a strikingly different physiology from that of the upper airway.

ACE2 is the canonical entry receptor used by SARS-CoV-2, yet is scarce in the alveoli.

It has been speculated that high levels of *ACE2* expression may be necessary for export

of mature virions [43] and because most virology assays measure viral titre, studies with

alveolar epithelial cell lines such as A549 have focused on increasing *ACE2* expression

through transfection. In contrast, alternate receptors such as NRP1 more abundant in

the lower airways have been understudied, and the effects of uptake via NRP1 vs. ACE2

on the viral life cycle is relatively unknown. Here, we populate our model with relevant

primary human cells and report low *ACE2* expression and high *NRP1* expression,

consistent with human alveolar physiology. Similarly, most infections of alveolar

epithelial cells are usually performed at high MOI[16]. Instead, we used an infectious dose

that better mimics either a direct inspiration of aerosols or the transfer of infectious

virions from the upper respiratory tract. These features, together with the ability to co-

culture epithelial cells (at an air-liquid interface) and endothelial cells (under flow)

enable the lung-on-chip model to be used to study aspects of early infection that cannot

be achieved with other systems.

Despite low ACE2 expression and low MOI, SARS-CoV-2 infection produced

pronounced physiological changes on-chip. This could be due to the high levels of

NRP1 and aided by upregulation of *TMPRSS2*, which is known to be important in

human SARS-CoV-2 infection. In contrast to models of productive replication, we

observe slow intracellular viral replication without significant release of infectious

virions, observations that are consistent with reports of recovery of viral RNA in

patients after they cease to be infectious[44]. Microscopy-based analyses also revealed

that responses to infection are highly heterogenous; for example, the few foci of heavily

infected cells might represent cells with constitutively higher levels of *ACE2* expression

at the time of infection. However, why alveolar epithelial cells show persistent but not

productive infection is still unclear.

Infection of the epithelial layer rapidly led to an infection of the endothelial layer.

Basolateral transmission cannot occur in typical monoculture experiments, but it has

not been reported to occur to a large extent in monocultures of airway epithelial cells

at the air-liquid interface. However, the basement membrane is much thinner in alveoli

than in the upper airway[45], with cell-to-cell contact between the epithelial and

endothelial cell layers that is recreated in the array of pores on-chip. Infection of lung

microvascular cells therefore might occur through uptake of vesicles as part of cell-to-

cell communication between the layers (a receptor-independent mechanism of uptake).

These observations are strengthened by the lack of observation of any signs of infection

in lung microvascular endothelial cells when directly exposed to the virus on their

apical surface. Alternatively, infection of endothelial cells when exposed to the virus

basolaterally might still occur due to receptor-mediated uptake, with differences

between apical and basolateral infection explained by differences in distribution of cell

surface receptors. Notably, NRP1 is likely to be located along the basal surface of the

endothelial cells as an integrin-binding protein[26], and SARS-CoV-2 has been shown to

infect vascularised organoids, albeit at high MOI[46]. The exact mechanisms remain to

be elucidated, but the consequences of this may extend downstream to the persistent

infection and endothelial-cell specific cell damage observed.

A striking feature of SARS-CoV-2 infections in the LoC model was the cell-type

specific morphological changes. Infection altered endothelial cell morphology in two

different but interconnected ways. First, the loss of tight junctions between cells is

indicative of increased vascular permeability that could lead to exudative alveolar

damage observed in autopsy reports. Exposed sub-endothelial collagen would likely

trigger the contact activation or intrinsic pathway of coagulation. This was also

accompanied by pro-coagulatory changes in the extrinsic pathway through a reduction

in expression of anti-coagulatory factors such as TFPI and thrombomodulin.

Interestingly, infection of Human Umbilical Vein Endothelial Cells (HUVECs) with

Dengue virus has been shown to have the opposite effect [47], with a strong increase in

thrombomodulin expression consistent with the disease manifestation as a

haemorrhagic fever. These results highlight the close link between coagulation and

inflammation in this cell type. Second, a subpopulation of endothelial cells completely

lost CD31 expression and adopted an altered morphology consisting of cell clusters.

These clusters could form either through enhanced proliferation or from cell migration.

The former is more likely, as a consistent increase in cell density was observed across

multiple LoCs, but both possibilities are consistent with the role of NRP1 as a growth

factor receptor or its role in VEGF signalling[48]. The loss of CD31 expression could

itself have a number of possible implications: CD31 itself plays important roles in

leukocyte migration[49], angiogenesis, in maintaining a anti-coagulatory environment[50,51]

as well as preventing endothelial cell death from immune cell mediated inflammation[52].

These CD31- cells could further contribute to a pro-coagulatory environment in the

microvasculature and lead to changes in vascular architecture reported in autopsies[11].

A loss of CD31 expression in endothelial cells will likely alter interactions with

neutrophils and platelets that who also express CD31, potentially contributing to the

formation of neutrophil extracellular traps and platelet hyperactivity reported in

COVID-19[53,54]. These findings are also consistent with the observation of elevated

serum levels of CD31 in patients with severe COVID-19[55]. The exact mechanisms for

the loss of CD31 and other tight junction markers remains to be elucidated, but a

possible mechanism could be through action of viral proteins themselves, as is

observed in the case of the K5 protein of the Kaposi Sarcoma herpesvirus[56] that causes

Kaposi Sarcoma.

Increased NF-KB inflammatory responses could also account for the vascular changes

observed, and the role of cytokine storms in COVID-19 pathogenesis is still unclear.

We observed that low levels of intracellular viral RNA generates an NF-KB mediated

pro-inflammatory response, with suppression of antiviral interferon responses,

consistent with reports of the effect of a number of viral proteins[57,58]. Within each cell

types, there were significant cell-to-cell differences in expression of inflammatory

markers, clearly identifiable with single-cell spatial resolution through the use of

RNAscope and confocal imaging in the LoC model. These differences could be due to

differences in the mode of uptake, or differences in the intracellular localization of viral

RNA and RNA replication, or differences in the levels of translation of viral proteins.

We observe instances of co-localisation of viral RNA and S protein with the nucleus,

which is consistent with bioinformatic predictions using algorithms for RNA

localisation[59], but has not been reported in studies using cell lines with productive

infections. These atypical sub-cellular localisations may be characteristic of persistent

infection and merit further investigation. Yet, once again the responses of epithelial and

endothelial cells contrasted sharply – inflammation in epithelial cells was transient,

whereas it was persistent in endothelial cells.

Lastly, given the persistent *IL6* expression in endothelial cells, we applied the model

and our microscopy-based approach to examine if Tocilizumab perfusion could the

occurrence of either increased vascular permeability or the formation of endothelial cell

clusters or both. We found that Tocilizumab *was* able to maintain cellular confluency

but did not prevent a subset of endothelial cells from over-proliferating and acquiring

an inflammatory phenotype. These observations at the single cell level offer insights

into reasons why outcomes with this drug in clinical trials have been modest and at

times contradictory[60] and there is insufficient evidence of the efficacy of Tocilizumab

in reducing mortality. The lung-on-chip model is therefore able to make clinically

testable predictions, highlighting the utility of this approach.

In sum, SARS-CoV-2 infection of a model of the alveolar space shows a set of unique

characteristics underlined by persistent infection, low viral replication, inflammatory

response, damage to endothelial barrier integrity, and the formation of a pro-

coagulatory environment that are in good agreement with reports of clinical disease.

The superior physiological mimicry LoC model (air-liquid interface, co-culture,

perfusion through a vascular channel) and high-resolution confocal imaging reveals

insights into the dynamics and mechanisms that underlie these observations that are

difficult to obtain in other experimental models and that will improve our understanding

of the pathogenesis of this multi-organ disease.

Acknowledgements

We gratefully acknowledge assistance from Dr Muhammet Fatih Gulen for the ELISA

assays. We are also grateful to the members of the BioImaging Core Facility (BIOP)

for assistance with confocal microscopy and to Prof Dr Thomas Hugle at the Centre

Hospitalier Universitaire Vaudois (CHUV(for donation of Tocilizumab.

Methods

<u>Cell culture</u>

Primary human alveolar epithelial cells (epithelial cells) and human lung microvascular

endothelial cells (endothelial cells) were obtained from a commercial supplier (Cell

Biologics, USA). Each vial of epithelial cells comprised a mix of Type I and Type II

cells at passage 3, and expression of cell-type specific markers (*AQP5, PDPN,* and

CAV1 for type I and *ABCA3* and *SFTPC* for type II) was verified via qRT-PCR on-

chip (Fig. S3D). Both cell types were cultured *in vitro* in complete medium comprising

base medium and supplements (Cell Biologics, USA) in 5% CO_2 at 37 °C. Passage of

epithelial cells lowered *ACE2* expression (Fig. S2A), therefore all chips were

reconstituted with epithelial cells seeded directly on the lung-on-chip (see below),

without any additional *in vitro* culture. In contrast, we observed no significant changes

in viral entry factor expression in endothelial cells upon passage, and so therefore these

cells were passaged between 3-5 times before seeding in the LoC devices.

<u>PBMC isolation and macrophage differentiation</u>

Peripheral blood mononuclear cells were obtained from buffy coat (Interregional Blood Transfusion SRC Ltd, Switzerland) obtained from anonymised donors. PBMCs were isolated using a Biocol Separation procedure as per the manufacturer's instructions. Isolated PBMCs were subsequently cryopreserved in a solution of 70% heat inactivated Fetal Bovine Serum (FBS, Gibco), 20% RPMI Medium (Gibco) and 10% dimethlysulfoxide (DMSO). One week prior to seeding the macrophages in the LoC devices, a cryopreserved aliquot was thawed and cultured in a T-75 flask (TPP, Switzerland) in RPMI supplemented with 10% FBS. CD14+ monocytes were isolated using positive selection (CD14 ultrapure isolation kit, Miltenyi Biosciences) and cultured in RPMI medium supplemented with 10% FBS and differentiated for 7 days with 20 ng/mL recombinant human Macrophage-Colony Stimulating Factor protein (M-CSF) (Thermo Fisher Scientific), and 100U/L of penicillin-streptomycin solution (Thermo Fisher Scientific) to avoid bacterial contamination. The cells were cultured in plastic petri dishes without pre-sterilisation (Greiner Bio-One) so that differentiated macrophages could be more easily detached.

Quantitative Real-Time PCR (qRT-PCR) for cell characterization

Epithelial cells obtained directly from the supplier were grown overnight in cell-culture microdishes (Ibidi) or T-25 cell culture flask (TPP, Switzerland). Passaged epithelial cells were grown to confluency in a T-75 cell culture flask (TPP). Growth media was removed from the flask, and the cells were incubated with the appropriate volume of RNA lysis buffer (RNAeasy Plus Micro Kit, Qiagen) and RNA isolated as per the manufacturer's instructions and resuspended in 25 µL of DEPC-treated water. Approximately 1 µg of RNA was then used to generate cDNA using the SuperScript®IV First-Strand Synthesis System with random hexamers (Invitrogen), which was stored at -20 °C. Specific primers for used for the qRT-PCR are listed in Table S1. qRT-PCR reactions were prepared with SYBR®Green PCR Master Mix (Applied Biosystems) with 500 nM primers, and 1 µL cDNA. Reactions were run as absolute quantification on ABI PRISM®7900HT Sequence Detection System (Applied Biosystems). Amplicon specificity was confirmed by melting-curve analysis.

Quantitative Real-Time PCR (qRT-PCR) for quantification of viral genomes in the supernatant

150 μL of cell-free supernatant was processed using a kit to extract viral RNA as per the manufacturer's instructions (NucleoSpin Dx Virus, Machery Nagel). For the vascular effluent, this was taken from the approximately 1.5 mL of effluent generated daily by the flow through the vascular channel. In the case of the apical wash, the 30 μL collected from the apical channel was further diluted in cell culture media up to 150 μL and subsequently processed. Viral RNA was eluted in 50mL of DEPC-treated water. 5 mL of RNA was then used in a one-step qRT-PCR kit for SARS-CoV-2 detection (2019-nCoV TaqMan RT-PCR kit, Norgen Biosciences) using primer and probe mixes corresponding to the nucleocapsid (N2) gene and the host RNAseP genes. Primer sequences are those listed by the US Centre for Disease Control (CDC).

Generation of SARS-CoV-2 stocks

VeroE6 cells and a clinical isolate of SARS-CoV-2 were a kind gift from the lab of Prof Carolyn Tapparel at the University of Geneva. SARS-CoV2/Switzerland/GE9586/2020 was isolated from a clinical specimen in the University Hospital in Geneva in Vero-E6. Cells were infected and supernatant was collected 3 days post infection, clarified, aliquoted and frozen at -80 °C and subsequently titrated by plaque assay in Vero-E6. Virus used for all experiments in this manuscript was at passage 3 in Vero E6 cells.

Human lung-on-chip (LoC) model

LoC made of polydimethylsiloxane (PDMS) were obtained from Emulate (Boston, USA). Extracellular matrix (ECM) coating was performed as per the manufacturer's instructions. Chips were activated using ER-1 solution (Emulate) dissolved in ER-2 solution at 0.5 mg/mL (Emulate) and exposed for 20 minutes under UV light. The chip was then rinsed with coating solution and exposed again to UV light for a further 20 minutes. Chips were then washed thoroughly with PBS before incubating with an ECM solution of 150 μg/mL bovine collagen type I (AteloCell, Japan) and 30 μg/mL fibronectin from human plasma (Sigma-Aldrich) in PBS buffered with 15 mM HEPES solution (Gibco) for 1-2 hours at 37°C. If not used directly, coated chips were stored at 4°C and pre-activated before use by incubation for 30 minutes with the same ECM solution at 37°C. Endothelial cells were cultured overnight at 37°C and 5% CO_2 in T-75 cell culture flasks, detached with 0.05% Trypsin, concentrated to 5-10 million cells/mL, and seeded on the bottom face of the PDMS membrane. The chip was then

incubated for a short period at 37 °C to allow the endothelial cells to spread and

subsequently seeded with epithelial cells. Epithelial cells were seeded directly from

cryopreserved vials received from the supplier, each vial of 0.5 million cells could be

used to seed two chips. To increase reproducibility, vials were pooled and used to seed

multiple chips. The chip was incubated overnight with complete epithelial and

endothelial media in the epithelial and endothelial channels, respectively, under static

conditions. The next day, the chip was washed and a reduced medium for the air-liquid

interface (ALI) was flowed through the vascular channel using syringe pumps

(Aladdin-220, Word Precision Instruments) at 60 μL/hour as described [61]. The

composition of the ALI media used was as described in [61] but with an FBS

concentration of 5%. The epithelial face was incubated with epithelial base medium

with 1 μM dexamethasone (Sigma Aldrich) without FBS supplementation to promote

tight junction formation and surfactant expression as reported in previous lung-on-chip

studies[61,62]. Flow was maintained over 2-3 days with daily replacement of the medium

on the epithelial face (with dexamethasone supplementation). At the end of this period,

in LoCs reconstituted with macrophages, CD14+ macrophages differentiated for 7 days

in M-CSF (described above) were detached from the petri dish using 2 mM

ethylenediaminetetraacetic acid (EDTA, Sigma Aldrich) in PBS at 4 °C as well as

mechanical scraping, centrifuged at 300 g for 5 minutes, and resuspended in a small

volume of epithelial cell media without dexamethasone. This solution containing

macrophages was introduced onto the epithelial face and incubated for 30 minutes at

37 °C and 5% CO_2 to allow macrophages to attach to the epithelial cells. Medium on

the epithelial face was then removed and the chip was maintained overnight at an air-

liquid interface (ALI). Chips were controlled to ensure that they successfully

maintained the ALI overnight and were then transferred to the biosafety level 3 (BSL-

3) facility for SARS-COV-2 infection. No antibiotics were used in any of the cell

culture media for setting up the LoC model.

<u>Tocilizumab perfusion in the LoC model</u>

Tocilizumab/Rho-Actemra (Roche, 20 mg/mL in saline) for intravenous administration

were a kind gift from Prof Thomas Hugle at the University Hospital in Lausanne,

Switzerland (CHUV). We simulated a single dose of 8 mg/kg given to a person of

bodyweight 70 kg and a blood volume of 5 litres to obtain a concentration of 56 μg/mL

in the cell culture media. Tocilizumab perfusion was started immediately after infection

and was flowed through the vascular channel of infected chips for a period of 2 days

post infection.

<u>Characterization of infection in LoCs</u>

Uninfected LoCs were maintained at an ALI for either 1 day (for total RNA extraction

(two LoCs without macrophages) or for up to 3 days for RNAscope characterization

(one LoC each were reconstituted with and without macrophages) after addition of

macrophages, during which time ALI medium was flowed through the endothelial

channel at 60 µL/hour. In the case of the former, total RNA from the epithelial and

endothelial layers was extracted separately by passing 350 µL of the lysis buffer

reconstituted as per the manufacturer's instructions (RNeasy Plus Micro Kit, Qiagen)

through easy layer. Complete detachment of the cells in each layer was verified via

optical microscopy. RNA was eluted in 25 µL of DEPC-treated nuclease free water and

used for subsequent qRT-PCR reactions.

In the case of LoCs processed for RNAscope, the chips were fixed for 1 hour using a

fresh solution of 4% paraformaldehyde in phosphate buffered saline (PBS), and

subsequently dehydrated in a sequence of washes with 50% EtOH, 70% EtOH, and 100

% EtOH. The chips were then stored in 100% EtOH at -20 ℃ until processed for

RNAscope at the Histology Cor Facility at EPFL.

<u>Immunofluorescence characterization</u>

For immunofluorescence labelling, the fixed LoCs were incubated with a blocking

solution of 2% Bovine Serum Albumin (BSA) in PBS ('blocking buffer'), followed by

overnight incubation with the primary antibody in the blocking buffer at 4℃. A list of

primary antibodies and concentrations used is included in Table S2. In addition, the

infected LoC shown in Fig. 4 E-G was immunostained with Rabbit anti-SARS-CoV

spike antisera [63], which was a kind gift of Prof Carolyn Machamer at John Hopkins

University. The following day the chip was washed extensively with fresh blocking

buffer before inoculation with the secondary antibody at a concentration of 1:300 for 1

hour at room temperature. The secondary antibodies used were: Donkey anti-mouse

Alexa Fluor 488 (A21206, Thermo Fisher), or Donkey anti-mouse Alexa Fluor 568

(A10037, Thermo Fisher), or Donkey anti-mouse Alexa Fluor 647 (A31573, Thermo

Fisher), and were chosen to complement the fluorophores already assigned to

RNAscope labelling. F-actin on both the epithelial and endothelial face was stained

using Sir-Actin dye in the far-red (Spherochrome) at 1 μM for 30 minutes concurrently

with Hoechst staining, as described above. Confocal images were obtained on a Leica

SP8 microscope in the inverted optical configuration at the EPFL BIOP core facility.

<u>Infection of the LoC with SARS-CoV-2</u>

LoCs for infection were assembled into a sealed Tupperware box modified to contain

inlets and outlets for the flow of medium through the vascular channel and was

maintained throughout the course of the experiment by use of a syringe pump. On the

day of the experiment, an aliquot of virus-containing supernatant was thawed from the

stock at -80 °C, and diluted approximately 35-fold in epithelial cell media without FBS

to generate the inoculum that corresponded to an infectious dose of 200 – 300 plaque

forming units (PFU) in a volume of 30 μL. 30 μL of this solution was then added to the

apical channel of each LoC, and the LoC was incubated for an hour at 37°C and 5%

CO_2 to enable infection. Thereafter, the medium in the apical channel was removed.

The LoC was returned to air-liquid interface and the inlets of the infected chip were

sealed with solid pins as a safety precaution. The media flowing through the vascular

channel of each LoC was collected in a separate, sealed 15 mL Falcon tube, and each

tube was emptied daily, and the effluent stored at -80 °C for further processing. At this

time, the metal pins sealing the apical channel of each chip were removed, and 30 μL

of epithelial cell media without FBS was briefly added to the channel before being

withdrawn and collected as the 'apical wash'. The metal pins were reinserted, and the

LoCs returned to the cell culture incubator. Both the vascular effluent and the apical

wash from each chip were collected for each day post infection.

Infection was terminated at specified time points either by treatment with the lysis

buffer for RNA extraction (one LoC reconstituted without macrophages) or by fixing

with freshly prepared 4% paraformaldehyde for a period of 1 hour (all other infected

LoCs) for RNAscope assay preparation. The chips were then dehydrated as described

above and immersed in 100 % EtOH before removal from the BSL-3 facility.

<u>RNAscope Assay</u>

RNAscope Multiplex Fluorescent V2 assay (Bio-techne, Cat. No. 323110) was
performed according to manufacturer's protocol directly on chips. They were
hybridized with the control Hs 3plex positive control (Bio-techne, Cat. No. 320861) or
a combination of the following probes V-nCoV2019-orf1ab-sense (Bio-techne, Cat.
859151), Hs-ACE2-C2 (Bio-techne, Cat. 848151-C2), V-nCoV2019-S-C3 (Bio-
techne, Cat. 848561-C3), HS-Il1b-C1 (Bio-techne, Cat. 310361), and Hs-Il6-C2 (Bio-
techne, Cat. No. 310371-C2) simultaneously at 40°C for 2 hours. The different channels
were revealed with TSA Opal570 (Akoya Biosciences, Cat. No. FP1488001KT), TSA
Opal650 (Akoya Biosciences, Cat. No. FP1488001KT) and TSA Opal690 (Akoya
Biosciences, Cat. No. FP1497001KT). Cells were counterstained with DAPI and
mounted with Prolong Gold Antifade Mountant (Thermo Fisher, P36930).

<u>ELISA assays</u>
Frozen vascular effluent from infected chips was thawed, incubated at 58 °C for 30
minutes to ensure sterilization, and then removed from the BSL-3 for further
processing. Levels of human IL-6 were determined by enzyme-linked immunosorbent
assays (BD Biosciences) as per the manufacturers' instructions.

<u>Confocal imaging and Image Analysis</u>
Infected and control LoCs were imaged using a Leica SP8 confocal microscope with a
white light laser. For chips processed via RNAscope and labelled with Opal 650 and
Opal 690, the excitation and emission windows were carefully chosen to minimise
overlap of signal. LoCs were imaged with a 25x water immersion objective (NA=0.95,
Leica), with standard settings across chips labelled the same way. Z stacks were
subsequently deconvolved using the Huygens Deconvolution Software (Scientific
Volume Imaging) and 3D views were rendered using the FIJI ClearVolume plugin[64].
Custom-written software in MATLAB was used to segment and identify the 3D
volume, mean intensity, and number of RNA dots in each field of view using the
nestedSortStruct algorithm for MATLAB written by the Hughey Lab
(https://www.github.com/hugheylab/nestedSortStruct, GitHub). Statistical analysis
was performed using Origin 9.2 (OriginLabs) and P-values were calculated using a
Kruskal-Wallis one-way ANOVA test, with the null hypothesis that the medians of
each population were equal.
Data availability statement

The datasets generated during and/or analysed during the current study are available from the corresponding authors on reasonable request. Software code used to analyse these datasets will be uploaded to Zenodo prior to publication

References

1. Bhatia, S. N. & Ingber, D. E. Microfluidic organs-on-chips. *Nat. Biotechnol.* **32**, 760–772 (2014).

2. Thacker, V. V *et al.* A lung-on-chip infection model reveals protective and permissive roles of alveolar epithelial cells in tuberculosis. *bioRxiv* 2020.02.03.931170 (2020) doi:10.1101/2020.02.03.931170.

3. Villenave, R. *et al.* Human Gut-On-A-Chip Supports Polarized Infection of Coxsackie B1 Virus In Vitro. *PLoS One* **12**, e0169412 (2017).

4. Ortega-Prieto, A. M. *et al.* 3D microfluidic liver cultures as a physiological preclinical tool for hepatitis B virus infection. *Nat. Commun.* **9**, 1–15 (2018).

5. Grassart, A. *et al.* Bioengineered Human Organ-on-Chip Reveals Intestinal Microenvironment and Mechanical Forces Impacting Shigella Infection. *Cell Host Microbe* **26**, 435-444.e4 (2019).

6. Torrelles, J. B. & Schlesinger, L. S. Integrating Lung Physiology, Immunology, and Tuberculosis. *Trends Microbiol.* **25**, 688–697 (2017).

7. Gattinoni, L. *et al.* COVID-19 does not lead to a "typical" acute respiratory distress syndrome. *American Journal of Respiratory and Critical Care Medicine* vol. 201 1299–1300 (2020).

8. Marini, J. J. & Gattinoni, L. Management of COVID-19 Respiratory Distress. *JAMA - Journal of the American Medical Association* vol. 323 2329–2330 (2020).

9. Yao, Y. *et al.* D-dimer as a biomarker for disease severity and mortality in

COVID-19 patients: A case control study. *J. Intensive Care* **8**, 49 (2020).

10.   Messner, C. B. *et al.* Ultra-High-Throughput Clinical Proteomics Reveals

Classifiers of COVID-19 Infection. *Cell Syst.* **11**, 11-24.e4 (2020).

11.   Ackermann, M. *et al.* Pulmonary vascular endothelialitis, thrombosis, and

angiogenesis in Covid-19. *N. Engl. J. Med.* **383**, 120–128 (2020).

12.   Carsana, L. *et al.* Pulmonary post-mortem findings in a series of COVID-19

cases from northern Italy: a two-centre descriptive study. *Lancet Infect. Dis.*

(2020) doi:10.1016/S1473-3099(20)30434-5.

13.   Teuwen, L. A., Geldhof, V., Pasut, A. & Carmeliet, P. COVID-19: the

vasculature unleashed. *Nature Reviews Immunology* vol. 20 389–391 (2020).

14.   Lang, M. *et al.* Pulmonary Vascular Manifestations of COVID-19 Pneumonia.

Radiol. Cardiothorac. Imaging **2**, e200277 (2020).

15.   Claudio, D. *et al.* Acute Lung injury evolution in Covid-19. *medRxiv*

2020.08.09.20170910 (2020) doi:10.1101/2020.08.09.20170910.

16.   Blanco-Melo, D. *et al.* Imbalanced Host Response to SARS-CoV-2 Drives

Development of COVID-19. *Cell* **181**, 1036-1045.e9 (2020).

17.   Hui, K. P. Y. *et al.* Tropism, replication competence, and innate immune

responses of the coronavirus SARS-CoV-2 in human respiratory tract and

conjunctiva: an analysis in ex-vivo and in-vitro cultures. *Lancet Respir. Med.* **8**,

687–695 (2020).

18.   Hou, Y. J. *et al.* SARS-CoV-2 Reverse Genetics Reveals a Variable Infection

Gradient in the Respiratory Tract. *Cell* **182**, 429-446.e14 (2020).

19.   Nawroth, J. C. *et al.* A Microengineered Airway Lung Chip Models Key

Features of Viral-induced Exacerbation of Asthma. *Am. J. Respir. Cell Mol.*

Biol. **63**, 591–600 (2020).

20.   Huang, J. *et al.* SARS-CoV-2 Infection of Pluripotent Stem Cell-derived

787   Human Lung Alveolar Type 2 Cells Elicits a Rapid Epithelial-Intrinsic

788   Inflammatory Response. *bioRxiv* 2020.06.30.175695 (2020)

789   doi:10.1101/2020.06.30.175695.

790   21.   Abo, K. M. *et al.* Human iPSC-derived alveolar and airway epithelial cells can

791   be cultured at air-liquid interface and express SARS-CoV-2 host factors.

792   *bioRxiv* 2020.06.03.132639 (2020) doi:10.1101/2020.06.03.132639.

793   22.   Youk, J. *et al.* Robust three-dimensional expansion of human adult alveolar

794   stem cells and SARS-CoV-2 infection. *bioRxiv* **2**, 2020.07.10.194498 (2020).

795   23.   Qi, F., Qian, S., Zhang, S. & Zhang, Z. Single cell RNA sequencing of 13

796   human tissues identify cell types and receptors of human coronaviruses.

797   *Biochem. Biophys. Res. Commun.* **526**, 135–140 (2020).

798   24.   Hikmet, F. *et al.* The protein expression profile of ACE2 in human tissues.

799   *Mol. Syst. Biol.* **16**, (2020).

800   25.   Song, J. *et al.* Upregulation of angiotensin converting enzyme 2 by shear stress

801   reduced inflammation and proliferation in vascular endothelial cells. *Biochem.*

802   *Biophys. Res. Commun.* **525**, 812–818 (2020).

803   26.   Valdembri, D. *et al.* Neuropilin-1/GIPC1 Signaling Regulates α5β1 Integrin

804   Traffic and Function in Endothelial Cells. *PLoS Biol.* **7**, e1000025 (2009).

805   27.   Daly, J. L. *et al.* Neuropilin-1 is a host factor for SARS-CoV-2 infection.

806   *bioRxiv* 2020.06.05.134114 (2020) doi:10.1101/2020.06.05.134114.

807   28.   Cantuti-Castelvetri, L. *et al.* Neuropilin-1 facilitates SARS-CoV-2 cell entry

808   and provides a possible pathway into the central nervous system. *bioRxiv*

809   2020.06.07.137802 (2020) doi:10.1101/2020.06.07.137802.

29.   Hoffmann, M. *et al.* SARS-CoV-2 Cell Entry Depends on ACE2 and

TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* **181**,

271-280.e8 (2020).

30. Chu, H. *et al.* Comparative tropism, replication kinetics, and cell damage

profiling of SARS-CoV-2 and SARS-CoV with implications for clinical

manifestations, transmissibility, and laboratory studies of COVID-19: an

observational study. *The Lancet Microbe* **1**, e14–e23 (2020).

31. Kim, D. *et al.* The Architecture of SARS-CoV-2 Transcriptome. *Cell* **181**, 914-

921.e10 (2020).

32. V'kovski, P. *et al.* Disparate temperature-dependent virus – host dynamics for

SARS-CoV-2 and SARS-CoV in the human respiratory epithelium. *bioRxiv*

2020.04.27.062315 (2020) doi:10.1101/2020.04.27.062315.

33. Tseng, C.-T. K. *et al.* Apical Entry and Release of Severe Acute Respiratory

Syndrome-Associated Coronavirus in Polarized Calu-3 Lung Epithelial Cells.

J. Virol. **79**, 9470–9479 (2005).

34. Zelaya, H., Rothmeier, A. S. & Ruf, W. Tissue factor at the crossroad of

coagulation and cell signaling. *J. Thromb. Haemost.* **16**, 1941–1952 (2018).

35. Mast, A. E. Tissue Factor Pathway Inhibitor: Multiple Anticoagulant Activities

for a Single Protein. *Arteriosclerosis, Thrombosis, and Vascular Biology* vol.

36 9–14 (2016).

36. Xia, H. *et al.* Evasion of Type I Interferon by SARS-CoV-2. *Cell Rep.* **33**,

108234 (2020).

37. Miorin, L. *et al.* SARS-CoV-2 Orf6 hijacks Nup98 to block STAT nuclear

import and antagonize interferon signaling. *Proc. Natl. Acad. Sci. U. S. A.* **117**,

28344–28354 (2020).

38. Lopez-Castejon, G. & Brough, D. Understanding the mechanism of IL-1β

secretion. *Cytokine and Growth Factor Reviews* vol. 22 189–195 (2011).

39.   Tanaka, T., Narazaki, M. & Kishimoto, T. Il-6 in inflammation, Immunity, And

disease. *Cold Spring Harb. Perspect. Biol.* **6**, 16295–16296 (2014).

40.   Zunke, F. & Rose-John, S. The shedding protease ADAM17: Physiology and

pathophysiology. *Biochimica et Biophysica Acta - Molecular Cell Research*

vol. 1864 2059–2070 (2017).

41.   Dreymueller, D. *et al.* Lung endothelial ADAM17 regulates the acute

inflammatory response to lipopolysaccharide. *EMBO Mol. Med.* **4**, 412–423

(2012).

42.   Tanaka, T., Narazaki, M., Ogata, A. & Kishimoto, T. A new era for the

treatment of inflammatory autoimmune diseases by interleukin-6 blockade

strategy. *Seminars in Immunology* vol. 26 88–96 (2014).

43.   Klein, S. *et al.* SARS-CoV-2 structure and replication characterized by in situ

cryo-electron tomography. *bioRxiv* 2020.06.23.167064 (2020)

doi:10.1101/2020.06.23.167064.

44.   Cevik, M. *et al.* SARS-CoV-2, SARS-CoV, and MERS-CoV viral load

dynamics, duration of viral shedding, and infectiousness: a systematic review

and meta-analysis. *The Lancet Microbe* (2020) doi:10.1016/S2666-

5247(20)30172-5.

45.   Weibel, E. R. On the Tricks Alveolar Epithelial Cells Play to Make a Good

Lung. *Am. J. Respir. Crit. Care Med.* **191**, 504–513 (2015).

46.   Monteil, V. *et al.* Inhibition of SARS-CoV-2 Infections in Engineered Human

Tissues Using Clinical-Grade Soluble Human ACE2 In Brief Clinical-grade

recombinant human ACE2 can reduce SARS-CoV-2 infection in cells and in

multiple human organoid models. Inhibition of SARS-CoV-2 Infections in

Engineered Human Tissues Using Clinical-Grade Soluble Human ACE2. *Cell*

181, 905-913.e7 (2020).

47.   Jiang, Z., Tang, X., Xiao, R., Jiang, L. & Chen, X. Dengue virus regulates the

expression of hemostasis-related molecules in human vein endothelial cells. *J.*

Infect. **55**, e23–e28 (2007).

48.   Guo, H. F. & Vander Kooi, C. W. Neuropilin functions as an essential cell

surface receptor. *Journal of Biological Chemistry* vol. 290 29120–29126

(2015).

49.   Newman, P. J. & Newman, D. K. Signal transduction pathways mediated by

PECAM-1: New roles for an old molecule in platelet and vascular cell biology.

Arteriosclerosis, Thrombosis, and Vascular Biology vol. 23 953–964 (2003).

50.   Zhang, J. J., Kelm, R. J., Biswas, P., Kashgarian, M. & Madri, J. A. PECAM-1

modulates thrombin-induced tissue factor expression on endothelial cells. *J.*

Cell. Physiol. **210**, 527–537 (2007).

51.   Falati, S. *et al.* Platelet PECAM-1 inhibits thrombus formation in vivo. *Blood*

107, 535–541 (2006).

52.   Cheung, K. *et al.* CD31 signals confer immune privilege to the vascular

endothelium. *Proc. Natl. Acad. Sci. U. S. A.* **112**, E5815–E5824 (2015).

53.   Zuo, Y. *et al.* Neutrophil extracellular traps in COVID-19. *JCI Insight* **5**,

(2020).

54.   Manne, B. K. *et al.* Platelet gene expression and function in patients with

COVID-19. *Blood* **136**, 1317–1329 (2020).

55.   Li, L. *et al.* Serum levels of soluble PECAM-1 in COVID-19 patients are

associated with disease severity. *J. Infect. Dis.* (2020)

doi:10.1093/infdis/jiaa642.

56.   Mansouri, M. *et al.* Kaposi sarcoma herpesvirus K5 removes CD31/PECAM

887   from endothelial cells. *Blood* **108**, 1932–1940 (2006).

888   57.   Hadjadj, J. *et al.* Impaired type I interferon activity and inflammatory responses

889   in severe COVID-19 patients. *Science (80-.).* eabc6027 (2020)

890   doi:10.1126/science.abc6027.

891   58.   Thoms, M. *et al.* Structural basis for translational shutdown and immune

892   evasion by the Nsp1 protein of SARS-CoV-2. *Science* (2020)

893   doi:10.1126/science.abc8665.

894   59.   Wu, K., Zou, J. & Chang, H. Y. RNA-GPS Predicts SARS-CoV-2 RNA

895   Localization to Host Mitochondria and Nucleolus. *bioRxiv Prepr. Serv. Biol.*

896   2020.04.28.065201 (2020) doi:10.1101/2020.04.28.065201.

897   60.   Parr, J. B. Time to Reassess Tocilizumab's Role in COVID-19 Pneumonia.

898   *JAMA Internal Medicine* (2020) doi:10.1001/jamainternmed.2020.6557.

899   61.   Hassell, B. A. *et al.* Human Organ Chip Models Recapitulate Orthotopic Lung

900   Cancer Growth, Therapeutic Responses, and Tumor Dormancy In Vitro. *Cell*

901   *Rep.* **21**, 508–516 (2017).

902   62.   Huh, D. *et al.* Reconstituting organ-level lung functions on a chip. *Science* **328**,

903   1662–8 (2010).

904   63.   McBride, C. E., Li, J. & Machamer, C. E. The Cytoplasmic Tail of the Severe

905   Acute Respiratory Syndrome Coronavirus Spike Protein Contains a Novel

906   Endoplasmic Reticulum Retrieval Signal That Binds COPI and Promotes

907   Interaction with Membrane Protein. *J. Virol.* **81**, 2418–2428 (2007).

908   64.   Royer, L. A. *et al.* ClearVolume: Open-source live 3D visualization for light-

909   sheet microscopy. *Nature Methods* vol. 12 480–481 (2015).

Figures

Fig. 1. SARS-CoV-2 infection in a human lung-on-chip model.

(**A**) Schematic of the LoC model for SARS-CoV-2. Maximum intensity projections show that confluent layers of epithelial and endothelial cells with strong expression of tight junction markers populate the top (**B**) and bottom faces (**C**) of the porous membrane. Actin, CD31, TJP-1 and nuclear labelling are shown in azure, spring green, bright pink, and electric indigo LUTs respectively. qRT-PCR characterization of the epithelial and endothelial layers on-chip for SARS-CoV-2 entry markers (*ACE2, TMPRSS2, NRP1*) for cells extracted from the apical and vascular channels from n=3 uninfected LoCs (**E**) and n=4 infected LoCs at 1 day-post infection (**F**). In both cases expression is normalized to *GAPDH* levels. *ACE2* levels were also characterized using RNAscope, a representative 3D view of a 232 x 232 μm^2 field of view of the epithelial face of an uninfected control at 3 days at air-liquid interface (3C) (**D**) and infected LoC

at 3 dpi (**G**) are shown. The nuclear labelling is indicated in electric indigo and *ACE2*

mRNA is labelled in spring green LUT. (**H**) *ACE2* expression decreases between 1and

3 dpi in the LoC model, quantified on a per-cell basis for 4-5 fields of view across one

control and one infected chip at each timepoint. (**I, J**) A plot of the fold change in

expression of host proteins required for viral entry at 1 day-post infection (n=4 LoCs)

and 3 days-post infection (n=2 LoCs) vs. uninfected controls (n=3 LoCs). The bars

represent the mean value, the solid line represents the median value, and the error bars

represent the standard deviation. P-values are calculated using a Kruskal-Wallis one-

way ANOVA test.

940
Fig. 2. Viral replication but lack of release of infectious virions characterizes

SARS-CoV-2 infection in the LoC model.

The kinetics of release of SARS-CoV-2 progeny was assessed from samples obtained

from a wash of the apical face of the chip (**A, B**), and the vascular effluent (**A**) collected

at each day-post infection. (**A**) The number of viral genomes in each sample was

quantified using a one-step qRT-PCR assay for the SARS-CoV-2 N gene. The starting

inoculum introduced to the apical side (labelled 'Inn') corresponded to 200-300 PFU.

Lines join datapoints from the same LoC, reconstituted with (green squares) and

without (dark blue circles) macrophages. Viral genome numbers within the apical wash

showed a declining trend over 3 days of infection. Viral genomes are not detected ('nd')

in the vascular effluent for all the chips shown. (**B**) Measurements of plaque forming

units (PFU) from the apical wash of LoCs reconstituted without macrophages (n=8 for

1 dpi, n=2 for 2 dpi) confirms the presence of infectious virions. (**C**) Quantification of

the intracellular levels of transcripts for the SARS-CoV-2 *N* and SARS-CoV-2 *E* genes

in the epithelial and endothelial layers of infected LoCs at 1 dpi (n=3) relative to

expression of the eukaryotic housekeeping gene *GAPDH*. (**D**) SARS-CoV-2 *N* : SARS-

CoV-2 *E* ratio in the epithelial and endothelial layers of the infected LoCs shown in

(**C**). In all plots, the bars represent the mean value and the error bars represent the

standard deviation, the solid black line in (**A**) represents the median.

Fig. 3. SARS-CoV-2 infection rapidly disrupts confluency of the endothelial layer.
(**A-D**) 3D views of representative 232 x 232 μm² (**A, B**) and 155 x 155 μm² (**C, D**)
fields of view from confocal imaging of an uninfected control (**A**), and infected chips
without macrophages at days 2 (**B**) and 3 (**C, D**) post infection respectively. Actin and
nuclear labelling are shown in azure and indigo, respectively. (**B**) At 2 dpi, adjacent
areas of endothelial cell clusters identified by a stronger nuclear stain and brighter actin
staining (yellow arrows) and cells with increased permeability and loss of confluence
(white arrows) are indicated. (**C**) At 3 dpi, a significant loss of barrier integrity is
observed in the endothelial layer, whereas the epithelial layer directly above this field
of view (**D**) is relatively intact. A small area of missing cells in the epithelial layer is
indicated with a white arrow in (**D**). Plots of density of nuclei (number of nuclei per
1E4 μm²) on the epithelial (**E**) and endothelial (**F**) faces of infected LoCs both with and
without macrophages for characterization of the cellular layer integrity. The bars

976 represent the mean value, the solid line represents the median and the error bars

977 represent the standard deviation from at least 4 fields of view each from two

978 independent LoCs for each timepoint. Data from uninfected controls corresponding to

979 the 3 dpi timepoint are labelled '3C'. The fields of view in (**C, D**) are shown also in

980 (**G, H**) respectively, SARS-CoV-2 spike (S) proteins labelled via immunofluorescence

981 are false-colored in amber. Zooms in (**I**) and (**J**) show clear evidence of intracellular

982 spike protein within both endothelial and epithelial cells, respectively. (**K**) A 3D view

983 of the field of views in (**C, D**) and (**G, H**) respectively highlights the relative damage

984 to the endothelial layer versus the epithelium, as well as the basolateral transmission of

985 S protein via junctions in the pores of the membrane (identified with white arrows). P-

986 values are calculated using a one-way Kruskal-Wallis ANOVA test.

987
Fig. 4. SARS-CoV-2 infection leads to a loss of tight junctions, generation of CD31-

endothelial cell clusters and a pro-coagulatory environment.

(**A-D**) Maximum intensity projects of 232 x 232 μm² fields of view from the epithelial

(**A**, **B**) and endothelial layers of an uninfected control LoC. Epithelial cells show strong

actin (**A**) and TJP1 expression (**B**, identified via antibody staining) at the cell junction

boundaries. Endothelial cells also show high levels of TJP-1 expression (**C**) together

with CD31 expression at cell junctions (**D**, identified via antibody labelling). Actin,

TJP-1, CD31, and nuclear labelling are indicated with the azure, bright pink, spring

green and electric indigo LUTs respectively. 3D views of one 232 x 232 μm² field of

view of the epithelial (**E**) and (**F-H**) three 232 x 232 μm² fields of view of the

endothelial layers of an infected LoC at 1 dpi. Areas with lower TJP1 expression in the

epithelial layer in (**E**) are indicated by white arrows. (**F-H**) Areas with loss of tight

junctions between cells on the endothelial surface is indicated by white arrows. The

proportion of the surface area covered by confluent cells decreases sequentially from

(**F**) to (**H**). Endothelial cell clusters with no CD31 expression are clearly visible and the

proportion of the surface area covered by these clusters increases sequentially from (**F**)

to (**H**). In (**F-H**), spike protein is identified via an anti-SARS-spike antibody and shown

in amber LUT, and von Willebrand Factor identified by antibody labelling is shown in

bring pink LUT. Zooms of two endothelial cell clusters (**I, J** and **K, L**) in the endothelial

layer of another infected LoC at 1 dpi confirm that little or no CD31 expression is

detected (**I**, **K**) despite high levels of actin and TJP-1 protein expression in these cells.

(**J**, **L**). (**M**) Plot of the fold change in expression of *TJP1* (epithelial and endothelial

layers) and *TJP1*, *CDH5*, and *CD31* (endothelial layers only) in infected LoCs at 1 dpi

(n=3) vs, uninfected controls (n=3). (**N**) Plot of the fold change in gene expression of

endothelial cell specific coagulation markers - Tissue Factor (*F3*), Tissue Factor

Pathway Inhibitor (*TFPI*), Thrombomodulin (*THBD*), Plasminogen Activator Inhibitor

(*SERPINE1*) and von Willebrand Factor (*VWF*) in infected LoCs at 1 dpi (n=3) vs.

uninfected controls (n=3). In all plots, bars represent the mean value and error bars

represent the standard deviation.

1017

Fig. 5. RNAscope analysis reveals intracellular localization and slow accumulation of intracellular viral RNA in the LoC model.

3D views of 232 x 232 μm² fields of view of the epithelial (**A, G**) and endothelial layer (**B, H**) from infected LoCs reconstituted without macrophages at 1 (**A, B**) and 3 dpi (**G, H**). *Orf1ab* antisense RNA, *S* RNA, *ACE2* mRNA, and nuclear staining with DAPI are false-colored pink, amber, spring green, and electric indigo, respectively. (**C, D**) Zooms corresponding to the region in (**A**) highlighted with white and yellow boxes, respectively. (**C**) An example of infection and intracellular replication in a cell with no detectable *ACE2* expression (white arrow) as well as nuclear localization of viral RNA (white arrowhead). (**D**) An example of an uninfected cell with *ACE2* mRNA expression (yellow arrow). (**E, F**) Zooms corresponding to the regions in (**B**) representing increased nucleic acid staining (hyperplasic) and normal levels of nucleic acid staining highlighted with white and yellow boxes, respectively. (**E**) Examples of infection of hyperplasic endothelial cells both with and without *ACE2* expression (**F**) Examples of

endothelial cell infection with no *ACE2* expression, nuclear localization of viral RNA

in an infected endothelial cell is indicated (white arrowheads). Quantification of viral

antisense RNA (**I, J**) and viral genomic RNA (**K, L**) from pairs of otherwise identical

LoCs reconstituted without (**I, K**) and with macrophages (**J, L**) analyzed at 1 and 3 dpi.

Plots show the number of spots normalized by the total number of cells in 4-6 field of

views detected using RNAscope and confocal imaging using identical imaging

conditions for all chips. Bars represent the mean value, the solid line represents the

median, and error bars represent the standard deviation. (**M**) Quantification of *ACE2*

expression per cell in epithelial cells and endothelial cells in LoCs reconstituted without

macrophages at 1 and 3 dpi respectively. P-values are calculated using a Kruskal-Wallis

One-Way ANOVA Test.

1044

Fig. 6. SARS-CoV-2 infection generates a persistent pro-inflammatory response in endothelial cells.

(**A**) Expression relative to *GAPDH* of pro-inflammatory cytokines (*TNFA, IL1B, IL6*), interferon genes (*IFNB, IFNL1, IFNL3*) and interferon stimulating genes (*IP10, ISG15*) in the epithelial and endothelial layer of infected LoCs at 1 dpi reconstituted without macrophages (n=3 for *IFNL1* and *IFNL3* in the epithelial layer, n=4 for all other data). (**B**) Fold-change in the markers in (**A**) relative to uninfected controls (n=3) (data in Fig. S12) at the same timepoint. The bar represents the mean and the error bars the standard deviation. (**C**) Enzyme-linked immunosorbent assay (ELISA) measurements for IL-6 in the vascular effluent for at least two infected LoCs at each timepoint reconstituted with and without macrophages. Uninfected controls for each condition are labelled with a 'C' and grey bars, and LoCs with Tocilizumab administration are indicated by ^. The height of the bars represents the mean, the median is represented by a solid line, and

the error bars represent the standard deviation. Each dot represents the mean of two

technical replicates. (**D**) Expression of *IL6R* and *ADAM17* relative to *GAPDH* in the

epithelial and endothelial layer of infected LoCs at 1 dpi (n=3 for the epithelial layer

and n=4 for endothelial layer). (***E***) Fold-change in expression relative to uninfected

controls (n=3 for both layers). (**F-I**) 3D views of representative 232 x 232 μm^2 fields of

view of the epithelial (**F, H**) and endothelial layer (**G, I**) from infected LoCs

reconstituted with macrophages at 1 (**F, G**) and 3 dpi (**H, I**). *IL6* mRNA, and nuclear

staining are indicated with the chartreuse and electric indigo LUTs respectively. (**K-N**)

Zooms corresponding to the regions in (**I**) highlighted with white (**K**) and yellow boxes

with solid (**L**) and dashed lines (**M, N**) respectively. In these panels, *orf1abantisense*

RNA (pink) and *S* RNA (amber) are also shown. The panels show examples of cells

with similar levels of viral infection but with no (**K**), intermediate (**L**) and high levels

(**M, N**) of *IL6* expression. (**J**) Quantification of *IL6* expression in epithelial and

endothelial cells from a pair of otherwise identical LoCs analyzed at 1 and 3 dpi,

respectively. Plots show the total number of spots normalized by the number of cells in

4-6 fields of view detected using RNAscope assay using identical imaging conditions

for all chips. Bars represent the mean value, the solid line represents the median, and

error bars represent the standard deviation. Data from uninfected controls for 3 days at

air-liquid interface is indicated by 'C'. The bar represents the mean and the error bars

the standard deviation. P-values are calculated using a Kruskal-Wallis One-Way

ANOVA Test.

1080
1081

Fig. 7. Tocilizumab treatment reduces vascular permeability but does not prevent formation of endothelial cell clusters with reduced *CD31* expression.

Maximum intensity projections of representative 232 x 232 μm^2 fields of view of the endothelial layer from an infected LoC reconstituted without macrophages at 2 dpi with (**A**) and without Tocilizumab treatment (**D**). Actin and nuclear staining are indicated with the azure and electric indigo LUTs respectively, the actin staining is saturated for the endothelial cell clusters to highlight cells with lower levels of actin. (**B, C**) Zooms corresponding to the region in (**A**) highlighted with white and yellow boxes, respectively. (**B**) A region with normal vascular profile, (**C**) A region of endothelial cell cluster formation with increased actin staining. (**E, F**) Zooms corresponding to the regions in (**D**) representing regions highlighted with white and yellow boxes, respectively. (**E**) An example of reduced tight junctions between endothelial cells; bare patches are indicated with arrows. (**F**) A region of endothelial cell clusters. (**G**) 3D views from two 232 x 232 mm2 field of views of the endothelial layer of a Tocilizumab treated infected LoC at 2 dpi, immunostaining for the tight junction protein CD31 is shown in the Spring Green LUT. Endothelial cell clusters with low CD31 expression

and altered actin staining are also visible and indicated by yellow arrows. (**I**) A plot of

the proportion of pixels (shown on a logarithmic scale) with regions of low F-actin

intensity identified via cut-off thresholds defined as a percentage of the maximum

intensity in 6-7 regions of interest each for LoCs with and without Tocilizumab

administration. ROIs were chosen from 232 x 232 μm^2 fields of view and were defined

to exclude the occurrence of hyperplasic cells. A significantly larger fraction of the

surface area of the ROIs for the untreated chips have low F-actin intensities. P-values

are calculated using a Kruskal-Wallis One-Way ANOVA Test.

1107
1108