Dissecting the microenvironment around biosynthetic scaffolds in murine skin wound healing

Chen Hu[a,#], Chenyu Chu[a,#], Li Liu[b], Shue Jin[c], Renli Yang[a], Shengan Rung[a],

Jidong Li[c], Yili Qu[a], Yi Man[a,*]

[a] *Department of Oral Implantology & State Key Laboratory of Oral Diseases & National Clinical Research Center for Oral Diseases, West China Hospital of Stomatology, Sichuan University, , Chengdu, Sichuan 610041, China*

[b] *State Key Laboratory of Biotherapy, West China Hospital, Sichuan University, Collaborative Innovation Center for Biotherapy, Chengdu, Sichuan 610041, China*

[c] *Research Center for Nano-Biomaterials, Analytical and Testing Center , Sichuan University , Chengdu 610064, China.*

[#] These authors contributed equally to this work

[*] Corresponding to Professor. Chair. **Yi Man**, E-mail: manyi780203@126.com

Corresponding author:

Yi Man, DDS, PhD

Professor and Chair

Department of Oral Implantology

State Key Laboratory of Oral Diseases, West China Hospital of Stomatology, Sichuan University, 14#, 3rd section, Renmin South Road

Chengdu 610041, China

Tel: 0086-28-85503571

Fax: 0086-28-85483678

E-mail: manyi780203@126.com

Abstract: Structural properties of biomaterials play critical roles in guiding cell behaviors and influence the immune response against them. We fabricated electrospun membranes with three types of surface topography (Random, Aligned, and Latticed). The aligned membranes showed immunomodulatory ability, and led to faster wound healing, reduced fibrotic response and enhanced regeneration of cutaneous appendages when used in skin wound repair. Based on that, we performed single-cell RNA sequencing analysis on cells from wounded mouse skin in the presence or absence of the Aligned scaffold. Keratinocytes, fibroblasts, and immune cells including neutrophils, monocytes, macrophages, dendritic cells, and T cells showed diverse cellular heterogeneity. More hair follicle progenitor cells, inner root sheath cells (anagen-related) and fibroblast subsets were found in the Aligned group, which corresponded to the improved regeneration of hair follicles and faster wound closure in the presence of scaffold. Immune responses towards the biomaterial differed from that of control group. In aligned samples, infiltrated macrophages and neutrophils were reduced, whereas more effector T cells were recruited. The time course of immune response was possibly advanced towards an adaptive immunity-dominant stage by the scaffold. The microenvironment around scaffold involved intricate interplay of immune cells and cutaneous cells, and wound healing was the comprehensive results of numerous influencing factors working together.

Key words: Surface topography, immune microenvironment, immunomodulatory ability, cellular heterogeneity

Introduction

Biomaterials and devices implanted in the body have a broad spectrum of clinical applications including tissue regeneration, cell transplantation, controlled drug release, and continuous monitoring of physiological conditions (1, 2). Among the components of a cell microenvironment, structural features (macroscale, microscale and nanoscale features) play critical roles in guiding cell behaviors (3). Electrospinning technology has been widely applied in preparing scaffolds due to its simplicity, capacity to form fibers on the micro- and nanoscale, structural control of electrospun membranes and cost-effectiveness (4). Electrospun nanofibers could be assembled into well-ordered nanofiber meshes with different morphology, e.g. parallel alignment or latticed patterns of nanofibers (5, 6), and have been preferentially applied to the regeneration of diverse tissues like skin and bone (6, 7). Upon implantation of a material, cells of both the innate and adaptive immune system have a role in the host response (8). Previous studies illustrated the type 1 (pro-inflammatory) immune polarization driven by T helper 1 (Th1) cells, and the induced pro-inflammatory M1 activation of macrophages (stimulated by IL-2 and IFN-γ from Th1 cells). By contrast, in type 2 immune response, T helper 2 (Th2) cells produce cytokines like IL-4 and IL-13, which regulate the polarization of macrophages towards an anti-inflammatory M2 activation (9, 10). More recently, a type 17 immune response was reported to promote chronic fibrosis in tissue around implants (11, 12). In summary, previous studies usually focused on certain types of immune cells, like macrophages, and explored their roles in the host response. However, the immune response is jointly regulated by

various immune cells, whose phenotype and function are dictated by external and

internal signals. An overview of different immune cells in the microenvironment will

aid in comprehensive understanding of the immune responses elicited by scaffolds.

Technological advances such as single-cell RNA sequencing (scRNA-seq) (13, 14)

have enabled cell populations, functions, and the nuances of their phenotypes in vivo

to be studied at a high resolution. By changing the collector, we developed

poly(lactic-co-glycolic acid)-fish collagen (PLGA-FC) hybrid electrospun scaffolds

with three types of surface topography, i.e. the group with randomly oriented fibers

(Random group), the group with mesh-like topography in macroscale and randomly

oriented fibers in microscale (Latticed group), and the group with aligned fibers

(Aligned group). We explored the regenerative outcomes of these scaffolds in

rat/mouse dorsal skin excisional wounds, and evaluated their immunomodulatory

properties. The scaffold with the best performance was further investigated.

Microenvironment around the scaffold was probed by scRNA-seq. Heterogeneity of

keratinocytes, fibroblasts, and immune cell populations, cellular functions, and their

interactions in vivo were explored.

Results

Evaluation of wound healing in a rat skin wound model

We fabricated scaffolds with random, aligned and latticed patterns of fibers (**Fig. 1A**),

and placed them below the full-thickness excisional wound (diameter=6mm) on rat

dorsal skin (SD rat). The Ctrl group received no scaffolds (**Fig. 1C**). Biophysical

properties of the scaffolds were summarized in **Fig. S1**. Workflow for evaluating

wound healing was summarized in Fig. 1B. Wound healing rate was significantly

accelerated by the aligned membranes (**Fig. 1D-E**). The Latticed group showed

delayed wound healing with the largest residual wound area left at day 7. On day 14,

basically all groups achieved complete closure of the wound. Above the scaffolds,

surrounding epithelium formed epithelial tongue as the first layer advancing towards

the wound (15). On day 7, the Aligned group presented the fastest coverage of the

wound, leaving the smallest gap width, whereas the latticed membrane seemed to

impede the advancement of surrounding epithelium (**Fig. 1F-G, Fig. S2**). On day 14,

re-epithelialization was completed in all groups except for the Latticed group. All

groups showed matured stratified epithelium (**Fig. 1F**). Immunofluorescent staining

for Krt5 (keratin secreted by keratinocytes in the basal layer) and Krt10 (keratin

secreted by differentiated keratinocytes in the supra-basal layers) showed that on day

7, the Aligned group had the largest area of newly formed keratinized epithelium, and

more keratinocytes were undifferentiated (Krt5-positive). On day 14, all groups

revealed formation of stratified epithelium (**Fig. 1H-I**) (15). For the dermis part,

wound space below epithelium was filled with granulation tissue on day 7. Collagen

kept being deposited and remodeled (**Fig. S2**). On day 28, the Aligned group had

more regenerated hair follicles and sebaceous glands than other treatment groups (**Fig.**

1F). The scaffolds were placed subcutaneously in rats to evaluate host response

against them. Thickness of fibrotic capsules around scaffolds was measured at day 3,

7 and 14. The Aligned group had the smallest fibrotic capsule thickness at all time

points (**Fig. 1J-K, Fig. S3**). The bulk-tissue RNA-Seq analysis for samples harvested

on day 7 (n=3 for each group) identified transcripts corresponding to 34459 genes,

distributed over 6 orders of magnitude of expression level. Principal component

analysis (PCA) of the data revealed that gene expression profiles of the Random and

Aligned samples were more similar, compared to that of Latticed and Ctrl samples

(**Fig. 1L**). In Gene ontology (GO) analysis, random and aligned scaffolds induced

up-regulated expression of genes associated with immune responses when compared

with the Ctrl samples, whereas the Latticed group did not (**Fig. 1M**). Kyoto

Encyclopedia of Genes and Genomes (KEGG) analysis also revealed gene enrichment

in an immune-related pathway "Cytokine-cytokine receptor interaction" (KEGGID

rno04060) (**Fig. S4A-B**) (16). Real-time Quantitative Polymerase Chain Reaction

(qPCR) confirmed these elevated genes observed in Aligned and Random groups (**Fig.**

S4C). The data suggested that random and aligned membranes were able to modulate

the local immune microenvironment. To explore whether the scaffolds had similar

performance in other species, we applied them in mouse skin wound model.

Wound healing in mouse

Workflow for evaluating wound healing in C57BL/6 mouse was summarized in Fig.

2A. We placed three types of scaffolds below the full-thickness excisional wound

(diameter=6mm) (**Fig. 2B**). The Ctrl group received no scaffolds. Wound coverage

was faster in the Aligned (residual wound area=29.29±4.81%) and Ctrl group

(20.80±4.66%) on day 7. On day 14, basically all groups achieved complete closure

of the wound (**Fig. 2C-D**). The Aligned group had the smallest gap width on day 7,

followed by the Ctrl group (**Fig. 2E-F**). On day 14, re-epithelialization was completed

in all groups except for some samples of the Latticed group. All groups showed

matured stratified epithelium. Regeneration of hair follicles was seen in Aligned

samples after 14 days of healing (**Fig. 2E**). Scaffolds were also placed subcutaneously

to evaluate the host response against them. The aligned membranes had the smallest

capsule thickness over the observation period (**Fig. 2G, Fig. S5**). Bulk-tissue

RNA-Seq for Aligned and Ctrl samples harvested on day 7 (n=3 for each group) was

performed. Compared with Ctrl, the Aligned group showed increased genes enriched

in inflammatory response, leukocyte chemotaxis and migration (**Fig. 2I**). KEGG

analysis revealed elevated gene expression in immuno-related signaling pathways for

the Aligned group (KEGGID mmu04657, mmu04668, mmu04060 and mmu04064)

(**Fig. 2J**). To summarize, in both rat and mouse models, the aligned membranes led to

faster wound healing, reduced fibrotic response and enhanced regeneration of

cutaneous appendages compared to other membranes. Meanwhile, they showed

immunomodulatory properties. We therefore sought to explore how aligned

membranes regulated peri-implant microenvironment. Here, we used scRNA-seq to

sequence cells from the wounded murine full-thickness skin 7 days post-wounding.

Single-cell transcriptome analysis of full-thickness skin after wounding and

scaffold placement

We isolated cells from the Aligned and Ctrl samples (n=4 biological replicates in each

group), and applied them to the 10X scRNA-seq platform (**Fig. 3A**). A total of 8,982

cells in the Ctrl group and 9,593 cells in the Aligned group were captured. After cell

filtering, 17,181 single cell transcriptomes were included in the final dataset (8,869

for the Aligned group and 8,312 for the Ctrl group) (**Fig. S6**). We first

computationally pooled cells from Ctrl and Aligned groups to create a virtual

aggregate. Unsupervised clustering using Seurat categorized the cells into 26 clusters

based on global gene expression patterns (Fig. S7), which were assigned to thirteen

main classes of cells (**Fig. 3B**): keratinocytes (KER), fibroblasts (FIB), sebocytes

(SEB), smooth muscle cells (SMC), endothelial cells (EC), Schwann cells (SC),

melanocytes (MEL), innate lymphoid cells (ILC), monocyte-macrophages (MAC), T

cells (TC), neutrophils (NEU), dendritic cells (DC) and B cells (BC). Marker genes

for each cell cluster were shown in the heatmap (**Fig. 3C**). The composition of each

cell cluster was listed so that the proportion of cells from two groups could be

identified across all cell clusters (**Fig. 3B**). As smooth muscle cells (57% from the

Aligned group and 43% from the Ctrl group) could hardly regenerate at 7 days

post-wounding, we regarded a proportion within 50±7% as equilibrium between two

groups. Genes related to macrophages (Itgam, Cd68, Arg1, Mrc1) were significantly

higher expressed by the Ctrl samples (**Fig. 3D**).

Sub-clustering of keratinocytes reveals a higher proportion of hair follicle

progenitor cells from the Aligned group

We selected cells that were in the first-level clustering defined as keratinocyte, and

subjected them to a second round of unsupervised clustering (**Fig. 4A**). Terminal

cornification of epidermis is achieved by keratinocytes passing through basal layers,

differentiated layers, and cornified layers (15). In our study, the inter-follicular

epithelium was composed of the basal layer cells (IFEB, $Krt5^{hi}Krt14^{hi}$), supra-basal

layer cells (Krt10^{hi}Krt1^{hi} IFED1, Krt10^{hi}Mt4^{hi} IFED2), and cornified layer (Lor^{+}

IFED3) cells (**Fig. 4A-B**). The location of keratinocyte subsets was marked in **Fig. 4C**.

Hair follicles in this study were in second telogen since the mice were 8 to 10 weeks

old when they were sacrificed (17). However, anagen hair follicle gene signatures

were also found, possibly resulting from hair follicle regeneration after wounding.

Upper hair follicle cells were separated into three subsets named uHF1

(Krt79^{hi}Krt17^{hi}), uHF2 (Klk10^{+}Krt79^{hi}), and uHFB (the upper hair follicle basal layer

cells, Sostdc1^{hi}Apoe^{hi}). The hair follicle progenitor cells (Krt28^{hi}Lhx2^{hi}Mki67^{hi} HFP)

were highly proliferative. Germinative layer cells (Mt2^{hi}Dcn^{hi}) belonging to anagen

hair follicles also expressed high cell proliferation related genes like Top2a and Birc5.

Meanwhile, they had a basal cell gene signature (Krt5^{hi}Krt14^{hi}). The inner root sheath

(IRS) and cortex cells were characterized by Krt28, Krt27, Krt73, and Krt25, markers

for the Henle and Huxley layers of anagen hair follicles (18). Cells of the inner bulge

layer (IB) and outer bulge layer (OB) expressed their typical gene signatures (**Fig.**

4A-C). When analyzing the inter-group differences, the Aligned group contributed to

a larger proportion of hair follicle progenitor cells (HFP) and highly proliferative

inner root sheath cells (IRS) (**Fig. 4A**). The overall difference analysis also revealed

up-regulation of genes (Krt28, Sox18) related to hair follicle stem cells in the Aligned

group (**Fig. 3D**). This might explain the enhanced hair follicle regeneration observed

in Aligned groups either on rat or mouse models.

Inter-group differences in fibroblasts suggest more active ECM formation in the

Aligned group

The dermis consists of several layers: the papillary dermis lies closest to the epidermis,

the underlying reticular dermis contains the bulk of the fibrillary extracellular matrix,

and beneath the reticular dermis lies the hypodermis (19). Fibroblasts from different

layers presented distinct gene signatures (**Fig. 4D**). There were four populations of

papillary fibroblasts (Crabp1$^+$Col23a1$^+$) (20, 21). Dermal fibroblasts expressed

increased Ccl11 and Dcn. The Gpx3$^+$Plac8^{hi} subset was identified as hypodermis

fibroblast located to the adventitia (HDF1), and the Gpx3$^+$Plac8^{lo} subset (HDF2) was

around the panniculus carnosus muscle (18). Contractile myo-fibroblasts (Acta2$^+$ MF)

expressed elevated genes associated with cell proliferation (Birc5^{hi}). Composition of

each subset revealed that more papillary fibroblasts (PF2), dermal fibroblasts (DF),

hypodermis fibroblasts (HDF2) and myo-fibroblasts (MF) were from the Aligned

group, suggesting more robust ECM formation in the presence of scaffolds.

Neutrophils and dendritic cells are more abundant in the Ctrl samples

After wounding, neutrophils close to the focus of injury migrate toward the nidus,

followed by those recruited from more than 200μm from the site of tissue injury (22).

Three neutrophil subsets were identified in our study (**Fig. 4E**). Peripheral blood

neutrophils (PBN) expressed typical gene signatures including Csf3r, Pglyrp1, Il1b,

and Retnlg. Neutrophil 2 (Neu2) expressed elevated Cd14 and Ccl3, and was

identified as antimicrobial phagocytic neutrophils. Neutrophil 1 (Neu1) expressed

elevated Ccr1, a chemokine receptor that mediates neutrophil migration.

Correspondingly, PBN showed gene enrichment in leukocyte chemotaxis and

migration; Neu1 expressed genes enriched in leukocyte/granulocyte chemotaxis and

migration, and Neu2 was enriched in anti-bacterial biological processes (**Fig. 4F**).

According to the correlation analysis, Neu 1 and Neu 2 were derived from PBN in

circulation (**Fig. 4G**). In Neu1 and Neu2, more cells belonged to the Ctrl group (**Fig.**

4E), suggesting more significant Neu1 and Neu2 infiltration in the Ctrl samples at a

proliferative stage (7 days post-wounding). Dendritic cells were classified into two

subpopulations (**Fig. 4H**). The subset derived from monocytes (DC) was

characterized by increased Cd207 and Cd86 (10, 23). The Langerhans cell subset bore

a keratinocyte gene signature, probably transferred from the resident

microenvironment (**Fig. 4I**) (24). Both DC and LC presented elevated major

histocompatibility complex (MHC) molecules, suggesting an antigen presenting

function of them (**Fig. 4J**). LCs contained more Ctrl-derived cells, indicating

differences in Langerhans cell infiltration between groups.

Macrophage heterogeneity and their down-regulation by scaffolds

To explore the heterogeneity of macrophages in vivo and their inter-group differences,

we subjected them to further unsupervised sub-clustering. Four subsets were

determined (**Fig. 5A**). The subset that showed increased anti-inflammatory genes

(Ccl8, Folr2, C1qa and Mrc1) was named anti-inflammatory macrophages (AIM) (25,

26). Another subset characterized by elevated expression of pro-inflammatory genes

(Ptgs2, Ccl3, Inhba, Nos2) was named pro-inflammatory macrophages (PIM) (27, 28).

The monocyte subset (Mono) showed higher Ly6c2, Plac8, Cd14, and Clec4e,

indicting inflammatory responses against lesions and microorganism (29, 30). The

Cytip^{hi}H2-Eb1^{hi} subset of cells was defined as monocyte-derived dendritic cells

(M-DC) (**Fig. 5B**) (31). We further found that canonical M1 and M2 markers were not

entirely consistent with computationally determined AIM and PIM. Arg1, a canonical

M2 marker, was expressed by basically all monocyte-macrophage subsets (AIM, PIM,

Mono), and was regarded as a pan-macrophage marker in this study (**Fig. 5B**).

Expression of another type 2 gene, Socs3, did not parallel Mrc1 expression either.

Similar disparity was found in the expression of canonical type 1 genes. M-DC, rather

than PIM, expressed more Cd86. Neither was the expression of Cd86 correlated with

Nfkbiz. Expression of other genes associated with fibrotic or regenerative

macrophage subsets in a scaffold immune microenvironment did not correspond with

these clusters either (**Fig. 5B**) (12). It was determined that Ly6c2, Arg1, Mrc1 and

Nos2 were sufficient to distinguish the computationally determined AIM, PIM and

Mono subsets. We performed flow cytometry on cells isolated from the Aligned and

Ctrl samples using Cd68 (a monocyte-macrophage marker, also expressed by some

neutrophils and dendritic cells) and the proposed markers (Ly6c2, Arg1, Mrc1 and

Nos2). The Cd68$^+$ cells were selected to create a t-distributed stochastic neighbor

embedding (tSNE) plot. We then identified Arg1$^+$ macrophages expressing the surface

markers Mrc1 and Nos2 in the gated dataset to represent AIM and PIM, respectively.

The Arg1$^+$Ly6c2$^+$ monocytes were also identified (Mono). The three terminal clusters

(AIM, PIM, and Mono) could be separated, indicating that the subsets can be

identified experimentally using flow cytometry (**Fig. 5C**). Pseudo-temporal trajectory

(Monocle 2) of the four subsets revealed that monocytes developed into M-DCs and

polarized macrophages (AIM and PIM). Although AIM and PIM expressed distinct

gene signatures, they were highly correlated (**Fig. 5D-E**). The Ctrl samples

contributed to a larger proportion of cells in all subsets (**Fig. 5A**). Therefore, the

scaffold might play a role in the reduction of macrophage infiltration at the

proliferative stage.

Sub-clustering of T cells revealed a novel T cell population and more effector T

cells in the Aligned group

T cells were clustered into four subsets (**Fig. 5F**). T cells characterized by increased

Cd7, Cd3g, and Areg were named as Early T cells (ET) (32). The subset adjacent to

ET expressed elevated Xcl1 and Sult2b1 (associated with T cell activation), and was

named Activated T cell 1 (AT1) (33). AT1 also expressed increased Areg, Ctla2a, and

Ctla2b, genes related to immune homeostasis and immuno-suppression (34, 35).

Another activated T cell subset (AT2) expressed up-regulated genes associated with

cytotoxic T cells (Cd8b1), Th1 cells (Ifng, Ptpn18), and Th17 cells (Il17a, Il17f) (36,

37). Therefore, AT2 might include multiple effector T cells. A novel T cell subset

connecting AT2 and ET was featured by high expression of Birc5, Mki67 and Stmn1,

markers for cell proliferation, and was named Proliferating T cell (PT) (**Fig. 5F-G**). In

GO analysis, both ET and AT1 were enriched in T cell activation. AT1 was also found

elevation of T cell receptor signaling pathway, suggesting that these T cells played a

role in antigen recognition (**Fig. 5G**). We found three terminally differentiated clusters

stemming from two precursors (ET and AT1) in pseudo-time analysis. PT was

differentiated from AT1 and ET after the first branch point, and might be a transitional

status between early T cells and effector T cells. At the second branch, cells

differentiated into two terminal clusters. One belonged to AT1 and ET, and another

one was the AT2 population (**Fig. 5H**). Correlation analysis showed that ET and AT1

were highly correlated (**Fig. 5I**). In the ET and AT1 populations, more cells were from

Ctrl samples, whereas a larger number of cells in AT2 and PT were from aligned

samples.

Receptor-ligand analysis reveals intricate interactions among immune cells,

keratinocytes and fibroblasts

To explore potential interactions among immune cells, keratinocytes and fibroblasts,

we ran CellChat analysis on these datasets (38). For anti-inflammatory macrophages

(AIM), broadly speaking, "pro-inflammatory" signals like Tnf, Visfatin, Rankl, EGFR

and C3a signaling, and "anti-inflammatory" signals including Spp1, Lgals9, Sema3,

Chemerin, Il2, Il13 and Il10 signaling were involved either in a paracrine or autocrine

way. Signals like Mif, Ccl, Csf, Edn, and Cxcl12 signaling exerted protective or

deleterious effects depending on their specific roles (**Fig. 6E**) (25, 39-50). Among T

cell populations (**Fig. 6F**), ET communicated with other T cells mainly through Tgfb

signaling, mediating ET cell tolerance and immune homeostasis (51). The novel PT

population secreted Wnt11 signals that bound to Fzd5 receptor on ET. Inner bulge

layer keratinocytes secreted Cxcl16 that targeted Cxcr6 on ET, chemoattracting ET to

the wound area (52). For fibroblast populations, HDF1 had the most frequent

interaction with other cells (**Fig. 6G**). In addition to the mentioned signals, OSM

signaling was also activated and induced the expression of the pro-inflammatory

cytokine IL-6 (53). T cells and macrophages secreted Tgfb1 and Tgfb2 to target

HDF1 and PF1, leading to proliferation and differentiation of fibroblasts into

myo-fibroblasts (54). Among keratinocytes, IFEB and IB1 had the most frequent

contact with other cells (**Fig. 6H, Fig. S8I-M**). IFEB received Tgfb1 and Tgfb2

(signals playing key roles in regulating epithelial-to-mesenchymal transition (EMT)

for keratinocytes) from T cells, macrophages and germinative layer (GL)

keratinocytes (55). Interactive plots for other types of cells were summarized in Fig.

S8. The communication between immune cells and cutaneous cells was complicated,

and wound healing was the comprehensive results of numerous influencing factors

working together.

Discussion

For this study, we fabricated electrospun membranes with three types of surface

topography (Random, Aligned, and Latticed). The aligned membranes led to faster

wound healing, reduced fibrotic response and enhanced regeneration of cutaneous

appendages compared to other scaffolds. Meanwhile, the aligned membranes

exhibited immunomodulatory properties (**Fig. 6A**). Based on that, we generated

single-cell transcriptomes from wounded mouse skin to investigate the

microenvironment around scaffolds. The skin wound repair process was classically

divided into four phases: hemostasis (hours), inflammation (days), proliferation (1-2

weeks), and remodeling (>2 weeks). 7 days post-wounding was a transitional time

point when the innate immune response subsided and activity of adaptive immune

cells increased (10). In the Ctrl samples, the infiltration of T cells achieved a similar

extent with macrophages (**Fig. 6B** Overall distribution). However, in Aligned group,

infiltrated macrophages were much fewer than T cells, and more terminally

differentiated effector T cells were present (**Fig. 6B**). According to the timeframe of

innate and adaptive immune responses (10), the process of innate immunity seemed to

be alleviated earlier, and adaptive immune response was advanced in the presence of

aligned scaffolds (**Fig. 6C**). In the immune microenvironment around aligned

scaffolds (**Fig. 6D**), damage-associated molecular patterns (DAMPs),

pathogen-associated molecular patterns (PAMPs) and antigens from cell debris,

pathogens, and foreign agents (scaffold) triggered innate and adaptive immune

responses. Neutrophils from circulation (PBN) quickly migrated to the wound area.

The infiltrated neutrophils (Neu1 and Neu2) phagocytosed dying cells and

microorganisms, and secreted chemo-attractants like Ccr1 and Ccl4 to recruit more

leukocytes and lymphoid cells (52). Circulating monocytes (Mono) were also

recruited to the wound area, and differentiated into pro-inflammatory macrophages

(PIM), clearing dying neutrophils and debris. Meanwhile, anti-inflammatory

macrophages (AIM) chemo-attracted more cells via chemokines like Ccl8 and Ccl6.

Part of monocytes differentiated into dendritic cells (M-DC). Tissue resident

Langerhans cells (LC), together with M-DC, functioned as antigen-presenting cells

(APCs). In the antigen specific signal, LC and DC processed antigens into peptides,

and presented them by MHC class II molecules (H2-Aa, H2-Eb1) on the cell surface.

T cells (ET and AT1) bound to the MHC molecules through surface receptors (TCR),

and differentiated into effector T cells ($Cd4^+$ AT2 cells). The co-stimulatory signal was

characterized by the engagement of Cd28 receptor on T cells (AT2, $Cd8b1^+Cd28^{hi}$)

with Cd86 ligands on APCs (Cd86[+] DC and LC) (10, 56). With regard to the

influences that immune microenvironment had on tissue generation, immune cells

sent out a variety of signals (like Grn, Tgfb, Areg/Hbegf) that modulated behaviors of

keratinocytes and fibroblasts (**Fig. 6**). On the other hand, keratinocytes and fibroblasts

secreted chemotactic, pro- or anti-inflammatory signals that regulated immune cell

polarization and function. Immune cells themselves also interact with each other via

numerous signals. The communication network of immune cells and cutaneous cells

was complicated, and wound healing was the comprehensive results of these factors.

Acknowledgement

This work was supported by Research and Develop Program, West China Hospital of

Stomatology Sichuan University (No.LCYJ2019-19); The Fundamental Research

Funds for The Central Universities (No. 2082604401239)

Statement of conflict of interest

There are no conflicts of interest related to this manuscript.

    Figure legends:

373   **Fig. 1 Evaluation of rat skin wound healing implanted with three types of**

374   **electrospun scaffolds at 7 days post-wounding.** (A) Surface topography of three

types of electrospun membranes. (B) Workflow for evaluating rat skin wound healing. (C) Surgical processes for the rat skin excisional wound model. (D, E) Residual wound area at each time point. (F) Histological view of Ctrl group and groups implanted with three types of scaffolds. (G) Semi-quantitative evaluation of gap width. (H) Immunofluorescent staining using Krt5 (red) and Krt10 (green). (I) Semi-quantitative evaluation of the fluorescent area. (J) Evaluation of foreign body reaction around biomaterials. FBGCs lined up on Random and Latticed membranes, whereas fewer of them appeared on Aligned ones. (K) Thickness of fibrotic capsules. (L-N) PCA and GO analysis revealed immunomodulatory effects for aligned and random scaffolds. ****$P < 0.0001$, **$P < 0.01$, and *$P < 0.05$ by ANOVA for data in (E), (G) and (K).

Fig. 2 Evaluation of mouse skin wound healing implanted with three types of

electrospun scaffolds at 7 days post-wounding. (A) Workflow for evaluating mouse

skin wound healing. (B) Surgical processes for mouse skin excisional wound model.

(C, D) Residual wound area at each time point. (E) Histological analysis of Ctrl group

and groups implanted with three types of scaffolds. (F) Semi-quantitative evaluation

of gap width. (G) Evaluation of foreign body reaction around biomaterials. (H)

Thickness of fibrotic capsules. (I, J) GO and KEGG analysis revealed

immunomodulatory effects for aligned scaffolds. ****$P < 0.0001$, ***$P < 0.001$, and

*$P < 0.05$ by ANOVA for data in (D), (F) and (H).

Fig. 3 Overview of the single-cell transcriptome analysis. (A) Workflow for single-cell experiment. (B) Cells were categorized into thirteen main classes. Cell populations in each class, number of cells (%), and composition of aligned and Ctrl cells were listed. (C) Marker genes for different cell classes. (D) Gene expression differences between Ctrl and Aligned groups.

Fig. 4 Further analysis of keratinocytes, fibroblasts, neutrophils and dendritic

cells. (A) Sub-clustering of keratinocytes revealed twelve subsets. Marker genes and

composition for each subset was listed. (B) Marker genes for each keratinocyte subset.

(C) Location of different keratinocyte subsets. Color of each cell corresponds with its

cell type in (A). (D) Sub-clustering of fibroblasts revealed eight subsets. The marker

genes and composition for each subset was listed. (E) Sub-clustering of neutrophils

revealed three subsets. (F) Marker genes for neutrophil subsets and their enriched

gene sets in GO analysis. (G) Correlation analysis of neutrophil subsets. (H)

Sub-clustering of dendritic cells revealed two subsets. (I) Marker genes for dendritic

cell subsets. (J) Expression of genes associated with antigen presenting in dendritic

cells.

Fig. 5

Fig. 5 Further analysis of macrophages and T cells. (A) Sub-clustering of

macrophages revealed four subsets. Marker genes and composition for each subset

was listed. (B) Marker genes for each macrophage/monocyte subset and their enriched

gene sets in GO analysis. Expression of canonical M1 and M2 markers and other

proposed markers were shown. (C) In vivo flow cytometry strategy using Cd68 to

identify monocytes and macrophages. Arg1+ macrophages expressing the surface

markers Mrc1 and Nos2 in the gated dataset represented AIM and PIM, respectively.

The Arg1+Ly6c2+ monocytes were also identified (Mono). (D) Pseudo-temporal

ordering of macrophage/monocytes, and the distribution of four subsets along the

trajectory. (E) Correlation analysis of macrophage/monocyte subsets. (F)

Sub-clustering of T cells revealed four subsets. (G) Marker genes for T cell subsets

and their enriched gene sets in GO analysis. AT2 subset expressed gene signatures of

cytotoxic T cells, Th1 cells, and Th17 cells. (H) Pseudo-temporal ordering of T cells,

and the distribution of four subsets along the trajectory. (I) Correlation analysis of T

cell subsets.

Fig. 6 Summary of the differences in wound healing and immune responses between the Aligned group and Ctrl group. (A) The aligned membranes had immunomodulatory properties, and led to faster wound healing, reduced fibrotic

response and enhanced regeneration of skin appendages. (B) Pie plot for overall

immune cells in two groups. For each cell type, differences in cell amount between

aligned and Ctrl groups. (C) Timeframe of innate and adaptive immune responses

occurred in Aligned and Ctrl groups. (D) The immune microenvironment around

aligned scaffolds. (E) Cell-cell interaction plots according to receptor-ligand analysis

via CellChat.

References

1. X. Li, B. Cho, R. Martin, M. Seu, C. Zhang, Z. Zhou, J. S. Choi, X. Jiang, L. Chen, G. Walia, J. Yan, M. Callanan, H. Liu, K. Colbert, J. Morrissette-McAlmon, W. Grayson, S. Reddy, J. M. Sacks, H. Q. Mao, Nanofiber-hydrogel composite-mediated angiogenesis for soft tissue reconstruction. *Sci Transl Med* **11**, eaau6210 (2019).

2. A. Mandal, A. V. Boopathy, L. K. W. Lam, K. D. Moynihan, M. E. Welch, N. R. Bennett, M. E. Turvey, N. Thai, J. H. Van, J. C. Love, P. T. Hammond, D. J. Irvine, Cell and fluid sampling microneedle patches for monitoring skin-resident immunity. *Sci Transl Med* **10**, eaar2227 (2018).

3. G. Huang, F. Li, X. Zhao, Y. Ma, Y. Li, M. Lin, G. Jin, T. J. Lu, G. M. Genin, F. Xu, Functional and Biomimetic Materials for Engineering of the Three-Dimensional Cell Microenvironment. *Chem Rev* **117**, 12764-12850 (2017).

4. H. Li, Y. Xu, H. Xu, J. Chang, Electrospun membranes: control of the structure and structure related applications in tissue regeneration and drug delivery. *J Mater Chem B* **2**, 5492-5510 (2014).

5. Y. Wang, H. Shi, J. Qiao, Y. Tian, M. Wu, W. Zhang, Y. Lin, Z. Niu, Y. Huang, Electrospun tubular scaffold with circumferentially aligned nanofibers for regulating smooth muscle cell growth. *ACS Appl Mater Interfaces* **6**, 2958-2962 (2014).

6. A. Martins, M. L. Alves da Silva, S. Faria, A. P. Marques, R. L. Reis, N. M.

Neves, The influence of patterned nanofiber meshes on human mesenchymal stem cell osteogenesis. *Macromol Biosci* **11**, 978-987 (2011).

7. K. T. Kurpinski, J. T. Stephenson, R. R. Janairo, H. Lee, S. Li, The effect of fiber alignment and heparin coating on cell infiltration into nanofibrous PLLA scaffolds. *Biomaterials* **31**, 3536-3542 (2010).

8. C. Chu, L. Liu, S. Rung, Y. Wang, Y. Ma, C. Hu, X. Zhao, Y. Man, Y. Qu, Modulation of foreign body reaction and macrophage phenotypes concerning microenvironment. *J Biomed Mater Res A* **108**, 127-135 (2020).

9. K. Sadtler, A. Singh, M. T. Wolf, X. Wang, D. M. Pardoll, J. H. Elisseeff, Design, clinical translation and immunological response of biomaterials in regenerative medicine. *Nat Rev Mater* **1**, 16040 (2016).

10. M. O. Dellacherie, B. R. Seo, D. J. Mooney, Macroscale biomaterials strategies for local immunomodulation. *Nat Rev Mater* **4**, 379-397 (2019).

11. L. Chung, D. R. Maestas, Jr., A. Lebid, A. Mageau, G. D. Rosson, X. Wu, M. T. Wolf, A. J. Tam, I. Vanderzee, X. Wang, J. I. Andorko, H. Zhang, R. Narain, K. Sadtler, H. Fan, D. Cihakova, C. J. Le Saux, F. Housseau, D. M. Pardoll, J. H. Elisseeff, Interleukin 17 and senescent cells regulate the foreign body response to synthetic material implants in mice and humans. *Sci Transl Med* **12**, eaax3799 (2020).

12. S. D. Sommerfeld, C. Cherry, R. M. Schwab, L. Chung, D. R. Maestas, P. Laffont, J. E. Stein, A. Tam, S. Ganguly, F. Housseau, J. M. Taube, D. M. Pardoll, P. Cahan, J. H. Elisseeff, Interleukin-36 gamma-producing

490   macrophages drive IL-17-mediated fibrosis. *Sci Immunol* **4**, eaax4783 (2019).

491   13.   T. Stuart, A. Butler, P. Hoffman, C. Hafemeister, E. Papalexi, W. M. Mauck,

492   3rd, Y. Hao, M. Stoeckius, P. Smibert, R. Satija, Comprehensive Integration of

493   Single-Cell Data. *Cell* **177**, 1888-1902 e1821 (2019).

494   14.   H. Wu, S. Han, B. Wu, X. Du, Z. Sheng, J. Lin, X. Chen, K. Zhao, V.

495   Bunpetch, Y. Chen, M. Zeng, E. V. Alakpa, Y. Ma, X. Lei, J. Huang, X. Zou, H.

496   Ouyang, Single-cell mass cytometry reveals in vivo immunological response

497   to surgical biomaterials. *Appl Mater Today* **16**, 169-178 (2019).

498   15.   I. Pastar, O. Stojadinovic, N. C. Yin, H. Ramirez, A. G. Nusbaum, A. Sawaya,

499   S. B. Patel, L. Khalid, R. R. Isseroff, M. Tomic-Canic, Epithelialization in

500   Wound Healing: A Comprehensive Review. *Adv Wound Care (New Rochelle)*

501   **3**, 445-464 (2014).

502   16.   R. Tokunaga, W. Zhang, M. Naseem, A. Puccini, M. D. Berger, S. Soni, M.

503   McSkane, H. Baba, H. J. Lenz, CXCL9, CXCL10, CXCL11/CXCR3 axis for

504   immune activation - A target for novel cancer therapy. *Cancer Treat Rev* **63**,

505   40-47 (2018).

506   17.   P. Myung, M. Ito, Dissecting the bulge in hair regeneration. *J Clin Invest* **122**,

507   448-454 (2012).

508   18.   S. Joost, K. Annusver, T. Jacob, X. Sun, T. Dalessandri, U. Sivan, I. Sequeira,

509   R. Sandberg, M. Kasper, The Molecular Anatomy of Mouse Skin during Hair

510   Growth and Rest. *Cell Stem Cell* **26**, 441-457.e447 (2020).

511   19.   C. Philippeos, S. B. Telerman, B. Oules, A. O. Pisco, T. J. Shaw, R. Elgueta, G.

512   Lombardi, R. R. Driskell, M. Soldin, M. D. Lynch, F. M. Watt, Spatial and

513   Single-Cell Transcriptional Profiling Identifies Functionally Distinct Human

514   Dermal Fibroblast Subpopulations. *J Invest Dermatol* **138**, 811-825 (2018).

515 20.   T. Tabib, C. Morse, T. Wang, W. Chen, R. Lafyatis, SFRP2/DPP4 and

516   FMO1/LSP1 Define Major Fibroblast Populations in Human Skin. *J Invest*

517   *Dermatol* **138**, 802-810 (2018).

518 21.   C. F. Guerrero-Juarez, P. H. Dedhia, S. Jin, R. Ruiz-Vega, D. Ma, Y. Liu, K.

519   Yamaga, O. Shestova, D. L. Gay, Z. Yang, K. Kessenbrock, Q. Nie, W. S. Pear,

520   G. Cotsarelis, M. V. Plikus, Single-cell analysis reveals fibroblast

521   heterogeneity and myeloid-derived adipocyte progenitors in murine skin

522   wounds. *Nat Commun* **10**, 650 (2019).

523 22.   W. M. Nauseef, N. Borregaard, Neutrophils at work. *Nat Immunol* **15**, 602-611

524   (2014).

525 23.   K. Takahara, Y. Omatsu, Y. Yashima, Y. Maeda, S. Tanaka, T. Iyoda, B. E.

526   Clausen, K. Matsubara, J. Letterio, R. M. Steinman, Y. Matsuda, K. Inaba,

527   Identification and expression of mouse Langerin (CD207) in dendritic cells.

528   *Int Immunol* **14**, 433-444 (2002).

529 24.   Q. Su, B. Z. Igyarto, Keratinocytes Share Gene Expression Fingerprint with

530   Epidermal Langerhans Cells via mRNA Transfer. *J Invest Dermatol* **139**,

531   2313-2323 e2318 (2019).

532 25.   S. S. Bohlson, S. D. O'Conner, H. J. Hulsebus, M. M. Ho, D. A. Fraser,

533   Complement, c1q, and c1q-related molecules regulate macrophage

polarization. *Front Immunol* **5**, 402 (2014).

26.   S. A. Islam, D. S. Chang, R. A. Colvin, M. H. Byrne, M. L. McCully, B.

Moser, S. A. Lira, I. F. Charo, A. D. Luster, Mouse CCL8, a CCR8 agonist,

promotes atopic dermatitis by recruiting IL-5+ T(H)2 cells. *Nat Immunol* **12**,

167-177 (2011).

27.   M. Carrion, S. Perez-Garcia, C. Martinez, Y. Juarranz, L. Estrada-Capetillo, A.

Puig-Kroger, R. P. Gomariz, I. Gutierrez-Canas, VIP impairs acquisition of the

macrophage proinflammatory polarization profile. *J Leukoc Biol* **100**,

1385-1393 (2016).

28.   J. E. Kanter, F. Kramer, S. Barnhart, M. M. Averill, A. Vivekanandan-Giri, T.

Vickery, L. O. Li, L. Becker, W. Yuan, A. Chait, K. R. Braun, S. Potter-Perigo,

S. Sanda, T. N. Wight, S. Pennathur, C. N. Serhan, J. W. Heinecke, R. A.

Coleman, K. E. Bornfeldt, Diabetes promotes an inflammatory macrophage

phenotype and atherosclerosis through acyl-CoA synthetase 1. *Proc Natl Acad*

Sci U S A **109**, E715-724 (2012).

29.   M. Clement, G. Basatemur, L. Masters, L. Baker, P. Bruneval, T. Iwawaki, M.

Kneilling, S. Yamasaki, J. Goodall, Z. Mallat, Necrotic Cell Sensor Clec4e

Promotes a Proatherogenic Macrophage Phenotype Through Activation of the

Unfolded Protein Response. *Circulation* **134**, 1039-1051 (2016).

30.   A. M. Soler-Rodriguez, H. Zhang, H. S. Lichenstein, N. Qureshi, D. W. Niesel,

S. E. Crowe, J. W. Peterson, G. R. Klimpel, Neutrophil activation by bacterial

lipoprotein versus lipopolysaccharide: differential requirements for serum and

CD14. *J Immunol* **164**, 2674-2683 (2000).

31.   L. Grosche, C. Drassner, P. Muhl-Zurbes, L. Kamm, V. T. K. Le-Trilling, M.

558   Trilling, A. Steinkasserer, C. S. Heilingloh, Human Cytomegalovirus-Induced

559   Degradation of CYTIP Modulates Dendritic Cell Adhesion and Migration.

560   *Front Immunol* **8**, 461 (2017).

561   32.   B. Gerby, E. Clappier, F. Armstrong, C. Deswarte, J. Calvo, S. Poglio, J.

562   Soulier, N. Boissel, T. Leblanc, A. Baruchel, J. Landman-Parker, P. H. Roméo,

563   P. Ballerini, F. Pflumio, Expression of CD34 and CD7 on human T-cell acute

564   lymphoblastic leukemia discriminates functionally heterogeneous cell

565   populations. *Leukemia* **25**, 1249-1258 (2011).

566   33.   S. J. Bensinger, M. N. Bradley, S. B. Joseph, N. Zelcer, E. M. Janssen, M. A.

567   Hausner, R. Shih, J. S. Parks, P. A. Edwards, B. D. Jamieson, P. Tontonoz,

568   LXR signaling couples sterol metabolism to proliferation in the acquired

569   immune response. *Cell* **134**, 97-111 (2008).

570   34.   D. M. W. Zaiss, W. C. Gause, L. C. Osborne, D. Artis, Emerging functions of

571   amphiregulin in orchestrating immunity, inflammation, and tissue repair.

572   *Immunity* **42**, 216-226 (2015).

573   35.   B. Mahata, X. Zhang, A. A. Kolodziejczyk, V. Proserpio, L. Haim-Vilmovsky,

574   A. E. Taylor, D. Hebenstreit, F. A. Dingler, V. Moignard, B. Gottgens, W. Arlt,

575   A. N. McKenzie, S. A. Teichmann, Single-cell RNA sequencing reveals T

576   helper cells synthesizing steroids de novo to contribute to immune

577   homeostasis. *Cell Rep* **7**, 1130-1142 (2014).

578   36.   P. Castro-Sanchez, R. Ramirez-Munoz, P. Roda-Navarro, Gene Expression
579   Profiles of Human Phosphotyrosine Phosphatases Consequent to Th1
580   Polarisation and Effector Function. *J Immunol Res* **2017**, 8701042 (2017).

581   37.   I. C. Boothby, J. N. Cohen, M. D. Rosenblum, Regulatory T cells in skin
582   injury: At the crossroads of tolerance and tissue repair. *Sci Immunol* **5**,
583   eaaz9631 (2020).

584   38.   S. Jin, C. F. Guerrero-Juarez, L. Zhang, I. Chang, P. Myung, M. V. Plikus, Q.
585   Nie, Inference and analysis of cell-cell communication using CellChat.
586   *bioRxiv*,   (2020).

587   39.   M. Croft, The role of TNF superfamily members in T-cell function and
588   diseases. *Nat Rev Immunol* **9**, 271-285 (2009).

589   40.   J. D. Klement, A. V. Paschall, P. S. Redd, M. L. Ibrahim, C. Lu, D. Yang, E.
590   Celis, S. I. Abrams, K. Ozato, K. Liu, An osteopontin/CD44 immune
591   checkpoint controls CD8+ T cell activation and tumor immune evasion. *J Clin
592   Invest* **128**, 5549-5560 (2018).

593   41.   C. Wu, T. Thalhamer, R. F. Franca, S. Xiao, C. Wang, C. Hotta, C. Zhu, M.
594   Hirashima, A. C. Anderson, V. K. Kuchroo, Galectin-9-CD44 interaction
595   enhances stability and function of adaptive regulatory T cells. *Immunity* **41**,
596   270-282 (2014).

597   42.   M. Hayashi, T. Nakashima, M. Taniguchi, T. Kodama, A. Kumanogoh, H.
598   Takayanagi, Osteoprotection by semaphorin 3A. *Nature* **485**, 69-74 (2012).

43.   D. S. Shouval, A. Biswas, J. A. Goettel, K. McCann, E. Conaway, N. S. Redhu,

I. D. Mascanfroni, Z. Al Adham, S. Lavoie, M. Ibourk, D. D. Nguyen, J. N.

Samsom, J. C. Escher, R. Somech, B. Weiss, R. Beier, L. S. Conklin, C. L.

Ebens, F. G. Santos, A. R. Ferreira, M. Sherlock, A. K. Bhan, W. Muller, J. R.

Mora, F. J. Quintana, C. Klein, A. M. Muise, B. H. Horwitz, S. B. Snapper,

Interleukin-10 receptor signaling in innate immune cells regulates mucosal

immune tolerance and anti-inflammatory macrophage function. *Immunity* **40**,

706-719 (2014).

44.   S. S. Jankauskas, D. W. L. Wong, R. Bucala, S. Djudjaj, P. Boor, Evolving

complexity of MIF signaling. *Cell Signal* **57**, 76-88 (2019).

45.   W. Lin, D. Xu, C. D. Austin, P. Caplazi, K. Senger, Y. Sun, S. Jeet, J. Young, D.

Delarosa, E. Suto, Z. Huang, J. Zhang, D. Yan, C. Corzo, K. Barck, S. Rajan,

C. Looney, V. Gandham, J. Lesch, W. C. Liang, E. Mai, H. Ngu, N. Ratti, Y.

Chen, D. Misner, T. Lin, D. Danilenko, P. Katavolos, E. Doudemont, H. Uppal,

J. Eastham, J. Mak, P. E. de Almeida, K. Bao, A. Hadadianpour, M. Keir, R. A.

D. Carano, L. Diehl, M. Xu, Y. Wu, R. M. Weimer, J. DeVoss, W. P. Lee, M.

Balazs, K. Walsh, K. R. Alatsis, F. Martin, A. A. Zarrin, Function of CSF1 and

IL34 in Macrophage Homeostasis, Inflammation, and Cancer. *Front Immunol*

10, 2019 (2019).

46.   S. F. Nie, L. F. Zha, Q. Fan, Y. H. Liao, H. S. Zhang, Q. W. Chen, F. Wang, T.

T. Tang, N. Xia, C. Q. Xu, J. Y. Zhang, Y. Z. Lu, Z. P. Zeng, J. Jiao, Y. Y. Li, T.

Xie, W. J. Zhang, D. Wang, C. C. Wang, J. J. Fa, H. B. Xiong, J. Ye, Q. Yang,

P. Y. Wang, S. H. Tian, Q. L. Lv, Q. X. Li, J. Qian, B. Li, G. Wu, Y. X. Wu, Y.

Yang, X. P. Yang, Y. Hu, Q. K. Wang, X. Cheng, X. Tu, Genetic Regulation of

the Thymic Stromal Lymphopoietin (TSLP)/TSLP Receptor (TSLPR) Gene

Expression and Influence of Epistatic Interactions Between IL-33 and the

TSLP/TSLPR Axis on Risk of Coronary Artery Disease. *Front Immunol* **9**,

1775 (2018).

47.   S. Avniel, Z. Arik, A. Maly, A. Sagie, H. B. Basst, M. D. Yahana, I. D. Weiss,

B. Pal, O. Wald, D. Ad-El, N. Fujii, F. Arenzana-Seisdedos, S. Jung, E. Galun,

E. Gur, A. Peled, Involvement of the CXCL12/CXCR4 pathway in the

recovery of skin following burns. *J Invest Dermatol* **126**, 468-476 (2006).

48.   A. Stofkova, Resistin and visfatin: regulators of insulin sensitivity,

inflammation and immunity. *Endocr Regul* **44**, 25-36 (2010).

49.   D. M. Hardbower, K. Singh, M. Asim, T. G. Verriere, D. Olivares-Villagomez,

D. P. Barry, M. M. Allaman, M. K. Washington, R. M. Peek, Jr., M. B.

Piazuelo, K. T. Wilson, EGFR regulates macrophage activation and function in

bacterial infection. *J Clin Invest* **126**, 3296-3312 (2016).

50.   M. C. Walsh, Y. Choi, Biology of the RANKL-RANK-OPG System in

Immunity, Bone, and Beyond. *Front Immunol* **5**, 511 (2014).

51.   J. C. Marie, D. Liggitt, A. Y. Rudensky, Cellular mechanisms of fatal

early-onset autoimmunity in mice with the T cell-specific targeting of

transforming growth factor-beta receptor. *Immunity* **25**, 441-454 (2006).

52.   S. K. Bromley, T. R. Mempel, A. D. Luster, Orchestrating the orchestrators:

chemokines in control of T cell traffic. *Nat Immunol* **9**, 970-980 (2008).

53. B. Le Goff, S. Singbrant, B. A. Tonkin, T. J. Martin, E. Romas, N. A. Sims, N. C. Walsh, Oncostatin M acting via OSMR, augments the actions of IL-1 and TNF in synovial fibroblasts. *Cytokine* **68**, 101-109 (2014).

54. V. Miguel, O. Busnadiego, M. Fierro-Fernandez, S. Lamas, Protective role for miR-9-5p in the fibrogenic transformation of human dermal fibroblasts. *Fibrogenesis Tissue Repair* **9**, 7 (2016).

55. S. Liarte, A. Bernabe-Garcia, F. J. Nicolas, Human Skin Keratinocytes on Sustained TGF-beta Stimulation Reveal Partial EMT Features and Weaken Growth Arrest Responses. *Cells* **9**, 255 (2020).

56. M. L. Kapsenberg, Dendritic-cell control of pathogen-driven T-cell polarization. *Nat Rev Immunol* **3**, 984-993 (2003).

Materials and methods

Electrospinning of polymer scaffolds

We used poly(lactic-co-glycolic acid) (PLGA) (LA/GA = 75:25, Mw = 105 kDa, dispersity is 1.897) produced by Jinan Daigang Biomaterial Co., Ltd. (Shandong, China) and FC (from fish scale and skin) obtained from Sangon Biotech Co., Ltd. (Shanghai, China) to fabricate scaffolds by electrospinning. The PLGA (20% w/v) and FC (2% w/v) solution (dissolved in 1,1,1,3,3,3-Hexafluoro-2-propanol (HFIP) solvent (Aladdin Co., Ltd. (Shanghai, China)) were loaded into a plastic syringe fitted with a flat-tipped 21G needle (inner diameter=0.5mm). A high voltage of 7kV and a distance of 16 cm were applied between the needle and the collector. For randomly oriented fibers (Random group), the electrostatically charged fiber was ejected toward the

grounded flat collector in the high electric field, forming a membrane deposited on the aluminium foil. For the Latticed group, an electroconductive chess-like wire net was used as the collector. For the Aligned group, the rotational speed of the collecting drum is set at 2800rpm. Finalized scaffolds were approximately 30–90 μm in thickness. To crosslink FC, the membranes were immersed in 50 mM of EDS/NHS and 10 mM of MES ethanol solution for 24 h at 4 °C. Then membranes were washed three times with 75% ethanol and dried in vacuum oven for 24 h. Subsequently, the prepared membranes were sterilized by γ-irradiation for in vitro and in vivo experiments.

Characterization of scaffolds

Scanning electron microscopy (SEM; JEOL, JSM-6510LV, Japan) was employed to observe the surface morphology of the electrospun membranes. Image-Pro Plus was applied to quantitatively measure the fiber diameter and distribution from the SEM images obtained. The surface wetting behavior of the membranes were characterized by measuring the water contact angles (Chengde Dingsheng, JY-82B, China). Five samples were tested for each type of membrane to obtain an average value. The tensile properties of the membranes were tested under a constant upper clamp at speed of 15 mm/min. All tensile tests follow the criteria of "Plastics-Determination of tensile properies of films" (GB/T 1040.3–2006, corresponding with ISO 1184–1983).

Cell culture and cell viability test

L929 mouse fibroblast cells and Human oral keratinocytes (HOK) were used for viability tests. Cells were cultured in medium containing RPMI 1640 medium

(HyClone) supplemented with 10% fetal bovine serum (Gibco), and were kept at 37□

in humidified 5% CO2/95% air. The cell viability was determined by Cell Counting

Kit-8 (CCK-8, Dojindo Laboratories, Kumamoto, Japan). Electrospun membranes

were cut into squares (edge length=5mm) and placed in the bottom of 96-well plates

(n=3 for each group). L929 cells and HOK cells were seeded onto membranes at

$4*10^4$ cells/ml. Cells were co-cultured with membranes for 1, 3, and 5 days. Blank

wells seeded with equal amount of cells were used as Ctrl. 10μl CCK-8 solution was

added to each well, and the plates were incubated at 37 °C for 1h. After incubation,

the Abs at 450nm was measured to determine the cell viability using a micro-plate

reader (Multiskan, Thermo, USA).

Experimental model

Excisional wound model

The protocol of the present experiment was approved by Institution Review Board of

West China Hospital of Stomatology (No.WCHSIRB-D-2017-033-R1) Animals

included Sprague Dawley male rat at ages from 7 to 8 weeks and C57BL/6 male mice

at ages from 7 to 9 weeks (Chengdu Dossy Experimental Animals Co., LTD.). Hair at

the surgical area was removed. Full-thickness circular excisional wound

(diameter=6mm) was created at the dorsal skin of rats/mice. Random, aligned and

latticed electrospun scaffolds were trimmed into circular shape (diameter=8mm), and

placed below the wound. The Control group did not receive any implants. A sterile

Tegaderm film (3M) was placed above the wound to protect the wound area. Then

annular silicone splints (inner diameter=8mm, outer diameter=12mm) were sutured

with the Tegaderm film and underlying skin in order to minimize the contraction of

the dorsal muscle. After healing for 1, 2 and 4 weeks, animals were euthanized for

sample harvest. Using the residual wound as center, a round skin sample (diameter

=10mm) containing all the layers of skin was harvested.

Model for subcutaneous implant placement

The surgical area on dorsal skin was shaved and aseptically prepared. Three

horizontal incisions of approximately 10 mm were made and subcutaneous pockets

were created for membrane implantation. Then random, align and lattice scaffolds

were placed into the pockets. After implantation, the incisions were sutured with

interrupted sutures. After recovering for 3, 7, and 14 days, samples including scaffolds

and the whole layer of skin at surgical sites were together harvested.

Specimen harvest for scRNA-seq

We obtained skin samples by cutting off skin at the wound area (circular,

diameter=10mm). Subcutaneous tissues were removed. A total of four samples were

harvested in each group. The tissues were washed in a 100 mm petri dish containing

20 ml of phosphate-buffered saline (PBS). Then they were transferred to a 50 mm

petri dish containing 100μL of Enzyme G (Epidermis Dissociation Kit mouse,

Miltenyi) and 3.9 ml of PBS buffer with the dermal side facing downwards. Tissues

were digested for 16 hours at 4°C. Then they were transferred into a 50 mm petri dish

containing 4mL of 1×Buffer S (Miltenyi). Epidermis was peeled off from the skin

using tweezers, and was cut into pieces. Enzyme mix containing 3.9 ml of 1×Buffer S,

100 μl of Enzyme P, and 20 μl of Enzyme A (Miltenyi) stored in a gentleMACS™ C

Tube was used to digest the epidermis pieces for 20 minutes at 37□. Then 4 ml of

PBS that contained 0.5% bovine serum albumin (BSA) was added. A gentleMACS

Dissociator (Miltenyi) was applied to automatically dissociate the epidermis (Program

B). The sample was passed through a 70μm cell strainer (Corning), centrifuged at

300×g for 10 minutes at room temperature, and resuspended with PBS that contained

0.5% BSA. Cells were gently washed twice and stored in an ice box. For the dermis

part, they were first cut into pieces (diameter < 1mm). The tissue was mixed with

10ml enzyme mix containing Type I Collagenase (3125u/mL) (Gibco) and 2.5ml

trypsin (Gibco), and poured into a gentleMACS™ C Tube. After dissociating the

tissue on gentleMACS Dissociator for 37s (Skin mode), another 10ml enzyme mix

was added. The sample was digested for 2.5 hours at 37□ in a rotary machine

(Peqlab). Then the dermis sample was passed through a 70μm cell strainer (Corning),

centrifuged at 300×g for 5 minutes at room temperature, and resuspended with 3ml

red blood cell lysis buffer (Solarbio). After 3 minutes, the cell suspension was

centrifuged and gently resuspended with RPMI 1640 medium (Hyclone). Cells were

gently washed twice with PBS containing 0.5% BSA and stored in an ice box. The

epidermis and dermis cell solutions were mixed together as a whole. The sample was

centrifuged, and resuspended with 100 μl Dead Cell Removal MicroBeads (Miltenyi).

After incubation for 15min at room temperature, the cell suspension was diluted in

3ml 1×Binding buffer (Miltenyi). LS columns (Miltenyi) and a magnetic stand

(Miltenyi) were used for removal of dead cells and debris. The negatively selected

live cells passed through the column, and were resuspended with PBS containing 0.05%

BSA. Finally, we proceeded with the 10x Genomics® Single Cell Protocol.

Single-cell encapsulation and library generation

Single cells were encapsulated in water-in-oil emulsion along with gel beads coated

with unique molecular barcodes using the 10× Genomics Chromium Single-Cell

Platform. For single-cell RNA library generation, the manufacturers' protocol was

performed. (10×Single Cell 3' v3) Sequencing was performed using Illumina 1.9

mode with 94574 reads per cell.

Quality control

We use fastp to analyze basic statistics on the quality of the raw reads.

If clean reads is indispensable, then raw read sequences produced by the Illumina

pipeline in FASTQ format were pre-processed through Trimmomatic software which

can be summarized as below:

(1)Remove low-quality reads: scan the reads with a 4-base wide sliding window, cut

when the average quality per base drops below 10 (SLIDINGWINDOW: 4:10)

(2)Remove trailing low quality or N bases (below quality 3) (TRAILING: 3)

(3)Remove adapters: there are two modes to remove the adapter sequence: a.

alignment with the adapter sequence, the number of matching bases was greater than

7 and mismatch=2; b. when read1 and read2 overlapping base scoring was greater

than 30, we removed non-overlapping portions (ILLUMINACLIP:adapter.fa:2:30:7)

(4)Drop reads below the 26 bases long

(5)Discard those reads that did not form pairs.

The remaining reads that passed all the filtering steps was counted as clean reads and

all subsequent analyses were based on this.

Generation and Analysis of Single-Cell Transcriptomes

Raw reads were demultiplexed and mapped to the reference genome by 10X

Genomics Cell Ranger pipeline

(https://support.10xgenomics.com/single-cell-gene-expression/software/pipelines/late

st/what-is-cell-ranger) using default parameters. All downstream single-cell analyses

were performed using Seurat. In brief, for each gene and each cell barcode (filtered by

CellRanger3.0), unique molecule identifiers were counted to construct digital

expression matrices. Secondary filtration by Seurat: A gene with expression in more

than 3 cells was considered as expressed, and each cell was required to have at least

200 expressed genes. We also filtered out some of the foreign cells.

The Seurat package was used in data normalization, dimension reduction, clustering,

and differential expression. We used the Seurat alignment method canonical

correlation analysis (CCA) [Nat. Biotechnol. 36, 411–420 (2018).] for integrated

analysis of datasets. For clustering, highly variable genes were selected and the

principal components based on those genes used to build a graph, which was

segmented with a resolution of 0.6.

RNA-seq analysis

The bulk tissue RNA-seq analysis was carried out by Novogene Corporation (Beijing,

China). RNA was extracted from tissues or cells using standard methods to make sure

samples were strictly controlled for quality. In terms of library construction and

quality control, mRNA can be obtained in two main ways: firstly, most eukaryotes'

mRNA has poly A-tailed structural, and poly A-tailed mRNA can be enriched by

Oligo (dT) magnetic beads. The other is the removal of ribosomal RNA from the total

RNA to obtain mRNA. Subsequently, the obtained mRNA was randomly interrupted

by divalent cations in NEB Fragmentation Buffer, and the database was constructed

according to the NEB general database construction method or chain specific database

construction method. Upon completion of library construction, a Qubit2.0

Fluorometer was used for initial quantification, and the library was diluted to 1.5ng/ul.

Then the insert size of the library was detected using Agilent 2100 bioanalyzer. After

the insert size met the expectation, the effective concentration of the library was

accurately quantified by qRT-PCR (the effective concentration of the library higher

than 2nM) to ensure library quality. Finally, the libraries were qualified for

sequencing, and Illumina sequencing was performed after pooling the different

libraries according to the requirements of effective concentration and target data

volume, of which the basic principle is Sequencing by Synthesis. Through

z-transformation of Fragments Per Kilobase of transcript per Million mapped reads

(fpkm) of the selected gene, gene expression was analyzed. Sample size for

conventional, bulk RNA-Seq libraries was fixed at 3 biological replicates.

Quantitative Real-Time Polymerase Chain Reaction (qPCR).

The harvested samples were cut into pieces, and homogenated in TRIzol™ Reagent

(Cat. #15596026, Invitrogen, Thermo Scientific). Concentration and ratio of total

RNA were detected by NanoPhotometer NP80 (Implen, Westlake Village, CA) at

wavelength of 260 nm and 280 nm. The cDNAs were synthesized using

PrimeScript™ RT reagent Kit with gDNA Eraser (Perfect Real Time) (Cat.

#RR047A), then amplified by qPCR with the specific primers (Table 1). PCR was

performed on QuantStudio 3 Real-Time PCR Systems (ThermoFisher Scientific,

Waltham, MA). Each 20 μL of PCR mixture contained 10μl of TB Green Premix Ex

Taq (Tli RNaseH Plus, Takara) (2X), 0.4μL of PCR Forward Primer (10μM), 0.4μl of

PCR Reverse Primer (10μM), 0.4μl of ROX Reference Dye (50X), 2μl Template and

6.8μl of Sterile purified water. Samples were incubated at 1 cycle of 95□ for 30 s

followed by 40 cycles of 95□ for 5 s and 60□ for 34 s, and ended up with a cycle

composing of 95□ for 15 s, 60□ for 1 min and 95□ for 15 s. Results were analyzed

using the comparative CT (ΔΔCT) method to calculate gene expression fold changes

normalized to the levels of Gapdh gene transcripts. The experiments were repeated for

three times independently (n=3).

Primer	Forward sequence	Primer	Reverse sequence
Gapdh forward	AGTGCCAGCCTCGTCTCATA	Gapdh reverse	GATGGTGATGGGTTTCCCGT
Cd68 forward	TGGGGCCTCTCTGTATTG	Cd68 reverse	TCTGATGTCGGTCCTGTTT
Tgm2 forward	CGGAAGGACACGGTTAGG	Tgm2 reverse	GTATGGGACCAGGAAACCC
Cd4 forward	AACCAAAAGGTGTCTGTGC	Cd4 reverse	ACCAGCAAACTGAAGGGA
Cd3e forward	TCTCCATCTCAGGAACCAG	Cd3e reverse	CAGGCAGGACTTTGTCATT

Table1 TB Green rat mRNA primers

Fluorescence activated Cell Sorting (FACS) analysis

The surface markers of macrophages and their phenotypes were examined by flow

cytometry to evaluate proportion and polarization of macrophages.

The in vivo specimens were first cut into pieces (diameter < 1mm). The tissue was

mixed with 10ml enzyme mix containing Type I Collagenase (3125u/mL) (Gibco) and

2.5ml trypsin (Gibco), and poured into a gentleMACS™ C Tube. After dissociating

the tissue on gentleMACS Dissociator for 37s (Skin mode), another 10ml enzyme mix

was added. The sample was digested for 2.5-3 hours at 37□ in a rotary machine

(Peqlab). Then sample was passed through a 70μm cell strainer (Corning), centrifuged

at 300×g for 5 minutes at room temperature. Cells were gently washed twice with

PBS containing 0.05% BSA and stored in an ice box. Then the cell solutions were

co-incubated with antibodies against iNOS (PE, NBP2-22119PE, Novus), CD68

(FITC, ab134351, Abcam), Arg1 (PE/Cyanine7, Cat# 369707, Biolegend), Ly6c

(Alexa Fluor 700, Cat# 128023, Biolegend) and Mrc1 (Alexa Fluor 647, ab195192,

Abcam) at 1:400 dilution in the dark for 1 h at 4°C (100μl per antibody for each

sample). All samples were centrifuged at 450RCF for 5 min at 4°C. Supernatants were

removed by aspiration, 1ml 1XPBS solution containing 0.05% BSA was used to wash

the cells twice. FACS analysis was performed on NovoCyte Flow Cytometers (ACEA

Biosciences®, San Diego, California) and FlowJo10.5.0. The experiments were

repeated for three times independently (n=3).

Histological and immunofluorescent staining

The sections were pretreated with 1% BSA in PBS containing 0.1% Triton X 100 for

1 h, incubated in 1% Tween 20 for 20 min and washed again in PBS. The sections

were subsequently analyzed for Krt10 and Krt5, according to the manufacturers'

instructions. Briefly, sections were incubated for 30 min in dark. The excessive dye

was rinsed off with PBS. Sections were incubated with antibody isotype to exclude

false positive staining. Double immunofluorescence staining with primary antibodies

against cytokeratin 10 (ab76318, Abcam, 1:150), cytokeratin 5 (ab52635, Abcam,

1:200) and secondary antibodies (GB25303, GB21303, Servicebio, 1:400) was

performed. The immunostained specimens were further subjected to Hoechst33258

staining (G1011, Servicebio). At least three parallel sections were observed with

fluorescence microscope (ZEISS SteREO Discovery.V20, Olympus). The

fluorescence area measurement was conducted on five random sights of regenerated

epithelium with CaseViewer 2.1 and Image Pro Plus 7.0 (n = 5).

Statistical Analysis

Statistical significance for in vivo and in vitro data of histological analysis, qPCR and

FACS were analyzed by analysis of variance (ANOVA) at the 95% confidence level,

which were performed in GraphPad Prism 8.0 (GraphPad Software, San Diego, CA,

USA) and P < 0.05 was considered statistically significant, while P > 0.05 was

considered having no statistical differences, which was marked with NS.

873