A comprehensive transcriptome signature of murine hematopoietic

stem cell aging

Arthur Flohr Svendsen[1], Daozheng Yang[1], KyungMok Kim[2], Seka Lazare[1], Natalia

Skinder[1], Erik Zwart[1], Anna Mura-Meszaros[2], Albertina Ausema[1], Björn von Eyss[2],

Gerald de Haan[1,3]*, Leonid Bystrykh[1,3]*

[1]European Research Institute for the Biology of Ageing, University Medical Center

Groningen, University of Groningen, Antonius Deusinglaan 1, 9713 AV, Groningen,

The Netherlands.

[2]Transcriptional Control of Tissue Homeostasis Lab, Leibniz Institute on Aging, Fritz

Lipmann Institute e.V., Beutenbergstr. 11,07745 Jena, Germany

[3] These authors contributed equally

* Correspondence: g.de.haan@umcg.nl, l.bystrykh@umcg.nl

14
15
16 Text word count: 3940

17 Abstract word count: 160

18 Number of figures: 6 and 6 supplemental

19 Number of Tables:1 and 3 supplemental

20 Number of references: 51

Abstract

We surveyed 16 published and unpublished data sets to determine whether a consistent pattern of transcriptional deregulation in aging murine hematopoietic stem cells (HSC) exists. Despite substantial heterogeneity between individual studies, we uncovered a core and robust HSC aging signature. We detected increased transcriptional activation in aged HSCs, further confirmed by chromatin accessibility analysis. Unexpectedly, using two independent computational approaches, we established that deregulated aging genes consist largely of membrane-associated transcripts, including many cell surface molecules previously not associated with HSC biology. We show that *Selp*, the most consistent deregulated gene, is not merely a marker for aged HSCs but is associated with HSC functional decline. Additionally, single-cell transcriptomics analysis revealed increased heterogeneity of the aged HSC pool. We identify the presence of transcriptionally "young-like" HSCs in aged bone marrow. We share our results as an online resource and demonstrate its utility by confirming that exposure to sympathomimetics, and deletion of Dnmt3a/b, molecularly resembles HSC rejuvenation or aging, respectively.

Key Points

1. A comprehensive transcriptome analysis of aged murine hematopoietic stem cells identifies an aging signature;

2. The HSC aging signature is highly enriched for cell membrane-related transcripts and identifies age-associated heterogeneity.

Introduction

Aging is a natural and time-dependent process, leading to the functional decline of many cell types and tissues, including those of the hematopoietic system. Aged hematopoietic stem cells (HSCs) have less regenerative potential and their overall contribution to formation of mature blood populations, at the clonal level, is restricted and lineage-skewed when compared to their young counterparts – suggesting stem cell exhaustion, one of the hallmarks of aging[1–9] . Transplantation of aged HSCs into young recipients does not improve HSC function[3,5], indicating that cell-intrinsic aspects play a pivotal role in HSC aging. The differential expression of genes is a prominent read-out of multiple cell-intrinsic factors, and various studies have compared the transcriptomes of young and aged HSCs[10,11,20–24,12–19]. These studies have suggested a plethora of distinct aging-associated mechanisms, including aberrant regulation of genes involved in lymphoid and platelet differentiation[14,22]; elevated inflammation and stress responses[12,17]; chromatin remodeling/DNA methylation[12,23–25]; changes in cell cycle and maintenance of quiescence[15,16,19]; and age-associated replicative stress in HSCs[13]. It remains unclear whether all suggested factors equally contribute to HSC aging, and why different studies report different mechanisms.

Although a substantial number of HSC aging transcriptome studies has been published, no cross-validation of all transcriptomic HSC aging data has yet been carried out. In fact, it is unknown whether a common HSC aging transcriptome exists, and if so, which genes are involved. In order to address this question, we selected 15 studies in which transcriptomes of young and aged murine HSCs were compared, and added to these our own recently published transcriptome analysis[26].

Despite uncovering considerable heterogeneity among different studies, we were able to identify a robust and biologically relevant HSC transcriptomic aging signature. This signature is mainly composed of genes that encode for membrane-associated proteins, many with unknown function(s) in HSC biology. A large fraction of the aging signature genes is upregulated in aged HSCs, indicating transcriptional activation in this population. We also show that aging increases transcriptional heterogeneity among single HSCs and provide evidence that transcriptionally 'young' HSCs exist in aged mice. Furthermore, we demonstrate that the top age-associated gene – *Selp* (P-selectin) – contributes to functional decline of HSCs. Finally, we provide an open

82 web-based resource to the community and use independent data sets to
83 demonstrate how our approach can be used to validate novel biological paradigms.
84

85 **METHODS**
86

Mice

All experiments were approved by the Central Commission for animal Testing and by
Animal Ethical Committee. Young (4-5 months) and aged (20-24 months) C57BL/6,
C57BL/6.SJL (10-14 weeks) and C57BL/6J-kitW-41J/kitW-41J (W41) (originally from
European division of Harlan) were obtained from the Central Animal Facility (CDP) at
the University Medical Center Groningen. TetOP-H2B-GFP mice provided by Prof.
Dr. Rob Coppes (UMCG, Groningen, Netherlands). H2B-GFP expression was
induced by providing mice (6 weeks) with food containing doxycycline (Dox) for 6
weeks. Mice were then fed with normal food for 14 weeks. Mice were kept in group
cages and housed under conventional in a temperature and day cycle controlled
conditions.
98

Flow Cytometry

Bone marrow was isolated from the tibia, femur, pelvis, sternum and spine by
crushing, and red blood cells were lysed with erylysis buffer. All samples were
analyzed on BD FACSCanto™ II (BD Biosciences) and sorted on MoFlo Astrios or
XDP cell sorters (Beckman Coulter). Detailed information of the antibodies used for
isolation and identifications can be found in supplementary text.
105

Retroviral constructs and transduction

Selp was amplified from C57BL/6J LT-HSC cDNA and then cloned into pSF91–
IRES–GFP vector (originally as pSF91 from C. Baum, Hannover Medical School,
Germany, further modified in our lab). Virus was produced by transfecting Platinum-
E retroviral packaging cells (Cell Biolabs) according to the manufacturer's protocol.
Viral supernatant was harvested 48 hours after transfection. LT-HSCs were
transduced as previously described[6]. Cells were harvested 20-22 hours after
transduction for transplantation.
114

BM transplantation

For SELP^{low} and SELP^{high} transplantation, 1000 SELP^{low} and SELP^{high} LT-HSCs

were isolated from old CD45.2$^+$ C57BL/6 mouse and were transplanted together with

2 million CD45.1$^+$ W41 bone marrow into lethally irradiated (9 Gy) CD45.1$^+$

C57BL/6.SJL recipients, respectively. For *Selp* overexpression transplantation, EV

(SF91–IRES–GFP empty) and SELP OE CD45.2$^+$ cells harvested after retroviral

transduction were transplanted together with 2 million CD45.1$^+$ W41 bone marrow

into lethally irradiated CD45.1$^+$ C57BL/6.SJL recipients, respectively.

ATAC-sequencing

25,000 LT-HSCs were sorted from young and aged mice in biological replicates.

Library preparation was performed according to Buenrostro *et al.,* 2013[27]. In brief,

cells were lysed and tagmented for 1h at 37°C. Samples were then PCR-amplified

using provided adapter primers for 11 or 12 cycles. Following, samples were size-

selected for fragments of 150-1000 bp. Libraries were pooled and sequenced to

obtain 30-50 10^6 reads using a NextSeq 500. Data are deposited under accession

number GSE GSE166674

Immunofluorescence

Images were acquired on a Leica Sp8 confocal microscope and quantified on Fiji

Image J.

RNA content

4000-6000 LT-HSCs were seeded onto spots on an adhesion immunofluorescent

slide (VWR). For RNA staining, cells were fixed with 100% methanol for 10 minutes

and washed with PBS. Cells were subsequently stained with 1:10000 SYTO RNA

Select Green Fluorescent Stain (ThermoFisher) for 20 minutes at room temperature.

After washing, coverslips were mounted with ProLong Diamond Antifade Mountant

with DAPI.

RNA pol II

For RNA-Polymerase II staining, cells were fixed and permeabilized with

Fixation/Permeabilization Solution Buffer (BD Bioscience) for 20 minutes on ice.

After washing, cells were blocked first with Endogenous Biotin-Blocking Kit (Thermo

Fisher) followed by 4% BSA for 30 minutes. Cells were stained with 1:100 biotin
mouse monoclonal RNA Polymerase II antibody (Novus Bio) at 4°C overnight. Cells
were then washed with 0.1% Triton-X-100 and stained with 1:1000 streptavidin
Alexa-488 secondary antibody. After washing, coverslips were mounted with
ProLong Diamond Antifade Mountant with DAPI.
154
155 **Further information**
156 More detailed information about the methods can be found in the supplementary
157 text.
158
159 **Additional Resources**
160 Aging signature WEB tool site: http://agingsignature.webhosting.rug.nl/
161 GitHub with all resources: https://github.com/LeonidBystrykh/data-for-manuscript
162 (will be accessible after manuscript acceptance)
163

Results
Systematic collection of publications reporting gene expression changes in
aged HSC
We took advantage of PubMed's API E-utility to select all informative studies on HSC
aging in C57BL/6 mice **(Figure 1A, Supplemental text)**. This search resulted in 16
published studies that met all criteria, and included our own recently published data
set[26] **(Figure 1B/Table S1)**. The various studies used different sequencing
platforms, ranging from microarray hybridization, bulk RNA-sequencing (bRNA-seq)
to single-cell RNA-sequencing (scRNA-seq)**(Figure 1B/Table S1)** and included
mainly 2 age groups **(Figure 1B/S1A)**. We employed two alternative approaches to
perform the analysis. First, we compared the self-reported, pre-analyzed lists of
DEGs from the selected manuscripts and referred to this as 'meta-analysis'. In
parallel, we re-analyzed all available raw data in the most uniform way; these are
referred to as 're-analysis' **(Figure 1A/Table S1)**.
178
Individual studies identify highly variable numbers of DEGs
Strikingly, whereas some studies reported thousands of genes to be differentially
expressed, others only reported dozens **(Figure 1C)**. When DEGs from the meta-

analysis and re-analysis approaches were compared, both demonstrated a
significant correlation between fold-change (FC) expression values, indicating that
re-analysis of the raw data did not deviate drastically from the originally reported
data sets **(Figure S1C).** The observed study-to-study variations might result from
differences in mouse gender, group size, exact age, HSC immunophenotyping,
intrinsic differences between expression platforms, or unknown environmental
effects, including differences in mouse facilities or breeding companies from which
the mice originated **(Table S1)**. Principal component analysis (PCA) performed in the
re-analyzed sets demonstrated that studies clustered according to their sequencing
platform **(Figure S1B)**, although degree of their separation was minor. Overall, this
analysis demonstrates that the variability between studies partially has a technical
origin, but the discrepancy between studies is not fully explained.
194

Most reported age-associated HSC genes have low reproducibility
Collectively these 16 studies reported more than 6000 genes being differentially
expressed between young and aged HSCs. Surprisingly, the vast majority of
reported genes demonstrated poor reproducibility as almost 80% of all putative age-
associated genes were reported in a single study only and not in any other **(Figure**
S1D). Moreover, pair-wise analysis of all different studies showed that some studies
had little to no overlapping genes with any other study **(Figure S1E/F).**
202

Identification of a HSC aging signature
Despite limited reproducibility between individual data sets, we wondered whether it
would be possible to identify consistently DEGs across all studies. To this end, we
ranked all reported genes by their occurrence in different studies (consistency)
(Figure 1D). This revealed a limited number of highly consistent genes that were
frequently documented to be differentially expressed upon HSC aging. The single
most consistently deregulated gene was *Selp.* We provide a complete gene list in
Table S2. We restricted further analysis to genes which were consistently reported in
≥ 4 studies, resulting in a list of 142 genes for the meta-analysis and 220 genes for
the re-analyzed data sets. We refer to this as the "aging signature" (AS) **(Figure**
S1G/ Table S2).
214

The aging signature is robust and provides higher resolution than individual studies

In order to assess the minimal number of publications required to generate a robust list of consistent aging genes, we modelled all possible combinations of most consistent genes from the re-analyzed data, starting from 1 data set at a time to all combinations of 11 out of 12 data sets **(Figure 2B)**. This analysis generated multiple consistencies for the same genes across different permutations. When the analysis was performed with a small number of data sets, we observed a high degree of variation, which started to diminish when the number of data sets increased. When >6 data sets were used, the number and consistency of genes stabilized **(Figure 1E/ supplemental text)**. Complementarily, we also modeled the predicted resolution – the number of aging genes found – for the same range of studies **(Figure 1F)**. The model predicted that the number of studies that we used for our analyses (11 for meta-analysis and 12 for re-analysis) have high resolution; with the current number of included studies, >90% of all the possible age-associated genes have been found. This also implies that the gene collection in the HSC AS remains stable and addition of novel HSC transcriptomes is unlikely to significantly alter the aging signature.

We evaluated the degree of similarity between the AS and the individual studies from which it is derived **(Figure 1G)**. All studies demonstrated approximately the same similarity to the AS, with no clear outliers. No single study encompassed all aging genes – but collectively they contribute to the robustness of the AS. We conclude that the AS can be used as a reference for DEGs upon HSC aging.

The aging signature reveals high enrichment for membrane-related processes in HSC transcriptional activation

We restricted subsequent analyses to the aging signature derived from the re-analyzed data and observed several remarkable features. First, approximately half of the aging genes encode for membrane-associated proteins. **(Figures 2A/B)**. These included many cell adhesion molecules or channels (*Selp, Alcam, Cd34, Dsg2, Itgb3, Clca3a1*) and cytokine receptors (*Csf2rb, Ebi3, Flt3, Ghr, Il1rapl2, Osmr),* many of which have not been previously associated with HSC biology.

Gene Ontology (GO) analysis identified a total of 46 enriched GO function terms **(Table S3)**. As expected, the most significant GO terms revolve around membrane associated functions **(Figure S2A)**. Among upregulated processes, we observed

significant enrichment of different GO terms associated with cell adhesion and
activation, and the organization of cell junction and projection **(Figure 2C)**. Within the
downregulated processes, we identified significant enrichment for DNA replication,
cytokine signaling, tissue homeostasis, and cell proliferation **(Figure 2D)**. Pathway
analysis also demonstrated analogous pathways **(Figure S2B)**.

We next collected all associated GO terms associated with every gene present in the
aging signature **(Figure S2C)**, resulting in more than 2000 GO terms. The majority of
these terms were very specific and contained less than 20 AS genes **(Figure S2C,**
lower panel). We then searched in the AS GO term database for key terms that
represent previously described HSC aging mechanisms, including "inflammation"[12],
"replicative stress"[13], "cell cycle"[15,16] and "epigenetic drift"[12,23,28] – and indeed found
these. However, the large majority of genes (149) were associated with "membrane"
(Figure S2D), denoting that significant changes occur in the HSC membrane upon
aging.

Secondly, we observed that a disproportional number (181 genes, ~80%) from the
AS were upregulated in aged HSCs **(Figure 3A).** We assessed whether this would
be reflected in global transcriptional activation. Comparing young and aged LT-HSCs
(LSK CD48$^-$ CD150$^+$) **(Figure S3A)**, we observed a ~3-fold higher overall RNA yield
per cell in aged HSCs **(Figure S3B).** We confirmed this by intracellular staining of
young and aged LT-HSCs with the RNA-specific dye SYTO **(Figure S3D)**. In order to
investigate if increased RNA abundance in aged LT-HSCs was directly linked to
increased transcription, we performed intracellular staining for RNA Polymerase II
(POL-II) in young and aged HSCs and indeed observed increased RNA POL-II
activity in aged cells **(Figure 3B)**.

Finally, we evaluated the number of chromatin accessible sites by ATAC-seq. In
agreement with the data above, aged HSCs showed an increased number of overall
accessible sites compared to young HSCs **(Figure 3SD).** In addition, differential
analysis of accessible sites demonstrated strong skewing towards more open sites in
aged HSCs **(Figure 3C)**. We then speculated that aged-specific accessible sites
could contribute to deregulated expression. To investigate this, we overlapped the
age-restricted accessible sites with the genes in the aging list – which would reflect
genes more accessible in aged HSCs and therefore upregulated – and performed a
GO analysis **(Figure 3D)**. This revealed high enrichment for transcription-related

processes, further reinforcing our hypothesis that aged HSCs are transcriptionally activated.

Taken together, the aging signature suggests that aged HSCs undergo adaptation at cellular and functional levels through membrane-related processes and that aged HSCs demonstrate higher transcriptional activation.

The aging signature as a reference for age-associated studies

We aimed to test whether the aging signature can be used as a reference to (re-) analyze existing and future HSC gene expression profiles. To this end, we developed two independent approaches where custom transcriptome sets can be compared to the aging signature.

First, we designed two scoring methods, the discrete Gene Set Enrichment (dGSE) algorithm and a Rank Score (RS), both of which are able to cope with semi-ordered (discrete) lists of genes **(Supplemental text)** and can be used for testing bulk RNA-seq sets. To benchmark both methods, we individually tested the 12 HSC aging transcriptome studies that comprise the aging signature **(Figure 4A)**. As expected, we observed that the different data sets had a wide range of enrichment scores when compared to the aging signature.

We next used transcriptomic data sets from three unrelated studies. The first independent data set compared young and physiologically aged HSCs (Kim *et al.*, manuscript in revision). The second involved HSCs that were "rejuvenated" by a sympathomimetic drug acting on adrenoreceptor β3[18]. The third study used expression data from HSCs that were derived from *Dmnt3a/b* deficient mice that show accelerated aging phenotypes[29]. As a negative control, we included a transcriptomic signature of embryonic stem cells (ES)[30]. As expected, the ES signature demonstrated low scores for aging genes. In contrast, the independent set of young and physiologically aged HSCs demonstrated enrichment when compared to the AS. When we compared the list of genes that were downregulated in "rejuvenated" old HSCs to the AS, we also found strong enrichment. The transcriptome of young HSCs isolated from Dmnt3a/b KO mice also scored high for HSC aging genes. These data suggest that both β3 sympathomimetic signaling and loss of *Dnmt3a/b* molecularly phenocopy normal HSC aging, additionally confirming the value of our method **(Figure 4B)**.

This approach allowed us to investigate the relationship of our HSC aging signature
to a previously established multi-organ inter-species transcriptomic signature[31].
Expectedly, we observed a mild enrichment between both signatures, denoting that
some more general aging-associated genes are also present in HSC aging **(Figure**
S4A). Genes present in both signatures are related to inflammatory response. We
also investigated the involvement of TGF-β signaling, which has been implicated in
HSC aging[21,23,32], in the aging signature. We compared different molecular
signatures of TGF-β signaling[33,34] to our aging signature, but none of the sets
demonstrated high scores. Furthermore, only 5 TGF-β target genes from all sets
(*Bmpr1a*, *Id2*, *Jun*, *Itgb3* and *Mef2c*) are present in the AS. Thus, TGF-β signaling,
from a transcriptional perspective, is not robustly involved in HSC aging. **(Figure**
S4A).
Interestingly, single-cell transplantations have revealed that the overall engraftment
potential of aged HSCs is substantially lower than that of young HSCs[2]. However,
some aged HSCs display engraftment performance similar to young cells,
suggesting that the aged HSC pool is functionally heterogenous. Thus, we wondered
whether the AS can be used to also identify intrinsic variation between cells from the
same population. To identify such "young-like" cells using single-cell transcriptomic
studies, we assigned an Aging Signature score (AS score) to individual cells
(Figures 4C/S4B). This revealed that separation of young and aged cells is in good
agreement with their assigned AS scores where aged cells score higher and young
cells lower, also predicting their age. Interestingly, some aged cells had a low AS
score and were close to the young cells cluster, suggesting that these cells are, at
least transcriptionally, "young-like".
Taken together the HSC aging signature demonstrated to be a *bode fine*
transcriptional reference to physiological aging and thus a valuable resource for the
scientific community. It allowed us to demonstrate that the aged HSC pool is
heterogeneous and that some aged cells are transcriptionally similar to young HSCs.
We provide both (bulk and single-cell) analyses as a resource to the community at
https://agingsignature.webhosting.rug.nl/.
345
Machine learning algorithms independently identify the best age-associated
gene predictors

Employing machine learning strategies in conjunction with scRNA-seq data have
allowed for prediction and detection of different hematopoietic populations[35,36]. We
envisioned that by applying this approach to the age-related scRNA-seq data sets
would allow for the identification of genes which best divide young and old HSCs. It
would also serve as an independent and alternative approach to DE analysis. We
trained and tested two different machine learning algorithms – ADAboost and
Random Forest (RNDforest) – for identification of the best gene(s) predictors.
Interestingly, both algorithms determined that *Nupr1*, *Selp*, and *Sult1a1* were the
best predictors **(Figure S5SA)**. We tried to restrict the aging-related list to the best
20 age-associated genes. With those 20 genes, both algorithms performed well on
identifying young and aged HSCs **(Figures 5A/B/S5B/C)** when compared to the full
list of AS genes. Attempts to further narrow down the age-predictors from 20 to 5
genes, however, showed considerable deterioration in prediction capacity compared
to larger lists of age-predictors **(Figure S5D)**. Out of 20 gene predictors, the majority
are also membrane-related **(Figure 5C)**.
363
The top aging gene – Selp – affects HSC functioning.
The most consistent gene from the AS – *Selp* – **(Table 2)** was also among the best
machine learning predictors **(Figure 5)**. *Selp*, which encodes for the membrane
adhesion molecule P-selectin, was confirmed to be differently expressed also at the
protein levels in young and aged hematopoietic progenitor subsets by flow
cytometry. **(Figures 6A/S6A)**. To investigate whether Selp affects HSC functioning
we competitively transplanted 1000 SELP^{low} or 1000 SELP^{high} aged LT-HSCs
(Figure S6B) into lethally irradiated young recipients. Whereas overall engraftment
levels were similar for SELP^{low} and SELP^{high} LT-HSCs, we found that SELP^{high} LT-
HSCs produced more myeloid cells, suggesting that SELP^{high} HSCs are myeloid
biased **(Figures 6B/S6C)**.
To determine the role of *Selp* in the functioning of LT-HSCs, we overexpressed *Selp*
in young LT-HSCs, and competitively transplanted these in young recipients.
Whereas *Selp* overexpression did not affect differentiation to leukocytes and
platelets, erythroid production was almost completely abolished **(Figure 6C).** We
also employed a Tet-On H2B-GFP reporter mouse to determine the dormancy or
activation status of SELP^{low} and SELP^{high} LT-HSCs. GFP fluorescence was

significantly higher in SELP^{low} LT-HSCs, indicating that myeloid-biased SELP^{high} LT-

HSCs were more actively cycling **(Figure S6D).**

Thus, higher levels of SELP in LT-HSCs functionally contribute to myeloid bias,

erythroid blockage and HSC activation. Collectively, this demonstrates that SELP is

not only a marker of aged HSCs but has the capacity to contribute to the functional

impairment of HSCs.

Discussion

In this study we demonstrate that individual studies have limited resolution to identify

genes altered in HSC aging, but collectively they provide an unprecedented profile of

transcriptomic alterations in HSC aging **(Figure 1D)**. From this extensive repertoire

of studies, we were able to identify a comprehensive, robust, and stable

transcriptomic signature of HSC aging, which we validated by 3 independent data

sets.

The HSC aging signature is highly enriched for membrane-associated transcripts

(Figure 2A). Several of these have been previously associated with aspects of the

HSC aging process and thus document the rigor of our approach **(Table 1)**.

However, the role of most other genes in HSC (aging) biology remains unknown.

The strong involvement of cell surface molecules does provide insight into the long-

standing discussion whether HSC aging results from cell-intrinsic or cell-extrinsic

molecular alterations. This enrichment of cell membrane-associated transcripts

suggests that physiologically aged HSCs communicate differently with their

immediate environment compared to their young counterparts.

Our study also provides novel insight into the transcriptional activation state of HSCs

(**Figure 3**). In line with recent reports that loss of heterochromatin leads to age-

associated phenotypes in HSCs[37], RNA-specific quantification, RNA POL-II activity

and chromatin accessibility analyses, all indicate that aged HSCs are, at a

population level, more transcriptional active than young HSCs.

The aging signature can be employed as a *bona fide* transcriptional reference for

physiological HSC aging for both bulk and single-cell transcriptomic studies **(Figure**

4). We provide protocols and a user-friendly and interactive website –

https://agingsignature.webhosting.rug.nl/ – to allow users to execute custom queries

and compare new data sets against the aging signature that we present here.

Our approaches also allowed powerful analyses and uncovered novel perspectives
into HSC aging from a single-cell perspective. By applying our AS score to single-cell
transcriptomic studies, we determined that HSCs isolated from aged mice contain a
fraction of cells which behave, at least transcriptionally, like young HSCs **(Figure
4C)**. Further studies are necessary to understand if such cells indeed are functionally
superior and also why such cells do not seem to "age". We also successfully
employed machine leaning to further identify best age-associated gene predictors
(Figure 5).

Both computational approaches demonstrated that *Selp* expression is strongly
associated with HSC aging. Previous data have shown that *Selp*-deficient HSCs
display a competitive advantage over wild-type HSCs[38]. Here we document that high
levels of SELP expression induced myeloid-biased, erythroid blockage and cell
activation, all hallmarks of HSC aging[8,9] **(Figure 6)**. Together with *Selp,* which is a
well-known inflammatory marker, *Nupr1* is one of the top-listed predictors and
belongs to a group of inflammatory and stress-responding transcription factors, not
earlier detected in relation to the aging phenotype. Although we were able to reduce
the list of age-related genes, it is clear that HSC aging cannot be reduced to a single
gene. Instead, our machine learning analysis suggests that multiple age-related
genes are involved in the HSC aging process. The idea of a single gene explaining
most of the aging phenotype sounds unlikely, and that multiple cell surface markers
are required to prospectively separate young and aged HSCs.

Nevertheless, our data and others demonstrate that (at least some of) the aging
genes that we identify are not merely markers of aged HSCs, but also are involved in
age-dependent functional decline. The identification of the HSC aging signature does
not resolve whether altered expression is the cause or consequence of loss of HSC
functioning during aging. Future functional studies are required to assess whether
perturbation of AS genes can improve functioning of aged stem cells. We believe
that this resource is valuable in assisting the aging research community to verify
involvement of genes in physiological HSC aging.

Acknowledgements

We thank T. Bijma, G.Mesander and J. Teunis from UMCG Flowcytometry Unit facilities for their assistance on cell sorting; Klaas Sjollema from UMCG Microscopy and Imaging Center (UMIC) for assistance with confocal microscopy; B Bakker, M Gerritsen, E Verovskaya and all members of the Ageing Biology and Stem Cells laboratory for discussions. This work was supported by the Netherlands Organization for Scientific Research/Mouse Clinic for Cancer and Aging, the Landsteiner Foundation for Blood Transfusion Research (LSBR1703), a China Student Council Fellowship awarded to D.Y., Marriage, a EU FP7 Marie Curie Initial Training Network (Contract number 316964) and ARCH, a European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 813091. B.v.E. was supported by grants from the DFG (EY 120/1-1), Else Kröner-Fresenius Foundation (2016_A58) and the German Cancer Aid (Deutsche Krebshilfe; 70113138). The FLI is a member of the Leibniz Association and is financially supported by the Federal Government of Germany and the State of Thuringia.

Author Contributions

Conceptualization, A.F.S., D.Y., S.S., G.d.H. and L.B.; Methodology, A.F.S., D.Y., and L.B.; Investigation, A.F.S., D.Y., S.S., A.A. K.K; A.M-M., B.v.E. and L.B.; Formal Analysis, A.F.S., D.Y., S.S. K.K; A.M-M., B.v.E. and L.B.; Data Curation, A.F.S., E.Z., L.B.; Resources, E.Z. and A.A.; Writing – Original Draft, A.F.S.; Writing – Review & Editing, A.F.S., D.Y., G.d.H. and L.B.; Supervision, G.d.H. and L.B.

Disclosure of Conflicts of Interest

The authors declare no competing interests.

References

1. de Haan G, Nijhof W, Van Zant G. Mouse strain-dependent changes in frequency and proliferation of hematopoietic stem cells during aging: correlation between lifespan and cycling activity. *Blood.* 1997;89(5):1543–50.

2. Dykstra B, Olthof S, Schreuder J, Ritsema M, de Haan G. Clonal analysis reveals multiple functional defects of aged murine hematopoietic stem cells. *J. Exp. Med.* 2011;208(13):2691–2703.

3. Morrison SJ, Wandycz AM, Akashi K, Globerson A, Weissman IL. The aging of hematopoietic stem cells. *Nat. Med.* 1996;2(9):1011–1016.

4. Rossi DJ, Jamieson CHM, Weissman IL. Stems Cells and the Pathways to Aging and Cancer. *Cell.* 2008;132(4):681–696.

5. Sudo K, Ema H, Morita Y, Nakauchi H. Age-associated characteristics of murine hematopoietic stern cells. *J. Exp. Med.* 2000;192(9):1273–1280.

6. Verovskaya E, Broekhuis MJC, Zwart E, et al. Asymmetry in skeletal distribution of mouse hematopoietic stem cell clones and their equilibration by mobilizing cytokines. *J. Exp. Med.* 2014;211(3):487–497.

7. Verovskaya E, Broekhuis MJC, Zwart E, et al. Heterogeneity of young and aged murine hematopoietic stem cells revealed by quantitative clonal analysis using cellular barcoding. *Blood.* 2013;122(4):523–532.

8. López-Otín C, Blasco MA, Partridge L, Serrano M, Kroemer G. The hallmarks of aging. *Cell.* 2013;153(6):1194.

9. Singh PP, Demmitt BA, Nath RD, Brunet A. The Genetics of Aging: A Vertebrate Perspective. *Cell.* 2019;177(1):200–220.

10. Beerman I, Rossi DJ. Epigenetic Control of Stem Cell Potential during Homeostasis, Aging, and Disease. *Cell Stem Cell.* 2015;16(6):613–625.

11. Bersenev A, Rozenova K, Balcerek J, et al. Lnk deficiency partially mitigates hematopoietic stem cell aging. *Aging Cell.* 2012;11(6):949–959.

12. Chambers SM, Shaw CA, Gatza C, et al. Aging hematopoietic stem cells decline in function and exhibit epigenetic dysregulation. *PLoS Biol.* 2007;5(8):1750–1762.

13. Flach J, Bakker ST, Mohrin M, et al. Replication stress is a potent driver of functional decline in ageing haematopoietic stem cells. *Nature.* 2014;512(7513):198–202.

505   14.   Grover A, Sanjuan-Pla A, Thongjuea S, et al. Single-cell RNA sequencing
506   reveals molecular and functional platelet bias of aged haematopoietic stem
507   cells. *Nat. Commun.* 2016;7:.

508   15.   Kirschner K, Chandra T, Kiselev V, et al. Proliferation Drives Aging-Related
509   Functional Decline in a Subpopulation of the Hematopoietic Stem Cell
510   Compartment. *Cell Rep.* 2017;19(8):1503–1511.

511   16.   Kowalczyk MS, Tirosh I, Heckl D, et al. Single-cell RNA-seq reveals changes
512   in cell cycle and differentiation programs upon aging of hematopoietic stem
513   cells. *Genome Res.* 2015;25(12):1860–1872.

514   17.   Mann M, Mehta A, de Boer CG, et al. Heterogeneous Responses of
515   Hematopoietic Stem Cells to Inflammatory Stimuli Are Altered with Age. *Cell
516   Rep.* 2018;25(11):2992-3005.e5.

517   18.   Maryanovich M, Zahalka AH, Pierce H, et al. Adrenergic nerve degeneration in
518   bone marrow drives aging of the hematopoietic stem cell niche. *Nat. Med.*
519   2018;24(6):782–791.

520   19.   Noda S, Ichikawa H, Miyoshi H. Hematopoietic stem cell aging is associated
521   with functional decline and delayed cell cycle progression. *Biochem. Biophys.
522   Res. Commun.* 2009;383(2):210–215.

523   20.   Norddahl GL, Pronk CJ, Wahlestedt M, et al. Accumulating mitochondrial DNA
524   mutations drive premature hematopoietic aging phenotypes distinct from
525   physiological stem cell aging. *Cell Stem Cell.* 2011;8(5):499–510.

526   21.   Quere R, Saint-Paul L, Carmignac V, et al. Tif1 regulates the TGF- 1 receptor
527   and promotes physiological aging of hematopoietic stem cells. *Proc. Natl.
528   Acad. Sci.* 2014;111(29):10592–10597.

529   22.   Rossi DJ, Bryder D, Zahn JM, et al. Cell intrinsic alterations underlie
530   hematopoietic stem cell aging. 2005.

531   23.   Sun D, Luo M, Jeong M, et al. Epigenomic profiling of young and aged HSCs
532   reveals concerted changes during aging that reinforce self-renewal. *Cell Stem
533   Cell.* 2014;14(5):673–688.

534   24.   Wahlestedt M, Norddahl GL, Sten G, et al. An epigenetic component of
535   hematopoietic stem cell aging amenable to reprogramming into a young state.
536   *Blood.* 2013;121(21):4257–4264.

537   25.   Beerman I, Bock C, Garrison BS, et al. Proliferation-dependent alterations of
538   the DNA methylation landscape underlie hematopoietic stem cell aging. *Cell

539   *Stem Cell.* 2013;12(4):413–425.

540   26.   Renders S, Svendsen AF, Panten J, et al. Niche derived netrin-1 regulates

541   hematopoietic stem cell dormancy via its receptor neogenin-1. *Nat. Commun.*

542   2021;12(1):1–15.

543   27.   Buenrostro JD, Giresi PG, Zaba LC, Chang HY, Greenleaf WJ. Transposition

544   of native chromatin for fast and sensitive epigenomic profiling of open

545   chromatin, DNA-binding proteins and nucleosome position. *Nat. Methods.*

546   2013;10(12):1213–8.

547   28.   Hidalgo I, Herrera-Merchan A, Ligos JM, et al. Ezh1 is required for

548   hematopoietic stem cell maintenance and prevents senescence-like cell cycle

549   arrest. *Cell Stem Cell.* 2012;11(5):649–662.

550   29.   Challen GA, Sun D, Mayle A, et al. Dnmt3a and Dnmt3b have overlapping and

551   distinct functions in hematopoietic stem cells. *Cell Stem Cell.* 2014;15(3):350–

552   364.

553   30.   Wong DJ, Liu H, Ridky TW, et al. Module Map of Stem Cell Genes Guides

554   Creation of Epithelial Cancer Stem Cells. *Cell Stem Cell.* 2008;2(4):333–344.

555   31.   de Magalhães JP, Curado J, Church GM. Meta-analysis of age-related gene

556   expression profiles identifies common signatures of aging. *Bioinformatics.*

557   2009;25(7):875–881.

558   32.   Blank U, Karlsson S. TGF-β signaling in the control of hematopoietic stem

559   cells. *Blood.* 2015;125(23):3542–3550.

560   33.   Subramanian A, Tamayo P, Mootha VK, et al. Gene set enrichment analysis: A

561   knowledge-based approach for interpreting genome-wide expression profiles.

562   *Proc. Natl. Acad. Sci. U. S. A.* 2005;102(43):15545–15550.

563   34.   Liberzon A, Birger C, Thorvaldsdóttir H, et al. The Molecular Signatures

564   Database Hallmark Gene Set Collection. *Cell Syst.* 2015;1(6):417–425.

565   35.   Hamey FK, Göttgens B. Machine learning predicts putative hematopoietic

566   stem cells within large single-cell transcriptomics data sets. *Exp. Hematol.*

567   2019;78:11–20.

568   36.   Fidanza A, Kingdom U, Stumpf P, et al. Single cell analyses and machine

569   learning define hematopoietic progenitor and HSC-like cells derived from

570   human PSCs Single cell analyses and machine learning define haematopoietic

571   progenitor and 4 - Centre for Integrative Systems Biology and Bioinformati.

572   *Blood.* 2021;

37. Keenan CR, Iannarella N, Naselli G, et al. Extreme disruption of
heterochromatin is required for accelerated hematopoietic aging. *Blood*.
2020;135(23):2049–2058.

38. Sullivan C, Chen Y, Shan Y, et al. Functional ramifications for the loss of p-
selectin expression on hematopoietic and leukemic stem cells. *PLoS One*.
2011;6(10):1–10.

39. Rapisarda V, Borghesan M, Miguela V, et al. Integrin Beta 3 Regulates
Cellular Senescence by Activating the TGF-β Pathway. *Cell Rep*.
2017;18(10):2480–2493.

40. Sanjuan-Pla A, Macaulay IC, Jensen CT, et al. Platelet-biased stem cells
reside at the apex of the haematopoietic stem-cell hierarchy. *Nature*.
2013;502(7470):232–236.

41. Pinho S, Marchand T, Yang E, et al. Lineage-Biased Hematopoietic Stem
Cells Are Regulated by Distinct Niches. *Dev. Cell*. 2018;44(5):634-641.e4.

42. Gulati GS, Zukowska M, Noh JJ, et al. Neogenin-1 distinguishes between
myeloid-biased and balanced Hoxb5+ mouse long-term hematopoietic stem
cells. *Proc. Natl. Acad. Sci. U. S. A.* 2019;116(50):25115–25125.

43. Jeannet R, Cai Q, Liu H, Vu H, Kuo Y-H. Alcam Regulates Long-Term
Hematopoietic Stem Cell Engraftment and Self-Renewal. *Stem Cells*.
2013;31(3):560–571.

44. Shimazu T, Iida R, Zhang Q, et al. CD86 is expressed on murine
hematopoietic stem cells and denotes lymphopoietic potential. *Blood*.
2012;119(21):4889–4897.

45. Thalheimer FB, Wingert S, De Giacomo P, et al. Cytokine-regulated
GADD45G induces differentiation and lineage selection in hematopoietic stem
cells. *Stem Cell Reports*. 2014;3(1):34–43.

46. Satoh Y, Yokota T, Sudo T, et al. The Satb1 Protein Directs Hematopoietic
Stem Cell Differentiation toward Lymphoid Lineages. *Immunity*.
2013;38(6):1105–1115.

47. Taiwo O, Wilson GA, Emmett W, et al. DNA methylation analysis of murine
hematopoietic side population cells during aging. *Epigenetics*.
2013;8(10):1114–1122.

48. Oguro H, Ding L, Morrison SJ. SLAM family markers resolve functionally
distinct subpopulations of hematopoietic stem cells and multipotent

607   progenitors. *Cell Stem Cell*. 2013;13(1):102–116.

608 49. Beerman I, Bhattacharya D, Zandi S, et al. Functionally distinct hematopoietic

609   stem cells modulate hematopoietic lineage potential during aging by a

610   mechanism of clonal expansion. *Proc. Natl. Acad. Sci. U. S. A.*

611   2010;107(12):5465–70.

612 50. Tadokoro Y, Ema H, Okano M, Li E, Nakauchi H. De novo DNA

613   methyltransferase is essential for self-renewal, but not for differentiation, in

614   hematopoietic stem cells. *J. Exp. Med.* 2007;204(4):715–722.

615 51. Challen GA, Mayle A, Sun D, et al. Dnmt3b Has Few Specific Functions In

616   Adult Hematopoietic Stem Cells But Shows Abnormal Activity In The Absence

617   Of Dnmt3a. *Blood.* 2013;122(21):734–734.

Tables

Table 1. Top 10 Aging signature genes with experimental validated function in HSC aging

Gene Symbol	Consistency (out of 12)	Average Log$_2$FC	Function	Phenotype	Reference
Itgb3	9	1.25±0.48	Cell adhesion; Signal transduction	Itgb3[high] identifies myeloid-biased HSCs	[17,39]
Vwf	8	1.37±0.66	Cell adhesion; Signal transduction	Vwf[high] identifies myeloid-biased HSCs	[14,40,41]
Neo1	8	1.84±1.18	Cell adhesion; Signal transduction	Neo1[high] identifies more dormant HSCs	[26,42]
Alcam	8	1.39±0.62	Cell adhesion	Alcam[+] identifies myeloid-biased HSCs and with reduced potential	[43]
CD86	6	-1.39±0.36	Signal transduction	Required for lymphoid differentiation	[44]
Gadd45g	6	1.54±0.99	Stress response	Gadd45g[-/-] show increased self-renewal and deficient Mk differentiation	[45]
Satb1	6	-1.21±0.50	Epigenetic effector	Required for lymphoid differentiation	[46]
Sdpr	6	1.88±1.61	Transcription regulation	Involved in PRC2 expression regulation	[47]
Slamf1	5	0.92±0.16	Cell adhesion Signal; transduction	Slamf1[high] identifies myeloid-biased HSCs and with reduced potential	[48,49]
Dnmt3b	4	-1.09±0.62	Epigenetic regulator	*Dnmt3a/b[-/-]* show differentiation blockage/ increased self-renewal	[29,50,51]

Figure Legends

Figure 1: Systematic analysis reveals heterogeneity in transcriptomic studies of young and aged murine HSCs.

A) Selection of young and aged HSC transcriptomic studies. Representation of decision tree used to acquire and analyze the different publications used for the systematic analysis.

B) Overview of the age groups used in each of the selected studies. The age group of mice used in each study (y-axis) is represented in months (x-axis). Color codes refer to the transcriptome platform used. Squares next to each study indicate in which kind of analysis each study was included.

C) Heterogeneity across different transcriptomic studies. Number of differentially expressed genes (DEGs) reported by each study. Left panel shows the number of DEGs as reported by the authors in their respective manuscripts. Right panel shows the number of DEGs identified by us using a uniform re-analysis. Red bars indicate the number of genes that are upregulated upon aging, while blue bars indicate downregulated genes. In the re-analysis panel, "h-Chambers" indicates half-Chambers meaning that just half of the DEGs were taken into account.

D) Ranking of DEGs across different studies. Horizontal bars represent the number of DEGs (x-axis, the actual number of DEG genes are shown in each bar), ranked by the number of studies in which they were found (occurrence in studies (consistency), y-axis). Purple and green bars indicate meta-analysis and re-analysis, respectively.

E) The list of genes gets more consistent with the increase number of studies added. Simulation of different list assembly using combinations of subsets of 12 studies (gradient color from blue to red). As the number of studies increased, the coefficient of variation (CV, y-axis) decreases. Dots represent the different

values and mean consistency for individual genes **(see supplemental text for**
further details).

F) Predicted resolution to discover consistently reported genes. The model
demonstrates that >90% of all HSC aging genes have been identified (y-axis) in
the two analytic approaches. Purple and green dashed lines indicate meta-
analysis and re-analysis, respectively.

G) Extent of overlap between all independent studies and the HSC aging signature
(centered red circle, AS). The size of the circles represents the number of DEGs
identified in each study; the number of overlapping DEGs with the aging
signature is represented next to each study identification; the distance from the
center represents the degree of similarity of each study to the HSC aging
signature. Studies that show a higher degree of similarity are closer to the AS.

Figure 2: The HSC aging signature is highly enriched for membrane-
associated genes
676

677   A) Classification of the proteins of the aging signature by cellular compartment. The
678   localization of each protein was determined using UNIPROT classification.
679   Colors of the bars represent upregulation (red) or repression (blue) of genes in
680   the AS. Numbers next to the bars indicate the frequency of genes found in each
681   category.
682

683   B) Network representation of the aging genes based on the cellular compartment
684   where the encoded proteins are predicted to be localized. The size of each circle
685   represents the number of genes found in each category. Connections between
686   each cellular compartment reflect the number of genes which are shared per
687   cellular compartment.
688

C/D) GO analysis of biological process for AS genes. Horizontal bar represent
individual GO terms (y-axis) and their FDR values (x-axis). Numbers in each bar
demonstrate the number of genes found per GO term. Red (panel C) and blue
(panel D) bars represent GO terms found to be significantly enriched in
upregulated and downregulated AS genes, respectively. GO terms are selected
based on an FDR < 0.05 cut-off.
695
696

697 **Figure 3: Skewing of HSC aging signature genes reflects transcriptional**
698 **activation in aged HSCs**
699

A) Skewing of HSC aging signature genes**.** Volcano plot depicting expression levels
of AS genes. 181 upregulated and 39 downregulated genes are represented.
Dots represent the average of either adjusted p-values (adjpval, y-axis) and fold-
changes (FC, x-axis) per gene in different studies.

B) Protein levels of RNA Pol-II in individual young (6 months) and aged (24 months)
LT-HSCs. Right panel shows the MFI quantification of young (n = 120 cells) and
aged HSCs (n = 139 cells). Scale bars in IF images are 5 µm. ±SD is shown. ****
p < 0.0001.

C) Aged HSCs demonstrate more age-specific chromatin accessible sites than
young HSCs. Volcano plot depicting differentially accessible sites (peaks)
between young and old HSCs measured by ATAC-seq. Dot colors demonstrate
peaks either in genomic regions overlapping with coding (red) or non-coding
(blue) annotations. Peaks with negative FC values are significantly more
accessible in young HSCs whereas peaks with positive FCs values are more
significantly accessible in aged HSCs. Gene symbols annotate peaks which
overlap with gene bodies from some AS genes.

718
D) Transcriptional program is associated with ATAC-seq of young and aged HSCs.
GO analysis of functional process for overlapping aged-specific ATAC-seq
accessible sites with genes from AL. Horizontal bar represent individual GO
terms (y-axis) and their FDR values (x-axis). Numbers in each bar demonstrate
the number of genes found per GO term. GO terms are selected based on a
FDR < 0.05 cut-off.

Figure 4: Employing the HSC aging signature as a *bona fide* transcriptomic reference

A) Using Rank Score (RS) and discrete gene set enrichment (dGSE) analyses to compare existing data sets to the HSC aging signature. Left panel: graphical representation of the dGSE enrichment analyses from our own generated HSC transcriptome data was compared to the HSC aging signature. Right panel: Plots of RS (upper) and dGSE (lower) enrichment analyses of all individual studies that comprise the AS. Numbers refer to the studies shown on the right.

B) Searching for an aging signature using RS and dGSE analyses in independent data sets. Upper left panel: Lack of enrichment for aging genes in an embryonic stem cell signature from *Wong et al.* Upper Right panel: Enrichment for the top250 genes (sorted by FDR value) of Kim *et al.;* Lower right panel: Significant enrichment for an HSC aging signature among genes downregulated in "rejuvenated" HSCs from Maryanovich *et al.* Lower Right panel: Significant enrichment for a HSC signature among differentially expressed genes in HSCs isolated from young DNMT3A/B KO mice from Challen *et al.* The right-most panels show combined RS and dGSE scores for each test set.

C) Single-cell AS scores. Separation of young (circles) and aged (triangles) single-cell transcriptomics and their respective AS score. From Grover scRNA-seq data (48 young and 68 aged cells), individual cells were separated by PC1 (x-axis) and PC2 (y-axis) and color-coded according to their individual aging signature score. Left column display only young cells; middle column shows just aged cells and right column shows both populations. Cells are colored score from low (blue) to high (yellow) AS scores. On the far-right panel, violin and boxplot demonstrate the overall AS score (y-axis) distribution of the young (grey) and aged (red) cells.

Figure 5: Machine learning identifies best 20 gene predictors for separating young and aged HSCs in scRNA-seq

A) Machine learning is able to predict with high accuracy young and aged HSCs. Top panel depicts a schematic representation of machine learning application into transcriptomic data from different scRNA-seq sets. Lower panel represents the output of one of the algorithm used (ADAboost) in individual cells from different sets. Young (circles) and aged (triangles) single-cell were separated by PC1 (x-axis) and PC2 (y-axis) and color-coded according to the match between the measured aged of the cells and the predicted aged measured by the algorithm (grey for young cells; red for aged cells and orange for mismatched cells).

B) Machine learning scores varies depending on which training set is used. Heatmap from ADAboost training depicting different overall scores for different scRNA-seq sets used. The overall score is color-coded from blue (lower scores) to yellow (higher scores). Sets on the x-axis (training sets) were used to train the algorithm and the following sets on the y-axis (test sets) were scored according to training.

C) The best machine learning genes predictors have also high enrichment for membrane-associated proteins. Horizontal stacked plot comparing the list of the 20 best predictors extracted from machine learning algorithms (best predictors) and the aging signature genes. The percentage of genes in each cellular location is represented as a percentage (x-axis) and divided by category (different colors). The 20 best predictors gene symbols are represented below.

Figure 6: *Selp* (P-Selectin) is the top aging signature candidate and contributes to HSC functional decline

A) SELP protein expression level across different hematopoietic stem and progenitors (HSPCs) in young (grey) and aged (red) mice. Panel shows fold-changes of aged (n = 3 mice) samples normalized to their young (n = 3 mice) counterparts. Dots show biological replicates while the bars represent the mean values for the given population. Flow cytometry plot for SELP expression in young and aged LT-HSCs population in the 3 biological replicates.

B) Selp^high^ LT-HSCs display a myeloid bias. The upper panel shows the experimental approach for competitive transplantation. SELP^low^ (blue) and SELP^high^ (orange) LT-HSCs were purified from aged mice and competitively transplanted (n = 6-7 mice for each group). Overall engraftment (WBC), and T cells levels were similar comparing both groups whereas Myeloid (My) and T-cells levels indicate that transplanted LT-HSC SELP^high^ have Myeloid-bias.

C) SELP OE recipients show impairment of erythroid differentiation. The upper panel shows the experimental approach for competitive transplantation of SELP OE and EV (empty vector) HSCs. Lower panel displays peripheral blood (PB) output of the GFP$^+$ (transduced cells) of both groups (n = 3 per group). Within the GFP$^+$ population, the number of Ter-119$^+$ population is significantly reduced.

For all panels, ±SD is shown. ns = non-significant; * $p < 0.05$; ** $p < 0.01$; n indicates biological replicates

Figure 1

Figure 2

A

membrane — 17 | 97
cytoplasm — 7 | 44
nucleus — 9 | 33
secreted — 9 | 18
endoplasmic reticulum — 2 | 8
extracellular — 6 |
golgi apparatus — | 7
lysosome — 3 |
mitochondrion — 2 |
cytoskeleton — 1 | 4
microsome — 1 | 3
endosome — | 3
melanosome — 1 | 2
desmosome — | 2
peroxisome — | 1
phagosome — 1 |

number of genes

B

peroxisome
mitochondria
endoplasmic reticulum
microsome
nucleus
lysosome
desmosome
membrane
phagosome
secretion
cytoplasm
cytoskeleton
golgi apparatus
extracllular matrix
melanosome
extracelullar
endosome
cillium

number of genes

C

upregulated GO terms

cell adhesion — 22/705
cell activation — 19/552
cell junction organization — 10/147
cell projection organization — 25/1020
regulation of neuron death — 14/348
tissue development — 32/1599
endocytosis — 15/424
regulation of transport — 34/1782
tube morphogenesis — 19/690
regulation of coagulation — 7/80

−log$_2$FDR

D

downregulated GO terms

cytokine−mediated signaling pathway — 7/313
response to other organism — 8/779
collagen−activated tyrosine kinase receptor signaling pathway — 2/9
angiogenesis — 5/303
peptidyl−tyrosine phosphorylation — 4/164
tissue homeostasis — 4/165
cell population proliferation — 6/565
DNA−dependent DNA replication — 3/95
DNA replication initiation — 2/24
positive regulation of neutrophil chemotaxis — 2/31

−log$_2$FDR

Figure 3

Figure 4

A

RENDERS_YvsO

Enrch.Score = 60.982
Rank.Score = 412

Enrichment Score

Rank Score

1.h-Chambers
2.Norddahl
3.Bersenev
4.Wahlestedt
5.Sun
6.Flach
7.Kowalczyk
8.Grover
9.Kirschner
10.Maryanovich
11.Mann
12.Renders
13.Aging Signature (AS)

B

WONG_ESC_CORE

Enrch.Score = −0.689
Rank.Score = 4

KIM *et al.;*

Enrch.Score = 39.997
Rank.Score = 674

MARYANOVICH_DO

Enrch.Score = 45.204
Rank.Score = 746

CHALLEN_DNMT3_KO

Enrch.Score = 6.891
Rank.Score = 256

WONG_ESC_CORE (-0.6)   KIM *et al.;*(40)
Challen (6.8)   MARYANOVICH_DO (45.2)
AS(95.47)

Enrichment Score

Challen (-415) WONG_ESC_CORE (-213)
KIM *et al.;*(674)
MARYANOVICH_DO (722)
AS(3171)

Rank Score

C

Young cells   Aged cells   Merge

AS score
older
1.00
0.75
0.50
0.25
0.00
younger

cell
● young
▲ aged

AS score
Young   Aged

Figure 5

Figure 6