Ebselen Reacts with SARS Coronavirus-2 Main Protease Crystals

Tek Narsingh Malla[1] , Suraj Pandey[1] , Ishwor Poudyal[1] , Denisse Feliz[2] , Moraima Noda[2] ,
George Phillips[3], Emina Stojkovic[2], Marius Schmidt[1]

University of Wisconsin-Milwaukee, Milwaukee, WI
Northeastern Illinois University, Chicago, IL
Rice University, Houston, TX
7

Abstract. The SARS coronavirus 2 main protease 3CLpro tailor cuts various essential virus
proteins out of long poly-protein translated from the virus RNA. If the 3CLpro is inhibited, the
functional virus proteins cannot form and the virus cannot replicate and assemble. Any compound
that inhibits the 3CLpro is therefore a potential drug to end the pandemic. Here we show that the
diffraction power of 3CLpro crystals is effectively destroyed by Ebselen. It appears that Ebselen
may be a widely available, relatively cost effective way to eliminate the SARS coronavirus 2.

Introduction.

Figure 1. SARS CoV-2 3CLpro in its functional, dimeric form[1]. The two subunits A and B are shown in dark and light blue. An α-ketoamide inhibitor is bound to the active site (red box, enlarged). The His-41/Cys-145 catalytic dyad is marked.

During the COVID-19 pandemic in the last half year, a large body of literature became available
to understand and combat the SARS coronavirus-2 (CoV-2). Due to its importance for virus protein
maturation the SARS CoV-2 main protease (Mpro), also called 3CLpro, became a major drug
target. In its functional form the 3CLpro is a homo dimer with His/Cys dyads (His-41 and Cys-
145) in the active centers[1]. The structure of the CoV-2 3CLpro has been solved recently (Fig. 1)
guided by high similarity to other coronavirus 3CLpros[1]. A large number (>500) of SARS CoV-2
3CLpro structures were recently deposited in the protein data bank mainly following a fragment
screening study at the synchrotron Diamond near Oxford, England. Binding of fragments not only
to the active center was observed. They all constitute a database of potential drugs to target the
3CLpro. Apart from the fragments, the most promising compounds are the α-ketoamides (Fig. 1)
which bind tightly to the 3CLpro[1-3]. As they are complicated to synthesize they carry a hefty price

tag (companies ask for 14,000 dollars/g as recently inquired by one of the authors). A less
expensive, but also less known compound that binds to the 3CLpro is Ebselen[4]. Ebselen is a
selenium compound (Scheme 1) currently tested for a number of diseases such as bipolar disorder
and hearing loss[4]. Selenium is an essential metal, but toxic in higher doses. Ebselen has been shown
to bind strongly to the CoV-2 3CLpro[4], but the structure of the complex is unknown. Here we
show what happens when Ebselen is added to 3CLpro crystals.

Methods.

Scheme 1. Chemical structure of Ebselen, an organo-metallic compound.

Expression. The CoV-2 3CLpro sequence was synthetized (GenScript) for optimized expression in E. coli according to sequence information published previously[1]. In short, the N-terminus of 3CLpro is fused to glutathione-S-transferase (GST). It further has a 6-His tag at the c-terminus. The N-terminal GST will be autocatalytically cleaved off after expression due to an engineered 3CLpro cleavage sequence. The His tag can cleaved off by a PreCission Protease. Overexpression and protein purification protocols were modified from previous reports. E. coli were grown to 0.8 OD_{600} at 37°

in terrific broth. Expression was induced by 1 mmol/L IPTG at 25° C. After 3 h of expression, the
culture was induced a second time (1 mmol/L IPTG), and shaken overnight at 20° C. The yield is
about 80 mg for a 6 L culture. Cells were resuspended in lysis buffer (20mM Tris Base, 150
mmol/L NaCl, pH 7.8.). After lysis of the bacterial cells, debris was centrifuged at 50,000 g for 1
hour. The lysate was let stand at room temperature for at least 3 h (overnight is also possible).
After this, the lysate was pumped through a column containing 15 mL of Talon Cobalt resin
(TAKARA). The resin was washed without using imidazole using a wash cycle consisting of low

Figure 2. Low resolution (~3.5 Å in the corner) diffraction patterns of SARS CoV-2 3CLpro. (a) no ligand, (b) with Ebselen. Inserts: crystal images w/o and w Ebselen, respectively

N-1

C-306

His164   Glu166

Cys145

His41

Figure 3. Structure of the SARS CoV-2 3CLpro without Ebselen as determined here. Electron density (1.2 sigma) is shown for the catalytic dyad. Other active site residues are also marked.

salt (20 mmol/L Tris Base, 50 mmol/L NaCl, pH 7.8), high salt (20 mmol/L Tris Base, 1 mol/L NaCl, pH 7.8) and low salt (as above) solutions (about 20 column volumes each). After the wash cycle was completed, the column was let stand for an additional 2 h at room temperature followed by another wash cycle. The final product was eluted by 300 mmol/L imidazole, dialyzed immediately in 20 mmol/L Tris base, 150 mmol/L NaCl, 0.1 mmol/L dithiotreitol (DTT), pH 7.8, and concentrated to 20 mg/mL. Note, the c-terminal 6-His tag was not cleaved off. Due to this one step purification protocol, only 24 h are required from cell lysis to the pure 3CLpro product. The product is within 1.7 Da of the theoretical molecular weight as determined by mass spectroscopy.

Crystallization. The concentrated 3CLpro (with the His-tag left on) was diluted to 4 mg/mL. 100 μL of the diluted 3CLpro was mixed (1:1) in batch mode with the same amount of 25 % PEG 3350, Bis-Tris 100 mmol/L, pH 6.5. A few days later, nicely shaped crystals with dimensions of about 200 x 30 x 30 μm^3 were appearing. Crystals were soaked in mother liquor by adding Ebselen in powder form as Ebselen is not very soluble in water. Enough Ebselen that would otherwise produce a 50 mmol/L solution was added. After 2 days of soaking, the crystals

completely maintained their morphology (Fig. 2b).

Data Collection and Structure determination. The crystals were mounted in Mitegen microloops
(30 - 50 μm) and directly frozen in pucks suspended in liquid nitrogen for automated (robotic) data
collection. The dewar with the pucks were driven to the Advanced Photon Source for robotic data
collection at Sector-19 (Structural Biology Center, SBC, beamline 19-ID-D). Data collection was
fully remote due to restriction of the COVID-19 pandemic. Fig. 2 shows a comparison of
diffraction patterns collected from untreated crystals (Fig. 2a), and crystals treated with the Ebselen
(Fig. 2b). As the untreated crystals diffracted beyond 2 Å, the treated crystals did not show any
Bragg reflections whatsoever, even at lowest resolution. They are completely amorphous, despite
the nice crystal-like shape. Accordingly, the structure of only the untreated 3CLpro can be solved.
A dataset to 2.2 Å was collected (0.5° rotation and 0.3 s exposure per detector readout for a total

of 180º). Data was processed with HKL3000[5]. Data statistics in shown in Tab. 1. Initially the
spacegroup was found to be C2 (monoclinic centered). It was further determined by the CCP4
program[6] *pointless* that spacegroup I2 (monoclinic body centered) is more suitable, in accordance
with published results[7]. The 3CLpro structure with pdb access code 6WQF[7] was used as initial
model. Molecular replacement was not necessary. The model fits immediately and can be used for
refinement. Refinement was done using refmac[8] (version 5.8.0238). The structure is shown in Fig.
3. Its mean square deviation from model 6WQF is about 0.4 Å with 0.5 Å standard deviation. This
means that the 3CLpro cryo structure with the His-tag and the room temperature structure are
identical within the resolution limit. The structure of the His-tag could not be determined. There
is, however, a large crystal cavity near the C-terminus that might accommodate (a very disordered)
6 histidine structure.

Results and Discussion. A very important result is that c-terminal PreCission His-tag cleavage is
not required to obtain 3CLpro crystals. This speeds up purification to the extent that now gram
sized 3CLpro preparations can be obtained in a relatively short period of time. This is a prerequisite
for mix-and-inject approaches at free electron lasers with gas-dynamic-virtual-nozzle type mixing
injectors[9,10]. At XFELs low concentrations of Ebselen can be mixed with the crystalline slurry
which is quickly injected into the X-ray beam. There is hope that crystal decomposition is slower
than the delay between mixing and injection. Then, the structure of the Ebselen-3CLpro complex
may be determined. In addition, microcrystals also tend to be unusually flexible and may survive
large unit cell changes[11], which might make these experiments even more conceivable.

Since Ebselen has this strong effect on 3CLpro crystals, it a clear indication for the tight affinity
of this compound to the 3CLpro. Binding of Ebselen to the catalytic dyad has been already
identified by mass-spectroscopy and published earlier[4]. However, also after the experiments
reported here, the 3CLpro-Ebselen structure remains elusive. As Ebselen is a potent drug (and it
is not too toxic even in higher concentrations[4]), it has to be seen whether it can be used as an
effective weapon against the SARS CoV-2, and maybe all other pathogenic coronaviruses.
Whether it is suitable for a drug must be determined by physicians in a clinical setting perhaps in
combination with a SARS CoV-2 RNA dependent RNA polymerase inhibitor such as remdesivir[12].
As crystallographers and structural biologists, though, we can deliver evidence for a potential
mechanism as shown recently by others for the α-ketoamides[1,2] and also remdesivir[12].

This manuscript is published without delay on BioRxiv to disseminate rapid methods for
preparation, purification and structure determination of 3CLpro to a wide community. In addition,
this manuscript intends to alert and increase awareness of the potential of Ebselen as a 3CLpro
inhibitor and its value perhaps in combination, as the authors are not aware that anything is
reported about it in the public press. The coordinates of the 100 K 3CLpro structure as determined
here are available on request.

Acknowledgement. This work is supported by NSF grant STC 1727290 (BioXFEL) and NSF
grant RAPID 2030466. Results shown in this report are derived from work performed at Argnonne
National Laboratory (ANL), Structural Biology Center (SBC) at the Advanced Photon Source
(APS), under U.S. Department of Energy, Office of Biological and Environmental Research

contract DE-AC02-06CH11357. The authors thank Darren A. Sherrell for setting up the
experiment at the beamline.

Table. 1 Data collection and refinement statistics*

	SARS CoV-2 3CLpro
Beamline	19-ID-D, APS
Resolution	2.2 Å
Temperature	110 K
Space group	I2
Unit-cell parameters (,º)	a = 44.7 Å b = 53.4 Å c= 112.3 Å $\alpha=90^{\circ}$ $\beta=100.0^{\circ}$ $\gamma=90^{\circ}$
No of unique reflections	9426
Redundancy	3.1 (1.9)
Completeness (%)	70.0
CC1/2 (%)	99.8 (47.6)
R_{merge} (%)	7.3 (76.5)
Refinement	
R_{cryst}/R_{free} (%)	19.1/26.3
RMSD to 6WQF[7]	0.39 +/- 0.53 Å

138   *last resolution shell in brackets

References.

Zhang, L. L. *et al.* Crystal structure of SARS-CoV-2 main protease provides a basis for design of improved alpha-ketoamide inhibitors. *Science* **368**, 409-+.

Zhang, L. L. *et al.* alpha-Ketoamides as Broad-Spectrum Inhibitors of Coronavirus and Enterovirus Replication: Structure-Based Design, Synthesis, and Activity Assessment. *Journal of Medicinal Chemistry* **63**, 4562-4578.

Dai, W. *et al.* Structure-based design of antiviral drug candidates targeting the SARS-CoV-2 main protease. *Science* **368**, 1331-1335, PMC7179937.

Jin, Z. M. *et al.* Structure of M-pro from SARS-CoV-2 and discovery of its inhibitors. *Nature* **582**, 289-+.

Minor, W., Cymborowski, M., Otwinowski, Z. & Chruszcz, M. HKL-3000: the integration of data reduction and structure solution - from diffraction images to an initial model in minutes. *Acta Crystallographica Section D-Structural Biology* **62**, 859-866.

Winn, M. D. *et al.* Overview of the CCP4 suite and current developments. *Acta Crystallogr D* **67**, 235-242.

Kneller, D. W. *et al.* Structural plasticity of SARS-CoV-2 3CL M-pro active site cavity revealed by room temperature X-ray crystallography. *Nat Commun* **11**.

Murshudov, G. N. *et al.* REFMAC5 for the refinement of macromolecular crystal structures. *Acta crystallographica. Section D, Biological crystallography* **67**, 355-367, 3069751.

Olmos, J. L., Jr. *et al.* Enzyme intermediates captured "on the fly" by mix-and-inject serial crystallography. *BMC Biol* **16**, 59, PMC5977757.

Calvey, G. D., Katz, A. M. & Pollack, L. Microfluidic Mixing Injector Holder Enables Routine Structural Enzymology Measurements with Mix-and-Inject Serial Crystallography Using X-ray Free Electron Lasers. *Anal Chem* **91**, 7139-7144.

Stagno, J. R. *et al.* Structures of riboswitch RNA reaction states by mix-and-inject XFEL serial crystallography. *Nature* **541**, 242-246.

Yin, W. C. *et al.* Structural basis for inhibition of the RNA-dependent RNA polymerase from SARS-CoV-2 by remdesivir. *Science* **368**, 1499-+.