A multi-hierarchical approach reveals D-serine as a hidden substrate of sodium-coupled monocarboxylate transporters

Pattama Wiriyasermkul[1,2,3], Satomi Moriyama[3], Masataka Suzuki[4], Pornparn Kongpracha[1,2], Nodoka Nakamae[3], Saki Takeshita[3], Yoko Tanaka[3], Akina Matsuda[4], Masaki Miyasaka[1,2], Kenji Hamase[5], Tomonori Kimura[6,7], Masashi Mita[8], Jumpei Sasabe[4], Shushi Nagamori[1,2,3]*

[1]Center for SI Medical Research, [2]Department of Laboratory Medicine, The Jikei University School of Medicine, Tokyo, Japan

[3]Department of Collaborative Research for Biomolecular Dynamics, Nara Medical University, Nara, Japan

[4]Department of Pharmacology, Keio University School of Medicine, Tokyo, Japan

[5]Graduate School of Pharmaceutical Sciences, Kyushu University, Fukuoka, Japan

[6]KAGAMI Project, [7]Reverse Translational Research Project, Center for Rare Disease Research, National Institutes of Biomedical Innovation, Health and Nutrition (NIBIOHN). Osaka, Japan

[8]KAGAMI Inc., Osaka, Japan

* Corresponding author: Shushi Nagamori

Email: snagamori@nagamori-lab.jp

Keywords

transporter, amino acid, kidney diseases, micronutrient, biomarker

Abstract

Transporter research primarily relies on the canonical substrates of well-established transporters. This approach has limitations when studying transporters for the low-abundant micromolecules, such as micronutrients, and may not reveal physiological functions of the transporters. While D-serine, a trace enantiomer of serine in the circulation, was discovered as an emerging biomarker of kidney function, its transport mechanisms in the periphery remain unknown. Here, using a multi-hierarchical approach from body fluids to molecules, combining multi-omics, cell-free synthetic biochemistry, and *ex vivo* transport analyses, we have identified two types of renal D-serine transport systems. We revealed that the small amino acid transporter ASCT2 serves as a D-serine transporter previously uncharacterized in the kidney and discovered D-serine as a noncanonical substrate of the sodium-coupled monocarboxylate transporters (SMCTs). These two systems are physiologically complementary, but ASCT2 dominates the role in the pathological condition. Our findings not only shed light on renal D-serine transport, but also clarify the importance of non-canonical substrate transport. This study provides a framework for investigating multiple transport systems of various trace micromolecules under physiological conditions and in multifactorial diseases.

Introduction

Membrane transport proteins are key molecules to control the homeostasis of micromolecules. To date, over 450 solute carriers (SLCs), 40 ATP-binding cassettes (ABCs), 60 ATPase pumps, and 300 ion channels have been identified in mammals (Ranjan et al., 2011; César-Razquin et al., 2015; Perland and Fredriksson, 2017; The UniProt Consortium et al., 2021). Approximately 30% of membrane transport proteins remain uncharacterized. While some membrane transport proteins can be classified based on sequence similarities and transport substrates of well-studied family members, this approach is limited when it comes to discovering transporters for some micromolecules, such as micronutrients and their metabolites. Many of these micromolecules have not been extensively studied and do not fit into traditional nutrient types. Currently, the importance of micronutrients and gut microbiota-derived metabolites in health and disease has gained a high impact (Shenkin, 2006; Vernocchi et al., 2016; Liu et al., 2022), yet their responsible transporters are not well understood. Understanding the absorption and reabsorption mechanisms of dietary micronutrients and gut microbiota-derived metabolites will provide a key molecular process for controlling their dynamics and molecular activities.

This study aims to establish an approach to investigate the physiological function and pathological significance of transporters, particularly transport systems for trace micromolecules. As a paradigm, we investigated renal transport systems for D-amino acids, micronutrients derived mostly from the diet and some from the gut microbiota. Amino acid transport systems have been well studied (Bröer, 2008a), but have focused on L-amino acids, which are macronutrients, leaving open the question of how D-amino acids are transported. We have developed two-dimensional high-performance liquid chromatography (2D-HPLC), which allows the segregation of L- and D-enantiomers in liquid biopsy (Miyoshi et al., 2009; Hamase et al., 2010). The importance of D-amino acids in renal function has received increasing

attention (Sasabe et al., 2014; Kimura et al., 2016; Sasabe and Suzuki, 2018; Nakade et al.,

2018; Hesaka et al., 2019; Kimura et al., 2020; Okushima et al., 2021; Iwata et al., 2022; Suzuki

et al., 2022). D-Serine in the body fluid was discovered as a promising biomarker for acute

kidney injury (AKI) and chronic kidney disease (CKD) due to its clear association with kidney

function in rodents and humans (Sasabe et al., 2014; Kimura et al., 2016; Nakade et al., 2018;

Hesaka et al., 2019; Okushima et al., 2021). Diseased-associated alterations in plasma and

urinary D-/L-serine ratio suggest that renal D-serine transport systems are different and distinct

from L-serine transport systems under physiological and pathological conditions. The transport

of D-serine has long been found to take place at the proximal tubules (Kragh-Hansen and

Sheikh, 1984; Sasabe and Suzuki, 2018; Shimomura et al., 1988; Silbernagl et al., 1999), yet

the corresponding transporter has not been clearly elucidated.

D-Serine is found at submillimolar levels in the brain (Hashimoto et al., 1992), where

it acts as an obligatory physiological co-agonist of N-methyl-D-aspartate receptors (NMDARs)

(Mothet et al., 2000; Basu et al., 2009; Papouin et al., 2012; Wolosker, 2018). Extensive studies

of D-serine in the brain show that D-serine is transported by four plasma membrane

transporters: ASCT1 (SLC1A4), ASCT2 (SLC1A5), Asc-1 (SLC7A10), and SNAT1

(SLC38A1) (Rosenberg et al., 2013; Foster et al., 2016; Kaplan et al., 2018; Bodner et al.,

2020). In contrast, D-serine is detected at low micromolar levels in the periphery, and little is

known about its dynamics as well as transporting systems. Mammals acquire D-serine by

biosynthesis via serine racemase function (Wolosker et al., 1999) and absorption from the diet

and gut microbiota presumably via intestinal transport system(s) (Sasabe and Suzuki, 2018;

Nakade et al., 2018; Gonda et al., 2023). Similar to the renal epithelia, a D-serine transporter

in the intestine is also unknown.

In this study, we integrated multiple analytical methods for the different biological

hierarchies. We applied 2D-HPLC to measure plasma and urinary enantiomers of 20 amino

acids of ischemia-reperfusion injury (IRI) mice, a model of AKI and AKI-to-CKD transition

(Sasabe et al., 2014; Fu et al., 2018). Membrane proteomics of the renal proximal tubules of

the IRI mouse models was performed to explore the key molecules responsible for the

alterations of D- and L-amino acid transport. With bioinformatics, cell-based screening and

cell-free transport analysis, we identified two types of D-serine transport systems in the kidney.

Subsequently, *ex vivo* transport analyses from the normal kidney and the IRI model explained

the transport mechanism of D-serine as a biomarker in kidney diseases.

Results

Amino acid metabolomics in plasma and urine of the renal IRI model

To understand kidney-associated alteration of enantiomeric amino acids, we measured L- and D-isomers of 20 amino acids in the body fluids of the IRI model (Figure supplements 1-2). Dynamics of L- and D- enantiomers in the plasma were evaluated from the ratios of D-/L-enantiomers (Figure 1A). Among amino acids, only the ratios of plasma D-/L-serine were increased in a time dependent-manner from 4 – 40 hours after the ischemia-reperfusion (Figure 1A-B). The elevation of plasma D-/L-serine ratios at early time points (4h – 8h) was due to a sharp decrease of the L-isomer, while the rise of D-/L-serine ratios at late time points (20h – 40h) was a result of a continuous acceleration of the D-isomer (Figure 1C-D). The results of plasma D-serine were consistent with previous observations (Sasabe et al., 2014). In addition to D-serine, we detected certain amounts of D-alanine, D-proline and D-methionine in the plasma (Figure supplement 1A). However, their D-/L- enantiomeric ratios were uncorrelated to the pathological conditions (Figure supplement 1A). Our knowledge of traditional amino acid transport suggests that alanine, serine and proline share the same transport systems (Bröer, 2008a). The noticeably different profile of plasma D- and L-serine from those of alanine and proline led to the speculation that D-serine transport was possibly mediated by unique transport systems, different from those of D-alanine, D-proline, and the L-isomers. Moreover, such transport systems may be sensitive to the IRI conditions.

We measured the enantiomeric amino acid profiles from urine samples of the IRI model (Figure supplement 2). While plasma is a reliable resource of metabolite biomarkers, urine is a meaningful non-invasive diagnostic indicator of kidney function. More importantly, the analysis of urinary metabolites is indispensable for elucidating the unknown reabsorption systems of certain substances in the kidney. From our results, the ratios of urinary D-/L-serine

were drastically decreased at early IRI (4h – 8h) (Figure 1E-F). Similar tendencies of D-/L-amino acid ratios were observed in alanine, proline, asparagine, and histidine (Figure 1E-H). These urinary amino acid profiles indicated the impairment in both D- and L-amino acid reabsorption since the early injury. We hypothesized that transporters play a major role in the dynamics of D- and L-serine in the kidney with respect to changes in the D-/L-serine ratio during pathology.

Proteomics for membrane transport proteins from renal brush border membranes of the IRI model

We aimed to identify the corresponding molecules for the dynamics of D- and L-serine. To obtain the candidates, we performed proteomic analysis of renal brush border membrane vesicles (BBMVs) (the membrane fraction enriching apical membrane proteins of proximal tubular epithelia) from the IRI model. We analyzed the BBMVs from 4 – 8 hours after IRI, the times of rigorous shift of plasma and urinary D-/L-serine ratio (Figure 1B, F). The proteomes of the IRI samples were calculated as ratios of the negative control group (sham), yielding two groups of comparative proteome data: 4h IRI/sham (4h) and 8h IRI/sham (8h). In total, 4,423 proteins were identified, of which 1,187 proteins (27% of the total detected proteins) were categorized as plasma membrane proteins and extracellular matrix proteins (Table supplement 1). We observed a significant increase of two well-known early AKI biomarkers: Ccn1 (Cyr61: matrix-associated heparin-binding protein; 4h IRI/sham = 41.4 folds; 8h IRI/sham = 29.9 folds) and Ngal (Lcn2: Neutrophil gelatinase-associated lipocalin; 4h IRI/sham = 1.8 folds; 8h IRI/sham = 9.5 folds). This finding confirmed the reliability of our proteomics for molecular analysis of early AKI (Table supplement 1) (Marx et al., 2018).

From the proteome, we detected 398 membrane transport proteins (325 transporters and 73 ion channels). Analysis of all membrane transport proteins by Ingenuity Pathway Analysis

(IPA) revealed alterations of the biological function in *Molecular Transport*, which are

categorized by the types of their canonical substrates (Table supplement 2). Transport of "all

micromolecules" was shown to be drastically decreased from 4h IRI (Figure 2A: Top). This

decrease was due to the dramatic reduction of transport systems for heavy metals and organic

compounds (Figure 2A). Figure 2B shows membrane transport proteins corresponding to the

annotation in Figure 2A. This proteomic result unveiled the whole picture of the molecular

targets of the injury. Furthermore, the results revealed the key membrane transport proteins

behind AKI metabolite biomarkers. It is noted that some transporters were upregulated,

suggesting the compensatory mechanism repair processes for cell survival from the injury

(Figure 2B).

ASCT2 is one of D-serine transporters at the apical membrane of renal proximal tubules

Given insight into transporters in the proteome, we observed the elevation of mouse

Slc1a5/Asct2, a known D-serine transporter in the brain, at both 4h and 8h IRI (Figure 2B:

Transport of Amino acids). Asct2 was previously detected in renal brush border membranes,

but its precise localization and function in the kidney have not been characterized (Avissar et

al., 2001; Scalise et al., 2018). We then examined the localization of Asct2 in the kidney by

using affinity-purified Asct2 antibodies that recognized its N-terminus (NT) or C-terminus

(CT) (Figure supplement 3A). Asct2 was partially co-immunostained with both Sglt2 (Slc5a2;

sodium/glucose cotransporter 2) and Agt1 (Slc7a13, aspartate/glutamate transporter 1), which

are apical membrane markers for S1 + S2 and S3 segments, respectively (Figure 3A-B)

(Nagamori et al., 2016a; Ghezzi et al., 2018). In contrast, Asct2 did not co-localize with

Na^+/K^+-ATPase, a basolateral membrane marker (Figure 3C). The results demonstrated that

Asct2 is localized at the apical side in all segments of proximal tubules.

To confirm the D-serine transport function of ASCT2 in human cells, we used wild-

type (WT) near-haploid human leukemic cancer cell line (HAP1) and HAP1 cells carrying

CRISPR/CAS-mediated *ASCT2* knockout (ASCT2-KO). D-serine uptake was significantly

decreased in the ASCT2-KO cells (Figure 3D). The function of ASCT2 was also verified in

human embryonic kidney HEK293 cells, which endogenously express ASCT2, by using

ASCT2-knockdown (Figure 3E).

Screening of candidate molecules for D-serine transporters

While we have shown that ASCT2 is a D-serine transporter at the apical membranes of

all proximal tubular segments, ASCT2 transports D-serine with high affinity (K_m of 167 µM in

oocyte system) but with weak stereoselectivity (Foster et al., 2016). These properties of ASCT2

differ from those of the reported D-serine transport in the kidney. First, D-serine transport

systems in S1 + S2 segments are reported to be different from those in the S3 segments and the

kinetics of both systems are in mM range (Kragh-Hansen and Sheikh, 1984; Silbernagl et al.,

1999). Silbernagl et al. also suggested that ASCT2 is not (or not only) a D-serine transporter at

S3 segment (Silbernagl et al., 1999). Second, the plasma and urinary D-/L-serine profiles from

CKD and AKI samples indicated the stereoselectivity of the transporter (Figure 1) (Sasabe and

Suzuki, 2018). Third, ASCT2 recognizes serine, alanine, and proline, but our enantiomeric

amino acid profile in plasma samples showed a distinct D-/L- ratio pattern of serine from alanine

and proline (Figure 1B). All data suggest the existence of (an)other D-serine transporter(s).

To search for the D-serine transporters, we chose the ratios of mouse Asct2 as the cutoff

values of our membrane proteomic data at both 4h and 8h IRI (Figure 4A). Membrane transport

proteins showing more alteration than Asct2 at both increased and decreased values were

selected because some transporters are known to switch roles upon pathology. Further, we

focused on the luminal transporters and omitted transporters that are reported to be located at

basolateral membranes and organelles as the BBMVs are enriched apical membrane fractions.

Membrane transport proteins that recognize only inorganic ions were also excluded. Finally,

ten candidates were selected (Table 1). Slc36a1/Pat1 and Slc6a18/B⁰at3, the known small

amino acid transporters, were included in the list, although their expressions only passed the

cutoff at only one-time point (either 4h or 8h IRI). Slc5a12/Smct2 was also included because

Smct1, another member of sodium-coupled monocarboxylate transporter (SMCT) family, was

selected as a candidate. Smct2 has a comparable role with Smct1, but both transporters are

localized at different segments of the proximal tubules.

To analyze the function of the candidate transporters, we selected HEK293 cell line for

the screening method because of its high transfection efficiency. Similar to other several

common cell lines, HEK293 cells express ASCT2 endogenously. However, we examined other

possible D-serine transport systems in HEK293 cells, aiming to reduce the transport

background. Proteomic analysis of the membrane fractions from HEK293 cells identified

amino acid transporters as shown in Table supplement 3. Transport activity of D-serine in

HEK293 cells showed Na^+ dependency and ASCT2-substrate mediated inhibition (L-Ser, L-

Thr, and L-Met) (Figure supplement 3B-C). Ben-Cys (S-benzyl-L-cysteine) and GPNA (L-γ-

glutamyl-p-nitroanilide) inhibited D-serine transport in HEK293 cells similar to that of ASCT2-

expressing cells (Bröer et al., 2016), but SLC38 inhibitor MeAIB (2-(methylamino)isobutyric

acid) had no effect (Figure supplement 3C). These results indicated that ASCT2 is the main D-

serine transporter in HEK293 cells.

Due to the absence of any ASCT2-specific inhibitor and the compensatory transport

mechanism that occurred in the ASCT2-deletion condition (Figure supplement 3E) (Bröer et

al., 2016), we avoided using the traditional cell-based transport assay for screening. Instead,

we established a new screening method based on cell growth. Apart from the studies on D-

serine biomarkers in kidney diseases, a previous study found that high concentrations (10 – 20

mM) of D-serine impaired the growth of a human proximal tubular cell line (Okada et al., 2017).

We then tested whether high concentrations of D-serine affect HEK293 cell growth. Unlike L-

serine which had no effect, D-serine reduced cell growth in a concentration-dependent manner

with IC_{50} of 17.4 ± 1.05 mM (Figure supplement 3D). In the ASCT2-knockdown cells, the

growth inhibition by D-serine was attenuated (Figure 4B), indicating that D-serine suppressing

cell growth was caused by ASCT2-mediated D-serine transport.

Based on this finding, we utilized cell growth determination assay as the screening

method even without ASCT2 knockdown. HEK293 cells were transfected with human

candidate genes. The transfected cells were treated with either 15 mM (near IC_{50} concentration)

or 25 mM (high concentration) D-serine for 2 days, and cell growth was determined. The

positive control Asc-1 transfected cells showed a reduction in cell growth, confirming the

effectiveness of this assay (Figure 4C-D). Among ten candidates, PAT1 (SLC36A1) and

B^0AT3 (SLC6A18) showed higher cell growth than the mock cells, suggesting that PAT1 and

B^0AT3 may reduce D-serine influx mediated by endogenous ASCT2. In contrast, SMCT1

(SLC5A8) and SMCT2 (SLC5A12) extended the growth suppression after treatment with both

15 and 25 mM D-serine (Figure 4C-D and supplement 4A-B). Thus, with the hypothesis that

the candidate molecule influxes D-serine and brings about growth reduction, we selected both

SMCT1 and SMCT2 for further analysis.

Characterization of D-serine transport in SMCTs

SMCTs are known to transport monocarboxylates such as lactate, propionate, and

nicotinate in a Na^+-dependent manner (Ganapathy et al., 2008). To characterize D-serine

transport in both SMCT1 and SMCT2, we generated Flp-In T-REx 293 cells stably expressing

human SMCT1 (FlpInTR-SMCT1) and SMCT2 (FlpInTR-SMCT2). Expression of SMCTs in

FlpInTR-SMCT1 and FlpInTR-SMCT2 was verified by Western blot using anti-FLAG

antibody to detect FLAG-tagged SMCTs (Figure supplement 5A). SMCT1 and SMCT2

expressions remained unchanged upon ASCT2 knockdown (Figure supplement 5A). The

function of SMCTs was confirmed by their canonical substrate [^{14}C]nicotinate transport in the

SMCT1- and SMCT2-stable cells (Figure supplement 5B-C).

The transport functions of SMCTs are inhibited by non-steroidal anti-inflammatory

drugs (NSAIDs; e.g., ibuprofen and acetylsalicylate) (Itagaki et al., 2006; Gopal et al., 2007).

To verify the contribution of SMCTs to D-serine-reduced cell growth, we added ibuprofen to

the growth assay. In contrast to mock, in which ibuprofen has no effect, the growth suppression

by D-serine treatment was gradually attenuated by ibuprofen in both FlpInTR-SMCT1 and

FlpInTR-SMCT2 cells, indicating that the cell growth suppression by D-serine treatment is the

result of SMCT1 and SMCT2 functions (Figure 5A).

We characterized SMCTs mediating D-[^{3}H]serine transport by using SMCT-stable cell

lines. Both FlpInTR-SMCT1 and FlpInTR-SMCT2 cells transported D-[^{3}H]serine over Mock

(Figure 5B) and the transport was enhanced when ASCT2 was knocked down (Figure 5C),

indicating the recognition of D-serine by both SMCT1 and SMCT2. The D-[^{3}H]serine transport

was inhibited by ibuprofen and acetylsalicylate, confirming the specific transport by SMCTs

(Figure 5D). The excess of non-radioisotope-labeled D-serine also inhibited D-[^{3}H]serine

uptake in both SMCTs. Still, its inhibitory effect in SMCT2 was much less than in SMCT1,

suggesting the lower affinity of D-serine in SMCT2 than in SMCT1 (Figure 5D).

D-Serine transport properties and kinetics in SMCT1

Kinetics of D-[^{3}H]serine transport in SMCT1 was measured in FlpInTR-SMCT1 cells

in the presence of ASCT2 knockdown. SMCT1 transported D-[^{3}H]serine in a concentration-

dependent manner and the curve fitted to Michaelis−Menten kinetics with the apparent K_m of

3.39 ± 0.79 mM and V_{max} of 18.23 ± 1.73 pmol/µg protein/min (Figure 5E and supplement 5D).

The substrate selectivity of SMCT1 was investigated using synthetic biochemistry to

avoid the interference of amino acid and carboxylate transports by endogenous transporters in

cells. We established a cell-free assay system using proteoliposomes, in which purified proteins

are reconstituted into liposomes and all substances in the system are controllable. We purified

human SMCT1 by affinity column and reconstituted SMCT1-proteoliposome (SMCT1-PL)

(Figure 6A). The function of SMCT1 in SMCT1-PL was verified by the transport of lactate,

propionate, and nicotinate but not urate (negative control) (Figure supplement 6A). The

kinetics of [^{14}C]nicotinate transport in SMCT1-PL revealed an apparent K_m of 442 ± 94 μM

(Figure supplement 6B), which is in a similar range to the K_m previously described by

electrophysiological experiments in *Xenopus* oocyte expression systems (K_m 390 μM) (Paroder

et al., 2006), confirming the functional properties of SMCT1 in our cell-free transport assay

system.

The transport of D-[^{3}H]serine in SMCT1-PL was Na$^+$-dependent and reached the

stationary phase at approximately 5 min (Figure 6B). The amount of D-serine transport was

slightly lower than lactate and propionate but higher than L-serine and the transport was

inhibited by ibuprofen, confirming SMCT1-mediated D-serine transport (Figure 6C). Amino

acid selectivity revealed that SMCT1 recognized both L- and D-serine over other small amino

acids (L- and D-alanine), acidic amino acids (L- and D-glutamate), and large neutral amino acids

(L- and D-tyrosine) (Figure 6D). Therefore, we concluded that serine is the substrate of SMCT1

and the recognition is more stereoselective for the D- than the L-isomer.

ASCT2 and SMCTs contribute to D-serine transport in renal proximal tubular epithelia

We next examined the contributions of ASCT2 and SMCTs to D-serine reabsorption in

the kidney using an *ex vivo* transport assay. ASCT2 is an antiporter, influx one amino acid with

efflux of another, while SMCTs are symporters (Ganapathy et al., 2008; Scalise et al., 2018).

Therefore, we measured D-serine transport in the presence or absence of L-Gln preloading and

with or without ibuprofen to distinguish the functions of mouse Asct2 and Smcts. The

enantiomeric amino acid profile showed D-serine concentrations of $4 - 10$ μM in the plasma

samples (Figure supplement 1) and $20 - 100$ μM in the urine samples (Figure supplement 2).

Accordingly, we tested 10 μM D-[^{3}H]serine transport in BBMVs derived from normal mice. In

the non-preloading condition, D-[^{3}H]serine uptake in Na^+ dependence gradually but

continuously increased and the uptake was remarkably inhibited by ibuprofen, suggesting that

the transport was mainly attained from Smcts functions (Figure 7A). In L-Gln preloaded

BBMVs, D-[^{3}H]serine uptake in Na^+-dependence arose quickly and reached the saturated point

at $1 - 2$ min (Figure 7B). The uptake at early time points (≤ 30 sec) was ibuprofen-insensitive

but became partly ibuprofen-sensitive from 1 min to the stationary phase (Figure 7B). These

results indicate that the D-serine transport was mediated by both Asct2 and Smcts. At early

time points, D-[^{3}H]serine was transported by Asct2, which occurred highly and rapidly.

Meanwhile, Smcts functions (ibuprofen-sensitive) initiated later but had prolonged functions

as seen in the non-preloading condition. Taken altogether, we suggested that, in the normal

kidney where renal proximal tubular epithelial cells contain intracellular L-Gln, D-serine

reabsorption is derived from combinational functions of both ibuprofen-sensitive (e.g. Smcts)

and ibuprofen-insensitive (e.g. Asct2) transporters.

The localization of Smct1 in the kidney was unclear. Gopal et al. reported the

localization of Smct1 at S3 and Smct2 in all segments, while single-nucleus RNA sequencing

(snRNA-Seq) showed Smct1 at S2 – S3 and Smct2 at S1 segment (Figure supplement 7:

Controls) (Gopal et al., 2007; Kirita et al., 2020). We generated an anti-Smct1 antibody and

examined the localization of Smct1. The result showed that Smct1 was mainly localized at the

S3 and slightly at the S1 + S2 segments (Figure supplement 8).

D-Serine transport in renal proximal tubular epithelia of IRI model

Under IRI conditions, we observed low urinary but high plasma levels of D-serine, suggesting high D-serine reabsorption (Figure supplements 1A, 2A). Our membrane proteomics revealed that Asct2 was increased while and Smct1 and Smct2 were decreased in both 4h and 8h IRI (Table 1). We then examined the functional contributions of Asct2 and Smcts on the D-serine reabsorption during the pathology. Similar to those of the normal BBMVs (Figure 7A-B), D-[^{3}H]serine transport in Sham operation showed ibuprofen-sensitive in non-preloading and partly ibuprofen-sensitive in L-Gln preloading conditions (Figure 7C: Sham). D-[^{3}H]Serine influx reduced sharply in the BBMVs of 4h IRI model, and the transport was ibuprofen-insensitive in both with and without L-Gln preloading (Figure 7C: 4h IRI). Likely, this feature was due to the slight increase of Asct2 and the dramatic decrease of Smcts as observed in the proteome data (Asct2: 4h IRI/sham = 1.29 fold, Smct1: 4h IRI/sham = 0.59 fold; Smct2: 4h IRI/sham = 0.86 fold; Table supplement 1). We verified that BBMVs from all conditions contain functional proteins by measuring L-[^{3}H]aspartate transport, which represented the function of Slc1a1/Eaac1 (Kanai and Hediger, 1992; Bailey et al., 2011). The assay confirmed that BBMVs derived from 4h IRI model were not fully damaged, although the low influx was likely from the loss of protein expression during the injury (Figure 2B: Amino acids, Figure supplement 9). In the BBMVs of 8h IRI model, D-[^{3}H]serine transport was exceedingly high in L-Gln preloading and it was ibuprofen-insensitive (Figure 7C: L-Gln preloaded). These results could be explained by the continuing increase of Asct2 and decrease of Smcts as revealed by our proteomics (Asct2: 8h IRI/sham = 1.64 fold; Smct1: 8h IRI/sham = 0.62 fold; Smct2: 8h IRI/sham = 0.78 fold) (Table 1, Table supplement 1). The slight increase in D-[^{3}H]serine transport in non-preloaded 8h IRI BBMVs may be due to an unknown transporter (Figure 7C: No preload). Our amino acid profiling showed that the amount of plasma and urinary L-Gln were not largely altered during 4h – 8h IRI (Figures 1B, 2B),

suggesting that renal epithelial cells maintain intracellular L-Gln. Most likely, the high

reabsorption of D-serine during IRI resulted from the increase of Asct2, despite the decreases

of Smct1 and Smct2.

For comparison with our proteome data, we analyzed the mRNA expressions of Asct2,

Smct1, and Smct2 in proximal tubule clusters at the early IRI from the open-source snRNA-

seq dataset (Kirita et al., 2020). Consistent with our proteome data, Smct1 and Smct2 mRNA

expressions were dramatically decreased, whereas Asct2 was increased since the early IRI

(Figure supplement 7). The correlation of proteomics and transcriptomics suggested that the

protein alterations of Asct2 and Smcts during the pathology emerged from transcriptional

mechanisms rather than post-translational protein processing or protein degradation.

Discussion

In this study, we presented an approach to investigate the transport systems for micronutrients and metabolites under physiological conditions and in multifactorial diseases. We selected AKI as a model study because AKI is known to target renal proximal epithelia and disrupt absorption/reabsorption systems that are the results of cooperative functions of multiple transporters. We utilized amino acid metabolomics to reveal a unique feature of the enantiomeric dynamics of serine in the AKI and analyzed membrane proteomes to obtain the D-serine transporter candidates. We then applied cell-based and cell-free transport assays to identify two D-serine transport systems; one was ASCT2 and the other consisted of unanticipated SMCTs (SMCT1 and SMCT2). Using *ex vivo* analysis of apical membrane-enriched renal proximal tubules, we showed that both transport systems contributed comparably to D-serine handling in the normal kidney, but ASCT2 became dominant in AKI, leading to the increase of D-serine in the blood. The alteration of two transport systems explained the dynamics of D-serine and suggested the transport mechanism behind the D-serine biomarker.

Several studies have reported the non-canonical substrates of membrane transport proteins, such as the recognition between amino acids, carboxylates and amines among SLCs (Metzner et al., 2005; Matsuo et al., 2008; Schweikhard and Ziegler, 2012; Wei et al., 2016). Serine consists of a hydroxypropionic acid and an amino group at the α-carbon. Possibly, the hydroxypropionic acid on serine is the main part to interact with SMCTs because hydroxypropionic acid shares similar moieties to lactic acid and propionic acid which are the high-affinity substrates of SMCTs. The apparent K_m of D-serine transport in SMCT1 is 3.39 mM (Figure 5E), which is within the range of known SMCT1 substrates (0.07 – 6.5 mM) (Ganapathy et al., 2008), suggesting that SMCT1 accepts D-serine in the same manner as other monocarboxylate substrates. Inhibition of D-[^{3}H]serine transport by non-radioisotope labeled

D-serine in SMCT2 was less effective than in SMCT1 (Figure 5D). Together with the previous

report that SMCT2 had lower inhibition affinities for nicotinate, lactate and butyrate than

SMCT1 (Srinivas et al., 2005), we suggest that SMCT2 recognizes D-serine with lower affinity

than SMCT1.

Kinetics analysis of D-serine transport revealed the high affinity by ASCT2 (K_m 167

µM) (Foster et al., 2016) and low affinity by SMCT1 (K_m 3.39 mM; Figure 5E). Besides

transport affinity, the expression level is another important factor in evaluating the transport

capacity. Regarding chromatogram intensities in the proteomics data, we found that the

intensities of Smct1 (2.9×10^9 AU) and Smct2 (1.6×10^8 AU) were remarkably higher than

Asct2 (1.5×10^7 AU) in the control mice (Table 1: abundance in Sham). Although it is not

accurate to compare intensities between different proteins in a mass spectrometry analysis, the

comparison may roughly reflect the protein amounts. Together with our proteome data, this

observation is consistent with the snRNA-seq result (Figure supplement 7: Controls),

suggesting that Asct2 is only expressed in minor cell populations, while Smct1 and Smct2

expressions are high and ubiquitous in the normal conditions. Thus, we proposed that ASCT2

transports D-serine with low capacity due to its low expression level, while SMCTs have high

transport capacity due to their abundance. Consequently, both ASCT2 and SMCTs are likely

to contribute to D-serine reabsorption in the normal kidney as seen in the *ex vivo* assay (Figure

7B).

Several pieces of evidence support SMCT1 at S3 segment and SMCT2 at S1 + S2

segments on renal D-serine handling. First, our enantiomeric amino acid profile from the IRI

model, which agrees with the previous reports in CKD patients and AKI animal models,

suggested the existence of a stereoselective transport system for serine (Figure 1) (Hesaka et

al., 2019; Kimura et al., 2020, 2016; Sasabe et al., 2014; Sasabe and Suzuki, 2018). The system

seems distinct from the known classical small amino acid transport systems and ASCT2.

Second, previous studies indicated renal D-serine transport takes place at the proximal tubules

by the distinct transport systems between S1 + S2 and S3 segments. Both systems exhibited

the characteristics of Na^+ dependency, electrogenicity, low affinity (mM range), and partial

stereoselectivity (Kragh-Hansen and Sheikh, 1984; Shimomura et al., 1988; Silbernagl et al.,

1999). These properties of the D-serine transport suit well to SMCTs and convince us of the

contribution of SMCTs to renal D-serine transport in addition to ASCT2. Moreover, continuous

administration of high doses of D-serine induces nephrotoxicity at the S3 segment of proximal

tubules in normal animals (Silbernagl et al., 1999; Morehead et al., 1945; Hasegawa et al.,

2019). It is most likely that the D-serine-induced proximal tubular damage is partly due to the

absorption of D-serine by SMCTs, in particular by SMCT1 at the S3 segment. It is noted that

SMCT2 is reported as a protein indicator for the kidney repair during the injury. Prolonged

SMCT2 down-expression indicates the failed repair of proximal tubular cells in IRI (Kirita et

al., 2020). The significance of SMCT2, in turn, supports the merit of D-serine usage in AKI

and CKD diagnosis.

Combining all the results, we propose the model of D-serine transport systems in renal

proximal tubules (Figure 8). In the normal kidneys, the expression levels of SMCT2 at S1 + S2

segments and SMCT1 at S3 segment are high, whereas ASCT2 is low. Both SMCTs and

ASCT2 are involved in D-serine reabsorption. Nonetheless, the net reabsorption levels appear

to be relatively low due to the low affinities of SMCTs. It is likely that reabsorption at the S3

segment is dominated by SMCT1 function. The reabsorbed D-serine in proximal tubular cells

is degraded into OH-pyruvate by D-amino acid oxidase (DAAO), thereby maintaining the low

plasma D-serine but high urinary D-serine (Figure 8: Normal). In pathological conditions such

as AKI and CKD, the expression levels of SMCT1 and SMCT2 decrease dramatically while

ASCT2 increases. The high affinity and ubiquitous expression of ASCT2 suggest the increased

D-serine reabsorption in all proximal tubular segments. Together with the genetic inactivation

of DAAO during the pathology (Sasabe et al., 2014), the excess reabsorbed D-serine may lead

to a high plasma D-serine level (Figure 8: IRI).

Enantiomeric profiles of serine showed low plasma D/L-serine ratios in the normal

control but high D/L-serine ratios in IRI regardless of the weak stereoselectivity of ASCT2

(Figure 1B). This data suggested the different renal handling between D-serine and the L-serine.

L-Serine reabsorption has been reported to be mediated by B^0AT3 (Singer et al., 2009), and we

showed SMCTs-mediated D-serine reabsorption in this study. Thus, we propose that the low

plasma D/L-serine ratio in the normal kidney is due to the high L-serine reabsorption by B^0AT3,

together with the DAAO function that degrades D-serine in the cells. Our enantiomeric amino

acid profile detected low plasma L-serine and high urinary L-serine in IRI (Figure supplements

1B, 2B) and the proteomics revealed B^0AT3 decrease (4h IRI/sham = 0.56 fold; 8h IRI/sham

= 0.65 fold; Table S1). It is possible that low L-serine reabsorption in IRI is a result of B^0AT3

reduction.

In proteomics, some transporters including ASCT2 were found to be increased in the

AKI conditions (Figure 2B), yet the mechanism behind this change is still unclear. snRNA-seq

study revealed that the cell clusters in all proximal tubular segments during early AKI were

drastically different from the healthy kidney suggesting that the new distinct cell clusters in

AKI were *de novo* synthesized for the repair process, as early AKI is a reversible condition

(Kirita et al., 2020). ASCT2 was found to be only one type of small amino acid transporter

which was increased during AKI conditions (Figure 2B). ASCT2 was reported to be highly

expressed in proliferative cells and stem cells and played a key role in the regulation of

mammalian target of rapamycin (mTOR) signaling pathway and glutamine-mediated

metabolism (Scalise et al., 2018; Formisano and Van Winkle, 2016; Kandasamy et al., 2018).

Thus, we postulate that ASCT2 is a nascent transporter to provide small amino acids and

glutamine for cell growth, as a molecular mechanism in *de novo* cellular synthesis for the

recovery processes.

Besides ASCT2 and SMCTs, other transporters may contribute to D-serine handling in

the kidney. Influxes of D-[^{3}H]serine in the Na$^+$-free condition of the normal mouse BBMVs

accounted for ~ 30% of total uptake (Figure 7A-B). These are results of the experimental

background of non-specific accumulation of radiotracers and probably also Na$^+$-independent

transporter(s). In BBMVs from 8h IRI model, D-[^{3}H]serine uptake without preloading

suggested the existence of Na$^+$-dependent transporters during the injury. In the D-serine

transporter screening, we found that PAT1 and B^{0}AT3 attenuated D-serine-mediated growth

inhibition (Figure 4C-D). PAT1 is reported to transport D-serine in the transfected cell and

oocyte models while the recognition of D-serine by B^{0}AT3 is unknown (Boll et al., 2002; Chen

et al., 2003). PAT1, a low-affinity proton-coupled amino acid transporter, was found at both

sub-apical membranes at the S1 segment and inside of the epithelia (The Human Protein Atlas:

https://www.proteinatlas.org; updated on Dec 7[th], 2022) (Sagné et al., 2001; Vanslambrouck et

al., 2010). It is suggested that PAT1 is less active at the apical membrane where the luminal

pH is neutral (Miyauchi et al., 2005; Bröer, 2008b). However, a future experiment is required

to address the significance of PAT1 on D-serine transport in pathological conditions where the

pH homeostasis could be disrupted (Nakanishi et al., 2012; Bouchard and Mehta, 2022; Imenez

Silva and Mohebbi, 2022). Despite the D-serine transport systems at the apical side, other

transport systems at the basolateral side are also necessary to complete the picture of the D-

serine transport systems in renal proximal tubules.

Transport of a micromolecule is contributed by multiple transporters that orchestrate

together to maintain homeostasis and to control transport compensation in an aberrant

condition (Bröer et al., 2016; Wiriyasermkul et al., 2020; Gauthier-Coles et al., 2021). The

molecular characteristics and physiological functions of transporters are largely dependent on

their microenvironments, such as electrochemical gradients and the presence of other

transporters for coordinating functions. In our proteomic analysis, a high population of

membrane transport proteins was obtained in the dataset by applying urea treatment to the

membrane preparation process (Figure 2, Table supplement 1) (Kongpracha et al., 2022). By

integrating data focused on specific targets, such as transporters or amino acids, from the

protein molecule level to the metabolome or proteome, we were able to reveal the subtle

changes in the body and identify the causes of those small changes. This integrative and multi-

hierarchical approach, emphasizing the coordinate functions of multiple transporters, holds the

potential for investigating the dynamics of a broad range of micromolecules under diverse or

intricate conditions within any tissues or organs.

Materials and Methods

Materials, animals, and graphical analysis

General chemicals and cell culture media were purchased from Wako Fujifilm and Nacalai tesque. Chemicals used in mass spectrometry were HPLC or MS grades. Flp-In T-Rex 293 cells, Expi293F cells, Expi293 Expression medium, Lipofectamine 3000, fluorescent-labeled secondary antibodies, and Tyramide SuperBoost kit were from Thermo. Amino acids, fetal bovine serum (FBS), anti-FLAG antibody, and anti-FLAG M2 column were from Sigma. Secondary antibodies with HRP conjugated were from Jackson ImmunoResearch. L-[^{3}H]Serine, D-[^{3}H]serine, L-[^{3}H]alanine, and D-[^{3}H]alanine were from Moravek. L-[^{3}H]Glutamic acid, D-[^{3}H]glutamic acid, L-[^{14}C]tyrosine, D-[^{14}C]tyrosine, DL-[^{3}H]lactic acid, [^{3}H]propionic acid, [^{14}C]nicotinic acid, [^{14}C]uric acid, and L-[^{3}H]aspartic acid were from American Radiolabeled Chemicals. Anti-SGLT2 (SC-393350) and anti-Na$^+$/K$^+$-ATPase (SC-21712) antibodies were obtained from Santa Cruz Biotechnology.

All animal experiments were carried out following institutional guidelines under the approval of the Animal Experiment Committees of The Jikei University School of Medicine, Nara Medical University, and Keio University, Japan. Eight weeks old C57BL/6J mice were purchased from Japan SLC and CLEA Japan. Frozen kidneys of the 8 weeks old C57BL/6J mice were purchased from Sankyo Labo Service.

Human genes/proteins were defined with all letters in uppercase while those of the mouse molecules were defined with the first letter in uppercase. Unless otherwise indicated, data shown in all figures are mean ± SEM of the representative data from 3 reproducible experiments. Statistical differences and p-values were determined using the unpaired Student t test. Graphs, statistical significance, and kinetics were analyzed and plotted by GraphPad Prism 8.4.

Plasmid construction

In this study, we used the cDNA of human SLC5A8/SMCT1 clone "NM_145913". We generated the clone NM_145913 from the clone AK313788 (NBRC, NITE, Kisarazu, Japan). At first, SMCT1 from AK313788 was subcloned into p3XFLAG-CMV14 (Sigma) via *Hind*III and *Bam*HI sites. The clone "pCMV14-SMCT1" NM_145913 was subsequently generated by mutagenesis of p3XFLAG-CMV14-SMCT1_AK313788 at I193V, T201A and I490M (variants between NM_145913 and AK313788) by HiFi DNA Assembly Cloning (NEB) and site-directed mutagenesis. Human SLC5A12/SMCT2 cDNA (NM_178498; Sino Biological Inc.) was amplified by PCR and cloned into p3XFLAG-CMV14 via *Kpn*I and *Bam*HI sites to generate "pCMV14-SMCT2". Human expression clones of SLC7A10/Asc-1 (NM_019849), SLC7A1/CAT1 (NM_003045), SLC36A1/PAT1 (NM_078483), SLC2A5/GLUT5 (NM_003039), SLCO4C1/OATP-H (NM_180991), SLC22A13/OAT10 (NM_004256), SLC22A7/OAT2 (NM_153320), SLC6A18/B^{0}AT3 (NM_182632), SLC15A2/PEPT2 (NM_021082), and TMEM27/Collectrin (NM_020665) were obtained from GenScript and RIKEN BRC through the National BioResource Project of the MEXT/AMED, Japan. Mouse Asct2 (NM_009201) TrueORF clone was obtained from OriGene. p3XFLAG-CMV14 empty vector was used for Mock production. pcDNA5-SMCT1 and pcDNA5-SMCT2 used for generation of SMCT1- and SMCT2-stable cell lines, respectively, were constructed by assembling the PCR products of SMCT1 or SMCT2 into pcDNA5/FRT/TO (Thermo) by HiFi DNA Assembly Cloning.

Antibody (Ab) production

Anti-mouse Smct1 Ab was custom-produced. Peptide antigen corresponding to the amino acid residues 596 – 611 of Smct1 and the corresponding anti-sera from the immunized rabbit was produced by Cosmo Bio. Anti-Smct1 Ab was purified using HiTrap Protein G HP

(GE Healthcare) following the manufacturer's protocol. After purification, the Ab was dialyzed

against PBS pH 7.4 and adjusted concentration to 1 mg/mL.

Anti-human ASCT2 and anti-mouse Asct2 Abs were custom-produced. First, we

generated pET47b(+)-GST by subcloning the GST from pET49b(+) template into pET47b(+)

via *Not*I and *Xho*I restriction sites. Then, human ASCT2 antigen (amino acid residues 7 – 20)

was amplified and cloned into pET47b(+)-GST to obtain GST-fused ASCT2 antigen. For the

Asct2 antigen, both N-terminal (Asct2(NT), amino acid residues 1 – 38) and C-terminal

(Asct2(CT), amino acid residues 521 – 553) fragments were fused with GST by cloning into

pET47b(+)-GST and pET49b(+), respectively (Cosmo Bio). The GST-fusion antigens were

expressed in *E. coli* BL21(DE3) and purified by Glutathione Sepharose 4B (GE Healthcare) as

described previously (Nagamori et al., 2016a). The antigens were used to immunize rabbits to

obtain anti-sera (Cosmo Bio).

Anti-ASCT2 Ab was purified from the anti-sera using HiTrap Protein G HP (GE

Healthcare) following the manufacturer's protocol. After purification, the Ab was dialyzed

against PBS pH 7.4 and adjusted concentration to 1 mg/mL.

Anti-Asct2(NT) and anti-Asct2(CT) Abs were purified by using two-step purifications.

First, affinity columns of GST-fused Asct2(NT)-antigen and GST-fused Asct2(CT)-antigen

were produced by conjugating the antigens with HiTrap NHS-activated HP (GE Healthcare).

To purify anti-Asct2(NT) Ab, the anti-sera was subjected to the first purification by using the

GST-fused Asct2(NT)-antigen column. The elution fraction was dialyzed and subsequently

subjected to the second column of GST-fused Asct2(CT)-antigen column. The flowthrough

fraction corresponding to the affinity-purified anti-Asct2(NT) Ab was obtained. Purification of

anti-Asct2(CT) Ab was performed in the same way as Asct2(NT) Ab but used GST-fused

Asct2(CT)-antigen column followed by GST-fused Asct2(NT)-antigen column.

Ischemia-reperfusion injury (IRI) model

C57BL/6J (CLEA Japan) male mice between 12 – 16 weeks old underwent

experimental procedures for the IRI model. Ischemia-reperfusion was performed as previously

described (Sasabe et al., 2014). Before IRI induction, right kidney was removed. After twelve

days, the mice were grouped by randomization. Ischemia was operated for 45 min by clamping

the vessel under anesthesia. After that, the vessel clamp was removed, and the abdomen was

closed. Sham-operated control mice were treated identically except for clamping. Urine and

blood samples were collected at 4, 8, 20 and 40 hours after reperfusion. The mice were

anesthetized with isoflurane and euthanized by perfusion with PBS pH 7.4. The kidney was

removed and stored at -80 °C until use.

Measurement of amino acid enantiomers by 2D-HPLC

Plasma and urine samples collected from IRI mice were prepared as described with

modifications (Miyoshi et al., 2009; Hamase et al., 2010). Briefly, 20-fold volumes of methanol

were added to the samples and an aliquot was placed in a brown shading tube and used for

NBD derivatization (1 μL of the plasma was used for the reaction). After drying the solution

under reduced pressure, 20 μL of 200 mM sodium borate buffer (pH 8.0) and 5 μL of

fluorescent-labeling reagent [40 mM 4-fluoro-7-nitro-2,1,3-benzoxadiazole (NBD-F) in

anhydrous acetonitrile (MeCN)] were added, and then heated at 60 °C for 2 min. An aqueous

0.1% (v/v) TFA solution (375 μL) was added, and 20 μL of the reaction mixture was applied

to 2D-HPLC.

Enantiomers of amino acids were quantified using the 2D-HPLC platform (Miyoshi et

al., 2009; Hamase et al., 2010). The NBD-derivatives of the amino acids were separated using

a reversed-phase column (Singularity RP column, 1.0 mm i.d. × 50 mm; designed by Kyushu

University and KAGAMI INC.) with the gradient elution using aqueous mobile phases

containing MeCN and formic acid. To determine D- and L-amino acids, the fractions of amino

acids were automatically collected using a multi-loop valve and transferred to the

enantioselective column (Singularity CSP-001S, 1.5 mm i.d. × 75 mm; designed by Kyushu

University and KAGAMI INC.). Then, D- and L-amino acids were separated in the second

dimension by the enantioselective column. The mobile phases are the mixed solutions of

MeOH–MeCN containing formic acid, and the fluorescence detection of the NBD-amino acids

was carried out at 530 nm with excitation at 470 nm. Target peaks were quantified by scaling

the standard peak shapes (Hamase et al., 2018).

Isolation of brush border membrane vesicles (BBMVs) for mass spectrometry analysis

BBMVs were prepared by the calcium precipitation method (Biber et al., 2007;

Kongpracha et al., 2022). Frozen kidneys were minced into fine powders by using a polytron-

type homogenizer (Physcotron, Microtec) in the homogenization buffer containing 20 mM

Tris-HCl, pH 7.6, 250 mM sucrose, 1 mM EDTA and cOmplete EDTA-free protease inhibitor

cocktail (Roche). After low-speed centrifugation at 1,000 ×g and 3,000 ×g, the supernatant was

collected and incubated with 11 mM $CaCl_2$ for 20 min on ice with mild shaking. The

supernatant was ultra-centrifuged at 463,000 ×g for 15 min at 4 °C. The pellet was resuspended

in the homogenization buffer and repeated the steps of $CaCl_2$ precipitation. Finally, the pellet

of BBMVs was resuspended in 20 mM Tris-HCl pH 7.6, and 250 mM sucrose. Membrane

proteins of BBMVs were enriched by the Urea Wash method (Kongpracha et al., 2022). The

urea-washed BBMV samples were subjected to sample preparation for mass spectrometry.

Cell culture, transfection, and generation of stable cell lines

HEK293 and Flp-In T-Rex 293 cells were cultured in DMEM supplemented with 10 %

(v/v) FBS, 100 units/mL penicillin G, and 100 μg/mL streptomycin (P/S), and routinely

maintained at 37 °C, 5 % CO_2 and humidity. For transfection experiments, the cells were

seeded in antibiotic-free media for one day prior to transfection to obtain approximately 40%

confluence. DNA transient transfection and ASCT2-siRNA (ID#s12916; Ambion) transfection

were performed by using Lipofectamine 3000 following the manufacturer's protocol. The ratio

of DNA : P3000 : Lipofectamine 3000 is 1.0 μg : 2.0 μL : 1.5 μL. The ratio of siRNA :

Lipofectamine 3000 is 10 pmol : 1 μL. The cells were further maintained in the same media

for two days prior to the assays.

Flp-In T-REx 293 stably expressing SMCT1 (FlpInTR-SMCT1) and SMCT2

(FlpInTR-SMCT2) were generated by co-transfection of pOG44 and pcDNA5-SMCT1 or

pcDNA5-SMCT2 and subsequently cultured in the media containing 5 mg/L blasticidin and

150 mg/L hygromycin B for positive clone selection. Mock cells were generated in the same

way using the empty plasmid. Expressions of SMCT1 in FlpInTR-SMCT1 and SMCT2 in

FlpInTR-SMCT2 were induced by adding 1 mg/L doxycycline hyclate (Dox; Tet-ON system)

at one day after seeding. The cells were further cultured for two days prior to performing

experiments.

Wild-type and *ASCT2*-knockout HAP1 cells (Human near-haploid cell line; Horizon

Discovery) were cultured in IMDM supplemented with 10 % (v/v) FBS, P/S, and routinely

maintained at 37 °C, 5 % CO_2 and humidity.

Expi293F cells were cultured in Expi293 Expression medium at 37 °C, 8 % CO_2, and

humidity. To express SMCT1 transiently, the cells were transfected with pCMV14-SMCT1

using PEI MAX pH 6.9 (MW 40,000; Polysciences) and cultured for two days.

Mass spectrometry and proteome data analysis

624   Proteomics of mouse BBMVs was analyzed as described (Kongpracha et al., 2022;

625 Uetsuka et al., 2015). After preparing urea-washed BBMVs, the samples were fractionated into

626 four fractions by SDB-XC StageTips and desalted by C18-StageTips. Mass spectrometry was

627 performed using the Q Exactive (Thermo) coupled with nano-Advance UHPLC (Michrom

628 Bioresources). The UHPLC apparatus was equipped with a trap column (L-column ODS, 0.3

x 5 mm, CERI) and a C18 packed tip column (0.1 x 125 mm; Nikkyo Technos). Raw data from four fractions were analyzed using Proteome Discoverer 2.2 (Thermo) and Mascot 2.6.2 (Matrix Science). Data from four fractions were combined and searched for identified proteins from the UniProt mouse database (released in March 2019). The maximum number of missed cleavages, precursor mass tolerance, and fragment mass tolerance were set to 3, 10 ppm and 0.01 Da, respectively. The carbamidomethylation Cys was set as a fixed modification. Oxidation of Met and deamidation of Asn and Gln were set as variable modifications. A filter (false discovery rate < 1%) was applied to the resulting data. For each mouse sample, the analysis was conducted twice and the average was used. One data set was composed of 3 samples from each operation condition (n = 3). Statistical analyses were determined using Proteome Discoverer 2.2 (Thermo). Statistical differences and p-values were determined using the unpaired Student *t* test. Data represented the median ± SEM (Table S1).

Mass spectrometry of HEK293 cells was analyzed from crude membrane fractions. Membrane fractions from HEK293 cells were prepared from 3-day cultured cells as described (Nagamori et al., 2016b). Membrane proteins were enriched by the Urea Wash method, and tryptic peptides were subject for analysis as described above.

Proteome of BBMVs from mouse kidneys after ischemia operation for 4 (4h) or 8 hours (8h) was normalized to that of sham operation. The identified proteins were then subjected to annotate the biological functions by Ingenuity Pathway Analysis (IPA, Qiagen). Molecules from the dataset that met the cutoff of the Ingenuity Knowledge Base were considered for the analysis. A right-tailed Fisher's Exact Test was used to calculate the p-value determining the probability (z-scores) of the biological function. Protein localization in kidney segments was evaluated based on related literature and The Human Protein Atlas (http://www.proteinatlas.org: updated Dec 7th, 2022).

Effect of D-serine on cell growth

HEK293 cells were seeded into 96-well-plate at 10,000 cells/well. Transient

transfection was performed 12 hours after seeding followed by L- or D-serine treatment at 12

hours after that. In the case of FlpInTR-stable cell lines, if needed, ASCT2 siRNA was

transfected 12 hours after seeding. Dox was added one day after seeding followed by treatment

with L- or D-serine (in the presence or absence of ibuprofen as indicated) 10 hours after adding

Dox. The cells were further maintained for two days. Cell growth was examined by XTT assay.

In one reaction, 50 µL of 1 mg/mL XTT (2,3-Bis-(2-Methoxy-4-Nitro-5-Sulfophenyl)-2H-

Tetrazolium-5-Carboxanilide) (Biotium) was mixed with 5 µL of 1.5 mg/mL phenazine

methosulfate. The mixture was applied to the cells and incubated for 4 hours at 37 °C in the

cell culture incubator. Cell viability was evaluated by measuring the absorbance at 450 nm.

Cell growth in serine treatment samples was compared to the control (without serine treatment).

For transporter screening, the growth of the transfected cells at a specific D-serine concentration

was compared to that of Mock after normalization with no treatment.

Transport assay in cultured cells

Transport assay in cells was performed as described previously with some

modifications (Wiriyasermkul et al., 2012). Briefly, for D-[^{3}H]serine transport in HEK293 cells,

the cells were seeded into poly-D-lysine-coated 24-well plates at 1.2 x 10^5 cells/well and

cultured for three days. Unless indicated elsewhere, uptake of 20 µM (100 Ci/mol) or 100 µM

(10 Ci/mol) D-[^{3}H]serine was measured in PBS pH 7.4 at 37 °C at the indicated time points.

After termination of the assay, the cells were lysed. An aliquot was subjected to measure

protein concentration, and the remaining lysate was mixed with Optiphase HiSafe 3

(PerkinElmer). The radioactivity was monitored using a β-scintillation counter (LSC-8000,

Hitachi). In the transport assay with or without Na$^+$, Na$^+$-HBSS, or Na$^+$-free HBSS (choline-

Cl substitution) were used instead of PBS.

Transport assay in FlpInTR-stable cell lines was performed in a similar way to HEK293

cells. After cell seeding for one day, SMCT1 and SMCT2 expression were induced by adding

Dox for two days. For the ASCT2 knockdown experiment, ASCT2-siRNA was transfected 12

hours prior to Dox induction. The time course of 100 µM D-[^{3}H]serine transport (10 Ci/mol)

was measured 37 °C for 5 – 20 min. Inhibition assay was performed by adding the inhibitors

at the same time with D-[^{3}H]serine substrate. Kinetics of D-[^{3}H]serine transport were examined

by the uptake of D-[^{3}H]serine at the concentration of 0.5 – 8 mM (0.125 – 2 Ci/mol) for 10 min

at 37 °C.

Transport of D-[^{3}H]serine in wild-type and *ASCT2*-knockout HAP1 was performed in

a similar way to HEK293 and FlpInTR-stable cells but without the process of transfection.

SMCT1 purification, proteoliposome reconstitution, and transport assay

Human SMCT1 was purified from SMCT1-expressing Expi293F cells. Membrane

fraction and purification processes were performed as previously described with small

modifications (Nagamori et al., 2016a). First, cell pellets were resuspended in 20 mM Tris-HCl

pH 7.4, 150 mM NaCl, 10% (v/v) glycerol, and protease inhibitor cocktail (Roche). The crude

membrane fraction was derived from sonication and ultracentrifugation (sonication method).

Membrane proteins were extracted from the crude membrane fraction with 2% (w/v) DDM

and ultracentrifugation. SMCT1 was purified by anti-FLAG M2 affinity column. Unbound

proteins were washed out by 20 mM Tris-HCl pH 7.4, 200 mM NaCl, 10% (v/v) glycerol, and

0.05% (w/v) DDM. Then SMCT1 was eluted by 3xFLAG peptide in the washing buffer.

Purified SMCT1 was concentrated by Amicon Ultra Centrifugal Filters 30K (Millipore).

The reconstitution of proteoliposomes was performed as described with minor

modifications (Lee et al., 2019). The purified SMCT1 was reconstituted in liposomes (made

from 5:1 (w/w) of type II-S PC : brain total lipid) at a protein-to-lipid ratio of 1:100 (w/w) in

20 mM MOPS-Tris pH 7.0 and 100 mM KCl.

Transport assay in proteoliposomes was conducted by the rapid filtration method (Lee

et al., 2019). Uptake reaction was conducted by dilution of 1 μg SMCT1 proteoliposomes in

100 μL uptake buffer (20 mM MOPS-Tris pH 7.0, 100 mM NaCl for Na^+-buffer or KCl for

Na^+-free buffer, 1 mM $MgSO_4$ and 1 mM $CaCl_2$) containing radioisotope-labeled substrates.

The reaction was incubated at 25 °C for an indicated time. The radioisotope-labeled substrates

were used as follows: 5 Ci/mol for [^{14}C]uric acid, L-[^{14}C]tyrosine and D-[^{14}C]tyrosine; 10

Ci/mol for DL-[^{3}H]lactic acid, L-[^{3}H]alanine, D-[^{3}H]alanine, L-[^{3}H]serine, D-[^{3}H]serine, L-

[^{3}H]glutamic acid and D-[^{3}H]glutamic acid; and 20 Ci/mol for [^{3}H]propionic acid. In the

inhibition assay, 1 mM ibuprofen was applied at the same time with the radioisotope-labeled

substrates. Kinetics of nicotinate transport were measured for 3 min using 0.01 – 2 mM

[^{14}C]nicotinate (0.2 – 50 Ci/mol) in Na^+-containing buffer, and then subtracted from the uptake

in Na^+-free buffer.

715   Transport assay in mouse BBMVs

Both left and right kidneys from eight weeks old male normal control (Japan SLC;

Sankyo Labo Service) or IRI model (sham, 4h IRI, and 8h IRI) were taken out after PBS

perfusion and frozen until use. After mincing and homogenizing the frozen kidneys in the

buffer containing 20 mM Tris-HCl pH 7.6, 150 mM mannitol, 100 mM KCl, 1 mM EDTA,

and protease inhibitor cocktail, the BBMVs were prepared as described in the above section

but using magnesium instead of calcium for BBMV precipitation. The pellet of BBMVs was

resuspended in the suspension buffer (10 mM Tris-HCl, pH 7.6, 100 mM mannitol, and 100

mM KCl). In the L-glutamine preloading experiment, the BBMVs were incubated with 4 mM

L-glutamine (L-Gln) or only buffer (no preload) on ice for 3 hours. External buffers were then

removed by centrifugation at 21,000 ×g for 20 min and the BBMVs were resuspended in the

suspension buffer.

727  Transport assay was performed by rapid filtration. Prior to initiating the reaction, 5 µM

728 valinomycin was added to the BBMV samples. Transport assay was examined by diluting 100

729 µg BBMVs in 100 µL of the uptake buffer (10 mM Tris-HCl pH 7.6, 150 mM NaCl for Na^+

730 condition or KCl for Na^+-free condition, 50 mM mannitol and 5 µM valinomycin) containing

731 radioisotope substrates, 10 µM D-[^{3}H]serine (100 Ci/mol) or 10 µM L-[^{3}H]aspartic acid (100

732 Ci/mol), as described in the figures. The reaction was incubated at 30 °C at an indicated time,

733 then terminated by the addition of ice-cold buffer containing 10 mM Tris-HCl pH 7.6 and 200

734 mM mannitol and filtered through 0.45 µm nitrocellulose filter (Millipore), followed by

735 washing with the same buffer once. The membranes were soaked in Clear-sol I (Nacalai

736 Tesque), and the radioactivity on the membrane was monitored. For the inhibition experiments,

737 the tested inhibitors were added into the D-[^{3}H]serine substrate solution at the same time.

738 **Western Blot analysis**

739  Expressions of targeting proteins from membrane fractions were verified by Western

740 blot analysis as described (Nagamori et al., 2016b). Membrane fractions were dissolved in 1%

741 w/v DDM prior to the addition of the SDS-PAGE sample buffer. Signals of chemiluminescence

742 (Immobilon Forte Western HRP substrate; Millipore) were visualized by ChemiDoc MP

743 Imaging system (Bio-Rad).

744 **Immunofluorescence staining of mouse kidneys**

745  The 8 weeks old male C57BL/6J mice (Japan SLC) were anesthetized and fixed by

746 anterograde perfusion via the aorta with 4% w/v paraformaldehyde in 0.1 M sodium phosphate

747 buffer pH 7.4. The kidneys were dissected, post-fixed in the same buffer for two days, and

748 cryoprotected in 10 %, 20 %, and 30 % w/v sucrose. Frozen kidney sections were cut at 7 µm

749 thickness in a cryostat (Leica) and mounted on MAS-coated glass slides (Matsunami). The

750 sections were placed in antigen retrieval buffer (10 mM citrate and 10 mM sodium Citrate),

autoclaved at 121 °C for 5 min and washed by TBS-T (Tris-buffered saline (TBS) with 0.1 %

v/v Tween 20). Immunostaining was done by serial incubation with each antibody as below.

For Asct2 and Sglt2 co-immunostaining, the samples were incubated with 3 %

hydrogen peroxide solution for 10 min, washed with TBS, and incubated in Blocking One

Histo (Nacalai tesque) for 15 min. The samples were then incubated with mouse anti-SGLT2

antibody diluted in immunoreaction enhancer B solution (Can Get Signal immunostain,

TOYOBO) overnight at 4 °C. Signal was enhanced by Alexa Fluor 568 Tyramide SuperBoost

(TSA) kit, goat anti-mouse IgG, following the manufacturer's instruction (Thermo). The

antibodies were then stripped by citrate/acetate-based buffer, pH 6.0, containing 0.3% w/v SDS

at 95 °C for 10 min (Buchwalow et al., 2018), washed by TBS, and incubated with Blocking

One Histo. Asct2 staining was done as described (Nagamori et al., 2016a). Briefly, the samples

were incubated with rabbit anti-Asct2(NT) antibody diluted in immunoreaction enhancer A

solution (Can Get Signal immunostain) overnight at 4 °C. After washing with TBS-T, the

specimens were incubated with Alexa Fluor 488-labeled donkey anti-rabbit IgG.

For Asct2 and Agt1 co-immunostaining, signals of both antibodies were enhanced by

TSA kit. First, the specimens were incubated with rabbit anti-Agt1(G) antibody (Nagamori et

al., 2016a) overnight at 4 °C followed by Alexa Fluor 568 TSA kit with goat anti-rabbit. The

antibodies were then stripped. The specimens were incubated with rabbit anti-Asct2(NT)

overnight at 4 °C and then repeated the steps of TSA kit using Alexa Fluor 488, goat anti-rabbit.

Staining of Asct2 and Na^+/K^+-ATPase was performed without TSA enhancement. After

blocking by Blocking One Histo, the samples were incubated with rabbit anti-Asct2(NT)

antibody diluted in immunoreaction enhancer A solution overnight. The samples were washed

with TBS-T, incubated with Alexa Fluor488-labeled donkey anti-rabbit IgG, and washed again.

Non-specific staining was blocked by Blocking One Histo and the specimens were then

incubated with mouse anti-Na^+/K^+-ATPase antibody diluted in immunoreaction enhancer B

solution overnight at 4 °C. The specimens were washed with TBS-T and incubated with Alexa

Fluor568-labeled goat anti-mouse IgG for 1 hour.

Smct1 and Sglt2 were co-stained. After blocking by Blocking One Histo, the samples

were incubated with rabbit anti-Smct1 antibody and mouse anti-SGLT2 diluted in

immunoreaction enhancer B solution overnight at 4 °C. The specimens were washed with TBS-

T and incubated with Alexa Fluor488-labeled donkey anti-rabbit IgG and Alexa Fluor568-

labeled goat anti-mouse IgG for 1 hour.

Prior to the co-staining of Smct1 and Agt1, anti-Agt1 antibody was conjugated with

Alexa Fluor 568 dye via succinimidyl ester reaction. The antibody was adjusted to pH 8.3 and

mixed with Alexa Fluor 568 NHS Ester (Thermo) for 2 hours. The reaction was stopped by

incubation with 1 M ethanolamine for 1 hour. The unconjugated dye was removed by size

exclusion using Bio-Spin P-30 gel columns (Bio-rad) equilibrated in PBS pH 7.4. For Smct1

and Agt1 co-staining, after the blocking step, the samples were firstly stained with rabbit anti-

Smct1 antibody diluted in immunoreaction enhancer A solution. Signal was enhanced by Alexa

Fluor 488 TSA kit with goat anti-rabbit IgG. Subsequently, the specimen was washed with

TBS-T and non-specific staining was blocked by Blocking One Histo. The samples were then

incubated with Alexa Fluor 568-conjugated anti-Smct1 antibody diluted in Blocking One Histo

overnight at 4 °C.

All specimens were washed with TBS-T and mounted with Fluoromount (Diagnostic

Biosystems). Imaging was detected using a KEYENCE BZ-X710 microscope. Images were

color-adjusted using ImageJ ver. 1.51 (NIH). Images in Figure 3B were subjected to

deconvolution process in Fiji/ImageJ2 ver. 2.9 prior to color adjustment.

Analysis of Smcts and Asct2 from the open-sourced dataset of single nucleus RNA

sequencing (snRNA-seq) of the IRI model

The dataset of snRNA-seq used is created from the mice after ischemia-reperfusion injury (IRI) and published by Kirita et al. (Kirita et al., 2020). The snRNA-seq dataset (GSE139107), which has already processed by zUMIs and SoupX, was downloaded from GEO, a public functional genomics data repository (Barrett et al., 2012). The dataset revealed percentage of mitochondrial UMI counts of each cell (calculated by *PercentageFeatureSet* function of Seurat) to be < 0.01 % in more than 94% cells in each cluster. From the "GSE139107_MouseIRI.metadata", we annotated the expression of Smct1, Smct2 and Asct2 in proximal tubule segments 1 – 3 (S1, S2, and S3) at the early stages of IRI (4 and 12 hours) compared to the control mice. Downstream analyses including normalization, scaling and visualization were performed by Seurat v3.2, an R package designed for Quality Control, analysis and exploration of single-cell RNA-seq data (Stuart et al., 2019). After selecting the cells in S1, S2, and S3 from the datasets (control, 4-hour IRI, and 12-hour IRI), we merged the data of the cells in the same cluster. The data were then calculated as log-normalized values by *NormalizeData* function and scaled by *ScaleData* function of Seurat. Visualization of the expression data was performed using Seurat plug-in of R software and ggplot2 plug-in of R software (Wickham, 2010). Dot plots of gene expression in each cluster after log-transformation and scaling were created by *Dotplot* function of Seurat.

Data and materials availability

All proteomics data have been deposited in Japan Proteome Standard Repository/Database: JPST000929 and JPST000931. Codes for snRNA sequencing reanalysis are available in GitHub repository at https://github.com/SN-PW/snRNA_seq.git.

Acknowledgments

We greatly appreciate Noriyoshi Isozumi for preliminary proteomic analysis, Yuki Mori, Junko Iwatani, and Yuika Shimo for experimental assistance, Hiroshi Imoto, Eiichi Negishi, Maiko Nakane, and Shoto Ishigo for optimization of enantiomeric amino acid analysis, Genro Kashino and Haruka Minowa for management of radioisotope facilities, Rikako Furuya for crucial discussion, Xuan Trang T Nguyen for her native English proofreading, and Sae Ochi and Tomokazu Matsuura for their tremendous supports. We are especially grateful to Yoshinori Moriyama for his critical reading and suggestion. This work is partly supported by JSPS KAKENHI (JP22K06150) to P.W. and (JP21H03365) to S.N.; research grants from Gout and uric acid Foundation, Nakatani Foundation, Shiseido Company, Ltd., AMED (JP21ek0310012), and AMED-CREST (JP21gm0810010) to S.N.; and unrestricted fund provided from Dr. Jin and Mrs. Keiko Hanyu (Jiseikai Kajigaya Clinic.) to S.N. and P.W.

Competing Interest Statement

A patent (WO/2021/132691) has been applied by KAGAMI Inc., Nara Medical University, and NIBIOHN with P.W., S.M., P.K., T.K., M.Mit., and S.N. as inventors based on this research. M.Mit. is a founder and CEO of KAGAMI Inc., a startup company working on chiral amino acids analysis and research for medical applications. The authors declare that nothing potential conflicts of interest.

Author contributions

Pattama Wiriyasermkul—Methodology, Investigation, Validation, Formal analysis, Data curation, Visualization, Funding acquisition, Writing—original draft, Writing—review and editing

Satomi Moriyama—Investigation, Formal analysis

Masataka Suzuki—Investigation, Formal analysis

Pornparn Kongpracha—Investigation, Formal analysis, Data curation

Nodoka Nakamae—Investigation

Saki Takeshita—Formal analysis, Data curation

Yoko Tanaka—Investigation

Akina Matsuda—Investigation

Masaki Miyasaka—Formal analysis, Validation

Kenji Hamase—Methodology, Investigation, Formal analysis

Tomonori Kimura—Methodology, Resources

Masashi Mita—Conceptualization, Funding acquisition

Jumpei Sasabe—Methodology, Investigation, Validation, Resources

Shushi Nagamori—Conceptualization, Methodology, Validation, Data curation, Supervision,

Resources, Funding acquisition, Project administration, Writing—original draft, Writing—

review and editing

All authors contributed to the final manuscript.

References

Avissar NE, Ryan CK, Ganapathy V, Sax HC. 2001. Na$^+$-dependent neutral amino acid transporter ATB^0 is a rabbit epithelial cell brush-border protein. *American Journal of Physiology-Cell Physiology* **281**:C963–C971. doi:10.1152/ajpcell.2001.281.3.C963

Bailey CG, Ryan RM, Thoeng AD, Ng C, King K, Vanslambrouck JM, Auray-Blais C, Vandenberg RJ, Bröer S, Rasko JEJ. 2011. Loss-of-function mutations in the glutamate transporter SLC1A1 cause human dicarboxylic aminoaciduria. *Journal of Clinical Investigation* **121**:446–453. doi:10.1172/JCI44474

Barrett T, Wilhite SE, Ledoux P, Evangelista C, Kim IF, Tomashevsky M, Marshall KA, Phillippy KH, Sherman PM, Holko M, Yefanov A, Lee H, Zhang N, Robertson CL, Serova N, Davis S, Soboleva A. 2012. NCBI GEO: archive for functional genomics data sets—update. *Nucleic Acids Research* **41**:D991–D995. doi:10.1093/nar/gks1193

Basu AC, Tsai GE, Ma C-L, Ehmsen JT, Mustafa AK, Han L, Jiang ZI, Benneyworth MA, Froimowitz MP, Lange N, Snyder SH, Bergeron R, Coyle JT. 2009. Targeted disruption of serine racemase affects glutamatergic neurotransmission and behavior. *Molecular Psychiatry* **14**:719–727. doi:10.1038/mp.2008.130

Biber J, Stieger B, Stange G, Murer H. 2007. Isolation of renal proximal tubular brush-border membranes. *Nature Protocols* **2**:1356–1359. doi:10.1038/nprot.2007.156

Bodner O, Radzishevsky I, Foltyn VN, Touitou A, Valenta AC, Rangel IF, Panizzutti R, Kennedy RT, Billard JM, Wolosker H. 2020. D-Serine Signaling and NMDAR-Mediated Synaptic Plasticity Are Regulated by System A-Type of Glutamine/D-Serine Dual Transporters. *Journal of Neuroscience* **40**:6489–6502. doi:10.1523/JNEUROSCI.0801-20.2020

Boll M, Foltz M, Rubio-Aliaga I, Kottra G, Daniel H. 2002. Functional Characterization of
Two Novel Mammalian Electrogenic Proton-dependent Amino Acid Cotransporters.
Journal of Biological Chemistry **277**:22966–22973. doi:10.1074/jbc.M200374200

Bouchard J, Mehta RL. 2022. Timing of Kidney Support Therapy in Acute Kidney Injury:
What Are We Waiting For? *American Journal of Kidney Diseases* **79**:417–426.
doi:10.1053/j.ajkd.2021.07.014

Bröer A, Rahimi F, Bröer S. 2016. Deletion of Amino Acid Transporter ASCT2 (SLC1A5)
Reveals an Essential Role for Transporters SNAT1 (SLC38A1) and SNAT2
(SLC38A2) to Sustain Glutaminolysis in Cancer Cells. *Journal of Biological Chemistry*
291:13194–13205. doi:10.1074/jbc.M115.700534

Bröer S. 2008a. Amino Acid Transport Across Mammalian Intestinal and Renal Epithelia.
Physiological Reviews **88**:249–286. doi:10.1152/physrev.00018.2006

Bröer S. 2008b. Apical Transporters for Neutral Amino Acids: Physiology and
Pathophysiology. *Physiology* **23**:95–103. doi:10.1152/physiol.00045.2007

Buchwalow I, Samoilova V, Boecker W, Tiemann M. 2018. Multiple immunolabeling with
antibodies from the same host species in combination with tyramide signal
amplification. *Acta Histochemica* **120**:405–411. doi:10.1016/j.acthis.2018.05.002

César-Razquin A, Snijder B, Frappier-Brinton T, Isserlin R, Gyimesi G, Bai X, Reithmeier RA,
Hepworth D, Hediger MA, Edwards AM, Superti-Furga G. 2015. A Call for Systematic
Research on Solute Carriers. *Cell* **162**:478–487. doi:10.1016/j.cell.2015.07.022

Chen Z, Fei Y, Anderson CMH, Wake KA, Miyauchi S, Huang W, Thwaites DT, Ganapathy
V. 2003. Structure, function and immunolocalization of a proton-coupled amino acid
transporter (hPAT1) in the human intestinal cell line Caco-2. *The Journal of Physiology*
546:349–361. doi:10.1113/jphysiol.2002.026500

Formisano TM, Van Winkle LJ. 2016. At Least Three Transporters Likely Mediate Threonine

Uptake Needed for Mouse Embryonic Stem Cell Proliferation. *Frontiers in Cell and*

Developmental Biology **4**. doi:10.3389/fcell.2016.00017

Foster AC, Farnsworth J, Lind GE, Li Y-X, Yang J-Y, Dang V, Penjwini M, Viswanath V,

Staubli U, Kavanaugh MP. 2016. D-Serine Is a Substrate for Neutral Amino Acid

Transporters ASCT1/SLC1A4 and ASCT2/SLC1A5, and Is Transported by Both

Subtypes in Rat Hippocampal Astrocyte Cultures. *PLoS ONE* **11**:e0156551.

doi:10.1371/journal.pone.0156551

Fu Y, Tang C, Cai J, Chen G, Zhang D, Dong Z. 2018. Rodent models of AKI-CKD transition.

American Journal of Physiology-Renal Physiology **315**:F1098–F1106.

doi:10.1152/ajprenal.00199.2018

Ganapathy V, Thangaraju M, Gopal E, Martin PM, Itagaki S, Miyauchi S, Prasad PD. 2008.

Sodium-coupled Monocarboxylate Transporters in Normal Tissues and in Cancer. *The*

AAPS Journal **10**:193. doi:10.1208/s12248-008-9022-y

Gauthier-Coles G, Vennitti J, Zhang Z, Comb WC, Xing S, Javed K, Bröer A, Bröer S. 2021.

Quantitative modelling of amino acid transport and homeostasis in mammalian cells.

Nature Communications **12**:5282. doi:10.1038/s41467-021-25563-x

Ghezzi C, Loo DDF, Wright EM. 2018. Physiology of renal glucose handling via SGLT1,

SGLT2 and GLUT2. *Diabetologia* **61**:2087–2097. doi:10.1007/s00125-018-4656-5

Gonda Y, Matsuda A, Adachi K, Ishii C, Suzuki M, Osaki A, Mita M, Nishizaki N, Ohtomo

Y, Shimizu T, Yasui M, Hamase K, Sasabe J. 2023. Mammals sustain amino acid

homochirality against chiral conversion by symbiotic microbes. *Proceedings of the*

National Academy of Sciences USA **120**:e2300817120. doi:10.1073/pnas.2300817120

Gopal E, Umapathy NS, Martin PM, Ananth S, Gnana-Prakasam JP, Becker H, Wagner CA,

Ganapathy V, Prasad PD. 2007. Cloning and functional characterization of human

SMCT2 (SLC5A12) and expression pattern of the transporter in kidney. *Biochimica et*

Biophysica   Acta   (BBA)   -   Biomembranes   **1768**:2690–2697.

doi:10.1016/j.bbamem.2007.06.031

Hamase K, Ikeda T, Ishii C, Ishigo S, Masuyama K, Akita T, Furusho A, Takahashi M, Ide T,

Mita M. 2018. Determination of Trace Amounts of Chiral Amino Acids in Complicated

Biological   Samples   Using   Two-Dimensional   High-Performance   Liquid

Chromatography   with   an   Innovative   "Shape-Fitting"   Peak

Identification/Quantification   Method.   *CHROMATOGRAPHY*   **39**:147–152.

doi:10.15583/jpchrom.2018.019

Hamase K, Miyoshi Y, Ueno K, Han H, Hirano J, Morikawa A, Mita M, Kaneko T, Lindner

W, Zaitsu K. 2010. Simultaneous determination of hydrophilic amino acid enantiomers

in mammalian tissues and physiological fluids applying a fully automated micro-two-

dimensional   high-performance   liquid   chromatographic   concept.   *Journal   of*

Chromatography A **1217**:1056–1062. doi:10.1016/j.chroma.2009.09.002

Hasegawa H, Masuda N, Natori H, Shinohara Y, Ichida K. 2019. Pharmacokinetics and

toxicokinetics of D-serine in rats. *Journal of Pharmaceutical and Biomedical Analysis*

162:264–271. doi:10.1016/j.jpba.2018.09.026

Hashimoto A, Nishikawa T, Hayashi T, Fujii N, Harada K, Oka T, Takahashi K. 1992. The

presence of free D-serine in rat brain. *FEBS Lett* **296**:33–36. doi:10.1016/0014-

5793(92)80397-y

Hesaka A, Sakai S, Hamase K, Ikeda T, Matsui R, Mita M, Horio M, Isaka Y, Kimura T. 2019.

D-Serine   reflects   kidney   function   and   diseases.   *Scientific   Reports*   **9**:5104.

doi:10.1038/s41598-019-41608-0

Imenez Silva PH, Mohebbi N. 2022. Kidney metabolism and acid–base control: back to the

basics. *Pflugers Arch - European Journal of Physiology* **474**:919–934.

doi:10.1007/s00424-022-02696-6

Itagaki S, Gopal E, Zhuang L, Fei Y-J, Miyauchi S, Prasad PD, Ganapathy V. 2006. Interaction

of Ibuprofen and Other Structurally Related NSAIDs with the Sodium-Coupled

Monocarboxylate Transporter SMCT1 (SLC5A8). *Pharmaceutical Research* **23**:1209–

1216. doi:10.1007/s11095-006-0023-1

Iwata Y, Nakade Y, Kitajima S, Yoneda-Nakagawa S, Oshima M, Sakai N, Ogura H, Sato K,

Toyama T, Yamamura Y, Miyagawa T, Yamazaki H, Hara A, Shimizu M, Furuichi K,

Mita M, Hamase K, Tanaka T, Nishida M, Muramatsu W, Yamamoto H, Shichino S,

Ueha S, Matsushima K, Wada T. 2022. Protective effect of D-alanine against acute

kidney injury. *American Journal of Physiology-Renal Physiology* **322**:F667–F679.

doi:10.1152/ajprenal.00198.2021

Kanai Y, Hediger MA. 1992. Primary structure and functional characterization of a high-

affinity glutamate transporter. *Nature* **360**:467–471. doi:10.1038/360467a0

Kandasamy P, Gyimesi G, Kanai Y, Hediger MA. 2018. Amino acid transporters revisited:

New views in health and disease. *Trends in Biochemical Sciences* **43**:752–789.

doi:10.1016/j.tibs.2018.05.003

Kaplan E, Zubedat S, Radzishevsky I, Valenta AC, Rechnitz O, Sason H, Sajrawi C, Bodner

O, Konno K, Esaki K, Derdikman D, Yoshikawa T, Watanabe M, Kennedy RT, Billard

J-M, Avital A, Wolosker H. 2018. ASCT1 (Slc1a4) transporter is a physiologic

regulator of brain D-serine and neurodevelopment. *Proceedings of the National*

Academy of Sciences USA **115**:9628–9633. doi:10.1073/pnas.1722677115

Kimura T, Hamase K, Miyoshi Y, Yamamoto R, Yasuda K, Mita M, Rakugi H, Hayashi T,

Isaka Y. 2016. Chiral amino acid metabolomics for novel biomarker screening in the

981 prognosis of chronic kidney disease. *Scientific Reports* **6**:26137.

982 doi:10.1038/srep26137

983 Kimura T, Hesaka A, Isaka Y. 2020. Utility of D-serine monitoring in kidney disease.

984 *Biochimica et Biophysica Acta (BBA) - Proteins and Proteomics* **1868**:140449.

985 doi:10.1016/j.bbapap.2020.140449

986 Kirita Y, Wu H, Uchimura K, Wilson PC, Humphreys BD. 2020. Cell profiling of mouse acute

987 kidney injury reveals conserved cellular responses to injury. *Proceedings of the*

988 *National Academy of Sciences USA* **117**:15874–15883. doi:10.1073/pnas.2005477117

989 Kongpracha P, Wiriyasermkul P, Isozumi N, Moriyama S, Kanai Y, Nagamori S. 2022. Simple

990 But Efficacious Enrichment of Integral Membrane Proteins and Their Interactions for

991 In-Depth Membrane Proteomics. *Molecular & Cellular Proteomics* **21**:100206.

992 doi:10.1016/j.mcpro.2022.100206

993 Kragh-Hansen U, Sheikh MI. 1984. Serine uptake by luminal and basolateral membrane

994 vesicles from rabbit kidney. *The Journal of Physiology* **354**:55–67.

995 doi:10.1113/jphysiol.1984.sp015361

996 Lee Y, Wiriyasermkul P, Jin C, Quan L, Ohgaki R, Okuda S, Kusakizako T, Nishizawa T, Oda

997 K, Ishitani R, Yokoyama T, Nakane T, Shirouzu M, Endou H, Nagamori S, Kanai Y,

998 Nureki O. 2019. Cryo-EM structure of the human L-type amino acid transporter 1 in

999 complex with glycoprotein CD98hc. *Nature Structural & Molecular Biology* **26**:510–

1000 517. doi:10.1038/s41594-019-0237-7

1001 Liu J, Tan Y, Cheng H, Zhang D, Feng W, Peng C. 2022. Functions of Gut Microbiota

1002 Metabolites, Current Status and Future Perspectives. *Aging and disease* **13**:1106.

1003 doi:10.14336/AD.2022.0104

1004 Marx D, Metzger J, Pejchinovski M, Gil RB, Frantzi M, Latosinska A, Belczacka I, Heinzmann

1005 SS, Husi H, Zoidakis J, Klingele M, Herget-Rosenthal S. 2018. Proteomics and

Metabolomics for AKI Diagnosis. *Seminars in Nephrology* **38**:63–87. doi:10.1016/j.semnephrol.2017.09.007

Matsuo H, Chiba T, Nagamori S, Nakayama A, Domoto H, Phetdee K, Wiriyasermkul P, Kikuchi Y, Oda T, Nishiyama J, Nakamura T, Morimoto Y, Kamakura K, Sakurai Y, Nonoyama S, Kanai Y, Shinomiya N. 2008. Mutations in Glucose Transporter 9 Gene SLC2A9 Cause Renal Hypouricemia. *The American Journal of Human Genetics* **83**:744–751. doi:10.1016/j.ajhg.2008.11.001

Metzner L, Kottra G, Neubert K, Daniel H, Brandsch M. 2005. Serotonin, L-tryptophan, and tryptamine are effective inhibitors of the amino acid transport system PAT1. *FASEB j* **19**:1468–1473. doi:10.1096/fj.05-3683com

Miyauchi S, Abbot EL, Zhuang L, Subramanian R, Ganapathy V, Thwaites DT. 2005. Isolation and function of the amino acid transporter PAT1 (slc36a1) from rabbit and discrimination between transport via PAT1 and system IMINO in renal brush-border membrane vesicles. *Molecular Membrane Biology* **22**:549–559. doi:10.1080/09687860500421779

Miyoshi Y, Hamase K, Tojo Y, Mita M, Konno R, Zaitsu K. 2009. Determination of D-serine and D-alanine in the tissues and physiological fluids of mice with various D-amino-acid oxidase activities using two-dimensional high-performance liquid chromatography with fluorescence detection. *Journal of Chromatography B* **877**:2506–2512. doi:10.1016/j.jchromb.2009.06.028

Morehead RP, Fishman WH, Artom C. 1945. Renal Injury in the Rat Following the Administration of Serine by Stomach Tube. *The American Journal of Pathology* **21**:803–815.

Mothet J-P, Parent AT, Wolosker H, Brady RO, Linden DJ, Ferris CD, Rogawski MA, Snyder

SH. 2000. D-Serine is an endogenous ligand for the glycine site of the *N*-methyl-D-

aspartate receptor. *Proc Natl Acad Sci USA* **97**:4926–4931. doi:10.1073/pnas.97.9.4926

Nagamori S, Wiriyasermkul P, Guarch ME, Okuyama H, Nakagomi S, Tadagaki K, Nishinaka

Y, Bodoy S, Takafuji K, Okuda S, Kurokawa J, Ohgaki R, Nunes V, Palacín M, Kanai

Y. 2016a. Novel cystine transporter in renal proximal tubule identified as a missing

partner of cystinuria-related plasma membrane protein rBAT/SLC3A1. *Proceedings of*

the National Academy of Sciences USA **113**:775–780. doi:10.1073/pnas.1519959113

Nagamori S, Wiriyasermkul P, Okuda S, Kojima N, Hari Y, Kiyonaka S, Mori Y, Tominaga

H, Ohgaki R, Kanai Y. 2016b. Structure–activity relations of leucine derivatives reveal

critical moieties for cellular uptake and activation of mTORC1-mediated signaling.

Amino Acids **48**:1045–1058. doi:10.1007/s00726-015-2158-z

Nakade Y, Iwata Y, Furuichi K, Mita M, Hamase K, Konno R, Miyake T, Sakai N, Kitajima

S, Toyama T, Shinozaki Y, Sagara A, Miyagawa T, Hara A, Shimizu M, Kamikawa Y,

Sato K, Oshima M, Yoneda-Nakagawa S, Yamamura Y, Kaneko S, Miyamoto T,

Katane M, Homma H, Morita H, Suda W, Hattori M, Wada T. 2018. Gut microbiota–

derived D-serine protects against acute kidney injury. *JCI Insight* **3**:e97957.

doi:10.1172/jci.insight.97957

Nakanishi N, Fukui M, Tanaka M, Toda H, Imai S, Yamazaki M, Hasegawa G, Oda Y,

Nakamura N. 2012. Low Urine pH Is a Predictor of Chronic Kidney Disease. *Kidney*

and Blood Pressure Research **35**:77–81. doi:10.1159/000330487

Okada A, Nangaku M, Jao T-M, Maekawa H, Ishimono Y, Kawakami T, Inagi R. 2017. D-

serine, a novel uremic toxin, induces senescence in human renal tubular cells via GCN2

activation. *Scientific Reports* **7**:11168. doi:10.1038/s41598-017-11049-8

Okushima H, Iwata Y, Hesaka A, Sugimori E, Ikeda T, Nakane M, Mita M, Hayashi T, Isaka Y, Kimura T. 2021. Intra-body dynamics of D-serine reflects the origin of kidney diseases. *Clinical and Experimental Nephrology* **25**:893–901. doi:10.1007/s10157-021-02052-5

Papouin T, Ladépêche L, Ruel J, Sacchi S, Labasque M, Hanini M, Groc L, Pollegioni L, Mothet J-P, Oliet SHR. 2012. Synaptic and Extrasynaptic NMDA Receptors Are Gated by Different Endogenous Coagonists. *Cell* **150**:633–646. doi:10.1016/j.cell.2012.06.029

Paroder V, Spencer SR, Paroder M, Arango D, Schwartz S, Mariadason JM, Augenlicht LH, Eskandari S, Carrasco N. 2006. Na$^+$/monocarboxylate transport (SMCT) protein expression correlates with survival in colon cancer: Molecular characterization of SMCT. *Proceedings of the National Academy of Sciences USA* **103**:7270–7275. doi:10.1073/pnas.0602365103

Perland E, Fredriksson R. 2017. Classification Systems of Secondary Active Transporters. *Trends in Pharmacological Sciences* **38**:305–315. doi:10.1016/j.tips.2016.11.008

Ranjan R, Khazen G, Gambazzi L, Ramaswamy S, Hill SL, Schürmann F, Markram H. 2011. Channelpedia: An Integrative and Interactive Database for Ion Channels. *Frontiers in Neuroinformatics* **5**. doi:10.3389/fninf.2011.00036

Rosenberg D, Artoul S, Segal AC, Kolodney G, Radzishevsky I, Dikopoltsev E, Foltyn VN, Inoue R, Mori H, Billard J-M, Wolosker H. 2013. Neuronal D-Serine and Glycine Release Via the Asc-1 Transporter Regulates NMDA Receptor-Dependent Synaptic Activity. *Journal of Neuroscience* **33**:3533–3544. doi:10.1523/JNEUROSCI.3836-12.2013

Sagné C, Agulhon C, Ravassard P, Darmon M, Hamon M, El Mestikawy S, Gasnier B, Giros B. 2001. Identification and characterization of a lysosomal transporter for small neutral

amino acids. *Proceedings of the National Academy of Sciences USA* **98**:7206–7211.

doi:10.1073/pnas.121183498

Sasabe J, Suzuki M. 2018. Distinctive Roles of D-Amino Acids in the Homochiral World:

Chirality of Amino Acids Modulates Mammalian Physiology and Pathology. *The Keio*

Journal of Medicine **68**:1–16. doi:10.2302/kjm.2018-0001-IR

Sasabe J, Suzuki M, Miyoshi Y, Tojo Y, Okamura C, Ito S, Konno R, Mita M, Hamase K, Aiso

S. 2014. Ischemic Acute Kidney Injury Perturbs Homeostasis of Serine Enantiomers in

the Body Fluid in Mice: Early Detection of Renal Dysfunction Using the Ratio of Serine

Enantiomers. *PLoS ONE* **9**:e86504. doi:10.1371/journal.pone.0086504

Scalise M, Pochini L, Console L, Losso MA, Indiveri C. 2018. The Human SLC1A5 (ASCT2)

Amino Acid Transporter: From Function to Structure and Role in Cell Biology.

Frontiers in Cell and Developmental Biology **6**:96. doi:10.3389/fcell.2018.00096

Schweikhard ES, Ziegler CM. 2012. Amino Acid Secondary Transporters: toward a common

transport mechanism. *Current Topics in Membranes* Elsevier. **70**:1–28.

doi:10.1016/B978-0-12-394316-3.00001-6

Shenkin A. 2006. Micronutrients in health and disease. *Postgraduate Medical Journal* **82**:559–

567. doi:10.1136/pgmj.2006.047670

Shimomura A, Carone FA, Peterson DR. 1988. Contraluminal uptake of serine in the proximal

nephron. *Biochimica et Biophysica Acta (BBA) - Biomembranes* **939**:52–56.

doi:10.1016/0005-2736(88)90046-6

Silbernagl S, Völker K, Dantzler WH. 1999. D-Serine is reabsorbed in rat renal pars recta.

American Journal of Physiology-Renal Physiology **276**:F857–F863.

doi:10.1152/ajprenal.1999.276.6.F857

Singer D, Camargo SMR, Huggel K, Romeo E, Danilczyk U, Kuba K, Chesnov S, Caron MG,

Penninger JM, Verrey F. 2009. Orphan Transporter SLC6A18 Is Renal Neutral Amino

Acid Transporter B^{0}AT3. *Journal of Biological Chemistry* **284**:19953–19960.

doi:10.1074/jbc.M109.011171

Srinivas SR, Gopal E, Zhuang L, Itagaki S, Martin PM, Fei Y-J, Ganapathy V, Prasad PD.

2005. Cloning and functional identification of slc5a12 as a sodium-coupled low-affinity

transporter for monocarboxylates (SMCT2). *Biochemical Journal* **392**:655–664.

doi:10.1042/BJ20050927

Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, Hao Y, Stoeckius M,

Smibert P, Satija R. 2019. Comprehensive Integration of Single-Cell Data. *Cell*

177:1888-1902.e21. doi:10.1016/j.cell.2019.05.031

Suzuki M, Shimizu-Hirota R, Mita M, Hamase K, Sasabe J. 2022. Chiral resolution of plasma

amino acids reveals enantiomer-selective associations with organ functions. *Amino*

Acids **54**:421–432. doi:10.1007/s00726-022-03140-w

The UniProt Consortium, Bateman A, Martin M-J, Orchard S, Magrane M, Agivetova R,

Ahmad S, Alpi E, Bowler-Barnett EH, Britto R, Bursteinas B, Bye-A-Jee H, Coetzee

R, Cukura A, Da Silva A, Denny P, Dogan T, Ebenezer T, Fan J, Castro LG, Garmiri

P, Georghiou G, Gonzales L, Hatton-Ellis E, Hussein A, Ignatchenko A, Insana G,

Ishtiaq R, Jokinen P, Joshi V, Jyothi D, Lock A, Lopez R, Luciani A, Luo J, Lussi Y,

MacDougall A, Madeira F, Mahmoudy M, Menchi M, Mishra A, Moulang K,

Nightingale A, Oliveira CS, Pundir S, Qi G, Raj S, Rice D, Lopez MR, Saidi R,

Sampson J, Sawford T, Speretta E, Turner E, Tyagi N, Vasudev P, Volynkin V, Warner

K, Watkins X, Zaru R, Zellner H, Bridge A, Poux S, Redaschi N, Aimo L, Argoud-Puy

G, Auchincloss A, Axelsen K, Bansal P, Baratin D, Blatter M-C, Bolleman J, Boutet E,

Breuza L, Casals-Casas C, de Castro E, Echioukh KC, Coudert E, Cuche B, Doche M,

Dornevil D, Estreicher A, Famiglietti ML, Feuermann M, Gasteiger E, Gehant S,

Gerritsen V, Gos A, Gruaz-Gumowski N, Hinz U, Hulo C, Hyka-Nouspikel N, Jungo

F, Keller G, Kerhornou A, Lara V, Le Mercier P, Lieberherr D, Lombardot T, Martin X, Masson P, Morgat A, Neto TB, Paesano S, Pedruzzi I, Pilbout S, Pourcel L, Pozzato M, Pruess M, Rivoire C, Sigrist C, Sonesson K, Stutz A, Sundaram S, Tognolli M, Verbregue L, Wu CH, Arighi CN, Arminski L, Chen C, Chen Y, Garavelli JS, Huang H, Laiho K, McGarvey P, Natale DA, Ross K, Vinayaka CR, Wang Q, Wang Y, Yeh L-S, Zhang J, Ruch P, Teodoro D. 2021. UniProt: the universal protein knowledgebase in 2021. *Nucleic Acids Research* **49**:D480–D489. doi:10.1093/nar/gkaa1100

Uetsuka S, Ogata G, Nagamori S, Isozumi N, Nin F, Yoshida T, Komune S, Kitahara T, Kikkawa Y, Inohara H, Kanai Y, Hibino H. 2015. Molecular architecture of the stria vascularis membrane transport system, which is essential for physiological functions of the mammalian cochlea. *European Journal of Neuroscience* **42**:1984–2002. doi:10.1111/ejn.12973

Vanslambrouck JM, Bröer A, Thavyogarajah T, Holst J, Bailey CG, Bröer S, Rasko JEJ. 2010. Renal imino acid and glycine transport system ontogeny and involvement in developmental iminoglycinuria. *Biochemical Journal* **428**:397–407. doi:10.1042/BJ20091667

Vernocchi P, Del Chierico F, Putignani L. 2016. Gut Microbiota Profiling: Metabolomics Based Approach to Unravel Compounds Affecting Human Health. *Frontiers in Microbiology* **7**:1144. doi:10.3389/fmicb.2016.01144

Wei L, Tominaga H, Ohgaki R, Wiriyasermkul P, Hagiwara K, Okuda S, Kaira K, Kato Y, Oriuchi N, Nagamori S, Kanai Y. 2016. Transport of 3-fluoro-L-α-methyl-tyrosine (FAMT) by organic ion transporters explains renal background in [^{18}F]FAMT positron emission tomography. *Journal of Pharmacological Sciences* **130**:101–109. doi:10.1016/j.jphs.2016.01.001

Wickham H. 2010. A Layered Grammar of Graphics. *Journal of Computational and Graphical*

Statistics **19**:3–28. doi:10.1198/jcgs.2009.07098

Wiriyasermkul P, Moriyama S, Nagamori S. 2020. Membrane transport proteins in

melanosomes: Regulation of ions for pigmentation. *Biochimica et Biophysica Acta*

(BBA) - Biomembranes **1862**:183318. doi:10.1016/j.bbamem.2020.183318

Wiriyasermkul P, Nagamori S, Tominaga H, Oriuchi N, Kaira K, Nakao H, Kitashoji T, Ohgaki

R, Tanaka H, Endou H, Endo K, Sakurai H, Kanai Y. 2012. Transport of 3-Fluoro-L-α-

Methyl-Tyrosine by Tumor-Upregulated L-Type Amino Acid Transporter 1: A Cause

of the Tumor Uptake in PET. *Journal of Nuclear Medicine* **53**:1253–1261.

doi:10.2967/jnumed.112.103069

Wolosker H. 2018. The Neurobiology of D-Serine Signaling. *Advances in Pharmacology*

82:325–348. doi:10.1016/bs.apha.2017.08.010

Wolosker H, Sheth KN, Takahashi M, Mothet J-P, Brady RO, Ferris CD, Snyder SH. 1999.

Purification of serine racemase: Biosynthesis of the neuromodulator D-serine.

Proceedings of the National Academy of Sciences USA **96**:721–725.

doi:10.1073/pnas.96.2.721

1168

Figures and Tables

Figure 1. Wiriyasermkul, et al.

Figure 1. Enantiomeric profiles of D- and L-amino acids in plasma and urine of the IRI model

Plasma and urine were collected from the mice after ischemia operation for 4, 8, 20, and 40 h

or sham operation (0 h). The concentrations of twenties amino acids were measured by 2D-

HPLC and plotted as mean ± SD. (**A**) Ratio of D-/L-amino acids from the plasma of the IRI

model. The graph of Met is shown separately. n = 3. (**B**) – (**D**) Plasma amino acid profiles of

serine, alanine, and proline from (**A**) were normalized with 0 h and plotted as ratios of D-/L-

enantiomers (**B**), relative concentrations of D-isomers (**C**), and relative concentrations of L-

isomers (**D**). (**E**) Ratio of D-/L-amino acids from the urine of the IRI model. n = 3 – 7. (**F**) –

(**H**) Urinary amino acid profiles of serine, alanine, and proline from (**E**) were normalized with

those at 0 h and plotted as ratios of D-/L- enantiomers (**F**), relative concentrations of D-isomers

(**G**), and relative concentrations of L-isomers (**H**). C/CC: cysteine or cystine.

Figure 2. *Molecular Transport* in renal BBMVs proteome of the IRI model

(**A**) Ingenuity Pathway Analysis (IPA) shows the heatmaps of *Molecular Transport* in the membrane proteome from BBMVs of the IRI model (4h IRI/sham (4h) or 8h IRI/sham (8h)). Transport functions are categorized by types of substrates. Transport of all micromolecules (top) is derived from the combination of inorganic (middle) and organic compounds (bottom). Area and colors represent -$\log_{10}$(p-value) and annotated functions (z-score), respectively.

Numbers inside the columns indicate the numbers of corresponding proteins. (**B**) Heatmaps of membrane transport proteins that mediate the transport of organic compounds. The proteins with ratios of more than 1.1-fold change are shown. The category "others" includes the transport of peptides, organic cations, organic anions, and drugs. Colors indicate $\log_2$ fold of 4h IRI/sham (4h) or 8h IRI/sham (8h).

Figure 3. ASCT2 is one of D-serine transporters at the apical membrane of renal proximal tubular epithelia.

Figure 3. ASCT2 is one of D-serine transporters at the apical membrane of renal proximal tubular epithelia.

(**A**) – (**C**) Localization of Asct2 in mouse kidney by immunofluorescence staining. Mouse kidney slides were co-stained with anti-Asct2(NT) antibody (Asct2; green) and protein markers for renal proximal tubule segments: anti-Sglt2 antibody (**A**: Sglt2, apical membrane marker of S1 + S2 segments), anti-Agt1 antibody (**B**: Agt1, apical membrane marker of S3 segment), and anti-Na$^+$/K$^+$-ATPase antibody (**C**: Na$^+$/K$^+$-ATPase, basolateral membrane marker). Merge images are shown in the right panel. Arrowheads indicate co-localization of the proteins. Scale bar = 20 μm. (**D**) Time course of 100 μM D-[^{3}H]serine transport in wild-type (WT) and *ASCT2-*

1205   knockout (ASCT2-KO) HAP1 cells measured in PBS pH 7.4. Dot plot = mean ± SEM; n = 3.

1206   E) Left: Western blot using anti-ASCT2 antibody verified the suppression of ASCT2 in

1207   ASCT2-siRNA transfected HEK293 cells. Right: Transport of 100 μM D-[^{3}H]serine (in PBS

1208   pH 7.4) was measured in ASCT2-knockdown (ASCT2-siRNA) in comparison to the Mock

1209   cells (control). Dot plot = mean ± SEM; n = 3.

1210

Figure 4. Identification of SMCT1 and SMCT2 as candidates of D-serine transporters

(**A**) Volcano plots of 398 membrane transport proteins identified from the BBMV proteome of

the IRI model. The Median of $\log_2$ fold of 4h IRI/sham (left) or 8h IRI/sham (right) were plotted

against $-\log_{10}$ of p-value. Three proteins with $\log_2$ fold more than 4.0 (see values in Table S1)

are omitted for the better view. The value of Asct2 (red dot) was set as a cut-off value to select

D-serine transporters candidates (both increased (red area) and decreased (blue area)

expressions). (**B**) Cell-growth measurement (XTT assay) of ASCT2-siRNA or without siRNA (control) transfected HEK293 cells treated with D-serine. Data represent percent cell growth compared to the non-treated cells. The graphs were fitted to inhibition kinetics (Dose-response – Inhibition). Dot plot = mean ± SEM; n = 5. (**C**) – (**D**) Candidates of D-serine transporters were screened by cell-growth measurement. HEK293 cells were transfected with various cDNA clones, as indicated. After transfection, the cells were treated with either 15 mM (**C**) or 25 mM (**D**) D-serine for two days and cell growth was examined. The growth effect by D-serine treatment was normalized with that of no treatment and then calculated as $\log_2$ fold change of Mock at the same D-serine concentration. Asc-1 is used as the positive control. Bar graph = mean ± SEM; n = 4; *$p < 0.05$; **$p < 0.01$.

Figure 5. Characterization of SMCT1 and SMCT2 as D-serine transporters using SMCT1- and SMCT2-stably expressing cells

(**A**) Inhibition effect of ibuprofen on D-serine-induced cell growth. FlpInTR-Mock (Mock), FlpInTR-SMCT1 (SMCT1), or FlpInTR-SMCT2 (SMCT2) cells were treated with D-serine for two days in the presence or absence of 0.5 mM ibuprofen. Cell growth was measured by XTT assay. For comparison, the maximum growth inhibition by 25 mM D-serine treatment was

set as 100 % inhibition, and no D-serine treatment was set as 0 % inhibition. The graphs were fitted to inhibition kinetics (Dose-response – Inhibition). Dot plot = mean ± SEM; n = 5. (**B**) Time course of 100 μM D-[^{3}H]serine uptake in FlpInTR-SMCT1 (SMCT1), FlpInTR-SMCT2 (SMCT2) and Mock cells. D-[^{3}H]Serine transport was measured in PBS pH 7.4. Dot plot = mean ± SEM; n = 4. (**C**) Time course of 100 μM D-[^{3}H]serine uptake was measured similarly to B), but the cells were subjected to ASCT2 knockdown (siRNA transfection) two days before the assay. Dot plot = mean ± SEM; n = 4. (**D**) Transport of 20 μM D-[^{3}H]serine by ASCT2-siRNA-transfected FlpInTR-SMCT1 or FlpInTR-SMCT2 stable cell lines were measured in the absence (-) or presence of 5 mM indicated inhibitors. The uptake was measured for 10 min in PBS pH 7.4. Graphs represented the uptake data subtracted from those of Mock cells. Bar graph = mean ± SEM; n = 3. IBU, Ibuprofen; ASA, Acetylsalicylic acid. (**E**) Concentration dependence of D-[^{3}H]serine transport in ASCT2-siRNA-transfected FlpInTR-SMCT1 cells. Uptake of D-[^{3}H]serine (0.5 – 8 mM) was measured for 10 min in PBS pH 7.4. Raw data was shown in Figure supplement 5D. The uptake data in FlpInTR-SMCT1 were subtracted from those of Mock cells and fitted to Michaelis-Menten plot with the apparent K_m of 3.39 ± 0.79 mM and V_{max} of 18.23 ± 1.73 pmol/μg protein/min. Dot plot = mean ± SEM; n = 3 – 4.

Figure 6. Characterization of SMCT1 as D-serine transporter using SMCT1-reconstituted proteoliposomes (SMCT1-PL)

(**A**) Left: Stain-free SDS-PAGE gel shows SMCT1 (purified SMCT1) purified from pCMV14-SMCT1-transfected Expi293F cells, reconstituted empty liposomes (Liposome) and SMCT1-reconstituted proteoliposomes (SMCT1-PL). (B) Time course of D-[^{3}H]serine transport in

SMCT1-PL. Uptake of 200 μM D-[^{3}H]serine was measured in Na$^+$-containing buffer (Na$^+$)

compared to Na$^+$-free buffer (K$^+$). Dot plot = mean ± SEM; n = 3. (**C**) Ibuprofen effect on the

uptake of [^{3}H]lactate, [^{3}H]propionate, L-[^{3}H]serine, and D-[^{3}H]serine in SMCT1-PL. Uptakes

of 50 μM radiolabeled substrates were measured for 5 min in Na$^+$-containing buffer (Na$^+$) or

Na$^+$-free buffer (K$^+$) in the presence or absence of 1 mM ibuprofen. Bar graph = mean ± SEM;

n = 3. (**D**) Amino acid selectivity of SMCT1-PL. Transport of 50 μM radiolabeled amino acids

was measured in SMCT1-PL for 5 min. The substrate uptake in Na$^+$-containing buffer was

subtracted from those in Na$^+$-free buffer and calculated as % lactate uptake. Bar graph = mean

± SEM; n = 4.

Figure 7. Characterization of D-serine transporters in BBMVs of normal mice and the IRI model

(**A**) Transport of 10 μM D-[³H]serine in renal BBMVs isolated from normal mice. The uptake was measured in Na⁺-free buffer (K⁺), Na⁺-containing buffer (Na⁺), or the presence of 1 mM ibuprofen (Na⁺ + Ibuprofen). Dot plot = mean ± SEM; n = 3. (**B**) Transport of 10 μM D-[³H]serine in renal BBMVs isolated from the normal mice was performed similarly to (**A**) but the BBMVs were preloaded with 4 mM L-Gln prior to the measurement. Dot plot = mean ± SEM; n = 4. (**C**). Transport of 10 μM D-[³H]serine in renal BBMVs isolated from IRI model. Prior to uptake measurement, the BBMVs were preloaded with 4 mM L-Gln or buffer (no preload). The uptake was measured for 1 min in Na⁺-free buffer (K⁺), Na⁺-containing buffer

1276   (Na$^+$), or the presence of 1 mM ibuprofen (Na$^+$ + Ibuprofen). Bar graph = mean ± SEM; n = 3

1277   − 4.

Figure 8. Proposed model of D-serine transport systems in renal proximal tubules

The model summarizes the contributions of ASCT2, SMCT1, and SMCT2 to D-serine transport in renal proximal tubules. Left: the physiological conditions (Normal kidney). SMCT2 at S1 + S2 segments and SMCT1 at segment S3 have high expression levels, whereas the expression of ASCT2 is low. The higher D-serine affinity of SMCT1 compared to SMCT2 supports that the D-serine reabsorption tends to be exceeded at the S3 segment. Right: the pathological conditions (IRI model). expressions of SMCTs decrease while that of ASCT2 increases. ASCT2 expresses ubiquitously and exhibits high D-serine affinity. Blue arrows anticipate total D-serine fluxes in each proximal tubular segment.

1289 **Table 1**. Candidate transporters from proteomics of the BBMVs from the IRI model. The list
1290 is ordered according to the fold change.

Transporters	Accession	log$_2$ fold 4h IRI/ sham	p-value of 4h IRI/ sham	log$_2$ fold 8h IRI/ sham	p-value of 8h IRI/ sham	Peptides	Score Mascot	Abundance in sham[a]
Increased								
Slc7a1/Cat1	Q09143	0.8	0.12	1.3	0.01	1	131	2.6E+06
Slc1a5/Asct2	P51912	0.4	0.10	0.7	0.09	4	767	1.5E+07
Decreased								
Slc36a1/Pat1	Q8K4D3	-1.2	0.08	-1.0	0.12	3	103	5.4E+06
Slc2a5/Glut5	Q9WV38	-1.2	0.01	-0.8	0.04	2	1,140	6.5E+07
Slco4c1/Oatp-m1	Q8BGD4	-1.1	0.00	-0.9	0.00	9	4,011	1.5E+08
Slc22a13/Oat10	Q6A4L0	-1.1	0.03	-0.8	0.03	15	5,948	5.6E+08
Slc22a7/Oat2	Q91WU2	-0.9	0.00	-0.6	0.02	13	6,100	4.3E+08
Slc6a18/B^{0}at3	O88576	-0.8	0.17	-0.6	0.25	11	4,310	2.1E+08
Slc5a8/Smct1	Q8BYF6	-0.8	0.05	-0.7	0.02	25	26,044	2.9E+09
Slc15a2/Pept2	Q9ES07	-0.7	0.01	-1.1	0.05	9	2,379	4.6E+07
Slc5a12/Smct2	Q49B93	-0.6	0.01	-0.9	0.01	12	4,725	1.6E+08
TMEM27/ Collectrin[b]	Q9ESG4	-0.4	0.01	-0.1	0.17	9	23,686	3.3E+09

[a]Median from n = 3
[b]Collectrin is a regulatory protein for B^{0}at3 function

1291