1   **Short Title**: Interplay between autophagy and lipid metabolism
2
3
4
*Corresponding author information:

Tamar Avin-Wittenberg

Department of Plant and Environmental Sciences,

Alexander Silberman Institute of Life Sciences,

The Hebrew University of Jerusalem,

9190401, Jerusalem, Israel

E-mail: tamar.avin-wittenberg@mail.huji.ac.il

Tel: +972 2 5494023

Wagner L. Araújo

Departamento de Biologia Vegetal,

Universidade Federal de Viçosa,

36570-900 Viçosa, Minas Gerais, Brazil

E-mail: wlaraujo@ufv.br

Tel: +55 31 3612.5358

25
26

Title:

Autophagy is required for lipid homeostasis during dark-induced senescence in

Arabidopsis

Authors: Jessica A. S. Barros[a,b], Sahar Magen[b], Taly Lapidot-Cohen[c], Leah Rosental[c],

Yariv Brotman[c], Wagner L. Araújo[a*] and Tamar Avin-Wittenberg[b*]

Affiliation:

[a] Departamento de Biologia Vegetal, Universidade Federal de Viçosa, 36570-900

Viçosa, Minas Gerais, Brazil

[b] Department of Plant and Environmental Sciences, Alexander Silberman Institute of

Life Sciences, Hebrew University of Jerusalem, Givat Ram, 9190401, Jerusalem, Israel

[c] Department of Life Sciences, Ben-Gurion University of the Negev, 8410501, Beer-

Sheva, Israel

*Author for Correspondence

One-sentence summary: Autophagy is required for the mobilization of membrane lipid

components and lipid droplet dynamics during extended darkness in *Arabidopsis*.

Author contributions:
J.A.S.B., T.A-W., and W.L.A. designed the research; J.A.S.B. performed most of the

research; T.A-W. and W.L.A. supervised the project; Y.B. contributed with new

reagents/analytic tools for lipid analysis; S.M contributed with transcript analysis; T.L-

C perfomed lipid profile; L.R. perfomed the free fatty acid analysis; J.A.S.B., T.A-W.,

and W.L.A. analyzed the data and wrote the article with contributions of all the authors.

Funding information:
60
This work was made possible through financial support from the Serrapilheira Institute

(grant Serra-1812-27067). Work at the TAW group is supported by the Israeli Science

Foundation, grant number 1899/16. Scholarships granted by the National Council for

Scientific and Technological Development (CNPq-Brazil) and Council for Higher

Education (Malag-Israel) to JASB and research fellowships awarded by CNPq-Brazil to

WLA are also gratefully acknowledged.

Abstract

Autophagy is an evolutionarily conserved mechanism that mediates the degradation of cytoplasmic components in eukaryotic cells. In plants, autophagy has been extensively associated with the recycling of proteins during carbon starvation conditions. Even tough lipids constitute a significant energy reserve, our understanding of the function of autophagy in the management of cell lipid reserves and components remains fragmented. To further investigate the significance of autophagy in lipid metabolism, we performed an extensive lipidomic characterization of Arabidopsis (*Arabidopsis thaliana*) autophagy mutants (*atg*) submitted to dark-induced senescence conditions. Our results revealed an altered lipid profile in *atg* mutants, suggesting that autophagy affects the homeostasis of multiple lipid components under dark-induced senescence. The acute degradation of chloroplast lipids coupled with the differential accumulation of triacylglycerols (TAGs) and plastoglobuli-related transcripts indicates an alternative metabolic reprogramming towards lipid storage in *atg* mutants. The imbalance of lipid metabolism compromises the production of cytosolic lipid droplets and the regulation of peroxisomal lipid oxidation pathways in *atg* mutants.

Introduction

The precise control of energy homeostasis is crucial for plant survival, particularly under biotic and abiotic stress conditions. Under low energy availability, plants require a strategy, other than carbohydrate oxidation, to fully sustain energy production. Therefore, proteins, lipids, and chlorophylls can be catabolized instead of sugars to provide alternative substrates for the tricarboxylic acid cycle. This assures the continued operation of the mitochondrial electron transport chain (Ishizaki et al., 2005; Kunz et al., 2009; Araujo et al., 2011; Hortensteiner & Krautler, 2011).

Among the alternative substrates of respiration in plants, lipids comprise a significant carbon reserve that represents an important alternative source of energy. Although in comparison to carbohydrates, the usage of lipids is less efficient, as assessed by the respiratory quotient, their intracellular storage and utilization are critical for the maintenance of cellular energy homeostasis (Araujo et al., 2011; Yang et al., 2018). Plants store lipids as the neutral lipid species TAG (Yang et al., 2018). In leaf mesophyll cells, TAGs are deposited in specialized storage compartments, namely lipid droplets (LDs) in the cytosol or plastoglobuli (PGs) inside chloroplasts (Van wick et al., 2017). During nutrient deprivation, cellular lipid reserves are hydrolyzed into fatty acids (FAs), which are further catabolized during β-oxidation. Accordingly, mutants with dysfunctional β-oxidation, triggered by the disruption of the PEROXISOMAL ABC TRANSPORTER 1 (PXA1), exhibit hypersensitivity to extended darkness (Kunz et al., 2009). Moreover, several transcripts of the β-oxidation pathway are up-regulated during developmental leaf senescence (Troncoso-Ponce et al., 2013).

Autophagy, one of the primary cellular catabolic process, is involved in the degradation and recycling of cytoplasmic constituents into primary metabolites (Avin-Wittenberg et al., 2018). The autophagy mechanism is governed by autophagy-related (ATG) genes that are highly conserved among eukaryotes (Marshall and Vierstra, 2018). Plants with genetic disruption of *ATG* genes (hereafter *atg* mutants) have been well characterized for sensitivity to several stresses, particularly nutrient starvation (Avin-Wittenberg, 2018). The importance of autophagy in plants has been mostly associated with the recycling of proteins to amino acids under starvation conditions (Li et al., 2014, Izumi et al., 2013; Barros et al., 2017: Hirota et al., 2018; Mcloughin et al., 2018, 2020). By contrast, in mammals and yeast, autophagy has been largely associated not only with proteins but also with lipid metabolism (Elander et al., 2018; Barros et al., 2020). Accordingly, it was shown that autophagy supplies FAs for LD biogenesis as

well as mediates the selective degradation of LDs in both mammals and yeast (Singh et
al., 2009; Rambolt et al., 2015; Ngyugen et al., 2017).

A growing body of evidence suggests a dual role of autophagy in the lipid
metabolism of alga and plants. Thus, in Chlamydomonas (*Chlamydomonas reinhardtii*),
it has been postulated that autophagy participates in both TAG biosynthesis and LD
formation (Couso et al., 2017; Pugkaew et al., 2017). In addition, microscopic evidence
pointed to the autophagy-mediated engulfment of LDs in the algae *Auxenochlorella*
protothecoides (Zhao et al., 2014) and *Micrasterias denticulata* (Schwarz et al., 2017).
Interestingly, recent data also suggested that autophagy may mediate LD formation
during an early stage of starvation, while likely operating in LD degradation at the later
stages of nitrogen starvation in Chlamydomonas (Tran et al., 2019).

In vascular plants, it was demonstrated that autophagy contributes to TAG
synthesis via the turnover of organellar membranes in Arabidopsis under normal growth
conditions (Fan et al., 2019). On the other hand, it was suggested the autophagy
operates in the degradation of LDs by a selective microautophagy mechanism under
extended darkness (Fan et al., 2019). Notably, this study was based on the analysis of
Aradiposis mutants overexpressing the LD marker OLE1-GFP and knockout for
SUGAR DEPENDENT-1 (SDP1) lipase, which are genotypes with altered LD and
TAG metabolism (Fan et al., 2013, 2017). In maize (*Zea mays*), *atg12-1* mutant
presented a higher LD number, which could reflect active LD assembly or disruption of
LD catabolism (Mcloughin et al., 2020). It remains contentious, however, whether
autophagy mediates the biogenesis or degradation of LDs in vegetative tissues (Elander
et al., 2018; Barros et al., 2020; Masclaux-Daubresse et al., 2020). Although recent
studies have clearly revealed an interplay between autophagy and lipid metabolism
(Mcloughin et al.,2018, 2020; Have et al., 2019; Fan et al., 2019), relatively little is
known about the role of autophagy in the mobilization of lipid components under
energy limitation conditions.

Here, we investigated the significance of autophagy in lipid metabolism during
dark-induced senescence in Arabidopsis. By performing an extensive lipidomic analysis
of three independent *atg* mutants, we observed an overall impact of autophagy
deficiency on all lipid classes analyzed and on the free fatty acids (FFA) profile. The
differential pattern of phospholipid levels and TAG accumulation was further
coordinated with changes in LD number and PG-related transcripts. In addition, analysis
of β-oxidation genes revealed an induction of lipid oxidation pathways in *atg* mutants.

Collectively, our results demonstrate that autophagy is an essential component in the
management of lipid reserves, affecting membrane lipid mobilization, the balance of LD
and PG formation, and the regulation of β-oxidation pathways under dark-induced
senescence.
172
173

Results

Differential lipid response of *atg* mutants under dark-induced senescence

To examine the involvement of autophagy in lipid remodeling during dark-

induced senescence, we analyzed three previously described Arabidopsis autophagy

mutants: *atg5-1* and *atg7-2* with full inhibition of autophagy (Chung et al., 2010); and

atg9-4 mutant that displayed a milder reduction of the autophagic process (Shin et al.,

2014). The experimental setup consisted of the submission of 4-week old plants to

extended darkness conditions, a system widely used to induce energetic stress in plants

(Ishizaki et al., 2005; Araujo et al., 2010; Barros et al., 2017).

To further investigate the impact of autophagy disruption on lipid homeostasis

during dark-induced senescence, we performed Liquid Chromatography–Mass

Spectrometry lipidomic analysis of *atg5-1*, *atg7-2*, *atg9-4,* and their corresponding

wild-type (WT) plants. We were able to identify 119 lipid species of the following

classes: Phosphatidylcholine (PC), Phosphatidylethanolamine (PE), Diacylglycerol

(DAG), Triacylglycerol (TAG), Monogalactosyldiacylglycerol (MGDG) and

Digalactosyldiacylglycerol (DGDG). We first performed a principal component analysis

(PCA) based on the lipid profiles of samples from control and 6 days (d) of darkness

(Fig. 1). Most of the variance of the dataset was covered by the first two principal

components, explaining 69.1% of the overall data variance (57.8% and 11.3% for PC1

and PC2, respectively). This fingerprint analysis allowed us to obtain a general picture,

discerning the main differences between the darkness treatment and the *atg* mutant

lines. Briefly, it revealed that under control conditions (before darkness treatment), all

genotypes clustered together; however, a clear separation according to the treatment and

genotype was observed following darkness, wherein *atg7-2* and *atg5-1* were

disconnected of the WT samples (Fig. 1). Interestingly, *atg9-4* presented an

intermediary distribution, sharing certain metabolic similarities with WT and the other

atg genotypes. The analysis of all data points also highlighted the delayed response of

atg9-4, which after 6d and 9d of darkness, was more closely related to *atg5-1* and *atg7-*

2 after 3d of darkness conditions (Supplementary Figure S1).

Our results are in close agreement with the milder impact of the *ATG9*

disruption on metabolic features previously reported under dark-induced senescence

(Barros et al., 2017). This response was related to residual autophagy activity observed

in *atg9* lines (Yoshimoto et al., 2004; Shin et al., 2014), which explains the intermediary

lipid response of *atg9-4* genotype. The information of the individual variables that

Figure 1 - Differential lipid response of *atg* mutants under dark-induced senescence.
Principal component analysis (PCA) was performed using the lipid data obtained in samples from 4-week-old Arabidopsis plants immediately before lights were turned off (0 d) and after further treatment for 6d in darkness. Three loss-of-function mutants of the autophagy pathway, namely *atg5-1*, *atg7-2*, *atg9-4* and their respective wild type (WT) were analyzed.

contributed most to PC1 and PC2 (i.e., the lipids with the main impact on the variance
of the dataset) is additionally available in Supplemental Table S1. Several lipids
accounted for the main changes observed in the PCA. Briefly, PC1 was mainly
comprised of MGDG 34:4, MGDG 34:3, PC 38:4(1) and MGDG 34:1 underlining the
separation of darkened and control samples. On the other hand, TAG 54:9, MGDG 32:3
(1), and TAG 52:6 were the main lipids responsible for changes in PC2 (Supplemental
Table 1), contributing to the differences between the genotypes under darkness (Fig. 1).
215

216   **The lack of autophagy affects the dynamics of phospholipids during dark-induced**
217   **senescence**

218   In order to obtain a more detailed characterization of the lipid composition of
219   *atg* mutants following extended darkness, we analyzed each lipid class separately. We

Figure 2 - *atg* mutants present altered phospholipid levels under extended darkness conditions
Hierarchical clustering of (A) Phosphatidylcholine (PC) and (B) Phosphatidylethanolamine (PE). Values plotted are log2 fold change. Distances were measured by Pearson correlation using MeV 4.9.0 software. Data were normalized to the mean response calculated for the 0-d dark treated leaves of the wild type (WT). Values presented are means ± SE of five-six biological replicates per genotype; an asterisk (*) designates values that were determined by Student's t-test to be significantly different (P < 0.05) from WT at each time point analyzed.

were able to identify the two dominant structural phospholipid species, namely PC and

PE. These lipids are components of the plasma membrane, in which PC is associated

with the outer leaflet, and PE is concentrated in the inner leaflet. PC and PE are also

precursors of lipid mediators, such as phosphatidic acid (PA) and diacylglycerol (DAG)

(Wang et al., 2006; Testerink and Munnik, 2011). Additionally, PE plays a crucial role

in autophagosome formation (Soto-Burgo et al., 2018; Barros et al., 2020).

Interestingly, a slight accumulation of both PC and PE was observed in *atg5-1* and *atg7-*

2 mutants before darkness treatment (control conditions, Fig. 2 – 0d). The difference of

PC levels under control conditions was marked by the accumulation of 36C PC in *atg5-*

1 and *atg7-2* mutants, whereas no significant changes were observed in *atg9-4* at this

time point (Fig. 2A – 0d). Extended darkness triggered a mixed response of different PC

species. Thus, the levels of 32C and 34C mostly increased in WT plants under darkness.

However, *atg5-1* and *atg7-2* presented lower accumulation of 34C PC species that

contain a combination of 18C and 16C polyunsaturated acyl species, such as 34:5-

(18:3-16:2-) and 34:6- (18:3-16:3) during dark treatment. The PC containing FAs with

longer chains, 36C and 38C, were generally reduced in response to extended darkness.

However, *atg5-1* and *atg7-2* presented a milder reduction of 36C PC species after 3d

and 6d of darkness, whereas the 38C species (38:3(2), 38:4, 38:5, 38:6) were more

reduced in these genotypes after 6d and 9d of darkness (Fig 2A).

Following dark treatment, the majority of PE species decreases in WT plants;

however, this reduction was milder in *atg* mutant lines after 6d of darkness (Fig. 2B).

On the other hand, after 9d of darkness, species such as PE 36:2, 34:2, and 36:3(2) were

highly reduced in *atg5-1* and *atg7-2* compared to WT. Surprisingly, the PE species 36:5

and 38:3 presented a contrasting pattern, accumulating after 6d and 9d of darkness in

atg mutants (Fig. 2B).

Impairment of autophagy accelerates the loss of chloroplast lipids

Since chloroplast lipids are subjected to constant turnover during senescence, we

next evaluated the predominant lipid constituents of chloroplast membranes, MGDG

and DGDG. An apparent reduction in more than 90% of the detected MGDGs and

DGDGs was observed in *atg* mutants exposed to dark treatment (Fig. 3). This response

was intensified throughout the dark treatment, and thus the levels of some species were

below the detection threshold after 9d. Although *atg9-4* mutants followed the same

pattern, the levels of MGDG and DGDG were less reduced in comparison to *atg5-1*

and *atg7-2* mutants (Fig. 3). The overall reduction of chloroplast lipids reported here is

consistent with the more severe phenotype described previously for *atg5-1* and *atg7-2*

mutants lines under extended darkness (Barros et al., 2017). Interestingly, slightly lower

levels of unsaturated MGDG and DGDG were also observed in *atg* mutants under

control conditions, highlighting the potential role of autophagy in basal chloroplast

maintenance, as recently verified in different studies using *atg* mutants (McLoughlin et

al., 2018, 2020; Have et al., 2019).

Disruption of autophagy compromises TAG and LD response under dark-induced

senescence

The analysis of neutral lipids revealed a general increase in TAG levels during

extended darkness. TAGs are composed of three FAs esterified to a glycerol backbone

(Yang et al., 2018). The dominating accumulated TAG species during dark-induced

senescence harbor three FA molecule with 18C with two or three double bonds (54:7 to

54:9 in Fig. 4A), or two FA molecules with 18C with two to three double bonds and

16:3 or 16:0 as a third FA (52:5 to 52:9 in Fig. 4A). Unsaturated C18 FAs (oleic acid,

18:1; linoleic acid, 18:2, and α-linolenic acid, 18:3), Palmitic acid (16:0) and

Figure 3 - Chloroplast lipids are substantially reduced in *atg* mutants during dark-induced senescence
Monogalactosyldiacylglycerol (MGDG) and Digalactosyldiacylglycerol (DGDG) values plotted are log2 fold change. Data were normalized to the mean response calculated for the 0-d dark treated leaves of the wild type (WT). Values presented are means ± SE of five-six biological replicates per genotype; an asterisk (*) designate values that were determined by the Student's t-test to be significantly different (P < 0.05) from WT at each time point analyzed.

unsaturated C16 FAs (16:1 and 16:3) are highly abundant in chloroplasts of

Arabidopsis (Hölzl and Dörmann, 2019). Our data indicate that the accumulated TAG

species are most likely derived from plastidial lipids. Interestingly, the increase in

chloroplast-derived TAG species was less pronounced in both *atg5-1* and *atg7-2*

mutants after 3d and 6d of darkness. Additionally, an increase of other TAG species

was observed from 6d of dark treatment onwards in *atg5-1* and *atg7-2* mutants,

resulting in general TAG accumulation during the late period of darkness (Fig. 4A).

DAG is a central metabolite in plant lipid metabolism, serving as an

intermediate of both TAG and galactolipids (Muthan et al., 2013). Similar to the

observed for TAGs, the four DAG molecules identified (34:3, 34:6, 36:5, 36:6) were

composed of unsaturated FAs, containing at least one unit of 16:2, 16:3 or 18:3 plastid-

derived FA (Fig. 4B). Although no significant differences were observed during the

early stages of dark treatment, DAGs started to accumulate 6d after the onset of dark

treatment in *atg5-1* and *atg7-2* mutants. At the same time, WT and *atg9-4* plants

showed a trend of reduction (Fig. 4B). The accumulation of unsaturated DAG species is

likely a result of the degradation of chloroplast membranes in *atg* mutants serving as a

precursor for the general TAG accumulation in the late stages of darkness.

TAG is the central lipid reserve of the plant cell, stored in LDs (Pyc et al.,

2017). To further investigate how the altered levels of TAGs in *atg7-2* and *atg5-1*

Figure 4 - Distinct accumulation of neutral lipids in *atg* mutants during dark-induced senescence
(A) Triacylglycerol (TAG) and (B) Diacylglycerol (DAG). Values plotted are log2 fold change. Data were normalized to the mean response calculated for the 0-d dark treated leaves of the wild type (WT). Values presented are means ± SE of five-six biological replicates per genotype; an asterisk (*) designates values that were determined by Student's t-test to be significantly different (P < 0.05) from WT each time point analyzed.

mutants affect LD metabolism during dark-induced senescence, we visualized the

abundance of LDs in leaf tissues by staining with the neutral-lipid-selective fluorescent

dye, BODIPY493/503. An increased number of LDs was observed after 3d of darkness

in WT plants (Fig. 5). However, no accumulation of LDs was observed in *atg5-1* and

atg7-2 mutants (Fig. 5). Notably, the multiple z-stacked confocal images revealed that

LDs are mostly accumulated in the cytosol and not inside chloroplasts (Fig. 5A). The

dynamics of LD accumulation corroborate the high levels of the plastid-derived TAG

species, which were subsequently reduced after 6d of darkness in WT. Furthermore,

atg5-1 and *atg7-2* mutants displayed reduced levels of plastid-derived TAG species that

can also be associated with the impairment of LD accumulation (Fig. 4A).

The disruption of autophagy triggers activation of PG metabolism

PG are specialized globular LDs composed by a membrane lipid monolayer

surrounding a core of neutral lipids present in the chloroplast membranes (Van Wijck an

Figure 5 - Autophagy deficiency affects Lipid Droplet (LD) dynamics during dark-induced senescence
(A) Representative confocal images of LDs (BODIPY 493/503 fluorescence, false color green) in wild type (WT), *atg5-1*, and *atg7-2* leaves. Red color depicts chloroplast autofluorescence. Images were collected at the same magnification, and are projections of Z-stacks of 20 optical sections Bar = 10 μm (B) Number of total LDs per image area. The number of LDs was quantified by ImageJ as BODIPY-stained lipid area in 2D projections of Z-stacks of 3 images of four biological replicates per genotype and time point.

Kessler, 2017). The production of PG is related to the accumulation of products from

chlorophyll and chloroplast lipid catabolism such as fatty acid phytyl esters (FAPEs),

TAGs and FAs (Gaude et al., 2007; Lippold et al., 2012). Given that *atg* mutants

presented a substantial reduction of chloroplast lipids under extended darkness, which

was not translated to an increase in TAGs or LD accumulation, we next investigated

PG-related pathways by quantitative reverse transcription (RT)-PCR. The genes phytyl

ester synthase 1 (*PES-1*) and *PES-2,* belonging to the esterase/lipase/thioesterase family,

are required for FAPE production in Arabidopsis PG during senescence (Lippold et al.,

2012). Accordingly, we observed an increase of *PES-1* and *PES-2* transcript levels

following dark-induced senescence (Fig. 6A-B). However, higher induction of *PES-1*

in *atg5-1* and *atg7-2*, as well as of *PES-2* in *atg7-2* mutant, was observed after 3d and

6d of darkness (Fig. 6A-B). The production of FAPE relies on both free FA released

from chloroplast membranes and phytol from chlorophyll (Lippold et al., 2012). We,

therefore, analyzed the chlorophyll degradation enzyme, pheophytin pheophorbide

hydrolase (*PPH*). As observed for *PES1* and *PES2*, the *PPH* transcript accumulated at

higher levels in *atg* mutants under dark-induced senescence in comparison to WT levels

(Fig. 6D).

PG serve not only as a lipids deposit but contains an actively associated

proteome (Lundquist et al., 2012). The presence of several Absence of Bc1 Complex

Kinase (ABC1K) proteins was described in PG, suggesting a central role of these

Figure 6 - PG related genes are up-regulated in *atg* mutants during dark-induced senescence
Transcript abundance is shown for genes associated with plastoglobuli (PG), including PES-1 (A), PES-2 (B), ABC1K7 (C), PPH (D). The y-axis values represent the gene expression level relative to the wild type (WT). Data were normalized with respect to the mean response calculated for the 0-d dark-treated leaves of the WT. Values presented are means ± SE of three independent biological replicates. An asterisk (*) indicates values that were determined by Student's t-test to be significantly different (P < 0.05) from the WT at each time point analyzed. A number sign (#) indicates values determined by Student's t-test to be significantly different (P < 0.05) from WT at 0d.

proteins in the regulation of PG metabolism (Ytterberg et al., 2006; Brehelin et al.,

2007). The protein ABC1K7 has been characterized as a PG core component with

marked accumulation during late senescence events (Lundquist et al., 2012; Bhuiyan et

al., 2016). Despite the unchanged transcript levels of *ABC1K7* in WT, this gene was

induced in *atg5-1* and *atg7-2* mutants following extended darkness (Fig. 6C). Although

our results demonstrate a boost of PG pathways in the absence of autophagy under dark-

induced senescence, *atg* mutants presented a slight but significant up-regulation for all

PG-related genes even under control conditions (Fig. 6), highlighting intrinsic crosstalk

between these processes.

***atg* Mutants induce lipid oxidation pathways in response to dark-induced**

senescence

FA oxidation may be of particular significance when the carbon and energy

status is rather low. During carbon starvation, the importance of lipid metabolism was

demonstrated by the characterization of β-oxidation loss-of-function mutants (Kunz et

al., 2009). In addition, autophagy association with peroxisomal protein abundance was

recently demonstrated (McLoughlin et al., 2018, 2020; Have et al., 2019). The general

reduction of chloroplast lipids, together with reduced LD numbers in *atg5-1* and *atg7-2*

Figure 7 - Changes in transcript levels in WT and *atg* mutants during dark-induced senescence.
Transcript abundance is shown for genes associated with lipid catabolic pathways, including ACX-4 (A), CSY-1(B) KAT-2 (C), MPF-2 (D), pMDH-1 (E), SDP-1 (F). The y-axis values represent the expression level relative to the wild type (WT). Data were normalized with respect to the mean response calculated for the 0-d dark-treated leaves of the WT. Values presented are means ± SE of three independent biological replicates. An asterisk (*) indicates values that were determined by Student's t-test to be significantly different (P < 0.05) from the WT at each time point analyzed. A number sign (#) indicates values determined by Student's t-test to be significantly different (P < 0.05) from WT at 0d.

mutants subjected to extended darkness, raised the question of whether these lipids were

being quickly oxidized to be used as energetic substrates in *atg* mutants. We, therefore,

evaluated the expression of genes related to LD catabolism, and peroxisomal pathways

of FA oxidation.. We observed that the transcript levels of the peroxisomal enzymes

Acyl-CoA oxidase (ACX-4), Multifunctional protein 2 (MPF-2), Ketoacyl-CoA thiolase

(KAT-2), Citrate synthase 1 (CSY-1) and peroxisomal Malate Dehydrogenase

(pMDH1) genes were generally higher in *atg5-1* and *atg7-2* under extended darkness

(Fig. 7 A-E). More specifically, *ACX-4* and *CSY-1* were slightly induced in WT

throughout the treatment. However, higher transcript levels were observed in *atg5-1* and

atg7-2 plants after 6d of darkness (Fig. 7A-B). Despite no major changes in *MPF-2* and

KAT-2 transcript abundance in WT plants, trends of increase were observed in *atg5-1*

and *atg7-2* after 9d of darkness (Fig. 7 C-D). We further confirmed that *pMDH-1* was

downregulated upon dark-induced senescence. *pMDH-1* operates in the reduction of

oxaloacetate to malate and enables the re-oxidation of NADH for continued β-oxidation

(Pracharoenwattana et al., 2007). Given that *pMDH-1* is not participating directly in β-

oxidation, it is possible that an alternative pathway of NADH oxidation becomes

operational under carbon starvation.

To investigate whether the reduced number of LDs is a result of the induction of catabolic pathways or impairment in biosynthesis, we next evaluated the expression of *SDP-1* gene. Despite being characterized as the primary lipase mediating LD catabolism in seeds, compelling evidence has suggested that *SDP-1* also operates in senescent and carbon-starved leaves (Fan et al., 2017, 2019). Indeed, an accumulation of *SDP-1* transcript levels was observed in WT plants during the early stages of darkness (Fig. 7F). Nonetheless, no significant difference in *SDP-1* transcript levels between WT plants and *atg5-1* and *atg7-2* was observed, suggesting that the reduced levels of LDs are likely more related to the impairment of LD biogenesis rather than an induction of LD degradation pathways.

Altered FFA response of *atg* mutants under dark-induced senescence

Since autophagy deficiency altered cytosolic LD dynamics and β-oxidation pathways, we next evaluated if this response also affected the pool of free FA (FFA). Figure 8 shows that long-chain FFA (between 13C-21C chain length) are the more affected by both genotype and darkness, while no major changes were observed for very long-chain FFA (> 22C). Higher levels of 16:3, 18:1, 18:2, 18:3 in *atg5-1* and *atg7-2* genotypes under control conditions were observed, reinforcing the functional role of autophagy in basal lipid homeostasis. Upon darkness treatment, the levels of C18 and C20 FFAs were reduced, and unsaturated C16 were accumulated in *atg5-1* and *atg7-2* mutants (Fig. 8). By contrast, WT plants presented an early increase of C16 and C18 FFAs, which were later reduced after 6 days of dark-induced senescence. Apart from 16:3 and 18:1 FFAs, which were continuously accumulated and reduced, respectively, throughout darkness (Fig. 8). Although unsaturated C16 and C18 FAs are typical components of chloroplast membranes, only 16:3 is exclusively found in the thylakoid lipid MGDG in plant species (e.g. Arabidopsis) and it is, therefore, a chloroplast lipid marker (Ohlrogge et al., 1991; Kunz et al., 2009; Hölzl and Dörmann, 2019). Thus, the overall accumulation of 16:3 is in good agreement with the reduction of galactolipids observed under dark-induced senescence. Additionally, this result suggests that the rate of 16:3 release exceeds its channeling to LD and PG. The higher accumulation of 16:3 in *atg* mutants strengthens our claim that this disbalance is more intense when autophagy is disrupted.

Figure 8- Changes in free Fatty Acid levels in WT and *atg* mutant plants under dark-induced senescence.

Values plotted are log2 fold change. Data were normalized to the mean response calculated for the 0-d dark treated leaves of the wild-type (WT). Values presented are means ± SE of five-six biological replicates per genotype; an asterisk (*) designate values that were determined by the Student's t-test to be significantly different (P < 0.05) from WT each time point analyzed.

Discussion

It is well established that plant autophagy plays a crucial role in the energetic metabolism mediating the supply of proteins and amino acids under carbon starvation conditions (Avin-Wittenberg et al., 2015; Barros et al., 2017; Hirota et al., 2018; McLoughlin et al., 2020). Furthermore, in both animals and yeast, autophagy is recognized as an important lipid catabolic pathway in starved cells (Singh et al., 2009). In plant leaves, however, little is currently known regarding the function of autophagy in managing cell lipid components and reserves (Avin-Wittenberg et al., 2015; McLoughlin et al., 2018,2020; Fan et al., 2019; Have et al., 2019). Here, we provide further evidence that autophagy broadly affects lipid and LD metabolism during dark-induced senescence in Arabidopsis.

By performing a detailed lipidomics analysis, we revealed that autophagy widely affects the lipid homeostasis of Arabidopsis during dark-induced senescence. The major differences reported here are related to the *atg5-1* and *atg7-2* mutant plants, characterized by a full inhibition of autophagy (Chung et al., 2010). By contrast, *atg9-4*, which is characterized by a partial reduction of autophagy, presented an intermediary

impact on lipid metabolism that was more evident during the late stages of extended
darkness (Fig. 1 and Supplemental Figure S1). The fact that *atg9-4* did not cluster with
atg5-1 and *atg7-2* implies that the lipid response is directly related to autophagy
intensity. Indeed, the characterization of Arabidopsis *ATG*5 and *ATG*7 knockout and
overexpression mutants showed a positive correlation between the level of autophagy
and seed FA content (Minina et al., 2018). We previously demonstrated that amino acid
catabolic pathways were less compromised in *atg9-4* mutants, contributing to its more
tolerant phenotype during dark-induced senescence (Barros et al., 2017). Additionally,
the limited reproductive phenotype of *atg* mutants is also dependent on autophagy
intensity since parameters related to seed and silique production were more
compromised in *atg* lines fully defective in autophagy (Barros et al., 2017). Taking into
account the fact that Arabidopsis is an oil seed (Li et al., 2006), the results reported here
and by others (Minina et al., 2018) suggest that the altered reproductive and carbon
starved phenotype of *atg* mutants is likely not only a consequence of impairment of
protein catabolic process but is also derived from an extensive reprogramming of lipid
metabolism.

The detailed analysis of the specific lipid classes revealed an altered lipid
phenotype in mutants lacking autophagy under non-stress conditions. These differences
are mostly attributed to galactolipids and the phospholipids PC and PE, whereas minor
differences were observed for TAGs (Figs 2, 3, and 4). The galactolipids MGDG and
DGDG are prevalent in the chloroplast envelope and the thylakoid membrane (Hölzl
and Dörmann, 2019). By contrast, PC and PE are highly abundant in the plasma
membrane, mitochondrial membrane, and endoplasmic reticulum (Liu et al., 2015).
Hence, the high levels of PC and low levels of MGDG and DGDG reveal an imbalance
of cytosolic and plastidial membrane lipids in *atg5-1* and *atg7-2* mutants under normal
growth conditions. Similar results were previously reported in Arabidopsis *atg5*
mutants, which displayed a general alteration of phospholipid levels under non-stressful
conditions (Have et al., 2019). The accumulation of lipid-breakdown products in maize
atg12 mutants also suggested an induced turnover of membrane components under
normal growth conditions (McLoughlin et al., 2018, 2020). In addition, Fan et al., 2019
reported that the disrupted membrane lipid turnover in *atg* mutants affected TAG
synthesis. The TAG profile detailed here reveals that only specific TAG species are
reduced in *atg5-1,* and *atg7-2* mutants under normal growth conditions, which seems to
contribute to the low total TAG content previously reported.

Collectively, our results support the notion that, beyond the role of autophagy in
basal lipid metabolism, autophagy is clearly required for lipid remodeling under
starvation conditions in Arabidopsis. Upon exposure to extended darkness, PC
metabolism is broadly rearranged, presenting different patterns of accumulation and
reduction (Fig. 2). In general, the majority of 32C and 34C PC species increases,
whereas 36C and 38C declines upon darkness (Fig. 2). However, an altered response of
PC accumulation and degradation was observed in *atg* mutants. The turnover of
membrane lipids is a recognized aspect of plant senescence in general (Troncoso-Ponce
et al., 2013). It seems that the disruption of autophagy leads to a differential degradation
rate of membrane lipid components, most likely affecting the supply of FAs to sustain
other cellular processes, including respiration. Alteration of PE levels was also observed
in *atg* mutants under our experimental conditions (Fig. 2). Noteworthy, PE is an
essential component of ATG8 conjugation during autophagosome membrane biogenesis
(Soto-Burgo et al., 2018; Barros et al., 2020). The less accentuated reductions and
accumulation of several PE species (PE 36:4, 36:3(1), 38:2, 36:5, 38:3) in *atg* mutants
might be related to the disrupted usage and recycling of these molecules for
autophagosome biogenesis, as previously suggested (Avin-Wittenberg et al., 2015;
Have et al., 2019).

Apart from changes in the extra-plastidial phospholipids, *atg7-2* and *atg5-1*
mutants also presented intense degradation of chloroplastic MGDG and DGDG during
dark-induced senescence (Fig. 5). This data is consistent with the hypersensitive
phenotype marked by chloroplast dismantling observed in *atg* mutants submitted to
dark-induced senescence(Thompson et al., 2005; Phillips et al., 2008; Barros et al.,
2017). Accordingly, autophagy has been characterized to mediate the degradation of
chloroplast proteins by different selective processes (Ishida et al., 2008; Wada et al.,
2009; Michaeli et al., 2014; Izumi et al., 2017; Nakamura et al., 2018). This fact aside,
the results shown here reinforce the hypothesis that other pathways likely mediate the
degradation of chloroplast lipid components in carbon-starved plants and that they are
likely to be more active when autophagy is absent. Moreover, this response is not
restricted to carbon limitation, as similar results were reported in both Arabidopsis and
maize *atg* mutants under low-nitrogen conditions (McLoughlin et al., 2018; Have et al.,
2019).

The breakdown of membrane lipid components releases FFAs for further
oxidation in peroxisomes. The overaccumulation of FFAs is detrimental to the plant

cell. Consequently, FFAs are first incorporated as neutral lipids, to then be directed to

peroxisomal FA catabolic pathways (Fan et al., 2017). TAGs stored in LDs are not

abundant in leaf mesophyll cells. However, their number substantially increases under

abiotic conditions and developmental senescence (Pyc et al., 2017; Yang et al., 2018).

In agreement with this, we observed a transient LD accumulation after 3d of darkness in

WT plants. In addition, a marked increase of highly unsaturated TAG levels was

observed after 3d of darkness in WT (Fig. 3). Despite an increase of some FFA species

after 3d of darkness, this increase was not overloaded to result in a lipotoxic effect,

since WT plants present favorable performance under dark-induced senescence (Barros

et al., 2017). Indeed, the pattern of initial accumulation and later reduction of FFA 16:2,

18:0, 18:2, 18:3 was similar to the observed for TAG and LD (Figs. 4, 5 and 8),

suggesting a dynamic equilibrium between membrane consumption, lipid reserve

formation and FA utilization in WT plants. By contrast, not only LD accumulation was

disrupted in *atg5-1* and *atg7-2* mutants, but they also showed lower levels of

unsaturated TAGs and higher accumulation of plastidial FFA 16:3 after 3d and 6d of

darkness. Collectively, these results demonstrate that autophagy is required for the

proper mobilization of chloroplast lipids destined for cytosolic LD biosynthesis in

Arabidopsis starved cells. A recent study of maize *atg12* mutants subjected to carbon

deprivation reported distinct LD dynamics (McLoughlin et al., 2020). It was observed

that *atg12* mutants contained more cytoplasmic LD under normal growth conditions. In

individually darkened leaves, LD numbers were not affected in WT and only slightly

reduced in the *atg12* mutant. A possible explanation for this discrepancy may stem from

the experimental systems utilized. While McLoughlin et al. (2020) analyzed

individually darkened leaves of maize, we examined Arabidopsis in a whole plant

darkening system. Plants have divergent metabolic strategies in response to different

darkening treatments (Law et al., 2018). It has been postulated that individually

darkened leaves still receive sugars from other parts of the plant during short-term dark

periods. However, the lipid catabolic response is not activated as long as carbohydrates

are available, which likely affects LD dynamics (Kunz et al., 2009; Law et al., 2018).

Not only LD responses diverge according to the darkness treatment imposed, but the

results reported until now also open new avenues regarding contrasting LD responses in

different plant species.

In Arabidopsis, autophagy operates in the mobilization of FAs from organelle

membranes to TAG synthesis under normal growth conditions (Fan et al., 2019). The

same study stated that autophagy has no influence on TAG formation during darkness, but instead operates in LD degradation by lipophagy mechanism. Despite that, general TAG accumulation was observed in *atg* mutants after 6d of darkness, which was not accompanied by cytosolic LD accumulation (Figs 4 and 5). Nonetheless, caution must be taken when comparing the results obtained here with previous findings (Fan et al., 2019) due to the differences of the genotypes analyzed (Barros et al., 2020). Thus, although Fan et al. (2019) demonstrated the existence of lipophagy in different mutant backgrounds, we analyzed plants solely lacking autophagy. When considered together, our results, coupled with the results from Fan and colleagues, highlight the potential role of autophagy as a versatile player in LD metabolism, particularly in starved cells.

Beyond the LDs in the cytosol, PG serve as subcompartments for lipid biosynthesis and storage in chloroplasts (Rottet et al., 2015; van Wijk and Kessler, 2017). The analysis of PG-related genes indicates a marked activation of PG metabolism in *atg5-1* and *atg7-2* mutants during dark-induced senescence (Fig. 6). There is a strong correlation between thylakoid dismantling and supersizing of PG during senescence, which is likely associated with the massive deposit of FAs and phytol, released from chloroplast dismantling (Lippold et al., 2012; Besagni et al., 2013; Rottet et al., 2015; van Wijk and Kessler, 2017). In fact, a positive correlation between PG core proteins and chlorophyll degradation enzymes was previously identified during developmental senescence (Lundquist et al., 2012). In light of this, we also verified an increase of *PPH* transcript in *atg5-1* and *atg7-2* mutants compared to WT under darkness (Fig. 6D). Thus, the reduced levels of chlorophyll (Barros et al., 2017) and the massive degradation of chloroplast lipids in *atg* mutants submitted to extended darkness (Fig. 3) may be correlated to the induction of PG pathways (Fig. 6). This response can be presumed as a strategic mechanism preventing the accumulation of lipotoxic intermediaries. Collectively, our results imply that the disruption of autophagy triggers a differential response prompting PG rather than cytosolic LD formation under dark-induced senescence. Nonetheless, whether PG can be remobilized to provide lipid-derived energetic substrates remains to be investigated.

It is already known that, during extended darkness, FAs are oxidized by β-oxidation, generating acetyl-CoA that, combined to oxaloacetate, produces citrate to be exported to mitochondria for ATP generation (Kunz et al., 2009). To further understand whether the released membrane lipid components were channeled to the maintenance of the energetic status, we paid particular attention to the transcriptional activation of lipid

oxidation pathways. Our results show an up-regulation of peroxisomal genes in plants
with disrupted autophagy after 6d and 9d of darkness (Fig. 7). Notably, the
transcriptional induction of β-oxidation has been correlated to mitochondrial activity in
darkened leaves (Law et al., 2018). Interestingly, the same *atg* mutants exhibited higher
rates of respiration and an elegant metabolic reprogramming under carbon starvation
conditions (Avin-Winttenberg et al., 2015; Barros et al., 2017).

Autophagy also plays role in peroxisome quality control by the selective
degradation of damaged peroxisomes, a process known as pexophagy (Marshall and
Vierstra, 2018). The accumulation of peroxisome-related transcripts/ proteins in *atg*
mutants has been reported lately, and it can represent both activation of peroxisomal
metabolism or accumulation of oxidized peroxisome (McLoughlin et al., 2018, 2020;
Have et al.,2019). Specifically, under carbon starvation conditions, the enrichment of
peroxisomal proteins was followed with an increase of glyoxysomal proteins
(McLoughlin et al., 2020). This response suggests that peroxisomes are most likely
converted to glyoxysomes to sustain high activation of FA ß-oxidation pathways during
starvation conditions (McLoughlin et al., 2020). The induction of peroxisomal pathways
reported here is positively correlated with the high levels of plastidial FFAs in *atg*
mutants, which can be interpreted as a response to control FFA levels concomitantly
with energy production. When taken together with our work , these studies suggest that
the primary metabolism of *atg* mutants is not only rearranged for amino acid catabolism
but is also possibly involved in supporting lipid oxidation pathways. In light of this,
further studies are clearly required (*i*) to investigate whether the disruption of LD
dynamics and differential membrane recycling of *atg* mutants impact the supply of lipid
substrates for FA oxidation pathways and (*ii*) to characterize the function of autophagy
under a range of environmental stresses during this process.

Conclusions

In the current work, we present compelling evidence regarding the importance of
autophagy to lipid homeostasis in Arabidopsis leaves. During carbon starvation, the
products of organellar membrane recycling are channeled to cytosolic LD production.
This enables the balance of FA production and turnover. The functional lack of
autophagy impacts proper membrane mobilization and activates a general chloroplast
lipid degradation program while failing to produce cytosolic LDs. Nonetheless, the
products of chloroplast membrane dismantling are possibly channeled to PG generation

Figure 9 - Schematic model of lipid remodeling response of *atg* mutants during extended darkness.
Under extended darkness, the plastidial and extraplastidial membranes, composed by MGDG/DGDG and phospholipids, respectively, are degraded, releasing FAs to be stored as TAGs in LD in the cytosol. LDs and transient FFA donate substrates for β-oxidation in peroxisomes. Meanwhile, in *atg* mutants, there is a substantial degradation of chloroplast lipids, while the production of cytosolic LDs is impaired. Part of the released plastidial FAs, together with phytol derived from chlorophyll degradation, are channeled to the production of PG in the chloroplast. β-oxidation pathways are highly induced to consume overloaded plastidial FFA and, possibly, PGs lipids (question mark). Abbreviations: Digalactosyldiacylglycerol (DGDG); Fatty acids (FA); Free Fatty Acids (FFA); Lipid Droplet (LD); Monogalactosyldiacylglycerol (MGDG).

in *atg* mutants chloroplasts (Fig. 9). We postulate that the increased activation of

peroxisomal lipid catabolic pathways is part of a complex metabolic response to

catabolize the FFAs released from chloroplast dismantling, and possibly, from PGs

mobilization.   Although our results highlight novel insights regarding the role of

autophagy in lipid metabolism, they certainly underestimate the complex crosstalk

between autophagy, peroxisome, mitochondria, and the endoplasmic reticulum . An

investigation of the interplay involving autophagy, membrane recycling, and organellar

maintenance is required to fully understand the implications of autophagy for the

regulation of lipid reserve dynamics and energy homeostasis. The combined results

presented here are of potential application for the breeding of crops with a higher yield,

increased resilience, and it seems reasonable to assume that there is a timely need to

accelerate our understanding of the mechanisms controlling autophagy and associated

processes in response to environmental conditions.

Methods

Plant Material and Dark Treatment

All *Arabidopsis thaliana* (L.) Heynh plants used in this study were of the Columbia ecotype (Col-0). The T-DNA mutant lines *atg9-4* (SALK_145980) (Shin et al., 2014), *atg5-1* (SAIL_129B079) (Yoshimoto et al., 2009), *atg7-2* (GK-655B06) (Hofius et al., 2009) were used in this study. Seeds were surface-sterilized and imbibed for 4 days at 4°C in the dark on 0.7% (w/v) agar plates containing half-strength Murashige and Skoog (MS) media (Sigma-Aldrich; pH 5.7). Seeds were subsequently germinated and grown at 22°C under short-day conditions (8 h light/16 h dark), 60% relative humidity with 150 μmol photons m^{-2} s^{-1}. For dark treatments and subsequent induction of dark-induced senescence, 10- to 14-day-old seedlings were transferred to soil and then grown at 22°C under short-day conditions (8 h light/16 h dark). Four-week-old plants were transferred to darkness in the same growth cabinet. Whole rosettes of two different plants representing an independent replicated by genotype were harvested at intervals of 0, 3, 6, 9 days after transition to darkness and immediately frozen in liquid nitrogen and stored at –80°C until further analysis.

Visualization and Analysis of LDs

To visualize LDs in Arabidopsis leaves, LDs were stained with BODIPY 493/503 (Invitrogen; from 4 mg/mL stock in DMSO). The sixth rosette leaf of 4-week-old plants was used for LD visualization and analysis. Leaf discs were stained with BODIPY solution (2μg/mL) in the dark and under vacuum for 10 min. Confocal images were acquired using the FV-1200 confocal microscope (Olympus, Japan) with a 60X/1.42 oil immersion objective. Excitation was performed at 488nm, and emission was collected between 500-540nm for BODIPY 493/503 and 650-750nm for the chlorophyll autofluorescence. The LD number was quantified using ImageJ (Schneider et al., 2012). Z-stack series of three images from four plants for each genotype were used for the quantification of LDs.

Lipid Profiling and Free Fatty Acid Analysis

Lipids and natural endogenic free fatty acid were extracted from six biological replicates using the MTBE method described by Salem et al. (2016). Vacuum-dried organic phases were processed using ultra-performance liquid chromatography (on a C8

reverse-phase column) coupled with Fourier transform mass spectrometry (exactive mass spectrometer; Thermo Fisher Scientific) in positive and negative ionization modes. The free fatty acid were analysed using using ultra-performance liquid chromatography (Waters, Acquity I Class System) coupled with mass spectrometer (Q-Exactive Thermo Fisher Scientific, H-ESI). Processing of chromatograms, peak detection, and integration were performed using REFINER MSH 10 (GeneData). Processing of mass spectrometry data included the removal of the fragmentation information, isotopic peaks, and chemical noise. Selected features were annotated using an in-house lipid database (Lapidot-Cohen et al., 2020). Five-six biological replicates were used for the analysis.

Expression analysis by RT-PCR

Total RNA was isolated using TRIzol reagent (Ambion, Life Technology) according to the manufacturer's recommendations. The total RNA was treated with TURBO DNA-free kit (Invitrogen). The integrity of the RNA was checked on 1% (w/v) agarose gels, and the concentration was measured using a Nanodrop spectrophotometer. Finally, 1 µg of total RNA was reverse transcribed with SensiFAST cDNA Synthesis Kit (Bioline) according to the manufacturer's recommendations. Real-time PCR reactions were performed in a 384-well microlitre using SensiFAST SYBR® No-ROX Kit (Bioline). The primers used here were designed using the open-source program QuantPrime-qPCR primer designed tool (Arvidsson et al., 2008) and are described in Table S2. The transcript abundance was calculated by the standard curves of each selected gene and normalized using the constitutively expressed genes ACTIN (AT2G37620). Three biological replicates were processed for each experimental condition.

Statistical analyses

The experiments were conducted in a completely randomized design with 3-6 replicates of each genotype. Data were statistically examined using analysis of variance and tested for significant ($P < 0.05$) differences using Student's t tests from algorithm embedded into Microsoft Excel. Principal component analysis (PCA) was performed to reduce the dataset and identify the variables that best explained the highest proportion of total variance using Metaboanalyst 4.0, an open source web-based tools for metabolomics data analysis (Chong et al., 2019).

Accession Numbers

The Arabidopsis Genome Initiative locus numbers for the major genes discussed in this

article are as follows: ATG5 (At5g17290), ATG7 (At5g45900), and ATG9

(At2g31260).

Supplemental Data

The following supplemental materials are available.

Supplemental Figure S1: Lipid response of *atg* mutants throughout dark-induced

senescence. Principal component analysis (PCA) was performed using the lipid data of

atg5-1, atg7-2, atg9-4 and WT genotypes after 0 (0d), 3 (3d), (6d) and 9 (9d) days of

dark treatment.

Supplemental Table S1: Summary of the main variables of the first three Principal

Component (PC) of *atg* mutants and WT genotypes after 0 and 6 days of dark

treatment

Supplemental Table S2: Primers used in the RT-PCR analyses performed in this study

Supplemental Table S3: Relative lipid content of leaves of Arabidopsis knockout

mutants *atg5-1, atg7-2, atg9-4* and WT after 0, 3, 6 and 9 days of dark treatment

Supplemental Table S4: Variables of the first three Principal Component (PC) of *atg*

mutants and WT genotypes after 0 and 6 days of dark treatment

ACKNOWLEDGEMENTS

We thank Prof. Oren Ostersetzer-Biran from The Hebrew University of Jerusalem for

providing the facilities allowing transcript analysis, as well as Corinne Best and Sofia

Shevtsov, for their assistance with qPCR analyses. We also thank Naomi Melamed-

Book from the microscopy core facility at The Hebrew University's Life Science core

facility for her technical support with confocal analysis.

FIGURE CAPTIONS

Figure 1 - Differential lipid response of *atg* mutants under dark-induced
senescence.
Principal component analysis (PCA) was performed using the lipid data obtained in
samples from 4-week-old Arabidopsis plants immediately before lights were turned off

(0 d) and after further treatment for 6d in darkness. Three loss-of-function mutants of
the autophagy pathway, namely *atg5-1, atg7-2, atg9-4* and their respective wild type
(WT) were analyzed.
699
Figure 2 - *atg* mutants present altered phospholipid levels under extended
darkness conditions
Hierarchical clustering of (A) Phosphatidylcholine (PC) and (B)
Phosphatidylethanolamine (PE). Values plotted are $\log_2$ fold change. Distances were
measured by Pearson correlation using MeV 4.9.0 software. Data were normalized to
the mean response calculated for the 0-d dark treated leaves of the wild type (WT).
Values presented are means ± SE of five-six biological replicates per genotype; an
asterisk (*) designates values that were determined by Student's t-test to be significantly
different ($P < 0.05$) from WT at each time point analyzed.
709
Figure 3 - Chloroplast lipids are substantially reduced in *atg* mutants during dark-
induced senescence
Monogalactosyldiacylglycerol (MGDG) and Digalactosyldiacylglycerol (DGDG)
values plotted are $\log_2$ fold change. Data were normalized to the mean response
calculated for the 0-d dark treated leaves of the wild type (WT). Values presented are
means ± SE of five-six biological replicates per genotype; an asterisk (*) designate
values that were determined by the Student's t-test to be significantly different ($P <$
0.05) from WT at each time point analyzed.
718
Figure 4 - Distinct accumulation of neutral lipids in *atg* mutants during dark-
induced senescence
(A) Triacylglycerol (TAG) and (B) Diacylglycerol (DAG). Values plotted are $\log_2$ fold
change. Data were normalized to the mean response calculated for the 0-d dark treated
leaves of the wild type (WT). Values presented are means ± SE of five-six biological
replicates per genotype; an asterisk (*) designates values that were determined by
Student's t-test to be significantly different ($P < 0.05$) from WT each time point
analyzed.
727
Figure 5 - Autophagy deficiency affects Lipid Droplet (LD) dynamics during dark-
induced senescence
(A) Representative confocal images of LDs (BODIPY 493/503 fluorescence, false color
green) in wild type (WT), *atg5-1,* and *atg7-2* leaves. Red color depicts chloroplast
autofluorescence. Images were collected at the same magnification, and are projections
of Z-stacks of 20 optical sections Bar = 10 μm (B) Number of total LDs per image area.
The number of LDs was quantified by ImageJ as BODIPY-stained lipid area in 2D
projections of Z-stacks of 3 images of four biological replicates per genotype and time
point.
737
Figure 6 - PG related genes are up-regulated in *atg* mutants during dark-induced
senescence
Transcript abundance is shown for genes associated with plastoglobuli (PG), including
PES-1 (A), PES-2 (B), ABC1K7 (C), PPH (D). The y-axis values represent the gene
expression level relative to the wild type (WT). Data were normalized with respect to
the mean response calculated for the 0-d dark-treated leaves of the WT. Values
presented are means ± SE of three independent biological replicates. An asterisk (*)
indicates values that were determined by Student's t-test to be significantly different (P

< 0.05) from the WT at each time point analyzed. A number sign (#) indicates values
determined by Student's t-test to be significantly different ($P < 0.05$) from WT at 0d.
748
Figure 7 - Changes in transcript levels in WT and *atg* mutants during dark-
induced senescence.
Transcript abundance is shown for genes associated with lipid catabolic pathways,
including ACX-4 (A), CSY-1(B) KAT-2 (C), MPF-2 (D), pMDH-1 (E), SDP-1 (F). The
y-axis values represent the expression level relative to the wild type (WT). Data were
normalized with respect to the mean response calculated for the 0-d dark-treated leaves
of the WT. Values presented are means ± SE of three independent biological replicates.
An asterisk (*) indicates values that were determined by Student's t-test to be
significantly different ($P < 0.05$) from the WT at each time point analyzed. A number
sign (#) indicates values determined by Student's t-test to be significantly different ($P <$
0.05) from WT at 0d.
760
Figure 8- Changes in free Fatty Acid levels in WT and *atg* mutant plants under
dark-induced senescence.
Values plotted are $\log_2$ fold change. Data were normalized to the mean response
calculated for the 0-d dark treated leaves of the wild-type (WT). Values presented are
means ± SE of five-six biological replicates per genotype; an asterisk (*) designate
values that were determined by the Student's t-test to be significantly different ($P <$
0.05) from WT each time point analyzed.
768
Figure 9 - Schematic model of lipid remodeling response of *atg* mutants during
extended darkness.
Under extended darkness, the plastidial and extraplastidial membranes, composed by
MGDG/DGDG and phospholipids, respectively, are degraded, releasing FAs to be
stored as TAGs in LD in the cytosol. LDs and transient FFA donate substrates for β-
oxidation in peroxisomes. Meanwhile, in *atg* mutants, there is a substantial degradation
of chloroplast lipids, while the production of cytosolic LDs is impaired. Part of the
released plastidial FAs, together with phytol derived from chlorophyll degradation, are
channeled to the production of PG in the chloroplast. β-oxidation pathways are highly
induced to consume overloaded plastidial FFA and, possibly, PGs lipids (question
mark). Abbreviations: Digalactosyldiacylglycerol (DGDG); Fatty acids (FA); Free
Fatty Acids (FFA); Lipid Droplet (LD); Monogalactosyldiacylglycerol (MGDG).
781
782

Parsed Citations

Araújo WL, Ishizaki K, Nunes-Nesi A, Larson TR, Tohge T, Krahnert I, Witt S, Obata T, Schauer N, Graham IA, Leaver CJ, Fernie AR (2010) Identification of the 2-hydroxyglutarate and isovaleryl-CoA dehydrogenases as alternative electron donors linking lysine catabolism to the electron transport chain of Arabidopsis mitochondria. Plant Cell 22: 1549-1563

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Araújo WL, Tohge T, Ishizaki K, Leaver CJ, Fernie AR (2011) Protein degradation: an alternative respiratory substrate for stressed plants. Trends Plant Sci 16: 489–498

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Avin-Wittenberg T, Bajdzienko K, Wittenberg G, Alseekh S, Tohge T, Bock R, Giavalisco P, Fernie AR (2015) Global analysis of the role of autophagy in cellular metabolism and energy homeostasis in Arabidopsis seedlings under carbon starvation. Plant Cell 27: 306–322

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Avin-Wittenberg T, Baluska F, Bozhkov PV, Elander PH, Fernie AR, Galili G, Hassan A, Hofius D, Isono E, Le Bars R et al (2018) Autophagy-related approaches for improving nutrient use efficiency and crop yield protection. J Exp Bot 69: 1335–1353

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Arvidsson S, Kwasniewski M, Riaño-Pachón DM, Mueller-Roeber B (2008) QuantPrime: a flexible tool for reliable high-throughput primer design for quantitative PCR. BMC Bioinformatics 9: 465

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Barros JAS, Cavalcanti JHF, Medeiros DB, Nunes-Nesi A, Avi-Wittenberg T, Fernie AR, Araújo WL (2017) Autophagy deficiency compromises alternative pathway 414 of respiration following energy deprivation. Plant Physiol 175: 62–76

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Barros JAS, Siqueira JAB, Cavalcanti JHF, Araújo WL, Avin-Wittenberg T (2020) Multifaceted Roles of Plant Autophagy in Lipid and Energy Metabolism. Trends Plant Sci. doi:10.1016/j.tplants.2020.05.004

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Besagni C, Kessler F (2013) A mechanism implicating plastoglobules in thylakoid disassembly during senescence and nitrogen starvation. Planta 237: 463-470

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Bhuiyan NH, Friso G, Rowland E, Majsec K, van Wijk KJ (2016) The plastoglobule-localized metallopeptidase PGM48 is a positive regulator of senescence in Arabidopsis thaliana. Plant Cell 28:3020–37

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Bréhélin C, Kessler F, van Wijk KJ (2007). Plastoglobules: Versatile lipoprotein particles in plastids. Trends Plant Sci. 12:260–266.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Chong J, Wishart DS, Xia J (2019) Using MetaboAnalyst 4.0 for Comprehensive and Integrative Metabolomics Data Analysis. Curr. Protoc. Bioinform. 68: 1–128

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Chung T, Phillips AR, Vierstra RD (2010) ATG8 lipidation and ATG8-mediated autophagy in Arabidopsis require ATG12 expressed from the differentially controlled ATG12a and ATG12b loci. Plant J. 62:483–93

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Couso I, Perez-Perez ME, Martinez-Force E, Kim HS, He Y, Umen JG, Crespo JL (2017) Autophagic flux is required for the synthesis of triacylglycerols and ribosomal protein turnover in Chlamydomonas. J Exp Bot 69:1355–1367.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Elander PH, Minina EA, Bozhkov PV (2018) Autophagy in turnover of lipid stores: trans-kingdom comparison. J. Exp. Bot. 69: 1301–1311.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Fan J, Yan C, Zhang X, Xu C (2013) Dual role for phospholipid: diacylglycerol acyltransferase: Enhancing fatty acid synthesis and diverting fatty acids from membrane lipids to triacylglycerol in Arabidopsis leaves. Plant Cell 25: 3506–3518.

Pubmed: Author and Title

Google Scholar: Author Only Title Only Author and Title

Fan J, Yu L, Xu C (2017) A central role for triacylglycerol in membrane lipid breakdown, fatty acid b-oxidation, and plant survival under extended darkness. Plant Physiol. 174: 1517–1530.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Fan J, Yu L, Xu CG (2019) Dual role of autophagy in lipid metabolism in Arabidopsis. Plant Cell 31: 1598–1613
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Gaude N, Bréhélin C, Tischendorf G, Kessler F, Dörmann P (2007) Nitrogen deficiency in Arabidopsis affects galactolipid composition and gene expression and results in accumulation of fatty acid phytyl esters. Plant J. 49: 729–739.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Havé M, Luo J, Tellier F, Balliau T, Cueff G, Chardon F, Zivy M, Rajjou L, Cacas JL, Masclaux-Daubresse C (2019) Proteomic and lipidomic analyses of the Arabidopsis atg5 autophagy mutant reveal major changes in endoplasmic reticulum and peroxisome metabolisms and in lipid composition. New Phytologist 223: 1461–1477.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Hirota T, Izumi M, Wada S, Makino A, Ishida H (2018) Vacuolar protein degradation via autophagy provides substrates to amino acid catabolic pathways as an adaptive response to sugar starvation in Arabidopsis thaliana. Plant Cell Physiol 59: 1363–458 1376
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Hofius D, Schultz-Larsen T, Joensen J, Tsitsigiannis DI, Petersen NH, Mattsson O, Jørgensen LB, Jones JD, Mundy J, Petersen M (2009) Autophagic components contribute to hypersensitive cell death in Arabidopsis. Cell 137: 773–783
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Hölzl G, Dörmann P (2019) Chloroplast lipids and their biosynthesis. Annu Rev Plant Biol 70: 51–81
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Hörtensteiner S, Kräutler B (2011) Chlorophyll breakdown in higher plants. Biochim. Biophys. Acta 1807: 977–988
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Ishida H, Yoshimoto K, Izumi M, Reisen D, Yano Y, Makino A, Ohsumi Y, Hanson MR, Mae T (2008) Mobilization of Rubisco and stroma-localized fluorescent proteins of chloroplasts to the vacuole by an ATG gene-dependent autophagic process. Plant Physiol 148: 142–155
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Ishizaki K, Larson TR, Schauer N, Fernie AR, Graham IA, Leaver CJ (2005) The critical role of Arabidopsis electron-transfer flavoprotein: Ubiquinone oxido reductase during dark-induced starvation. Plant Cell 17: 2587-2600
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Izumi M, Hidema J, Makino A, Ishida H (2013) Autophagy contributes to nighttime energy availability for growth in Arabidopsis. Plant Physiol 161: 1682-1693
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Izumi M, Ishida H, Nakamura S, Hidema J (2017) Entire photodamaged chloroplasts are transported to the central vacuole by Autophagy. Plant Cell 29: 377–394
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Kunz HH, Scharnewski M, Feussner K, Feussner I, Flügge UI, Fulda M, Gierth M (2009) The ABC transporter PXA1 and peroxisomal b-oxidation are vital for metabolism in mature leaves of Arabidopsis during extended darkness. Plant Cell 21: 2733–2749.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Lapidot-Cohen T, Rosental L, Brotman Y (2020) Liquid chromatography–mass spectrometry (LC-MS)-based analysis for lipophilic compound profiling in plants. Curr Protoc Plant Biol. 5 e20109. doi: 10.1002/cppb.20109
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Law SR, Chrobok D, Juvany M, Delhomme N, Lindén P, Brouwer B, Ahad A, Moritz T, Jansson S, Gardestrom P, Keech Olivier (2018) Darkened leaves use different metabolic strategies for senescence and survival. Plant Physiol 177: 132–150.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Li F, Chung T, Vierstra RD (2014) AUTOPHAGY-RELATED11 plays a critical role in general autophagy and senescence-induced mitophagy in Arabidopsis. Plant Cell 26: 788-807.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Li YH, Beisson F, Pollard M, Ohlrogge J (2006) Oil content of Arabidopsis seeds: The influence of seed anatomy, light and plant-to-plant variation. Phytochemistry 67: 904–915.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Lippold F, vom Dorp K, Abraham M, Hölzl G, Wewer V, Yilmaz JL, Lager I, Montandon C, Besagni C, Kessler F, Stymne S, Dörmanna P (2012) Fatty acid phytyl ester synthesis in chloroplasts of Arabidopsis. Plant Cell 24(5): 2001–14

Liu Y,Wang G,Wang X (2015) Role of aminoalcoholphosphotransferases 1 and 2 in phospholipid homeostasis in Arabidopsis. Plant Cell 27:1512–28

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Lundquist PK, Poliakov A, Bhuiyan NH, Zybailov B, Sun Q, van Wijk KJ (2012). The functional network of the Arabidopsis plastoglobule proteome based on quantitative proteomics and genome-wide coexpression analysis Plant Physiol.158: 1172-1192

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Marshall RS,Vierstra RD (2018) Autophagy: the master of bulk and selective recycling Annu. Rev. Plant Biol. 69: 173-208

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Masclaux-Daubresse C, d'Andrea S, Bouchez I, Cacas J-L (2020). Reserve lipids and plant autophagy. J Exp Bot. doi.org/10.1093/jxb/eraa082

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

McLoughlin F, Augustine RC, Marshall RS, Li F, Kirkpatrick LD, Otegui MS, Vierstra RD (2018) Maize multi-omics reveal roles for autophagic recycling in proteome remodelling and lipid turnover. Nature Plants 4: 1056–1070.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

McLoughlin F, Marshall RS, Ding X, Chatt EC, Kirkpatrick LD, Augustine RC, Li F, Otegui MS, Vierstra RD (2020) Autophagy Plays Prominent Roles in Amino Acid, Nucleotide, and Carbohydrate Metabolism During Fixed-carbon Starvation in Maize. Plant Cell. DOI: 10.1105/tpc.20.00226

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Michaeli S, Honig A, Levanony H, Peled-Zehavi H, Galili G (2014) Arabidopsis ATG8-INTERACTING PROTEIN1 is involved in autophagy-dependent vesicular trafficking of plastid proteins to the vacuole. Plant Cell 26: 4084–4101

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Minina EA, Moschou PN, Vetukuri RR, Sanchez-Vera V, Cardoso C, Liu Q, Elander PH, Dalman K, Beganovic M, Yilmaz JL, Marmon S, Shabala L, Suarez MF, Ljung K, Novák O, Shabala S, Stymne S, Hofius D, Bozhkov PV (2018) Transcriptional stimulation of rate-limiting components of the autophagic pathway improves plant fitness. J Exp Bot 69: 1415–1432.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Muthan B, Roston RL, Froehlich JE, Benning C (2013) Probing Arabidopsis chloroplast diacylglycerol pools by selectively targeting bacterial diacylglycerol kinase to suborganellar membranes. Plant Physiol. 163: 61–74.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Nakamura S, Hidema J, Sakamoto W, Ishida H, Izumi M (2018) Selective elimination of membrane-damaged chloroplasts via microautophagy. Plant Physiol. 177: 1007–1026.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Nguyen TB, Louie SM, Daniele JR, Tran Q, Dillin A, Zoncu R, Nomura DK, Olzmann, JA (2017) DGAT1-dependent lipid droplet biogenesis protects mitochondrial function during starvation-induced autophagy. Dev. Cell 42: 9–21

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Ohlrogge JB, Browse J, Somerville CR (1991) The genetics of plant lipids. Biochim. Biophy. Acta 1082: 1–26.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Phillips AR, Suttangkakul A, Vierstra RD (2008) The ATG12-conjugating enzyme ATG10 is essential for autophagic vesicle formation in

Arabidopsis thaliana. Genetics 178: 1339–1353

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Pracharoenwattana I, Cornah JE, Smith SM (2007) Arabidopsis peroxisomal malate dehydrogenase functions in b-oxidation but not in the glyoxylate cycle. Plant J 50: 381–390

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Pugkaew W, Meetam M, Ponpuak M, Yokthongwattana K, Pokethitiyook P (2017) Role of autophagy in triacylglycerol biosynthesis in Chlamydomonas reinhardtii revealed by chemical inducer and inhibitors. J. Appl. Phycol. 30: 15–22.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Pyc M, Cai Y, Greer MS, Yurchenko O, Chapman KD, Dyer JM, Mullen RT (2017) Turning over a new leaf in lipid droplet biology. Trends Plant Sci 22: 596–609

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Rambold AS, Cohen S, Lippincott-Schwartz J (2015) Fatty acid trafficking in starved cells: Regulation by lipid droplet lipolysis, autophagy, and mitochondrial fusion dynamics. Dev. Cell 32: 678–692.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Rottet S, Besagni C, Kessler F (2015) The role of plastoglobules in thylakoid lipid remodeling during plant development. Biochim. Biophys. Acta 1847: 889–899

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Salem MA, Juppner J, Bajdzienko K, Giavalisco P (2016) Protocol: a fast, comprehensive and reproducible one-step extraction method for the rapid preparation of polar and semi-polar metabolites, lipids, proteins, starch and cell wall polymers from a single sample. Plant Methods 12: 45.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image analysis. Nat. Methods 9: 671–675.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Schwarz V, Andosch A, Geretschläger A, Affenzeller M, Lütz-Meindl U (2017) Carbon starvation induces lipid degradation via autophagy in the model alga Micrasterias. J Plant Physiol 208:115–127.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Shin KD, Lee HN, Chung T (2014) A revised assay for monitoring autophagic flux in Arabidopsis thaliana reveals involvement of AUTOPHAGY-RELATED9 in autophagy. Mol Cells 37: 399–405

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Singh R, Kaushik S, Wang Y, Xiang Y, Novak I, Komatsu M, Tanaka K, Cuervo AM, Czaja MJ (2009) Autophagy regulates lipid metabolism. Nature 458: 1131-1135

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Soto-Burgos J, Zhuang X, Jiang L, Bassham DC (2018) Dynamics of autophagosome formation. Plant Physiol. 176: 219–229.

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Testerink C, Munnik T (2011) Molecular, cellular, and physiological responses to phosphatidic acid formation in plants. J. Exp. Bot. 62: 2349–2361

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Thompson AR, Doelling JH, Suttangkakul A, Vierstra RD (2005) Autophagic nutrient recycling in Arabidopsis directed by the ATG8 and ATG12 conjugation pathways. Plant Physiol 138: 2097-2110

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Tran QG, Yoon HR, Cho K, Lee SJ, Crespo JL, Ramanan R, Kim HS (2019) Dynamic Interactions between Autophagosomes and Lipid Droplets in Chlamydomonas reinhardtii. Cells: 8:992

Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Troncoso-Ponce MA, Cao X, Yang Z,Ohlrogge JB (2013) Lipid turnover during senescence. Plant Sci 205-206: 13–19

Pubmed: Author and Title

Google Scholar: Author Only Title Only Author and Title

van Wijk KJ, Kessler F (2017) Plastoglobuli: Plastid microcompartments with integrated functions in metabolism, plastid developmental transitions, and environmental adaptation. Annu Rev Plant Biol 68: 253–289
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Wada S, Ishida H, Izumi M, Yoshimoto K, Ohsumi Y, Mae T, Makino A (2009) Autophagy plays a role in chloroplast degradation during senescence in individually darkened leaves. Plant Physiol 149: 885–893
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Wang X, Devaiah SP, Zhang W, Welti R (2006) Signaling functions of phosphatidic acid. Prog Lipid Res. 5: 250–78
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Yang Y, Benning C (2018) Functions of triacylglycerols during plant development and stress. Curr. Opin. Biotechnol 49: 191–198
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Yoshimoto K, Hanaoka H, Sato S, Kato T, Tabata S, Noda T, Ohsumi Y (2004) Processing of ATG8s, ubiquitin-like proteins, and their deconjugation by ATG4s are essential for plant autophagy. Plant Cell.16: 2967–2983.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Yoshimoto K, Jikumaru Y, Kamiya Y, Kusano M, Consonni C, Panstruga R, Ohsumi Y, Shirasu K (2009) Autophagy negatively regulates cell death by controlling NPR1-dependent salicylic acid signaling during senescence and the innate immune response in Arabidopsis. Plant Cell 21: 2914–2927
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Ytterberg AJ, Peltier JB, van Wijk, KJ (2006). Protein profiling of plastoglobules in chloroplasts and chromoplasts. A surprising site for differential accumulation of metabolic enzymes. Plant Physiol. 140: 984–997.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title

Zhao L, Dai J, Wu Q (2014) Autophagy-like processes are involved in lipid droplet degradation in Auxenochlorella protothecoides during the heterotrophy–autotrophy transition. Front Plant Sci 5: 400.
Pubmed: Author and Title
Google Scholar: Author Only Title Only Author and Title