Reelin deficiency contributes to long-term behavioral abnormalities induced by chronic

adolescent exposure to Δ9-tetrahydrocannabinol in mice

Running Title: Influence of Reelin deficiency on THC adolescent exposure

Attilio Iemolo[1], Aisha Nur[2], Patricia Montilla-Perez[1], Victoria B Risbrough[3], Francesca Telese[1]

[1]Department of Medicine, University of California, San Diego, La Jolla, CA 92093, USA.

[2]Division of Biological Sciences, University of California, San Diego, La Jolla, CA 92093, USA.

[3]Department of Psychiatry, University of California, San Diego, La Jolla, CA 92093, USA;

Center for Excellence in Stress and Mental Health, Veterans Affairs San Diego Healthcare

System, San Diego, CA 92161, USA; Research Service, Veterans Affairs San Diego Healthcare

System, San Diego, CA 92161, USA.

*Correspondence should be addressed to ftelese@ucsd.edu

Number of pages: 35

Number of Figures: 12 figures, (1 table)

Number of Words: Abstract (244), Introduction (624), Discussion (1,497)

Conflict of Interest statement:

The authors declare no competing financial interests.

Acknowledgments

This work was supported by the National Institute of Drug Abuse [DP1DA042232 to FT].

We thank R.F. Hechinger and S. Roige-Sanchez for helpful discussions and critical reading of

the manuscript; A. Turner, H. Taylor, S. Nolan, A. Apelian, and J. Hightower for technical

assistance.

Abstract

Heavy and frequent use of cannabis during adolescence increases the risk of developing

psychiatric disorders. However, the neurobiological mechanisms underlying this vulnerability

remain largely unknown. Here, we explore whether adolescent vulnerability to long-term

behavioral effects of cannabis is modulated by *Reelin*, a gene implicated in the development of

the brain and of psychiatric disorders. To this aim, heterozygous Reeler (HR) mice, that express

reduced level of *Reelin*, were chronically exposed during adolescence to high doses (10mg/kg) of

Δ9-tetrahydrocannabinol (THC), a major psychoactive component of cannabis. Mice were tested

in early adulthood with multiple behavioral assays, including working memory, social

interaction, locomotor activity, anxiety-like responses, stress reactivity, and pre-pulse inhibition.

Compared to wild-type (WT), HR mice treated with THC showed, in a sex-specific manner,

impaired social behaviors, elevated disinhibitory phenotypes and increased responsiveness to

aversive situations. Independent of THC exposure, HR mice also spent more time exploring

unfamiliar objects, indicating that Reelin modulates novelty seeking behavior. To identify the

neuronal ensemble underlying this elevated novelty seeking in HR mice, we mapped the regional

brain expression of the immediate early gene, *Fos,* in mice exposed to novel objects. HR mice

exhibited reduced neuronal activation in the lateral septum, a subcortical brain structure

implicated in emotions, cognition and reward processes. Overall, these findings show that (1)

Reelin deficiency influences behavioral abnormalities caused by heavy consumption of THC

during adolescence, and (2) that Reelin plays a role in the neurobiological mechanisms

underlying disinhibitory behaviors, such as novelty seeking.

Significant Statement

The link between cannabis abuse and the development psychiatric disorders, especially in

adolescents, makes understanding the neurobiological mechanisms underlying cannabis effects

on the brain a significant biomedical problem. Reelin is a key signaling molecule in the

development of the adolescent brain and of psychiatric disorders, but its role in modulating the

behavioral changes induced by cannabis remain unknown. Here, we report an interaction

between *Reelin* deficiency and chronic adolescent exposure to THC, a major psychoactive

component of cannabis in mice, which led to cognitive deficits, disinhibitory behaviors and

altered emotional reactivity in a sex-specific manner. These experiments are the first to establish

a link between Reelin signaling and the endocannabinoid system targeted by THC.

Introduction

Heavy and frequent cannabis use by adolescents has been linked epidemiologically to increased

risk of developing psychiatric conditions, including schizophrenia, psychosis, and substance use

disorders (Volkow, 2016). Similarly, animal studies show that administration of synthetic

cannabinoids (e.g. THC) during adolescence perturbs a wide range of behaviors, including

memory, social interaction, anxiety, and sensorimotor gating by targeting the endocannabinoid

(eCB) system (Rubino et al., 2015). Despite the evidence of possible detrimental health

outcomes associated with adolescent cannabis use, a recent survey in the US revealed a

substantial increase in daily use of cannabis and a decreased perception of the risks associated

with its regular use by adolescents (Johnston, 2018). Further, the increasing legalization of

recreational cannabis use has led to calls to understand whether such policies put adolescents at

higher risk of developing psychiatric disorders. All this emphasizes the need to better understand

the neurobiological mechanisms associated with heavy consumption on cannabis during

adolescence (Wilkinson et al., 2016).

During adolescence, the brain undergoes continuous remodeling of its structure,

connectivity, and plasticity (Sturman and Moghaddam, 2011; Arain et al., 2013). In addition,

substantial hormonal changes during adolescence influence not only reproductive functions, but

also the emergence of sex differences in cognitive, social, and emotional behaviors (Schulz and

Sisk, 2016). Thus, the adolescence is considered a critical period wherein brain development

may be altered by the exposure to psychoactive drugs, which can lead to sex-specific behavioral

abnormalities and increased risk for psychopathology in adulthood (Cousijn et al., 2018; Lisdahl

et al., 2018).

However, the neurobiological mechanisms underlying the increased vulnerability of the

adolescent brain to the effects of THC remain poorly understood. The goal of this study is to

examine the potential role of Reelin signaling in modulating the behavioral effects of cannabis

on the adolescent brain.

Reelin is a protein of the extracellular matrix that is predominately expressed in neuronal

cells and plays a key role in brain development and synaptic plasticity (D'Arcangelo et al., 1995).

During embryonic stages, Reelin activates an extensive signaling cascade that is critical for the

proper migration and cell positioning of cortical neurons (Sekine et al., 2014). During

adolescence, Reelin signaling promotes the development of the synaptic excitation/inhibition

(E/I) balance within the prefrontal cortex (Iafrati et al., 2014; Bouamrane et al., 2016). In the

mature brain, Reelin is required for learning and memory by regulating the N-methyl-D-aspartate

receptors (NMDA-R) function and the expression of neuronal activity-dependent genes (Weeber

94 et al., 2002; Qiu et al., 2006; Niu et al., 2008; Rogers et al., 2011; Telese et al., 2015). In

95 humans, *Reelin* deficiency has been linked to the development of psychiatric disorders (Ishii et

96 al., 2016). Thus, the HR mice, that expressed lower level of Reelin, have been proposed as a

97 valid animal model to study neurodevelopmental psychiatric disorders (Lossi et al., 2019).

98 Whether there is a functional relationship between *Reelin* deficiency and the consequences of

99 early-life exposure to high levels of THC remains unknown.

100  To this aim, we examined the long-lasting behavioral outcomes of chronic adolescent

101 exposure to high doses of THC (10mg/kg) in female and male HR mice. We compared HR mice

102 to their WT littermate controls in a battery of behavioral tests exploring different facets of

103 cognitive and emotional responsiveness, including working memory (Sannino et al., 2012),

104 social interaction (Yang et al., 2011), anxiety-like responses (Bailey and Crawley, 2009), stress

105 reactivity (Can et al., 2012), and pre-pulse inhibition (Geyer et al., 2002). To further characterize

106 the neuronal ensembles underlying behavioral differences between WT and HR mice, we

107 examined the expression of the immediate early gene, *Fos,* in four brain regions that were

108 activated after exposure to unfamiliar objects. This is the first study to investigate the

109 relationship between *Reelin* deficiency and the effect of adolescent exposure to THC in mice.

111 **Materials and Methods**

113 **Animals.** All experimental procedures were approved by the institutional animal care and use

114 committee at University of California, San Diego. Mice were housed (3-4 per cage) under a 12h

115 light/12h dark cycle and provided with food and water ad libitum. HR mice were bred in house

116 using the B6C3Fe a/a-Relnrl/J line (The Jackson Laboratory, #000235), which carry a null

mutation in the Reelin gene (D'Arcangelo et al., 1995). HR express ~40% reduced level of

Reelin mRNA in brain tissues (**Suppl. Fig. 1A-B**).

Drug treatment protocol and experimental design for behavioral analysis. THC was

provided by the U.S. National Institute on Drug Abuse and was dissolved in a vehicle solution

consisting of ethanol, tween, and 0.9% saline (1:1:18) on the day of administration. Vehicle or

THC (10 mg/kg) were administered daily to adolescent mice by intraperitoneal injections from

post-natal day (PND) 28 to PND 48, which cover the adolescent period in mice (Laviola et al.,

2003) (**Suppl. Fig. 1C**). This dose was based on previous studies using low to high ranges of

THC in mice (Trexler et al., 2018; Kasten et al., 2019). Moreover, this dose used has been shown

to elevate plasma levels of THC in rodents (> 100 ng/mL) similar to those found in humans

smoking cannabis (Huestis, 2007; Zuurman et al., 2008; Nguyen et al., 2016). To examine the

long-term effects of chronic adolescent exposure to THC, mice were tested 15 days after a drug

abstinence period, starting at PND 63. We used 9 cohorts of mice to perform multiple behavioral

assays. Cohorts 1 to 5 were subjected to locomotor activity, open field (OF), six-different objects

(6-DOT), light-dark (LD), three-chamber social approach, and tail suspension (TS) tests. Cohorts

6 to 9 were subjected to locomotor activity, acoustic startle response (ASR) and pre-pulse

inhibition (PPI) tests. Mice were tested between 10:00 am and 5:00 pm. Behavioral assays were

conducted on separate days and all behavioral tests were performed once on each mouse. The

number of mice used in each behavioral assay is reported in Table 1. To prevent bias due to

olfactory cues, the behavioral apparatus was cleaned with diluted ethanol solution in between

mice.

Six different objects test. We used the 6-DOT to study short-term working memory in mice, as

previously described (Sannino et al., 2012). First, mice were left free to explore an empty arena

(60x40 x35 cm) during a 10 min habituation trial. Afterwards, mice were exposed to six different

objects for a total time of 10 min (familiarization trial). After an inter-trial interval of 1 minute,

mice were exposed to identical copies of the familiar objects, but one object was substituted with

a novel object (test trial). The exploration time across different trials was measured with

Anymaze and was used to calculate a discrimination index (DI) as follows: (time spent exploring

novel object – average time spent exploring familiar objects) / Total time spent exploring novel +

familiar objects. The time exploring the novel object and the discrimination index (DI) were used

as indexes of novelty-induced exploratory activity and working memory, respectively.

Three-chamber social approach test. The three-chamber social approach test was used to

measure social behaviors, as previously described (Yang et al., 2011). The apparatus comprised

of three-chambered box with dividing walls with small openings to allow free exploration of the

three chambers, such as one empty central chamber, one side chamber containing an empty small

wire cage (novel object) and one side chamber containing a stranger mouse inside a small wire

cage (novel mouse). The target mouse was first placed in the center chamber and allowed to

explore the apparatus for 15 min. After introducing the novel mouse and the novel object in the

side chambers, the mouse was allowed to explore for 10 min. The placement of the novel mouse

or novel object in the left or right chambers was systematically alternated in between trials. The

time spent in each compartment and the time spent actively sniffing the novel mouse or the novel

object were manually scored. Longer time spent with or exploring the novel mouse versus the

novel object was considered an index of sociability. Sociability index (SI) was calculated as

[(time spent exploring or sniffing novel mouse - time spent exploring or sniffing novel object) /

(total time spent exploring or sniffing novel mouse and novel object)].

Tail suspension test. The TS test was used to measure motor responses under aversive

conditions (Can et al., 2012). Mice were suspended by their tails with tape to a bar in a position

that they could not escape or hold on to nearby surfaces. The test lasted for 6 minutes and the

mobility time (sec) was manually scored for each minute. We analyzed was the mobility time

(sec) as the sum of the final 5 minutes.

Open field test. The OF test was used to measure anxiety-like behavior(Bailey and Crawley,

2009). Mice were placed randomly in one of the 4 corners of an open plexiglass arena (60x40x35

cm) for 5 minutes. The total time spent in and the latency to entry the center of the arena (s) were

recorded and scored using Anymaze.

Light-dark test. The LD test was used to measure anxiety-like behavior(Bailey and Crawley,

2009). The animals were tested for 10 min in a light–dark rectangular box (60x40x35 cm) in

which the aversive light compartment (40x40x35 cm) was illuminated by a 100 lux light. The

dark side (20x40x35 cm) had an opaque cover and ~ 0 lux of light. The two compartments were

connected by an open doorway, which allowed the subjects to move freely between the two

compartments. The test began by placing the animal in the dark compartment. The time spent in

the light compartment (s), the latency to enter in the light chamber (s), and the total number of

transitions, were measured using Anymaze.

Acoustic startle response and pre-pulse inhibition test. The ASR and PPI tests were used to assess stress reactivity and sensorimotor gating functions. The tests were performed with a startle reflex measuring apparatus (SR-LAB; San Diego Instruments, San Diego, CA), as previously described (Toth et al., 2013). The system comprises a piezoelectric unit that transduces vibrations into signals when mice startle inside the plexigas cylinder. First, mice are placed in a plexiglas cylinder with background noise (65 decibel [db]) for 5 min (acclimation phase). Then, mice were subjected to a total of 179 trials, for a total of 25 min, including: (a) startle trials (40 milliseconds [ms] with 80, 90, 100, 110 and 120 db acoustic pulses), (b) prepulse+startle trials: 20 ms with acoustic prepulses of 3 (68), 6 (71) and 12 (77) db above background noise followed, 100ms later, by a 40 ms 120-db startling pulse. Startle amplitude was measured every 1ms over a 65ms period beginning at the onset of the startle stimulus. Average startle amplitude over the sampling period was taken as the dependent variable. Percent PPI at each pre-pulse intensity was calculated as 100 - [(startle response for prepulse/startle response for startle-alone trials) x 100].

Locomotor activity. Locomotor activity was measured using the video tracking system Anymaze (Ugo Basile, Varese, Italy). Mice were placed in an empty open field (60x40x35 cm) for 20 minutes. The distance traveled (m) was recorded in 5 minutes intervals and used as an index of locomotor activity in a novel environment.

Body weight measurements. Body weight (g) was measured through the course of the drug administration protocol and at PND63 when the behavioral assays began. The change in body weight was calculated as the difference between body weight at any given day and body weight at PND 28.

Immunohistochemistry (IHC). To examine Fos expression by IHC in WT and HR mice, we

used $n = 6$ male WT, $n = 6$ male HR, $n = 4$ female WT, and $n = 9$ female HR littermates. Mice

were subjected to the 3 trials of the 6-DOT test, as described above. To confirm that the exposure

to the behavioral task triggers Fos expression, a separate group of 4 WT mice was used,

including $n = 1$ mouse for each trial and $n = 1$ mouse without any behavioral exposure (home

cage). Thirty minutes after the last trial, mice were anesthetized with isoflurane and fixed via

transcardial perfusion with 4% paraformaldehyde (PFA) in phosphate buffered saline (PBS).

Brain tissues were processed as previously described (Iemolo et al., 2020). Briefly, 30-µm thick

coronal slices were stained with a primary antibody recognizing Fos (1:1000, ABE457, EMD

Millipore, RRID:AB_2631318) and a secondary anti-rabbit antibody (donkey anti-rabbit Alexa

Fluor 488 (1:1000, A21206, Life Technologies). Images were acquired with a fluorescent

microscope (BZX800, Keyence Corporation, Osaka, Japan). Different brain areas were identified

using a mouse brain atlas as a reference (Paxinos G, 2001). For the medial prefrontal cortex

(mPFC), 5 sections from bregma + 2.46 to + 1.10 mm, ± 340 µm apart, were stained and 4

images for each section were captured for each mouse. For the CA1 and dentate gyrus (DG)

subregions of the hippocampus (HP), 4 sections from bregma -1.46 to - 2.46 mm, ± 250 µm

apart, were stained and 4 images for each section were captured for each animal. For the lateral

septum (LS), 1 section from bregma 1.42 mm to 1.10 mm was stained and 1 image for each

section was captured for each mouse. The number of Fos+ cells was counted by two examiners

blind to the experimental conditions. All numbers are presented as mean ± SEM.

RNA fluorescent in situ hybridization (FISH). To examine the expression of various

transcripts in specific cell types of the lateral septum, we performed RNA FISH using the

RNAscope Multiplex Fluorescent Reagent Kit v2 (ACD, #323100) and following the

instructions for 'fixed-frozen tissue sample' of the user manual (ACD, USM-323100). Tissue

sections corresponding to the LS (bregma 0.26-0.50 mm) from $n = 1$ mouse exposed to the 3

trials of the 6-DOT, as described above, were hybridized with a mix of three probes; Reelin

(ACD, # 405981) + Slc32a1 (ACD, #319191-C3) + Chat (ACD, # 408731) or Fos (ACD,

#316921) + Slc32a1 + Chat. To assess both tissue RNA integrity and assay procedure, a separate

group of sections were incubated with probes (data not shown). Images were acquired with the

Keyence fluorescent microscope.

Statistical analysis. To examine how factors (treatment, genotype, sex) affected mice behavior

or Fos expression, we used linear mixed models (LMM) in JMP pro v. 15.0 (SAS Institute, Inc).

LMM allow modeling of both fixed and random effects (which subsume repeated measures)

(Quinn GP, 2002; Zuur AF, 2016). We incorporated the categorical predictor variables of interest

as fixed effects and included all possible interactions. We included mouse cohorts and individual

subjects as random effects to account for possible non-independence of the data. We ensured

assumptions of approximate normality and variance homogeneity were met by inspecting plots

of residuals versus predicted values, and by inspecting quantile-quantile plots with 95%

confidence limit curves. When residual plots indicated that it was appropriate for repeated

measures, we used a covariance structure that allowed variances to differ across the levels of the

repeated variable (Garrett M. Fitzmaurice, 2011). We analyzed the untransformed data in all but

few traits where a fourth root transformation mitigated variance heterogeneity (e.g. latency to

light, latency to center, Vmax of acoustic startle, body weight change). When significant

interactions were found, planned post-hoc pairwise comparisons were performed to identify

differences among specific genotype and treatment groups. We report *P*-values that remained

significant after controlling for multiple comparisons by holding the 'false-discovery rate' to

0.05 using the Benjamini-Hochberg method (Hochberg, 1995). Outliers were detected using the

Huber M-estimation method (Huber, 1973) in JMP pro v. 15.0 and removed when appropriate (*n*

= 3 in latency to light, *n* = 2 latency to center, *n* = 2 time spent sniffing, *n* = 1 sociability index, *n*

= 1 in % PPI).

Factor analysis of behavioral assays. Factor analysis is a data reduction method for

understanding underlying relationships among variables (Bartholomew, 2008). We performed

factor analysis using the variables from multiple behavioral assays, such as 6-DOT (DI, total

exploration time in test trial [T3]) , social interaction (time sniffing novel mouse, difference time

sniffing novel mouse vs novel object), OF (time in center), LD (time in light), TS (mobility

time). Factor analysis was computed in JMP 15.0 Pro and was conducted with varimax rotation

with a factor-loading cutoff of 0.3 (Manly and Navarro Alberto; Stephens, 1996). The number of

factors retained in our model was selected by inspecting the 'elbow' on the scree plot curve with

factors retained if their eigenvalues were greater than 1 (Cattell, 1966). With these settings, a

three-factor model was generated for our dataset. The loadings of the observed variables on the

extracted factors are shown in Fig. 6. Factor scores for individual mice were extracted and used

as variables for subsequent LMM analysis to identify their associations with treatment, sex, and

genotype.

Results

Working memory was impaired after chronic adolescent exposure to THC, and in male HR

mice

To assess the long-term effects of chronic adolescent exposure to THC on working

memory in WT and HR mice, we used a modified novel object recognition test that uses 6

instead of 2 objects (6-DOT) (Sannino et al., 2012; Olivito et al., 2016). This test evaluates

recognition memory under conditions of high loads of information processing, which is referred

to as memory span and is considered a form of working memory. A discrimination index (DI)

was calculated to determine the amount of time mice explored the novel object compared to the

familiar objects. A preference for the novel object is considered as a sign that mice remember the

familiar objects (Sannino et al., 2012). Higher rates of exploratory activity in response to the

novel object is also interpreted as a sign of novelty seeking behavior (Flagel et al., 2014).

Chronic THC exposure during adolescence decreased the DI (working memory) of mice

across all groups by ~31% compared to the vehicle-treated control group (**Fig. 1A-B**, treatment

effect, $F_{1,86} = 16.4$, $P < 0.0001$). HR mice also showed decreased DI, but the magnitude of the

genotype effect varied among sexes (**Fig. 1C**, sex x genotype interaction, $F_{1,86} = 5.7$, $P = 0.019$).

Compared to WT, female HR mice did not show impaired working memory, but male HR mice

showed a decreased DI that was similar to the level of impairment caused by THC treatment in

male WT mice (**Fig. 1C,** paired t test, $t = 2.9$, $P_{adj} = 0.019$).

This assay also revealed that HR mice displayed a ~25% increase in novelty-induced

exploratory activity compared to WT controls (**Fig. 1D-E**, genotype effect, $F_{1,84} = 6.1$, $P =$

0.015). This effect was triggered by the novel object, as it was revealed during the test phase

while the total exploratory activity toward objects did not vary among groups in the initial

familiarization phase (**Fig. 1F**, main effects and interactions Ps > 0.3).

To further evaluate potential confounding effects of general locomotion or body weight

having an impact on exploratory activity in this task, we analyzed changes in spontaneous

locomotor activity in an empty arena or in body weight in WT and HR mice (**Suppl. Fig. 2A-B**).

There was no effect of THC treatment, genotype, or sex on locomotor activity or body weight at

the time of testing, suggesting that the deficits in working memory observed in WT and HR mice

were not driven by changes in locomotion or body weight.

Social behaviors were impaired by chronic adolescent exposure to THC in HR male mice

To assess the long-term effects of chronic adolescent exposure to THC on social

interaction behavior in WT and HR mice, we used the three-chamber social approach test in

which a mouse is given the choice to spend time interacting with a novel mouse or a novel

inanimate object, which are referred to as "exploration targets" in our analysis (Yang et al.,

2011). We considered two aspects of social interaction: (1) time spent in the chamber with the

novel mouse versus the novel object, and (2) time actively sniffing the novel mouse versus the

novel object, which is a more specific measure of social interaction (Yang et al., 2011). Higher

time spent in the chamber with or sniffing the novel mouse compared to the novel object was

considered an index of "sociability".

All mice explored the novel mouse more than the inanimate object, confirming that the

behavioral assay can detect sociability across all groups (**Fig. 2A-2D**, exploration target effect,

$F_{1,119} = 49.5$, $P < 0.0001$; $F_{1,113} = 83.3$, $P < 0.0001$, respectively). There were no significant

effects of treatment, genotype and sex on time spent in the chamber ($P > 0.1$). In contrast, when

analyzing time sniffing, the effect of chronic adolescent exposure to THC varied among the two

exploration targets (**Fig. 2E**, treatment x exploration target interaction, $F_{1,113} = 4.1$, $P = 0.046$).

Precisely, THC had no significant effect on time spent sniffing the novel inanimate object

compared to vehicle (**Fig. 2E**, paired *t* test, $t = 0.06$, $P_{adj} = 1$); however, THC caused a ~20%

decrease in time spent sniffing the novel mouse compared to vehicle across all groups (**Fig. 2E,** *t*

$= 3.1$, $P_{adj} = 0.003$). The suppressive effect of THC on social interaction depended on the

genotype (**Fig. 2F**, treatment x genotype interaction, $F_{1,108} = 6.1$, $P = 0.015$); a post-hoc analysis

revealing that only HR mice treated with THC exhibited a significant reduction in overall time

sniffing when compared to vehicle treated HR mice (paired *t* test, $t = 3.2$, $P_{adj} = 0.006$). However,

when a sociability index was calculated, we did not detect any effect of treatment or genotype

(all effects and interaction *Ps* >0.1, **Suppl. Fig. 3A-B**).

Chronic adolescent exposure to THC increased anxiety-like behavior in female HR mice

To evaluate how impaired Reelin signaling influences anxiety-like behavior in mice

chronically exposed to THC during adolescence, we compared WT and HR mice in the LD and

OF tests. These behavioral assays examine anxiety-like responses by which animals avoid

illuminated or open areas (Crawley and Goodwin, 1980; Bailey and Crawley, 2009). Decreased

anxiety (increased disinhibition) is represented by mice spending more time in the light or open

compartments, and by having a shorter latency to enter the brightly lit or center chambers.

In the LD test, THC increased the overall time spent in the light compartment (**Fig. 3A,**

treatment effect, $F_{1,102} = 8.5$, $P = 0.0043$), but the strength of these effects varied among

genotypes and sexes (**Fig. 3B**, treatment x genotype x sex interaction, $F_{1,97.5} = 4.4$, $P = 0.038$).

This effect was mainly driven by female HR mice treated with THC, which spent 43% longer

time in the light compared to the vehicle-treated group (**Fig. 3B,** paired *t* test, t = 3.6, $P_{adj} =$

0.004). A separate analysis revealed that the latency to enter the light chamber was also

influenced by genotype and sex (**Fig. 3D-E**, sex x genotype interaction, $F_{1,47} = 5$, $P = 0.0197$).

Female HR mice showed ~18% increase in latency compared to female WT, indicating an

increased baseline anxiety-like responses (paired t test, $t = 2.5$, $P_{adj} = 0.03$). However, the latency

to enter the light compartment was not significantly influenced by THC (treatment effect, $F_{1,97} =$

0.02, $P = 0.88$).

In the open field test, HR mice spent ~18% more time in the center (**Fig. 3F-G**, genotype

effect, $F_{1,102} = 3.7$, $P = 0.057$ marginally significant), suggesting a reduced anxiety-like behavior.

Time spent in center also varied among sexes and treatment groups (treatment x genotype x sex

interaction, $F_{1,99} = 6.1$, $P = 0.016$); however, despite this significant overall variability, a post-

hoc analysis did not identify significant differences between specific pairs of groups. Latency to

enter the open area did not vary among groups (**Fig. 3H**, all $P > 0.2$).

Male HR mice showed enhanced ability to strive against stress following chronic adolescent

exposure to THC

To test the effect of adolescent exposure to THC on reactivity to aversive conditions in

WT and HR mice, we performed the TS test, an assay commonly used to screen anti-depressive

drugs and to measure behavioral despair in mice (Can et al., 2012). An increased mobility has

been associated with enhanced ability to strive against stress and is broadly related to active

coping, impulsive and aggressive behaviors (Strekalova et al., 2004; Brockhurst et al., 2015).

We quantified total mobility time in a 6-minute tail suspension assay in WT and HR

mice. THC treatment influenced total mobility time, but the effect varied among sexes and

genotypes (**Fig. 4A**, treatment x sex x genotype interaction, $F_{1,158} = 12.8$, $P = 0.0005$). While

THC did not significantly change the mobility time of either male or female WT mice compared

to the vehicle controls (paired t test, $t = 1.9$, $P_{adj} = 0.07$; $t = 0.5$, $P_{adj} = 0.6$, respectively), it

induced a 19% increase in mobility time for male HR mice (**Fig. 4B**, paired t test, $t = 2.6$, $P_{adj} =$

0.04). We also observed sex differences in baseline mobility time for which female HR mice

showed a 21% increase in mobility time compared to male HR mice (**Fig. 4B,** paired t test, $t = 3$,

$P_{adj} = 0.03$).

Reelin deficiency led to enhanced startle responses and pre-pulse inhibition

To evaluate psychotic-like behaviors in WT and HR mice following chronic adolescent

exposure to THC, we quantified the ASR and the % PPI. The ASR is a reflexive reaction to a

sudden acoustic stimulus (Swerdlow et al., 2001; Geyer et al., 2002). The % PPI is used as a

measure of sensorimotor gating, which occurs when a weak pre-pulse stimulus suppresses the

response to the subsequent startling stimulus. PPI is impaired in schizophrenia patients and

serves as an animal model of schizophrenia (Cadenhead et al., 1993).

We first examined the differences in ASR of WT and HR mice at various acoustic

intensities (65, 80, 90, 100, 110, 120 db, **Fig. 5A**). THC did not influence ASR (treatment effect,

$F_{1,451} = 0.7$, $P = 0.4$). Compared to WT, HR mice exhibited an overall ~3% higher startle

response (genotype effect, $F_{1,451} = 7.2$, $P = 0.008$), which varied by the acoustic intensity of the

pulse (**Fig. 5B,** genotype x db interaction, $F_{5,408} = 2.8$, $P = 0.016$). The increased startle tended to

be stronger at 120db (**Fig. 5B,** paired t test, t = 2, $P_{adj} = 0.08$ marginally significant).

When examining the % PPI at three pre-pulse intensities (**Fig. 5C,** 68, 71 and 77 db), HR

mice showed an overall ~13% increase in % PPI compared to WT mice (**Fig. 5D**, genotype

effect, $F_{1,482} = 8.6$, $P = 0.004$). Adolescent exposure to THC also had an effect on % PPI that

varied among sexes (**Fig. 5E,** treatment x sex interaction, $F_{1,482} = 8$, $P = 0.005$). Female mice

treated with THC tended to show ~12 % reduction in % PPI compared to vehicle-treated mice

(paired *t* test, t = 2, P_{adj} = 0.05 marginally significant).

THC and Reelin deficiency influenced behavioral domains that were associated with

cognition and disinhibition

We performed a factor analysis to determine whether the multiple behavioral responses

observed in our study reflect a smaller number of synthetic behavioral domains that are

influenced by THC and *Reelin* deficiency.

The factor analysis identified three unique behavioral domains (factors), which explained

57% of total variance (**Fig. 6A**). The first factor explained 23.8% of the total variance and

contained outcomes (positive loadings) from the social interaction and working memory tests

(e.g. time sniffing novel mouse, discrimination index). As factor 1 largely reflected measures of

cognitive functions, we named it "*cognition*". The second factor (17% of the total variance) was

named "*disinhibition*" as it contained behaviors that reflected increased exploratory or

disinhibited behavior (e.g. time in the center of the open field, novel object exploration). The

third factor explained 15.7% of the total variance and was named "*stress reactivity*" because the

main outcome that contributed to this cluster was the mobility time of the TS test.

To determine whether these behavioral domains differed between treatments, genotypes,

or sexes, we analyzed the rotator factor scores for each subject (**Fig. 6B-C**). We found that the

behavioral domains were significantly influenced by treatment or genotype (treatment x

behavioral domains interaction, $F_{2,142}$ = 5.7, P = 0.004; genotype x behavioral domains

interaction, $F_{2,142}$ = 3.4, P = 0.035; respectively). Post-hoc analysis revealed that *cognition* was

significantly decreased by THC (**Fig. 6B,** paired *t* test, t = 3.1, P_{adj} = 0.006), and *disinhibition*

was significantly increased by *Reelin* deficiency (**Fig. 6C,** paired *t* test, t = 2.8, P_{adj} = 0.014).

Neither treatment, genotype, nor sex significantly influenced the third behavioral domain, *stress*

reactivity.

Overall, this exploratory data analysis is consistent with the results of the separate

behavioral tests by indicating that adolescent exposure to THC detrimentally affected cognitive

functions, and it revealed that reduced expression of *Reelin* led to disinhibitory behaviors, which

were intensified by THC treatment in a sex-specific manner.

Novelty seeking in HR mice was associated with reduced neuronal activity in the lateral

septum.

In response to different environmental stimuli, the expression of *Fos* is rapidly induced in

specific neuronal populations allowing the identification of relevant neuronal ensembles

encoding specific behaviors (Curran and Morgan, 1995; Chaudhuri, 1997). To explore the

neuronal ensemble underlying the enhanced novelty seeking behavior observed in the HR mice

(**Fig. 1E**), we mapped the regional brain expression of Fos protein by IHC in juvenile mice

subjected to the 6-DOT (**Fig. 1A**).

In response to this behavioral task, WT mice activated the expression of Fos in four brain

areas, including mPFC, LS, DG and CA1 sub-regions of the HP (**Suppl. Fig. 4**). Thus, we

selected these areas for measuring the number of Fos-expressing (Fos+) cells in WT and HR

mice, thirty minutes after the end of the task (**Fig. 7A-B**). There were no significant differences

in Fos expression between genotypes or sex in the mPFC, DG or CA1 (**Fig. 7F,H-I**). In contrast,

compared to WT, HR mice showed a ~28% reduction in number of Fos+ cells in the LS (**Fig.**

7G, genotype effect, $F_{1,13}$ = 5, P = 0.04). To examine the expression of *Reelin* and *Fos* in the LS,

we used RNA FISH and showed that *Reelin*-expressing and *Fos*-expressing cells are

predominately located in the LS caudal (LSc), with fewer positive cells present in the LS rostral

(LSr) (**Fig. 8A,C**). Consistent with the expression of *Reelin* by GABAergic neurons (Lossi et al.,

2019), *Reelin*+ cells also co-expressed the vesicular GABA transporter (*Scl32a1*), a marker of

GABAergic neurons (**Fig. 8B**). To further characterize the *Fos*-positive ensemble, we co-labeled

Fos+ cells with markers of GABAergic (*Scl32a1*) and cholinergic neurons (*Chat*), which

represent the major neuronal cell types of the LS. We found that *Fos* transcript was activated in

both cell types and that the location of *Fos*+ cells showed a preferential location in the LSc,

similarly to those expressing *Reelin* (**Fig. 8D**).

Discussion

The present study revealed for the first time that reduced levels of *Reelin* influences

behavioral abnormalities caused by heavy consumption of THC during adolescence, and that

Reelin plays a role in the neurobiological mechanisms underlying novelty seeking by modulating

neuronal ensembles in the LS.

Influence of THC and Reelin deficiency on cognitive functions

We found that chronic adolescent exposure to THC impaired working memory, in line

with previous reports in rodent models and clinical studies in humans (Renard et al., 2014;

Rubino and Parolaro, 2016; Hurd et al., 2019). Additionally, HR mice showed lower working

memory in comparison to the WT littermates and to a similar extent as did chronic adolescent

exposure to THC. Although it is the first time that HR mice are tested in a task that measures

memory span, such as the 6-DOT, this finding is consistent with studies demonstrating the

crucial role of Reelin in mechanisms underlying learning and memory (Weeber et al., 2002;

Rogers et al., 2013; Iafrati et al., 2014; Telese et al., 2015).

We found that social interaction was impaired by chronic adolescent exposure to THC

with the strongest effects in HR mice. These observations are in line with evidence in human and

preclinical studies showing social deficits induced by THC (Long et al., 2010). Only few studies

have examined social behaviors of the HR mice and did not report prominent social deficits

(Podhorna and Didriksen, 2004; Macri et al., 2010; Michetti et al., 2014). Thus, our study

revealed for the first time that *Reelin* deficiency exacerbates social deficits induced by THC.

Consistently, the factor analysis revealed that variables from both working memory and

social interaction tests contributed to a behavioral domain, named '*cognition*', which was

negatively influenced by adolescent exposure to THC.

Reelin deficiency is associated with disinhibitory phenotypes

We speculate that multiple behavioral responses exhibited by HR mice may reflect general

behavioral disinhibition. In rodents, behavioral disinhibition is measured as a function of

increased exploratory activity towards unfamiliar objects or environments, and has been

associated with compulsive drug taking (Davis et al., 2008; Flagel et al., 2010; Belin et al.,

2011). In humans, behavioral disinhibition reflects personality traits that encompasses

impulsivity, risk-taking and novelty seeking phenotypes; these behavioral patterns are more

pronounced in adolescence and have been linked to addiction susceptibility (Young et al., 2009).

Here, we found that, compared to WT, both male and female HR mice explored the novel objects

for longer time in the 6-DOT. Additionally, female HR mice treated with THC showed signs of

reduced inhibitory control as they spent more time in the center chamber of the OF test or in the

brightly lit compartment of the LD test. Consistently, variables from these tests contributed to the

same behavioral domain ('*disinhibition*') revealed by the factor analysis, which was positively

influenced by *Reelin* deficiency. These observations are also in line with a previous study

showing decreased anxiety of HR mice in the elevated plus maze and increased motor

impulsivity (Ognibene et al., 2007).

Collectively, these results raise the hypothesis that impairment in Reelin signaling may

underlie addiction traits by disrupting inhibitory control. Further supporting this hypothesis, in a

recent genome-wide association study (GWAS), a variant of the *Reelin* gene was associated with

higher likelihood to consume alcohol ($P = 4 \times 10^{-9}$, variant and risk allele rs756747-T) (Karlsson

Linner et al., 2019).

Reelin deficiency leads to abnormal responses to aversive conditions

We observed abnormal behavior in HR mice in response to aversive conditions. First,

male HR mice treated with THC showed prolonged mobility time in the TS test compared to

vehicle group. This behavior has been linked to proactive coping in response to a threat

(Strekalova et al., 2004; Brockhurst et al., 2015). Second, HR mice exhibited enhanced startle

reactivity to acoustic stimuli compared to WT mice, which was not influenced by THC

treatment. In a previous study, the startling response of HR mice did not change following a

single acoustic pulse of 105db; however, in our study, we used five startling pulses from 80db to

120db that likely increased the sensitivity of the task. Elevated ASR has been observed in mental

conditions associated with impaired emotional reactivity, such as posttraumatic stress disorders

(Orr et al., 1995) and obsessive-compulsive disorders (Kumari et al., 2001). These observations

led us speculate that reduced expression of *Reelin* may lead to altered emotional reactivity. This

hypothesis deserves further examination of the HR mice using tasks specifically designed to

assess impulsivity and aggression.

Reelin as a susceptibility factors for psychiatric disorders

The array of behavioral phenotypes exhibited by HR mice in our study, encompassing memory impairments, social deficits, poor inhibitory control, and altered stress responses, is reminiscent of the behaviors characterizing numerous psychiatric conditions, ranging from autism spectrum disorders (ASD) to schizophrenia and substance use disorders. Consistently, a role of Reelin in the development of these disorders is supported by several lines of evidence. In particular, whole exome sequencing studies identified *de novo* mutations in the *Reelin* gene in individuals with ASD (Neale et al., 2012; Iossifov et al., 2014; Wang et al., 2014). Further support for Reelin involvement in psychiatric disorders is provided by the observation of reduced expression of *Reelin* transcript or protein in postmortem brains of individuals affected by ASD (Fatemi et al., 2001), and schizophrenia (Guidotti et al., 2000; Ruzicka et al., 2007; Habl et al., 2012). These observations suggest that altered Reelin signaling may be a vulnerability factor for psychiatric disorders and that the HR mice represent a valuable animal model for translational research. However, we did not observe any PPI deficits in HR mice. In contrast, a previous study showed that a single *in vivo* injection of Reelin protein increases % PPI (Rogers et al., 2013). It is possible that, in our study, the elevated startle response observed in HR mice may act as a confounding factor for accurately assessing the effect of *Reelin* deficiency. It is also possible that impaired Reelin signaling contributes mainly to the cognitive, social and emotional deficits associated with schizophrenia, which are referred to as negative symptoms, as opposed to positive symptoms that include psychotic-like behaviors (Kirkpatrick et al., 2006).

Notably, THC reduced % PPI in female WT mice, suggesting that a history of drug exposure leads to psychotic-like behaviors. Conflicting results have been reported concerning the effects of adolescent chronic exposure to THC on psychotic-like behaviors in rodents (Rodriguez

531 et al., 2017; Todd et al., 2017; Ibarra-Lecue et al., 2018). Differences in THC doses, mice genetic

532 background and specific experimental conditions among different research groups may explain

533 some of these conflicts.

Sex differences in behavioral responses associated with Reelin haploinsufficiency

Our study revealed numerous sex differences in the behavioral abnormalities associated

with _Reelin_ deficiency and/or adolescent exposure to THC. Male HR mice were more sensitive

to working memory impairments, as well as to emotional reactivity in the TS test. In contrast,

female mice, showed reduced anxiety-like behaviors (HR) or reduced % PPI (WT) in response to

THC. Consistently, previous finding in rodents showed that chronic adolescent exposure to

cannabinoids is associated with numerous behavioral sex differences; however, the knowledge of

the underlying mechanisms remains limited (Craft et al., 2013). Given the critical role of Reelin

brain development, it is possible that Reelin influences the development of the endocannabinoid

system in a sex-specific manner, which can lead to sex differences to the behavioral effects of

THC.

The novelty seeking phenotype observed in HR mice is associated with reduced Fos expression
in the LS

We found that the prolonged exploratory activity of novel objects exhibited by HR mice

in the 6-DOT was associated with reduced neuronal activity in the LS. Although the LS has not

been linked directly to novelty seeking, it has been implicated in the modulation of rewarding

experiences that are correlated to novelty seeking (Luo et al., 2011; Harasta et al., 2015; McGlinchey

and Aston-Jones, 2018). Moreover, several behaviors influenced by _Reelin_ deficiency in our study

have also been linked to the activity of the LS, such as social interaction, memory and emotional

processes associated with aversive situations (Singewald et al., 2011; Leroy et al., 2018) (Shin et

al., 2018). Notably, *Fos* and *Reelin* expression are detected in the LSc, a region implicated in the

regulation of drug-seeking behavior via a population of GABAergic neurons that project to the

ventral tegmental area (VTA) (Luo et al., 2011). We speculate that *Reelin* deficiency may alter

LS-VTA pathway underlying motivated behavior by disrupting GABA inhibition in the LSc and,

hence, disinhibiting dopaminergic neurons in the VTA. In our previous work, we showed that

Reelin triggers a signaling cascade via the intracellular domain of its receptor LRP8 and directly

activates transcriptional enhancers of immediate early genes, including *Fos* (Telese et al., 2015).

Thus, it is possible that Reelin signaling modulates the transcriptional and epigenomic landscape

of specific neuronal populations in the LS.

In conclusion, our study suggests that elucidating Reelin signaling will improve our

understanding of neurobiological mechanisms underlying behavioral traits relevant to the

development of psychiatric conditions.

Acknowledgments

We thank R.F. Hechinger for commenting on text and analysis; S. Sanchez-Roige for helpful

discussions and critical reading of the manuscript; Patricia Montilla-Perez and Andrew Turner

for assistance in genotyping. H. Taylor for help with animal care; S. Nolan for assistance with

video scoring; Alik Apelian for technical support with the Keyence microscope; J. Hightower for

assistance with figure preparation. This work was supported by the National Institute of Drug

Abuse [DP1DA042232 to FT].

References

Arain M, Haque M, Johal L, Mathur P, Nel W, Rais A, Sandhu R, Sharma S (2013) Maturation of the
576   adolescent brain. Neuropsychiatr Dis Treat 9:449-461.
577 Bailey KR, Crawley JN (2009) Anxiety-Related Behaviors in Mice. In: Methods of Behavior Analysis in
578   Neuroscience (nd, Buccafusco JJ, eds). Boca Raton (FL).
579 Bartholomew DJ (2008) Analysis of multivariate social science data, 2nd Edition. Boca Raton: CRC Press.
580 Belin D, Berson N, Balado E, Piazza PV, Deroche-Gamonet V (2011) High-novelty-preference rats are
581   predisposed to compulsive cocaine self-administration. Neuropsychopharmacology 36:569-579.
582 Bouamrane L, Scheyer AF, Lassalle O, Iafrati J, Thomazeau A, Chavis P (2016) Reelin-Haploinsufficiency
583   Disrupts the Developmental Trajectory of the E/I Balance in the Prefrontal Cortex. Front Cell
584   Neurosci 10:308.
585 Brockhurst J, Cheleuitte-Nieves C, Buckmaster CL, Schatzberg AF, Lyons DM (2015) Stress inoculation
586   modeled in mice. Transl Psychiatry 5:e537.
587 Cadenhead KS, Geyer MA, Braff DL (1993) Impaired startle prepulse inhibition and habituation in
588   patients with schizotypal personality disorder. Am J Psychiatry 150:1862-1867.
589 Can A, Dao DT, Terrillion CE, Piantadosi SC, Bhat S, Gould TD (2012) The tail suspension test. J Vis
590   Exp:e3769.
591 Cattell RB (1966) The Scree Test For The Number Of Factors. Multivariate Behav Res 1:245-276.
592 Chaudhuri A (1997) Neural activity mapping with inducible transcription factors. Neuroreport 8:iii-vii.
593 Cousijn J, Luijten M, Feldstein Ewing SW (2018) Adolescent resilience to addiction: a social plasticity
594   hypothesis. Lancet Child Adolesc Health 2:69-78.
595 Craft RM, Marusich JA, Wiley JL (2013) Sex differences in cannabinoid pharmacology: a reflection of
596   differences in the endocannabinoid system? Life Sci 92:476-481.
597 Crawley J, Goodwin FK (1980) Preliminary report of a simple animal behavior model for the anxiolytic
598   effects of benzodiazepines. Pharmacol Biochem Behav 13:167-170.
599 Curran T, Morgan JI (1995) Fos: an immediate-early transcription factor in neurons. J Neurobiol 26:403-
600   412.
601 D'Arcangelo G, Miao GG, Chen SC, Soares HD, Morgan JI, Curran T (1995) A protein related to
602   extracellular matrix proteins deleted in the mouse mutant reeler. Nature 374:719-723.
603 Davis BA, Clinton SM, Akil H, Becker JB (2008) The effects of novelty-seeking phenotypes and sex
604   differences on acquisition of cocaine self-administration in selectively bred High-Responder and
605   Low-Responder rats. Pharmacol Biochem Behav 90:331-338.
606 Fatemi SH, Stary JM, Halt AR, Realmuto GR (2001) Dysregulation of Reelin and Bcl-2 proteins in autistic
607   cerebellum. J Autism Dev Disord 31:529-535.
608 Flagel SB, Waselus M, Clinton SM, Watson SJ, Akil H (2014) Antecedents and consequences of drug
609   abuse in rats selectively bred for high and low response to novelty. Neuropharmacology 76 Pt
610   B:425-436.
611 Flagel SB, Robinson TE, Clark JJ, Clinton SM, Watson SJ, Seeman P, Phillips PE, Akil H (2010) An animal
612   model of genetic vulnerability to behavioral disinhibition and responsiveness to reward-related
613   cues: implications for addiction. Neuropsychopharmacology 35:388-400.
614 Garrett M. Fitzmaurice NML, and James H. Ware (2011) Applied Longitudinal Analysis, 2nd Edition.
615 Geyer MA, McIlwain KL, Paylor R (2002) Mouse genetic models for prepulse inhibition: an early review.
616   Mol Psychiatry 7:1039-1053.
617 Guidotti A, Auta J, Davis JM, Di-Giorgi-Gerevini V, Dwivedi Y, Grayson DR, Impagnatiello F, Pandey G,
618   Pesold C, Sharma R, Uzunov D, Costa E (2000) Decrease in reelin and glutamic acid
619   decarboxylase67 (GAD67) expression in schizophrenia and bipolar disorder: a postmortem brain
620   study. Arch Gen Psychiatry 57:1061-1069.

Habl G, Schmitt A, Zink M, von Wilmsdorff M, Yeganeh-Doost P, Jatzko A, Schneider-Axmann T, Bauer M,
Falkai P (2012) Decreased reelin expression in the left prefrontal cortex (BA9) in chronic
schizophrenia patients. Neuropsychobiology 66:57-62.
Harasta AE, Power JM, von Jonquieres G, Karl T, Drucker DJ, Housley GD, Schneider M, Klugmann M
(2015) Septal Glucagon-Like Peptide 1 Receptor Expression Determines Suppression of Cocaine-
Induced Behavior. Neuropsychopharmacology 40:1969-1978.
Hochberg Ba (1995) Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple
Testing. Journal of the Royal Statistical Society Series B (Methodological) 57:289-300.
Huber PJ (1973) Robust Regression: Asymptotics, Conjecture, and Monte Carlo. Annals of Statistics
1:799–821.
Huestis MA (2007) Human cannabinoid pharmacokinetics. Chem Biodivers 4:1770-1804.
Hurd YL, Manzoni OJ, Pletnikov MV, Lee FS, Bhattacharyya S, Melis M (2019) Cannabis and the
Developing Brain: Insights into Its Long-Lasting Effects. J Neurosci 39:8250-8258.
Iafrati J, Orejarena MJ, Lassalle O, Bouamrane L, Gonzalez-Campo C, Chavis P (2014) Reelin, an
extracellular matrix protein linked to early onset psychiatric diseases, drives postnatal
development of the prefrontal cortex via GluN2B-NMDARs and the mTOR pathway. Mol
Psychiatry 19:417-426.
Ibarra-Lecue I, Mollinedo-Gajate I, Meana JJ, Callado LF, Diez-Alarcia R, Uriguen L (2018) Chronic
cannabis promotes pro-hallucinogenic signaling of 5-HT2A receptors through Akt/mTOR
pathway. Neuropsychopharmacology : official publication of the American College of
Neuropsychopharmacology 43:2028-2035.
Iemolo A, Montilla-Perez P, Lai IC, Meng Y, Nolan S, Wen J, Rusu I, Dulcis D, Telese F (2020) A cell type-
specific expression map of NCoR1 and SMRT transcriptional co-repressors in the mouse brain.
The Journal of comparative neurology.
Iossifov I et al. (2014) The contribution of de novo coding mutations to autism spectrum disorder.
Nature 515:216-221.
Ishii K, Kubo KI, Nakajima K (2016) Reelin and Neuropsychiatric Disorders. Front Cell Neurosci 10:229.
Johnston LD, Miech, R. A., O'Malley, P. M., Bachman, J. G., Schulenberg, J. E., & Patrick, M.E. (2018)
Monitoring the Future national survey results on drug use, 1975-2017: Overview, key findings on
adolescent drug use. . Ann Arbor: Institute for Social Research, The University of Michigan
Karlsson Linner R et al. (2019) Genome-wide association analyses of risk tolerance and risky behaviors in
over 1 million individuals identify hundreds of loci and shared genetic influences. Nat Genet
51:245-257.
Kasten CR, Zhang Y, Boehm SL, 2nd (2019) Acute Cannabinoids Produce Robust Anxiety-Like and
Locomotor Effects in Mice, but Long-Term Consequences Are Age- and Sex-Dependent. Front
Behav Neurosci 13:32.
Kirkpatrick B, Fenton WS, Carpenter WT, Jr., Marder SR (2006) The NIMH-MATRICS consensus statement
on negative symptoms. Schizophr Bull 32:214-219.
Kumari V, Kaviani H, Raven PW, Gray JA, Checkley SA (2001) Enhanced startle reactions to acoustic
stimuli in patients with obsessive-compulsive disorder. Am J Psychiatry 158:134-136.
Laviola G, Macri S, Morley-Fletcher S, Adriani W (2003) Risk-taking behavior in adolescent mice:
psychobiological determinants and early epigenetic influence. Neurosci Biobehav Rev 27:19-31.
Leroy F, Park J, Asok A, Brann DH, Meira T, Boyle LM, Buss EW, Kandel ER, Siegelbaum SA (2018) A circuit
from hippocampal CA2 to lateral septum disinhibits social aggression. Nature 564:213-218.
Lisdahl KM, Sher KJ, Conway KP, Gonzalez R, Feldstein Ewing SW, Nixon SJ, Tapert S, Bartsch H, Goldstein
RZ, Heitzeg M (2018) Adolescent brain cognitive development (ABCD) study: Overview of
substance use assessment methods. Dev Cogn Neurosci 32:80-96.

Long LE, Chesworth R, Huang XF, McGregor IS, Arnold JC, Karl T (2010) A behavioural comparison of
acute and chronic Delta9-tetrahydrocannabinol and cannabidiol in C57BL/6JArc mice. The
international journal of neuropsychopharmacology 13:861-876.
Lossi L, Castagna C, Granato A, Merighi A (2019) The Reeler Mouse: A Translational Model of Human
Neurological Conditions, or Simply a Good Tool for Better Understanding Neurodevelopment? J
Clin Med 8.
Luo AH, Tahsili-Fahadan P, Wise RA, Lupica CR, Aston-Jones G (2011) Linking context with reward: a
functional circuit from hippocampal CA3 to ventral tegmental area. Science 333:353-357.
Macri S, Biamonte F, Romano E, Marino R, Keller F, Laviola G (2010) Perseverative responding and
neuroanatomical alterations in adult heterozygous reeler mice are mitigated by neonatal
estrogen administration. Psychoneuroendocrinology 35:1374-1387.
Manly BFJ, Navarro Alberto JA Multivariate statistical methods : a primer, Fourth edition. Edition.
McGlinchey EM, Aston-Jones G (2018) Dorsal Hippocampus Drives Context-Induced Cocaine Seeking via
Inputs to Lateral Septum. Neuropsychopharmacology 43:987-1000.
Michetti C, Romano E, Altabella L, Caruso A, Castelluccio P, Bedse G, Gaetani S, Canese R, Laviola G,
Scattoni ML (2014) Mapping pathological phenotypes in reelin mutant mice. Front Pediatr 2:95.
Neale BM et al. (2012) Patterns and rates of exonic de novo mutations in autism spectrum disorders.
Nature 485:242-245.
Nguyen JD, Aarde SM, Vandewater SA, Grant Y, Stouffer DG, Parsons LH, Cole M, Taffe MA (2016)
Inhaled delivery of Delta(9)-tetrahydrocannabinol (THC) to rats by e-cigarette vapor technology.
Neuropharmacology 109:112-120.
Niu S, Yabut O, D'Arcangelo G (2008) The Reelin signaling pathway promotes dendritic spine
development in hippocampal neurons. J Neurosci 28:10339-10348.
Ognibene E, Adriani W, Granstrem O, Pieretti S, Laviola G (2007) Impulsivity-anxiety-related behavior
and profiles of morphine-induced analgesia in heterozygous reeler mice. Brain Res 1131:173-
180.
Olivito L, Saccone P, Perri V, Bachman JL, Fragapane P, Mele A, Huganir RL, De Leonibus E (2016)
Phosphorylation of the AMPA receptor GluA1 subunit regulates memory load capacity. Brain
Struct Funct 221:591-603.
Orr SP, Lasko NB, Shalev AY, Pitman RK (1995) Physiologic responses to loud tones in Vietnam veterans
with posttraumatic stress disorder. J Abnorm Psychol 104:75-82.
Paxinos G FK (2001) The Mouse Brain in Stereotaxic Coordinates, Second Edition. San Diego: Academic
Press.
Podhorna J, Didriksen M (2004) The heterozygous reeler mouse: behavioural phenotype. Behav Brain
Res 153:43-54.
Qiu S, Zhao LF, Korwek KM, Weeber EJ (2006) Differential reelin-induced enhancement of NMDA and
AMPA receptor activity in the adult hippocampus. J Neurosci 26:12943-12955.
Quinn GP KM (2002) Experimental design and data analysis for biologists. Cambridge University Press,
Cambridge, UK.
Renard J, Krebs MO, Le Pen G, Jay TM (2014) Long-term consequences of adolescent cannabinoid
exposure in adult psychopathology. Front Neurosci 8:361.
Rodriguez G, Neugebauer NM, Yao KL, Meltzer HY, Csernansky JG, Dong H (2017) Delta9-
tetrahydrocannabinol (Delta9-THC) administration after neonatal exposure to phencyclidine
potentiates schizophrenia-related behavioral phenotypes in mice. Pharmacology, biochemistry,
and behavior 159:6-11.
Rogers JT, Rusiana I, Trotter J, Zhao L, Donaldson E, Pak DT, Babus LW, Peters M, Banko JL, Chavis P,
Rebeck GW, Hoe HS, Weeber EJ (2011) Reelin supplementation enhances cognitive ability,
synaptic plasticity, and dendritic spine density. Learn Mem 18:558-564.

Rogers JT, Zhao L, Trotter JH, Rusiana I, Peters MM, Li Q, Donaldson E, Banko JL, Keenoy KE, Rebeck GW,
Hoe HS, D'Arcangelo G, Weeber EJ (2013) Reelin supplementation recovers sensorimotor gating,
synaptic plasticity and associative learning deficits in the heterozygous reeler mouse. J
Psychopharmacol 27:386-395.
Rubino T, Parolaro D (2016) The Impact of Exposure to Cannabinoids in Adolescence: Insights From
Animal Models. Biol Psychiatry 79:578-585.
Rubino T, Prini P, Piscitelli F, Zamberletti E, Trusel M, Melis M, Sagheddu C, Ligresti A, Tonini R, Di Marzo
V, Parolaro D (2015) Adolescent exposure to THC in female rats disrupts developmental changes
in the prefrontal cortex. Neurobiol Dis 73:60-69.
Ruzicka WB, Zhubi A, Veldic M, Grayson DR, Costa E, Guidotti A (2007) Selective epigenetic alteration of
layer I GABAergic neurons isolated from prefrontal cortex of schizophrenia patients using laser-
assisted microdissection. Mol Psychiatry 12:385-397.
Sannino S, Russo F, Torromino G, Pendolino V, Calabresi P, De Leonibus E (2012) Role of the dorsal
hippocampus in object memory load. Learn Mem 19:211-218.
Schulz KM, Sisk CL (2016) The organizing actions of adolescent gonadal steroid hormones on brain and
behavioral development. Neurosci Biobehav Rev 70:148-158.
Sekine K, Kubo K, Nakajima K (2014) How does Reelin control neuronal migration and layer formation in
the developing mammalian neocortex? Neurosci Res 86:50-58.
Shin S, Pribiag H, Lilascharoen V, Knowland D, Wang XY, Lim BK (2018) Drd3 Signaling in the Lateral
Septum Mediates Early Life Stress-Induced Social Dysfunction. Neuron 97:195-208 e196.
Singewald GM, Rjabokon A, Singewald N, Ebner K (2011) The modulatory role of the lateral septum on
neuroendocrine and behavioral stress responses. Neuropsychopharmacology 36:793-804.
Stephens D (1996) Hearing rehabilitation in a psychosocial framework. Scand Audiol Suppl 43:57-66.
Strekalova T, Spanagel R, Bartsch D, Henn FA, Gass P (2004) Stress-induced anhedonia in mice is
associated with deficits in forced swimming and exploration. Neuropsychopharmacology
29:2007-2017.
Sturman DA, Moghaddam B (2011) The neurobiology of adolescence: changes in brain architecture,
functional dynamics, and behavioral tendencies. Neurosci Biobehav Rev 35:1704-1712.
Swerdlow NR, Geyer MA, Braff DL (2001) Neural circuit regulation of prepulse inhibition of startle in the
rat: current knowledge and future challenges. Psychopharmacology (Berl) 156:194-215.
Telese F, Ma Q, Perez PM, Notani D, Oh S, Li W, Comoletti D, Ohgi KA, Taylor H, Rosenfeld MG (2015)
LRP8-Reelin-Regulated Neuronal Enhancer Signature Underlying Learning and Memory
Formation. Neuron 86:696-710.
Todd SM, Zhou C, Clarke DJ, Chohan TW, Bahceci D, Arnold JC (2017) Interactions between cannabidiol
and Delta(9)-THC following acute and repeated dosing: Rebound hyperactivity, sensorimotor
gating and epigenetic and neuroadaptive changes in the mesolimbic pathway. European
neuropsychopharmacology : the journal of the European College of Neuropsychopharmacology
27:132-145.
Toth M, Ziegler M, Sun P, Gresack J, Risbrough V (2013) Impaired conditioned fear response and startle
reactivity in epinephrine-deficient mice. Behav Pharmacol 24:1-9.
Trexler KR, Nass SR, Crowe MS, Gross JD, Jones MS, McKitrick AW, Siderovski DP, Kinsey SG (2018) Novel
behavioral assays of spontaneous and precipitated THC withdrawal in mice. Drug Alcohol
Depend 191:14-24.
Volkow ND (2016) Effects of Cannabis Use on Human Behavior-Reply. JAMA Psychiatry 73:996.
Wang Z, Hong Y, Zou L, Zhong R, Zhu B, Shen N, Chen W, Lou J, Ke J, Zhang T, Wang W, Miao X (2014)
Reelin gene variants and risk of autism spectrum disorders: an integrated meta-analysis. Am J
Med Genet B Neuropsychiatr Genet 165B:192-200.

Weeber EJ, Beffert U, Jones C, Christian JM, Forster E, Sweatt JD, Herz J (2002) Reelin and ApoE
receptors cooperate to enhance hippocampal synaptic plasticity and learning. J Biol Chem
277:39944-39952.
Wilkinson ST, Yarnell S, Radhakrishnan R, Ball SA, D'Souza DC (2016) Marijuana Legalization: Impact on
Physicians and Public Health. Annu Rev Med 67:453-466.
Yang M, Silverman JL, Crawley JN (2011) Automated three-chambered social approach task for mice.
Curr Protoc Neurosci Chapter 8:Unit 8 26.
Young SE, Friedman NP, Miyake A, Willcutt EG, Corley RP, Haberstick BC, Hewitt JK (2009) Behavioral
disinhibition: liability for externalizing spectrum disorders and its genetic and environmental
relation to response inhibition across adolescence. J Abnorm Psychol 118:117-130.
Zuur AF IE (2016) A protocol for conducting and presenting results of regression-type analyses. .
Methods in Ecology and Evolution 7:636-645.
Zuurman L, Roy C, Schoemaker RC, Hazekamp A, den Hartigh J, Bender JC, Verpoorte R, Pinquier JL,
Cohen AF, van Gerven JM (2008) Effect of intrapulmonary tetrahydrocannabinol administration
in humans. J Psychopharmacol 22:707-716.

Figure legends

Figure 1: Six-different objetcs test in WT and HR after chronic adolescent exposure to

THC. (A) 6-DOT with habituation, familiarization, and test trials. **(B)** Discrimination index (DI)

is shown as mean ± 95% confidence intervals. **(C)** THC reduced the DI (mean ± SEM) across all

groups (main treatment effect, $P < 0.0001$, LMM). **(D)** Male HR mice showed a reduced DI

(mean ± SEM) compared to male WT mice (sex x genotype interaction, $P = 0.02$, LMM). **(E)**

Exploration time (s) in test trial is expressed as mean ± SEM. **(F)** HR mice spent longer time

(mean ± SEM) exploring the objects in the test trial (main genotype effect, $P = 0.016$, LMM).

(G) Exploration time (mean ± SEM) in familiarization trial did not change across groups.

Adjusted $P < 0.05$ for planned post-hoc pairwise t tests are reported.

Figure 2: Social interaction behavior in WT and HR after chronic adolescent exposure to

THC. (A) Three-chamber interaction test diagram. Time (s) spent in chamber **(B)** or sniffing **(D)**

is expressed as mean ± 95% confidence intervals. **(C)** Mice spent more time (mean ± SEM) in

the chamber or **(D)** more time sniffing (mean ± SEM) the novel mouse compared to the novel

object (main exploration target effect, $P < 0.0001$, LMM). **(E)** THC reduced social investigation

(post-hoc pairwise t test $P_{adj} = 0.003$; treatment x exploration target interaction, $P = 0.05$, LMM).

(G) HR exposed to THC during adolescence showed reduced social interaction behavior (post-

hoc pairwise t test $P_{adj} = 0.006$; treatment x genotype interaction, $P = 0.015$, LMM).

Figure 3: Anxiety-like responses in WT and HR following chronic adolescent exposure to

THC. (A) Light-dark test schematic. **(B)** Time (s) spent in the light compartment is expressed as

mean ± SEM. **(C)** Female HR treated with THC during adolescence showed reduced anxiety

(treatment x genotype x sex interaction, $P = 0.04$, LMM). **(E)** Female HR mice showed increased

latency (s) to enter the light compared to WT controls (genotype x sex interaction, $P = 0.02$,

LMM). **(F)** Open field test schematic. **(G)** Time (s) spent in the center of the arena is expressed

as mean ± SEM. **(H)** HR mice showed reduced anxiety (main genotype effect, $P = 0.057$, LMM).

(I) Latency (s) to enter the center compartment is expressed as mean ± SEM and did not change

across groups. Adjusted $P < 0.05$ for planned post-hoc pairwise t tests are reported.

Figure 4: Reactivity to aversive conditions in WT and HR mice following chronic

adolescent exposure to THC. (A) Tail suspension test diagram. **(B)** Mobility time (s) is

expressed as mean ± SEM. **(C)** Male HR mice treated with THC during adolescence showed

increased mobility (treatment x genotype x sex interaction, $P = 0.0005$, LMM). Adjusted $P <$

0.05 for planned post-hoc pairwise t tests are reported.

Figure 5: Acoustic startle reflex of pre-pulse inhibition in WT and HR mice following

chronic adolescent exposure to THC. (A) Pre-pulse inhibition of acoustic startle test diagram**.**

(B) Startle amplitude is expressed as Vmax ± SEM. **(C)** HR mice show increased startle response

(genotype x db interaction, P = 0.016, LMM). **(D)** Pre-pulse inhibition is expressed as % PPI ±

SEM. **(E)** HM mice exhibited higher % PPI compared to WT controls (main genotype effect, P =

0.004, LMM). **(F)** Female treated with THC during adolescence showed reduced %PPI

compared to vehicle (treatment x sex interaction, P = 0.005, LMM). Adjusted $P < 0.05$ for

planned post-hoc pairwise t tests are reported.

Figure 6: Factor analysis of behavioral tests. (A) Factor loadings for each named behavioral

domain are reported for several behavioral variables. LMM analysis for rotated factor scores

revealed a **(B)** treatment x behavioral domain interaction (P = 0.04), and a **(C)** genotype x

behavioral domain interaction (P = 0.035). Adjusted $P < 0.05$ for planned post-hoc pairwise t

tests are reported.

Figure 7: Fos expression is reduced in the LS of HR mice in response to novel object

exploration. (A) Brain schematic illustrating the target areas for Fos quantification: mPFC, LS,

CA1, DG. **(B-E)** Schematic brain palette with bregma levels (top) and representative images

(bottom) of the mPFC, LS, CA1, and DG in WT (n = 6 male, n = 4 female) and HR mice (n = 6

male, n = 9 female). Scale bar, 50um.

Figure 8: Distribution of different cell types of the the LS. (A) RNA FISH showing cells

expressing *Reelin* and *Slc32a1* relative to Dapi-stained nuclei in the LS at bregma +0.50 mm. **(B)**

Higher magnification images illustrate the co-localization (white arrow) of *Reelin* and

*Slc32a1*mRNA. **(C)** RNA FISH showing cells expressing *Fos* and *Slc32a1* relative to Dapi-

stained nuclei in the LS at bregma +0.26 mm. **(D)** Higher magnification images illustrate the co-

localization of *Fos* and *Slc32a1* (white arrow) or *Fos* and *Chat)* mRNA. Dotted lines indicate the

caudal (LSc) and rostral (LSr) regions of the LS, and the fornix (fx) fibers. Scale bars are 100µm

in **(A)** and 20µm in **(B-C).**

Supplementary Figure 1: HR mice are used for chronic adolescent administration of THC.

(A) RT-PCR indicate Reelin mRNA expression in brain tissues isolated from WT and HR mice.

Level of Reelin expression are normalized with the housekeeping gene ACTB. **(B)** Reelin

mRNA is reduced in HR mice compared to WT littermates (genotype effect, $P = 0.0007$, LMM).

(C) Schematic diagram illustrates the daily i.p. administration of THC (10 mg/Kg) during

adolescence (PND28-48), and the day when behavioral assessment began (PDN63) after 15 days

of washout.

Supplementary Figure 2: Locomotor activity and body weight for WT and HR mice. (A)

Twenty-minute locomotor activity is expressed as mean distance traveled (m) ± SEM for every

5-minute intervals. **(B)** Body weight (g) change is expressed as mean ± SEM for the THC

administration course (day 1 to 14), and for day 63 when the behavioral testing started.

Supplementary Figure 3: Sociability index in WT and HR mice following chronic

adolescent exposure to THC. (A) Sociability index calculated using time spent in the chamber

with novel mouse versus novel object is shown as mean ± 95% confidence intervals. **(B)**

Sociability index calculated using time sniffing novel mouse versus novel object is shown as

mean $\pm$ 95% confidence intervals. Treatment, genotype and sex had no significant effect on both

sociability indices.

Supplementary Figure 4: Fos activation in response to novelty. **(A)** Schematic of behavioral

manipulation, including home cage (control), habituation, familiarization, and test phases of the

6-DOT. **(B-C)** Representative images of Fos expression by IHC in different brain regions,

including LS, mPFC, CA1 and DG. Dotted lines illustrate the line separating the LS from the

striatum (St) and the lateral ventricle (LV). Scale bars are 100μm in **(B, E)** and 50μm in **(C, D).**

Table 1. Number of mice used for behavioral analysis*

BEHAVIORAL ASSAYS	MALE WT		FEMALE WT		MALE HET		FEMALE HET	
	VEH	THC	VEH	THC	VEH	THC	VEH	THC
Locomotor	19	20	19	18	17	21	19	23
Tail suspension	24	24	20	18	18	22	20	22
Open field	18	16	13	12	10	11	16	14
Light-dark	18	16	13	12	10	11	16	14
Social	19	20	17	15	10	11	16	14
6-DOT	12	13	9	9	10	11	16	14
PPI (no MK-801)	23	24	24	23	20	22	25	26
ASR	23	24	25	23	20	22	25	26

872
873   * The number of outliers for specific variables and removed from analysis of are reported in the
874   method section
875

A

Three-Chamber Approach Test

Novel mouse

Novel object

B

Genotype

WT   HR

Time Spent in Chamber (sec)

Female

Male

Novel mouse
Novel object

Veh THC   Veh THC
Treatment

C

Exploration Target
Effect (*P* < 0.0001)

Time Spent in Chamber (sec)

Novel object   Novel mouse
Exploration Target

D

Genotype

WT   HR

Time Sniffing (sec)

Novel mouse
Novel object

Female

Male

Veh THC   Veh THC
Treatment

E

Exploration Target Effect
(*P* = 0.0001)

Time Sniffing (sec)

Novel object   Novel mouse
Exploration Target

F

Treatment x Exploration
Target (*P* = 0.05)

Time Sniffing (sec)

Novel object
Novel mouse

Vehicle vs THC
● *P*adj = 0.003

Veh   THC
Treatment

G

Treatment x Genotype
Interaction (*P* = 0.015)

Time Sniffing (sec)

WT
HR

Vehicle vs THC
● *P*adj = 0.003

Veh   THC
Treatment

A Tail suspension test

B Genotype

WT   HR

Female

Male

Sex

Mobility Time (sec)

Veh   THC   Veh   THC

Treatment

C Treatment x Genotype x Sex Interaction ($P = 0.0005$)

Female

Male

Sex

Mobility Time (sec)

○ WT   ● HR

● HR

Vehicle vs THC
$P_{adj} = 0.04$

Veh   THC

Treatment

A Pre-pulse Inhibition of Acoustic Startle Response

A

	Cognition	Stress reactivity	Disinhibition
T sniffing mouse			
T in light			
T in center			
Mobility T			
ExplorationT3			
Difference sniffing			
DI			

Factor Loadings

B Treatment x Behavioral Domain Interaction (*P* = 0.004)

○ Veh
● THC

Rotated Factors Scores

Veh vs THC P$_{adj}$ = 0.006

Cognition   Stress reactivity   Disinhibition

Behavioral Domain

C Genotype x Behavioral Domain Interaction (*P* = 0.035)

○ WT
● HR

Rotated Factors Scores

WT vs HRM P$_{adj}$ = 0.014

Cognition   Stress reactivity   Disinhibition

Behavioral Domain

A Brain areas of c-Fos activation

mPFC   CA1

LS   DG

B mPFC   **C** LS   **D** CA1   **E** DG

bregma (+2.46 ~ +1.10)   bregma (+1.42 ~ +1.10)   bregma (−1.46 ~ −2.46)   bregma (−1.46 ~ −2.46)

WT

50 µm

HRM

50 µm

F Genotype   **G** Genotype   **H** Genotype   **I** Genotype

Female / Male

Sex

cFos+ cells/mm²

WT   HR

Genotype effect $P = 0.04$

A Genotype

B Genotype effect
(*P* = 0.0007)

C

THC (10 mg/kg) or vehicle (i.p.)
21 days

15 days washout

Behavioral battery

PND28

PND48

PND63

A

Home cage | Habituation | Familiarization | Test

B

St   LV   LS

100 µm

C mPFC

50 µm

D CA1

50 µm

E DG

100 µm