Title:

Phosphatidylserine prevents the generation of a protein-free giant plasma membrane domain

in yeast

Running title:

Protein-free PM domains in yeast

Authors:

Tetsuo Mioka[1*], Guo Tian[1], Wang Shiyao[1], Takuma Tsuji[2], Takuma Kishimoto[1], Toyoshi

Fujimoto[2], and Kazuma Tanaka[1*]

Author affiliations:

[1] Division of Molecular Interaction, Institute for Genetic Medicine, Hokkaido University

Graduate School of Life Science, Sapporo, Hokkaido, Japan

[2] Laboratory of Molecular Cell Biology, Research Institute for Diseases of Old Age, Juntendo

University Graduate School of Medicine, Tokyo, Japan.

*Corresponding author: Tetsuo Mioka and Kazuma Tanaka

E-mail: mioka@igm.hokudai.ac.jp (TM) and k-tanaka@igm.hokudai.ac.jp (KT)

Summary statement

Yeast cells lacking phosphatidylserine generate protein-free plasma membrane domains, and vacuoles contact with this domain. This is the first report of micron-scale plasma membrane domains in living cells.

Abstract

Membrane phase separation accompanied with micron-scale domains of lipids and proteins occurs in artificial membranes; however, a similar large phase separation has not been reported in the plasma membrane of the living cells. We demonstrate here that a stable micron-scale protein-free region is generated in the plasma membrane of the yeast mutants lacking phosphatidylserine. We named this region the "void zone". Transmembrane proteins, peripheral membrane proteins, and certain phospholipids are excluded from the void zone. The void zone is rich in ergosterol and requires ergosterol and sphingolipids for its formation. These characteristics of the void zone are similar to the properties of the cholesterol-enriched domain in phase-separated artificial membranes. We propose that phosphatidylserine prevents the formation of the void zone by preferentially interacting with ergosterol. We also found that void zones were frequently in contact with vacuoles, in which a membrane domain was also formed at the contact site.

Introduction

The fluid mosaic model describing the dynamic distribution of proteins at the plasma membrane has been largely modified and developed to date (Singer and Nicolson, 1974; Nicolson, 2014; Kusumi A *et al.,* 2012). Lateral diffusion of proteins is not free and is influenced by protein interaction with other plasma membrane proteins and cytoskeletal

elements. In cholesterol-rich domains, such as lipid rafts, certain proteins can accumulate due

to protein-protein or protein-lipid interactions (Lingwood and Simons, 2010). The plasma

membrane is currently considered to be a nanoscale heterogeneous structure. In addition,

several macroscopic diffusion barriers have been detected, and some of the barriers represent

membrane compartmentalization due to interactions between the cytoskeleton and membrane

proteins (Kusumi *et al.,* 2012; Trimble and Grinstein, 2015). In artificial membranes, such as

giant unilamellar vesicles (GUVs) and giant plasma membrane vesicles (GPMVs), membrane

phase separation leads to the formation of even larger domains of proteins and lipids (Veatch

and Keller, 2003; Baumgart *et al.,* 2007; Elson *et al.,* 2010; Carquin *et al.,* 2016). Phase

separation in artificial membranes has been well studied and is often compared to the

nanoscale membrane domains found in the cells; however, large-scale phase separation is not

observed in the plasma membranes of the living cells due to unknown reasons.

The plasma membranes are composed of diverse lipid species, and the role of

phosphatidylserine (PS) and phosphatidylinositol phosphates (PIPs) in various cellular

functions has been studied (Uchida *et al.,* 2011; Cho *et al.,* 2015; Middel *et al.,* 2016;

Tsuchiya *et al.,* 2018; Michell, 2008; Balla, 2013). However, it is poorly understood how

individual phospholipids influence the membrane environment.

Although PS is essential for growth of mammalian cells, yeast mutant cells lacking *CHO1*,

the only PS synthase in the budding yeast, can grow (Arikketh *et al.,* 2008; Atkinson *et al.,*

1980). To explore a new role for PS, we have analysed PS-deficient *cho1Δ* yeast cells. In this

study, we show that stable large protein-free membrane domains are detected in the plasma

membrane of PS-deficient *cho1Δ* cells, which we named the "void zone". Transmembrane

proteins, peripheral membrane proteins, and certain phospholipids are excluded from the void

zone. This property is very similar to the cholesterol-enriched membrane domain in phase-

separated artificial membranes. Our results suggest that PS suppresses the development of

large-scale phase separation in the plasma membrane of the living cells and consequently

ensures the distribution of proteins and lipids throughout the plasma membrane. Furthermore,

we found that vacuoles, the lysosomal organelle of yeast, contact with the void zone on the

plasma membrane.

Results

PS-deficient cells show a protein-free region, "void zone", in the plasma membrane.

GFP-Snc1-pm, a mutant of v-SNARE Snc1, is uniformly distributed throughout the plasma

membrane due to a defect in its endocytosis (Lewis *et al.* 2000). In PS-deficient *cho1Δ* cells

grown at 37°C, GFP-Snc1-pm was heterogeneously distributed on the plasma membrane

(Figure 1A). A GFP-Snc1-pm-deficient region was barely detectable at 30°C, was frequently

present during incubation at 37°C for over 6 hours, and was not detected after heat shock at

42°C for 20 min (Figure 1B). This Snc1-pm-deficient region of the plasma membrane is

referred to as "void zone" in the present study. The shape of the void zone observed on the

cell surface was irregular and did not correspond to a smooth circle, and some cells had

multiple void zones (Figure 1C). When *cho1Δ* cells were observed immediately after staining

with FM4-64 lipophilic dye, FM4-64 was distributed throughout the plasma membrane

including the void zone (Figure 1D), suggesting that the plasma membrane is not lost or

significantly damaged in cells harbouring the void zone. To examine whether the void zone

influences the distribution of other transmembrane proteins, four different transmembrane

proteins, Pma1, Pdr5, Pdr12, and Sfk1, were compared with Snc1-pm (Figure 1E). Pma1 is

the major plasma membrane H^+-ATPase (Serrano *et al.,* 1986). Pdr5 and Pdr12 are the ATP-

binding cassette (ABC) transporters involved in the multidrug resistance and the weak organic

acid resistance, respectively (Bauer *et al.,* 1999). Sfk1 regulates phospholipid asymmetry in

conjunction with the flippase complex Lem3-Dnf1/2 and is involved in proper localization of

a phosphatidylinositol-4-kinase Stt4 (Audhya and Emr, 2002; Mioka *et al.,* 2018). The results

indicate that all these proteins showed void zones in the same region with void zones detected

by Snc1-pm; the percentage of overlapping void zones was more than 82% in all cases

(Figure 1E; percentage was determined from 70-110 cells with void zones). These results

suggest that the void zone is a membrane protein-free region common to all other membrane

proteins (see below), and the void zone can represent an abnormal lipid domain that inhibits

the lateral movement of the transmembrane proteins into the domain.

We also investigated the distribution of eisosomes, large immobile protein complexes that

form furrow-like invaginations in the fungal plasma membrane (Douglas and Konopka,

2014). Eisosome components, Pil1 and Sur7, were not distributed in the void zone (Figure

S1). The void zone was also devoid of the eisosome structure.

To further examine whether transmembrane proteins are completely absent in the void zone,

electron microscopy combined with the freeze-fracture replica method was applied (Fujita *et*

al., 2010; Tsuji *et al.,* 2017). In this method, transmembrane proteins are detected as the

granular structures called intramembrane particles (IMPs) (Figure 1F). In the protoplasmic

face of the plasma membrane in *cho1Δ* cells, most of the regions were IMP-rich although a

submicron-sized smooth region without IMPs was detectable (Figure 1G). To examine

whether this IMP-deficient area is the void zone, we labelled Pma1-GFP with an anti-GFP

antibody and colloidal gold-conjugated protein A (Figure 1F). As expected, only the IMP-rich

area was stained for Pma1-GFP and there was no labelling in the IMP-deficient area (Figure

1H). This result is consistent with the fluorescence microscopy images (Figure 1E) and

indicates that the void zone corresponds to the IMP-deficient area. Thus, there are no

transmembrane proteins in the void zone.

Figure 1

Figure 1 A protein-free region, void zone, is present in the plasma membrane of the PS-

deficient yeast cells

(A) Representative images of the void zone. Cells expressing GFP-Snc1-pm were grown

overnight in YPDA medium at 37°C. Arrowheads indicate the void zone. Scale bar: 5 μm. (B)

The percentage of cells with the void zone. *cho1Δ* cells expressing GFP-Snc1-pm were grown

under the indicated conditions to 0.8-1.5 OD_{600}. The incidence of the void zone was examined

(n > 100 cells, five independent experiments) and is shown as a box plot. Asterisks indicate

significant differences from the data obtained at 30°C according to the Tukey–Kramer test

(***, p < 0.001; n.d., not detected). (C) Representative images of the void zone in the cell

surface. Cells were prepared as in (A). Scale bar: 5 μm. (D) The lipophilic dye FM4-64 can be

distributed in the void zone. *cho1*Δ cells were prepared as in (A) and stained with FM4-64

just before the observation. Fluorescence intensities of GFP-Snc1-pm and FM4-64 around the

cell (arrows) are plotted on the right. Arrowheads indicate the void zone. Scale bar: 5 μm. (E)

The void zone is common to various transmembrane proteins. Cells expressing the indicated

proteins were prepared as in (A). Arrowheads indicate the void zone. Fluorescence intensities

around the cell (arrows) were plotted as in (D). Scale bar: 5 μm. (F) A scheme of freeze-

fracture replica labelling method. (G, H) Freeze-fracture EM images of the plasma membrane

of *cho1*Δ cells (G) or Pma1-GFP-expressing *cho1*Δ/P_{GAL1}-*CHO1* diploid cells (H). Cells were

grown at 37°C. The enlarged image of the area indicated by the square is shown on the right

(G) or at the bottom (H). Colloidal gold particles indicate Pma1-GFP labelled by anti-GFP

antibody (H). Scale bars: 0.2 μm. (IMP, intramembrane particles; Es, eisosome)

Characterization of the dynamics of the void zone.

We next examined the stability of the void zone by time-lapse imaging. Most of the void

zones were detected over tens of minutes, and some void zones were detectable for longer

period of time over 90 min. Bud growth and cytokinesis appeared to proceed normally in the

void zone-containing cells, suggesting that the void zone does not influence cell cycle

progression (Figure 2A). Consistently, *cho1*Δ cells showed no significant growth defects at

37°C compared to growth at 30°C (Figure 2B). The void zone rarely formed at 30°C (Figure

1B), but once formed, the void zone was largely maintained even after 90 min of incubation at

30°C (Figure. 2C). These results suggest that the void zone is a very stable lipid domain. On

the other hand, the void zone was essentially undetectable in a saturated culture with high cell

density and very low cell growth. Thus, we investigated whether the frequency of the void

zone formation is associated with the growth phase (Figure 2D). Yeast cells show rapid

growth in the presence of abundant carbon sources such as glucose. When glucose is depleted,

the cells switch to a slower growth rate using ethanol as a metabolic carbon source via a

diauxic shift and consequently enter the stationary phase (Gray *et al.,* 2004). As shown in

Figure 2D, the frequency of the void zone detection was increased early in the logarithmic

growth phase and began to decrease starting from the middle phase. Remarkably, the void

zone rapidly disappeared when the cells entered the diauxic shift, and no changes were

observed after 12 hours, suggesting that the void zone is formed and maintained only in the

presence of abundant carbon sources. To investigate the disappearance of the void zone, we

examined whether certain stresses induce the disappearance. *cho1*Δ cells grown at 37°C were

incubated for additional 3 h at 37°C under three stress conditions, including ATP depletion,

glucose starvation, and translation inhibition; then, the cell numbers and the frequency of the

void zone were measured. Cell growth was almost completely blocked by all these stresses,

and the frequency of the void zone detection was significantly reduced by ATP depletion and

glucose starvation, but not by translation inhibition (Figure 2E). These results suggest that the

maintenance of the void zone is energy-dependent and does not necessarily require cell

growth.

We also noticed that the void zone tends to occur in the mother cells rather than the daughter

cells (Figure 2F). To confirm this, the distribution of the void zone was categorized into the

following groups; cells with no bud or small bud (type 1), cells with mid-large bud in which

the void zone occurs only in the mother cell (type 2), and cells in which the void zone occurs

in the mother and daughter cells (type 3). As a result, almost all void zones appeared only in

the mother cells before or during budding (type 1 and 2); very few type 3 cells were detected,

and no cells with void zones formed only in the daughter cells were observed (Figure 2F).

This biased distribution was observed even under the condition of $OD_{600} = 4.0$-6.0, which

shows high frequency of the void zone. Exocytosis and endocytosis frequently occur in a bud,

which is a polarized growth site, compared to a mother cell. These results suggest that the

void zone is generated in a more static membrane.

Figure 2

Figure 2 The void zone is stably maintained in an energy-dependent manner

(A) Time-lapse imaging of the void zone. *cho1Δ* cells expressing GFP-Snc1-pm were grown

in YPDA medium at 37°C. Two representative examples are shown; a cell with a growing

bud (upper panels) and a dividing cell (lower panels) with the void zones. Arrowheads

indicate the void zone. Scale bar: 5 μm. (B) Normal growth of *cho1Δ* cells at 37°C. Serial

dilutions of cultures were spotted onto the YPDA plates followed by incubation at 30°C or

37°C for 1 d. (C) The void zone formed at 37°C remains stable at 30°C. *cho1Δ* cells

expressing GFP-Snc1-pm were grown in YPDA at 37°C, and then shifted to 30°C for 90 min.

The percentage of cells with the void zone before and after incubation at 30°C was measured.

Data of three independent experiments (n > 100 cells each) are shown as the mean and SD.

(D) Frequency of the void zone generation at various growth stages. *cho1Δ* cells expressing

GFP-Snc1-pm were grown in YPDA at 37°C, and the percentage of the cells with void zones

(blue bars) and the OD_{600} (orange line) were determined. Data of three independent

experiments (n > 100 cells each) are shown as the mean and SD. (E) Disappearance of the

void zone under stress conditions. *cho1Δ* cells expressing GFP-Snc1-pm were grown to the

early log phase at 37°C and were shifted to YPDA medium containing 20 mM sodium azide

(ATP depletion), YPA medium lacking glucose (glucose starvation), and YPDA medium

containing 10 μg/ml cycloheximide (translation inhibition). After incubation for 3 h at 37°C,

OD_{600} and the incidence of the void zone (n > 100 cells) were determined. Data of three

independent experiments are shown as the mean and SD. Asterisks indicate significant

differences determined by the Tukey–Kramer test (***, $p < 0.001$). (F) Biased formation of

the void zone in the mother cell. *cho1Δ* cells expressing GFP-Snc1-pm were prepared as in

(A). A representative pattern of the void zone taken on the different Z focal planes is shown in

the upper panel; the void zone occurs in the mother cell (M) but not in the daughter cell (D).

Arrowheads indicate the void zone. Scale bar: 5 μm. The percentage of the cells with the void

zone in the indicated pattern types (n > 200 cells each) is shown in the lower panel.

Void zone formation is specific to PS-depletion and is a reversible process.

To examine whether the void zone can be detected in the lipid mutants other than *cho1Δ*, we

investigated the mutants of the genes involved in the synthesis of phosphatidylethanolamine

(PE) and phosphatidylcholine (PC). In PE-, and PC-deficient cells, GFP-Snc1-pm was evenly

distributed in the plasma membrane, and the void zone was not detected (Figure 3A). We next

examined whether the recovery of PS levels dissipates the void zone by adding lyso-PS to the

culture media. Incorporated lyso-PS is rapidly converted to PS in the endoplasmic reticulum

(ER) via a *CHO1*-independent pathway (Fairn *et al.,* 2011; Maeda *et al.,* 2013). The recovery

of PS was verified by expressing a PS-specific biosensor Lact-C2 (Yeung *et al.,* 2008). In

*cho1*Δ cells without lyso-PS, the frequency of the void zone did not change for 60 min and

mRFP-Lact-C2 remained diffuse in the cytosol. On the other hand, the addition of lyso-PS

significantly reduced the incidence of the void zone and resulted in the localization of mRFP-

Lact-C2 to the plasma membrane (Figure 3B). These results indicate that the void zone

formation is a reversible process. After the addition of lyso-PS, void zones did not disappear

uniformly, but rather appeared to be gradually repaired from the boundaries (Figure 3C). In

some *cho1*Δ cells, mRFP-Lact-C2 was distributed outside of the void zone (Figure 3D). This

void zone is gradually repaired by PS around the void zone. Our results suggest that the void

zone is a lipid domain that consists of a lipid (lipids), and random distribution of these lipids

in the plasma membrane is facilitated by interaction with PS.

Figure 3

Figure 3 Void zone formation is specific to PS depletion and is a reversible process.

(A) The generation of the void zone is specific to the PS-deficient cells. All strains expressing

GFP-Snc1-pm were grown at 37°C. *cho1Δ* cells were grown in SD medium containing 1 mM

ethanolamine. *psd1Δ psd2Δ* cells were grown in SD medium containing 2 mM choline. *cho2Δ*

opi3Δ cells were grown in SD medium. Percentage of the cells with the void zone is indicated

as the mean and SD (n > 100 cells, three independent experiments). Arrowheads indicate the

void zone. Scale bar: 5 μm. (B) Regeneration of PS by lyso-PS supplementation dissipates the

void zone. Wild-type and *cho1Δ* cells expressing GFP-Snc1-pm and mRFP-Lact-C2 are

shown in the upper left panel. *cho1Δ* cells were grown in YPDA medium at 37°C in the

presence or absence of lyso-PS (20 μM). Percentage of the cells with the void zone was

examined at 20 and 60 min and shown as the mean and SD (n > 100 cells, three independent

experiments) (upper right panel). Representative images are shown in the lower panel.

Arrowheads indicate the void zone. Scale bars: 5 μm. (C) Time-lapse imaging of the void

zone disappearance induced by lyso-PS addition. *cho1Δ* cells expressing GFP-Snc1-pm were

prepared as in (B). Cell suspension was placed on the agarose gel pad immediately after

mixing with lyso-PS and time-lapse imaging was started. Three examples are shown in the

pseudo-colour. Arrowheads indicate the void zone. Scale bar: 5 μm. (D) Void zones are not

rapidly dissipated by PS. *cho1Δ* cells expressing GFP-Snc1-pm and mRFP-Lact-C2 were

prepared as in (B). Images 20 min after supplementation with lyso-PS are shown.

Fluorescence intensities of GFP and mRFP in the cell periphery (arrows) are plotted on the

right. Arrowheads indicate the void zone. Scale bar: 5 μm.

Void zone restricts lateral diffusion of the inner leaflet-anchored proteins, but has no

effect on a GPI-anchored protein.

We next examined whether the plasma membrane-localized peripheral membrane proteins

can be distributed in the void zone. Three types of proteins were tested (Figure 4A). Ras2 is a

small GTPase with the C-terminal lipid moiety inserted into the cytoplasmic leaflet of the

plasma membrane. Ras2 undergoes farnesylation and palmitoylation, and the latter is required

for membrane localization (Bhattacharya *et al.,* 1995). Gap1C is the C-terminal cytosolic

region of the amino acid permease Gap1. Gap1 localizes to the plasma membrane, and Gap1C

without the transmembrane domain is also localized at the plasma membrane (Popov-

Čeleketić *et al.,* 2016). Gap1C is palmitoylated; however, its membrane localization is due to

its amphipathic helix structure and not to the palmitoyl anchor (Popov-Čeleketić *et al.,* 2016).

Gas1 is a cell wall protein with glycosylphosphatidylinositol (GPI)-anchor inserted into the

extracellular leaflet of the plasma membrane (Nuoffer *et al.,* 1991). All these proteins were

mainly localized to the plasma membrane in the wild-type cells (Figure 4B). In *cho1Δ* cells

harbouring the void zone, Ras2 and Gap1C were absent from the void zone similar to Snc1-

pm (Figure 4B). In contrast, Gas1 was uniformly distributed in the plasma membrane

regardless of the void zone (Figure 4B). These results suggest that the void zone restricts the

lateral diffusion of the inner leaflet-associated proteins and the transmembrane proteins but

does not influence the diffusion of proteins anchored to the outer leaflet. Normally, PS is

abundantly distributed in the inner leaflet of the plasma membrane. These results suggest that

formation of the void zone is mainly due to lipid changes in the inner leaflet rather than that in

the outer leaflet.

Figure 4

Figure 4 Distribution of peripheral membrane proteins in the void zone.

(A) Localization of three peripheral membrane proteins. (B) Ras2 and Gap1C were excluded

from the void zone, but Gas1, a GPI-AP, was not excluded. Cells expressing GFP-Ras2 under

the control of the *GAL1* promoter and mRFP-Snc1-pm were grown in YPGA at 37°C. Cells

expressing GFP-Gap1C and mRFP-Snc1-pm or GFP-Snc1-pm and mRFP-Gas1 were grown

in SDA-U medium containing 1 mM ethanolamine at 37°C. Fluorescence intensities of GFP

and mRFP in the cell periphery (arrows) are plotted on the right. Arrowheads indicate the

void zone. Scale bars: 5 μm. (C) Cells observed as in (B) are categorized by the distribution of

each protein in the void zones detected by Snc1-pm (n > 50 cells with the void zone).

Void zone requires ergosterol and sphingolipid for its formation.

Molecular dynamics computer simulation, the giant unilamellar vesicle (GUV) assay, and

giant plasma membrane vesicle (GPMV) assay have shown that transmembrane helices and

transmembrane proteins are excluded from cholesterol-enriched lipid domains (Sengupta *et*

al., 2008; Schäfer *et al*., 2011). Transmembrane proteins are also excluded from ergosterol-

enriched raft-like domains that are generated in the vacuolar membrane of yeast in stationary

phase (Toulmay and Prinz, 2013; Wang *et al.,* 2014; Tsuji *et al.,* 2017). Therefore, we

examined whether the void zone is a sterol-enriched lipid domain. To assess the distribution

of ergosterol, a major sterol in yeast, cells were stained with the sterol-binding dye filipin.

Filipin showed a punctate distribution in the plasma membrane of the wild-type cells as

reported previously (Grossmann *et al.,* 2007), whereas in *cho1Δ* cells, filipin mainly

accumulated at the void zones (Figure 5A), indicating that the void zone is rich in ergosterol.

Similar to sterols, sphingolipids are important for the formation of specialized membrane

domains (Dietrich *et al.,* 2001; Baumgart *et al.,* 2003; de Almeida *et al.,* 2003; Veatch and

Keller, 2003). We therefore examined whether the biosynthesis of ergosterol and sphingolipid

is required for the formation of the void zone. Erg2, Erg3, and Erg4 are essential enzymes for

the synthesis of ergosterol (Silve *et al.,* 1996; Arthington *et al.,* 1991; Lai *et al.,* 1994). Ipt1 is

an inositolphosphotransferase required for the synthesis of mannosyl-

diinositolphosphorylceramide (M(IP)2C), the most abundant sphingolipid in yeast (Dickson *et*

al., 1997). Elo2 and Elo3, fatty acid elongases, participate in the long chain fatty acid

biosynthesis of sphingolipids (Oh *et al.,* 1997). Sur2 and Scs7 are hydroxylases involved in

the hydroxylation of sphingolipids (Haak *et al.,* 1997). Disruption of any of these genes

essentially abolished the formation of the void zone (Figure 5B). Consistently, an inhibitor of

ergosterol synthesis, fluconazole, blocked the formation of the void zone (Figure 5C). These

results indicate that ergosterol and sphingolipids are necessary for the formation of the void

zone and suggest that the void zone may be a novel lipid domain composed of abundant

ergosterol and sphingolipids.

We further investigated whether the formation of the void zone results from an increase in

the ergosterol level. The ergosterol level was increased in the *cho1Δ* cells grown at 37°C

compared to that in the wild-type cells grown at 30°C. However, a similar increase in the

ergosterol level was observed in the wild-type cells grown at 37°C and *cho1Δ* cells grown at

30°C, in which the void zone was not formed (Figure 5D). Therefore, the generation of the

void zone is not closely correlated with an increase in ergosterol.

The finding that the void zone is an ergosterol-rich lipid domain prompted us to investigate

the distribution of other lipids that can be visualized using the corresponding probes. To

examine the distribution of phosphatidylinositol 4-phosphate (PI(4)P), phosphatidylinositol

4,5-bisphosphate (PI(4,5)P2), and phosphatidic acid (PA), the PH domains of yeast Osh2 and

human PLCδ and the PA-binding domain of yeast Spo20 were used as the probes,

respectively (Roy and Levine, 2004; Lemmon *et al.,* 1995; Watt *et al.,* 2002; Nakanishi *et al.,*

2004). These probes were uniformly distributed in the plasma membrane in the wild-type

cells; however, none of the probes were detected in the void zone (Figure 5E). This result

suggests that PI(4)P, PI(4,5)P2, and PA cannot enter the void zone by lateral diffusion. The

majority of yeast phospholipids have mono- or di-unsaturated fatty acids (Schneiter *et al.,*

1999; Ejsing *et al.,* 2009; Klose *et al.,* 2012). In artificial membrane systems, these

unsaturated lipids have also been found to be separated from cholesterol-enriched membrane

domains (Baumgart *et al.,* 2003; Baumgart *et al.,* 2007; Risselada and Marrink, 2008;

Lingwood and Simons, 2010).

Our results show that the void zone is a lipid domain with an unusual assembly of ergosterol

that cooperates with sphingolipids and thereby has novel properties with limited lateral

diffusion of the peripheral membrane proteins and certain types of glycerophospholipids.

Figure 5

Figure 5 Ergosterol is accumulated in the void zone.

(A) Ergosterol accumulation in the void zone. Cells expressing GFP-Snc1-pm were grown in

YPDA medium at 37°C, fixed, and stained with filipin. The filipin staining patterns of *cho1Δ*

cells containing the void zone (n = 100) were categorized and are shown in the lower left

panel. Arrowheads indicate the void zone. Scale bars: 5 μm. (B) The formation of the void

zone was dependent on ergosterol and sphingolipids. Cells expressing GFP-Snc1-pm were

grown overnight in YPDA medium at the indicated temperature. The incidence of the void

zone (n > 100 cells) was examined in three independent experiments, and the data are shown

as the mean and SD. Asterisks indicate significant differences from the Cho1-depleted cells

grown at 37°C according to the Tukey–Kramer test (***, p < 0.001). (C) The void zone was

not generated after ergosterol depletion by fluconazole. *cho1Δ* cells expressing GFP-Snc1-pm

were precultured in YPDA medium at 30°C and incubated at 37°C for 6 h with or without 50

μM fluconazole. The incidence of the void zone (n > 100 cells) was examined and the data are

shown as in (B). Asterisks indicate a significant difference according to the Student's t-test

(**, p < 0.005). (D) The total levels of ergosterol in the wild-type and *cho1Δ* cells. Cells were

grown overnight in YPDA medium at 30°C or 37°C, and total cellular lipids were extracted.

Ergosterol contents were analysed by TLC. The ratio of total ergosterol to total phosphate

(mol/mol) is shown as the mean and SD of four independent experiments. Asterisks indicate

significant differences according to the Tukey–Kramer test (***, p < 0.001 compared to the

wild-type at 30°C). (E) PIPs and PA were excluded from the void zone. Cells expressing

mRFP-Snc1-pm and either Osh2-2xPH-3xGFP or GFP-PLCδ-2XPH were grown and

observed as in (B). Cells expressing GFP-Snc1-pm and mCherry-Spo20-PABD were grown

in SDA-U medium containing 1 mM ethanolamine at 37°C. Fluorescence intensities of GFP

and either mRFP or mCherry around the cells (arrows) are plotted on the right. Arrowheads

indicate the void zone. Scale bar: 5 μm. The distribution of each probe in the void zones

detected by Snc1-pm (n > 50 cells with the void zone) is shown in the lower panel.

Vacuole, a lysosome-like organelle in yeast, contacts the void zone

Observations using the lipid probes suggest phase separation in the plasma membrane; the

lipid composition of the void zone is different from that in the other plasma membrane

regions. The plasma membrane and the ER form the membrane contact sites (MCSs) via ER-

resident tethering proteins that interact with phospholipids of the plasma membrane (Saheki

and De Camilli, 2017). To test whether the void zone can influence this ER-PM contact, we

examined two ER marker proteins, Hmg1 and Rtn1 (Koning *et al.,* 1996; De Craene *et al.,*

2006). The cortical ER (cER) was clearly absent at the void zone in the *cho1Δ* cells (Figure

6A; 89.2% of Hmg1, 92.9% of Rtn1, n > 100 cells, respectively). The ER-PM tethering

proteins Tcb1/2/3, and Ist2 were reported to bind to PS and PI(4,5)P2, respectively (Schulz

and Creutz, 2004; Fischer *et al.,* 2009). Since PS is not synthesized in the *cho1Δ* cells and

PI(4,5)P2 is not distributed in the void zone (Figure 5E), the association of cER with the

plasma membrane may be lost in the void zone.

To investigate whether the void zone influences the distribution or morphology of other

organelles, several organelle markers were examined in the *cho1Δ* cells. Surprisingly, we

found that vacuoles were in contact with the void zones (Figure 6B). The observations using

Snc1-pm and FM4-64 indicate that the contact between the void zone and the vacuole was

detected in 45.1% of the *cho1Δ* cells with the void zone (45.1 ± 7.0%, five independent

experiments, n > 50 cells each). The proximity of the non-void zone regions and vacuoles was

observed only in 7.8% of the *cho1Δ* cells (7.8 ± 1.7%, three independent experiments, n > 100

cells each) compared to that in the wild-type cells (10.1 ± 2.1%, three independent

experiments, n > 100 cells each). These data suggest that frequent contact of the vacuoles

with the plasma membrane is specific to the void zone. The fact that not all void zones are in

contact with the vacuoles indicates that the exclusion of the transmembrane proteins from the

void zones was not caused by the vacuole contact (Figure 6B). On the other hand, the *trans-*

Golgi network (TGN), lipid droplets, and mitochondria did not contact the void zone (Figure

S2), suggesting that only vacuoles interact with the void zone. This type of contact between

the plasma membrane and the vacuoles or lysosomes has not been reported, and we refer to

the contact between the void zone and vacuoles as the "V-V contact".

In stationary phase yeast cells, a raft-like domain is formed in the vacuolar membrane, where

lipophagy, the uptake of the lipid droplets, occurs (Toulmay and Prinz, 2013; Wang *et al.,*

2014; Tsuji *et al.,* 2017). The properties of this vacuolar microdomain are very similar to the

properties of the void zone, including absence of the transmembrane proteins and ergosterol

enrichment. To test whether this vacuolar microdomain is present in the void zone contact

area, we examined three vacuolar transmembrane proteins, Vph1, Cot1, and Zrc1 (Manolson

et al., 1992; Li and Kaplan, 1998). These proteins were uniformly distributed in the vacuolar

membrane of the wild-type cells; however, in some vacuoles in contact with the void zone of

the *cho1Δ* cells, these proteins were excluded from the contact area (Figure 6C). One of these

proteins, Vph1, had the highest frequency of segregation on vacuoles in contact with the void

zone (73.5% of Vph1, 41.5% of Cot1, and 30.0% of Zrc1, n > 50). In contrast, the exclusion

of FM4-64 dye in vacuoles of the cells with the V-V contacts was not detected (n > 50; Figure

6B). These data suggest that certain vacuoles form a membrane domain at the V-V contact

site.

To understand the dynamics of the V-V contact, we used time-lapse imaging with GFP-

Snc1-pm and Vph1-mRFP (Movie 1) and detected two patterns. First, the movement of a

vacuole into the void zone was observed (Figure 6D, shown as "a"). The speed of the vacuole

migration to the void zone was highly variable between individual cells. However, some void

zones had no contact with the vacuoles during the 1 h observation. Second, the V-V contact

lasted over half an hour (Figure 6D, shown as "b"). In one hour of observation of the cells

with the void zone, 60% of the cells had the V-V contact over 30 min (Figure 6E). The

disappearance of the void zone after its formation was rarely observed, and this phenomenon

appears to be unrelated to the presence or absence of the V-V contacts. In addition, no

vacuoles left the void zone while the void zone was maintained. The molecular basis of the V-

V contact is unknown although this contact appears to be stable. Vacuoles in contact with the

void zone underwent fission and fusion (Figure S3).

Figure 6

Figure 6 Vacuoles contact the void zone.

(A) Cortical ER is disassociated from the void zone. Cells expressing mRFP-Snc1-pm and

either Hmg1-GFP or Rtn1-GFP were grown in YPDA medium at 37°C. The mRFP-Snc1-pm

and ER marker proteins are shown in green and magenta, respectively. Arrowheads indicate

the void zone. Scale bar: 5 μm. (B) Vacuoles contact the void zone. Cells expressing GFP-

Snc1-pm were grown in YPDA medium at 37°C and stained with FM4-64 for 20 min.

Arrowheads indicate the void zone. Scale bar: 5 μm. (C) Separation of the vacuolar protein in

the void zone contact region. Cells expressing mRFP-Snc1-pm and either Vph1-GFP or GFP-

Cot1 and cells expressing GFP-Snc1-pm and mRFP-Zrc1 were grown in YPDA medium at

37°C. Snc1-pm and vacuolar proteins are shown in green and magenta, respectively.

Arrowheads indicate the void zone. Scale bar: 5 μm. (D) Time-lapse imaging of the vacuole-

void zone contact. *cho1Δ* cells expressing GFP-Snc1-pm and Vph1-mRFP were grown in

YPDA medium at 37°C. Three examples with two distinctive patterns (a, b) are shown. In the

bottom left scheme, black regions in the plasma membrane indicate the void zone. Numbers

indicate time in min. Scale bar: 5 μm. (E) The vacuole-void zone contact is stable. Cells

observed by time-lapse imaging as in (D) are categorized by the presence or absence of the V-

V contact and the void zone behaviour (n > 100 cells with the void zone). Estimation was

based on the void zone or the V-V contact lasting more than 30 min during 1 h observation.

Identification of the genes required for the formation of the void zone

To further understand the mechanism of the void zone formation, we created a series of

deletion mutants on the background of the *cho1Δ* or glucose-repressible *P$_{GAL1}$-3HA-CHO1*

mutations and examined their effect on the generation of the void zone (Figure 7A). Various

genes involved in sterol trafficking were tested, and the results indicate that *kes1Δ* (*osh4Δ*)

mildly and *arv1Δ* significantly reduced the void zone formation. Kes1 is one of the yeast

oxysterol-binding proteins that exchanges sterols for PI(4)P between the lipid membranes

(Jiang, *et al.,* 1994; de Saint-Jean *et al.,* 2011). Arv1 was implicated in the GPI-anchor

biosynthesis and transport and in intracellular sterol distribution (Kajiwara *et al.,* 2008; Beh

and Rine, 2004). Both Kes1 and Arv1 are involved in the sterol transport; however, their

contribution to the sterol transport to the plasma membrane is very low (Georgiev *et al.,* 2011;

Georgiev *et al.,* 2013). These proteins regulate sterol organization in the plasma membrane;

the mutations influence the sensitivity to the sterol-binding drugs and sterol-extraction

efficiency of MβCD (Georgiev *et al.,* 2011; Georgiev *et al.,* 2013). These differences in sterol

organization may influence the formation of the void zone. Npc2 is an orthologue of

Niemann-Pick type C protein and plays an essential role together with Ncr1 in sterol insertion

into the vacuolar membrane from the inside of the vacuole, which is required for the

formation of the raft-like vacuolar domain during lipophagy in the stationary phase (Tsuji *et*

al., 2017). However, deletion of Npc2 did not influence generation of the void zone or

formation of the V-V contacts accompanied with protein-free vacuolar domain (Figure S4).

This result suggests that the void zone is formed independently of Ncr1/Npc2-mediated sterol

transport possibly because Ncr1/Npc2 are not involved in sterol organization in the plasma

membrane. The vacuolar domains detected at the V-V contact region and generated during the

stationary phase lipophagy are similar in appearance; however, the processes of their

formation may be different.

The void zone formation was observed in the absence of Pep4, a major vacuolar protease,

and Vac17, a myosin adapter required for inheritance of vacuoles. However, the void zone

formation was strongly suppressed by deletion of Vma2 and Fab1.

Vma2 is a subunit of the V-ATPase that regulates pH homeostasis and functions as a pH

sensor (Marshansky *et al.,* 2014). Consistent with this observation, other genes involved in

pH homeostasis (*NHA1*, *NHX1*, *RIM21*, and *RIM101*) were required for the void zone

formation (Figure 7A, pH regulation) (Sychrová *et al.,* 1999; Brett *et al.,* 2005; Obara *et al.,*

2012). Thus, we examined the effect of pH of the medium on the formation of the void zone.

The formation of the void zone was strongly suppressed by increasing pH in the medium from

6.6 to 7.5. (Figure 7B). On the other hand, the low pH medium (pH 4.0) slightly increased the

formation of the void zone. Interestingly, the frequency of the V-V contacts was significantly

reduced in the low pH medium (Figure 7C). The mechanism of these pH-dependent

phenomena is unclear although they may be important in assessment of the molecular basis of

the formation of the void zones and V-V contacts.

Fab1 is a phosphatidylinositol 3-phosphate (PI(3)P) 5-kinase that generates

phosphatidylinositol 3,5-bisphosphate (PI(3,5)P2) (Cooke *et al.,* 1998). PI(3,5)P2 functions as

a signal lipid in intracellular homeostasis, adaptation, and retrograde membrane trafficking

(Jin *et al.,* 2016). We speculated that the defects in the retrograde transport may indirectly

influence the void zone formation in the plasma membrane and thus examined various genes

involved in the membrane transport. Strikingly, conserved protein complexes, retromer, class

C core vacuole/endosome tethering (CORVET), homotypic fusion and vacuole protein sorting

(HOPS), and endosomal sorting complexes required for transport (ESCRT) were required for

the void zone formation (Figure 7A and D). As the name implies, the Vps proteins belonging

to these complexes were identified using mutants defective in vacuolar protein sorting (VPS)

(Robinson *et al.,* 1988; Rothman *et al.,* 1989). Dysfunction of these complexes perturbs

intracellular vesicle trafficking (Schmidt and Teis, 2012; Balderhaar and Ungermann, 2013;

Burd and Cullen, 2014), which may influence the plasma membrane recycling of cargo and

lipids involved in the void zone formation. Similarly, proteins involved in the retrograde

transport, such as the SNAREs Pep12 and Tlg2, the epsin-like adapter Ent3/Ent5, and the

clathrin adapter Gga1/Gga2, were required for the void zone formation. A dynamin-like

GTPase Vps1, Arf-like GTPase Arl1, and Rab6 GTPase homologue Ypt6 are known to be

closely related to the membrane trafficking (Vater *et al.,* 1992; Li and Warner, 1996;

Rosenwald *et al.,* 2002). Consistent with this notion, *vps1Δ, arl1Δ,* and *ypt6Δ* inhibited the

void zone formation. Impaired membrane transport in the inner membrane system can be

manifested as the changes in the plasma membrane lipid organization and/or defects in the pH

control. However, Apl2 and Apl1, the subunits of the adaptor complexes AP-1 and AP-2,

respectively, had little contribution to the void zone formation presumably because these

mutants had insignificant disruption of the membrane trafficking compared to the effects of

ent3Δ, ent5Δ and *gga1Δ gga2Δ* (Yeung *et al.,* 1999; Sakane *et al.,* 2006; Morvan *et al.,*

2015). Deletion of *APL5*, which encodes the subunit of AP-3 responsible for the transport

from the Golgi to the vacuole (Dell'Angelica, 2009), slightly reduced the void zone formation,

suggesting the importance of the vacuolar functions for the void zone formation.

Deletion of the autophagy-related genes (*atg1Δ*, *atg10Δ*, *atg12Δ*, and *atg15Δ*) did not

influence the void zone formation. *ATG1* is one of the core ATG genes (Mizushima *et al.,*

2011). The void zone was generated and the V-V contact with the vacuolar microdomain was

observed in the absence of Atg1 (Figure 7A, and Figure S4). This result suggests that the void

zone formation is independent of autophagy consistent with our notion that direct or indirect

effects on lipid organization in the plasma membrane (e.g., *via* the Vps pathway) are

influencing the void zone formation.

We also examined the effect of mutations in the flippase-related proteins. The deficiency of

Cdc50 or Any1/Cfs1 localized in endosomes and the TGN had little effect on the void zone

formation (Saito *et al.,* 2004; van Leeuwen *et al.,* 2016; Yamamoto *et al.,* 2017); however,

disruption of Lem3 localized in the plasma membrane completely suppressed the generation

of the void zone (Kato *et al.,* 2002). The Lem3-Dnf1/2 flippase complexes translocate

glycerophospholipids, but not ceramides and sphingolipids, to the cytoplasmic leaflet of the

lipid bilayer (Pomorski *et al.,* 2003; Saito *et al.,* 2004; Furuta *et al.,* 2007). We assumed that

disruption of the phospholipid asymmetry by *lem3Δ* may influence ergosterol behaviour in

the plasma membrane. To test this hypothesis, we examined the sensitivity to an antifungal

ergosterol-binding drug, amphotericin B (AmB) (Kamiński, 2014). The results indicate that

lem3Δ is highly sensitive to AmB, which is detected in the *cho1Δ* background cells, and this

effect was cancelled by addition of *erg6Δ* that causes defects in ergosterol biosynthesis

(Figure 7E). Thus, the disruption of phospholipid asymmetry alters the ergosterol distribution

in the plasma membrane, thereby suppressing the void zone formation (see discussion).

Although it is not yet clear about the mechanisms that some of these additive mutations

inhibit the void zone formation, these findings may help to elucidate the nature of the void

zone in the future.

Figure 7

Figure 7 A search for the genes required for the void zone formation

(A) Frequency of the void zone in the mutant cells generated on *cho1Δ* background or under

Cho1-depleted conditions. The incidence of the void zone was determined (n > 100 cells,

three independent experiments) and is shown as the mean and SD. The mutations responsible

for the low incidence (under 20%) are shown in red. (B) The void zone formation is

suppressed by high pH. *cho1Δ* cells expressing GFP-Snc1-pm were grown overnight in

YPDA at the indicated pH at 37°C. The incidence of the void zone was determined (n > 100

cells, five independent experiments) and is shown as a box plot. Asterisks indicate significant

differences according to the Tukey–Kramer test (**, $p < 0.005$; ***, $p < 0.001$). (C) Low pH

decreases the frequency of the V-V contact. *cho1Δ* cells expressing GFP-Snc1-pm and Vph1-

mRFP were grown in YPDA at the indicated pH at 37°C. The incidence of the V-V contacts

was determined (n > 100 cells with the void zone, three independent experiments) and is

shown as the mean and SD. Asterisks indicate a significant difference according to the

Student's t-test (***, $p < 0.001$). (D) A scheme of the membrane trafficking. Proteins and

protein complexes required for the formation of the void zone are shown in red. (E) The loss

of phospholipid asymmetry in the plasma membrane results in high sensitivity to

amphotericin B. Serial dilutions of the cultures were spotted onto YPDA plates containing 1.0

μM amphotericin B and incubated at 30°C for 2 d.

566　**Discussion**

In the artificial membranes, phase separation causes micron-scale separation of proteins and

lipids; such large separation does not occur in the plasma membrane of the living cells. The

mechanism that enables random distribution of proteins and lipids throughout the plasma

membranes on the macroscopic scale by preventing large-scale phase separation has not been

understood. We found that in PS-deficient cells grown at high temperatures, the protein-free

membrane domain "void zone" develops in the plasma membrane and exhibits properties

similar to phase-separated artificial membranes (Figure 8). We propose a new role of PS in

membrane organization and suggest that PS prevents the occurrence of abnormal membrane

domains due to phase separation and ensures macroscopic homogeneity of the distribution of

the molecules on the plasma membrane.

Mechanisms of the generation and disappearance of the void zone

PS is a phospholipid mainly distributed in the inner leaflet of the plasma membrane and has a

relatively high affinity for cholesterol (Maekawa and Fairn, 2015; Nyholm *et al.,* 2019).

Therefore, loss of PS may alter relative affinity of the phospholipids for ergosterol in the inner

leaflet and may also influence the transbilayer distribution of ergosterol. Almost all yeast

phospholipids, including PS, have at least one unsaturated fatty acid; however, a small

fraction of phosphatidylinositol (PI) has only saturated fatty acids (Schneiter *et al.,* 1999;

Ejsing *et al.,* 2009; Klose *et al.,* 2012). Molecular species of PS, PE, PC, and PI were

characterized in the isolated plasma membrane; interestingly, 29.2% of PI have two saturated

fatty acids (Schneiter *et al.,* 1999). Relative affinity of ergosterol to various phospholipid

species is unclear. One hypothesis is that ergosterol is not clustered due to the interactions

with unsaturated PS in the wild-type cells; however, in *cho1Δ* cells, saturated PI becomes the

most dominant interaction partner of ergosterol in the inner leaflet of the plasma membrane

thus creating a driving force for phase separation. When PS is resynthesized in the *cho1Δ*

cells after the addition of lyso-PS, the void zone may disappear because PS becomes the

predominant interaction partner of ergosterol (Figure 3B and C).

Generation of the void zone requires PS deficiency and high temperature conditions. In

artificial membrane systems, phase separation occurs at a temperature below the miscibility

transition temperature, which is dependent on the lipid composition of the membrane (Veatch

and Keller. 2002; Baumgart *et al.,* 2007; Leventable et al., 2009). Contrary to these phenomena,

the void zone is rarely formed at 30°C and occurs by incubation at 37°C (Figure 1B); thus, we

speculate that a change in miscibility transition temperature due to remodeling of lipid

composition at 37°C may be one of the triggers for void zone formation in *cho1Δ* cells. In

yeast, high temperature reduces the degree of unsaturation and increases the acyl chain length

of glycerophospholipids (Klose *et al.,* 2012), and these events promote phase separation in the

liposomes (Engberg *et al.,* 2016). In addition, our results that several hours of incubation at

37°C was required to generate the void zone (Figure 1B), and that the developed void zone

was maintained for 90 min after the temperature was returned to 30°C (Figure 2C), are

consistent with this idea and suggest that remodeling of the lipid composition takes a long

time. A combination of PS deficiency and high temperature-induced lipid remodelling may

create a specific membrane environment that results in the void zone development.

The void zones rapidly disappeared when the *cho1Δ* cells undergo a diauxic shift (Figure

2D) or when glucose or ATP is depleted from the medium (Figure 2E). During glucose

starvation, activities of the plasma membrane H^+-ATPase Pma1 and V-ATPase are reduced

(Young *et al.,* 2010; Dechant *et al.,* 2010). Therefore, the disappearance of the void zone may

be caused by a disturbance of pH homeostasis consistent with the suppression of the void

zone formation by mutations in the genes involved in pH homeostasis, including *VMA2*

(Figure 7A). Reduced activities of Pma1 and V-ATPases lead to acidification of the cytosol,

and the void zone formation is also inhibited in the medium with elevated pH (Figure 7B).

The relationship between pH and phase separation is largely unknown; however, it has been

reported that phase separation occurs at low pH and not at high pH in the artificial membranes

that mimic human stratum corneum (Plasencia *et al.,* 2007).

Our results indicate that long chain fatty acids and hydroxylation of sphingolipids are

necessary for the void zone formation (Figure 5B). It has been reported that GUVs prepared

from the yeast total lipids have extensive phase separation, which depends on long fatty acid

elongation and hydroxylation of sphingolipids (Klose *et al.,* 2010). M(IP)2C with a C26 fatty

acid is the most common yeast sphingolipid species and its levels are significantly reduced in

the *elo2Δ* and *elo3Δ* mutants (Oh *et al.,* 1997; Ejsing *et al.,* 2009). A recent study using

asymmetric GUVs suggested that C24 SM and not C16 SM has a role in cholesterol retention

in the inner leaflet of the lipid bilayer (Courtney *et al.,* 2018). Consistently,

dehydroergosterol, a closely related fluorescent analogue of ergosterol, is mainly located in

the inner leaflet of the plasma membrane, and this asymmetry is maintained by sphingolipids

in yeast (Solanko *et al.,* 2018). This type of interaction of ergosterol with sphingolipids may

be necessary for the void zone formation.

Analysis using several mutant strains on the *cho1* background revealed that retrograde

intracellular trafficking has a significant effect on the void zone formation (Figure 7A).

Perturbed membrane transport may alter the distribution of the proteins and the key lipids as

described above. Moreover, *ARV1* and *LEM3* are important for the void zone formation. In

arv1Δ cells and Δtether cells that lack the ER-PM contacts, the transport of ergosterol to the

plasma membrane is essentially unchanged, whereas there were significant changes in

ergosterol accessibility to the compounds, such as increased efficiency of ergosterol

extraction with MβCD and increased sensitivity to the sterol-binding drug nystatin (Georgiev

et al. 2013; Quon *et al.,* 2018). Similarly, *lem3Δ* does not change the total ergosterol level

(Mioka *et al.,* 2018), and Dnf1 and Dnf2 do not contribute to the asymmetric distribution of

the ergosterol analogues across the plasma membrane bilayer (Solanko *et al.,* 2018).

However, *lem3Δ* cells are highly sensitive to AmB (Figure 7E). Thus, similar to the *arv1Δ*

and Δtether mutations, *lem3Δ* may alter the ergosterol distribution in the plasma membrane

thus preventing the formation of the void zone.

Is the void zone a liquid-ordered domain?

To date, there are no known examples of micron-scale phase separation in the plasma

membrane of living cells, and its reason is not clear (Lee *et al.,* 2015; Levental *et al.,* 2020).

Thus, the void zone may be the first report of a micron-scale phase separation in the plasma

membrane in vivo. The following features of the void zone are similar to those of the liquid-

ordered domain in phase-separated artificial membranes: 1) absence of the transmembrane

proteins, 2) absence of certain peripheral membrane proteins, 3) exclusion of certain types of

glycerophospholipids, and 4) enrichment in sterols and the contribution of sphingolipids

(Baumgart *et al.,* 2007; Sengupta *et al.,* 2008; Kaiser *et al.,* 2009; Schäfer *et al.,* 2011). On the

other hand, the formation of void zones requires lipid asymmetry and their maintenance is

energy-dependent (Figure 2E and 7A), which is a very different feature from the membrane

domains in artificial membrane systems. Differences between living and artificial membranes

regarding phase separation have been reported previously; macroscopic phase separation does

not occur in living wild-type yeast cells (Figure 1A and C), but occurs in GUVs formed with

lipids extracted from the wild-type strains (Klose et al., 2010). Similarly, whereas no phase

separation is observed in living mammalian cells over a wide range of temperatures, GPMVs

prepared from the same cells shows macroscopic phase separation (Lee *et al.*, 2015). Further

studies of the void zone would help to bridge the gap between the artificial and biological

membranes. Besides the PS-sterol interaction, there may be other underlying mechanisms by

which living cells avoid macroscopic phase separation.

Membrane contact between the void zone and vacuoles

We have detected the contact between the void zone and the vacuoles (Figure 6B and D,

Figure 8A), and the time-lapse imaging revealed that this contact lasted for at least 30 min in

most cases (Figure 6D and E). These results suggest that the contact vacuoles may not

contribute to the rapid degradation and disappearance of the void zone. The contact vacuoles

are more likely to play a role in sealing the void zone to protect the cells. Recent studies

reported that in the phase-separated membranes with the liquid-disordered and liquid-ordered

domains, membrane permeability is increased at the interface between the two domains

(Cordeiro, 2018). A similar result indicated that the ER-PM contact sites are important for the

maintenance of the integrity of the plasma membrane (Omnus *et al.,* 2016; Collado *et al.,*

2019). In this study, we found that void zone has Lo phase-like properties and that cortical ER

is dissociated from the void zone. Therefore, it is possible that the void zone induces high

local permeability on its borders and low local integrity; thus, the void zone may be a fragile

region of the plasma membrane. The V-V contact may represent a protective cellular response

to these plasma membrane abnormalities.

Our results also suggest that membrane domains may be formed on the vacuolar

membranes in contact with the void zone (Figure 6C). A contact between the membrane

domains accompanied by protein separation has been detected in the nucleus-vacuole junction

(NVJ), which is one of the MCSs that excludes certain proteins from the nuclear and vacuolar

membranes (Pan *et al.,* 2000; Dawaliby and Mayer, 2010; Toulmay and Prinz, 2013). The

transmembrane proteins appear to be completely excluded from the void zone (Figure 1H);

hence, the lipid-protein interactions via protein tethering may be present in the V-V contact

similar to that detected in the ER-PM contact sites.

Figure 8

A

B

Figure 8 Summary and a model of the void zone

(A) Characteristics of the void zone revealed in this study. Our results suggest the following

properties of the void zone: (a) transmembrane proteins and peripheral membrane protein

distributed in the inner leaflet cannot enter the void zone by lateral diffusion; (b) a GPI-AP,

Gas1, localized in the outer layer of the plasma membrane is not influenced by the void zone;

and (c) vacuoles move to form a stable contact with the void zone, and a vacuolar membrane

domain also appears to be formed at this contact site. (B) The putative structure of the void

zone (see discussion).

Materials and Methods

Chemicals, Media, and Genetic Manipulation

Chemicals were purchased from Wako Pure Chemicals Industries (Osaka, Japan) unless

indicated otherwise. Standard genetic manipulations and plasmid transformation of yeast were

performed as described previously (Elble, 1992; Guthrie and Fink, 2002). Yeast strains were

cultured in rich YPDA medium containing 1% yeast extract (Difco Laboratories, MI, USA),

2% Bacto peptone (Difco), 2% glucose, and 0.01% adenine, or synthetic dextrose (SD)

medium containing 0.17% yeast nitrogen base w/o amino acids and ammonium sulfate

(Difco), 0.5% ammonium sulfate, 2% glucose, and the required amino acids or nucleic acid

bases. To induce the *GAL1* promoter, 3% galactose and 0.2% sucrose were used as carbon

sources instead of glucose (YPGA medium). To deplete PS in the *P$_{GAL1}$-3HA-CHO1*

background cells, the cells were grown on YPDA plates for more than 1 day before

inoculation into YPDA medium. Strains carrying *URA3*-harbouring plasmids were cultured in

SD medium containing 0.5% casamino acids (Difco), 0.03% tryptophan, and 0.01% adenine

(SDA-U). When *cho1Δ* or Cho1-depleted strains were cultured in SD or SDA-U medium,

ethanolamine was added to the final concentration of 1 mM. For serial dilution spot assays,

cells were grown to early log phase in an appropriate medium and adjusted to a concentration

of 1.0×10^7 cells/ml. After serial tenfold dilution, 4-µl drops were spotted onto appropriate

plates.

Strains and Plasmids

Yeast strains and plasmids used in this study are listed in Tables S1 and S2, respectively.

Standard molecular biological techniques were used for the construction of the plasmids, PCR

amplification, and DNA sequencing (Sambrook and Russell, 2001). PCR-based procedures

were used to construct the gene deletions and gene fusions with GFP, mRFP, 3HA, and the

GAL1 promoter (Longtine *et al*., 1998). All constructs produced by the PCR-based procedure

were verified by colony PCR to confirm that the replacement occurred at the expected locus.

Sequences of the PCR primers are available upon request.

To construct the *LEU2::mRFP-GAS1* strain, pMF608 was linearized and integrated into the

LEU2 locus. To construct the pRS316-mRFP-GAS1 (pKT2191), the SacII-XmaI fragment

from pMF608 was inserted into the SacII-SmaI gap of pRS316. To construct pRS416-GFP-

GAP1C (pKT2205), the C-terminal cytoplasmic region of Gap1 corresponding to 552-602

amino acids (Popov-Čeleketić *et al.,* 2016) and 257 bp downstream of the *GAP1* coding

region were amplified by PCR, and *PEP12* region of pRS416-GFP-PEP12 (pKT1487) (Furuta

et al., 2007) was replaced with the PCR product. To construct pRS316-mCherry-

SPO20PABD (pKT2206), the DNA fragments of mCherry and the PA-binding region of

Spo20 corresponding to 51-91 amino acids (Nakanishi *et al*., 2004) were inserted into

pRS316 along with P$_{TPI1}$ and T$_{ADH1}$.

Microscopic Observations

Cells were observed using a Nikon ECLIPSE E800 microscope (Nikon Instec, Tokyo, Japan)

equipped with an HB-10103AF super-high-pressure mercury lamp and a 1.4 numerical

aperture 100× Plan Apo oil immersion objective lens (Nikon Instec) with appropriate

fluorescence filter sets (Nikon Instec) or differential interference contrast optics. Images were

acquired using a cooled digital charge-coupled device (CCD) camera (C4742-95-12NR;

Hamamatsu Photonics, Hamamatsu, Japan) and the AQUACOSMOS software (Hamamatsu

Photonics).

GFP-, mRFP-, or mCherry-tagged proteins were observed in the living cells grown in early

to mid-logarithmic phase, harvested, and resuspended in SD medium. Cells were immediately

observed using a GFP bandpass (for GFP) or a G2-A (for mRFP and mCherry) filter set.

Observations were compiled based on the examination of at least 100 cells. For

supplementation of 18:1 lyso-PS (Sigma-Aldrich, St. Louis, MO, USA), a stock solution (10

mg/mL in 0.1% Nonidet P-40) was added to the culture media to the final concentration of 20

μM. For sterol staining, cells were fixed with 5% formaldehyde, washed with PBS, and

labelled for 10 min in 0.5 mg/ml filipin (Sigma-Aldrich) in PBS. For staining of the plasma

membrane by FM4-64 (Thermo Fisher Scientific, MA, USA), cell suspensions were mixed

with an equal volume of 100 μM FM4-64 on a glass slide and observed immediately. For

staining of the vacuolar membranes, cells were labelled for 15 min in 5μM FM4-64, washed

with SD medium, and immediately observed. Fluorescence of filipin and FM4-64 was

observed using a UV and a G2-A filter set, respectively. For the time-lapse imaging, cell

suspension was spotted onto a thin layer of SD medium containing 1 mM ethanolamine and

2% agarose on a glass slide, which was quickly covered with a coverslip. During the time-

lapse imaging, the sample was maintained at 37°C by a Thermo Plate (Tokai Hit, Fujinomiya,

Japan). The incidence of the void zone was assessed based on three or more images of

different focal planes, and based on single focal plane in the time-lapse imaging.

Freeze-fracture replica labelling

Yeast cells sandwiched between a 20-μm–thick copper foil and a flat aluminium disc

(Engineering Office M. Wohlwend, Sennwald, Switzerland) were quick-frozen by high-

pressure freezing using an HPM 010 high-pressure freezing machine according to the

manufacture's instruction (Leica Microsystems, Wetzlar, Germany). The frozen specimens

were transferred to the cold stage of a Balzers BAF 400 apparatus and fractured at −115° to

−105°C under vacuum at ~ 1×10^{-6} mbar.

For genuine morphological observation, samples were exposed to electron-beam evaporation

of platinum-carbon (Pt/C) (1–2 nm thickness) at an angle of 45° to the specimen surface

followed by carbon (C) (10–20 nm) at an angle of 90º. After thawing, the replicas were treated

with household bleach to digest biological materials before mounting on the EM grids for

observation.

For labelling, freeze-fractured samples were subjected to a three-step electron-beam

evaporation: C (2–5 nm), Pt/C (1–2 nm), and C (10–20 nm) as described previously (Fujita *et*

al., 2010). Thawed replicas were treated with 2.5% SDS in 0.1 M Tris-HCl (pH 8.0) at 60°C

overnight; with 0.1% Westase (Takara Bio) in McIlvain buffer (37 mM citrate and 126 mM

disodium hydrogen phosphate, pH 6.0) containing 10 mM EDTA, 30% fetal calf serum, and a

protease inhibitor cocktail for 30 min at 30°C; and with 2.5% SDS in 0.1 M Tris-HCl (pH 8.0)

at 60°C overnight. For labelling of GFP-Pma1, replicas were incubated at 4°C overnight with

a rabbit anti-GFP antibody in PBS containing 1% BSA followed by colloidal gold (10 nm)-

conjugated protein A (University of Utrecht, Utrecht, The Netherlands) for 60 min at 37°C in

1% BSA in PBS. Replicas were observed and imaged with a JEOL JEM-1011 EM (Tokyo,

Japan) using a CCD camera (Gatan, Pleasanton, CA, USA).

Lipid analysis

Total lipids were extracted basically by the Bligh and Dyer method (Bligh and Dyer, 1959).

Cells were grown in 100-200 ml of YPDA medium to 0.8-1.0 OD_{600}/ml at 30°C or 37°C. The

cells were collected and resuspended in 3.8 ml of chloroform-methanol-0.1 M HCl/0.1 M KCl

(1:2:0.8) and lysed by vortexing with glass beads for 1 min. Then, 1.0 ml each of chloroform

and 0.1 M HCl/0.1 M KCl were added followed by centrifugation, isolation of the lipid-

containing phase, and evaporation of the solvent. The extracted lipids were dissolved in an

appropriate volume of chloroform. Total phospholipids were determined by the phosphorus

assay (Rouser *et al*., 1970).

For ergosterol analysis, the samples containing 20 nmol phosphate were subjected to TLC on

an HPTLC plate (Silica gel 60; Merck Millipore, MA, USA) in the solvent system

hexane/diethyl ether/acetic acid (80:20:1) (Dodge and Phillips, 1967). After migration, the

plates were dried and sprayed with a 10% (w/v) cupric sulfate solution in 8% (w/v)

orthophosphoric acid. Plates were heated in an oven at 180°C for 20 min. Plates were scanned

with a CanoScan 8800F image scanner (Canon, Tokyo, Japan) and the acquired images were

quantified using the ImageJ software.

Statistical Analysis

Significant differences in Figures 5C and 7C were determined using a two-sided Student's *t*

test. Significant differences for all other figures were determined by the Tukey–Kramer test.

Acknowledgements

We thank Masahiko Watanabe (Hokkaido University, Sapporo, Japan) for providing the

rabbit anti-GFP antibody for freeze-fracture replica labelling. We thank Takehiko Yoko-o

(AIST, Tokyo, Japan) for providing the pMF608 plasmid. We thank our colleagues in the

Tanaka laboratory for valuable discussions and Eriko Itoh for technical assistance. This work

was supported by Japan Society for the Promotion of Science (JSPS) KAKENHI grants

18K14645 (T.M.), 18K06104 (T.K.), and 19K06536 (K.T.). This work was partly supported

by the Photo-excitonix Project at Hokkaido University.

Competing interests

The authors have no conflict of interests to declare.

Reference

Arikketh, D., Nelson, R., & Vance, J. E. (2008). Defining the importance of

phosphatidylserine synthase-1 (PSS1): unexpected viability of PSS1-deficient mice. *J*

Biol Chem, 283(19), 12888-12897. doi:10.1074/jbc.M800714200

Arthington, B. A., Bennett, L. G., Skatrud, P. L., Guynn, C. J., Barbuch, R. J., Ulbright, C. E.,

& Bard, M. (1991). Cloning, disruption and sequence of the gene encoding yeast C-5

sterol desaturase. *Gene, 102*(1), 39-44. doi:10.1016/0378-1119(91)90535-j

Atkinson, K. D., Jensen, B., Kolat, A. I., Storm, E. M., Henry, S. A., & Fogel, S. (1980).

Yeast mutants auxotrophic for choline or ethanolamine. *J Bacteriol, 141*(2), 558-564.

Audhya, A., & Emr, S. D. (2002). Stt4 PI 4-kinase localizes to the plasma membrane and

functions in the Pkc1-mediated MAP kinase cascade. *Dev Cell, 2*(5), 593-605.

doi:10.1016/s1534-5807(02)00168-5

Balderhaar, H. J., & Ungermann, C. (2013). CORVET and HOPS tethering complexes -

coordinators of endosome and lysosome fusion. *J Cell Sci, 126*(Pt 6), 1307-1316.

doi:10.1242/jcs.107805

Balla, T. (2013). Phosphoinositides: tiny lipids with giant impact on cell regulation. *Physiol

Rev, 93*(3), 1019-1137. doi:10.1152/physrev.00028.2012

Bauer, B. E., Wolfger, H., & Kuchler, K. (1999). Inventory and function of yeast ABC

proteins: about sex, stress, pleiotropic drug and heavy metal resistance. *Biochim

Biophys Acta, 1461*(2), 217-236. doi:10.1016/s0005-2736(99)00160-1

Baumgart, T., Hammond, A. T., Sengupta, P., Hess, S. T., Holowka, D. A., Baird, B. A., &

Webb, W. W. (2007). Large-scale fluid/fluid phase separation of proteins and lipids in

giant plasma membrane vesicles. *Proc Natl Acad Sci U S A, 104*(9), 3165-3170.

doi:10.1073/pnas.0611357104

Baumgart, T., Hess, S. T., & Webb, W. W. (2003). Imaging coexisting fluid domains in

biomembrane models coupling curvature and line tension. *Nature, 425*(6960), 821-

824. doi:10.1038/nature02013

Beh, C. T., & Rine, J. (2004). A role for yeast oxysterol-binding protein homologs in

endocytosis and in the maintenance of intracellular sterol-lipid distribution. *J Cell Sci,

117*(Pt 14), 2983-2996. doi:10.1242/jcs.01157

Bhattacharya, S., Chen, L., Broach, J. R., & Powers, S. (1995). Ras membrane targeting is

essential for glucose signaling but not for viability in yeast. *Proc Natl Acad Sci U S A,*

92(7), 2984-2988. doi:10.1073/pnas.92.7.2984

Bligh, E. G., & Dyer, W. J. (1959). A rapid method of total lipid extraction and purification.

Can J Biochem Physiol, 37(8), 911-917. doi:10.1139/o59-099

Brett, C. L., Tukaye, D. N., Mukherjee, S., & Rao, R. (2005). The yeast endosomal

Na+K+/H+ exchanger Nhx1 regulates cellular pH to control vesicle trafficking. *Mol*

Biol Cell, 16(3), 1396-1405. doi:10.1091/mbc.e04-11-0999

Burd, C., & Cullen, P. J. (2014). Retromer: a master conductor of endosome sorting. *Cold*

Spring Harb Perspect Biol, 6(2). doi:10.1101/cshperspect.a016774

Carquin, M., D'Auria, L., Pollet, H., Bongarzone, E. R., & Tyteca, D. (2016). Recent progress

on lipid lateral heterogeneity in plasma membranes: From rafts to submicrometric

domains. *Prog Lipid Res, 62*, 1-24. doi:10.1016/j.plipres.2015.12.004

Cho, K. J., van der Hoeven, D., Zhou, Y., Maekawa, M., Ma, X., Chen, W., . . . Hancock, J.

F. (2015). Inhibition of Acid Sphingomyelinase Depletes Cellular Phosphatidylserine

and Mislocalizes K-Ras from the Plasma Membrane. *Mol Cell Biol, 36*(2), 363-374.

doi:10.1128/mcb.00719-15

Collado, J., Kalemanov, M., Campelo, F., Bourgoint, C., Thomas, F., Loewith, R., . . .

Fernández-Busnadiego, R. (2019). Tricalbin-Mediated Contact Sites Control ER

Curvature to Maintain Plasma Membrane Integrity. *Dev Cell, 51*(4), 476-487.e477.

doi:10.1016/j.devcel.2019.10.018

Cooke, F. T., Dove, S. K., McEwen, R. K., Painter, G., Holmes, A. B., Hall, M. N., . . .

Parker, P. J. (1998). The stress-activated phosphatidylinositol 3-phosphate 5-kinase

Fab1p is essential for vacuole function in S. cerevisiae. *Curr Biol, 8*(22), 1219-1222.

doi:10.1016/s0960-9822(07)00513-1

Cordeiro, R. M. (2018). Molecular Structure and Permeability at the Interface between Phase-

Separated Membrane Domains. *J Phys Chem B, 122*(27), 6954-6965.

doi:10.1021/acs.jpcb.8b03406

Courtney, K. C., Pezeshkian, W., Raghupathy, R., Zhang, C., Darbyson, A., Ipsen, J. H., . . .

Zha, X. (2018). C24 Sphingolipids Govern the Transbilayer Asymmetry of

Cholesterol and Lateral Organization of Model and Live-Cell Plasma Membranes.

Cell Rep, 24(4), 1037-1049. doi:10.1016/j.celrep.2018.06.104

Dawaliby, R., & Mayer, A. (2010). Microautophagy of the nucleus coincides with a vacuolar

diffusion barrier at nuclear-vacuolar junctions. *Mol Biol Cell, 21*(23), 4173-4183.

doi:10.1091/mbc.E09-09-0782

de Almeida, R. F., Fedorov, A., & Prieto, M. (2003).

Sphingomyelin/phosphatidylcholine/cholesterol phase diagram: boundaries and

composition of lipid rafts. *Biophys J, 85*(4), 2406-2416. doi:10.1016/s0006-

3495(03)74664-5

De Craene, J. O., Coleman, J., Estrada de Martin, P., Pypaert, M., Anderson, S., Yates, J. R.,

3rd, . . . Novick, P. (2006). Rtn1p is involved in structuring the cortical endoplasmic

reticulum. *Mol Biol Cell, 17*(7), 3009-3020. doi:10.1091/mbc.e06-01-0080

de Saint-Jean, M., Delfosse, V., Douguet, D., Chicanne, G., Payrastre, B., Bourguet, W., . . .

Drin, G. (2011). Osh4p exchanges sterols for phosphatidylinositol 4-phosphate

between lipid bilayers. *J Cell Biol, 195*(6), 965-978. doi:10.1083/jcb.201104062

Dechant, R., Binda, M., Lee, S. S., Pelet, S., Winderickx, J., & Peter, M. (2010). Cytosolic pH

is a second messenger for glucose and regulates the PKA pathway through V-ATPase.

Embo j, 29(15), 2515-2526. doi:10.1038/emboj.2010.138

Dell'Angelica, E. C. (2009). AP-3-dependent trafficking and disease: the first decade. *Curr*

Opin Cell Biol, 21(4), 552-559. doi:10.1016/j.ceb.2009.04.014

Dickson, R. C., Nagiec, E. E., Wells, G. B., Nagiec, M. M., & Lester, R. L. (1997). Synthesis

of mannose-(inositol-P)2-ceramide, the major sphingolipid in Saccharomyces

cerevisiae, requires the IPT1 (YDR072c) gene. *J Biol Chem, 272*(47), 29620-29625.

doi:10.1074/jbc.272.47.29620

Dietrich, C., Bagatolli, L. A., Volovyk, Z. N., Thompson, N. L., Levi, M., Jacobson, K., &

Gratton, E. (2001). Lipid rafts reconstituted in model membranes. *Biophys J, 80*(3),

1417-1428. doi:10.1016/s0006-3495(01)76114-0

Dodge, J. T., & Phillips, G. B. (1967). Composition of phospholipids and of phospholipid

fatty acids and aldehydes in human red cells. *J Lipid Res, 8*(6), 667-675.

Douglas, L. M., & Konopka, J. B. (2014). Fungal membrane organization: the eisosome

concept. *Annu Rev Microbiol, 68*, 377-393. doi:10.1146/annurev-micro-091313-

103507

Ejsing, C. S., Sampaio, J. L., Surendranath, V., Duchoslav, E., Ekroos, K., Klemm, R. W., . . .

Shevchenko, A. (2009). Global analysis of the yeast lipidome by quantitative shotgun

mass spectrometry. *Proc Natl Acad Sci U S A, 106*(7), 2136-2141.

doi:10.1073/pnas.0811700106

Elble, R. (1992). A simple and efficient procedure for transformation of yeasts.

Biotechniques, 13(1), 18-20.

Elson, E. L., Fried, E., Dolbow, J. E., & Genin, G. M. (2010). Phase separation in biological

membranes: integration of theory and experiment. *Annu Rev Biophys, 39*, 207-226.

doi:10.1146/annurev.biophys.093008.131238

Engberg, O., Hautala, V., Yasuda, T., Dehio, H., Murata, M., Slotte, J. P., & Nyholm, T. K.

M. (2016). The Affinity of Cholesterol for Different Phospholipids Affects Lateral

Segregation in Bilayers. *Biophys J, 111*(3), 546-556. doi:10.1016/j.bpj.2016.06.036

Fairn, G. D., Hermansson, M., Somerharju, P., & Grinstein, S. (2011). Phosphatidylserine is

polarized and required for proper Cdc42 localization and for development of cell

polarity. *Nat Cell Biol, 13*(12), 1424-1430. doi:10.1038/ncb2351

Fischer, M. A., Temmerman, K., Ercan, E., Nickel, W., & Seedorf, M. (2009). Binding of

plasma membrane lipids recruits the yeast integral membrane protein Ist2 to the

cortical ER. *Traffic, 10*(8), 1084-1097. doi:10.1111/j.1600-0854.2009.00926.x

Fujita, A., Cheng, J., & Fujimoto, T. (2010). Quantitative electron microscopy for the

nanoscale analysis of membrane lipid distribution. *Nat Protoc, 5*(4), 661-669.

doi:10.1038/nprot.2010.20

Furuta, N., Fujimura-Kamada, K., Saito, K., Yamamoto, T., & Tanaka, K. (2007). Endocytic

recycling in yeast is regulated by putative phospholipid translocases and the

Ypt31p/32p-Rcy1p pathway. *Mol Biol Cell, 18*(1), 295-312. doi:10.1091/mbc.e06-05-

0461

Georgiev, A. G., Johansen, J., Ramanathan, V. D., Sere, Y. Y., Beh, C. T., & Menon, A. K.

(2013). Arv1 regulates PM and ER membrane structure and homeostasis but is

dispensable for intracellular sterol transport. *Traffic, 14*(8), 912-921.

doi:10.1111/tra.12082

Georgiev, A. G., Sullivan, D. P., Kersting, M. C., Dittman, J. S., Beh, C. T., & Menon, A. K.

(2011). Osh proteins regulate membrane sterol organization but are not required for

sterol movement between the ER and PM. *Traffic, 12*(10), 1341-1355.

doi:10.1111/j.1600-0854.2011.01234.x

Gray, J. V., Petsko, G. A., Johnston, G. C., Ringe, D., Singer, R. A., & Werner-Washburne,

M. (2004). "Sleeping beauty": quiescence in Saccharomyces cerevisiae. *Microbiol*

Mol Biol Rev, 68(2), 187-206. doi:10.1128/mmbr.68.2.187-206.2004

Grossmann, G., Opekarová, M., Malinsky, J., Weig-Meckl, I., & Tanner, W. (2007).

Membrane potential governs lateral segregation of plasma membrane proteins and

lipids in yeast. *Embo j, 26*(1), 1-8. doi:10.1038/sj.emboj.7601466

Guthrie, C., Fink, GR. (2002). Guide to Yeast Genetics and Molecular Biology, San Diego:

958   Academic Press.

959 Haak, D., Gable, K., Beeler, T., & Dunn, T. (1997). Hydroxylation of Saccharomyces

960   cerevisiae ceramides requires Sur2p and Scs7p. *J Biol Chem, 272*(47), 29704-29710.

961   doi:10.1074/jbc.272.47.29704

962 Jiang, B., Brown, J. L., Sheraton, J., Fortin, N., & Bussey, H. (1994). A new family of yeast

963   genes implicated in ergosterol synthesis is related to the human oxysterol binding

964   protein. *Yeast, 10*(3), 341-353. doi:10.1002/yea.320100307

965 Jin, N., Lang, M. J., & Weisman, L. S. (2016). Phosphatidylinositol 3,5-bisphosphate:

966   regulation of cellular events in space and time. *Biochem Soc Trans, 44*(1), 177-184.

967   doi:10.1042/bst20150174

968 Kaiser, H. J., Lingwood, D., Levental, I., Sampaio, J. L., Kalvodova, L., Rajendran, L., &

969   Simons, K. (2009). Order of lipid phases in model and plasma membranes. *Proc Natl*

970   *Acad Sci U S A, 106*(39), 16645-16650. doi:10.1073/pnas.0908987106

971 Kajiwara, K., Watanabe, R., Pichler, H., Ihara, K., Murakami, S., Riezman, H., & Funato, K.

972   (2008). Yeast ARV1 is required for efficient delivery of an early GPI intermediate to

973   the first mannosyltransferase during GPI assembly and controls lipid flow from the

974   endoplasmic reticulum. *Mol Biol Cell, 19*(5), 2069-2082. doi:10.1091/mbc.e07-08-

975   0740

976 Kamiński, D. M. (2014). Recent progress in the study of the interactions of amphotericin B

977   with cholesterol and ergosterol in lipid environments. *Eur Biophys J, 43*(10-11), 453-

978   467. doi:10.1007/s00249-014-0983-8

979 Kato, U., Emoto, K., Fredriksson, C., Nakamura, H., Ohta, A., Kobayashi, T., . . . Umeda, M.

980   (2002). A novel membrane protein, Ros3p, is required for phospholipid translocation

981   across the plasma membrane in Saccharomyces cerevisiae. *J Biol Chem, 277*(40),

982   37855-37862. doi:10.1074/jbc.M205564200

Klose, C., Ejsing, C. S., García-Sáez, A. J., Kaiser, H. J., Sampaio, J. L., Surma, M. A., . . .

Simons, K. (2010). Yeast lipids can phase-separate into micrometer-scale membrane

domains. *J Biol Chem, 285*(39), 30224-30232. doi:10.1074/jbc.M110.123554

Klose, C., Surma, M. A., Gerl, M. J., Meyenhofer, F., Shevchenko, A., & Simons, K. (2012).

Flexibility of a eukaryotic lipidome--insights from yeast lipidomics. *PLoS One, 7*(4),

e35063. doi:10.1371/journal.pone.0035063

Koning, A. J., Roberts, C. J., & Wright, R. L. (1996). Different subcellular localization of

Saccharomyces cerevisiae HMG-CoA reductase isozymes at elevated levels

corresponds to distinct endoplasmic reticulum membrane proliferations. *Mol Biol Cell,*

7(5), 769-789. doi:10.1091/mbc.7.5.769

Kusumi, A., Fujiwara, T. K., Chadda, R., Xie, M., Tsunoyama, T. A., Kalay, Z., . . . Suzuki,

K. G. (2012). Dynamic organizing principles of the plasma membrane that regulate

signal transduction: commemorating the fortieth anniversary of Singer and Nicolson's

fluid-mosaic model. *Annu Rev Cell Dev Biol, 28*, 215-250. doi:10.1146/annurev-

cellbio-100809-151736

Lai, M. H., Bard, M., Pierson, C. A., Alexander, J. F., Goebl, M., Carter, G. T., & Kirsch, D.

R. (1994). The identification of a gene family in the Saccharomyces cerevisiae

ergosterol biosynthesis pathway. *Gene, 140*(1), 41-49. doi:10.1016/0378-

1119(94)90728-5

Lee, I. H., Saha, S., Polley, A., Huang, H., Mayor, S., Rao, M., & Groves, J. T. (2015). Live

cell plasma membranes do not exhibit a miscibility phase transition over a wide range

of temperatures. *The journal of physical chemistry. B,* 119(12), 4450–4459.

doi.org/10.1021/jp512839q

Lemmon, M. A., Ferguson, K. M., O'Brien, R., Sigler, P. B., & Schlessinger, J. (1995).

Specific and high-affinity binding of inositol phosphates to an isolated pleckstrin

homology domain. *Proc Natl Acad Sci U S A, 92*(23), 10472-10476. doi:10.1073/pnas.92.23.10472

Levental, I., Byfield, F. J., Chowdhury, P., Gai, F., Baumgart, T., & Janmey, P. A. (2009). Cholesterol-dependent phase separation in cell-derived giant plasma-membrane vesicles. *Biochem J, 424*(2), 163-167. doi:10.1042/bj20091283

Levental, I., Levental, K. R., & Heberle, F. A. (2020). Lipid Rafts: Controversies Resolved, Mysteries Remain. *Trends in cell biology, 30*(5), 341–353. doi.org/10.1016/j.tcb.2020.01.009

Lewis, M. J., Nichols, B. J., Prescianotto-Baschong, C., Riezman, H., & Pelham, H. R. (2000). Specific retrieval of the exocytic SNARE Snc1p from early yeast endosomes. *Mol Biol Cell, 11*(1), 23-38. doi:10.1091/mbc.11.1.23

Li, B., & Warner, J. R. (1996). Mutation of the Rab6 homologue of Saccharomyces cerevisiae, YPT6, inhibits both early Golgi function and ribosome biosynthesis. *J Biol Chem, 271*(28), 16813-16819. doi:10.1074/jbc.271.28.16813

Li, L., & Kaplan, J. (1998). Defects in the yeast high affinity iron transport system result in increased metal sensitivity because of the increased expression of transporters with a broad transition metal specificity. *J Biol Chem, 273*(35), 22181-22187. doi:10.1074/jbc.273.35.22181

Lingwood, D., & Simons, K. (2010). Lipid rafts as a membrane-organizing principle. *Science, 327*(5961), 46-50. doi:10.1126/science.1174621

Longtine, M. S., McKenzie, A., 3rd, Demarini, D. J., Shah, N. G., Wach, A., Brachat, A., . . . Pringle, J. R. (1998). Additional modules for versatile and economical PCR-based gene deletion and modification in Saccharomyces cerevisiae. *Yeast, 14*(10), 953-961. doi:10.1002/(sici)1097-0061(199807)14:10<953::aid-yea293>3.0.co;2-u

Maeda, K., Anand, K., Chiapparino, A., Kumar, A., Poletto, M., Kaksonen, M., & Gavin, A.

1033   C. (2013). Interactome map uncovers phosphatidylserine transport by oxysterol-

1034   binding proteins. *Nature, 501*(7466), 257-261. doi:10.1038/nature12430

1035 Maekawa, M., & Fairn, G. D. (2015). Complementary probes reveal that phosphatidylserine

1036   is required for the proper transbilayer distribution of cholesterol. *J Cell Sci, 128*(7),

1037   1422-1433. doi:10.1242/jcs.164715

1038 Manolson, M. F., Proteau, D., Preston, R. A., Stenbit, A., Roberts, B. T., Hoyt, M. A., . . .

1039   Jones, E. W. (1992). The VPH1 gene encodes a 95-kDa integral membrane

1040   polypeptide required for in vivo assembly and activity of the yeast vacuolar H(+)-

1041   ATPase. *J Biol Chem, 267*(20), 14294-14303.

1042 Marshansky, V., Rubinstein, J. L., & Grüber, G. (2014). Eukaryotic V-ATPase: novel

1043   structural findings and functional insights. *Biochim Biophys Acta, 1837*(6), 857-879.

1044   doi:10.1016/j.bbabio.2014.01.018

1045 Michell, R. H. (2008). Inositol derivatives: evolution and functions. *Nat Rev Mol Cell Biol,*

1046   *9*(2), 151-161. doi:10.1038/nrm2334

1047 Middel, V., Zhou, L., Takamiya, M., Beil, T., Shahid, M., Roostalu, U., . . . Strähle, U.

1048   (2016). Dysferlin-mediated phosphatidylserine sorting engages macrophages in

1049   sarcolemma repair. *Nat Commun, 7*, 12875. doi:10.1038/ncomms12875

1050 Mioka, T., Fujimura-Kamada, K., Mizugaki, N., Kishimoto, T., Sano, T., Nunome, H., . . .

1051   Tanaka, K. (2018). Phospholipid flippases and Sfk1p, a novel regulator of

1052   phospholipid asymmetry, contribute to low permeability of the plasma membrane. *Mol*

1053   *Biol Cell, 29*(10), 1203-1218. doi:10.1091/mbc.E17-04-0217

1054 Mizushima, N., Yoshimori, T., & Ohsumi, Y. (2011). The role of Atg proteins in

1055   autophagosome formation. *Annu Rev Cell Dev Biol, 27*, 107-132.

1056   doi:10.1146/annurev-cellbio-092910-154005

Morvan, J., de Craene, J. O., Rinaldi, B., Addis, V., Misslin, C., & Friant, S. (2015). Btn3

regulates the endosomal sorting function of the yeast Ent3 epsin, an adaptor for

SNARE proteins. *J Cell Sci, 128*(4), 706-716. doi:10.1242/jcs.159699

Nakanishi, H., de los Santos, P., & Neiman, A. M. (2004). Positive and negative regulation of

a SNARE protein by control of intracellular localization. *Mol Biol Cell, 15*(4), 1802-

1815. doi:10.1091/mbc.e03-11-0798

Nicolson, G. L. (2014). The Fluid-Mosaic Model of Membrane Structure: still relevant to

understanding the structure, function and dynamics of biological membranes after

more than 40 years. *Biochim Biophys Acta, 1838*(6), 1451-1466.

doi:10.1016/j.bbamem.2013.10.019

Nuoffer, C., Jenö, P., Conzelmann, A., & Riezman, H. (1991). Determinants for

glycophospholipid anchoring of the Saccharomyces cerevisiae GAS1 protein to the

plasma membrane. *Mol Cell Biol, 11*(1), 27-37. doi:10.1128/mcb.11.1.27

Nyholm, T. K. M., Jaikishan, S., Engberg, O., Hautala, V., & Slotte, J. P. (2019). The Affinity

of Sterols for Different Phospholipid Classes and Its Impact on Lateral Segregation.

Biophys J, 116(2), 296-307. doi:10.1016/j.bpj.2018.11.3135

Obara, K., Yamamoto, H., & Kihara, A. (2012). Membrane protein Rim21 plays a central role

in sensing ambient pH in Saccharomyces cerevisiae. *J Biol Chem, 287*(46), 38473-

38481. doi:10.1074/jbc.M112.394205

Oh, C. S., Toke, D. A., Mandala, S., & Martin, C. E. (1997). ELO2 and ELO3, homologues of

the Saccharomyces cerevisiae ELO1 gene, function in fatty acid elongation and are

required for sphingolipid formation. *J Biol Chem, 272*(28), 17376-17384.

doi:10.1074/jbc.272.28.17376

Omnus, D. J., Manford, A. G., Bader, J. M., Emr, S. D., & Stefan, C. J. (2016).

Phosphoinositide kinase signaling controls ER-PM cross-talk. *Mol Biol Cell, 27*(7),

1170-1180. doi:10.1091/mbc.E16-01-0002

Pan, X., Roberts, P., Chen, Y., Kvam, E., Shulga, N., Huang, K., . . . Goldfarb, D. S. (2000).

Nucleus-vacuole junctions in Saccharomyces cerevisiae are formed through the direct

interaction of Vac8p with Nvj1p. *Mol Biol Cell, 11*(7), 2445-2457.

doi:10.1091/mbc.11.7.2445

Plasencia, I., Norlén, L., & Bagatolli, L. A. (2007). Direct visualization of lipid domains in

human skin stratum corneum's lipid membranes: effect of pH and temperature.

Biophys J, 93(9), 3142-3155. doi:10.1529/biophysj.106.096164

Pomorski, T., Lombardi, R., Riezman, H., Devaux, P. F., van Meer, G., & Holthuis, J. C.

(2003). Drs2p-related P-type ATPases Dnf1p and Dnf2p are required for phospholipid

translocation across the yeast plasma membrane and serve a role in endocytosis. *Mol*

Biol Cell, 14(3), 1240-1254. doi:10.1091/mbc.e02-08-0501

Popov-Čeleketić, D., Bianchi, F., Ruiz, S. J., Meutiawati, F., & Poolman, B. (2016). A Plasma

Membrane Association Module in Yeast Amino Acid Transporters. *J Biol Chem,*

291(31), 16024-16037. doi:10.1074/jbc.M115.706770

Quon, E., Sere, Y. Y., Chauhan, N., Johansen, J., Sullivan, D. P., Dittman, J. S., . . . Menon,

A. K. (2018). Endoplasmic reticulum-plasma membrane contact sites integrate sterol

and phospholipid regulation. *PLoS Biol, 16*(5), e2003864.

doi:10.1371/journal.pbio.2003864

Risselada, H. J., & Marrink, S. J. (2008). The molecular face of lipid rafts in model

membranes. *Proc Natl Acad Sci U S A, 105*(45), 17367-17372.

doi:10.1073/pnas.0807527105

Robinson, J. S., Klionsky, D. J., Banta, L. M., & Emr, S. D. (1988). Protein sorting in

Saccharomyces cerevisiae: isolation of mutants defective in the delivery and

processing of multiple vacuolar hydrolases. *Mol Cell Biol, 8*(11), 4936-4948.

doi:10.1128/mcb.8.11.4936

Rosenwald, A. G., Rhodes, M. A., Van Valkenburgh, H., Palanivel, V., Chapman, G., Boman,

A., . . . Kahn, R. A. (2002). ARL1 and membrane traffic in Saccharomyces cerevisiae.

Yeast, 19(12), 1039-1056. doi:10.1002/yea.897

Rothman, J. H., Howald, I., & Stevens, T. H. (1989). Characterization of genes required for

protein sorting and vacuolar function in the yeast Saccharomyces cerevisiae. *Embo j,*

8(7), 2057-2065.

Rouser, G., Fkeischer, S., & Yamamoto, A. (1970). Two dimensional then layer

chromatographic separation of polar lipids and determination of phospholipids by

phosphorus analysis of spots. *Lipids, 5*(5), 494-496. doi:10.1007/bf02531316

Roy, A., & Levine, T. P. (2004). Multiple pools of phosphatidylinositol 4-phosphate detected

using the pleckstrin homology domain of Osh2p. *J Biol Chem, 279*(43), 44683-44689.

doi:10.1074/jbc.M401583200

Saheki, Y., & De Camilli, P. (2017). Endoplasmic Reticulum-Plasma Membrane Contact

Sites. *Annu Rev Biochem, 86*, 659-684. doi:10.1146/annurev-biochem-061516-044932

Saito, K., Fujimura-Kamada, K., Furuta, N., Kato, U., Umeda, M., & Tanaka, K. (2004).

Cdc50p, a protein required for polarized growth, associates with the Drs2p P-type

ATPase implicated in phospholipid translocation in Saccharomyces cerevisiae. *Mol*

Biol Cell, 15(7), 3418-3432. doi:10.1091/mbc.e03-11-0829

Sakane, H., Yamamoto, T., & Tanaka, K. (2006). The functional relationship between the

Cdc50p-Drs2p putative aminophospholipid translocase and the Arf GAP Gcs1p in

vesicle formation in the retrieval pathway from yeast early endosomes to the TGN.

Cell Struct Funct, 31(2), 87-108. doi:10.1247/csf.06021

Sambrook, J., Russell, DW. (2001). Molecular Cloning: A Laboratory Manual, Cold Spring

Harbor, NY: Cold Spring Harbor Laboratory Press.

Schmidt, O., & Teis, D. (2012). The ESCRT machinery. *Curr Biol, 22*(4), R116-120.

doi:10.1016/j.cub.2012.01.028

Schneiter, R., Brügger, B., Sandhoff, R., Zellnig, G., Leber, A., Lampl, M., . . . Kohlwein, S.

D. (1999). Electrospray ionization tandem mass spectrometry (ESI-MS/MS) analysis

of the lipid molecular species composition of yeast subcellular membranes reveals

acyl chain-based sorting/remodeling of distinct molecular species en route to the

plasma membrane. *J Cell Biol, 146*(4), 741-754. doi:10.1083/jcb.146.4.741

Schulz, T. A., & Creutz, C. E. (2004). The tricalbin C2 domains: lipid-binding properties of a

novel, synaptotagmin-like yeast protein family. *Biochemistry, 43*(13), 3987-3995.

doi:10.1021/bi036082w

Schäfer, L. V., de Jong, D. H., Holt, A., Rzepiela, A. J., de Vries, A. H., Poolman, B., . . .

Marrink, S. J. (2011). Lipid packing drives the segregation of transmembrane helices

into disordered lipid domains in model membranes. *Proc Natl Acad Sci U S A, 108*(4),

1343-1348. doi:10.1073/pnas.1009362108

Sengupta, P., Hammond, A., Holowka, D., & Baird, B. (2008). Structural determinants for

partitioning of lipids and proteins between coexisting fluid phases in giant plasma

membrane vesicles. *Biochim Biophys Acta, 1778*(1), 20-32.

doi:10.1016/j.bbamem.2007.08.028

Serrano, R., Kielland-Brandt, M. C., & Fink, G. R. (1986). Yeast plasma membrane ATPase

is essential for growth and has homology with (Na+ + K+), K+- and Ca2+-ATPases.

Nature, 319(6055), 689-693. doi:10.1038/319689a0

Silve, S., Dupuy, P. H., Labit-Lebouteiller, C., Kaghad, M., Chalon, P., Rahier, A., . . .

Loison, G. (1996). Emopamil-binding protein, a mammalian protein that binds a series

of structurally diverse neuroprotective agents, exhibits delta8-delta7 sterol isomerase

activity in yeast. *J Biol Chem, 271*(37), 22434-22440. doi:10.1074/jbc.271.37.22434

Singer, S. J., & Nicolson, G. L. (1972). The fluid mosaic model of the structure of cell

membranes. *Science, 175*(4023), 720-731. doi:10.1126/science.175.4023.720

Solanko, L. M., Sullivan, D. P., Sere, Y. Y., Szomek, M., Lunding, A., Solanko, K. A., . . .

Wüstner, D. (2018). Ergosterol is mainly located in the cytoplasmic leaflet of the yeast

plasma membrane. *Traffic, 19*(3), 198-214. doi:10.1111/tra.12545

Sychrová, H., Ramírez, J., & Peña, A. (1999). Involvement of Nha1 antiporter in regulation of

intracellular pH in Saccharomyces cerevisiae. *FEMS Microbiol Lett, 171*(2), 167-172.

doi:10.1111/j.1574-6968.1999.tb13428.x

Trimble, W. S., & Grinstein, S. (2015). Barriers to the free diffusion of proteins and lipids in

the plasma membrane. *J Cell Biol, 208*(3), 259-271. doi:10.1083/jcb.201410071

Tsuchiya, M., Hara, Y., Okuda, M., Itoh, K., Nishioka, R., Shiomi, A., . . . Umeda, M. (2018).

Cell surface flip-flop of phosphatidylserine is critical for PIEZO1-mediated myotube

formation. *Nat Commun, 9*(1), 2049. doi:10.1038/s41467-018-04436-w

Tsuji, T., Fujimoto, M., Tatematsu, T., Cheng, J., Orii, M., Takatori, S., & Fujimoto, T.

(2017). Niemann-Pick type C proteins promote microautophagy by expanding raft-like

membrane domains in the yeast vacuole. *Elife, 6*. doi:10.7554/eLife.25960

Uchida, Y., Hasegawa, J., Chinnapen, D., Inoue, T., Okazaki, S., Kato, R., . . . Arai, H.

(2011). Intracellular phosphatidylserine is essential for retrograde membrane traffic

through endosomes. *Proc Natl Acad Sci U S A, 108*(38), 15846-15851.

doi:10.1073/pnas.1109101108

van Leeuwen, J., Pons, C., Mellor, J. C., Yamaguchi, T. N., Friesen, H., Koschwanez, J., . . .

Boone, C. (2016). Exploring genetic suppression interactions on a global scale.

Science, 354(6312). doi:10.1126/science.aag0839

Vater, C. A., Raymond, C. K., Ekena, K., Howald-Stevenson, I., & Stevens, T. H. (1992). The

VPS1 protein, a homolog of dynamin required for vacuolar protein sorting in

Saccharomyces cerevisiae, is a GTPase with two functionally separable domains. *J*

Cell Biol, 119(4), 773-786. doi:10.1083/jcb.119.4.773

Veatch, S. L., & Keller, S. L. (2002). Organization in lipid membranes containing cholesterol. *Physical review letters*, 89(26), 268101. doi.org/10.1103/PhysRevLett.89.268101

Veatch, S. L., & Keller, S. L. (2003). Separation of liquid phases in giant vesicles of ternary mixtures of phospholipids and cholesterol. *Biophys J, 85*(5), 3074-3083. doi:10.1016/s0006-3495(03)74726-2

Wang, C. W., Miao, Y. H., & Chang, Y. S. (2014). A sterol-enriched vacuolar microdomain mediates stationary phase lipophagy in budding yeast. *J Cell Biol, 206*(3), 357-366. doi:10.1083/jcb.201404115

Watt, S. A., Kular, G., Fleming, I. N., Downes, C. P., & Lucocq, J. M. (2002). Subcellular localization of phosphatidylinositol 4,5-bisphosphate using the pleckstrin homology domain of phospholipase C delta1. *Biochem J, 363*(Pt 3), 657-666. doi:10.1042/0264-6021:3630657

Yamamoto, T., Fujimura-Kamada, K., Shioji, E., Suzuki, R., & Tanaka, K. (2017). Cfs1p, a Novel Membrane Protein in the PQ-Loop Family, Is Involved in Phospholipid Flippase Functions in Yeast. *G3 (Bethesda), 7*(1), 179-192. doi:10.1534/g3.116.035238

Yeung, B. G., Phan, H. L., & Payne, G. S. (1999). Adaptor complex-independent clathrin function in yeast. *Mol Biol Cell, 10*(11), 3643-3659. doi:10.1091/mbc.10.11.3643

Yeung, T., Gilbert, G. E., Shi, J., Silvius, J., Kapus, A., & Grinstein, S. (2008). Membrane phosphatidylserine regulates surface charge and protein localization. *Science, 319*(5860), 210-213. doi:10.1126/science.1152066

Young, B. P., Shin, J. J., Orij, R., Chao, J. T., Li, S. C., Guan, X. L., . . . Loewen, C. J. (2010). Phosphatidic acid is a pH biosensor that links membrane biogenesis to metabolism. *Science, 329*(5995), 1085-1088. doi:10.1126/science.1191026