Jasmonic Acid coordinates with Light, Glucose and Auxin signalling in

Regulating Branching Angle of *Arabidopsis* Lateral Roots

Manvi Sharma[a,1], Mohan Sharma[a,1,#], Muhammed Jamsheer K[a,b] & Ashverya Laxmi[a*]

[a]National Institute of Plant Genome Research, Aruna Asaf Ali Marg, New Delhi-110067,

India

[b]Present address: Department of Biochemistry, University of Lausanne, Lausanne,

Switzerland

[1]The authors contributed equally to this work.

*Corresponding author details:

Ashverya Laxmi

Lab 203, National Institute of Plant Genome Research, Aruna Asaf Ali Marg, New Delhi-

110067, India. Tel: 91-11-26741612, 14, 17 Ext. 180. Email: ashverya_laxmi@nipgr.ac.in

Keywords: Jasmonate; phytohormones, TOR kinase, lateral root development, crosstalk

Abstract

The role of jasmonates (JAs) in primary root growth and development and in plant response

to external stimuli is already known. However, its role in lateral root (LR) development

remains to be explored. Our work identified methyl jasmonate (MeJA) as a key

phytohormone in determining the branching angle of *Arabidopsis* LRs. MeJA inclines the

LRs to a more vertical orientation, which was dependent on the canonical *JAR1-COI1-MYC2,*

3, 4 signalling. Our work also highlights the dual roles of light in governing LR angle. Light

signalling enhances JA biosynthesis, leading to erect root architecture; whereas, glucose

(Glc) induces wider branching angles. Combining physiological and molecular assays, we

revealed that Glc antagonizes the MeJA response via TARGET OF RAPAMYCIN (TOR)

signalling. Moreover, physiological assays using auxin mutants, MYC2-mediated

transcriptional activation of *LAZY2, LAZY4* and auxin biosynthetic gene *CYP79B2*, and

asymmetric distribution of *DR5::GFP* and *PIN2::GFP* pinpointed the role of an intact auxin

mechanism required by MeJA for vertical growth of LRs. We also demonstrated that light

perception and signalling are indispensable for inducing vertical angles by MeJA. Thus, our

investigation highlights antagonism between light and Glc signalling and how they interact

with JA-auxin signals to optimize the branching angle of LRs.

Introduction

The plant root system is highly plastic in nature and is influenced by exogenous and

endogenous cues to modify its architecture. Because of their sedentary nature, plants are

limited to their immediate surroundings for acquisition of water and nutrients (Malamy 2005;

Cuesta, Wabnik & Benková 2013) and therefore, depend heavily on their root system. While

the primary root maintains a nearly vertical orientation (orthogravitropism), lateral roots

(LRs) form a non-vertical growth orientation, away from the main root axis, establishing a

distinct gravitropic set-point angle (GSA) (Digby & Firn 1995). After their emergence, LRs

adopt a perpendicular angle with respect to gravity, but as the LRs continues to develop, their

GSA changes over time and exhibit de novo formation of an elongation zone (Kiss, Miller,

Ogden & Roth 2002; Rosquete *et al.* 2013). The transient expression of PIN3 in LR

columella cells defines asymmetric auxin distribution as well as differential elongation rates

in stage II LRs (Rosquete *et al.* 2013), leading to the establishment of primary GSA of LR

(Guyomarc'h *et al.* 2012) and is exemplified by an asymmetric growth towards gravity at a

slower rate than the primary roots (Rosquete *et al.* 2013). As a result, the primary GSA of

LRs is established. Subsequently, PIN3 repression in columella cells of stage III (SIII) LRs

coincides with symmetric elongation, maintaining this primary GSA. Thus, the primary growth direction of LRs is of paramount significance, as it directs plant anchorage and uptake of water and nutrients, ultimately contributing to root system architecture.

In plants, phytohormones have profound effects on determining GSA. For example, as mentioned above, recent findings have identified the fundamental role of auxin in governing the GSA of shoot branches and LRs of *Arabidopsis thaliana* and other vascular plants, such as rice and beans (Rosquete *et al.* 2013; Rosquete, Waidmann & Kleine-Vehn 2018; Roychoudhry, Del Bianco, Kieffer & Kepinski 2013; Roychoudhry *et al.* 2017, 2019). Recently, Waidmann et al (2019) showed that CK signalling leads to reduced growth at the upper flank of the LR, thereby working as an antigravitropic component and hence preventing downward growth. In this way, a CK-dependent mechanism allows the root system to radially explore its surroundings (Waidmann *et al.* 2019). Jasmonic acid (JA) and methyl jasmonate (MeJA), collectively called jasmonates (JAs), are cyclopentenone compounds that are known to primarily modulate a number of vital physiological processes like root development, gravitropism, reproductive development, senescence; wound responses; and defence responses against pathogens and insects (Wasternack & Hause 2013). Findings of JA on regulation of main root gravitropism has already been reported (Gutjahr *et al.* 2005). It was found that the JA receptor CORONATINE-INSENSITIVE1 (COI1) is required for JA-mediated LR formation, positioning, and emergence of root bends in *Arabidopsis* (Raya-González, Pelagio-Flores & López-Bucio 2012). Additionally, investigations in rice have revealed the role of MeJA in modifying lamina joint inclination (Gan *et al.* 2015). Nonetheless, the role of JAs in steering LR GSA has not yet been demonstrated.

Besides phytohormones, factors such as temperature, nutrient status, and light play a vital role in governing branching angles (Digby & Firn 2002; Bai, Murali, Barber & Wolverton 2013; Trachsel, Kaeppler, Brown & Lynch 2013; Roychoudhry *et al.* 2017). Previous reports have implied the role of sucrose (Suc) in influencing the gravitropic behaviour of stems in *Paspalum vaginatum* (Willemoes, Beltrano & Montalbi 1988). Sugars such as Suc and glucose (Glc) are not only important nutrients, but also act as key signalling molecules (Ramon, Rolland & Sheen 2008; Eveland & Jackson 2012; Li & Sheen 2016). A recent report suggests that light and Suc act antagonistically to influence root length, but additively affect root hair emergence and elongation (García-González, Lacek & Retzer 2021). OsERF2 has been shown to influence rice root growth by fine tuning the genes involved in ABA and ethylene signalling pathways and sucrose metabolism (Xiao *et al.* 2016). Sucrose also modulates auxin metabolism, transport and signalling to regulate elongation growth responses (Lilley, Gee, Sairanen, Ljung & Nemhauser 2012; Sairanen *et al.* 2013; Stokes, Chattopadhyay, Wilkins, Nambara & Campbell 2013; Lastdrager, Hanson & Smeekens 2014). In *Arabidopsis*, there are three distinct Glc signalling pathways: (1) the HEXOKINASE1 (HXK1)-dependent pathway that is governed by HXK1-mediated signalling function, in which Glc is possibly sensed by a module independent of the catalytic activity of HXK1; (2) the G-protein-coupled receptor signalling by REGULATOR OF G-PROTEIN SIGNALLING 1 (RGS1) and GPA1 which are implicated in sensing extracellular glucose and signalling through THF1, located in the plastids (Huang *et al.* 2006; Urano *et al.* 2012a); and (3) a glycolysis-dependent pathway that works through the antagonistic interaction between SUCROSE NONFERMENTING RELATED KINASE 1 (SnRK1) and TARGET OF RAPAMYCIN (TOR) (Baena-González, Rolland, Thevelein & Sheen 2007; Baena-González 2010; Xiong & Sheen 2015; Song *et al.* 2017). Previous studies have suggested the involvement of Glc in various aspects of early seedling development (Mishra, Singh,

Aggrawal & Laxmi 2009; Kircher & Schopfer 2012; Yuan, Xu, Zhang, Guo & Lu 2014). Glc

and phytohormones have been extensively shown to interact with each other to enhance plant

fitness. The interplay of Glc with various phytohormones has been shown to modulate root

directional responses in *Arabidopsis* seedlings (Singh, Gupta & Laxmi 2014a b). However,

very few reports link JA and sugar signalling (Song *et al.* 2017; Vleesschauwer *et al.* 2017;

Guo *et al.* 2018). Based on physiological and pharmacological studies, an intimate cross-talk

occurs between TOR and JA signalling pathways at multiple levels of JA signal transduction

(Vleesschauwer *et al.* 2017), but the JA-Glc signal crosstalk that regulates branching angle of

Arabidopsis LRs still remain obscure.

Despite the obvious importance of directional LR growth for plant propagation; the role of

factors other than CK and auxin are largely unexplored. In this study, we have characterized

the interaction of JA and Glc in the regulation of branching angle of LRs in *Arabidopsis*.

Using genetic, physiological, molecular and cell biological approaches, we show that two

opposing signals (JA and Glc) counteract each other in order to set directional LR growth.

Additionally, knowledge gained from this study can expand our insight on root architecture

establishment and may contribute to the regulation of GSA in the field of plant breeding.

Materials and methods

Plant materials. *Arabidopsis thaliana* ecotype*s* of Col-0, Ws and Ler were used as wild-

type controls. The following seed stocks were obtained from the *Arabidopsis* Biological

Resource Center (ABRC) at Ohio State University (http://www.*Arabidopsis*.org/abrc/):

DR5::GFP; *phyA-201* (CS6291); *phyB-5* (CS6213); *phyA201B5* (CS6224); *hy5-1* (CS71);

rgs1-1 (CS6537); *rgs1-2* (CS6538); *gpa1-3* (CS6533); *gpa1-4* (CS6534); *tir1-1* (CS3798);

axr1-3 (CS3075); *aux1-7* (CS3074); *axr2-1* (CS3077); *eir1-1* (CS8058); *myc2/jin1-9*

(SALK_017005C); *coi1* (SALK_095916C); *jar1-11* (CS67935); *jaz1* (SALK_011957C);

jaz2 (SALK_012782C); *jaz4* (SALK_141628C); *jaz6* (SALK_017531); *sweet11sweet12*

(CS68845). Following lines were obtained from the original published source as:

myc2myc3myc4 (Schweizer et al., 2013); *mdr1-1* (Noh et al., 2001); *lax3* (Swarup et al.,

2008); *pin3-4, pin4-3* (Friml et al., 2002b); *pin7-2* (Friml et al., 2003); *tor35-7 RNAi* (Deprost

et al. 2007); *p35S::Jas9-N7-VENUS* (Larrieu et al., 2015); *35S::MYC2::GFP* (Jung et al.,

2105); *PIN2::PIN2-eGFP* (NASC); *PIN3::PIN3-eGFP* (NASC). All mutant lines were in the

Col-0 background except the following: *mdr1-1* was derived from Ws background; and *hy5-*

1, phyA-201, phyB-5, phyA201B5 were in the Ler background.

Growth conditions. Seeds were surface sterilized and stratified at 4°C for 48 hours. The

imbibed seeds were grown vertically on square petri dishes containing 1/2 Murashige and

Skoog (MS) medium supplemented with 1% sucrose (29.13 mM; w/v) and solidified with

0.8% agar (w/v). Seed germination and plant growth were carried out in climate-controlled

growth rooms under long day conditions (16 hr light and 8 hr darkness), with 22°C ± 2°C

temperature and 60 µmol/sec/m^2 light intensity. To study branching angle of LRs, five-day-

old MS grown seedlings were transferred to hormone/inhibitor/sugar treatment media with

their root tips marked and grown vertically under above mentioned growth conditions, unless

otherwise stated. All chemicals were purchased from Sigma (St. Louis, MO, USA) except

specified otherwise. MeJA was prepared as 50 mM stock solution in 100% (v/v) ethanol.

Epibrassinolide was prepared as 10^{-2} M stock solution in 50% (v/v) ethanol. The following

were prepared as 10^{-2} M stock solutions in dimethyl sulfoxide: NPA, BAP, ABA, and GA3.

ACC was prepared as a sterile 10^{-2} M aqueous stock solution.

Physiological analyses. Seedlings were grown on 1/2 MS, 0.8% agar, and 1%Suc-containing

medium for five days. They were then transferred to 1/2 MS medium, supplemented with

different combinations of Glc [0.5% (27.75 mM; w/v) and 3% (166.52 mM; w/v)] and MeJA

(10 µM) or 1/2 MS medium supplemented with 1% sucrose and MeJA (1 µM, 5 µM, 10 µM)

or combinations of Glc (1%) and Man (1%, 2%, 3%, 4%) with and without 10 µM MeJA and

their root tips were marked. Dark experiments were carried out by transferring light grown,

five-day-old WT and *myc2myc3myc4* seedlings to treatment media in light for 24 hours,

followed by complete darkness for the next 5 days. For measurement of branching angle, the

angle formed between each LR and primary root was measured. The average branching angle

was determined by summing the angle formed by each LR of all the seedlings divided by the

total number of LRs. Also, the angular response was measured by distributing the branching

angle in three categories viz. <40°, 40°-70°, >70° and expressed as percentage of LR. ImageJ

program from NIH was used to quantify the branching angle as well epidermal cell length of

LRs (http://rsb.info.nih.gov/ij/). Digital images were captured from Nikon Cool pix camera

on 12th day of seedling growth.

Laser Confocal Scanning Microscopy. To observe auxin distribution, PIN2 and PIN3

expression in stage II LRs, five-day-old light grown *DR5::GFP*, *PIN2::PIN2-eGFP* and

PIN3::PIN3-eGFP expressing seedlings were transferred to 1/2 MS and 1/2 MS + 10 µM

MeJA for 6-7 days for imaging. GFP fluorescence was imaged under a Leica TCS SP8

AOBS Laser Confocal Scanning Microscope (Leica Microsystems). To image GFP, the 488

nm line of the argon laser was used for excitation and emission was detected at 520 nm. The

laser, pinhole and gain settings of the confocal microscope were kept identical among

different treatments. Images were assembled using Photoshop (Adobe Systems). For optical

sectioning under SP8 microscope, Z-stacks were scanned for 6 sections every 4 µm in

thickness and maximum projections were generated. At least two biological replicates, with

each replicate having 15 seedlings, were performed for all the experiments.

Gene Expression Analysis. For gene expression analysis, RT-qPCR was performed.

Imbided seeds of WT, *myc2myc3myc4*, and *hy5-1* were sown on 1/2 MS medium

supplemented with 1% (w/v) Suc and 0.8% (w/v) agar and grown vertically in culture room

conditions. Five-day-old light grown seedlings of WT and *myc2myc3myc4* were harvested

and stored in -80°C. WT and *hy5-1* were grown in light (long day regime) and continuous

dark for 5 days. For dark to light transition, WT and *hy5-1* were grown in continuous dark for

5 days followed by light treatment for 6 hours. To check, JA related gene expression, 5-d–old

Col-0 seedlings were first starved under 0% Glc for 24 h in the dark. After starvation,

seedlings were treated without or with Glc and a combination of 10 µM MeJA. Afterwards,

whole seedlings were flash frozen in liquid nitrogen and stored at -80°C. Total RNA was

isolated from frozen tissue using the RNeasy Plant Mini Kit (Qiagen) following the

manufacturer's protocol. RNA was quantified and tested for quality before it was used for

subsequent analyses. First-strand cDNA was synthesized by reverse transcription using 2 µg

of total RNA with a high-capacity cDNA Reverse Transcription Kit (Applied Biosystems).

Primers were designed by identifying a sequence stretch with a unique sequence. All

candidate gene primers were designed using the software Primer Express (version 3.0;

Applied Biosystems). An ABI Prism 7900 HP fast real-time PCR system (Applied

Biosystems) was used. For the normalization of variance among samples, UBIQUITIN10

(UBQ10) was used as a reference control. The fold-change for each candidate gene in

different experimental conditions was determined using the quantitative ΔΔCT method. All

primers used are mentioned in the Supplemental Table S1. Gene expression analysis was

performed 3 times unless otherwise stated.

Chromatin Immunoprecipitation. ChIP assays were performed by following the protocol

of Saleh et al. (2008) with minor modifications. *Arabidopsis* Col-0 and *35S::MYC2::GFP*

seedlings were grown for seven days in 1/2 MS containing 1%Suc and used for the ChIP

assay. Briefly, 1 gram (g) tissue of each sample was cross-linked with 1% formaldehyde to

fix protein-DNA complexes. The samples was crushed in liquid N2 and homogenized in

nuclei isolation and nuclei lysis buffers followed by sonication. Sonication of chromatin was

done in 4°C water sonicator (Diagenode Bioruptor Plus). Sonicated samples were first

precleared with Protein A Agarose beads (Millipore #16-157) and then mixed with

antibodies. Antibodies against GFP (Cat. no. ab290) were purchased from Abcam.

Approximately 200 bp upstream promoter region of *LAZY2, LAZY4* and *CYP79B2* containing

the G- and E-box elements which are known binding sites of MYC2 was enriched in the ChIP

assay. All primers used in the ChIP assays are mentioned in the Supplementary Table S1.

Protein extraction and immunoblot assay. Extraction of soluble proteins was performed

on seven day old seedlings of p35S::Jas9-VENUS. Seven day old light grown seedlings of

p35S::Jas9-N7-VENUS were treated in sugar-free liquid 1/2 MS in dark for 24 h to deplete

internal sugars. This was followed by treating the seedlings with 0% (w/v) Glc and 3%

(w/v) Glc containing 1/2 MS liquid medium for 3 hours. Following Glc treatment, 10 μM

MeJA or 10 μM AZD-8055 or both were added to all seedlings for 3 hours in dark.

Seedlings were harvested, frozen in liquid nitrogen and then ground in pre chilled mortar

pestle using liquid nitrogen. The powder was resuspended in cold extraction buffer (137

mM of NaCl, 2.7 mM of KCl, 4.3 mM of Na_2HPO_4, 1.47 mM of KH_2PO_4, 10% glycerol

[1.1 M], and 1 mM phenylmethylsulfonyl fluoride [PMSF]) supplemented with plant protease

inhibitor cocktail (Sigma-Aldrich, http://www.sigmaald rich.com/). This was followed by two

rounds of centrifugation (15 min, 13,000 rpm at 4 °C) to remove cell debris. The

samples were boiled for 10 min at 95 °C before being loaded onto a gel (40 μl; 80-100

μg per lane). SDS–PAGE was performed on 10% polyacrylamide gel. After transfer onto a

PVDF membrane, protein amounts in each lane were checked using Ponceau staining

(0.1%, 1.5 mM). Immunoblots were detected using a primary rabbit polyclonal anti-GFP

antibody (ab290, Abcam, diluted 1:5,000), primary rabbit polyclonal anti HSP90-2

antibody (diluted 1:5,000) and a secondary anti-rabbit IgG-HRP (diluted 1:10,000).

Proteins were visualized using the enhanced chemiluminescence kit.

Microarray analysis. For microarray of *Arabidopsis* WT seedlings under Glc and MeJA

treatment, 5-d–old seedlings were first starved under 0% Glc for 24 h in the dark. After

starvation, seedlings were treated with and without Glc and a combination of 10 μM

MeJA. Harvested samples were outsourced for transcriptome profiling. The data were

analyzed by the Transcriptome Analysis Console (v3.0; Affymetrix) with default

parameters. Two biological replicates were used for the microarray experiment.

Statistical analyses. All physiological experiments yielding similar results were repeated as

mentioned in the figure legends, in which each experiment was considered as an independent

biological replicate consisting of at least 25 seedlings. Immunoblot assays were performed 4

times unless otherwise stated. ChIP assays were performed as mentioned in the figure

legends. Statistical differences between control/treatment and WT/mutant pair were analyzed

using Student's *t* test with paired two-tailed distribution. *P*-value cutoff was taken at $P < 0.05$.

All data were managed and analyzed using Microsoft Excel. The graphs were made using

Microsoft Excel and Instant Clue. End point physiological analyses were carried out on 12[th]

day of seedling growth.

Accession Numbers

Arabidopsis Genome Initiative locus identifiers for the genes mentioned in this article are:

phyA-201, AT1G09570; *phyB-5,* AT2G18790; *hy5-1,* AT5G11260; *tor35-7 RNAi*

AT1G50030; *rgs1-1,* AT3G26090; *rgs1-2* AT3G26090; *gpa1-3* AT2G26300; *gpa1-4,*

AT2G26300; *tir1-1,* AT3G62980; *axr1-3,* AT1G05180; *aux1-7,* AT2G38120; *axr2-1,*

AT3G23050; *eir1-1,* AT5G57090; *myc2/jin1-9*, AT1G32640; *coi1,* AT2G39940; *jar1-11*,

AT2G46370; *jaz1*, AT1G19180; *jaz2,* AT1G74950; *jaz4,* AT1G48500; *jaz6*, AT1G72450;

mdr1-1, AT3G28860; *lax3,* AT1G77690; *pin4-3,* AT2G01420; *pin7-2,* AT1G23080; *pin3-4*,

AT1G70940; *sweet11swet12*, (AT3G48740, AT5G23660).

Results

MeJA modulates branching angle of *Arabidopsis* roots. To study the role of MeJA in

regulating growth angles, the branching angle of *Arabidopsis* WT LRs was analysed by two

methods: 1) calculating the angle formed by all LRs of wild-type (WT) seedlings with respect

to the main root and averaging the values, and 2) distributing the angles of LRs in three

categories (<40°, 40°-70° and >70°). We observed that MeJA treatment decreases the

branching angle in a dose-dependent manner (Fig. 1A-C). LRs of WT showed an average

branching angle of 67.02° on control media, but a more vertical angle of 40.21° when grown

on the high MeJA concentration (10 µM) (Fig. 1B). Approximately 63% of LRs adopted

angles between 0-40° and very few with angles >70° when grown on 1/2 MS supplemented

with 10 µM MeJA (Fig. 1C). Since the effect of auxin and CK in governing GSA is already

explored (Rosquete *et al.* 2013, 2018; Roychoudhry *et al.* 2013, 2019; Waidmann *et al.*

2019), we wanted to identify the role of other hormones in modulating this response. The

Arabidopsis WT seedlings did not display any changes in branching angle when grown in

brassinosteroids (BR), gibberellic acid (GA$_3$), 1-aminocyclopropane-1-carboxylic acid (ACC)

or abscisic acid (ABA) (Fig. 1D). Altogether, our data indicates that MeJA impacts the

angular growth of LRs.

MeJA decreases branching angle in a SCF^{COI1} and MYC2, 3, 4 dependent pathway. In

order to elucidate how JA biosynthesis and signalling are involved, several mutants defective

in the JA-biosynthesis and signalling pathways were assessed. JA perception by COI1 is the

first committed step of JA signalling. Before perception, JA is converted to its biologically

active form, jasmonoyl-isoleucine (JA-Ile), by the enzyme JASMONATE RESISTANT 1 (JAR1) (Staswick & Tiryaki 2004; Staswick 2009). Neither *jar1-11* nor *coi1* responded to MeJA treatment, and both exhibited an overall horizontal orientation of LRs (Fig. 1E-G). Also, very few LRs of these mutants adopted angles <40°, as compared to WT (Fig. 1E-G). The average branching angles of *jar1-11* and *coi1* indicated a complete resistance to the MeJA treatment (Fig. S1A). MYC2 is a bHLH transcription factor that is considered the master regulator of jasmonate and light responses (Lorenzo, Chico, Sá Nchez-Serrano & Solano; Dombrecht *et al.* 2007). Therefore, we assessed whether JA-dependent transcription factors indeed have an impact on LR growth in the WT. The *jasmonate insensitive 1 (jin1-9),* a mutant of *MYC2,* was less responsive to MeJA treatment as compared to WT, as its LR showed broader angles (Fig. 1E, H; Fig. S1B). Also, *myc2myc3myc4* displayed diminished sensitivity, as a large percentage of LRs acquired angles >70° as compared to the single mutant and WT (Fig. 1E-I; Fig. S1B), suggesting that *MYC3* and *MYC4* act additively with *MYC2* in regulating this response. T-DNA insertion lines of *JASMONATE-ZIM-DOMAIN* (*JAZ1, JAZ2, JAZ4,* and *JAZ6*) did not show any major changes to increasing MeJA doses which can be attributed to redundancy of *JAZ* genes (Fig. S1C).

Since our in vitro study allowed only two-dimensional analysis of root growth, we further assessed angular growth of LRs in three dimensional systems. To allow three dimensional root expansion in vitro, we cultivated twenty-days-old WT, *jar1-11* and *myc2myc3myc4* plants in cylindrical tubes containing 1/2 MS with 1% phytagel, and found broader LR angles when compared to WT (Fig. S2A and B), suggesting that our two-dimensional in vitro screen was highly suitable for identifying mutants with diverging GSA values. Moreover, these results suggest that the defects in endogenous JA biosynthesis and signalling machinery leads to broader LR angle. Altogether, this set of data confirms that JA signalling utilizing receptor COI1 and TF MYC2 regulates angular growth of LRs.

Glc production via photosynthesis negatively influences MeJA-modulated branching angle. Previously, Digby and Firn (2002) demonstrated that light effects on GSA of organs can be brought about both via the action of photosynthesis and via phytochrome reception and signalling (Digby & Firn 2002). Previous reports have established the role of Glc in primary root gravitropism in *Arabidopsis* (Singh *et al.* 2014a, b). In order to explore the role of Glc and its interaction with MeJA in altering branching angles, the WT seedlings were treated with MeJA (10 µM) in combination with different concentrations of Glc (0.5%Glc; 28 mM and 3%Glc; 167 mM). The 3% concentration of Glc widened the branching angle as compared to the 0.5% concentration (Fig. 2A-B; Fig. S3). The LRs attained more vertical orientation when treated with a combination of 0.5%Glc+MeJA. However, when both MeJA and 3%Glc were added together, MeJA still showed a strong effect of narrowing root angle in the presence of 3% Glc, but the overall percentage of root angles did not reach the same level as they do in the 0.5% treatment (Fig. 2A-B; Fig. S3A). The above results suggest an antagonistic interaction between Glc and MeJA in controlling this response. We also used 5%Glc to check whether higher Glc concentration causes more widening of branching angle. However, we did not observe any significant changes in the LR angle between 0.5%Glc and 5%Glc, thereby suggesting that 3%Glc is the optimal concentration for plant growth that widens LR angle (Fig. S3B-C). Also, higher Glc concentrations is shown to cause developmental arrest in WT seedlings such as anthocyanin accumulation, less LR production and shorter root length (Moore *et al.* 2003) and therefore is considered inhibitory for overall growth and development of plants. Next, we wanted to assess whether the observed phenotype is due to changes in the osmoticum of the growth media or specific to Glc. To test this, we used mannitol (Man), a non-metabolizable sugar-alcohol that initiates very few LRs (Gupta, Singh & Laxmi 2015). Hence, to induce LR production, a small amount of Glc (1%) was also combined with increasing concentrations of Man (1%, 2%, 3% and 4%). We

observed that the effects previously induced by the supply of Glc were not produced by Man.

Increased doses of Man induced a vertically oriented LRs as compared to Glc (Fig. 2C).

Thus, osmotic changes do have an effect on branch angle, however it appears to be the

opposite effect than that of Glc. SUGARS WILL EVENTUALLY BE EXPORTED

TRANSPORTERS (SWEET) are bidirectional sugar uniporters that are capable of

transporting Suc, Glc and Fruc (Chen *et al.* 2010). In order to test whether changes in

endogenous sugar level in root alters the LR angle, we used the *sweet11sweet12* mutants

defective in sugar transport from shoot to root. For this, we transferred 5-day-old 1/2 MS

grown seedlings on low (0.5%) and moderate (1%) Glc concentration and observed that

sweet11sweet12 showed steeper branching angles as compared to WT in both Glc

concentrations (Fig. 2D-E). This result indicates that depletion of endogenous Glc in root

alters the LR branching angle suggesting the role of Glc in inducing wider branching angle in

Arabidopsis.

Next, to understand the crosstalk between Glc and JA signalling in influencing LR angle at

the molecular level, we first examined the protein level of JAZ9, a negative regulator of JA

signalling, in the presence and absence of Glc. For this, we used Jas9-VENUS, a fluorescent

marker widely used for the perception of bioactive JA, showing dynamic changes in JA

levels (Larrieu *et al.* 2015). As shown in Figure 2F-G, we observed very low levels of Jas9-

VENUS protein in seedlings treated with 0%Glc and 0%Glc+MeJA. Interestingly, as the

concentration of Glc was increased from 0 to 0.5% and 3%, Jas9-VENUS fusion protein was

accumulated at higher levels (Fig. 2F-G; Fig. S4A), which was found to be degraded when

subjected to MeJA (3%Glc+MeJA). We also found that even a low amount of Glc was

sufficient for MeJA mediated degradation of Jas9-VENUS (Fig. 2F-G; Fig. S4A). To test

whether MeJA can be perceived properly in the absence of sugar in the media, we compared

the expression of various JA-related genes (*JAZ1, JAZ2, JAZ3, JAZ6, JAZ9, AOS, ERF1,*

ORA59 and *MYC2*) in WT seedlings subjected to MeJA treatment without Glc using RT-

qPCR and found out that these genes were upregulated in MeJA treatment even without Glc

(Fig. S4B). This suggests that MeJA is actively perceived within the system even in the

absence of sugar. (Fig. 2F-G; Fig. S4A). Thus, all these results suggest that Glc promotes the

accumulation as well as MeJA-mediated degradation of Jas9-VENUS protein.

Glc mediated TOR-kinase antagonizes JA response. To further understand how the

antagonistic interaction of Glc and MeJA occurs at the molecular level, we investigated the

involvement of different components of Glc signal transduction in this response. Since TOR

is a master regulator of Glc and energy signalling cascades (Sheen 2014) and that it regulates

photosynthesis and phytohormone signalling pathways including JA signalling pathway

(Dong *et al.* 2015), we wanted to assess the role of TOR-kinase in this response and for this

we used *tor35-7 RNAi* line (Deprost *et al.* 2007). The RNAi line showed decrease in

branching angle of LRs in response to independent and combined treatments of Glc and

MeJA, as compared to WT seedlings (Fig. 2H-I; Fig. S5A). The *REGULATOR OF G-*

PROTEIN SIGNALLING (*RGS1*)-dependent Glc pathway involves a G-protein signalling

component, wherein RGS1 possibly acts as a Glc sensor (Urano *et al.* 2012b). To elucidate

the role of *RGS1*-dependent Glc signalling components, Glc/JA regulation of branching

angles in the *RGS1*-dependent signalling mutants *rgs1-1*, *rgs1-2*, *gpa1-3,* and *gpa1-4*. The

RGS1-dependent signalling mutants showed no major change in LR angle as compared with

their respective WT (Fig. S5B-C), suggesting that Glc does not utilise this pathway to

regulate LR angle. In order to further substantiate the role of TOR signalling in this response,

we treated Jas9-VENUS with Glc, alone and in combinations with MeJA and AZD-8055 (a

potent inhibitor of TOR kinase activity) (Montané & Menand 2013). Intriguingly, we

observed that Glc mediated accumulation of Jas9-VENUS was perturbed in seedlings treated

with AZD (Fig. 2J-K; Fig. S6). Altogether, a TOR-dependent energy signalling mechanism

is involved in the regulation of branching angle, and any perturbation to the mechanism leads

to an altered JA response.

To further dissect the relationship between JA and Glc signalling at the whole genome

transcriptome level, microarray analysis was employed. Most of the core JA signalling genes

were down-regulated in the presence of 3%Glc and 3%Glc+MeJA when compared with

0%Glc+MeJA (Fig. S7). The microarray analysis thus, falls in agreement with our

physiological and molecular data, confirming that antagonism operates between JA and Glc

signalling pathways in regulating LR angle.

Auxin transport and signalling lies downstream to MeJA-mediated modulation of

branching angle. Recent reports have shed light on the role of PIN auxin efflux activity and

the TIR1/AFB-Aux/IAA-ARF-dependent auxin signalling module for the establishment of

GSA in young LRs (Rosquete *et al.* 2013, 2018; Roychoudhry *et al.* 2013, 2019). We,

therefore, assessed the involvement of auxin transport and signalling to govern this

physiological response. For this, the polar auxin transport inhibitor 1-N-naphthylphthalamic

acid (NPA) was applied to the WT seedlings. NPA increased the Glc-induced branching

angle (Fig. 3A). When applied in combination, NPA inhibited the MeJA response (Fig. 3A).

To further investigate whether a functional auxin transport machinery is needed for MeJA to

reduce branching angles, auxin transport defective mutants were examined for changes in

branching angle. The auxin influx-defective mutant *auxin resistant 1 (aux1-7)* showed

resistance to MeJA treatment and displayed expansive branching angles as compared to WT,

with a greater percentage of LRs exhibiting angles >70° (Fig. 3B-C; Fig. S8A). The *lax3 (like*

aux1 3) mutant showed WT-like responses to MeJA treatment (Fig. S8B). Auxin-efflux

defective mutants such as *ethylene insensitive root 1* (*eir1-1*), a mutant of *PIN2* and *multiple*

drug resistance 1 (*mdr1-1*), a mutant of *PGP1,* exhibited LRs with significantly broader

angles compared with WT in the presence of Glc and in combination with MeJA (Fig. 3C-E; Fig. S8C). Since PIN3 is a major player in SII LR angle regulation (Rosquete *et al.* 2013), we examined the phenotype of *pin3-4* and observed no difference in LR angles when compared with WT (Fig. S8D). Other weak alleles of the *pinoid* mutants (*pin4-3* and *pin7-2*) exerted a WT like response to MeJA (Žádníková *et al.* 2010) (Fig. S8D). The branching angle distribution of the auxin receptor mutant *transport inhibitor response 1* (*tir1-1*) had more LRs in the >70° category compared to WT (Fig. 4A-B; Fig. S8E). We also used *auxin resistant 1* (*axr1-3*) and *axr2-1,* a mutant of *IAA7,* which show constitutive downregulation of auxin responses (Leyser *et al.* 1993; Nagpal *et al.* 2000). Seedlings carrying the weak allele of *axr1-3* had near normal WT like responses to MeJA treatment (Fig. 4A, C; Fig. S8F). However, this response was completely abolished in *axr2-1* (Fig. 4A, C; Fig. S8F). Altogether, these results indicate that an intact auxin transport and signalling machinery is required for MeJA to set the angular growth of LRs.

MYC2 controls the transcription of *CYP79B2, LAZY2* and *LAZY4*. A former report has shown that jasmonate mediates the regulation of auxin biosynthesis (Sun *et al.* 2009). *CYP79B2*, a cytochrome P450 mono-oxygenase that forms indole-3-acetaldoxime and acts as a precursor for auxin biosynthesis (Zhao *et al.* 2002). Additionally, other molecular components are involved in modifying plant architecture, including the *LAZY* gene family, which controls both root and shoot gravitropism (Taniguchi *et al.* 2017; Yoshihara & Spalding 2017). *LAZY2 (DRO3/NGR1), LAZY3 (DRO2/NGR3),* and *LAZY4 (DRO1/NGR2)* have been reported to cause auxin redistribution and transport in *Arabidopsis* LRs; thus, leading to an overall vertical root architecture (Taniguchi *et al.* 2017; Yoshihara & Spalding 2017). Mutants defective in *LAZY2* and *LAZY4* have agravitropic LRs (Taniguchi *et al.* 2017; Yoshihara & Spalding 2017), prompted us to study jasmonate effects on the LAZY2 and

LAZY4-mediated LR angles. Given the established role of JA in transcriptionally activating

CYP79B2, and that MYC2 is the master regulator of many JA responses, it is reasonable to

hypothesize that MYC2 might control the transcription of these genes, enhancing LR

response to gravity. To test this hypothesis, we examined the MYC2-depenedent expression

of *CYP79B2, LAZY2* and *LAZY4.* As shown in Figure 5A-C, *CYP79B2, LAZY2* and *LAZY4*

transcript levels were downregulated in *myc2myc3myc4.* To understand how MYC2 regulates

the expression of these genes, we scanned the promoters of *CYP79B2*, *LAZY2* and *LAZY4,*

and found G- and E-box elements. Using chromatin immunoprecipitation-quantitative PCR

(ChIP-qPCR), we found that *CYP79B2*, *LAZY2,* and *LAZY4* promoters were highly enriched

with MYC2 protein in *35S::MYC2-GFP* when compared to WT (Fig. 5D-F). We also

examined the binding of MYC2 to the promoters of *ORA59,* which is a known MYC2 target

and served as a positive control for our experiments (Zhai *et al.* 2013). *ATXR6* was used as a

negative control because it does not possess any MYC2 binding sites in its promoter region

(Fig. 5D-F). The results suggest that MYC2 binds and controls the transcription of genes that

can influence the vertical orientation of LRs.

MeJA controls the direction of auxin transport that may regulate LR angle. There are

various factors that control the GSA of LRs via the regulation of auxin flow from LR tips

(Claudia-Anahı´ Pèrez-Torres *et al.* 2008; Roychoudhry *et al.* 2017; Taniguchi *et al.* 2017).

These reports led us to investigate whether JA signalling influences auxin distribution at the

Stage II (SII) LR tips. For this, we analysed the effect of MeJA on *DR5::GFP* expression.

Most of the 1/2 MS-treated LRs displayed near symmetrical *DR5::GFP* expression with a

faint green signal streak towards the lower side of the LR tip (Fig. 6A). However, after the

addition of MeJA, the *DR5::GFP* expression started to disappear from the upper side and the

green signal became more apparent on the margins of the lower side (Fig. 6A). These

observations suggested that JA signalling directs the flow of auxin towards the lower side,

which might lead to vertical orientation of SII LRs. Under control conditions (1/2 MS), the

expression of *PIN3::PIN3-eGFP* was strongly expressed in the columella cells of SII LRs.

However, in seedlings treated with MeJA, there was an expansion of GFP signal towards the

stelar region of SII LRs but the signal intensity was very low. Also, there weren't any

changes in the differential distribution of PIN3::PIN3-GFP upon MeJA treatment. Upon

quantification of the signal intensity, we did not find any significant difference in its

expression between control and MeJA treatment, suggesting that JA signalling might not

involve PIN3 mediated auxin redistribution (Fig. S9A-B). In addition to PIN3 activity in the

columella, the action of PIN2 in the epidermis is required to drive the basipetal flow of auxin

away from the root tip (Roychoudhry *et al.* 2019). We, therefore, studied *PIN2::PIN2-eGFP*

expression in LRs growing at their GSA to check for any differential expression of PIN2 that

might contribute to the regulation of growth angle. The *PIN2::PIN2-eGFP* signal intensity

was strong and intact in 1/2 MS-treated seedlings. Also, there was not any difference in PIN2

expression between upper and lower halves of LRs (Fig. 6B). However, we observed that

MeJA diminished PIN2 redistribution in the upper epidermal cells of the LR (Fig. 6B).

Additionally, we found that the length of the 1[st] two upper epidermal cells of LR treated with

MeJA was longer than that in the control (Fig. 6B-C). It has been previously revealed that

differential elongation is a major factor controlling LR gravitropism (Rosquete *et al.* 2013)

and therefore, these results suggest that the JA signalling via PIN2 plays a potential role in

differential cell elongation, leading to vertical orientation of LRs.

MeJA integrates environmental cues into angular growth of LRs. Our data highlights a

role for JA signalling in modulating angular growth of LRs. Also, light can modify the GSA

of organs (Digby & Firn 2002; Roychoudhry *et al.* 2017). To investigate whether light affects

MeJA mediated branching angle, 5-day-old light-grown WT and *myc2myc3myc4* seedlings

were transferred to 0.5%Glc, 0.5%Glc+MeJA, 3%Glc, and 3%Glc+MeJA under long-day

regime (16 hours light/8 hours dark) and under continuous dark conditions. In the dark,

seedlings showed elongated hypocotyls and significantly more horizontally placed LRs than

those grown in light conditions (Fig. S10A-B). In both WT and *myc2myc3myc4*, the effect of

3%Glc was more prominent in dark conditions, with LRs displaying >70° angles. MeJA

treatment did not decrease the angles in dark conditions. Also, LRs of *myc2myc3myc4*

showed wider angles with MeJA treatment in dark conditions as compared to light (Fig.

S10A-B), thus suggesting that light is required for an optimal MeJA response. Next, to

investigate whether light affects auxin distribution in controlling branching angle, we treated

WT and *aux1-7* seedlings in light and dark regimes and observed that WT seedlings showed

broader angles in dark as compared to light (Fig. S11). Interestingly, we found that *aux1-7*

showed broader branching angle under both light and dark conditions, suggesting that light

modulates auxin distribution and ultimately affects branching angle. We also observed that

aux1-7 was less responsive to MeJA treatment under both light and dark conditions (Fig.

S11). Collectively, these results suggest that light and JA signalling are prerequisite to

influence auxin distribution in governing branching angle.

Light perception and signalling intersect with the action of jasmonates to affect plant

development and defence (Dombrecht *et al.* 2007). We, therefore, explored the involvement

of phytochrome (PHY) mediated light signalling in regulating this response. Photoreceptor

single mutants *phyA-201* and *phyB-5* showed broader angles when compared with WT,

whereas the *phyA201phyB5* showed a much stronger response and exhibited completely

altered Glc and MeJA sensitivity, with LRs showing horizontal angles at all concentrations

(Fig. 7A-B). Hence, *PHYA* and *PHYB* act redundantly in controlling the phenotype. To

further substantiate the role of light signalling in regulating this response, we investigated the

involvement of *HY5*, a major downstream positive regulator of phytochrome signalling (Gangappa & Botto 2016). The *hy5-1* mutant seedlings, which are already known to show wider LR angles, (Oyama, Shimura & Okada 1997) displayed horizontally positioned LRs upon JA treatment as compared to the WT (Fig. 7A-B). Previous reports of ChIP-seq data showed that the binding of HY5 to the promoter of *LIPOXYGENASE3 (LOX3)*, a JA biosynthetic gene (Lee *et al.* 2007). To further confirm whether light signalling is involved in maintaining an optimal JA response by increasing JA biosynthesis, we treated WT and *hy5-1* seedlings in long day and continuous dark conditions for 6 days. We also shifted 6-day-old continuous dark grown WT and *hy5-1* seedlings to light for 6 hours and checked for the expression of *LOX3*. Indeed, we found a significant increase in the expression of *LOX3* in long-day treatment when compared to total darkness (Fig. 7C). Also, WT seedlings exposed to a 6 hours dark to light transition exhibited enhanced LOX3 expression (Fig. 7D). In contrast, *LOX3* expression was significantly downregulated in *hy5-1* seedlings, suggesting that the response is *HY5*-dependent. Collectively, the above data suggest that JA signalling integrates environmental factor, such as light, into GSA establishment of LRs.

Discussion

The angular growth of LRs (GSA) represents an important element in the adaptability of the root system to its environment. Both nutrient and hormonal signals act locally to regulate GSA (Bai *et al.* 2013; Rosquete *et al.* 2013, 2018; Roychoudhry *et al.* 2013, 2017, 2019; Trachsel *et al.* 2013). The net effect of this adaptive response is an increase in the surface area of the plant root system for resource capture (e.g. horizontal LRs for phosphorus uptake), or the securing of anchorage (Lynch & Brown 2001; Trachsel *et al.* 2013). Similarly, shoots with vertical lateral branches show enhanced efficiency of light capture, allowing higher density planting and higher yields (Sakamoto *et al.* 2006; Vriet, Russinova & Reuzeau

2012). The main aim of this study is to advance our understanding of GSA regulation and to

elucidate interconnections between environmental and hormonal signals in fine-tuning the

branching angle of *Arabidopsis* roots.

In this work, we demonstrated that MeJA reduced the branching angle of roots in a dose-

dependent manner, thereby resulting in an overall vertical orientation. Analysis of mutants

defective in JA biosynthesis and signalling revealed that an intact JA biosynthesis and

signalling is prerequisite to bring about changes in the branching angle. The branching angles

of *jar1-11*, *jin1-9,* and *myc2myc3myc4* mutants were defective in perceiving endogenous JA,

even under control conditions (1/2 MS), suggesting that, in nature, JA signalling plays an

important developmental role in governing root branching angle. The weaker phenotype in

jin1-9 compared to that of *myc2myc3myc4* suggested that *MYC3* and *MYC4* act additively

with *MYC2* in regulating this response. Since *MYC3* and *MYC4* are weakly expressed in the

roots of young seedlings unlike *MYC2* (Fernández-Calvo *et al.* 2011), and *MYC2* is the major

regulator of many JA governed biological responses (Lorenzo, Chico, Sánchez-Serrano &

Solano 2004; Dombrecht *et al.* 2007), we postulate that this response is majorly mediated by

MYC2.

Apart from affecting various parameters of RSA (Gupta, Singh, Mishra, Kushwah & Laxmi

2009; Gupta *et al.* 2015; Mishra *et al.* 2009; Singh *et al.* 2014a b), sugars can also influence

the gravitropic behaviour of lateral organs (Willemoes *et al.* 1988). Previous studies have

shown that 3%Glc causes more production of LRs and root growth and is therefore,

considered optimal for overall plant growth and development (Mishra *et al.* 2009; Singh *et al.*

2014a; Gupta *et al.* 2015). In this study, 3%Glc caused a significant shift in the branching

angle towards a more horizontal orientation. This effect of Glc on growth can favour plant

propagation because it allows plants to explore adjacent territories. The diagravitropic growth

of LRs upon Glc treatment was consistent with a report by Willemoës *et al.* (1998),

suggesting that sugars are responsible for maintaining a non-vertical angular growth of

organs (Willemoes *et al.* 1988). There are also reports citing crosstalk between sugars and

GA_3 to regulate growth direction of organs (Montaldi 1969; Willemoes *et al.* 1988).

Compared with well-studied interactions between Glc and other hormones, relatively little is

known about Glc and JA (Song *et al.* 2017; Vleesschauwer *et al.* 2017; Guo *et al.* 2018).

Most of these studies demonstrate an antagonistic interaction between these two signalling

pathways in both dicots and monocots (Song *et al.* 2017; Vleesschauwer *et al.* 2017).

Moreover, transcriptomic analysis revealed the regulation of JA signalling pathway by TOR-

kinase (Dong *et al.* 2015). In our study, microarray analysis, immunoblot assays and

physiological studies have confirmed that Glc and JA are two crucial signals that work

antagonistically via TOR kinase pathway to regulate the branching angle of *Arabidopsis*

roots. This observation falls in agreement with the previous report citing antagonistic

interaction between JA and TOR kinase (Song *et al.* 2017). Transcriptome analysis of rice

cells treated with the TOR-specific inhibitor rapamycin revealed that TOR, apart from

dictating transcriptional reprogramming of extensive gene sets involved in central and

secondary metabolism, cell cycle, and transcription, also suppresses many defence-related

genes and antagonizes the action of JA (Vleesschauwer *et al.* 2017). We hypothesize that

this antagonism occurs in nature in order to fine-tune the response and achieve the optimum

angle required for growth. However, more detailed molecular dissection is necessary to

obtain a deeper understanding of this crosstalk and its regulators.

Of all the plant hormones, auxin plays a central role in LR GSA control. Transient expression

of *PIN3* with strong repression of *PIN4* and *PIN7* in young LRs limits auxin redistribution

and hence explains the reduced gravitropic competence of laterals (Rosquete *et al.* 2013,

2018). JAs contribute to the regulation of transport of IAA by inducing the expression of

PIN1 and *PIN2* (Sun *et al.* 2009) and modulate the accumulation of PIN2 in the plasma

membrane and its recycling via endocytosis, in a dose-dependent manner (Sun *et al.* 2011). In

light of the above reports claiming recent interconnections between JA and auxin signalling,

and the modulation of JA homeostasis and signal transduction mimicking auxin effects on

root development, we show that MeJA requires auxin machinery to control branching angle.

Our results suggest that genetic disruption of auxin transport and signalling nullifies the

effect of MeJA-mediated control of branching angle. Additionally, due to differences in the

phenotypic strength of the mutants, MDR1-mediated auxin transport and TIR1-AXR1-

mediated auxin signalling show much fewer effects than AUX1-PIN2 and AXR2-mediated

transport and signalling (Tiryaki & Staswick 2002; Quint, Barkawi, Fan, Cohen & Gray

2009). JA signalling via MYC2 also induces the transcription of *CYP79B2*, *LAZY2* and

LAZY4. We suppose that this transcriptional activation might promote auxin biosynthesis and

its transport and redistribution, ultimately leading to vertical angles. Also, the modulation of

the asymmetric auxin transport in LR columella cells as observed by DR5::GFP expression,

diminished PIN2 activity from the upper epidermal cell profile, and the differential cell

elongation caused by MeJA may correlate with the vertical orientation of the LRs.

Light is one of the many diverse signals that changes the sugar/energy status in plants.

Studies till date have given tremendous insights into how light not only drives photosynthesis

but also acts as an environmental cue that informs plants about their environment. Previous

literature have stated that light can increase root gravitropism by influencing the direction of

polar auxin transport (Buer & Muday 2004; Laxmi, Pan, Morsy & Chen 2008; Sassi *et al.*

2012). Consistently, our results demonstrate that light acts as a stimulus to change auxin

distribution to modulate branching angle of *Arabidopsis* LRs. Also, the exogenous

application of MeJA could not restore the vertical orientation of branching angle in *aux1-7*

neither in dark nor in light, further suggesting that auxin distribution is indispensable for

MeJA modulation of branching angle in light. Light signalling also influences auxin transport

in the roots by vacuolar degradation of PIN proteins (Korbei & Luschnig 2013) and is

essential for regulation of root gravitropism. Light via COP1/CSN complex controls the

vacuolar targeting of PIN2-GFP (Laxmi *et al.* 2008; Sassi *et al.* 2012).

Previous reports have demonstrated that light quality governs the gravitropic behaviour of

lateral organs (Kiss *et al.* 2002). *Arabidopsis* LRs showed negative phototropism in the

presence of white light, and positive phototropism in response to red light (Kiss *et al.* 2002).

In the present study, we observed that light, acting via the redundant functions of PHYA-

PHYB and HY5 induced a more vertical LR orientation, supporting results reported by Kiss

et al. (2002). Light quality also influences formation of lateral root primordia by

accumulation of HY5. Also, HY5-mediated light signalling influences root development by

regulating auxin transport and signalling (van Gelderen *et al.* 2018). In our experiments,

exogenous application of MeJA could not induce vertical branching angles of LRs in dark

conditions. This suggests that light is prerequisite to MeJA-induced changes in branching

angle. We assume that the alteration in the angle is not only achieved by the biosynthesis of

JA, but also by the influence of another factor or signalling event that is currently unknown.

ChIP-seq data showed the binding of *HY5* at the promoter of *LOX3* (Lee *et al.* 2007).

Consistent with this observation, we found decreased expression levels of *LOX3* in *hy5-1*

under light conditions as well as dark to light transition. This suggests a likelihood that HY5

regulates JA levels in light. Other studies have shown that light environment and circadian

rhythms are crucially involved in modulating plant responses to JA biosynthesis (Radhika,

Kost, Mithofer & Boland 2010; Goodspeed, Chehab, Min-Venditti, Braam & Covington

2012). JA levels have been shown to rise during the day, reaching a maximum at midday, and

then declining again in the afternoon (Goodspeed *et al.* 2012). MYC2 protein levels were also

shown to rise during the day and under continuous light conditions (Shin, Heidrich, Sanchez-

Villarreal, Parker & Davis 2012). Moreover, ChIP-seq analyses show the binding of HY5 on

the promoter of *MYC2* (Lee *et al.* 2007). Considering previous literature and our current

findings, we show that light either directly or indirectly can dramatically affect plant root

growth and development. We have, therefore, in this study have unraveled the

interconnection of light signalling and its modulation by glucose (a downstream product of

light driven photosynthesis) and phytohormones like JA and auxin in controlling root

branching angle.

Based on our investigation and previous findings, we propose a testable model in Figure 7E.

We hypothesize that light works via two branches to optimize the branching angle of

Arabidopsis LRs. Contrary to the general notion that light and sugars should have the same

influence on developmental outputs, there are many reports that suggest that light and Glc

signalling have opposite effects on the growth and development of *Arabidopsis* seedlings

(Moore 2003; Eckstein, Zieba & Gabryś 2012). Light working via phytochrome signalling

(*PHYA/PHYB-HY5*) increases JA levels, which ultimately lead to the activation of JA

signalling and this physiological response is shown to be mainly mediated by *MYC2*. Our

results also suggest that auxin works further downstream as auxin transport and signalling

mutants displayed abrogated responses to JA regulation of branching angle. Additionally,

MYC2 regulates the expression of *CYP79B2*, *LAZY2* and *LAZY4* by binding to their

promoters that might promote auxin biosynthesis and its transport and redistribution,

ultimately leading to vertical angles. Glc, on the other hand, is produced by photosynthesis

promotes radial expansion of the root architecture, antagonistically interacting with JA

signalling via the *TOR*-mediated pathway and affecting the accumulation of JAZ9 to regulate

this developmental process.

We hypothesize that under normal conditions, roots grow in the soil without light. Upon light

exposure, the stress hormone JA is induced that further regulates auxin transport and

signalling to accelerate the vertical growth of LRs to avoid light. However, under sufficient

energy (3%Glc), roots do not feel stressed and hence, do not seem to avoid light and grow

horizontally. Though, horizontal angles favour plant propagation and explore adjacent

territories better but do not play a significant role in root anchorage (due to less anchorage

force) (Ennos 2000; Mickovski & Ennos 2003), Similarly, steeper LRs show less soil column

weight and hence low uprooting resistance. Thus, to maintain a balance, MeJA and Glc

signalling act in concert to maintain an optimal angle where LRs act as guy ropes, holding the

vertical root in position that not only favours plant propagation but also provides maximum

anchorage to prevent uprooting of plants (Ennos 2000). Apart from the root architecture,

signals from the environment and endogenous cues act together to optimize growth angles of

lateral branches. Depending upon the strength of the signal, the angle of the laterals is

decided. The present study in *Arabidopsis* can be used for crop improvement strategies that

can lead to crop species with improved water and nutrient acquisition capability and hence is

of broad relevance to plant and crop scientists. Future work will be directed at identifying

substantial mechanistic insights into hormonal-environmental crosstalk and characterizing

novel molecular components that are involved in shaping the overall root architecture.

Acknowledgements

The authors would like to thank Dr. Aditi Gupta for her advice and discussions and Ms

Harshita Bharti Saksena for reading the manuscript. The authors acknowledge NIPGR

Confocal Facility for their assistance, Prof. Philippe Reymond for providing seeds of

myc2myc3myc4, Dr. Nam-Hai Chua for providing seeds of *35S-MYC2-GFP* and Dr. Laurent

Laplaze for providing seeds of *p35S::Jas9-N7-VENUS*. The authors are thankful to DBT-

eLibrary Consortium (DeLCON) for providing access to e-resources. This work was

financially supported by the Core Grant from the National Institute of Plant Genome

Research to A.L., University Grant Commission, Government of India and Department of

Biotechnology, Government of India. M.S. acknowledges University Grant Commission,

Government of India for research fellowship, M.S. acknowledges Department of

Biotechnology, Government of India and MJK acknowledges Department of Science and

Technology (INSPIRE Faculty Programme Grant IFA18-LSPA110).

Author Contributions

M.S. and A.L. conceived and designed the experiments. M.S. performed physiology and

microarray. M.S, M.S[#] and M.J.K performed confocal experiments. M.S and M.S[#] performed

real time assays, western blot and ChIP-qPCR. M.S. wrote the article. M.S[#] and M.J.K.

assisted in microarray analysis and preparing the manuscript; A.L. supervised and

complemented the article.

Conflict of Interest statement

The authors declare no conflict of interest.

References

Baena-González E. (2010) Energy signaling in the regulation of gene expression during

stress. *Molecular Plant* **3**, 300–313.

Baena-González E., Rolland F., Thevelein J.M. & Sheen J. (2007) A central integrator of

transcription networks in plant stress and energy signalling. *Nature* **448**, 938–42.

Bai H., Murali B., Barber K. & Wolverton C. (2013) Low phosphate alters lateral root

setpoint angle and gravitropism. *American Journal of Botany* **100**, 175–182.

Buer C.S. & Muday G.K. (2004) The transparent testa4 mutation prevents flavonoid

synthesis and alters auxin transport and the response of arabidopsis roots to gravity and

light. *Plant Cell*.

Chen L.Q., Hou B.H., Lalonde S., Takanaga H., Hartung M.L., Qu X.Q., … Frommer W.B.

(2010) Sugar transporters for intercellular exchange and nutrition of pathogens. *Nature*.

Cuesta C., Wabnik K. & Benková E. (2013) Systems approaches to study root architecture

dynamics. *Frontiers in Plant Science* **4**, 1–11.

Deprost D., Yao L., Sormani R., Moreau M., Leterreux G., Nicolaï M., … Meyer C. (2007)

The Arabidopsis TOR kinase links plant growth, yie... [EMBO Rep. 2007] - PubMed

result. *EMBO reports*.

Digby J. & Firn R.D. (1995) The gravitropic set-point angle (GSA): the identification of an

important developmentally controlled variable governing plant architecture. *Plant, cell*

& environment **18**, 1434–40.

Digby J. & Firn R.D. (2002) Light modulation of the gravitropic set-point angle (GSA).

Journal of experimental botany **53**, 377–381.

Dombrecht B., Xue G.P., Sprague S.J., Kirkegaard J.A., Ross J.J., Reid J.B., … Kazan K.

(2007) MYC2 Differentially Modulates Diverse Jasmonate-Dependent Functions in

Arabidopsis. *the Plant Cell Online* **19**, 2225–2245.

Dong P., Xiong F., Que Y., Wang K., Yu L., Li Z. & Ren M. (2015) Expression profiling and

functional analysis reveals that TOR is a key player in regulating photosynthesis and

phytohormone signaling pathways in Arabidopsis. *Frontiers in Plant Science*.

Eckstein A., Zieba P. & Gabryś H. (2012) Sugar and Light Effects on the Condition of the

Photosynthetic Apparatus of Arabidopsis thaliana Cultured in vitro. *Journal of Plant*

Growth Regulation.

Ennos A.R. (2000) The mechanics of root anchorage. *Advances in Botanical Research*.

Eveland A.L. & Jackson D.P. (2012) Sugars, signalling, and plant development. *Journal of*

Experimental Botany **63**, 3367–3377.

Fernández-Calvo P., Chini A., Fernández-Barbero G., Chico J.-M., Gimenez-Ibanez S., Geerinck J., … Solano R. (2011) The Arabidopsis bHLH transcription factors MYC3 and MYC4 are targets of JAZ repressors and act additively with MYC2 in the activation of jasmonate responses. *The Plant cell* **23**, 701–715.

Gan L., Wu H., Wu D., Zhang Z., Guo Z., Yang N., … Zhu C. (2015) Methyl jasmonate inhibits lamina joint inclination by repressing brassinosteroid biosynthesis and signaling in rice. *Plant Science* **241**, 238–245.

Gangappa S.N. & Botto J.F. (2016) The Multifaceted Roles of HY5 in Plant Growth and Development. *Molecular Plant* **9**, 1353–1365.

García-González J., Lacek J. & Retzer K. (2021) Dissecting Hierarchies between Light, Sugar and Auxin Action Underpinning Root and Root Hair Growth. *Plants*.

van Gelderen K., Kang C., Paalman R., Keuskamp D., Hayes S. & Pierik R. (2018) Far-red light detection in the shoot regulates lateral root development through the HY5 transcription factor. *Plant Cell*.

Goodspeed D., Chehab E.W., Min-Venditti A., Braam J. & Covington M.F. (2012) Cozzarelli Prize Winner: Arabidopsis synchronizes jasmonate-mediated defense with insect circadian behavior. *Proceedings of the National Academy of Sciences* **109**, 4674–4677.

Guo R., Shen W., Qian H., Zhang M., Liu L. & Wang Q. (2018) Jasmonic acid and glucose synergistically modulate the accumulation of glucosinolates in Arabidopsis thaliana. **64**, 5707–5719.

Gupta A., Singh M. & Laxmi A. (2015) Interaction between glucose and brassinosteroid during the regulation of lateral root development in arabidopsis. *Plant Physiology* **168**, 307–320.

Gupta A., Singh M., Mishra B.S., Kushwah S. & Laxmi A. (2009) Role of glucose in spatial

distribution of auxin regulated genes. **4**, 862–863.

Gutjahr C., Riemann M., Müller A., Düchting P., Weiler E.W. & Nick P. (2005) Cholodny–

Went revisited: a role for jasmonate in gravitropism of rice coleoptiles. *Planta* **222**, 575–

585.

Guyomarc'h S., Léran S., Auzon-Cape M., Perrine-Walker F., Lucas M. & Laplaze L. (2012)

Early development and gravitropic response of lateral roots in Arabidopsis thaliana.

Philosophical Transactions of the Royal Society B: Biological Sciences **367**, 1509–1516.

Huang J., Taylor J.P., Chen J.-G., Uhrig J.F., Schnell D.J., Nakagawa T., … Jones A.M.

(2006) The plastid protein THYLAKOID FORMATION1 and the plasma membrane G-

protein GPA1 interact in a novel sugar-signaling mechanism in Arabidopsis. *The Plant*

cell **18**, 1226–1238.

Kircher S. & Schopfer P. (2012) Photosynthetic sucrose acts as cotyledon-derived long-

distance signal to control root growth during early seedling development in Arabidopsis.

Proceedings of the National Academy of Sciences of the United States of America **109**,

11217–21.

Kiss J.Z., Miller K.M., Ogden L. a & Roth K.K. (2002) Phototropism and gravitropism in

lateral roots of Arabidopsis. *Plant & cell physiology* **43**, 35–43.

Korbei B. & Luschnig C. (2013) Plasma membrane protein ubiquitylation and degradation as

determinants of positional growth in plants. *Journal of Integrative Plant Biology*.

Larrieu A., Champion A., Legrand J., Lavenus J., Mast D., Brunoud G., … Laplaze L. (2015)

A fluorescent hormone biosensor reveals the dynamics of jasmonate signalling in plants.

Nature Communications **6**, 1–9.

Lastdrager J., Hanson J. & Smeekens S. (2014) Sugar signals and the control of plant growth

and development. *Journal of Experimental Botany*.

Laxmi A., Pan J., Morsy M. & Chen R. (2008) Light plays an essential role in intracellular

distribution of auxin efflux carrier PIN2 in Arabidopsis thaliana. *PLoS ONE.*

Lee J., He K., Stolc V., Lee H., Figueroa P., Gao Y., … Xing W.D. (2007) Analysis of

transcription factor HY5 genomic binding sites revealed its hierarchical role in light

regulation of development. *Plant Cell.*

Leyser H.M.O., Lincoln C.A., Timpte C., Lammer D., Turner J. & Estelle M. (1993)

Arabidopsis auxin-resistance gene AXR1 encodes a protein related to ubiquitin-

activating enzyme E1. *Nature* **364**, 161–164.

Li L. & Sheen J. (2016) Dynamic and diverse sugar signaling. *Current Opinion in Plant*

Biology **33**, 116–125.

Lilley J.L.S., Gee C.W., Sairanen I., Ljung K. & Nemhauser J.L. (2012) An endogenous

carbon-sensing pathway Triggers increased auxin flux and hypocotyl elongation. *Plant*

Physiology.

Lorenzo O., Chico J.M., Sánchez-Serrano J.J. & Solano R. (2004) JASMONATE-

INSENSITIVE1 encodes a MYC transcription factor essential to discriminate between

different jasmonate-regulated defense responses in arabidopsis. *Plant Cell.*

Lorenzo W.O., Chico J.M., Sá Nchez-Serrano J.J. & Solano R. JASMONATE-

INSENSITIVE1 Encodes a MYC Transcription Factor Essential to Discriminate

between Different Jasmonate-Regulated Defense Responses in Arabidopsis.

Lynch J.P. & Brown K.M. (2001) Topsoil foraging—an architectural adaptation of plants to

low phosphorus availability. *Plant and Soil* **237**, 225–237.

Malamy J.E. (2005) Intrinsic and environmental response pathways that regulate root system

architecture. *Plant, Cell and Environment* **28**, 67–77.

Mickovski S.B. & Ennos A.R. (2003) Anchorage and asymmetry in the root system of Pinus

peuce. *Silva Fennica.*

Mishra B.S., Singh M., Aggrawal P. & Laxmi A. (2009) Glucose and auxin signaling

799 interaction in controlling arabidopsis thaliana seedlings root growth and development.

800 *PLoS ONE* **4**.

801 Montaldi E.R. (1969) Gibberellin-sugar interaction regulating the growth habit of

802 bermudagrass (Cynodon dactylon (L) Pers). *Experientia* **25**, 91–92.

803 Montané M.H. & Menand B. (2013) ATP-competitive mTOR kinase inhibitors delay plant

804 growth by triggering early differentiation of meristematic cells but no developmental

805 patterning change. *Journal of Experimental Botany*.

806 Moore B. (2003) Role of the Arabidopsis Glucose Sensor HXK1 in Nutrient, Light, and

807 Hormonal Signaling. *Science* **300**, 332–336.

808 Moore B., Zhou L., Rolland F., Hall Q., Cheng W.H., Liu Y.X., … Sheen J. (2003) Role of

809 the Arabidopsis glucose sensor HXK1 in nutrient, light, and hormonal signaling.

810 *Science*.

811 Nagpal P., Walker L.M., Young J.C., Sonawala A., Timpte C., Estelle M. & Reed J.W.

812 (2000) AXR2 encodes a member of the Aux/IAA protein family. *Plant physiology* **123**,

813 563–74.

814 Oyama T., Shimura Y. & Okada K. (1997) The Arabidopsis HY5 gene encodes a bZIP

815 protein that regulates stimulus- induced development of root and hypocotyl. *Genes and*

816 *Development*.

817 Quint M., Barkawi L.S., Fan K.T., Cohen J.D. & Gray W.M. (2009) Arabidopsis IAR4

818 Modulates Auxin Response by regulating auxin homeostasis 1[OA]. *Plant Physiology*.

819 Radhika V., Kost C., Mithofer A. & Boland W. (2010) Regulation of extrafloral nectar

820 secretion by jasmonates in lima bean is light dependent. *Proceedings of the National*

821 *Academy of Sciences* **107**, 17228–17233.

822 Ramon M., Rolland F. & Sheen J. (2008) Sugar sensing and signaling. *The Arabidopsis book*

823 */ American Society of Plant Biologists* **6**, e0117.

Raya-González J., Pelagio-Flores R. & López-Bucio J. (2012) The jasmonate receptor COI1

plays a role in jasmonate-induced lateral root formation and lateral root positioning in

Arabidopsis thaliana. *Journal of Plant Physiology* **169**, 1348–1358.

Rosquete M.R., Waidmann S. & Kleine-Vehn J. (2018) PIN7 auxin carrier has a preferential

role in terminating radial root expansion in Arabidopsis thaliana. *International Journal*

of Molecular Sciences **19**, 1–11.

Rosquete M.R., Wangenheim D. Von, Barbez E., Stelzer E.H.K., Maizel A. & Benkova E.

(2013) Report An Auxin Transport Mechanism Restricts Positive Orthogravitropism in

Lateral Roots. **23**, 817–822.

Roychoudhry S., Del Bianco M., Kieffer M. & Kepinski S. (2013) Auxin controls gravitropic

setpoint angle in higher plant lateral branches. *Current Biology* **23**, 1497–1504.

Roychoudhry S., Kieffer M., Del Bianco M., Liao C.-Y., Weijers D. & Kepinski S. (2017)

The developmental and environmental regulation of gravitropic setpoint angle in

Arabidopsis and bean. *Scientific Reports* **7**, 42664.

Roychoudhry S., Sageman-Furnas K., Wolverton C., Goodman H.L., Grones P., Mullen J.,

… Kepinski S. (2019) Antagonistic and auxin-dependent phosphoregulation of

columella PIN proteins controls lateral root gravitropic setpoint angle in Arabidopsis.

bioRxiv, 594838.

Sairanen I., Novák O., Pěnčík A., Ikeda Y., Jones B., Sandberg G. & Ljung K. (2013) Soluble

carbohydrates regulate auxin biosynthesis via PIF proteins in arabidopsis. *Plant Cell*.

Sakamoto T., Morinaka Y., Ohnishi T., Sunohara H., Fujioka S., Ueguchi-Tanaka M., …

Matsuoka M. (2006) Erect leaves caused by brassinosteroid deficiency increase biomass

production and grain yield in rice. *Nature Biotechnology* **24**, 105–109.

Sassi M., Lu Y., Zhang Y., Wang J., Dhonukshe P., Blilou I., … Xu J. (2012) COP1 mediates

the coordination of root and shoot growth by light through modulation of PIN1- and

PIN2-dependent auxin transport in Arabidopsis. *Development (Cambridge)*.

Sheen J. (2014) Master regulators in plant glucose signaling networks. *Journal of Plant Biology*.

Shin J., Heidrich K., Sanchez-Villarreal A., Parker J.E. & Davis S.J. (2012) TIME FOR COFFEE represses accumulation of the MYC2 transcription factor to provide time-of-day regulation of jasmonate signaling in Arabidopsis. *The Plant cell* **24**, 2470–82.

Singh M., Gupta A. & Laxmi A. (2014a) Glucose and phytohormone interplay in controlling root directional growth in Arabidopsis. *Plant Signaling and Behavior* **9**, 2–7.

Singh M., Gupta A. & Laxmi A. (2014b) Glucose control of root growth direction in Arabidopsis thaliana. *Journal of Experimental Botany* **65**, 2981–2993.

Song Y., Zhao G., Zhang X., Li L., Xiong F., Zhuo F., … Li F. (2017) The crosstalk between Target of Rapamycin (TOR) and Jasmonic Acid (JA) signaling existing in Arabidopsis and cotton. *Scientific Reports* **7**, 45830.

Staswick P.E. (2009) The Tryptophan Conjugates of Jasmonic and Indole-3-Acetic Acids Are Endogenous Auxin Inhibitors. *PLANT PHYSIOLOGY* **150**, 1310–1321.

Staswick P.E. & Tiryaki I. (2004) The oxylipin signal jasmonic acid is activated by an enzyme that conjugates it to isoleucine in Arabidopsis. *The Plant cell* **16**, 2117–27.

Stokes M.E., Chattopadhyay A., Wilkins O., Nambara E. & Campbell M.M. (2013) Interplay between sucrose and folate modulates auxin signaling in Arabidopsis. *Plant Physiology*.

Sun J., Chen Q., Qi L., Jiang H., Li S., Xu Y., … Li C. (2011) Jasmonate modulates endocytosis and plasma membrane accumulation of the arabidopsis pin2 protein. *New Phytologist* **191**, 360–375.

Sun J., Xu Y., Ye S., Jiang H., Chen Q., Liu F., … Li C. (2009) Arabidopsis ASA1 is important for jasmonate-mediated regulation of auxin biosynthesis and transport during lateral root formation. *The Plant cell* **21**, 1495–511.

Taniguchi M., Furutani M., Nishimura T., Nakamura M., Fushita T., Iijima K., … Morita

M.T. (2017) The Arabidopsis LAZY1 Family Plays a Key Role in Gravity Signaling

within Statocytes and in Branch Angle Control of Roots and Shoots. *The Plant Cell*,

tpc.00575.2016.

Tiryaki I. & Staswick P.E. (2002) An Arabidopsis mutant defective in jasmonate response is

allelic to the auxin-signaling mutant axr1. *Plant Physiology*.

Trachsel S., Kaeppler S.M., Brown K.M. & Lynch J.P. (2013) Maize root growth angles

become steeper under low N conditions. *Field Crops Research* **140**, 18–31.

Urano D., Phan N., Jones J.C., Yang J., Huang J., Grigston J., … Jones A.M. (2012a)

Endocytosis of the seven-transmembrane RGS1 protein activates G-protein-coupled

signalling in Arabidopsis.

Urano D., Phan N., Jones J.C., Yang J., Huang J., Grigston J., … Jones A.M. (2012b)

Endocytosis of the seven-transmembrane RGS1 protein activates G-protein-coupled

signalling in Arabidopsis. *Nature Cell Biology* **14**, 1079–1088.

Vleesschauwer D. De, Filipe O., Hoffman G., Seifi H.S., Haeck A., Canlas P., … Hofte M.

(2017) Target of rapamycin signaling orchestrates growth – defense trade-offs in plants.

Vriet C., Russinova E. & Reuzeau C. (2012) Boosting Crop Yields with Plant Steroids. *The*

Plant Cell **24**, 842–857.

Waidmann S., Ruiz Rosquete M., Schöller M., Sarkel E., Lindner H., LaRue T., … Kleine-

Vehn J. (2019) Cytokinin functions as an asymmetric and anti-gravitropic signal in

lateral roots. *Nature Communications* **10**.

Wasternack C. & Hause B. (2013) Jasmonates: Biosynthesis, perception, signal transduction

and action in plant stress response, growth and development. An update to the 2007

review in Annals of Botany. *Annals of Botany* **111**, 1021–1058.

Willemoes J.G., Beltrano J. & Montalbi R. (1988) Gibberellic acid-reversion of

paclobutrazol-inhibited growth on in vitro cultured potato nodal segments. 3–5.

Xiao G., Qin H., Zhou J., Quan R., Lu X., Huang R. & Zhang H. (2016) OsERF2 controls

rice root growth and hormone responses through tuning expression of key genes

involved in hormone signaling and sucrose metabolism. *Plant Molecular Biology*.

Xiong Y. & Sheen J. (2015) Novel links in the plant TOR kinase signaling network. *Current*

Opinion in Plant Biology **28**, 83–91.

Yoshihara T. & Spalding E.P. (2017) LAZY genes mediate the effects of gravity on auxin

gradients and plant architecture. *Plant Physiology*.

Yuan T.T., Xu H.H., Zhang K.X., Guo T.T. & Lu Y.T. (2014) Glucose inhibits root meristem

growth via ABA INSENSITIVE 5, which represses PIN1 accumulation and auxin

activity in Arabidopsis. *Plant, Cell and Environment* **37**, 1338–1350.

Žádníková P., Petrášek J., Marhavý P., Raz V., Vandenbussche F., Ding Z., … Benková E.

(2010) Role of PIN-mediated auxin efflux in apical hook development of Arabidopsis

thaliana. *Development*.

Zhai Q., Yan L., Tan D., Chen R., Sun J., Gao L., … Li C. (2013) Phosphorylation-Coupled

Proteolysis of the Transcription Factor MYC2 Is Important for Jasmonate-Signaled Plant

Immunity. *PLoS Genetics*.

Zhao Y., Hull A.K., Ecker J.R., Normanly J., Chory J. & Celenza J.L. (2002) Trp-dependent

auxin biosynthesis in. *Genes & Development*, 3100–3112.

Legends

Figure 1. MeJA decreases branching angle of Arabidopsis roots. **(A)** Phenotype of light
grown 12-day-old Arabidopsis Col-0 seedlings grown on different doses of MeJA. **(B)**
Average branching angle of 12-day-old Col-0 seedlings on different doses of MeJA. Data are
mean-SE of 3 biological replicates with 25 seedlings. **(C)** Distribution of branching angle of
12-day-old Col-0 Arabidopsis seedlings on different doses of MeJA. Data are mean-SE of 3
biological replicates with 25 seedlings. **(D)** Comparison of the effects of different
phytohormones (ACC, ABA, BR, GA) on Col-0 seedlings to determine their roles in
branching angle. Data are mean-SE of 3 biological replicates with 25 seedlings. **(E)**

Phenotype of light grown 12-day-old Arabidopsis Col-0, *jar1-11*, *coi1*, *jin1-9* and *myc2myc3myc4* seedlings grown on different doses of MeJA. **(F)** Distribution of branching angle in Col-0, *jar1-11* seedlings grown on different doses of MeJA. Data are mean-SE of 3 biological replicates with 25 seedlings. **(G)** Distribution of branching angle in seedlings of Col-0, *coi1* mutants grown on different doses of MeJA. Data are mean-SE of 4 biological replicates with 25 seedlings. **(H-I)** Distribution of branching angle in seedlings of Col-0, *jin1-9, myc2myc3myc4* mutants grown on different doses of MeJA. Data are mean-SE of 3 biological replicates with 25 seedlings. In A-I, 5-day-old 1/2 MS grown Col-0 seedlings were transferred to different doses of MeJA and phenotypes were analysed at 12th day. Asterisks indicate a significant difference in the studied parameter (P <0.05, paired two-tailed student's t-test; * control vs treatment and 'a' WT vs mutant).

Figure 2. Glc antagonizes MeJA-induced branching angle in *Arabidopsis* roots. (A) Phenotype of Col-0 seedlings grown in different concentrations of Glc (0.5%; 28 mM and 3%Glc; 167 mM) and in combination with 10 μM MeJA. **(B)** Distribution of branching angle in Col-0 seedlings grown in different concentrations of Glc and in combination with MeJA. The graph represents the average of 7 biological replicates having 25 plants in each replicate and error bars represent SE. **(C)** Distribution of branching angle in Col-0 seedlings grown in different concentrations of Man in combination with 1%Glc. The graph represents the average of 3 biological replicates consisting of 25 seedlings and error bars represent SE. **(D)** Phenotype of Col-0 and *sweet11sweet12* seedlings grown in different concentrations of Glc. **(E)** Distribution of branching angle in Col-0 and *sweet11sweet12* seedlings grown in different concentrations of Glc. The graph represents the average of 3 biological replicates having 50 plants in each replicate and error bars represent SE. In **A-E**, 5-day-old 1/2 MS grown seedlings were transferred to Glc and MeJA containing treatment media and phenotypes were analysed at 12th day. Asterisks indicate a significant difference in the studied parameter (*P* <0.05, Student's t-test; * control vs treatment and 'a' WT vs mutant. **(F)** Western blot analysis of total protein extracts of Jas9-VENUS seedlings treated for 3 hours with 0%, 0.5% and 3%Glc in combination with 10 μM MeJA and probed with an anti-GFP antibody. The experiment was performed four times. **(G)** Quantification of Jas9-VENUS protein levels. The graph represents average of 4 biological replicates. **(H)** Phenotype of Col-0 and *TOR* signalling mutant *tor35-7 RNAi* seedlings grown in different concentrations of Glc alone and in combination with 10 μM MeJA. **(I)** Distribution of branching angle in seedlings of Col-0 and *tor35-7 RNAi* grown in different concentrations of Glc alone or in combination with 10 μM MeJA. 5-day-old 1/2 MS grown Col-0 and *tor35-7* seedlings were transferred to treatment media and phenotypes were analysed at 12th day. The graph represents the average of 4 biological replicates consisting of 25 seedlings and error bars represent SE. Asterisks indicate a significant difference in the studied parameter (*P* <0.05, Student's t-test; * control vs treatment, and 'a' WT vs mutant). **(J)** Western blot analysis of total protein extracts of Jas9-VENUS seedlings treated for 3 hours with 0%, 0.5% and 3%Glc alone or in combination with 10 μM MeJA and 10 μM AZD-8055 for 3h and probed with an anti-GFP antibody. The experiment was performed four times. **(K)** Quantification of Jas9-VENUS protein levels. The graph represents average of 4 biological replicates. Asterisks indicate a significant difference in the studied parameter (P <0.05, paired two-tailed student's t-test; * control vs treatment and ** 0%Glc+AZD vs 3%Glc+AZD.

Figure 3. Components of auxin biosynthesis and transport lie downstream to MeJA-mediated branching angle. (A) Distribution of branching angle in Col-0 seedlings grown in 3%Glc alone or in combination with MeJA (10 μM) and different doses of NPA (100 nM to 1 μM). The graph represents the average of 4 biological replicates consisting of 25 seedlings

and error bar represent SE. **(B)** Distribution of branching angle in seedlings of auxin influx
defective mutant *aux1-7* grown in 3%Glc alone or in combination with different doses of
MeJA. The graph represents the average of 3 biological replicates consisting of 25 seedlings
and error bars represent SE. **(C)** Phenotype of light grown 12-day-old *Arabidopsis* Col-0,
aux1-7 and *eir1-1* seedlings grown in 3%Glc alone or in combination with different doses of
MeJA. **(D)** Phenotype of light grown 12-day-old *Arabidopsis* Ws and *mdr1-1* seedlings
grown in 3%Glc alone or in combination with different doses of MeJA. **(E)** Distribution of
branching angle in seedlings of WT (Col-0 and Ws) and auxin efflux defective mutants *eir1-1*
and *mdr1-1* grown in 3%Glc alone or in combination with different doses of MeJA. The
graph represents the average of 5 biological replicates consisting of 25 seedlings and error
bars represent SE. 5-day-old 1/2 MS grown Col-0 and mutant seedlings were transferred to
treatment media and phenotypes were analysed at 12th day. Asterisks indicate a significant
difference in the studied parameter (P <0.05, paired two-tailed student's t-test; * control vs
treatment and 'a' WT vs mutant).

Figure 4. Components of auxin signalling lie downstream to MeJA-mediated branching
angle. (A) Phenotype of light grown 12-day-old Arabidopsis Col-0, *tir1-1*, *axr1-3* and *axr2-1*
seedlings grown in 3%Glc alone or in combination with different doses of MeJA. **(B-C)**
Distribution of branching angle in seedlings of auxin signalling defective mutants *tir1-1*,
axr1-3 and *axr2-1* grown in 3%Glc alone or in combination with different doses of MeJA.
The graph represents the average of 4 biological replicates consisting of 25 seedlings and
error bars represent SE. 5-day-old 0.5X MS grown Col-0 and mutant seedlings were
transferred to treatment media and phenotypes were analysed at 12th day. Asterisks indicate a
significant difference in the studied parameter (P <0.05, paired two-tailed student's t-test; *
control vs treatment and 'a' WT vs mutant).

Figure 5. Transcriptional regulation of *CYP79B2*, *LAZY2* and *LAZY4* by MYC2. (A-C)
RT-qPCR showing the expression of *CYP79B2, LAZY2* and *LAZY4* in 1/2 MS media in Col-0
and *myc2myc3myc4* seedlings. The graph represents the average of 3 biological replicates
consisting of 25 seedlings and error bars represent SE. **(D)** ChIP-qPCR showing the
enrichment of *CYP79B2* promoter fragments with 35S::MYC2::GFP in all two regions. The
plot shows average of 2 biological replicates. **(E)** ChIP-qPCR showing the enrichment of
LAZY2 promoter fragments with MYC2 in all two regions. **(F)** ChIP-qPCR showing the
enrichment of *LAZY4* promoter fragments with MYC2. Fold enrichment of promoter
fragments was calculated by comparing samples treated without or with anti-GFP antibody.
Untransformed Col-0 was taken as a negative genetic control. The plot shows average of 3
biological replicates. *ORA59* and *ATXR6* were used as positive and negative controls,
respectively. Asterisks indicate a significant difference in the studied parameter. Error bars
represent SE. (P <0.05, paired two-tailed student's t-test; * WT vs overexpression).

Figure 6. MeJA controls lateral auxin redistribution in LRs. (A) Stage II LR of
DR5::GFP seedling treated with 1/2 MS and in combination with MeJA (10 µM). 5-day-old
1/2 MS grown *Arabidopsis* DR5::GFP seedlings were transferred to 1/2 MS alone or in
combination with MeJA. At 12th day, stage II lateral root were assessed under confocal
microscope. Scale bar: 20 µm. The data was repeated three times, with 8-9 roots imaged
every time. Solid white arrowheads depict strong GFP signal at the LR tip and dotted white
arrowheads depict reduced GFP signal. **(B)** Upper panel showing stage II LR of PIN2::PIN2-
eGFP seedlings treated with 1/2 MS and in combination with MeJA (10 µM). 5-day-old 1/2
MS grown Col-0 seedlings were transferred to treatment media. Lower panel showing
enlarged view of 1/2 MS and MeJA treated PIN2::PIN2-eGFP LR with diminished PIN2

activity from the upper epidermal cell profile. **(C)** The graph represents the cell length of 1st
two upper and lower epidermal cells of LRs treated with 1/2 MS alone and in combination
with 10 µM MeJA. The data shown is an average of two biological replicates. (P <0.05,
paired two-tailed student's t-test; * upper and lower epidermal cell profile; ** control vs
treatment). The data was repeated two times with 10-12 roots imaged.
1032

Figure 7. Light modulation of *Arabidopsis* root branching angle. (A) Phenotype of light
grown 12-day-old *Arabidopsis* Col-0, *phyA201phyB5* and *hy5-1* seedlings treated with
0.5%Glc and 3%Glc alone or in combination with 10 µM MeJA. **(B)** Distribution of
branching angle in seedlings of Col-0 and light signalling mutants *phyA-201, phyB-5,*
phyA201B5 and *hy5-1* grown in different concentrations of Glc (0.5%, 3%) alone or in
combination with MeJA (10 µM). 5-day-old 1/2 MS grown Ler and mutant seedlings were
transferred to treatment media and phenotypes were analysed at 12th day. The graph
represents the average of 5 biological replicates consisting of 25 seedlings and error bars
represent SE. Asterisks indicate a significant difference in the studied parameter (P <0.05,
paired two-tailed student's t-test; * control vs treatment and 'a' WT vs mutant). **(C)** RT-
qPCR showing the expression *LOX3* in 1/2 MS media under continuous light and dark for 6
days in Ler and *hy5-1* seedlings. **(D)** RT-qPCR showing the expression *LOX3* in 1/2 MS
media under dark (D) and dark (D) to light (L) transition for 6 hours in Ler and *hy5-1*
seedlings. The graph represents the average of 3 biological replicates, error bars represent SE.
Asterisks indicate a significant difference in the studied parameter (P <0.05, paired two-tailed
student's t-test; * control vs treatment and ** WT vs mutant). **(E)** Model depicting interaction
between light, Glc, JA and auxin signalling in regulating branching angle of *Arabidopsis*
LRs.
1051

1052 **Supplementary data**
1053
Figure S1. MeJA regulation of branching angle of *Arabidopsis* roots. (A and B) Average
branching angle of 12-day-old Col-0 and JA signalling mutants grown in different doses of
MeJA. The data represents the average of 4 biological replicates consisting of 25 seedlings
and error bars represent SE. **(C)** Distribution of branching angle of Col-0 and different *jaz*
seedlings grown in different doses of MeJA. The data represents the average of 3 biological
replicates consisting of 25 seedlings and error bars represent SE. 5-day-old 1/2 MS grown
Col-0 and mutant seedlings were transferred to treatment media and phenotypes were
analysed at 12th day. Asterisks indicate a significant difference in the studied parameter
(P <0.05, paired two-tailed student's t-test; * control vs treatment and   WT vs mutant).
1063
Figure S2. MeJA regulation of branching angle of *Arabidopsis* roots. (A) A representative
image of 20 days after germination (DAG) of Col-0 and JA biosynthesis and signalling
mutants *jar1-11* and *myc2myc3myc4* grown in 1/2 MS medium in cylindrical tubes. **(B)**
Average branching angle of 20 DAG of Col-0, *jar1-11* and *myc2myc3myc4* grown in 1/2 MS
medium in cylindrical tubes. The data represents images from 2 biological replicates, each
having 5 plants (P <0.05, paired two-tailed student's t-test; * WT vs mutant).
1070

1071 **Figure S3. MeJA-Glc regulation of branching angle of *Arabidopsis* roots. (A)** Average
1072 branching angle of 12-day-old Col-0 seedlings grown in different concentrations of Glc
1073 (0.5%, 3%) and in combination with 10 µM MeJA. The data represents the average of 7
1074 biological replicates consisting of 25 seedlings and error bars represent SE (P<0.05, Student's

t-test; * control (0.5%Glc) vs treatment. **(B)** Phenotype of Col-0 seedlings grown in different concentrations of Glc (0.5%; 28 mM and 5%Glc; 278 mM) and in combination with 10 µM MeJA. (C) Distribution of branching angle of seedlings of Col-0 grown in different different concentrations of Glc (0.5%, 3% and 5%) and in combination with MeJA. The data represents the average of 4 biological replicates consisting of 25 seedlings and error bars represent SE. Asterisks indicate a significant difference in the studied parameter ($P<0.05$,Student'st-test;*control (0.5%) vs treatment).

Figure S4. MeJA-Glc regulation of branching angle of *Arabidopsis* roots. (A) 3 individual biological replicates of Jas9-VENUS seedlings treated with increasing concentrations of Glc (0%Glc, 0.5%Glc, 3%Glc) along with 10 µM MeJA. **(B)** RT-qPCR showing the expression JA signalling genes (*JAZ1, JAZ2, JAZ3, JAZ6, JAZ9, AOS, MYC2, ERF1* and *ORA59*) in Col-0 seedlings grown in 0%Glc and 0%Glc+ 10 µM MeJA. 5-day-old 1/2 MS grown Col-0 seedlings were first starved for 24 hrs and transferred to liquid treatment media containing 10 µM MeJA and analysed after 3hrs. The data represents the average of 4 biological replicates consisting of 25 seedlings and error bars represent SE ($P <0.05$, paired two-tailed student's t-test; * control (0%Glc) vs treatment (0%Glc+MeJA).

Figure S5. Role of different components of Glc signalling in regulation of branching angle of *Arabidopsis* roots. **(A)** Average branching angle of 12-day-old Col-0 and Glc-dependent TOR signaling mutant *tor35-7RNAi* grown in different concentrations of Glc (0.5%, 3%) and in combination with MeJA. The data represents the average of 4 biological replicates consisting of 25 seedlings and error bars represent SE. ($P <0.05$, Student's t-test; * control vs treatment ** 0.5%Glc vs 3%Glc+10MeJA and ▲ wild-type vs mutant). **(B and C)** Distribution of branching angle in seedlings of Col-0 and RGS-dependent signalling mutants *rgs1-1, rgs1-2, gpa1-3* and *gpa1-4* grown in different concentrations of Glc in combination with MeJA. The data represents the average of 4 biological replicates consisting of 25 seedlings and error bars represent SE.

Figure S6. Glc regulates branching angle of LRs via TOR-mediated signalling pathway. 3 individual biological replicates of Jas9-VENUS seedlings treated without and with Glc (3%Glc) along with 10 µM MeJA, 10 µM AZD and a combination of 10 µM MeJA and 10 µM AZD (related to Figure 2J, K).

Figure S7. Heat map of JA signalling genes upon treatment with 0%Glc+10µM MeJA, 3%Glc and 3%Glc+10µM MeJA. The fold change value of genes (+/-1.5 fold) treated with 0%Glc+10µM MeJA was considered as 1. The fold change values of genes in 3%Glc and 3%Glc+ 10 µM MeJA was normalised (as % change) with those of 0%Glc+10µM MeJA. * indicates expression values of genes that are significantly altered. The data is the average of 2 biological replicates. Heat map was generated using MeV.

Figure S8. Role of auxin machinery in controlling JA-mediated branching angle. (A) Average branching angle of auxin influx defective mutant *aux1-7* grown in 3%Glc alone or in combination with MeJA. **(B)** Distribution of branching angle of auxin influx defective mutant *lax3* grown in 3%Glc alone or in combination with different doses of 10 µM MeJA **(C)** Average branching angle of auxin efflux defective mutants *eir1-1* and *mdr1.1* grown in 3%Glc alone or in combination with 10 µM MeJA. **(D)** Distribution of branching angle of auxin efflux defective mutants *pin3-4, pin4-3* and *pin7-2* grown in 3%Glc alone or in combination with different doses of MeJA. **(E and F)** Average branching angle of auxin

signalling mutants *tir1- 1*, *axr1-3* and *axr2-1* grown in 3%Glc alone or in combination with MeJA. 5-day-old 1/2 MS grown Col-0 and mutant seedlings were transferred to treatment media and phenotypes were analysed at 12th day. The data represents the average of 4 biological replicates consisting of 25 seedlings and error bars represent SE. Asterisks indicate a significant difference in the studied parameter (P <0.05, paired two-tailed student's t-test; * control vs treatment and ▲WT vs mutant).

Figure S9. Role of PIN3 in JA mediated regulation of branching angle. (A) Stage II LR of PIN3::PIN3-eGFP seedlings treated with 1/2 MS alone or in combination with 10 μM MeJA. **(B)** Quantification of fluorescence of stage II PIN3::PIN3-eGFP LRs treated with 1/2 MS alone or in combination with 10 µM MeJA. The graph represents average of 2 biological replicates with 10 roots imaged in each treatment.

Figure S10. Role of light signalling in controlling JA-mediated branching angle. (A) Phenotype of light and dark adapted Col-0 and *myc2myc3myc4* seedlings in different concentrations of Glc (0.5%, 3%) alone or in combination with MeJA (10 µM). **(B)** Distribution of branching angle in light and dark adapted Col-0 and *myc2myc3myc4* seedlings in different concentrations of Glc (0.5%, 3%) alone or in combination with MeJA (10 µM). The data represents the average of 6 biological replicates consisting of 25 seedlings and error bars represent SE. Asterisks indicate a significant difference in the studied parameter (P <0.05, paired two-tailed student's t-test; * control vs treatment and ▲WT vs mutant).

Figure S11. Role of light on auxin distribution to regulate branching angle. Distribution of branching angle in light and dark adapted Col-0 and *aux1-7* seedlings on ½ MS alone and in combination with MeJA (10µM). The data represents average of 4 biological replicates consisting of 25 seedlings and error bars represent SE. Asterisks indicate a significant difference in the studied parameter (*P* <0.05, Student's t-test; *control vs treatment and ▲WT vs mutant).

Supplementary Table 1: Primers used in the study.

Figure 1. MeJA decreases branching angle of Arabidopsis roots. (A) Phenotype of light grown 12-day-old Arabidopsis Col-0 seedlings grown on different doses of MeJA. **(B)** Average branching angle of 12-day-old Col-0 seedlings on different doses of MeJA. Data are mean-SE of 3 biological replicates with 25 seedlings. **(C)** Distribution of branching angle of 12-day-old Col-0 Arabidopsis seedlings on different doses of MeJA. Data are mean-SE of 3 biological replicates with 25 seedlings. **(D)** Comparison of the effects of different phytohormones (ACC, ABA, BR, GA) on Col-0 seedlings to determine their roles in branching angle. Data are mean-SE of 3 biological replicates with 25 seedlings. **(E)** Phenotype of light grown 12-day-old Arabidopsis Col-0, *jar1-11*, *coi1*, *jin1-9* and *myc2myc3myc4* seedlings grown on different doses of MeJA. **(F)** Distribution of branching angle in Col-0, *jar1-11* seedlings grown on different doses of MeJA. Data are mean-SE of 3 biological replicates with 25 seedlings. **(G)** Distribution of branching angle in seedlings of Col-0, *coi1* mutants grown on different doses of MeJA. Data are mean-SE of 4 biological replicates with 25 seedlings. **(H-I)** Distribution of branching angle in seedlings of Col-0, *jin1-9, myc2myc3myc4* mutants grown on different doses of MeJA. Data are mean-SE of 3 biological replicates with 25 seedlings. In A-I, 5-day-old 1/2 MS grown Col-0 seedlings were transferred to different doses of MeJA and phenotypes were analysed at 12[th] day. Asterisks

indicate a significant difference in the studied parameter (P <0.05, paired two-tailed student's t-test; * control vs treatment and 'a' WT vs mutant).

Figure 2. Glc antagonizes MeJA-induced branching angle in Arabidopsis roots. (A) Phenotype of Col-0 seedlings grown in different concentrations of Glc (0.5%; 28 mM and 3%Glc; 167 mM) and in combination with 10 µM MeJA. **(B)** Distribution of branching angle in Col-0 seedlings grown in different concentrations of Glc and in combination with MeJA. The graph represents the average of 7 biological replicates having 25 plants in each replicate and error bars represent SE. **(C)** Distribution of branching angle in Col-0 seedlings grown in different concentrations of Man in combination with 1%Glc. The graph represents the average of 3 biological replicates consisting of 25 seedlings and error bars represent SE. **(D)** Phenotype of Col-0 and *sweet11sweet12* seedlings grown in different concentrations of Glc. **(E)** Distribution of branching angle in Col-0 and *sweet11sweet12* seedlings grown in different concentrations of Glc. The graph represents the average of 3 biological replicates having 50 plants in each replicate and error bars represent SE. In **A-E**, 5-day-old 1/2 MS grown seedlings were transferred to Glc and MeJA containing treatment media and phenotypes were analysed at 12th day. Asterisks indicate a significant difference in the studied parameter (*P* <0.05, Student's t-

test; * control vs treatment and 'a' WT vs mutant. **(F)** Western blot analysis of total protein extracts of Jas9-VENUS seedlings treated for 3 hours with 0%, 0.5% and 3%Glc in combination with 10 µM MeJA and probed with an anti-GFP antibody. The experiment was performed four times. **(G)** Quantification of Jas9-VENUS protein levels. The graph represents average of 4 biological replicates. **(H)** Phenotype of Col-0 and *TOR* signalling mutant *tor35-7 RNAi* seedlings grown in different concentrations of Glc alone and in combination with 10 µM MeJA. **(I)** Distribution of branching angle in seedlings of Col-0 and *tor35-7 RNAi* grown in different concentrations of Glc alone or in combination with 10 µM MeJA. 5-day-old 1/2 MS grown Col-0 and *tor35-7* seedlings were transferred to treatment media and phenotypes were analysed at 12[th] day. The graph represents the average of 4 biological replicates consisting of 25 seedlings and error bars represent SE. Asterisks indicate a significant difference in the studied parameter ($P < 0.05$, Student's t-test; * control vs treatment, and 'a' WT vs mutant). **(J)** Western blot analysis of total protein extracts of Jas9-VENUS seedlings treated for 3 hours with 0%, 0.5% and 3%Glc alone or in combination with 10 µM MeJA and 10 µM AZD-8055 for 3h and probed with an anti-GFP antibody. The experiment was performed four times. **(K)** Quantification of Jas9-VENUS protein levels. The graph represents average of 4 biological replicates. Asterisks indicate a significant difference in the studied parameter ($P < 0.05$, paired two-tailed student's t-test; * control vs treatment and ** 0%Glc+AZD vs 3%Glc+AZD.

Figure 3. Components of auxin biosynthesis and transport lie downstream to MeJA-mediated branching angle. (A) Distribution of branching angle in Col-0 seedlings grown in 3%Glc alone or in combination with MeJA (10 µM) and different doses of NPA (100 nM to 1 µM). The graph represents the average of 4 biological replicates consisting of 25 seedlings and error bar represent SE. (B) Distribution of branching angle in seedlings of auxin influx defective mutant *aux1-7* grown in 3%Glc alone or in combination with different doses of MeJA. The graph represents the average of 3 biological replicates consisting of 25 seedlings and error bars represent SE. (C) Phenotype of light grown 12-day-old Arabidopsis Col-0, *aux1-7* and *eir1-1* seedlings grown in 3%Glc alone or in combination with different doses of MeJA. (D) Phenotype of light grown 12-day-old Arabidopsis Ws and *mdr1-1* seedlings grown in 3%Glc alone or in combination with different doses of MeJA. (E) Distribution of branching angle in seedlings of WT (Col-0 and Ws) and auxin efflux defective mutants *eir1-1* and *mdr1-1* grown in 3%Glc alone or in combination with different doses of MeJA. The graph represents the average of 5 biological replicates consisting of 25 seedlings and error bars represent

SE. 5-day-old 1/2 MS grown Col-0 and mutant seedlings were transferred to treatment media and phenotypes were analysed at 12th day. Asterisks indicate a significant difference in the studied parameter (P <0.05, paired two-tailed student's t-test; * control vs treatment and 'a' WT vs mutant).

Figure 4. Components of auxin signalling lie downstream to MeJA-mediated branching angle.
(A) Phenotype of light grown 12-day-old Arabidopsis Col-0, *tir1-1*, *axr1-3* and *axr2-1* seedlings grown in 3%Glc alone or in combination with different doses of MeJA. **(B-C)** Distribution of branching angle in seedlings of auxin signalling defective mutants *tir1-1*, *axr1-3* and *axr2-1* grown in 3%Glc alone or in combination with different doses of MeJA. The graph represents the average of 4 biological replicates consisting of 25 seedlings and error bars represent SE. 5-day-old 0.5X MS grown Col-0 and mutant seedlings were transferred to treatment media and phenotypes were analysed at 12th day. Asterisks indicate a significant difference in the studied parameter (P <0.05, paired two-tailed student's t-test; * control vs treatment and 'a' WT vs mutant).

Figure 5. Transcriptional regulation of *CYP79B2*, *LAZY2* and *LAZY4* by MYC2. (A-C) RT-qPCR showing the expression of *CYP79B2, LAZY2* and *LAZY4* in 1/2 MS media in Col-0 and *myc2myc3myc4* seedlings. The graph represents the average of 3 biological replicates consisting of 25 seedlings and error bars represent SE. **(D)** ChIP-qPCR showing the enrichment of *CYP79B2* promoter fragments with 35S::MYC2::GFP in all two regions. The plot shows average of 2 biological replicates. **(E)** ChIP-qPCR showing the enrichment of *LAZY2* promoter fragments with MYC2 in all two regions. **(F)** ChIP-qPCR showing the enrichment of *LAZY4* promoter fragments with MYC2. Fold enrichment of promoter fragments was calculated by comparing samples treated without or with anti-GFP antibody. Untransformed Col-0 was taken as a negative genetic control. The plot shows average of 3 biological replicates. *ORA59* and *ATXR6* were used as positive and negative controls, respectively. Asterisks indicate a significant difference in the studied parameter. Error bars represent SE. (P <0.05, paired two-tailed student's t-test; * WT vs overexpression).

Figure 6. MeJA controls lateral auxin redistribution in LRs. (**A**) Stage II LR of DR5::GFP seedling treated with 1/2 MS and in combination with MeJA (10 µM). 5-day-old 1/2 MS grown Arabidopsis DR5::GFP seedlings were transferred to 1/2 MS alone or in combination with MeJA. At 12[th] day, stage II lateral root were assessed under confocal microscope. Scale bar: 20 µm. The data was repeated three times, with 8-9 roots imaged every time. Solid white arrowheads depict strong GFP signal at the LR tip and dotted white arrowheads depict reduced GFP signal. (**B**) Upper panel showing stage II LR of PIN2::PIN2-eGFP seedlings treated with 1/2 MS and in combination with MeJA (10 µM). 5-day-old 1/2 MS grown Col-0 seedlings were transferred to treatment media. Lower panel showing enlarged view of 1/2 MS and MeJA treated PIN2::PIN2-eGFP LR with diminished PIN2 activity from the upper epidermal cell profile. (**C**) The graph represents the cell length of 1[st] two upper and lower epidermal cells of LRs treated with 1/2 MS alone and in combination with 10 µM MeJA. The data shown is an average of two biological replicates. (P <0.05, paired two-tailed student's t-test; * upper and lower epidermal cell profile; ** control vs treatment). The data was repeated two times with 10-12 roots imaged.

Figure 7. Light modulation of Arabidopsis root branching angle. (A) Phenotype of light grown 12-day-old Arabidopsis Col-0, *phyA201phyB5* and *hy5-1* seedlings treated with 0.5%Glc and 3%Glc alone or in combination with 10 µM MeJA. (B) Distribution of branching angle in seedlings of Col-0 and light signalling mutants *phyA-201, phyB-5, phyA201B5* and *hy5-1* grown in different concentrations of Glc (0.5%, 3%) alone or in combination with MeJA (10 µM). 5-day-old 1/2 MS grown Ler and mutant seedlings were transferred to treatment media and phenotypes were analysed at 12th day. The graph represents the average of 3 biological replicates consisting of 25 seedlings and error bars represent SE. Asterisks indicate a significant difference in the studied parameter (P <0.05, paired two-tailed student's t-test; * control vs treatment and 'a' WT vs mutant). (C) RT-qPCR showing the expression *LOX3* in 1/2 MS media under continuous light and dark for 6 days in Ler and *hy5-1* seedlings. (D) RT-qPCR showing the expression *LOX3* in 1/2 MS media under dark (D) and dark (D) to light (L) transition for 6 hours in Ler and *hy5-1* seedlings. The graph represents the average of 3 biological replicates, error bars represent SE. Asterisks indicate a significant difference in the studied parameter (P <0.05, paired two-tailed student's t-test; * control vs treatment and ** WT vs mutant). (E) Model depicting interaction between light, Glc, JA and auxin signalling in regulating branching angle of Arabidopsis LRs.