1   **Red deer *Cervus elaphus* blink more in larger groups**

3   Zeke W. Rowe*, Joseph H. Robins* and Sean A. Rands†

5   School of Biological Sciences, University of Bristol, Life Sciences Building, Tyndall Avenue,

6   Bristol BS8 1TQ, United Kingdom

8   ORCID: ZWR: 0000-0002-9617-5539

9   JHR: 0000-0001-8980-8150

10   SAR: 0000-0002-7400-005X

* JR and ZR are considered as joint first authors

† author for correspondence: sean.rands@bristol.ac.uk

ABSTRACT

Most animals need to spend time being vigilant for predators, at the expense of other

activities such as foraging. Group-living animals can benefit from the shared vigilance effort

of other group members, with individuals reducing personal vigilance effort as group size

increases. Behaviours like active scanning or head lifting are usually used to quantify

vigilance, but may not be accurate measures as the individual could be conducting them for

other purposes. We suggest that measuring an animal's blinking rate gives a meaningful

measure of vigilance: increased blinking implies reduced vigilance, as the animal cannot

detect predators when its eyes are closed. We demonstrate that as group size increases in

red deer, individuals increase their blink rate, confirming the prediction that vigilance should

decrease. Blinking is a simple non-invasive measure, and offers a useful metric for

assessing the welfare of animals experiencing an increase in perceived predation risk or

other stressors.

INTRODUCTION

Most animal species spend some part of their lives aggregated together in groups, and many benefits have been proposed and tested for this behaviour [1,2]. For prey species, grouping behaviour can offer protection from predators through both the dilution of individual risk if an attack occurs [3–5] and an increase in the chance of successfully detecting an approaching predator due to the combined vigilance effort of the group [5–7]. If an animal is being actively vigilant, it may be unable to conduct other important behaviours (like foraging or resting) at the same time as the vigilance behaviour. Group membership means that vigilance can be pooled among the group members, which could mean that each individual can spend less time being vigilant and more time conducting other fitness-enhancing behaviours. A rich body of theory and research has explored how group size and individual vigilance effort are related [8–11], with much of it focussing on the prediction that individual vigilance effort will decrease as the group becomes larger. This prediction requires each individual to show a trade-off between vigilance and other behaviours, where being actively vigilant either cannot occur at the same time as other behaviours, or leads to a reduction in the efficiency of other behaviours that are conducted at the same time as being vigilant.

Vigilance is usually assumed to be occurring when an animal is actively scanning its surrounding environment with its head upwards, although there is no obvious consensus in how vigilance is defined in any particular species (see [12] for discussion of this problem in studies on primates). Although scanning behaviour is likely to stop an animal from actively collecting food, this head-up activity may not completely interfere with simultaneous conducted behaviours, such as chewing or social interaction. If a behaviour that is recorded as vigilance allows an individual to do other things at the same time, then we may be falsely assuming that this behaviour incurs the time and attention costs that are associated with vigilance [13]. Without careful experimentation, it is difficult to assess how much of an individual's attention is devoted to vigilance when we observe scanning or other forms of vigilance-like behaviour, which may add to the huge variation (*e.g.* [8]) in whether a study demonstrates that individual vigilance is related to group size or not.

Although it is difficult to define exactly when an individual is being vigilant, we may instead be able to define when it is *not* able to be vigilant. Blinking (the temporary closure of both eyes, involving movements of the upper and lower eyelids [14]) is a good example of an activity where an individual is momentarily unable to visually scan the environment. It is an

essential maintenance behaviour to keep the eyes moist and clean [15], and is conducted

tens of times every minute in some species of diurnal mammals [16,17] and birds [18]. This

frequency means that although a blink takes only a fraction of a second, the sum of this loss

of visual information could be substantial for the individual. In humans, spontaneous blinking

is accompanied by attentional suppression, where the individual experiences a blackout in

visual attention for the duration of the blink, meaning that there is no awareness of the

temporary blindness and lack of visual information whilst the blinking is occurring [19,20].

Blinking suppresses activity in both the visual cortex and other areas of the brain that are

associated with awareness of environmental change [21]. If we assume that other animals

show similar attentional suppression, then they are essentially blind and unaware of

changes in their visual environment during each blink, which in turn means that they cannot

be vigilant for predators.

An individual's blink rate therefore presents a trade-off between the physiological benefits of

blinking and the loss of visual information during the blink [15]. If an animal needs to

dedicate more time to vigilance in a risky environment, then it has to reduce or suppress

blinking to accommodate this increased vigilance. This is anecdotally demonstrated in

American crows *Corvus brachyrhynchos*, which reduce their blink rates when looking at

potentially dangerous stimuli [22], and in horses *Equus caballus*, which decrease their

spontaneous blink rate in response to stress-inducing stimuli [23]. This link between blink

rate and vigilance implies that blink rate will also be related to group size. As group size

increases, theory predicts that individual vigilance can be reduced [5], and so any

requirement to suppress blinking will be relaxed. Blink rate may therefore increase with an

increase in group size. Evidence supporting this prediction is anecdotal: a comparison of

chickens *Gallus gallus* feeding solitarily or in pairs showed a non-significant increase in blink

rate in the group-feeding birds [24], while a comparison of the blink rates of olive baboons

Papio anubis showed individuals in a small group blinked less than those in a large group

(although the two groups were studied in different years). Here, we test this hypothesis by

observing the blink rate in a herd of red deer *Cervus elaphus*. Red deer are a prey species

that spend most of their lives in dispersed groups, and females have previously been shown

to reduce their vigilance behaviour as group size increases [25,26]. Because they show an

increase in vigilance in smaller groups, we would therefore predict that they should also

reduce their blink rate to avoid losing visual information about their environment.

METHODS

Study area, time and subjects

This observational study was conducted on the herd of red deer within the 40.5 hectare deer park in Ashton Court Estate, Bristol, England (51.4440° N, 2.6378° W), which is composed mainly of open grassland, with scattered forestry and a small area of water. The herd is managed by the Bristol City Council and consisted of c. 110 individuals of varying age and sex. The enclosure is open to the public outside of the rutting season, so the deer are habituated to both dogs and humans. Our observations were conducted during the red deer rutting season; therefore, observations were restricted between 1200 and 1630 h so they were outside of the dawn and dusk peaks of regular rutting activity [27].

Behavioural sampling and observations

A random individual was selected as carried out in previous research on this herd [28]. A random number was drawn between 1-99, with the focal deer being selected by counting linearly from leftmost or rightmost visible deer (the direction was determined by a coin toss). If the random number was higher than the number of visible deer, it was discarded, and the process started again. The selected individual was observed and dichotomously aged (mature or young) and sexed (male or female). The individual was sexed by the presence or absence of antlers, as after one year of age only males have antlers [29]. Age identification was done by the size and morphology of the individual (larger individuals were older). If the individual was observed suckling then its selection was discarded and randomisation was done again; as these young individuals are hard to sex and exhibit behaviours uncommon to the rest of the herd [27]. An estimation of the abundance of each demographic was also recorded, this was done by every individual being aged, sexed and counted on three different days. The average of these demographics was then calculated, and used in the pseudoreplication analyses presented.

Prior to the observations being conducted, the observers were trained in identifying the recorded behaviours. A blink was defined as a rapid full closure of the eye. Group size was defined as the individuals aggregated in no more than five body lengths away from each other. Before starting any set of observations, the observers waited 10 minutes at the site to habituate the deer. All observations were conducted approximately 10-100 m of the focal individuals through a 30× zoom spotting scope (Avian ED82 Magnesium Scope). The

observations for a single selected individual were recorded for a maximum of 10 minutes. At

the start of each minute the group size was counted, with blink rate (blinks per minute) being

continuously recorded for each minute. If the deer's eye(s) were obstructed or there was a

human/animal disturbance the observations were stopped with the current minute's

observations being disregarded. A total of 75 observations were collected using this method.

Statistical analysis

Data were recorded as blinks per minute with group size recorded for that minute.

Individuals were recorded a mean of 4.9 (± 2.0 SD) minutes before the data collection had to

be discontinued due to the observer's view of the eyes being obstructed. For each individual,

the mean number of blinks in a minute was calculated, as was the mean group size per

minute over the observed period. Analyses described below were conducted using these

mean blinks per minute and mean group sizes. To compensate for any unevenness caused

by some mean values being based on more observations than others, we conducted the

same analyses using just the first minute of data for all individuals (see the Supplementrary

Material): these data gave qualitatively similar results to those reported for the analysis

involving mean group sizes, and are not discussed further.

All analyses were conducted in R 3.6.1 [30]. A linear model was constructed where the

natural logarithm of mean blink rate was described by the natural logarithm of group size,

and the maturity and sex of the focal individual. Logarithms were used to satisfy model

conditions. A full model including interactions was initially considered, but no interaction

terms were significant and so the basic additive model with the three explanatory variables

was used.

Although attempts were made to avoid replicating observations on the same individual

during an observational day, it is likely that the 75 datapoints include some repeat

observations of the same individual, given that the herd had 95 observable individuals (of

which 11 were estimated to be mature males, 23 were young males, 26 were young females

and 37 were mature females), and individuals within each class could not be individually

identified accurately. To explore the potential effects of this pseudoreplication on the results,

we conducted simulation analyses where identities were randomly allocated to individuals in

each age class, and all but one datapoint for each 'assumed individual' was removed from

the dataset (code presented in the Supplementary Information). A linear model was then run

on this filtered subsample of the dataset, and the significance value was harvested. By

conducting 100,000 independent repeats of this resampling, we could identify how likely our

dataset was to give a significant result (assuming significance was set at the standard

threshold of $p = 0.05$) if we had only sampled each individual once.

RESULTS

We randomly selected individual deer within a herd of c. 110 individuals, and recorded their

blink rate over a maximum of ten minutes. At the same time, we recorded the mean size of

the local group that they were in, as well as the sex and maturity (young or adult) of the

observed individual. A linear model explained the data well ($F_{3,71} = 59.17$, $p < 0.001$,

adjusted $r^2 = 0.702$, parameter estimates given in Supplementary Material). Blink rate

increased with group size ($F_{1,72} = 172.94$, $p < 0.001$; figure 1), and adults blinked more than

young deer ($F_{1,72} = 4.47$, $p = 0.038$; figure 1). There was no relationship between sex and

blinking ($F_{1,72} = 0.98$, $p = 0.755$).

189
190

Figure 1. Scatterplot showing that blink rate increases with group size and maturity (where

orange triangles and the fitted solid line represent young individuals, and black circles and

the fitted dashed line represent adults).

Considering pseudoreplication of unidentifiable individuals, resampling showed that group

size remained significant (at $p < 0.001$) in 100% of simulations. There was a significant (at p

< 0.05) effect of sex in only 0.27% of simulations, whilst maturity was significant in 42.97% of

simulations. So, the effect of maturity observed could potentially have been an effect of

resampling the same individuals, but the increasing blink rate in response to increasing

group size is unlikely to have been affected by any pseudoreplication caused by repeated

sampling of the same individual.

DISCUSSION

Our results demonstrate that blinking increases as group size increases. Given that blinking

interferes with vigilance behaviour, and that individual vigilance is predicted to decrease as

groups get larger, this supports our argument that the blink rate represents a trade-off

between gaining visual information through vigilance and the physiological benefits of

blinking.

We argued earlier that observed behaviours that are typically recorded as vigilance (such as

holding a head upright or active scanning) may not be conducted solely for vigilance.

Similarly, blinking may not solely be a maintenance behaviour that is traded-off against

being able to collect visual information. Blinking may include a social element, and rhesus

macaques *Macaca mulatta* are able entrain their blink rate in response to social cues [31].

However, we suggest that the opposite may be true in red deer: scanning behaviour has

been shown to only change in response to group size in females [25,26], but we found that

both males and females increased their blink rate as group size increased. In this species,

scanning behaviour may be influenced by social behaviour as well as vigilance, but our

blinking results may demonstrate that the proportion of scanning time that an individual

allocates to vigilance is related to the size of the pooled group effort. Other aspects of social

behaviour may also be important for determining blink rates, such as position within the

group. Theory predicts that individuals on the outside of the group should be more vigilant

than those in the middle, and anecdotal evidence from olive baboons suggests that

peripheral individuals may blink less [32]. It may be possible to test many of the standard

predictions connecting vigilance and group size (e.g. [8–11]) using blink rate as a proxy for

vigilance.

The blink rate may also be influenced by factors other than the size of the group, including

rainfall and wind (which have been shown to influence blink rate in captive grackles

Quiscalus mexicanus [33,34]). Similarly, the behaviour that an individual conducts

simultaneously to the blink may be important. Experiments in humans a mechanism
controlling the timing of blinks, which occurs to minimise the chance of missing crucial
information [15,35–38], with evidence of similar behaviours in rhesus macaques *Macaca*
mulatta [31]. Peafowl *Pavo cristatus* also time their blinks to coincide with gaze shifts [39]
while chickens blink more when feeding when compared to scanning [24], both minimising
the time where visual information cannot be collected from the environment. Therefore, if an
individual is moving, the timing and frequency of its blinks may reflect this movement.
Similarly, animals in different attentional states may change their frequency of blinking, such
as during sleep in herring gulls *Larus argentatus* [40].
242
Our results suggest that the measurement of blinking presents a simple and non-invasive
technique for observing attention that can be conducted remotely. Although we conducted
our sampling in the field, this could be done using video footage. Being able to analyse video
footage means that information about blink duration can also be collected, and previous
studies have demonstrated that this additional metric, which gives another measure of the
length of time when an animal cannot be vigilant, can also vary between individuals and
species [17,24,32–34]. Given that blinking has been shown to decrease under stressful
conditions [22,23], this simple technique could help us to understand the welfare
requirements of managed animals that normally live in social groups.
252

253
254
255
256 **ACKNOWLEDGMENTS**
257

258 This project was conducted by ZWR and JHR as their final year project in the University of
259 Bristol School of Biological Science's BSc degree programme. We thank the teaching lab
260 team in the University of Bristol School of Biological Sciences for help with equipment.
261 Ethical permission to conduct the study was given by the University of Bristol Animal Welfare
262 and Experimental Review Board (#UB/16/062).

263
264
265 **FUNDING**
266
267 SAR was supported a Natural Environment Research Council (UK) standard grant
268 (NE/P012639/1).
269
270

271 **AUTHOR CONTRIBUTIONS**
272
273 ZWR: Conceptualisation, experimental design, data collection, drafting, and editing.
274 JHR: Conceptualisation, experimental design, data collection, drafting, and editing.
275 SAR: Conceptualisation, supervision, experimental design, analysis, original draft, and
276 editing.
277
278
279 **SUPPLEMENTARY MATERIAL**
280
281 Two files are included: 'blinkdata.txt' is the raw data analysed, and details the mean blinks
282 per minute, mean group size, sex (f=female, m=male), and age (m=mature adult, y=young)
283 for each datapoint. 'analysis.R' presents the annotated R code used for analysis.
284
285
286 **REFERENCES**

1. Krause J, Ruxton GD. 2002 *Living in groups*. Oxford: Oxford University Press.

2. Ward A, Webster M. 2016 *Sociality: the behaviour of group-living animals*. Switzerland: Springer.

3. Bednekoff PA, Lima SL. 1998 Re–examining safety in numbers: interactions between risk dilution and collective detection depend upon predator targeting behaviour. *Proc R Soc B* **265**, 2021–2026. (doi:10.1098/rspb.1998.0535)

4. Hamilton WD. 1971 Geometry for the selfish herd. *J Theor Biol* **31**, 295–311. (doi:10.1016/0022-5193(71)90189-5)

5. Pulliam HR. 1973 On the advantages of flocking. *J Theor Biol* **38**, 419–422. (doi:10.1016/0022-5193(73)90184-7)

6. Lima SL. 1990 The influence of models on the interpretation of vigilance. In *Interpretation and explanation in the study of animal behavior, volume 2: explanation, evolution and adaptation* (eds M Bekoff, D Jamieson), pp. 246–267. Boulder: Westview Press.

7. McNamara JM, Houston AI. 1992 Evolutionarily stable levels of vigilance as a function of group size. *Anim Behav* **43**, 641–658.

8. Beauchamp G. 2019 On how risk and group size interact to influence vigilance. *Biol Rev* **94**, 1918–1934. (doi:10.1111/brv.12540)

9. Beauchamp G. 2008 What is the magnitude of the group-size effect on vigilance? *Behav Ecol* **19**, 1361–1368. (doi:10.1093/beheco/arn096)

10. Beauchamp G. 2003 Group-size effects on vigilance: a search for mechanisms. *Behav Process* **63**, 111–121.

11. Elgar MA. 1989 Predator vigilance and group size in mammals and birds: a critical review of the empirical evidence. *Biol Rev* **64**, 13–33. (doi:10.1111/j.1469-185X.1989.tb00636.x)

12. Allan ATL, Hill RA. 2018 What have we been looking at? A call for consistency in studies of primate vigilance. *Am J Phys Anthropol* **165**, 4–22. (doi:10.1002/ajpa.23381)

13. Treves A. 2000 Theory and method in studies of vigilance and aggregation. *Anim Behav* **60**, 711–722.

14. Blount WP. 1927 Studies of the movements of the eyelids of animals: blinking. *Q J Exp Physiol* **18**, 111–125. (doi:10.1113/expphysiol.1927.sp000426)

15. Ranti C, Jones W, Klin A, Shultz S. 2020 Blink rate patterns provide a reliable measure of individual engagement with scene content. *Sci Rep* **10**, 8267. (doi:10.1038/s41598-020-64999-x)

16. Stevens JR, Livermore A. 1978 Eye blinking and rapid eye movement: pulsed photic stimulation of the brain. *Exp Neurology* **60**, 541–556. (doi:10.1016/0014-4886(78)90009-2)

17. Tada H, Omori Y, Hirokawa K, Ohira H, Tomonaga M. 2013 Eye-blink behaviors in 71
species of primates. *PLoS One* **8**, e66018. (doi:10.1371/journal.pone.0066018)

18. Kirsten SJ, Kirsten EB. 1983 Spontaneous blink rates of birds. *Condor* **85**, 92–93.

19. Volkmann FC, Riggs LA, Moore RK. 1980 Eyeblinks and visual suppression. *Science*
207, 900–902. (doi:10.1126/science.7355270)

20. Riggs LA, Volkmann FC, Moore RK. 1981 Suppression of the blackout due to blinks.
Vision Res **21**, 1075–1079. (doi:10.1016/0042-6989(81)90012-2)

21. Bristow D, Haynes J-D, Sylvester R, Frith CD, Rees G. 2005 Blinking suppresses the
neural response to unchanging retinal stimulation. *Curr Biol* **15**, 1296–1300.
(doi:10.1016/j.cub.2005.06.025)

22. Cross DJ, Marzluff JM, Palmquist I, Minoshima S, Shimizu T, Miyaoka R. 2013 Distinct
neural circuits underlie assessment of a diversity of natural dangers by American crows.
Proc R Soc B **280**, 20131046. (doi:10.1098/rspb.2013.1046)

23. Merkies K, Ready C, Farkas L, Hodder A. 2019 Eye blink rates and eyelid twitches as a
non-invasive measure of stress in the domestic horse. *Animals* **9**, 562.
(doi:10.3390/ani9080562)

24. Beauchamp G. 2017 Half-blind to the risk of predation. *Front Ecol Evol* **5**, 131.
(doi:10.3389/fevo.2017.00131)

25. Childress MJ, Lung MA. 2003 Predation risk, gender and the group size effect: does elk
vigilance depend upon the behaviour of conspecifics? *Anim Behav* **66**, 389–398.
(doi:10.1006/anbe.2003.2217)

26. Lung MA, Childress MJ. 2007 The influence of conspecifics and predation risk on the
vigilance of elk (*Cervus elaphus*) in Yellowstone National Park. *Behav Ecol* **18**, 12–20.
(doi:10.1093/beheco/arl066)

27. Clutton-Brock TH, Guinness FE, Albon SD. 1982 *Red deer: behavior and ecology of two*
sexes. Chicago: Chicago University Press.

28. Rands SA, Muir H, Terry NL. 2014 Red deer synchronise their activity with close
neighbours. *PeerJ* **2**, e344. (doi:10.7717/peerj.344)

29. Mitchell B, Staines BW, Welch D. 1977 *Ecology of red deer: a research review relevant*
to their management in Scotland. Cambridge: Institute of Terrestrial Ecology.

30. R Development Core Team. 2019 *R: a language and environment for statistical*
computing. Vienna: R Foundation for Statistical Computing.

31. Ballesta S, Mosher CP, Szep J, Fischl KD, Gothard KM. 2016 Social determinants of
eyeblinks in adult male macaques. *Sci Rep* **6**, 38686. (doi:10.1038/srep38686)

32. Matsumoto-Oda A, Okamoto K, Takahashi K, Ohira H. 2018 Group size effects on inter-
blink interval as an indicator of antipredator vigilance in wild baboons. *Sci Rep* **8**, 10062.
(doi:10.1038/s41598-018-28174-7)

33. Yorzinski JL, Argubright S. 2019 Wind increases blinking behavior in great-tailed
grackles (*Quiscalus mexicanus*). *Front Ecol Evol* **7**, 330. (doi:10.3389/fevo.2019.00330)

34. Yorzinski JL. 2020 Blinking behavior in great-tailed grackles (*Quiscalus mexicanus*)
increases during simulated rainfall. *Ethology* **126**, 519–527. (doi:10.1111/eth.13003)

35. Nakano T, Yamamoto Y, Kitajo K, Takahashi T, Kitazawa S. 2009 Synchronization of
spontaneous eyeblinks while viewing video stories. *Proc R Soc B* **276**, 3635–3644.
(doi:10.1098/rspb.2009.0828)

36. Shin YS, Chang W, Park J, Im C-H, Lee SI, Kim IY, Jang DP. 2015 Correlation between
inter-blink interval and episodic encoding during movie watching. *PLoS One* **10**,
e0141242. (doi:10.1371/journal.pone.0141242)

37. Wiseman RJ, Nakano T. 2016 Blink and you'll miss it: the role of blinking in the
perception of magic tricks. *PeerJ* **4**, e1873. (doi:10.7717/peerj.1873)

38. Maffei A, Angrilli A. 2019 Spontaneous blink rate as an index of attention and emotion
during film clips viewing. *Physiol Behav* **204**, 256–263.
(doi:10.1016/j.physbeh.2019.02.037)

39. Yorzinski JL. 2016 Eye blinking in an avian species is associated with gaze shifts. *Sci
Rep* **6**, 32471. (doi:10.1038/srep32471)

40. Amlaner CJ, McFarland DJ. 1981 Sleep in the herring gull (*Larus argentatus*). *Anim
Behav* **29**, 551–556. (doi:10.1016/S0003-3472(81)80118-2)