Integrative inference of transcriptional networks in Arabidopsis yields novel ROS signalling regulators

Inge De Clercq[1,2+*], Jan Van de Velde[1,2,3+], Xiaopeng Luo[1,2], Li Liu[1,2,3], Veronique Storme[1,2], Michiel Van Bel[1,2], Robin Pottie[1,2], Dries Vaneechoutte[1,2,3], Frank Van Breusegem[1,2], Klaas Vandepoele[1,2,3*]

(1) Department of Plant Biotechnology and Bioinformatics, Ghent University, Technologiepark 71, 9052 Ghent, Belgium

(2) VIB Center for Plant Systems Biology, Technologiepark 71, 9052 Ghent, Belgium

(3) Bioinformatics Institute Ghent, Ghent University, Technologiepark 71, 9052 Ghent, Belgium

+ both authors contributed equally

* corresponding authors: Klaas.Vandepoele@psb.vib-ugent.be and Inge.DeClercq@psb.vib-ugent.be

ABSTRACT

Gene regulation is a dynamic process in which transcription factors (TFs) play an important

role to control spatiotemporal gene expression. While gene regulatory networks describe the

interactions between TFs and their target genes, our global knowledge about the complexity of

TF control for different genes and biological processes is incomplete. To enhance our global

understanding of regulatory interactions in *Arabidopsis thaliana*, different regulatory input

networks capturing complementary information about DNA motifs, open chromatin, TF binding

and expression-based regulatory interactions, were combined using a supervised learning

approach, resulting in an integrated gene regulatory network (iGRN) covering 1,491 TFs and

31,393 target genes (1.7 million interactions). This iGRN outperforms the different input

networks to predict known regulatory interactions and has a similar performance to recover

functional interactions compared to state-of-the-art experimental methods like yeast one-

hybrid and ChIP-seq. The iGRN correctly inferred known functions for 681 TFs and predicted new

gene functions for hundreds of unknown TFs. For regulators predicted to be involved in reactive

oxygen species stress regulation, we confirmed in total 75% of TFs with a function in ROS and/or

physiological stress responses. This includes 13 novel ROS regulators, previously not connected

to any ROS or stress function, that were experimentally validated in our ROS-specific phenotypic

assays of loss- or gain-of-function lines. In conclusion, the presented iGRN offers a high-quality

starting point to enhance our understanding of gene regulation in plants by integrating different

experimental data types at the network level.

INTRODUCTION

Accurate spatiotemporal gene expression is of key importance for plant development and molecular responses to environmental cues. Key factors controlling this process are transcription factors (TFs) which regulate their target genes by recognizing and binding short *cis*-regulatory DNA sequences called TF binding sites (TFBSs or DNA motifs) [1]. The full set of regulatory interactions between TFs and their target gene(s) forms a gene regulatory network (GRN) and depending on the underlying source of data, GRNs are composed of physical and/or regulatory protein-DNA interactions. Efficiently identifying regulatory interactions is pivotal to understand how a gene's transcriptional activity is controlled through the interplay of one or multiple regulators. Driven by different modes of gene duplication, specific TF families have been expanded within the green plant lineage, with the model species *Arabidopsis thaliana* containing around 1700 TFs [2]. Although specific regulatory networks and the associated TFs have been studied and are often conserved across species [3], still for hundreds of *Arabidopsis* TFs experimental information is missing [4].

Several studies have used high-throughput assays to systematically identify GRNs under specific conditions or in specific developmental and/or anatomical contexts in *Arabidopsis.* Examples include yeast one-hybrid (Y1H) screens to map process-specific GRNs as well as TF chromatin immunoprecipitation (ChIP) experiments describing regulatory interactions between TFs and target genes involved in flower development, light and clock signaling, hormone response, and (a)biotic stress response [5]. ChIP data revealed the high complexity of plant gene regulation and suggested that a single gene can potentially be regulated by up to 75 different TFs [6,7]. Open chromatin profiling, either using DNase-Seq or Assay for Transposase-Accessible Chromatin using sequencing (ATAC-Seq), allows studying the organization and dynamics of accessible chromatin regions, and provides a template to infer GRNs in different organs, in response to specific treatments or in different cell types [8-12]. Whereas all these methods aid the construction of GRNs by reporting physical protein-DNA interactions, a major challenge remains the identification of functional interactions, referring to the binding of a specific TF to a cis-regulatory sequence leading to a regulatory effect on a target genes' expression.

In this paper we present a network-based approach based on supervised learning for large-scale functional data integration to enhance our understanding of TF gene regulation in *Arabidopsis*, hence referred to as an integrative GRN (iGRN). Supervised learning approaches make efficient use of the available prior knowledge to train a machine learning algorithm, called

a classifier, for handling new data. Compared to unsupervised learning approaches, supervised

learning is considered more powerful when a large amount of existing knowledge is available to

guide the classification task, which here is the identification of functional TF-target

interactions[13]. Construction and validation of the iGRN, leveraging a diverse set of

complementary experimental datasets, revealed a strong enrichment for functional interactions

and a high predictive power to correctly infer functions for both functionally characterized and

uncharacterized TFs involved in a variety of biological processes. The latter is demonstrated in a

case study by the experimental validation of novel iGRN-based function predictions for TFs

involved in plant responses to oxidative stresses.

RESULTS

An integrated gene regulatory network for *Arabidopsis thaliana*

To generate an integrated gene regulatory network (*Arabidopsis* iGRN), different types of

complementary regulatory datasets were first processed, resulting in different input networks

(Table 1). In total seven input networks, encompassing information about DNA motifs, open

chromatin, TF binding and expression-based regulatory interactions, were generated by applying

different network inference and data analysis approaches (see Methods for underlying datasets;

Figure 1a). Subsequently, through the application of a supervised learning approach, a classifier

was built to identify those interactions from the input networks that most likely represent

functional interactions.

The first network is based on the presence of a TF motif, modelled using Position Weight

Matrices (PWMs), in the promoter of a target gene. A total of 1,260 high quality PWMs, covering

874 TFs, were mapped onto the *Arabidopsis* genome resulting in a TF - PWM target gene

network (called "PWM"). Using the genomic locations of publicly available DNaseI datasets the

PWM network was filtered resulting in a second open chromatin regulatory input network

(called "PWM in DHS"). DNaseI hypersensitive sites (DHS) are considered to be a proxy for open

chromatin and potential protein binding to the DNA [14]. As a third network, the set of PWMs was

also evaluated for evolutionary conservation across 13 dicotyledonous genomes using a

comparative motif mapping algorithm [15], which resulted in an evolutionarily conserved

regulatory network ("Conserved PWM"). To integrate experimental data, a fourth "TF ChIP"

network was created by combining experimentally determined TF binding occupancy profiles

from 27 TFs [7]. Next to a physical TF binding network, two co-expression-based regulatory

networks were also generated. Co-expression approaches measure similarity at the output level

of transcriptional regulation and as such can provide a complementary view compared to

promoter-centered regulatory networks. First, the Pearson correlation between a TF and a

target gene was used to build co-expression clusters around each TF, resulting in a fifth "CoE TF-

target" input network. Secondly, GENIE3, a reverse engineering tool for reconstructing

regulatory networks based on expression data, was used to build a sixth input network [16].

GENIE3 was among the best performing tools in reconstructing regulatory interactions in a

large-scale benchmark study [17]. The last network was generated using a hybrid approach of co-

expression and PWM based methods ("COE+PWM"). Initially a co-expression cluster was

constructed for each gene and PWM enrichment was subsequently performed. Only genes with

significantly enriched PWMs in their co-expression cluster were retained, resulting in a seventh

network where TF binding site information is combined with co-regulatory information (see

Methods).

As the different input networks capture different aspects of gene regulation, we next applied

supervised learning to construct, starting from these input networks, an integrative regulatory

network reporting functional interactions. A score for each possible TF-target interaction was

devised as a function of the feature-specific weights provided by the seven input networks. For

building a classifier, we compiled a gold standard dataset comprising interactions based on all

validated interactions in AtRegNet [18], an exhaustive literature-based network reporting

interactions in secondary cell wall development [19], and a large-scale data mining effort on 974

peer reviewed articles [20]. This gold standard entails interactions for 47 TF families involved in

different biological processes, with on average 11 interactions per TF (Table 1 and Supplemental

Table S1). Eighty percent of the interactions present in this gold standard were retained as

training dataset for the machine learning classification algorithms while the remaining twenty

percent was used as an evaluation set. Initially different classifiers were learned and their

performance to identify functional interactions in the evaluation set was assessed

(Supplemental Table S2). The classifier based on gradient boosting model using decision trees

showed the best performance, with an area under the curve (AUC) value of 0.77, and Precision

and Recall of 71% and 46% on the evaluation set, respectively (Supplemental Table S2). Applying

this classifier to the complete set of interactions present in the seven input networks resulted in

an integrative network of 1,709,871 interactions for 1,491 TFs, containing 31,393 target genes

(29,137 protein-coding and 2,256 non-coding genes; see Data file 1).

In order to obtain an unbiased performance estimate of the input networks compared to the

iGRN, all input networks were compared to the evaluation dataset of known interactions (20%

of the gold standard) and benchmarked measuring the recovery of true edges (referred to as the

recall, Figure 1b). Furthermore, we also determined the enrichment of a network towards the

evaluation dataset, which functions as a proxy for specificity [21]. Focusing on the different input

networks, Figure 1b shows that the network based on evolutionary conserved TF binding sites

("Conserved PWM") has the highest enrichment (6.4 fold, Supplemental Table 3). Evolutionary

conservation is known to be a strong predictor of functional regulatory elements explaining this

high enrichment [22,23]. The TF ChIP-seq network also performs well with both a high recovery

(29%) and enrichment (6.0 fold). This overlap can partially be explained by the fact that a subset

of the validated AtRegNet interactions in the gold standard are based on ChIP studies [7]. Other TF

motif mapping strategies recover larger fractions of true edges but suffer from low specificity,

resulting in low enrichment values. The simple mapping of TF motifs ("PWM") is known to result

in many false positives because TF binding sites are often short and typically contain some level

of degeneracy [11,24]. Both co-expression based networks do not consider TF binding site

information and are therefore able to also identify regulatory interactions for TFs lacking DNA

binding motif information. Both these networks perform relatively well: correlation-based co-

expression between TF and target genes, implemented in the "CoE TF-target" network, has the

third highest enrichment towards the evaluation dataset (10% recovery and 3.0 fold

enrichment), while GENIE3 displays a good balance between recovery of true edges and

specificity (Supplemental Table S3). Focusing on the integrative network (iGRN) shows it has the

strongest enrichment (8.8 fold) as well as a high recovery for true regulatory relationships (46%

recovery of interactions in the evaluation set), clearly outperforming the different

computational and experimental approaches used to construct the input networks.

Apart from determining the power of the iGRN network to recover known functional

interactions, we also measured the contribution of each of the seven input networks to the

iGRN. Ideally, a good integration approach should favor interactions from the input networks

with high enrichment values towards the evaluation set, as these would boost the recovery of

correct interactions. As reported in Figure 1b (brown and red bar charts), the evolutionary

conserved motifs ("Conserved PWM") and the "TF ChIP" network contributed the largest part of

their edges to the integrative network (91% and 80%, respectively), while networks showing low

specificity, such as "PWM", only contributed a small fraction of interactions (7%). This result

corroborates that the applied supervised approach successfully identified functional interactions

coming from high-quality input networks. Next, we measured which iGRN interactions are

supported by the different input networks (Figure 1b, red bar plot). Overall, the integrative

network has 70% of the edges supported by a PWM mapping, meaning that most regulatory

interactions are supported by the presence of TF binding site(s). Furthermore, 36-48% of the

interactions are supported by at least one of the two co-expression networks, indicating for

nearly half of all interactions a regulatory effect at the transcript level. Overall, 92% of the iGRN

interactions are supported by more than one input network (Figure S1), confirming both the

commonality and complementarity of the input networks.

iGRN interactions are complementary to TF regulatory events identified from

experimental methods

To assess whether the iGRN identifies functional TF-target gene interactions that are

supported by experimental data, detailed comparisons of the iGRN interactions were made

against three different datasets, capturing a variety of TFs and biological processes, which were

not used during the iGRN construction (see Methods). These datasets cover i) a set of 2,759 Y1H

interactions reflecting direct *in vitro* interactions ii) 28 TF ChIP-seq experiments collected from

different studies (148,949 interactions, see also Methods) reflecting primarily direct *in vivo*

binding and iii) a set of 40 differentially expressed (DE) gene sets upon genetic TF perturbation

(51,178 interactions, see Methods). The significance of the observed overlaps was examined

through comparison to a background distribution of expected overlaps generated by

randomizing the iGRN (see Methods). First, we detected a weak but significant overlap between

edges in the integrative network and regulatory edges determined by Y1H (Figure 2a). An

overlap of 336 edges (12%) was observed reflecting a 1.6 higher overlap than expected by

chance (Figure 2b). TF ChIP-seq interactions displayed an overlap of 15,487 interactions (10%)

with a fold enrichment of 2.4. Only considering TFs that occur in both the iGRN and this ChIP-

Seq dataset, for 25/28 regulators there was a significant overlap between the iGRN and ChIP

defined target genes (Supplemental Table S4). Overall, we found that TF ChIP-seq data

confirmed 39% of predicted interactions in the iGRN, demonstrating that a large fraction of iGRN

interactions are supported by *in vivo* TF binding. This fraction is comparable with the

concordance between TF ChIP-based regulatory interactions and altered target gene transcript

levels after genetic TF perturbation, an approach that is frequently used to identify bound target

genes that are also regulated [25]. Next to TF-DNA binding assays, we also investigated the

recovery of directly and indirectly regulated target genes based on differential target gene

expression. The iGRN recovered 9,672 differentially expressed interactions (19%) resulting in a

3.6 fold enrichment (Figure 2a and 2b). Taken together, both the comparison with binding and

regulatory datasets demonstrate that the iGRN represents high-quality interactions supported

by experimental data.

To evaluate the relevance of the regulatory interactions uniquely predicted by the iGRN and

hence not overlapping with the three experimental datasets (Y1H, TF ChIP-Seq, and DE), a

functional metric was used to measure whether a TF's target genes participate in similar

biological processes. This metric starts from the assumption that a set of bona fide target genes

will have a higher enrichment for functional annotations than randomized networks [21]. For each

TF, the enriched Gene Ontology (GO) Biological Process terms based on its target genes were

determined, summarized in a distribution and compared to the background distribution based

on randomized networks. In Figure 2c, the iGRN-predicted interactions were first divided in

different sets based on their overlap with the TF ChIP, Y1H and DE experimental datasets (as in

Figure 2a), respectively. Next, for each subset the fold enrichments for functional GO Biological

Process terms were assessed. Whereas the grey series in Figure 2c show the regulatory

interactions shared between the iGRN and the experimental dataset, the green series shows the

interactions unique to the iGRN. Likewise, the blue, yellow and red series show the regulatory

interactions present in the ChIP, Y1H and DE dataset not overlapping with the iGRN.

Focusing on the overlap with TF ChIP dataset revealed that the functional enrichment was

similar for the interactions unique to the iGRN compared to those shared between the iGRN and

TF ChIP dataset (2.63 and 2.55, respectively). However, the regulatory interactions unique to the

TF ChIP datasets showed lower fold enrichments (1.94), indicating that these show less

functional coherence. For the Y1H comparison, the interactions uniquely predicted by the iGRN

had a similar fold enrichment as the interactions only reported by the Y1H assays (2.41 and

2.61). The interactions that are reported by both approaches had a considerable higher

enrichment compared to the other networks (5.55; though note the small set size, n=336). The

comparison with the DE dataset showed a similar result with an increase in enrichment fold

from uniquely predicted edges in the iGRN to the intersection and uniquely predicted edges by

the DE gene sets (2.33, 2.96 and 3.23, respectively). Taken together, these results indicate that

the predicted interactions, including those unique to the iGRN, convey functional gene

regulatory events.

Network architecture reveals different levels of gene regulatory complexity affecting

plant phenotypes

Starting from the complete iGRN comprising 1.7 million interactions, we investigated the

structure and complexity of the network and determined if network architecture can be linked

to specific biological functions[26]. On average, each gene in the network has 54 incoming TFs

(median 32), while each TF is linked to 1,147 target genes (median 802; Supplemental Figure S2).

There was a large range in connectivity with half of the genes being regulated by 8-79 TFs

(interquartile range), and a small number of TFs (107/1491, or 7%) regulating more than 3,000

target genes. Focusing on the top 1% genes with the largest number of incoming TFs revealed a

significant bias towards genes with transcription factor activity, corroborating previous findings

reporting that regulators themselves are complexly regulated by other regulators [7,27]. In

contrast, 2172 genes having only one incoming TF showed enrichment for RNA binding and

translation, suggesting that these functions are less complexly regulated by TFs.

Next, we examined if network structure can offer new insights in the role of individual

regulators. Concretely, we investigated whether the degree (number of target genes) and

betweenness centrality (a measure for the importance based on number of shortest paths in the

network graph) of TFs in the iGRN can be predictive for potential master regulators showing

prominent phenotypes upon genetic perturbation. Whereas degree refers to the number of

interactions a node, or gene, has in the network, betweenness centrality measures how

important a node is to the shortest paths through the network. Nodes with a high centrality are

assumed to be more important in the network as more information is passing through these

nodes. After processing large-scale phenotyping data from RARGE II, an integrated phenotype

database of Arabidopsis mutants scoring traits using a controlled vocabulary for 14,540 genes,

phenotypes were recorded for 2,224 genes [28]. When splitting all TFs present in the iGRN and

represented in the RARGE II dataset in four quartiles (Q1-Q4) based on their degree values, we

observed that 17.4% of the high-degree (Q3+Q4) and 15.1% of the low-degree (Q1+Q2) TFs had

a reported phenotype (p-value 0.43; two-tailed Mann-Whitney U test). When comparing the

quartiles of TFs based on betweenness centrality, a significant difference was found when

scoring regulators with a reported phenotype (19.6% for Q3+Q4 and 12.4% for Q1+Q2,

respectively; p-value 0.0088; two-tailed Mann-Whitney U test; Figure 3a). This pattern is

confirmed when comparing the betweenness centrality for TFs with and without phenotypes, as

regulators showing RARGE II phenotypes have more of the shortest network paths going

through their nodes (log2 median value of 17.50 and 16.46 for TFs with and without reported

phenotypes, respectively; p-value 1e-05; two-tailed Mann-Whitney U test; Figure 3b). iGRN TFs

within the top 10% genes with highest centrality for which mutants have reported phenotypes

(19/91) are significantly biased towards post-embryonic development and include ATGTL1, PIF4,

SEP3, IDD4 and IDD8, ANAC056, DOF5.6, LFY3 and TCP4. After assigning regulators to three

levels based on their network hierarchy [29], we observed that top-level TFs high in the hierarchy

(n=313) have higher degree and centrality values compared to bottom-level TFs (n=704, median

degree for top and bottom TFs is 1923 and 529, respectively, while log2 median betweenness

centrality for top and bottom TFs is 18.87 and 16.10, respectively). Top-level TFs were not

significantly enriched for specific TF families. Interestingly, both for TFs with and without

phenotypes, a similar number of TFs in their target genes was found (9.45% and 9.44%,

respectively), indicating that TFs showing a phenotype do not regulate more TFs and thus are

not higher in the TF hierarchy.

As transcriptional regulation is mediated through interactions between TFs, co-activators or

co-repressors, and the general transcriptional machinery, we determined whether the iGRN has

predictive power to identify cooperative TF regulation. Therefore, we measured if regulators

sharing a large number of target genes are more likely to interact with each other, based on

experimentally determined protein-protein interactions (PPIs). We found that TF pairs sharing

many target genes (Jaccard index >0.3) have a 15-fold higher occurrence of PPIs compared to TF

pairs having less shared target genes (Jaccard index <0.3), which is highly significant (2.71%

versus 0.31%; p-value <2.2e-16, hypergeometric distribution). Whereas most of these PPIs cover

TFs from the same gene family, probably due to the high similarity of their corresponding

binding sites yielding similar sets of target genes, their potential to act as heterodimers and/or

potential functional redundancy, also cases of interacting TF pairs from different gene families

were found. Examples of the latter case include HY5-PIF3/PIF4 involved in far-red signaling,

MYB21-RGL1/RGL2 controlling hormone-mediated cell proliferation and differentiation, and

BES1-BIM1/BIM2 acting synergistically to control brassinosteroid signaling (Supplemental Figure

S3). The highly correlated expression profiles of these TFs sharing many target genes further

suggest they exert their regulatory roles in a cooperative manner.

Network-based identification of gene functions for Arabidopsis TFs

Based on the validation results indicating the functional relevance of regulatory edges in the

iGRN, we also investigated if individual TF functions can be inferred. Therefore, we performed

GO enrichment analysis for all target genes of each TF, only considering experimental Biological

Process annotations and excluding very general GO terms (see Methods). Next, we determined

if the enriched Biological Process terms were already known for that specific TF or whether a

novel functionality was predicted (Data file 2. TF_function_all). In total we could infer functions

for 1,301 TFs and for 54% (681) of the 1,259 TFs with experimental Biological Process

annotations we could correctly infer one or more of the known functions. For 268 TFs lacking

experimental annotations, novel iGRN-based functions were predicted. Reciprocally, per

Biological Process term, on average 21% of the known regulators were recovered, while for

several distinct processes the recovery was 50% or higher. Examples of processes with high

recovery rates include response to wounding (14/20, 70%), plant-type secondary cell wall

biogenesis (18/28, 64%), response to hormone (174/343, 51%), chloroplast organization (6/12,

50%) and DNA replication (3/6, 50%).

The overall recovery of known TF functions based on the iGRN was lower compared to the

probabilistic functional gene network AraNet v2 [30], which recovered known functions for 739 TFs

(28% average recovery of known regulators per GO BP term). However, both approaches are

highly complementary, as 207 and 265 TFs had known functions only recovered using iGRN and

AraNet, respectively (Supplemental Figure S4). Examples of known TF functions uniquely

recovered through the iGRN cover EIN3 (ethylene-activated signaling pathway), APETALA2

(flower development), LFY (flower development, gibberellic acid-mediated signaling pathway),

FUS3 (somatic embryogenesis), TINY (multidimensional cell growth), and AMS, ATWRKY34 and

CRF12 (pollen development). Focusing on specific GO terms, biological processes with a higher

recovery of known regulators in the iGRN compared to AraNet cover response to biotic stimulus

(47% versus 35%), chloroplast organization (50% versus 0%), response to abiotic stimulus (47%

versus 26%), pollination (33% versus 0%), and root hair cell differentiation (35% versus 6%). For

the latter process we correctly identified the regulators ZPF5, RSL2 and RSL4, RHD6, LRL2 and

LRL3, as well as MYB36. MYB36 regulates the developmental transition in the root endodermis,

where the differentiated endodermis forms a protective waxy barrier called the Casparian strip.

In our network, MYB36 is predicted to regulate genes involved in cell wall organization such as

CASP1-5, PER64 and ESB1, which are known to be involved in Casparian strip formation and

have been shown to be differentially expressed in the *myb36* mutant [31]. Furthermore, the iGRN

also predicts MYB36 to be involved in root hair elongation, which agrees with the longer root

hairs observed in *myb36* mutants [31].

Identification of novel TFs in reactive oxygen species (ROS) stress regulation

Apart from recovering known functions of TFs involved in development and stress responses,

the predictive power of the iGRN was used to identify novel TF functions. As a test case, we

extracted TFs from the iGRN with predicted oxidative stress signaling and/or responsiveness

functions. As an increase in ROS levels is a common theme during abiotic and biotic stresses,

oxidative stress responsiveness was used to validate novel regulators of plant stress responses.

One hundred twenty-four ROS-related TFs were extracted from the iGRN by identifying those

TFs that entailed an enrichment for core ROS responsive transcripts (ROS marker genes) in their

target genes [32]. Among these TFs, 17 had an already reported function in plant responses to ROS

or ROS-causing agents (hydrogen peroxide [H_2O_2], methyl viologen [MV] and 3-amino triazole [3-

AT]), here referred to as 'KNOWN_ROS' TFs (Table 2, Supplemental Table S5). Fifty-eight ROS TFs

were previously reported to function in plant responses to environmental stresses, such as

salinity, drought, heavy metals, cold, hypoxia, high light and infection by bacterial and fungal

pathogens (and that we refer to as 'KNOWN_STRESS' TFs, Supplemental Table S5). The

remaining 49 identified ROS TFs had previously not been linked to any role in

ROS/environmental stress (referred to as 'NOVEL' TFs in Table 2), at the time the iGRN was

constructed (see Methods). However, five of the 49 novel TFs were validated for a role in

environmental stress responses by studies that were published after the iGRN construction

(Table 2 and Supplemental Table S5): ANAC102 and BME3 (high light [33-35]), NAC103 (genotoxic

[36]), ZAT18 (drought [37]) and ERF019 (drought, bacterial and fungal pathogens [38,39]).

For experimental validation, we selected the top 32 novel TFs, ranked based on enrichment

for ROS marker target genes[32], for which at least one *Arabidopsis* loss- or gain-of-function

transgenic line was available (Supplemental Table S6). Plant performance under oxidative stress

was assessed based on rosette growth of the mutant lines compared to the wild type when

grown at low and high MV and 3-AT concentrations, respectively, using automated image

analysis (see Methods), and, in addition, chlorosis symptoms were visually scored.

For 13 of the 32 phenotyped novel ROS TFs we found at least one mutant allele with

significant oxidative stress-induced changes in rosette area relative to the wild type (i.e.

genotype effect dependent on the oxidative stress treatment and/or genotype-by-treatment

effect under at least one of the tested stress treatments; Figure 4; Table 2). For the other novel

TFs, no significant changes were detected by the genotype under all tested conditions, either

there was an overall genotype effect under all tested conditions (similar changes were

observed) or the genotype effect was independent of the stress treatments (e.g. WRKY47;

Figure 4). For SCL13, WRKY45, BEH2 and PHL1, experimental evidence is provided by at least

two mutant alleles. *WRKY45* loss-of-function (T-DNA mutant RATM11−0634−1_H and RNAi line

AS3.1) affected the rosette growth during low and high 3-AT stress compared to the wild type in

a genotype-by-treatment dependent matter, whereas, *WRKY45* OE (line 15.1) caused an

increased rosette yield that was dependent on stress treatments low 3-AT and low MV (Figure

5). Two *SCL13* overexpression alleles (1.11 and 2.12) displayed decreased rosette growth

dependent on both MV conditions in a genotype-by-treatment-dependent manner (Figure 5).

Similarly, knockdown of *SCL13* (RNAi lines AS112D and AS28C) negatively affected rosette

growth under MV stress, but no genotype-by-treatment interaction effect was observed. *PHL1*

T-DNA and *PHL1/PHR1* double T-DNA lines displayed altered growth on low and high 3-AT,

respectively, the latter dependent on the genotype-by-treatment. *BEH2* mutation altered the

rosette area that was dependent on the stress treatments high 3-AT and low MV for line

SALK_079430 and dependent on the genotype-by-treatment for line SAIL_76B06.

We further demonstrated that ANAC102, BME3, MYB51, WRKY55, AT2G33710, ERF022,

AT5G57150 and ERF115 have a function in oxidative stress responses based on one mutant

allele (for *ERF022* and *ERF115* only one line was available) (Figure 4 and Table 2). For example,

ANAC102 OE line 3.4 (strong OE; ~ 100-fold) displayed a larger rosette area that was

independent of the stress treatment, while rosette area changes by *ANAC102* OE2.8 (weak OE;

~4.5-fold) were dependent on the genotype-treatment interaction (Figure 5). *ERF115* OE

affected plant performance in a manner that was dependent on the stress treatment, with a

decreased rosette area under low 3-AT (dependent on the genotype-by-stress treatment) and

severe bleaching under high MV and 3-AT concentrations (Figure 5).

Apart from validating the identified novel ROS TFs based on an oxidative stress-related growth

phenotype, we evaluated the correct prediction of ROS responsive target genes. Therefore, we

compared the predicted target genes of the highest ranked novel ROS TF ANAC102 with

genome-wide target genes identified by ChIP-Seq[27]. Among the 543 ANAC102 target genes

predicted by the iGRN, 185 (34%) were confirmed by ChIP-Seq, indicating a significant overlap

(2.28 fold enrichment, p-value<2e-26). Meta-analysis of transcript profiles during various

oxidative stress conditions shows that 103 of the 185 confirmed target genes are responsive to a

wide range of ROS-inducing conditions[32] (Supplemental Table S8, Supplemental Figure S5) and

include 20 out of the 35 ROS marker genes predicted to be ANAC102 targets. Among the 543

predicted ANAC102 target genes are also 62 TFs of which 28 are confirmed by ChIP-Seq. Among

these ChIP-confirmed TFs are the ROS marker genes *WRKY40, ZAT6, ZAT10, ZAT12* and *ERF5*,

indicating extensive TF-TF regulation of ROS responses downstream of ANAC102. Furthermore,

to be a true regulatory interaction, the spatial-temporal expression characteristics of the TF and

its target genes must be similar. To assess this, we explored transcriptome profiles of different

rosette and root tissues from Genevestivator[40] to compare tissue-specific expression profiles of

ANAC102 with its ChIP-Seq-confirmed target genes. We observed common expression

signatures, namely *ANAC102* and its target genes being expressed in juvenile, adult and

senescent leaves as well as in different primary root zones and in the lateral root (Supplemental

Figure S5). Taken together, using ANAC102 as a test case, we confirmed the prediction of true

target genes, and consistency in oxidative stress responsiveness and spatial-temporal (basal)

expression profiles between *ANAC102* and its targets.

Finally, the expression profiles of KNOWN_ROS and validated novel ROS TFs during different

oxidative stress conditions were analyzed and compared [32]. Fourteen out of the 17 known ROS

TFs were identified as responsive to a wide range of oxidative stress conditions ($|\log_2$ fold

change$| > 1$ and FDR < 0.01; Figure 6; Supplemental Table S7). Among the 13 newly validated

novel ROS TFs, only half were ROS responsive at the transcript level. *WRKY55, MYB122* (for

which the triple mutant *myb34/51/122* shows a phenotype, but not the *myb122* single mutant),

ERF022, *BEH2*, *PHL1*, AT5G57150 and *ERF115* expression levels are not regulated by oxidative

stress, or only regulated in a small number or specific set of oxidative stress-related conditions,

and thus, would not have been predicted from studies solely relying on differential expression at

the whole plant or organ level. For example, *ERF115* has tissue-specific expression patterns,

confined to the stem cell niche within the root meristem [41], indicating that transcript analysis on

the whole plant level might not reveal stress-induced alterations in this specific tissue [42].

Interestingly, we observed that novel ROS TFs showing a phenotype under oxidative stress have

a 2-fold higher degree and 5-fold higher betweenness centrality compared to novel ROS TFs

lacking such phenotype (median degree of 1670 and 749, median betweenness centrality of

699,318 and 142,777, for novel ROS TFs with and without oxidative stress phenotype,

respectively).

Overall, among the 124 novel ROS TFs extracted from the iGRN, 75 (60%) were validated by

previously published studies (before the iGRN was constructed) to be involved in the plant's

responses to ROS perturbation and/or (a)biotic stresses. By analyzing mutant lines perturbed in

one of the analyzed 32 novel ROS TFs, 13 additional regulators were validated for a function in

ROS responses in our phenotyping assays. Among the novel ROS TFs newly identified by our

phenotyping assays, WRKY55, WRKY47, AT2G33710 [ERF] and AT5G57150 [bHLH] had previously

not been associated with any function. Other validated novel ROS TFs are known to function in

high-light stress tolerance or acclimation responses (NAC102, BME3 [GATA family] [33-35]), age-

triggered senescence (WRKY45, ERF022 [AP2/EREBP]; [43,44]), response to phosphate starvation

(WRKY45, PHL1 [G2 family]; [45,46]), submergence stress (WRKY45; [47]), glucosinolate (antimicrobial

compounds involved in plant immunity) biosynthesis (MYB51, MYB122; [48]), brassinosteroid

signaling (BEH2 [BES1 TF family]; [49]), phytochrome-dependent light signaling (SCL13 [GRAS

family]; [50]), hypocotyl growth (BEH2 [BES1 family]; [51]), seed germination (BME3; [52]) and somatic

embryogenesis (ERF022; [53]). Overall, among the novel ROS TFs validated for an oxidative stress

function by literature evidence or in our phenotyping assays, the AP2-EREBP (5), NAC (5) and

WRKY (5) families are highly represented (Supplemental Table S5). These families are typically

associated with abiotic, biotic and/or oxidative stress responses [54] and the iGRN identified

additional members with an oxidative stress phenotype. In addition, the iGRN correctly

predicted an oxidative stress response function for one or more members of the bHLH, ZAT (or

C2H2), HSF, MYB and bZIP families that are also often associated with functions in plant stress

responses. The iGRN also identified novel ROS TFs from families that were known to be mainly

involved in plant growth/developmental processes, such as GRAS (SCL13), BES1 (BEH2) and

GATA (BME3) ([49,55,56]) . So far there is very little evidence for the involvement of GATA TFs in

plant stress responses except for reported functions for rice and soybean GATA TFs in low

nitrogen stress tolerance [57,58], the identification of salt and drought stress-induced alternative

splice products for rice GATA TFs [59], and the recent characterization of BME3 in systemic

acquired acclimation to high light stress, downstream of a systemic signaling cascade mediated

by ROS signaling [34,35]. Consistent with recently published studies [34,35], we show that BME3 is one

of the first GATA transcription factors with a proven function in the regulation of stress

responses.

DISCUSSION

The comprehensive experimental mapping of plant GRNs is a major challenge and different

technologies, either *in vitro* or *in vivo*, each provide partial snapshots of the regulatory logic in

Arabidopsis. While different integrative approaches have been applied to identify regulatory

networks controlling specific biological processes in plants [12,60-63], in this study we leveraged a

wide variety of regulatory datasets capturing different properties of TF-mediated transcriptional

control. Through the application of an advanced machine learning approach for large-scale

functional data integration, we performed for the first time a systematic regulatory annotation

for nearly all *Arabidopsis* genes. Although the presented iGRN is generic and not tailored

towards specific plant organs or growth stages, extensive validation using different types of

experimental datasets, measuring TF binding and/or regulation, revealed a strong enrichment

for functional interactions. In addition, the iGRN outperformed all input networks when

assessing the predictive power using a gold standard evaluation set, demonstrating the success

of the supervised learning approach. An assessment of the functional coherence of a TF's target

genes suggests that the iGRN regulatory interactions are of equally high quality compared to

those obtained through experimental methods. Taken together, our validation results indicate

that the iGRN is strongly enriched for functional interactions and that the applied integrative

approach clearly complements GRNs inferred through a single experimental assay.

Investigating the relationship between network architecture and the complexity of TF control

confirmed that TFs themselves are complexly regulated by other TFs. In addition, through the

integration of large-scale phenotyping data, a significant association was found between a TF's

network centrality (but not degree) and observing a phenotype when knocking out that TF. A

similar pattern was observed for the novel ROS TFs showing an oxidative stress phenotype. This

finding indicates that these TFs occupy key positions in the GRN and loss of their gene products

has major consequences for plant growth and development. Furthermore, TFs showing a

phenotype when screening mutant lines were not having more TFs among their target genes

compared to TFs lacking phenotypes, suggesting that master regulators high in the TF network

hierarchy are not more frequently showing phenotypes compared to lower-tier TFs. Obviously,

these findings depend on the global quality of the iGRN but complement previous results

reporting that higher-tier TFs in ABA networks do not show more differential expression during

ABA response [27]. Finally, TF pairs sharing many target genes are indicative of combinatorial

control, where TFs physically interacting with each other form higher-order protein complexes

mediating increased DNA-binding specificity [64]. Consequently, TF pairs with many shared target

genes in our integrative network which also show correlated expression profiles are good

candidates for future research to shed light on the role of cooperative TF control in specific

biological processes in plants.

Predicted TF functions based on the iGRN regulatory interactions indicated that for various

biological processes many known regulators could be recovered. Starting from our systematic

regulatory annotation, known functional annotations were confirmed for 681 TFs. For several

biological processes our predicted TF functions are highly complementary to the AraNet

probabilistic functional gene network and for 268 TFs lacking experimental annotations, new

functions were inferred. To validate the power of our integrative network, novel regulators

predicted to play a role in oxidative stress signaling were experimentally validated using ROS-

specific phenotyping of loss- or gain-of-function lines. By analyzing transgenic lines perturbed in

one of the 32 novel ROS TFs, an additional 13 (41%) regulators were validated for a function in

ROS responses by our phenotyping studies. In addition to TF families that have often been

associated with oxidative or other environmental stress responses, the iGRN identified novel

ROS TFs from the GRAS, BES1 and GATA families that are not, or to a lesser extent, known to be

involved in stress responses. Notable, compared to previous studies that predicted gene

functions or prioritized candidate regulators based on (co-)expression analysis [65], our method

enables the identification of novel regulators that would have been missed when solely relying

on (differential) expression information. To conclude, our integrative network offers a high-

quality resource to study TF control for individual target genes and to identify, through the

interrogation of specific or condition-specific gene sets, novel regulators controlling different

plant traits.

METHODS

Input datasets

We assembled a comprehensive compendium of genome-wide datasets that were used as

primary data sources to construct the seven input networks and to build the integrative gene

regulatory network. Starting from the TAIR10 Arabidopsis gene annotation, we added a set of

5711 non-coding RNAs described in [66] resulting in a dataset covering 38,966 genes. A first

dataset consisted of 1,260 PWM binding site profiles for 874 transcription factors. This dataset

contained 108 and 623 positional frequency matrices that were obtained from protein binding

microarray studies performed by [67] and [68], respectively, and 529 positional frequency matrices

obtained from DAP-seq experiments [69]. A second dataset contained all available DNaseI

hypersensitive data obtained from the plant regulome data resource

(http://www.plantregulome.org/, [14]). All separate conditions were merged into a single open

chromatin dataset. A third dataset, composed of evolutionary conserved binding sites for 874

TFs, was obtained using the comparative motif mapping algorithm as described [15], starting from

the input motif dataset as described above.

A fourth dataset comprised gene expression profiles generated by RNA-seq. The RNA-Seq

expression compendium was built with public datasets from NCBI's Sequence Read Archive

(SRA) using Curse [70]. The compendium contains gene-level expression values for 40 manually

selected samples (Supplemental Table S9) of different treatment and tissue combinations (in

total >2,7 billion reads). SRA files for each sequencing run were downloaded from the SRA and

converted to the FASTQ format using fastq-dump (v2.4.4) from the SRA toolkit. FASTQ files from

runs of the same sample were concatenated. Paired-end reads were unpaired by randomly

selecting either the forward or reverse read and processing it as single-end. FastQC (v0.9.1) was

used to detect overrepresented adapter sequences, which were subsequently clipped with

fastx_clipper from the FASTX toolkit (v0.0.13). Nucleotides with Phred quality scores lower than

20 were trimmed with fastq_quality_trimmer from the FASTX toolkit. Reads shorter than 20

nucleotides after quality trimming were discarded. To obtain raw read counts for each transcript

in the TAIR10 annotation, Sailfish (v0.6.3) [71] was run with a k-mer length of 20. For genes with

multiple transcripts, the raw read counts of its transcripts were summed to get a gene-level read

count. Counts were then normalized for the entire compendium with the Variance Stabilizing

Transformation (VST) from the DESeq R package (v1.14.0) [72]. VST was chosen since it results in

correlation coefficients between genes that are most comparable to those obtained with

microarray data [73]. A fifth dataset consisted of a compendium of 27 genome-wide TF binding

datasets obtained through ChIP-seq [7].

Network inference for the different input datasets

We derived input networks from each dataset and a weight was assigned to each TF-target

gene interaction reflecting the certainty of the regulatory interaction. The motif target gene

network (PWM) was created by mapping all binding site data on an extended locus for each

gene in the *Arabidopsis thaliana* genome spanning 2kb upstream, intron and 1kb downstream

regions, with the exons masked. All binding site specificity matrices were mapped genome-wide

using Cluster-Buster with C parameter set to zero and M parameter set to 5 [74]. Both the C and M

scores were used as weights, which refer to the cluster and motif score of Cluster-Buster

(Supplemental Table S3). Due to the short length of motifs, many instances of motifs in a single

genome are expected to be coincidental matches that may not be bound *in vivo* or have no

functional regulatory effect. Therefore, the resulting set of binding sites was filtered using

DNaseI hypersensitive sites [14] in order to obtain an open chromatin filtered motif network

(PWM in DHS). The signalValue reported by the authors was retained as weight for motif

matches in open chromatin, this is a measurement of overall enrichment for the region to

DNaseI hypersensitivity. A second filtering approach was based on the evolutionarily

conservation of these binding sites resulting in a conserved motif network (Conserved PWM).

Weights in the conserved motif network are defined by the species conservation score according

to the Comparative Motif Mapping (CMM) method as described in [23]. Co-expression clusters

were built around all TFs obtained from PlantTFDB [75] using the Pearson correlation coefficient

and only the top 1000 most co-expressed genes were retained as putative target genes (CoE TF-

target). GENIE3 was run using standard parameters with the same list of regulators, only

predicted interactions with a variance in expression explained larger than 0.005 were retained

(GENIE3) [16]. A k-nearest neighbor (kNN) co-expression cluster of size 100 was built around each

gene in the *Arabidopsis thaliana* genome using the RNA-seq compendium as defined above.

PWM motif enrichment was performed on this co-expression cluster using the hypergeometric

distribution and significantly enriched motifs (p-value < 0.001 after Benjamini–Hochberg

correction for multiple hypotheses testing) were assigned to the members of the co-expression

cluster in order to obtain the kNN motif enrichment network (CoE + PWM). The enrichment fold

of the motif in the co-expression cluster is used as a weight. The peak scores for each TF bound

region assigned to the closest gene were used as weight in the ChIP-seq network (TF ChIP) [7].

Integrative network inference

The integrative network inference method is based on a binary classifier, where the class label

represents the presence or absence of an interaction. The set of true interactions (positives)

used in the training set was obtained from all validated interactions in AtRegNet [18], a review

article on known interactions in secondary cell wall development [19] and large scale data mining

effort on 974 peer reviewed articles called ATRM [20], and has 5,732 edges, connecting 522 TFs

with 3,738 targets. The remaining interactions between these TFs and potential target genes are

considered to be negatives. In other words, while no true "negatives" are present in the dataset,

we assign negative labels to TF–target pairs where the TF appears in the TFs of one of the

positive pairs and this TF-target pair is not in the positive pairs. The negative labels reflect the

fact that the TF has been studied and since no interaction is reported for it, it is more likely to

represent a negative interaction. This dataset was split into a dataset for learning (80%) and a

dataset for independent evaluation (20%). As such, the training set consists of 1,951,236 TF–

target gene pairs, out of which 4,586 are positives (true interactions) and 1,946,650 negatives.

The remaining gene pairs for TFs absent from the learning/evaluation dataset are considered to

be negatives. Using the weights derived from the input networks we tested a number of

different classification methods (gene pairs missing in specific input networks received a weight

of zero in the matrix used for learning). We extracted the 4,586 validated interacting pairs from

the true dataset and obtained the negative pairs for training by selecting a random subset of

13,758 (three times the number of positives; 3x 4,586) negatively labeled interacting pairs. Using

this labeled training set different classifiers were built using the weights of the input networks as

features. All features were normalized to have zero mean and unit variance. Classifiers were

evaluated using the 20% evaluation dataset. In this analysis the gradient boosting using decision

trees, random forests, and logistic regression were compared. The gradient boosting model

made use of 1,000 trees, shrinkage of 0.01 and an interaction depth of 3. The random forest

model used 500 trees selecting a random set of features of size square root (features) was

sampled in each iteration. Logistic regression was used with a binomial distribution suitable for

binary classification. All classifiers were compared using the area under the curve (Supplemental

Table S2).

The final network was generated using the gradient boosting classifier, as this was the best

performing throughout the training and validation steps. A total set of 60,714,507 unlabeled

interactions was classified. We included all interactions that are predicted to interact with a

probability of 0.35 or higher in the integrative network as this cutoff maximized the F1 measure

on the test dataset (Supplemental Table S2). The full iGRN can be mined using overlap and TF

enrichment analysis at http://bioinformatics.psb.ugent.be/webtools/iGRN/. Code, datasets and

documentation describing the supervised learning approach are available at

https://github.com/VIB-PSB/iGRN.

The overlap between the iGRN and the evaluation dataset was evaluated by counting how

many interactions from these experimental interactions were present in the integrative

network. The enrichment between two networks was defined as the number of interactions that

are present in both networks divided by the number of interactions expected by chance. The

number of common interactions expected by chance is given by the mean overlap of 10,000

randomized networks [21]. Network randomization was done by permuting the labels of all TFs

and the labels of all genes, which preserves the network structure.

624   Functional evaluation using experimental datasets

Three experimentally obtained gene regulatory networks were used for further validation of

the predicted network. An accumulation of publicly available Y1H data from literature was used

to generate a Y1H network containing 2,759 interactions [76-79]. A set of differentially expressed

genes upon TF perturbation was obtained from the CORNET database, containing 51,178

interactions [80]. An independent set of interactions determined by ChIP-seq was obtained from

literature, reporting 148,949 interactions [27,81-84]. From Song et al. 2016 only non-ABA conditions

were considered.

The overlap between the iGRN and three experimentally determined gene regulatory

networks was evaluated by counting how many interactions from the experimental networks

were present in the integrative network. The enrichment between two networks was defined as

the number of interactions that are present in both networks divided by the number of

interactions expected by chance, as explained above. Network randomization was done by

permuting the labels of all TFs and the labels of all genes, which preserves the network structure

and assures that the observed enrichment is not due to potential biases arising from structural

properties of the network (Supplemental Table S3).

Functional enrichment was determined for each sub network (intersection, FN, FP) using

biological metrics as described in [23] using Gene Ontology information. Note that no functional

GO information was used as input for the iGRN construction.

Analysis of network architecture and overlap with plant phenotypes

Based on the target genes for all TFs present in the iGRN, degree and betweenness centrality

were computed using the igraph R package, considering directed edge information. Three levels

of network hierarchy were calculated using the R code available from

http://archive.gersteinlab.org/proj/hinet/ [29]. The Jaccard index, defined as the intersection over

union, was used to compute the overlap of target genes between two TFs. Based on bulk

downloads of the RARGE II database ([28]; download November 2018), which reports ontology-

based phenotype information for 48,077 lines (17,671 RIKEN Arabidopsis Ds transposon mutant

lines, 16,337 CSHL Arabidopsis genetrap mutant lines, 14,069 RIKEN Arabidopsis full-length

cDNA overexpressed Arabidopsis lines). The presence of phenotypes in the RARGE II dataset was

scored for different sets of regulators, defined based on degree and betweenness centrality.

TF function prediction

For each TF, iGRN target genes were subjected to GO enrichment analysis using the

hypergeometric distribution combined with Benjamini–Hochberg (B&H) correction for multiple

hypotheses testing. A q-value cutoff 1e-03 was applied since this resulted in the best F1 score, a

metric that combines precision and recall to recover known GO annotations. Only experimental

and curated GO BP annotations were considered (version August 2018), GO annotations were

propagated to include parental GO terms, and a set of 18 general GO terms, corresponding with

terms located at the root of the GO hierarchy, were excluded (GO:0008150, GO:0009987,

GO:0008152, GO:0044237, GO:0071704, GO:0050896, GO:0065007, GO:0032502, GO:0050789,

GO:0032501, GO:0007275, GO:0050794, GO:0006355, GO:0045893, GO:0045892).

To predict ROS TFs, core genes that are part of the ROS wheel clusters IV-VIII, covering genes

related to chemical and genetic perturbation of cellular ROS/redox, were used to define a

custom category of ROS marker genes [32]. Subsequently, the over-representation of these ROS

markers in the set of iGRN target genes per TF, was used to predict novel ROS TFs and rank them

based on the q-value for enrichment (hypergeometric distribution, q-value cutoff 1e-05). The

assignment of ROS TFs to the categories KNOWN_ROS, KNOWN_STRESS and NOVEL_TF was

based on the information available in the literature at the moment the initial iGRN was

constructed (December 2016).

Phenotyping candidate ROS TFs

T-DNA insertion mutants were obtained from the European Arabidopsis Stock Centre (NASC;

http://arabidopsis.info/), GABI-Kat (http://www.gabi-kat.de/db/), Versailles Arabidopsis Stock

Center (http://publiclines.versailles.inra.fr/tdna/index) (Supplemental Table S6). Homozygous T-

DNA insertion lines were selected by PCR analysis (Supplemental Table S10). Mutant lines

previously published were provided by the authors (Supplemental Table S6 (overview lines)).

Overexpression lines were generated by Gateway cloning in the binary destination vector

pK7WG2D (18056864) (Supplemental Table S10). Plants were grown on half-strength Murashige

and Skoog (1/2MS) medium (Duchefa) supplemented with 1% (w/v) sucrose, 0.75% (w/v) agar,

and B5 vitamins, pH 5.7, at 21°C and 60 µE m^{-2} s^{-1} light intensity in a 16-h/8-h light/dark

photoperiod. To induce oxidative stress, the 1/2MS medium was supplemented with 15 or 20

nM MV; or 1 or 1.5 µM 3-AT, causing on average 15% (15 nM MV), 30% (20 nM MV), 20% (1 µM

3-AT) and 55% (1.5 µM 3-AT) growth reduction compared to the nontreated seedlings. Rosette

area was determined by automated area measurements of individual plant rosettes using

ImageJ. Raw phenotyping results are available in Data file 3. Presence or absence of

chlorosis/bleaching was monitored by visual assessment. For each mutant line, the rosette area

of mutant seedlings was compared with wild-type plants grown in the same Petri plate. Data

were obtained from at least two biological replicates per experiment and from independent

experiments.

Statistical analysis leaf phenotyping data ROS TFs

A mixed model was fitted to the rosette area with genotype and treatment as fixed effects as

well as the interaction term with the mixed procedure from SAS (SAS Institute Inc. 2018.

SAS/STAT® 15.1, Cary, NC). The factor plate was added as random effect in the model to take

into account the correlation between observations made on the same plate. The Kenward-

Rogers method was used to calculate the denominator degrees of freedom of the fixed effects.

The null hypotheses of interest were an equal genotype effect upon treatment in the mutant

line and the wild-type line and an equal treatment versus MS effect between the mutant line

and the wild-type line (i.e. no interaction effect). A significance level of 0.05 was taken. To

correct for multiple testing, the maxT procedure was used as implemented in the plm

procedure.

REFERENCES

1   Franco-Zorrilla, J. M. & Solano, R. Identification of plant transcription factor target
sequences. *Biochim Biophys Acta* **1860**, 21-30, doi:10.1016/j.bbagrm.2016.05.001
(2017).
2   Lehti-Shiu, M. D., Panchy, N., Wang, P., Uygun, S. & Shiu, S. H. Diversity, expansion, and
evolutionary novelty of plant DNA-binding transcription factor families. *Biochim Biophys*
Acta Gene Regul Mech **1860**, 3-20, doi:10.1016/j.bbagrm.2016.08.005 (2017).
3   Jones, D. M. & Vandepoele, K. Identification and evolution of gene regulatory networks:
insights from comparative studies in plants. *Curr Opin Plant Biol* **54**, 42-48,
doi:10.1016/j.pbi.2019.12.008 (2020).
4   Rhee, S. Y. & Mutwil, M. Towards revealing the functions of all genes in plants. *Trends*
Plant Sci **19**, 212-221, doi:10.1016/j.tplants.2013.10.006 (2014).
5   Gaudinier, A. & Brady, S. M. Mapping Transcriptional Networks in Plants: Data-Driven
Discovery of Novel Biological Mechanisms. *Annu Rev Plant Biol* **67**, 575-594,
doi:10.1146/annurev-arplant-043015-112205 (2016).
6   Ferrier, T., Matus, J. T., Jin, J. & Riechmann, J. L. Arabidopsis paves the way: genomic and
network analyses in crops. *Curr Opin Biotechnol* **22**, 260-270,
doi:10.1016/j.copbio.2010.11.010 (2011).
7   Heyndrickx, K. S., Van de Velde, J., Wang, C., Weigel, D. & Vandepoele, K. A functional
and evolutionary perspective on transcription factor binding in Arabidopsis thaliana.
Plant Cell **26**, 3894-3910, doi:10.1105/tpc.114.130591 (2014).
8   Lu, Z., Hofmeister, B. T., Vollmers, C., DuBois, R. M. & Schmitz, R. J. Combining ATAC-seq
with nuclei sorting for discovery of cis-regulatory regions in plant genomes. *Nucleic*
Acids Res **45**, e41, doi:10.1093/nar/gkw1179 (2017).
9   Maher, K. A. *et al.* Profiling of Accessible Chromatin Regions across Multiple Plant
Species and Cell Types Reveals Common Gene Regulatory Principles and New Control
Modules. *Plant Cell* **30**, 15-36, doi:10.1105/tpc.17.00581 (2018).
10   Sijacic, P., Bajic, M., McKinney, E. C., Meagher, R. B. & Deal, R. B. Changes in chromatin
accessibility between Arabidopsis stem cells and mesophyll cells illuminate cell type-
specific transcription factor networks. *Plant J* **94**, 215-231, doi:10.1111/tpj.13882
(2018).
11   Kulkarni, S. R., Jones, D. M. & Vandepoele, K. Enhanced Maps of Transcription Factor
Binding Sites Improve Regulatory Networks Learned from Accessible Chromatin Data.
Plant Physiol **181**, 412-425, doi:10.1104/pp.19.00605 (2019).
12   Pajoro, A. *et al.* Dynamics of chromatin accessibility and gene regulation by MADS-
domain transcription factors in flower development. *Genome biology* **15**, R41 (2014).

13   Lee, I., Ambaru, B., Thakkar, P., Marcotte, E. M. & Rhee, S. Y. Rational association of
740   genes with traits using a genome-scale gene network for Arabidopsis thaliana. *Nat*
741   *Biotechnol* **28**, 149-156, doi:10.1038/nbt.1603 (2010).

742   14   Sullivan, A. M. *et al.* Mapping and dynamics of regulatory DNA and transcription factor
743   networks in A. thaliana. *Cell reports* **8**, 2015-2030 (2014).

744   15   Van de Velde, J., Van Bel, M., Vaneechoutte, D. & Vandepoele, K. A Collection of
745   Conserved Noncoding Sequences to Study Gene Regulation in Flowering Plants. *Plant*
746   *Physiol* **171**, 2586-2598, doi:10.1104/pp.16.00821 (2016).

747   16   Huynh-Thu, V. A., Irrthum, A., Wehenkel, L. & Geurts, P. Inferring regulatory networks
748   from expression data using tree-based methods. *PLoS One* **5**,
749   doi:10.1371/journal.pone.0012776 (2010).

750   17   Marbach, D. *et al.* Wisdom of crowds for robust gene network inference. *Nat Methods*
751   **9**, 796-804, doi:10.1038/nmeth.2016 (2012).

752   18   Yilmaz, A. *et al.* AGRIS: the Arabidopsis Gene Regulatory Information Server, an update.
753   *Nucleic Acids Res* **39**, D1118-1122, doi:10.1093/nar/gkq1120 (2011).

754   19   Hussey, S. G., Mizrachi, E., Creux, N. M. & Myburg, A. A. Navigating the transcriptional
755   roadmap regulating plant secondary cell wall deposition. *Front Plant Sci* **4**, 325,
756   doi:10.3389/fpls.2013.00325 (2013).

757   20   Jin, J. *et al.* An Arabidopsis Transcriptional Regulatory Map Reveals Distinct Functional
758   and Evolutionary Features of Novel Transcription Factors. *Mol Biol Evol* **32**, 1767-1773,
759   doi:10.1093/molbev/msv058 (2015).

760   21   Marbach, D. *et al.* Predictive regulatory models in Drosophila melanogaster by
761   integrative inference of transcriptional networks. *Genome Res* **22**, 1334-1349,
762   doi:10.1101/gr.127191.111 (2012).

763   22   Kheradpour, P., Stark, A., Roy, S. & Kellis, M. Reliable prediction of regulator targets
764   using 12 Drosophila genomes. *Genome Res* **17**, 1919-1931, doi:10.1101/gr.7090407
765   (2007).

766   23   Van de Velde, J., Heyndrickx, K. S. & Vandepoele, K. Inference of transcriptional
767   networks in Arabidopsis through conserved noncoding sequence analysis. *The Plant Cell*
768   **26**, 2729-2745 (2014).

769   24   Tompa, M. *et al.* Assessing computational tools for the discovery of transcription factor
770   binding sites. *Nat Biotechnol* **23**, 137-144, doi:10.1038/nbt1053 (2005).

771   25   O'Maoileidigh, D. S., Graciet, E. & Wellmer, F. Gene networks controlling Arabidopsis
772   thaliana flower development. *New Phytol* **201**, 16-30, doi:10.1111/nph.12444 (2014).

773   26   Luscombe, N. M. *et al.* Genomic analysis of regulatory network dynamics reveals large
774   topological changes. *Nature* **431**, 308-312, doi:10.1038/nature02782 (2004).

775   27   Song, L. *et al.* A transcription factor hierarchy defines an environmental stress response
776   network. *Science* **354**, aag1550 (2016).

777   28   Akiyama, K. *et al.* RARGE II: an integrated phenotype database of Arabidopsis mutant
778   traits using a controlled vocabulary. *Plant Cell Physiol* **55**, e4, doi:10.1093/pcp/pct165
779   (2014).

780   29   Cheng, C. *et al.* An approach for determining and measuring network hierarchy applied
781   to comparing the phosphorylome and the regulome. *Genome Biol* **16**, 63,
782   doi:10.1186/s13059-015-0624-2 (2015).

783   30   Lee, T. *et al.* AraNet v2: an improved database of co-functional gene networks for the
784   study of Arabidopsis thaliana and 27 other nonmodel plant species. *Nucleic Acids Res* **43**,
785   D996-1002, doi:10.1093/nar/gku1053 (2015).

31   Kamiya, T. *et al.* The MYB36 transcription factor orchestrates Casparian strip formation.
Proc Natl Acad Sci U S A **112**, 10533-10538, doi:10.1073/pnas.1507691112 (2015).
32   Willems, P. *et al.* The ROS Wheel: Refining ROS Transcriptional Footprints. *Plant Physiol*
171, 1720-1733, doi:10.1104/pp.16.00420 (2016).
33   D'Alessandro, S., Ksas, B. & Havaux, M. Decoding beta-Cyclocitral-Mediated Retrograde
Signaling Reveals the Role of a Detoxification Response in Plant Tolerance to
Photooxidative Stress. *Plant Cell* **30**, 2495-2511, doi:10.1105/tpc.18.00578 (2018).
34   Fichman, Y. *et al.* MYB30 Orchestrates Systemic Reactive Oxygen Signaling and Plant
Acclimation. *Plant Physiol* **184**, 666-675, doi:10.1104/pp.20.00859 (2020).
35   Zandalinas, S. I., Sengupta, S., Burks, D., Azad, R. K. & Mittler, R. Identification and
characterization of a core set of ROS wave-associated transcripts involved in the
systemic acquired acclimation response of Arabidopsis to excess light. *Plant J* **98**, 126-
141, doi:10.1111/tpj.14205 (2019).
36   Ryu, T. H. *et al.* SOG1-dependent NAC103 modulates the DNA damage response as a
transcriptional regulator in Arabidopsis. *Plant J* **98**, 83-96, doi:10.1111/tpj.14201 (2019).
37   Yin, M. *et al.* The Arabidopsis Cys2/His2 zinc finger transcription factor ZAT18 is a
positive regulator of plant tolerance to drought stress. *J Exp Bot* **68**, 2991-3005,
doi:10.1093/jxb/erx157 (2017).
38   Huang, P. Y. *et al.* NINJA-associated ERF19 negatively regulates Arabidopsis pattern-
triggered immunity. *J Exp Bot* **70**, 1033-1047, doi:10.1093/jxb/ery414 (2019).
39   Scarpeci, T. E., Frea, V. S., Zanor, M. I. & Valle, E. M. Overexpression of AtERF019 delays
plant growth and senescence, and improves drought tolerance in Arabidopsis. *J Exp Bot*
68, 673-685, doi:10.1093/jxb/erw429 (2017).
40   Hruz, T. *et al.* Genevestigator v3: a reference expression database for the meta-analysis
of transcriptomes. *Adv Bioinformatics* **2008**, 420747, doi:10.1155/2008/420747 (2008).
41   Heyman, J. *et al.* ERF115 controls root quiescent center cell division and stem cell
replenishment. *Science* **342**, 860-863, doi:10.1126/science.1240667 (2013).
42   Kong, X. *et al.* PHB3 Maintains Root Stem Cell Niche Identity through ROS-Responsive
AP2/ERF Transcription Factors in Arabidopsis. *Cell Rep* **22**, 1350-1363,
doi:10.1016/j.celrep.2017.12.105 (2018).
43   Chen, L., Xiang, S., Chen, Y., Li, D. & Yu, D. Arabidopsis WRKY45 Interacts with the DELLA
Protein RGL1 to Positively Regulate Age-Triggered Leaf Senescence. *Mol Plant* **10**, 1174-
1189, doi:10.1016/j.molp.2017.07.008 (2017).
44   Hinckley, W. E., Keymanesh, K., Cordova, J. A. & Brusslan, J. A. The HAC1 histone
acetyltransferase promotes leaf senescence and regulates the expression of ERF022.
Plant Direct **3**, e00159, doi:10.1002/pld3.159 (2019).
45   Wang, H. *et al.* Arabidopsis WRKY45 transcription factor activates PHOSPHATE
TRANSPORTER1;1 expression in response to phosphate starvation. *Plant Physiol* **164**,
2020-2029, doi:10.1104/pp.113.235077 (2014).
46   Bustos, R. *et al.* A central regulatory system largely controls transcriptional activation
and repression responses to phosphate starvation in Arabidopsis. *PLoS Genet* **6**,
e1001102, doi:10.1371/journal.pgen.1001102 (2010).
47   Meng, X. *et al.* Mitochondrial signalling is critical for acclimation and adaptation to
flooding in Arabidopsis thaliana. *Plant J* **103**, 227-247, doi:10.1111/tpj.14724 (2020).
48   Frerigmann, H. & Gigolashvili, T. MYB34, MYB51, and MYB122 distinctly regulate indolic
glucosinolate biosynthesis in Arabidopsis thaliana. *Mol Plant* **7**, 814-828,
doi:10.1093/mp/ssu004 (2014).

49　Yin, Y. *et al.* A new class of transcription factors mediates brassinosteroid-regulated
gene expression in Arabidopsis. *Cell* **120**, 249-259, doi:10.1016/j.cell.2004.11.044
(2005).
50　Torres-Galea, P., Huang, L. F., Chua, N. H. & Bolle, C. The GRAS protein SCL13 is a
positive regulator of phytochrome-dependent red light signaling, but can also modulate
phytochrome A responses. *Mol Genet Genomics* **276**, 13-30, doi:10.1007/s00438-006-
0123-y (2006).
51　Lachowiec, J., Mason, G. A., Schultz, K. & Queitsch, C. Redundancy, Feedback, and
Robustness in the Arabidopsis thaliana BZR/BEH Gene Family. *Front Genet* **9**, 523,
doi:10.3389/fgene.2018.00523 (2018).
52　Liu, P. P., Koizuka, N., Martin, R. C. & Nonogaki, H. The BME3 (Blue Micropylar End 3)
GATA zinc finger transcription factor is a positive regulator of Arabidopsis seed
germination. *Plant J* **44**, 960-971, doi:10.1111/j.1365-313X.2005.02588.x (2005).
53　Nowak, K., Wojcikowska, B. & Gaj, M. D. ERF022 impacts the induction of somatic
embryogenesis in Arabidopsis through the ethylene-related pathway. *Planta* **241**, 967-
985, doi:10.1007/s00425-014-2225-9 (2015).
54　Hoang, X. L. T., Nhi, D. N. H., Thu, N. B. A., Thao, N. P. & Tran, L. P. Transcription Factors
and Their Roles in Signal Transduction in Plants under Abiotic Stresses. *Curr Genomics*
18, 483-497, doi:10.2174/1389202918666170227150057 (2017).
55　Hirsch, S. & Oldroyd, G. E. GRAS-domain transcription factors that regulate plant
development. *Plant Signal Behav* **4**, 698-700, doi:10.4161/psb.4.8.9176 (2009).
56　Behringer, C. & Schwechheimer, C. B-GATA transcription factors - insights into their
structure, regulation, and role in plant development. *Front Plant Sci* **6**, 90,
doi:10.3389/fpls.2015.00090 (2015).
57　Hudson, D. *et al.* Rice cytokinin GATA transcription Factor1 regulates chloroplast
development and plant architecture. *Plant Physiol* **162**, 132-144,
doi:10.1104/pp.113.217265 (2013).
58　Zhang, C. *et al.* Genome-wide survey of the soybean GATA transcription factor gene
family and expression analysis under low nitrogen stress. *PLoS One* **10**, e0125174,
doi:10.1371/journal.pone.0125174 (2015).
59　Gupta, P., Nutan, K. K., Singla-Pareek, S. L. & Pareek, A. Abiotic Stresses Cause
Differential Regulation of Alternative Splice Forms of GATA Transcription Factor in Rice.
Front Plant Sci **8**, 1944, doi:10.3389/fpls.2017.01944 (2017).
60　Brooks, M. D. *et al.* Network Walking charts transcriptional dynamics of nitrogen
signaling by integrating validated and predicted genome-wide interactions. *Nat*
Commun **10**, 1569, doi:10.1038/s41467-019-09522-1 (2019).
61　Gaudinier, A. *et al.* Transcriptional regulation of nitrogen-associated metabolism and
growth. *Nature* **563**, 259-264, doi:10.1038/s41586-018-0656-3 (2018).
62　Zander, M. *et al.* Integrated multi-omics framework of the plant response to jasmonic
acid. *Nat Plants* **6**, 290-302, doi:10.1038/s41477-020-0605-7 (2020).
63　Reynoso, M. A. *et al.* Evolutionary flexibility in flooding response circuitry in
angiosperms. *Science* **365**, 1291-1295, doi:10.1126/science.aax8862 (2019).
64　Smaczniak, C., Muino, J. M., Chen, D., Angenent, G. C. & Kaufmann, K. Differences in
DNA Binding Specificity of Floral Homeotic Protein Complexes Predict Organ-Specific
Target Genes. *Plant Cell* **29**, 1822-1835, doi:10.1105/tpc.17.00145 (2017).
65　Ransbotyn, V. *et al.* A combination of gene expression ranking and co-expression
network analysis increases discovery rate in large-scale mutant screens for novel

880   Arabidopsis thaliana abiotic stress genes. *Plant Biotechnol J* **13**, 501-513,
881   doi:10.1111/pbi.12274 (2015).

882   66   Liu, J. *et al.* Genome-wide analysis uncovers regulation of long intergenic noncoding
883   RNAs in Arabidopsis. *Plant Cell* **24**, 4333-4345, doi:10.1105/tpc.112.102855 (2012).

884   67   Franco-Zorrilla, J. M. *et al.* DNA-binding specificities of plant transcription factors and
885   their potential to define target genes. *Proceedings of the National Academy of Sciences*
886   **111**, 2367-2372 (2014).

887   68   Weirauch, M. T. *et al.* Determination and inference of eukaryotic transcription factor
888   sequence specificity. *Cell* **158**, 1431-1443, doi:10.1016/j.cell.2014.08.009 (2014).

889   69   O'Malley, R. C. *et al.* Cistrome and Epicistrome Features Shape the Regulatory DNA
890   Landscape. *Cell* **165**, 1280-1292 (2016).

891   70   Vaneechoutte, D. & Vandepoele, K. Curse: building expression atlases and co-expression
892   networks from public RNA-Seq data. *Bioinformatics* **35**, 2880-2881,
893   doi:10.1093/bioinformatics/bty1052 (2019).

894   71   Patro, R., Mount, S. M. & Kingsford, C. Sailfish enables alignment-free isoform
895   quantification from RNA-seq reads using lightweight algorithms. *Nat Biotechnol* **32**, 462-
896   464, doi:10.1038/nbt.2862 (2014).

897   72   Anders, S. & Huber, W. Differential expression analysis for sequence count data.
898   *Genome Biol* **11**, R106, doi:10.1186/gb-2010-11-10-r106 (2010).

899   73   Giorgi, F. M., Del Fabbro, C. & Licausi, F. Comparative study of RNA-seq- and microarray-
900   derived coexpression networks in Arabidopsis thaliana. *Bioinformatics* **29**, 717-724,
901   doi:10.1093/bioinformatics/btt053 (2013).

902   74   Frith, M. C., Li, M. C. & Weng, Z. Cluster-Buster: Finding dense clusters of motifs in DNA
903   sequences. *Nucleic Acids Res* **31**, 3666-3668 (2003).

904   75   Jin, J., Zhang, H., Kong, L., Gao, G. & Luo, J. PlantTFDB 3.0: a portal for the functional and
905   evolutionary study of plant transcription factors. *Nucleic Acids Res* **42**, D1182-1187,
906   doi:10.1093/nar/gkt1016 (2014).

907   76   Brady, S. M. *et al.* A stele-enriched gene regulatory network in the Arabidopsis root. *Mol*
908   *Syst Biol* **7**, 459, doi:10.1038/msb.2010.114 (2011).

909   77   Li, B. *et al.* Promoter-based integration in plant defense regulation. *Plant Physiol* **166**,
910   1803-1820, doi:10.1104/pp.114.248716 (2014).

911   78   Taylor-Teeples, M. *et al.* An Arabidopsis gene regulatory network for secondary cell wall
912   synthesis. *Nature* **517**, 571-575 (2015).

913   79   Tian, C. *et al.* An organ boundary-enriched gene regulatory network uncovers regulatory
914   hierarchies underlying axillary meristem initiation. *Mol Syst Biol* **10**, 755,
915   doi:10.15252/msb.20145470 (2014).

916   80   De Bodt, S., Hollunder, J., Nelissen, H., Meulemeester, N. & Inze, D. CORNET 2.0:
917   integrating plant coexpression, protein-protein interactions, regulatory interactions,
918   gene associations and functional annotations. *New Phytol* **195**, 707-720,
919   doi:10.1111/j.1469-8137.2012.04184.x (2012).

920   81   Nagel, D. H. *et al.* Genome-wide identification of CCA1 targets uncovers an expanded
921   clock network in Arabidopsis. *Proc Natl Acad Sci U S A* **112**, E4802-4810,
922   doi:10.1073/pnas.1513609112 (2015).

923   82   Verkest, A. *et al.* A generic tool for transcription factor target gene discovery in
924   Arabidopsis cell suspension cultures based on tandem chromatin affinity purification.
925   *Plant Physiol* **164**, 1122-1133, doi:10.1104/pp.113.229617 (2014).

83   Birkenbihl, R. P., Diezel, C. & Somssich, I. E. Arabidopsis WRKY33 is a key transcriptional
regulator of hormonal and metabolic responses toward Botrytis cinerea infection. *Plant*
Physiol **159**, 266-285, doi:10.1104/pp.111.192641 (2012).

84   Merelo, P. *et al.* Genome-wide identification of KANADI1 target genes. *PLoS One* **8**,
e77341, doi:10.1371/journal.pone.0077341 (2013).

85   Qureshi, M. K., Sujeeth, N., Gechev, T. S. & Hille, J. The zinc finger protein ZAT11
modulates paraquat-induced programmed cell death in Arabidopsis thaliana. *Acta*
Physiologiae Plantarum **35**, 1863–1871 (2013).

86   Liu, X. M. *et al.* ZAT11, a zinc finger transcription factor, is a negative regulator of nickel
ion tolerance in Arabidopsis. *Plant Cell Rep* **33**, 2015-2021, doi:10.1007/s00299-014-
1675-7 (2014).

87   Rizhsky, L., Davletova, S., Liang, H. & Mittler, R. The zinc finger protein Zat12 is required
for cytosolic ascorbate peroxidase 1 expression during oxidative stress in Arabidopsis. *J*
Biol Chem **279**, 11736-11743, doi:10.1074/jbc.M313350200 (2004).

88   Davletova, S., Schlauch, K., Coutu, J. & Mittler, R. The zinc-finger protein Zat12 plays a
central role in reactive oxygen and abiotic stress signaling in Arabidopsis. *Plant Physiol*
139, 847-856, doi:10.1104/pp.105.068254 (2005).

89   Scarpeci, T. E., Zanor, M. I., Mueller-Roeber, B. & Valle, E. M. Overexpression of
AtWRKY30 enhances abiotic stress tolerance during early growth stages in Arabidopsis
thaliana. *Plant Mol Biol* **83**, 265-277, doi:10.1007/s11103-013-0090-8 (2013).

90   Perez-Salamo, I. *et al.* The heat shock factor A4A confers salt tolerance and is regulated
by oxidative stress and the mitogen-activated protein kinases MPK3 and MPK6. *Plant*
Physiol **165**, 319-334, doi:10.1104/pp.114.237891 (2014).

91   Vanderauwera, S. *et al.* AtWRKY15 perturbation abolishes the mitochondrial stress
response that steers osmotic stress tolerance in Arabidopsis. *Proc Natl Acad Sci U S A*
109, 20113-20118, doi:10.1073/pnas.1217516109 (2012).

92   Mittler, R. *et al.* Gain- and loss-of-function mutations in Zat10 enhance the tolerance of
plants to abiotic stress. *FEBS Lett* **580**, 6537-6542, doi:10.1016/j.febslet.2006.11.002
(2006).

93   Sewelam, N. *et al.* Ethylene response factor 6 is a regulator of reactive oxygen species
signaling in Arabidopsis. *PLoS One* **8**, e70289, doi:10.1371/journal.pone.0070289 (2013).

94   Moffat, C. S. *et al.* ERF5 and ERF6 play redundant roles as positive regulators of JA/Et-
mediated defense against Botrytis cinerea in Arabidopsis. *PLoS One* **7**, e35995,
doi:10.1371/journal.pone.0035995 (2012).

95   Dubois, M. *et al.* Ethylene Response Factor6 acts as a central regulator of leaf growth
under water-limiting conditions in Arabidopsis. *Plant Physiol* **162**, 319-332,
doi:10.1104/pp.113.216341 (2013).

96   Mahmood, K., Xu, Z., El-Kereamy, A., Casaretto, J. A. & Rothstein, S. J. The Arabidopsis
Transcription Factor ANAC032 Represses Anthocyanin Biosynthesis in Response to High
Sucrose and Oxidative and Abiotic Stresses. *Front Plant Sci* **7**, 1548,
doi:10.3389/fpls.2016.01548 (2016).

97   Allu, A. D., Brotman, Y., Xue, G. P. & Balazadeh, S. Transcription factor ANAC032
modulates JA/SA signalling in response to Pseudomonas syringae infection. *EMBO Rep*
17, 1578-1589, doi:10.15252/embr.201642197 (2016).

98   Babitha, K. C. *et al.* Co-expression of AtbHLH17 and AtWRKY28 confers resistance to
abiotic stress in Arabidopsis. *Transgenic Res* **22**, 327-341, doi:10.1007/s11248-012-9645-
8 (2013).

99   Chen, X. *et al.* Overexpression of AtWRKY28 and AtWRKY75 in Arabidopsis enhances
resistance to oxalic acid and Sclerotinia sclerotiorum. *Plant Cell Rep* **32**, 1589-1599,
doi:10.1007/s00299-013-1469-3 (2013).
100 Wu, L., Zhong, G., Wang, J., Li, X. & Yang, Y. Functional analysis of an arabidopsis
transcription factor WRKY28 in resistance to alternaria brassicicola. *Journal of*
Agricultural Science and Technology **14**, 65-71 (2012).
101 De Clercq, I. *et al.* The membrane-bound NAC transcription factor ANAC013 functions in
mitochondrial retrograde regulation of the oxidative stress response in Arabidopsis.
Plant Cell **25**, 3472-3490, doi:10.1105/tpc.113.117168 (2013).
102 Kim, J. B., Kang, J. Y. & Kim, S. Y. Over-expression of a transcription factor regulating
ABA-responsive gene expression confers multiple stress tolerance. *Plant Biotechnol J* **2**,
459-466, doi:10.1111/j.1467-7652.2004.00090.x (2004).
103 Van Aken, O. *et al.* Mitochondrial and Chloroplast Stress Responses Are Modulated in
Distinct Touch and Chemical Inhibition Phases. *Plant Physiol* **171**, 2150-2165,
doi:10.1104/pp.16.00273 (2016).
104 Ng, S. *et al.* A membrane-bound NAC transcription factor, ANAC017, mediates
mitochondrial retrograde signaling in Arabidopsis. *Plant Cell* **25**, 3450-3471,
doi:10.1105/tpc.113.113985 (2013).
105 Dombrecht, B. *et al.* MYC2 differentially modulates diverse jasmonate-dependent
functions in Arabidopsis. *Plant Cell* **19**, 2225-2245, doi:10.1105/tpc.106.048017 (2007).
106 Verma, D., Jalmi, S. K., Bhagat, P. K., Verma, N. & Sinha, A. K. A bHLH transcription
factor, MYC2, imparts salt intolerance by regulating proline biosynthesis in Arabidopsis.
FEBS J **287**, 2560-2576, doi:10.1111/febs.15157 (2020).
107 Song, S. *et al.* Interaction between MYC2 and ETHYLENE INSENSITIVE3 modulates
antagonism between jasmonate and ethylene signaling in Arabidopsis. *Plant Cell* **26**,
263-279, doi:10.1105/tpc.113.120394 (2014).
108 Wu, A. *et al.* JUNGBRUNNEN1, a reactive oxygen species-responsive NAC transcription
factor, regulates longevity in Arabidopsis. *Plant Cell* **24**, 482-506,
doi:10.1105/tpc.111.090894 (2012).
109 Shahnejat-Bushehri, S., Nobmann, B., Devi Allu, A. & Balazadeh, S. JUB1 suppresses
Pseudomonas syringae-induced defense responses through accumulation of DELLA
proteins. *Plant Signal Behav* **11**, e1181245, doi:10.1080/15592324.2016.1181245
(2016).
110 Ebrahimian-Motlagh, S. *et al.* JUNGBRUNNEN1 Confers Drought Tolerance Downstream
of the HD-Zip I Transcription Factor AtHB13. *Front Plant Sci* **8**, 2118,
doi:10.3389/fpls.2017.02118 (2017).
111 Nishizawa, A. *et al.* Arabidopsis heat shock transcription factor A2 as a key regulator in
response to several types of environmental stress. *Plant J* **48**, 535-547,
doi:10.1111/j.1365-313X.2006.02889.x (2006).
112 Nishizawa-Yokoi, A., Yabuta, Y. & Shigeoka, S. The contribution of carbohydrates
including raffinose family oligosaccharides and sugar alcohols to protection of plant cells
from oxidative damage. *Plant Signal Behav* **3**, 1016-1018, doi:10.4161/psb.6738 (2008).
113 Zhang, L., Li, Y., Xing, D. & Gao, C. Characterization of mitochondrial dynamics and
subcellular localization of ROS reveal that HsfA2 alleviates oxidative damage caused by
heat stress in Arabidopsis. *J Exp Bot* **60**, 2073-2091, doi:10.1093/jxb/erp078 (2009).
114 Olate, E., Jimenez-Gomez, J. M., Holuigue, L. & Salinas, J. NPR1 mediates a novel
regulatory pathway in cold acclimation by interacting with HSFA1 factors. *Nat Plants* **4**,
811-823, doi:10.1038/s41477-018-0254-2 (2018).

115 Enomoto, T. *et al.* STOP1 regulates the expression of HsfA2 and GDHs that are critical for
low-oxygen tolerance in Arabidopsis. *J Exp Bot* **70**, 3297-3311, doi:10.1093/jxb/erz124
(2019).
116 Hsieh, T. H. *et al.* Heterology expression of the Arabidopsis C-repeat/dehydration
response element binding factor 1 gene confers elevated tolerance to chilling and
oxidative stresses in transgenic tomato. *Plant Physiol* **129**, 1086-1094,
doi:10.1104/pp.003442 (2002).
117 Jaglo-Ottosen, K. R., Gilmour, S. J., Zarka, D. G., Schabenberger, O. & Thomashow, M. F.
Arabidopsis CBF1 overexpression induces COR genes and enhances freezing tolerance.
Science **280**, 104-106, doi:10.1126/science.280.5360.104 (1998).
118 Hsieh, T. H., Lee, J. T., Charng, Y. Y. & Chan, M. T. Tomato plants ectopically expressing
Arabidopsis CBF1 show enhanced resistance to water deficit stress. *Plant Physiol* **130**,
618-626, doi:10.1104/pp.006783 (2002).
1034

ACKNOWLEDGEMENTS

The authors wish to thank Cordelia Bolle (SCL13), Xuemei Chen (ARF3), Yi-Fang Chen

(WRKY45), Donna Fernandez (AGL15), Henning Frerigmann (MYB51, MYB122), Małgorzata Gaj

(ERF022), Tamara gigolashvili (MYB51), Dwayne Hegedus (SCL15), Lars Hennig (AGL19), Ildoo

Hwang (BME3), Zhongchi Liu (TSO1), Lars Østergaard (ARF3), Javier Paz-Ares (PHL1) and Olivier

Van Aken (WRKY45) for kindly providing Arabidopsis mutant lines. We thank Zoé Joly-Lopez and

Olivia Wilkins for providing in-depth comments on an earlier version of the manuscript, as well

as Camilla Ferrari and Indeewari Dissanayake for proofreading. I.D.C. is supported by the

Research Foundation Flanders (postdoctoral fellowship 12N2415N and grant for a long stay

abroad V400620N). J.V.d.V. is indebted to the Agency for Innovation by Science and Technology

in Flanders for a predoctoral fellowship. X.L. and L.L. are supported by the China Scholarship

Council for a PhD fellowship (201706910099 and 201808530499). F.V.B. is supported by the

Fonds Wetenschappelijk Onderzoek – Vlaanderen (FWO) and the Fonds de la Recherche

Scientifique – FNRS under EOS Project No. 30829584 and the Research Foundation Flanders

(project nos. G0D7914N and G055416N).

AUTHOR CONTRIBUTIONS STATEMENT

J.V.d.V., I.D.C. and K.V. designed the research. I.D.C., X.L., L.L. and R.P. performed the

phenotype screening. V.S. performed statistical data analysis. D.V. generated the expression

compendium. J.V.d.V. and K.V. performed network analysis and function prediction. J.V.d.V.,

I.D.C., F.V.B. and K.V. wrote the manuscript.

COMPETING INTERESTS STATEMENTS

The authors declare no competing interests.

 FIGURE LEGENDS

 Figure 1. Construction and evaluation of the integrated gene regulatory network

 (iGRN).

 a. Overview of the different input networks used to construct the supervised iGRN. Whereas

 colored circles indicate the type of data that is used to construct the input network, the

 information below each network depicts how edges in that network are weighted. b. Overview

 of the sizes of the different networks together with the recall and enrichment for edges in the

 evaluation test dataset (black and white bars, respectively). The brown bars report the fraction

 of each input network that is present in the iGRN while the red bars report which fraction of the

 supervised iGRN edges is supported by this input network.

 Figure 2. Overlap between iGRN interactions and different experimental data sources

 describing gene regulatory and functional information.

 a. Venn diagrams showing the number of shared interactions between the iGRN and TF ChIP,

 Y1H and differentially expressed (DE) genes after TF perturbation (depicted in blue, yellow and

 red, respectively). b. Histograms showing the expected overlap based on 10,000 randomized

 networks, together with the observed overlap indicated by the black arrow. The p-values for

 observing the measured overlap by chance are also reported (permutation based statistics, one-

 sided). c. Fold enrichment for a functional metric describing how well target genes of a specific

 TF are involved in similar biological processes (average values for all enriched GO terms and all

 TFs present in the different data series are reported). Whereas the grey bars indicate the

 functional enrichment for shared interactions, blue, red, yellow and green depict fold

 enrichment for interactions unique to the ChIP, DE, Y1H and iGRN, respectively.

 Figure 3. Phenotypes reported after TF perturbation for TFs with different network

 centrality properties.

 a. RARGE-II phenotypes reported for iGRN TFs with different betweenness centrality. Q1-Q4

 refer to the four quartiles of all iGRN TFs (sorted based on increasing betweenness centrality)

 and values in parenthesis refer to the median betweenness centrality value together with the

 number of lines available in the RARGE II dataset for that quartile. b. Betweenness centrality

1087 distribution for TFs with and without phenotypes reported in RARGE-II (n= 153 and 1589 TFs,

1088 respectively).

1089 Figure 4. Oxidative stress phenotyping results of novel ROS TFs.

Heatmap presenting phenotypes associated with mutant alleles of iGRN extracted novel ROS TFs

under different oxidative stress conditions. Mutant lines with a statistically significant

phenotype under oxidative stress (1 µM and 1.5 µM 3-AT and 15 and 20 nM MV) and/or normal

growth conditions (MS) are displayed. Results of other mutant lines that were tested in this

study can be found in Supplemental Table S5 and S6. Statistically significant differences relative

to wild type (Col-0) are indicated by asterisk; *, $P < 0.05$ and **, $P < 0.001$ as determined by the

hypothesis tests on the fixed effects of the mixed model (two-sided, see Methods). Data were

obtained from at least two biological replicates per experiment and from independent

experiments. Color scale indicates the ratio of the rosette area of 14-day-old mutant alleles

relative to wild type (Col-0). Black and grey borders indicate presence or absence of a bleaching

phenotype of the mutant plants, respectively. Small grey circle indicates a statistically significant

genotype-by-stress treatment interaction effect.

1102 Figure 5. Representative images of oxidative stress phenotypes for a selection of

1103 mutants.

1104 Representative images of novel ROS TFs with a phenotype during oxidative stress. 17-days-old

1105 seedlings of selected mutant lines compared to wild type (Col-0). Scale bars, 1 cm.

1106 Figure 6. Expression of KNOWN and NOVEL ROS TFs during various conditions

1107 provoking cellular ROS/redox imbalances.

Heatmap showing the expression response of ROS TFs extracted from the iGRN during a

selection of conditions provoking cellular ROS production or redox imbalances (a more elaborate

set of oxidative stress conditions is displayed in Supplemental Table S7). TFs with previously

known function in ROS responses and TFs with no functional link with any oxidative stress

and/or (a)biotic stress responses (NOVEL) that were tested in our phenotyping assays are

displayed. NOVEL TFs validated in our study are shown using blue and red circles, indicating a

positive and negative effect by the TF on oxidative stress tolerance, respectively.

TABLES

Table 1. Overview of different networks used to construct the iGRN.

Type	Network name	#interactions	#TFs	Median #interactions (average)
Input network	PWM	17,053,785	874	21,797 (19,512)
	Conserved PWM	340,893	874	247 (390)
	PWM in DHS	5,231,258	874	6,195 (5,985)
	COE + PWM	324,864	703	290 (462)
	TF ChIP	46,619	27	1,026 (1,727)
	COE TF-target	1,547,002	1547	1,000 (1,000)
	GENIE3	2,588,312	1261	1,985 (2,053)
Gold standard	Gold standard	5,732	522	2 (11)
Evaluation	Test dataset	1,146	257	1 (4)
Output network	iGRN	1,709,871	1491	802 (1,147)

1117

Table 2. Overview phenotype information for known and validated novel ROS regulators.

RANK iGRN	Symbol	q-val	Status (1)	Known oxidative stress phenotype from literature	Reference	Known other (a/biotic) stress phenotype from literature	Reference	Validated oxidative stress phenotype from this study
iGRN1	ANAC102	1.84E-35	**NOVEL**			high light	[33]	**MV**
iGRN3	ZAT11	2.81E-25	KNOWN_ROS	MV	[85]	heavy metal	[86]	
iGRN5	ZAT12	8.70E-25	KNOWN_ROS	MV	[87]	osmotic/salt/high light/heat	[88]	
iGRN6	WRKY30	1.42E-24	KNOWN_ROS	MV	[89]	salt	[89]	
iGRN11	HSFA4A	5.09E-23	KNOWN_ROS	MV	[90]	salt, anoxia	[90]	
iGRN12	WRKY15	7.46E-23	KNOWN_ROS	H2O2	[91]	salt/osmotic	[91]	
iGRN16	STZ	4.90E-22	KNOWN_ROS	MV	[92]	osmotic/salt/heat	[92]	
iGRN19	SCL13	3.76E-21	**NOVEL**					MV
iGRN22	MYB51	1.84E-20	**NOVEL**					**3-AT, MV**
iGRN27	BME3	2.01E-18	**NOVEL**					MV
iGRN29	WRKY55	6.69E-18	**NOVEL**					**3-AT, MV**
iGRN36	ERF6	3.13E-15	KNOWN_ROS	H2O2	[93]	high light, B. cinerea, osmotic	[93 94 95]	

iGRN50	WRKY45	5.93E-13	NOVEL(*)			submergence stress	[47]	3-AT, MV
iGRN52	ANAC032	1.26E-12	KNOWN_ROS	3-AT	[96]	P. syringae, salt/high light	[96 97]	
iGRN55	AT2G33710	9.71E-12	NOVEL					3-AT, MV
iGRN58	MYB122	1.42E-10	NOVEL					3-AT, MV
iGRN60	WRKY28	1.70E-10	KNOWN_ROS	MV, H2O2	[98]	salt/osmotic/ drought, S. sclerotiorum, A. brassicicola	[98 99 100]	
iGRN62	ANAC013	1.93E-10	KNOWN_ROS	MV	[101]			
iGRN65	ABF3	8.06E-10	KNOWN_ROS	MV	[102]	cold/freezing/heat /drought	[102]	
iGRN66	ERF022	9.55E-10	NOVEL					MV
iGRN68	BEH2	2.75E-09	NOVEL					3-AT, MV
iGRN69	ANAC017	3.04E-09	KNOWN_ROS	MV	[103]	drought/high light	[104]	
iGRN81	PHL1	2.76E-08	NOVEL					3-AT, MV
iGRN85	JAM1	4.07E-08	KNOWN_ROS	MV, H2O2	[98]	salt/osmotic/ drought	[98]	
iGRN93	MYC2	4.19E-07	KNOWN_ROS	MV	[105]	salt, B. Cinerea	[106 107]	
iGRN100	AT5G57150	1.50E-06	NOVEL					3-AT, MV
iGRN102	ERF115	1.73E-06	NOVEL					3-AT, MV
iGRN110	JUB1	3.20E-06	KNOWN_ROS	H2O2	[108]	salt/heat, P. syringae, drought	[108 109 110]	
iGRN111	HSFA2	3.21E-06	KNOWN_ROS	MV	[111]	high light, heat, cold, hypoxia	[111 112 113 114 115]	
iGRN118	CBF1	5.32E-06	KNOWN_ROS	MV	[116]	salt, drought, freezing	[116 117 118]	

(1) Asterisk indicates NOVEL TFs that were validated for a function in (a)biotic stress (*) responses by literature evidence after 2016.

OVERVIEW OF SUPPLEMENTAL FIGURES AND TABLES

Figure S1. Overview of the iGRN interactions supported by different input networks.

Figure S2. Overview of the in- and out-degree for the complete iGRN.

Figure S3. Examples of predicted cooperative TFs supported by protein-protein interactions.

Figure S4. Recovery of known GO annotations for TFs belonging to different biological processes.

Figure S5. Oxidative stress-responsive and tissue-specific expression profiles of ANAC102 and its ChIP-confirmed target genes.

Supplemental Table S1. Overview of TF families in gold standard.

Supplemental Table S2. Overview of data types, training methods and performance results for different machine learning classification algorithms.

Supplemental Table S3. Overlap and enrichment between different networks.

Supplemental Table S4. Overlap of target genes between iGRN and ChIP-Seq data.

Supplemental Table S5. Overview of all TFs predicted by the iGRN with a function in ROS responses.

Supplemental Table S6. Overview of mutant lines used for phenotypic analysis.

Supplemental Table S7. Expression of KNOWN_ROS and NOVEL ROS TFs during various conditions provoking cellular ROS/redox imbalances.

Supplemental Table S8. Expression of ANAC102 and its ChIP-confirmed target genes during various conditions provoking cellular ROS/redox imbalances.

Supplemental Table S9. Overview of samples included in the RNA-Seq expression compendium.

Supplemental Table S10. Primers used in this study.

DATA AVAILABILITY

Source data for Figs. 1,4 and S3 are provided with the paper.

Data file 1. Data_file1_iGRN_supervised_network_support.txt All TF-target interactions in the iGRN.

Columns report: TF, target genes, score, rank, support input networks.
Overview of generated, analysed and used iGRN regulatory interaction data for Fig. 1.

Data file 2. TF_function_all.txt reporting predicted functions for all TFs present in the iGRN.

The sheet README explains the different columns and offers 3 examples how this file can be explored for specific TFs, functions or a specific target gene.
Overview of generated, analysed and used iGRN TF function data for Fig. S3.

Data file 3. Raw phenotype results that were used as input for the statistical analysis.

Overview of raw phenotyping measurements for Fig. 4.

a

PWM	PWM in DHS	Conserved PWM	TF ChIP

CoE TF-target | GENIE3 | CoE + PWM

Motif score & Cluster score

DNase HSS score

Conservation score

Peak score

Pearson Correlation Coefficient

Variance explained

Enrichment PWM in COE cluster

⬤ TF motif   ⬤ TF ChIP binding

⬤ open chromatin   ⬤ co-expression

⬤ conserved non-coding sequences

Gold standard

Final Network iGRN

Classifier → 20% evaluation data
→ 80% training data

b

Approach	Number of interactions	Recall of true positives	Enrichment evalation set	Network contribution	Percentage of iGRN
TF ChIP	46,619	29 %	6.0	80 %	2 %
PWM	17,053,785	60 %	1.0	7 %	70 %
Conserved PWM	340,893	11 %	6.4	91 %	17 %
PWM in DHS	5,231,258	34 %	1.5	17 %	49 %
CoE + PWM	324,864	3 %	2.2	18 %	3 %
CoE TF- target	1,547,002	10 %	3.0	40 %	36 %
GENIE3	2,588,312	17 %	1.7	32 %	48 %
Supervised Learning iGRN	1,709,871	46 %	8.8		

log2 fold change

-3   -2   -1   0   1   2   3

flu (2h D-L)
H2O2 (1h)
AA (3h)
MV (3h)
PNO8 (3h)
UV 310nm (3h)
DBMIB (2h)
O3 125ppb (3h)
flg22 (1h)
flu (30min D-L)
UV 310nm (30min)
GO5 (5h 380ppm CO2)
RbohF (cat2)
UV 310nm (24h)
O3 125ppb (48h)
cs26 (LD)
ssi2

KNOWN_ROS

iGRN3_ZAT11
iGRN5_ZAT12
iGRN6_WRKY30
iGRN11_HSFA4A
iGRN12_WRKY15
iGRN16_STZ
iGRN36_ERF6
iGRN52_ANAC032
iGRN60_WRKY28
iGRN62_ANAC13
iGRN65_ABF3
iGRN69_ANAC017
iGRN85_JAM1
iGRN93_MYC2
iGRN110_JUB1
iGRN111_HSFA2
iGRN118_CBF1

NOVEL

iGRN1_ANAC102
iGRN19_SCL13
iGRN20_AT5G59450
iGRN22_MYB51
iGRN27_BME3
iGRN29_WRKY55
iGRN33_GT3a
iGRN37_AGL19
iGRN44_HAM4
iGRN47_AT2G38250
iGRN50_WRKY45
iGRN53_AT2G22760
iGRN54_BEH3
iGRN55_AT2G33710
iGRN57_ANAC090
iGRN58_MYB122
iGRN66_ERF022
iGRN68_BEH2
iGRN72_AT1G25550
iGRN76_ZFN2
iGRN79_AT4G32800
iGRN80_ZAT18
iGRN81_PHL1
iGRN83_ANAC036
iGRN87_WRKY47
iGRN90_ARF10
iGRN91_BZIP25
iGRN96_AGL15
iGRN99_TSO1
iGRN100_AT5G57150
iGRN102_ERF115
iGRN104_ARF3