How widespread was the Tapanuli orangutan and what led to its decline?

Erik Meijaard[1,2,3,*], Safwanah Ni'matullah[1], Rona Dennis[1], Julie Sherman[1,4], Onrizal[5] and Serge

A. Wich[6]

[1] Borneo Futures, Bandar Seri Begawan, Brunei Darussalam

[2] Durrell Institute of Conservation and Ecology, University of Kent, Canterbury, UK

[3] School of Biological Sciences, University of Queensland, St Lucia, Australia

[4] Wildlife Impact. PO Box 31062, Portland, Oregon, USA

[5] University of Sumatera Utara, Medan, Indonesia

[6] Liverpool John Moores University, Liverpool, UK

* Corresponding author

E-mail: emeijaard@gmail.com

Abstract

The Tapanuli orangutan (*Pongo tapanuliensis*) is the most threatened great ape species in the

world. It is restricted to an area of about 1,000 km^2 of mostly hill forest where fewer than 800

animals survive in three declining subpopulations. Through a historical ecology approach

involving analysis of colonial-era and other literature, we demonstrate that historically the

Tapanuli orangutan lived in a much larger area, and across a much wider range of habitat types

compared to now. Its current range is about 10-20% of the range it had some 150-200 years

ago. A combination of historic fragmentation of forest habitats, mostly for small-scale

agriculture, and unsustainable hunting likely drove various populations to the south, east and

west of the current population to extinction prior to the industrial-scale forest conversion that

started in the 1970s. Our findings indicate how sensitive orangutans are to the combined effects

of habitat fragmentation and unsustainable mortality or rescue and translocation rates. Saving

this species will require prevention of any further fragmentation and any killings or other removal

of animals from the remaining population. Without concerted action to achieve this, the

remaining populations of Tapanuli orangutan are doomed to become extinct within several

orangutan generations.

Introduction

Determining the key drivers of population decline is a primary objective in conservation biology

and wildlife management. Many wildlife species are threatened by a range of different and often

interacting factors, and developing effective conservation strategies requires unravelling how

these threats interact [1]. This is rarely easy, because species operate in complex socio-

ecological systems in which different components are affected by a range of anthropomorphic

factors such as habitat loss and fragmentation or unsustainable harvest. Evidence-based

conservation seeks to address this by quantifying the relationships between conservation

actions, change in threat severity and change in conservation status [2, 3]. Collecting evidence

is, however, time-consuming, and when conservation problems are "wicked", i.e., the problems

change as solutions are found [4], a stable solution may not be found to a particular

conservation problem [5]. This often means that scientific evidence does not support clear-cut

conclusions in value-driven debates that characterize conservation [6]. Nevertheless,

conservation advocates often seek simple narratives to convince the public of the urgency of

environmental problems and the need to support it. A good example is the polarized debate

around oil palm (*Elaeis guineensis*) and orangutans (*Pongo* sp.) where both proponents and

opponents of oil palm cherry-pick facts to drive home particular messages, i.e., oil palm is either

the greatest evil destroyer of tropical rainforests or a blessing to impoverished rural communities

in tropical countries [7]. A similarly polarized debate currently rages between opponents and

proponents of a hydropower project in the range of the Tapanuli orangutan (*Pongo*

tapanuliensis), an endemic great ape of the Indonesian island of Sumatra [8] (Fig 1).

One way to bring more clarity in muddled or polarized debates is to be more specific about the

system in which a particular problem plays out. For example, if the system boundaries are

limited to oil palm as an ecological threat to orangutan survival [9], a simple solution would be to

ban palm oil use and to stop its production, preventing further deforestation. If the system

boundaries are extended to include smallholder farmers who produce palm oil for their own

needs as well as international markets, a ban on palm oil would encompass broader ethical

connotations as it would affect people's livelihoods [7]. The use of different perspectives in

complex conservation contexts may not make it easier to solve them but can provide helpful

insights about the boundaries of a particular problem. Are they, for example, mostly ecological,

or do they involve human threats, such as hunting, or societal ethics? One such perspective is

history. Looking back in time on the development of a particular problem may provide insights

about the underlying drivers of that problem. The historical ecology approach builds on this

approach by using historical knowledge in the management of ecosystems or species [10].

Referring to historical evidence has, for example, provided valuable understanding about the

ecology of orangutans and what likely caused their decline during the Late Pleistocene and

Holocene, which informs their management today [11]. Here we apply an analysis of historical

ecology to one particular species of orangutan, the Tapanuli Orangutan, by analysing rarely

used colonial-era literature to better understand the historic distribution of the species.

Indonesia's colonial literature on natural history was mostly written in Dutch and German, and is

not commonly used by conservation scientists working in Indonesia.

The Tapanuli Orangutan was described in 2017 as a third species of orangutan [12], 20 years

after this orangutan population was formally reported to modern science [13]. The species is

restricted to three areas of mostly upland forest in the Batang Toru area in North Sumatra (Fig

1), totalling approximately 1,023 km^2 [14, 15]. This orangutan population had been largely

overlooked by science, despite having been tentatively described in the colonial literature [16].

The estimated total number of wild Tapanuli Orangutans is currently 767 [95%: 213-1,597, 14]

making this the great ape species with the lowest number of individuals in the wild and perhaps

the most threatened in the world [8].

Fig 1. Map of the island of Sumatra, showing the distribution ranges of *P. tapanuliensis*

and *P. abelii*, and the main locations (cities, districts and other geographic features)

mentioned in the text. The inset shows the area of Fig 2.

The species is currently under threat from habitat loss for agriculture, hunting and conflict killing,

and development in the area, for infrastructure, gold mining, and geothermal and hydro-energy,

which threaten to further reduce and fragment remaining habitat, reduce dispersal opportunities

for the orangutans between subpopulations, and undermine population viability through

unsustainable mortality rates [14, 17-20]. Due to its restricted current distribution mostly centred

around higher elevations (834.4±219.3m asl) compared to 701.7±454.8 m asl for the Sumatran

orangutan (*P. abelii*) and 170.6±187.0 m asl for the Bornean orangutan (*P. pygmaeus*) [21], it

has been argued that the individuals of the species have adapted specifically to the uplands that

cover most of its current distribution [17]. What is not clear is whether the currently existing

altitudinal differences between these orangutan species are the result of ecological

specializations to highland ecological conditions, or whether the highland species now occur at

higher altitudes because their previous lowland habitats no longer exist or because the species

became extinct there. The fossil record for Sumatran orangutans confirms that the genus *Pongo*

was once more widespread. Extensive remains from the Late Pleistocene and Holocene have

been excavated from a range of caves in the Padang Highlands, some 300 km south of the

current range [22] (Fig 1). Why the species disappeared from that part of Sumatra remains

unclear but unsustainable hunting is one of the possible explanations, because until recently

large areas of suitable forest habitat remained in areas where orangutans are now extinct [23].

Given that, in the past, forest cover was also much more widespread in the range of the

Tapanuli Orangutan, it is important to determine whether historically (ca. past 500 years)

orangutans did occur in those areas. This would help establish whether the Tapanuli orangutan

has indeed evolved to only live in the highlands and estimate what its past distribution could

have been.

The aim of this paper is to compile reports of orangutans occurring to the south of the Toba lake

(Fig 1) with the focus on determining how reliable these are, and, where feasible, provide a

location for the occurrence of orangutans to assess whether these are predominantly highland

sites, and to assess which factors could have led to their disappearance in those areas.

Methods
110

We compiled records of orangutans from historic sources by searching natural history books,

scientific papers, and historic newspapers from before 1940. We searched databases with

location specific keywords such as Sumatra, Batang Toeroe, and Tapanoeli, using Dutch

spelling. We combined this with searches for terms specifically referring to orangutans: Orang

oetan, orang-oetan, orangutan, and also mawas, mias and maias (local names for orangutan

commonly used in historic literature), using a variety of spellings. For the period since 1940, we

used the sources from the review in Rijksen and Meijaard [23] as well as scientific papers and

personal communications. To determine the locations of the historic sightings or captures we

consulted the online Leiden University Library colonial map repository

(http://maps.library.leiden.edu/apps/s7). In some cases, rivers or villages were indicated which

made it feasible to estimate the location of the sightings quite accurately. In other cases, the

area of the sighting or captures was indicated in a broader area which reduced accuracy (tens

of kilometers).

We assessed the likely vegetation types that Tapanuli Orangutans would have occurred in, and

determined the altitude at which they were reported from the global SRTM layer (30m

accuracy). For this, we digitized the first official forest cover map of Indonesia [24], which is

likely based on maps produced by the Netherlands-Indies cartographic service from the 1930s

and 1940s. We carried out the digitization of the 1950 Indonesia forest cover map using

ArcScan extension in ArcGIS. ArcScan is an extension in ArcGIS that converts scanned raster

images into vector-based GIS layers [25]. Prior to digitization, the Indonesia forest cover map

was georeferenced. This is because scanned raster maps mostly do not contain spatial

reference information. Georeferencing helps to assign the coordinate systems of the map and

output of ArcScan. Digitization of raster map via ArcScan resulted in polylines. These polylines

were useful in constructing vegetation polygons of the area of interest. Therefore, it is important

to clear any unrelated polylines generated in the ArcScan output to ensure polygons could be

properly identified and constructed. We completed the construction of polygons using advance

editing tool in ArcGIS. This tool closed and converted polylines into polygons. Polygons

constructed represent different vegetation types in the area. We then classified the polygons in

accordance to their vegetation type with reference to the Indonesia forest cover map. Before

vegetation types assignment, we ensured that different vegetation types were represented with

separate polygons.

The exact location of the historic orangutan sightings cannot be determined with certainty but

the descriptions often provide sufficient detail through names of rivers and villages to estimate

the location on a map. Due to this uncertainty the altitude was extracted as a range for some

locations. Altitude was determined from the altitude layer in Google Earth Pro. We used the

vegetation map for Sumatra [26] in combination with knowledge gained by co-author SW during

surveys in the region to assign one of the forest categories to an estimated historic location.

Results

Historic accounts

We report the various historic accounts of orangutan sightings or specimens from outside the

currently known range in chronological order, starting with Schlegel and Mueller [27] who

reported on two orangutan crania that were obtained by a military doctor in the surroundings of

Jambi, some 650 km south-east of the current closest orangutan population. It is unclear

whether these animals were obtained in Jambi from the wild or whether they were in captivity

and possibly originated from northern Sumatra. Schlegel and Mueller reported that the two

skulls were housed in the "Rijks-Museum", which presumably refers to the Leiden Museum of

Natural History. Furthermore, they wrote that on the west coast of Sumatra, especially north of

the equator, the orangutan was known by the name *mawej*, although in areas further south such

as Indrapura and Bengkulu, the names *orang-panda* or *orang-pandak* were used. The authors

acknowledged that the orangutan was especially common in the north-east of Sumatra, but that

occasionally they were encountered further south and along the western shores of Sumatra.

Also, the species had been reported from Indrapura (in present-day Riau Province) and near

Padang (in West Sumatra, Fig 1), although the descriptions by Schlegel and Mueller are

insufficient to confirm that these reports referred to orangutans. Co-author Onrizal, remembers

growing up east of Padang (Sungai Dareh) listening to stories about *orang-pendek*, human-like

creatures living in the forest, which reportedly ceased to exist in the area in the 1970s. Stories of

such *orang-pendek* abound in other parts of southern Sumatra and there has been speculation

for over a century that these could be remnant orangutan populations [28, 29], although firm

evidence remains lacking. For the purpose of the current paper we do not focus on the *orang-*

pendek narratives.

In a description of their travels across Sumatra, Müller and Horner [30] wrote that orangutans

were not unknown in the Tapanuli area and especially common in "Taroemon", i.e., present-day

Trumon in the Singkil area (Fig 1), which is part of the *Pongo abelii* range. They report that

people distinguished between two types of orangutans, the *maweh baroet* (baroet meaning

monkey in the local language) and *maweh orang* (the 'human' orangutan). Von Rosenberg [31]

did not provide much detail but similarly mentioned that orangutans were present north of

"Tapanoeli" (what is now Sibolga) to Singkil, indicating presence of the species in the coastal

lowlands west of Lake Toba (no. 1 in Fig 2). He saw two orangutans but did not clarify where he

saw them.

Fig 2. Map showing points where orangutans where historically reported and the forest

cover in 1950 before large-scale commercial forest exploitation began.

Kramm [32] reported on a hunting expedition near "Soeroe Mantinggi", where he found several

orangutans and observed them for several hours. The location likely referred to Sayur Matinggi

(no. 2 in Fig 2), which is currently located in the Batang Gadis area, some 50 km south of the

current range of *P. tapanuliensis*. Kramm mentioned that Soeroe Mantinggi is located at a

distance of 22 "*palen*" from Padang Sidempoean. A "paal" was a measurement used in the

Netherlands-Indies, equalling 1852 m on Sumatra, indicating a distance of about 40 km for 22

"*palen*". Sayur Matinggi is currently located some 26 km from Padangsidempuan as the crow

flies which indicates that indeed this is likely to be the location where Kramm observed

orangutans. Kramm was familiar with orangutans which he reported to have also encountered in

"Loemoet" and "Batang-Taro". We believe that the former refers to Lumut (no. 3 in Fig 2), just

south of Sibolga, and that Batang-Taro is an older name for the Batang Toru area, where *P.*

tapanuliensis occurs until today.

Orangutans also seem to have occurred northeast of the current range of the Tapanuli

Orangutan. Neumann [33] described the species from "Hadjoran", which was located in the

watershed of the "Batang Si Ombal" and "Aek Hiloeng", and for which the following coordinates

were given: N 2°1'25" and E 99°29', in the current district of Padang Lawas (Fig 1, no. 4 in Fig

2). This is about 50 km northeast of the most eastern current range of Tapanuli Orangutan. The

detailed description, however, suggests that the species was very rare there, and the people of

"Hadjoran" had not seen the orangutan there before. The animal was shot, with descriptions of

the local people suggesting it was at least 1 m tall, possibly indicating an adult male, which are

known to roam far from breeding populations. Neumann writes that he travelled extensively

through forest areas in the Padang Lawas area searching for orangutans but never managed to

encounter one.

Hagen [34, p. 66] stated that orangutans were known from the west coast between Tapanuli

and Singkil (no. 5 in Fig 2), although Singkil is in the range of *Pongo abelii* and it is not clear

whether the coastal Tapanuli reference referred to the area of the current range of *P.*

tapanuliensis, or whether it referred to what is now the Central Tapanuli District which extends

to the southern part of Singkil, west of Lake Toba. Interestingly, he referred to an orangutan

from the interior of Padang (in reference to an article by S. Jentink, Aardrijkskundig Weekblad,

1881, No. 44, p. 287 – not seen by the authors), in west Sumatra that ended up in the

Rotterdam Zoo where it died of a bone deformation disease (the skull is kept in the Natural

History museum in Leiden, the Netherlands: RMNH.MAM.544).

Miller [35, p. 483] in his account about the mammals collected by W.L. Abbott on the west coast

of Sumatra in 1901 and 1902 mentioned the following about orangutans: "The orang utan exists,

but not abundantly, about Tapanuli Bay. Two miles up the Jaga Jaga River (no. 7 in Fig 2) some

nibong palms were seen that had been broken off by orangs, and also an old sarong (shelter),

but the traces were old. There were said to be more a few miles farther inland, particularly up

the Berdiri River (no. 6 in Fig 2). The natives say they always go about in pairs." Miller described

the Jaga Jaga River as "a stream near the south end of the Tapanuli or Sibolga Bay". We

located the Berdiri River on an old map under the name "Bardari River", and we located "Djaga

Djaga" as well. It seems clear from this that orangutans occurred in the area inland from the

southern part of Sibolga Bay (Fig 1) where there are currently no longer any orangutans.

Beccari [36] reported orangutans around Rambung, in the Tapanuli region, and in the hinterland

of Sibolga, where he collected a specimen. We were unable to determine the location of

Rambung but there is a Rambong north of the Singkil area, and thus well in *P. abelli* range. The

hinterland of Sibolga could either refer to the current *P. tapanuliensis* population or the historic

range. He further stated that "In the Zoological Museum at Florence is the skeleton of a young

orang-utan, described as coming from Palembang (Fig 1), on the east coast of Sumatra", some

800 km southeast of the nearest current orangutan population. We contacted the curator of the

Florence museum, who wrote in response that the specimen was indeed present under

specimen number MZUF-12: "The specimen was purchased in 1889 in London (G.A. Frank, 9

Haverstock Hill, London). It is a subadult male. The place of origin is Palembang, but may have

been captured elsewhere. There are no manifest connections with O. Beccari." Gustav Adolf

Frank was a well-known natural history trader based in Amsterdam and London, and he

probably had a good network of local suppliers. Description of the skeleton is provided by

Agnelli, Becchi [37] is inconclusive as to what species it belongs to. We can therefore not know

for sure whether the animal was caught near Palembang and transported to Europe from there,

or whether it originated from northern Sumatra (either of the two known species).

Volz [38] wrote about the distribution of orangutans on Sumatra, although it is not clear to what

extend the information is informed by Volz' own surveys or interpretation of secondary

information. Volz suggested that there were no orangutans east of the Langkat River, which he

thought was likely the remnant of a large bay or sea connection that once separated north and

south Sumatra approximately in a line from Sibolga to north of Medan. He expanded on this in

his work a few years later [39], in which he also described additional orangutan sightings. This

included a sighting in the upland area west of Lake Toba at an elevation of 1,400 m asl.

Referring to the same area, Kohler [40] described a visit to Sibolga where the guest had a

young orangutan which had also been caught in the forest on the west of the Toba lake,

indicating a breeding population there. Volz [38] also described a sighting of an orangutan east

of Lake Toba in the upper Kualu River area (ca. N 2°26' E 99°32', no. 8 in Fig 2). Again,

however, the description of a large ape that moved 'slowly and ponderously' may suggest an

adult male, and because people there are not familiar with the species, possibly a wandering

male outside the range of a breeding population.

Delmont [41] described a hunting expedition on the upper Musi River, near "Sekajoe" in the

foothills of the Barisan mountain range in what is now South Sumatra Province. His informant,

Mr Ghoba Ramah, told him that orangutans were particularly common in the area and were

raiding the crops of local farmers. After four weeks, they managed to catch seven orangutans.

They then moved to a location four hours rowing upstream, where they quickly observed a

female orangutan with young. They set out cages with fruit bait for capturing orangutans, but the

first morning after arrival they had only managed to catch some monkeys and a pig. After that

they were more successful and claimed to have caught one male orangutan and a female with

young, and over the next few days they caught several more orangutans. Delmont's stories are

intriguing but strike us as somewhat fantastical, as it is unlikely that anyone could catch

significant numbers of orangutans with baited cages. More likely these could be Pig-tailed

Macaques, *Macaca nemestrina*, who indeed move about in groups, raid crops and can be

trapped in cages. We therefore do not consider this source to be reliable.

The various historic accounts above where summarized in a map drawn by van Heurn [42]

which shows that clearly the conservation community was aware of the existence of orangutans

south, west and east of Lake Toba. Interestingly, though this map depicts the current Batang

Toru population to be part of the range where the species had become extinct, while the only

extant population is a narrow band to the east of Lake Toba in the Asahan District (Fig 1), where

the species is not currently known. It suggests that information about orangutan distribution was

still rudimentary in the 1930s, which may have the reason for a request to Carpenter [43] who

conducted a survey on behalf of the Nederlandsch-Indische Vereeniging tot

Natuurbescherming. He worked mostly in the northern parts of Sumatra but sent questionnaires

to Dutch soldiers stationed in areas where orangutans could potentially occur. Carpenter

assumed that orangutan did not occur south of a line drawn from Singkil to Sumatran east

coast, thus overlooking much of the historic evidence of orangutans south of Lake Toba.

Carpenter's questionnaires included two reports of orangutans outside the known range. The

first is from Captain H.J. Kloprogge who had been based in Siak, Indrapura and Pekanbaru (Fig

1) and claimed to have seen orangutans several times during forest patrols. Second, Captain M.

Kooistredie reported seeing 12 orangutans near Jambi (Muara Tembesi) (Fig 1) of which he

shot two and later another one. Again, we treat these reports as unreliable also because

Carpenter did not actually refer to them in his report and presumably did not assume the reports

to be accurate.

There seems to be a gap in records between the 1930s and 1970s, but in the early 1970s, the

Indonesian forester K. S. Depari reported orangutan sightings along the Batang Toru River and

in the Rimbu Panti Wildlife Reserve (no. 9 in Fig 2) [16]. Borner [44] also noted that in Rimbu

Panti one orangutan had been shot, while Wilson and Wilson [45] confirmed the presence of

orangutans in South Tapanuli, while also reporting them around Pekanbaru, in Riau Province.

Finally, the presence of orangutans was indicated by a botanist and a wildlife researcher on

Gunung Talamau (no. 10 in Fig 2; Laumonier, pers. comm.). Presence in this region was

confirmed by a survey in 1996 by Rijksen, Meijaard and van Schaik [13], when several nests

were found on the edge of this Reserve, but follow up surveys by SW could not confirm this

report and suggested that the nests may have been eagle nests. More recently, Wich et al. [46]

found several orangutan nests in the peat swamp forests near Lumut (no. 1 in Fig 2) and heard

a male long call in the same area. Local community members mentioned that they had also

seen orangutans in the area [46]. Approximately 2 km south of the Batang Toru River

(southeast of the village of Batang Toru), a geologist (Martin Jones) spotted a solitary orangutan

in the forest in 2004 (no. 12 in Fig 2). Finally, Bradshaw, Isagi [47] reported on orangutans in the

Barumun Wildlife Reserve in the Padang Lawas District (no. 13 in Fig 2). Nests were reportedly

encountered and one staff of the local conservation department reported a direct encounter with

an orangutan in 2009.

While there is significant spatial inaccuracy in the historic records of *P. tapanuliensis* outside the

current range, we can still make an educated guess of the different habitats and altitudes in

which these populations occurred (Table 1). Habitats in which the species once occurred

included tall peat swamp forest, freshwater swamp forest mosaic and secondary forest, forest

on limestone, hill forest between 300 and 1,000 a.s.l., and submontane forest between 1,000

and 1,800 m a.s.l., indicating the full range of habitats that is also used by *P. abelii* [16].

Table 1. Orangutan records, most likely *P. tapanuliensis*, but outside the current range,

that we consider to be reliable.

Number	Location	Source	Vegetation	Likely status
1[a]	Tapanoeli	Von Rosenberg 1878	Forest 300-1000m (logged)	Breeding population
2	Sayur Matinggi	Kramm 1879	Limestone/Secondary Forest 300 - 1000m	Breeding populaton
3	Loemoet	Kramm 1879	Tall Peat Swamp Forest	Breeding population
4	Hadjoran	Neumann 1885	Forest 300-1000m (Sibolangit)	Roaming male
5	West Coast	Hagen 1890	Forest 300-1000m (logged)	Unclear
6	Berdiri	Miller 1903	Secondary Mosaic <300m	Breeding population
7	Jaga Jaga	Miller 1903	Tall Swamp Forest (Shallow Peat)	Breeding population
8	Upland areas west of Lake Toba	Volz 1912	1400 m	Breeding population
9	Kualu River	Volz 1912	Forest <300m	Roaming male
10	Talamau Mountain	Laumonier 1983	Submontane Forest	Unclear

			1000 - 1800m	
11	Rimbu Panti	Meijaard 1997	Forest 300 - 1000m (Singkel)	Unclear
3	Lumut	Nests observed by Serge Wich and Tine Geurts in June 2001	Tall Peat Swamp Forest	Breeding population
3	Nests near Lumut	Nests observed by Serge Wich and Tine Geurts in June 2001	Tall Peat Swamp Forest	Breeding population
12	South of Batang Toru	Orangutan observed by Martin Jones (geologist) in 2004	Secondary Mosaic <300m	Unclear
13	Barumun	Bradshaw et al. 2007		Unclear

313 [a] Point numbers in the table correspond to the numbers in Fig 1. Vegetation information was

314 derived from Laumonier [26].

Discussion

Possible drivers of historical declines

Our historical ecology analysis of *P. tapanuliensis* has revealed that the species occurred

beyond its current range until quite recently. Breeding populations occurred in the Batang Gadis

area (Fig 1), probably through much of today's South Tapanuli and Mandailing Natal Regencies.

The historic records to the north of the current *P. tapanuliensis* range make it difficult to judge

whether these are part of *P. abelii* or *P. tapanuliensis*. For example, the records of orangutans

west of Lake Toba, could also refer to populations that still occur in the Pakpak Bharat Regency

[48] and the Batu Ardan and Siranggas forest blocks in the Dairi Regency [49], which genetically

are closer to *P. abelii* than *P. tapanuliensis* [50]. It is not clear whether the records in the

Padang Highlands, Rimbu Panti and Padang Lawas referred to itinerant males or breeding

populations but the scarcity of records could indicate that breeding populations became extinct

there earlier.

Because of the uncertainties in population status (breeding or not) and taxonomy, it is not

currently possible on the basis of the historic data to determine how large the former range of *P.*

tapanuliensis was. We know from evidence of Late Pleistocene orangutan fossils in the Padang

area (Lida Ajer, Ngalau Sampit, Ngalau Gupin) [51], that orangutans lived in this part of Sumatra

at least until some 50,000 years ago. What we don't know is whether this was *P. abelii*, *P.*

tapanuliensis, or a species different from both, as suggested by Drawhorn [52]. No specimens

of Tapanuli Orangutan were collected by any of the historic sources, except those reported by

Hagen [34] and Beccari [36], but these have not been genetically analysed yet. There is

therefore no robust evidence as to whether the orangutans reported from outside the current

Tapanuli Orangutan range were *P. tapanuliensis* or *P. abelii*. Further genetic study of the

specimens reportedly originating from Padang (RMNH.MAM.544) and Palembang (MZUF-12),

and also of fossil teeth from the Padang Caves area could shed light on the taxonomic status of

the orangutans in Central Sumatra, and their relationship to *P. tapanuliensis*. Based on

distribution range patterns, with *P. abelii* clearly restricted to the northern parts of Sumatra [12,

46], we consider it most likely that all historic orangutan populations south and south-east of the

current range of *P. tapanuliensis* were indeed *P. tapanuliensis*. This would indicate that the

species currently retains some 10-20% of the range it occupied some 150 to 200 years ago.

This raises the question why the species declined.

It is unclear when the populations outside the current range became extinct. Some populations

such as those in Lumut, seen by Neumann in 1885 and by SW in 2001, but not seen since, may

have become extinct quite recently because of forest loss. Other populations likely disappeared

sometime in the 20[th] century. There have been no recent records from areas west or southwest

of Lake Toba, nor from the Batang Gadis region, south of the current populations. Also, while

there are a number of alleged records from the area east of Padang City (nos 12 and 13 in Fig

1), there are no confirmed recent records. It thus appears that a lot of these populations

disappeared around a time when forests were still extensive and the commercial exploitation of

forest for timber (starting in the 1970s) or their conversion to plantations (starting in the 1990s)

had not yet decimated available habitats. Nevertheless, there had been significant historic

deforestation prior to 1950 as shown in Fig 1, mostly for small-holder agriculture and livestock,

firewood and timber, and as result of wars and fires [53, 54]. For example, the colonial-era

district of *Tapanoeli* (now North, South and Central Tapanuli) had an estimated forest cover of

19000 km^2 in a total area of 39,481 km^2 (48% forested) in the 1930s [53]. Some populations

therefore appear to have become isolated in historic times, when early agricultural development

created large grassland areas. So, why did these populations become extinct? This appears to

have been a combination of habitat loss and population fragmentation, and mortality rates that

exceeded reproductive replacement rates.

Several authors have suggested that orangutan density and range on both Borneo and Sumatra

were primarily determined by the ability of people to access areas and hunt orangutans [55, 56].

For example, Jentink (1889) writes that orangutans in Sumatra are only common in swamp

areas like those in Singkil which are so inaccessible that there are rarely "stepped on by human

feet", apparently quoting von Rosenberg [31], who had made a similar statement a decade

earlier. Wallace (1869) similarly argued that orangutans were common in swamp forest, not

because these were particularly suitable ecologically but rather because human hunters rarely

went there. Such hunting was certainly common in the orangutan's range in Sumatra. Schneider

[57], for example, writes that Batak people hunt orangutans with blow pipe, spears or shotguns,

while young animals are often caught and sold to plantation owners.

Batang Gadis was populated by Loeboe people [58, p. 327] who were nomadic tribes that also

occurred in Padang Lawas and "Groot-Mandailing" [59-61]. Another nomadic tribe, the Oeloes

occurred around Muara Si Pongi and Pahantan (now Pakantan) [59, 62]. Similar to other

nomadic people such as the Kubu further south in Sumatra [63] or the Punan of Borneo [64],

nomadic people often prefer primate meat over other meat sources. Kreemer [61] mentioned

that the Loeboe people consider primate meat a delicacy. They hunted primates, including

Siamang *Symphalangus syndactylus* with blowpipes, and used snares for pigs and deer. Still,

there are to our knowledge no specific accounts of people from the historic range of *P.*

tapanuliensis hunting and eating orangutans. Nevertheless, we consider it likely that, similar to

Borneo, orangutans would have been hunted for food. Van den Burg [65], in a general account

about orangutans, describes how generally orangutans were shot with poison darts, after which

they fell out of the trees and were killed with spears. Alternatively, they were caught alive and

killed later. The whitish meat was generally grilled over a fire, and was described as soft and

sweet [65]. This is also suggested by the use in local language of *juhut bontar*, or white meat, to

describe orangutan or pig-tailed macaque *Macaca nemestrina*, while such descriptions of sweet

meat were similarly recorded by EM on Borneo [23]. Orangutan fat, especially from adult males,

was often saved for later use in the preparation of other dishes [65].

Marshall et al. [66] examined population viability models with 1%, 2%, and 3% additional

mortality in all age classes, running 500 iterations with populations of 250 orangutans. In the

best quality orangutan habitats, i.e., mosaic landscapes of swamp, riparian and hill forests [23],

annual hunting rates of 1% did not cause population extinction, but did decrease population

size. In less than optimal habitats, e.g., forests at higher elevation, a 1% level of hunting caused

declines to extinction irrespective of initial population size. Higher rates of hunting were

unsustainable even in the highest quality habitats [66]. These models were conducted for

Bornean orangutan *P. pygmaeus*, but the authors thought that hunting effects would be similar

for Sumatran orangutans. The best orangutan habitats would like be those with the highest soil

fertility, which at levels of intermediate rainfall would also be the best areas for agriculture [67]. It

is thus likely that historically most Tapanuli orangutans would occur in suboptimal habitats,

where the removal of one animal from a population of 100 per year, would drive such a

population to extinction. Given the available information, we consider it most likely that the

Tapanuli orangutan was hunted to extinction in the increasingly fragmented parts of its former

range and only survived in the remote and rugged Batang Toru mountains which may have

provided orangutans with a refuge from hunting.

Implications for species conservation

What do our findings mean for conservation? The Tapanuli orangutan's remaining three

subpopulations are in apparent decline, threatened by conflict killing and hunting, and loss of

lowland habitat [14, 19, 20, 68]. Our insights from past population declines, driven by habitat

loss and fragmentation and probably unsustainable mortality rates, indicate that without

preventing further losses to the population, even if in the single numbers per year, the last

remaining populations of the species are doomed to rapidly decline within several orangutan

generation lengths [estimated at 25 years, 21]. Current killing rates of *P. tapanuliensis* already

meet or exceed this threshold. Two wild-captured infant Tapanuli orangutan were reported thus

far in 2020, with one confiscated from the owner and the other illegally released to avoid legal

repercussions [69, 70]. Obtaining wild orangutan infants necessitates killing the mother in nearly

all cases [23, 71], hence these infants are assumed to represent two adult females killed in the

first six months of 2020 alone. Such records are indicative of a lowest minimum number of

killings, as they represent only criminal acts that have been detected and acted upon, which is a

fraction of the total orangutan-related wildlife crime [72, 73]. Records of an adult male killed in

2013 (OIC pers. comm. 2020), a male severely injured by humans in 2019 [74], and another

male captured and translocated twice in the past 12 months due to complaints about crop

raiding from local community members [75] suggest that killings have been ongoing in recent

years, although prior to 2017 most detected incidents would have been recorded as *P. abelii*.

While translocation has been used as a response to Tapanuli orangutans in conflict with

humans, translocated animals are not monitored beyond a few days following release, and their

long-term survival is not known. Behavioural traits of female site fidelity and male territoriality,

and adaption issues of Bornean orangutans released in unfamiliar habitats indicate that

translocation risks are high and survival rates may be low [72, 76].

Long-term protection of the Tapanuli Orangutan requires that mortality rates of <1% per year

are maintained over long (decadal) time frames across the species' range. This also means that

that all subpopulations have to remain connected, because once connections between

populations are lost this should result in higher extinction risks for the remaining subpopulations,

as was modeled for Bornean orangutan *P. pygmaeus* [77]. Within the subpopulations, the

prevention of killing is urgently needed, which requires innovative management of crop conflicts

[78, 79], and effective law enforcement and awareness campaigns. Such campaigns have so far

had insufficient impact on reducing orangutan losses and new approaches may be required

[80]. This could include, for example, direct conditional payments to rural communities for

maintaining habitats and preventing any deaths or harm, i.e., orangutan guardians [81] or

support for "buffer gardens" to concentrate crop losses from orangutan foraging into areas

acceptable to communities [82].

Given the high extinction risks, it is important that a comprehensive plan of action is developed

for the species that accurately determines how many animals remain, the level of gene flow

between subpopulations, current loss rates (including removal of animals in rescues and

translocations), and works towards full and permanent protection of all remaining habitat. Such

a conservation management plan would need clarity about long-term funding, organizational

responsibilities, and clear, science-based plan to allow the Tapanuli orangutan population to

stop declining, or better, increase to safer population numbers. Without such concerted and

coordinated action, the remaining populations of the Tapanuli orangutan are doomed to follow

their historic predecessors on their path to rapid extinction.

452 Acknowledgements

453 We thank XXX reviewers for their constructive comments on an earlier version of this

454 manuscript.

455 References

1. Doherty TS, Dickman CR, Nimmo DG, Ritchie EG. Multiple threats, or multiplying the

threats? Interactions between invasive predators and other ecological disturbances.

Biological Conservation. 2015;190:60-8. doi:10.1016/j.biocon.2015.05.013.

2. Kapos V, Balmford A, Aveling R, Bubb P, Carey P, Entwistle A, et al. Outcomes, not

implementation, predict conservation success. Oryx. 2009;43(3):336-42.

doi:10.1017/S0030605309990275.

3. Sutherland W. Evidence-based Conservation. Conservation in Practice. 2003;4(3):39-

42. doi:10.1111/j.1526-4629.2003.tb00068.x.

4. Balint PJ, Stewart RE, Desai A, Walters LC. Wicked environmental problems: managing

uncertainty and conflict. Washington: Island Press; 2011.

5. Game E, Meijaard E, Sheil D, McDonald-Madden E. Conservation in a wicked complex

world; challenges and solutions. Conservation Letters. 2014;7(3):271-7.

doi:10.1111/conl.12050.

6. Adams WM, Sandbrook C. Conservation, evidence and policy. Oryx. 2013;47(3):329-35.

doi:10.1017/S0030605312001470.

7.   Meijaard E, Sheil D. The Moral Minefield of Ethical Oil Palm and Sustainable

Development. Frontiers in Forests and Global Change. 2019;2(22).

doi:10.3389/ffgc.2019.00022.

8.   Laurance WF, Wich SA, Onrizal O, Fredriksson G, Usher G, Santika T, et al. Tapanuli

orangutan endangered by Sumatran hydropower scheme. Nature Ecology & Evolution.

2020. doi:10.1038/s41559-020-1263-x.

9.   Gaveau DLA, Wich S, Epting J, Juhn D, Kanninen M, Leader-Williams N. The future of

forests and orangutans (Pongo abelii) in Sumatra: predicting impacts of oil palm

plantations, road construction, and mechanisms for reducing carbon emissions from

deforestation. Environmental Research Letters. 2009;4(3):34013. doi:10.1088/1748-

9326/4/3/034013.

10.   Swetnam TW, Allen CD, Betancourt JL. Applied historical ecology: Using the past to

manage for the future. Ecological Applications. 1999;9(4):1189-206. doi:10.1890/1051-

0761(1999)009[1189:AHEUTP]2.0.CO;2.

11.   Spehar SN, Sheil D, Harrison T, Louys J, Ancrenaz M, Marshall AJ, et al. Orangutans

venture out of the rainforest and into the Anthropocene. Science Advances.

2018;4:e1701422. doi:10.1126/sciadv.1701422

12.   Nater A, Mattle-Greminger MP, Nurcahyo A, Nowak MG, de Manuel M, Desai T, et al.

Morphometric, Behavioral, and Genomic Evidence for a New Orangutan Species.

Current Biology. 2017. doi:10.1016/j.cub.2017.09.047.

13.   Meijaard E. A survey of some forested areas in South and Central Tapanuli, North

Sumatra; new chances for orangutan conservation. Wageningen: Tropenbos and the

Golden Ark; 1997.

14.   Wich SA, Fredriksson G, Usher G, Kuhl H, Nowak M. The Tapanuli orangutan: Status,

threats and steps for improved conservation. Conservation Science and Practice. 2019.

doi:10.1111/csp2.33.

15.   Wich SA, Singleton I, Nowak MG, Utami Atmoko SS, Nisam G, Arif SM, et al. Land-
cover changes predict steep declines for the Sumatran orangutan (*Pongo abelii*).
Science Advances. 2016;2(3):e1500789. doi:10.1126/sciadv.1500789

16.   Rijksen HD. A fieldstudy on Sumatran orang utans (*Pongo pygmaeus abelii* Lesson
1827). Ecology, behaviour and conservation [PhD thesis]. Wageningen, The
Netherlands: Agricultural University Wageningen; 1978.

17.   IUCN Section on Great Apes. Batang Toru Hydropower Project. Factcheck and
References on Key Issues. IUCN/SSC Primate Specialist Group; 2020.

18.   Wich S, Fredriksson G, Usher G, Peters H, Priatna D, Basalamah F, et al. Hunting of
Sumatran orang-utans and its importance in determining distribution and density.
Biological Conservation. 2012;146(1):163-9.

19.   Nasution A, Perwitasari-Farajallah D, Utami-Atmoko SS. Declining Orangutans
Population in the Unprotected Forest of Batang Toru. Trop Life Sci Res. 2018;29(2):77-
87. doi:10.21315/tlsr2018.29.2.6.

20.   Samsuri, Surati Jaya IN, Kusmana C, Murtilaksono K. Connectivity and ecological
indicators analysis of tropical forest landscape in Batang Toru watershed - Indonesia.
Agriculture, Forestry and Fisheries. 2014;3(3):147-54. doi:10.11648/j.aff.20140303.12.

21.   Nowak MG, Rianti P, Wich SA, Meijaard E, Fredriksson G. *Pongo tapanuliensis*. The
IUCN Red List of Threatened Species 2017. 2017:e.T120588639A62.
doi:10.1111/csp2.33.

22.   Hooijer DA. Prehistoric teeth of man and of the orang utan from Central Sumatra, with
notes on the fossil orang utan from Java and Southern China. Zoölogische
Mededelingen Rijksmuseum Leiden. 1948;29:175 - 83.

23.   Rijksen HD, Meijaard E. Our vanishing relative. The status of wild orang-utans at the
close of the twentieth century. Dordrecht, The Netherlands: Kluwer Academic
Publishers; 1999. 480 p.

24.   Planning Department of the Forest Service. Vegetation map of Indonesia. Scale 1:

524   2500000. Under supervision of L.W. Hannibal. 1950.

525   25.   ESRI. What is ArcScan?. ArcGIS Desktop. Retrieved on 17 June 2020 from

526   https://desktop.arcgis.com/en/arcmap/latest/extensions/arcscan/what-is-arcscan-.htm.

527   2019.

528   26.   Laumonier Y. The vegetation of Sumatra. Bogor, Indonesia: SEAMEO-BIOTROP; 1997.

529   154 p.

530   27.   Schlegel H, Müller S. Bijdragen tot de natuurlijke historie van den orang-oetan (*Simia*

531   *satyrus*). In: Temminck CJ, editor. Verhandelingen over de natuurlijke geschiedenis der

532   Nederlandsche overzeesche bezittingen: Zoologie. Leiden, the Netherlands: Van den

533   Koning; 1841. p. 28.

534   28.   Martyr D. The other orang. BBC Wildlife. 1993 October 1993:35 - 6.

535   29.   Westenenk LC. Orangpandak (boschmensen) op Sumatra. Tropische Natuur.

536   1919;7:108- 10.

537   30.   Müller DS, Horner DL. Fragmenten uit de Reizen en Onderzoekingen in Sumatra

538   (Vervolg van Deel III, blz. 72). Bijdragen tot de taal-, land- en volkenkunde / Journal of

539   the Humanities and Social Sciences of Southeast Asia. 1855;3(1):193-228.

540   doi:10.1163/22134379-90001160.

541   31.   von Rosenberg H. Der Malayische Archipel. Land und Leute. Leipzig1878.

542   32.   Kramm W. Tochtjes in Tapanoeli. Sumatra-Courant. 1879;20(180 (Thursday 30

543   October)):1-2.

544   33.   Neumann JB. Het Pane- en Bila-stroomgebied op het eiland Sumatra. Tijdschrift

545   Nederlandsch Aardrijkskundig Genootschap. 1885;2(II):2 and 1-133 (also III, 1-99, 215-

546   314 &457-543; IV, 1-110&217-319).

34. Hagen B. Die Pflanzen- und Thierwelt von Deli auf der Ostkuste Sumatra's.

Naturwissenschaftliche Skizzen und Beitraege. Tijdschrift Nederlandsch Aardrijkskundig

Genootschap. 1890;7(1):1-240.

35. Miller GS. Mammals collected by Dr. W.L. Abbott on the coast and islands of northwest

Sumatra. Proceedings US National Museum, Washington. 1903;26:437-84.

36. Beccari O. Wanderings in the great forests of Borneo. From the English translation

published by Archibald Constable & Co. Ltd, London, reprinted in 1986. Oxford, UK:

Oxford University Press; 1904. 424 p.

37. Agnelli P, Becchi C, Cecconi S, Fontani S, Guimaraes S, Moggi I. Indagine storico e

studio morfometrico dell'esemplare di *Pongo pygnaeus* no MZUF 12. Museo di Storia

Naturale dell'Univercità degli Studi di Firenze. Sezione di Zoologia "La Specola"; 2002.

38. Volz W. Uber die Verbreitung einiger anthropoiden Affen in Sumatra. Biologische

CentralBlatt. 1904;24(14):475-6.

39. Volz W. Nord-Sumatra; bericht über eine im auftrage der Humboldt-stiftung der Königlich

preussischen akademie der wissenschaften zu Berlin in den jahren 1904-1906

augeführte Forschungreise. Bd. II. Die Gajolander. Berlin: Dietrich Reimer (E. Vohsen);

1912.

40. Köhler HJ. Habinsaran. Het land van den zonnestraal. Mijn leven onder de Bataks.

Zutphen: W.J. Thieme & Cie; 1926.

41. Delmont J. Levende Natuur: Het Vangen Van Wilde Dieren. De liefde van een Orang-

Oetan-wijfje. Niewsblad Van Het Noorden. 1930(December 27, 1930):3.

42. van Heurn FC. De anthropoide apen, de rhinocerossen en het baardzwijn. The

Netherlands: Overgedrukt uit de Mededeeling van de Nederlandsche Commissie for

Internationale Natuurbescherming, Biologische Aanteekeningen over eenige

belanggrijke diersoorten van Java, Sumatra en Borneo; 1935. Report No.: No.10

Supplement.

43.   Carpenter CR. A survey of wildlife conditions in Atjeh, North Sumatra. Mededelingen

Nederlandsche Commissie voor Internationale Natuurbescherming. 1938;12:1-33.

44.   Borner M. Sumatra's orang-utans. Oryx 1976;13:290-3.

doi:10.1017/S0030605300013752.

45.   Wilson CC, Wilson WL. Behavioral and morphological variation among primate

populations in Sumatra. Yearbook of Physical Anthropology. 1977;20:207-23.

46.   Wich SA, Singleton I, Utami-Atmoko SS, Geurts ML, Rijksen HD, van Schaik CP. The

status of the Sumatran orang-utan *Pongo abelii*: an update. Oryx. 2003;37(1):49-54.

doi:10.1017/S0030605303000115.

47.   Bradshaw CJA, Isagi Y, Kaneko S, Brook BW, Bowman DMJS, Frankham R. Low

genetic diversity in the bottlenecked population of endangered non-native banteng in

northern Australia. Molecular Ecology. 2007;16(14):2998-3008. doi:10.1111/j.1365-

294X.2007.03365.x.

48.   Sembiring J, Azvi TS. Estimation of orangutan populations in Sibongkaras Pakpak

Bharat village forest area. Journal of Islamic Science and Technology. 2019;6(1):133-45.

doi:10.22373/ekw.v6i1.5549.

49.   Utami Atmoko SS, Sihite J, Nuzuar, Singleton I. Perkembangan terbaru dan sabaran

orangutan sumatera (*Pongo abelii*). In: Susilo HD, Utami Atmoko SS, Sihite J, editors.

Workshop Proceedings: International Workshop on Orangutan Conservation, Sanur,

Bali, Indonesia 20102012. p. 105-10.

50.   Nater A, Arora N, Greminger MP, van Schaik CP, Singleton I, Wich SA, et al. Marked

Population Structure and Recent Migration in the Critically Endangered Sumatran

Orangutan (Pongo abelii). Journal of Heredity. 2013;104(1):2-13.

doi:10.1093/jhered/ess065.

51.   Louys J, Zaim Y, Rizal Y, Aswan, Puspaningrum M, Trihascaryo A, et al. Sumatran

orangutan diets in the Late Pleistocene as inferred from dental microwear texture

analysis. in press.

52.   Drawhorn GM. The systematics and Paleodemography of fossil Orangutans (Genus

Pongo) [PhD Dissertation]: University of California; 1994.

53.   van Steenis CGGJ. Maleische Vegetatieschetsen I: Toelichting bij de

Plantengeografische Kaart van Nederlandsch Oost Indië. In: Brill EJ, editor. Tijdschrift

van het Koninklijk Nederlandsch Aardrijkskundig Genootschap, Tweede Serie Deel Lii.

Leiden, The Nederlandsch1935. p. 1-56.

54.   Endert FH. Boschbouwkundige aanteekeningen over een reis in Atjeh, Sum. Oostkust,

Tapanuli en Sumatra' Westkust. Tectona. 1925;18:1-60.

55.   Jentink JA. On a collection of mammals from East-Sumatra. Proceedings of the

Zoological Society of London. 1889;11:17-30.

56.   Wallace AR. The Malay Archipelago. reprinted in 1986 ed. Oxford: Oxford University

Press; 1869. 625 p.

57.   Schneider G. Ergebnisse zoologischer Forschungsreisen in Sumatra. Erster Teil.

Säugetiere (Mammalia). Zoologischer Jahrbücher Abteilung für Systematik, Geographie

und Biologie der Tiere. 1906;23:1-172.

58.   Reitsma SA. Van Stockums's travellers' handbook for the Dutch East Indies: Van

Stockum; 1930. XI, 613 p. p.

59.   Joustra M. Batakspiegel: S.C. van Doesburgh; 1926. XVI, 382 p p.

60.   van Dijk PALE. Rapport over de Loeboe-bevolking in de onder afdeeling Groot-

Mandheling en Batang Natal. Bijdragen tot de taal-, land- en volkenkunde van

Nederlandsch-Indië. 1884;8:151-61.

61.   Kreemer J. De Loeboes in Mandailing. Bijdragen tot de Taal-, Land- en Volkenkunde

van Nederlandsch-Indië. 1912;66:303-36.

62.   Ris H. De Onderafdeeling Klein Mandailing Oeloe en Pahantan en Hare Bevolking met

624   Uitzondering van de Oeloe's. Bijdragen tot de Taal-, Land- en Volkenkunde van

625   Nederlandsch-Indië. 1896;46(3):441-534.

626   63.   Veth PJ. Midden-Sumatra. Reizen en Onderzoekingen der Sumatra Expeditie 1877-

627   1879. Derde Deel. Volksbeschrijving en Taal. Leiden: E.J. Brill; 1892.

628   64.   Stolk JJ. Opsporing van den zwervenden stam der Penjaboeng-Poenan's, op de

629   waterscheiding der Barito met de Mahakam en Kapoeas (Midden-Borneo), in Oct. 1905.

630   Koninklijk Nederlands Aardrijkskundig Genootschap. 1907;24.

631   65.   van den Burg CL. De Voeding in Nederlandsch-Indië. Amsterdam: J.H. de Bussy; 1904.

632   546 p.

633   66.   Marshall AJ, Lacy R, Ancrenaz M, Byers O, Husson S, Leighton M, et al. Orangutan

634   population biology, life history, and conservation. Perspectives from population viability

635   analysis models. In: Wich S, Atmoko SU, Mitra Setia T, van Schaik CP, editors.

636   Orangutans: geographic variation in behavioral ecology and conservation. Oxford, UK:

637   Oxford University Press; 2009. p. 311-26.

638   67.   Santika T, Ancrenaz M, Wilson KA, Spehar S, Abram N, Banes GL, et al. First

639   integrative trend analysis for a great ape species in Borneo. Scientific Reports.

640   2017;7(1):4839. doi:10.1038/s41598-017-04435-9.

641   68.   Nowak MG, Rianti P, Wich SA, Meijaard E, Fredriksson G. Pongo tapanuliensis. The

642   IUCN Red List of Threatened Species2017.

643   69.   Karokaro AS. Call for prosecution of Indonesian politician who kept baby orangutan as

644   pet. https://news.mongabay.com/2020/02/tapanuli-orangutan-indonesia-sumatra-official-

645   nikson-nababan-pet/. Mongabay. 2020(26 February 2020).

646   70.   Pasaribu O. Anak Orangutan Tapanuli Masuk Perkampungan, Ditemukan Warga dalam

647   Kondisi Lemah. https://regional.kompas.com/read/2020/06/19/13323731/anak-

orangutan-tapanuli-masuk-perkampungan-ditemukan-warga-dalam-kondisi. Kompas.

2020;19 June 2020.

71.   Russon AE. Orangutan rehabilitation and reintroduction. In: Wich S, Atmoko SU, Setia

TM, van Schaik CP, editors. Orangutans: Geographic variation in behavioral ecology

and conservation. Oxford, UK: Oxford University Press; 2009. p. 327-50.

72.   Sherman J, Ancrenaz M, Meijaard E. Shifting apes: Conservation and welfare outcomes

of Bornean orangutan rescue and release in Kalimantan, Indonesia. Journal for Nature

Conservation. 2020;55. doi:10.1016/j.jnc.2020.125807.

73.   Freund C, Rahman E, Knott C. Ten years of orangutan-related wildlife crime

investigation in West Kalimantan, Indonesia. American Journal of Primatology.

2017;79(11):22620. doi:10.1002/ajp.22620.

74.   Gunawan A. Malnourished, injured Tapanuli orangutan rehabilitated and released.

https://www.thejakartapost.com/news/2019/12/10/malnourished-injured-tapanuli-

orangutan-rehabilitated-and-released.html. The Jakarta Post. 2019;10 December 2019.

75.   KSDAE. Penanganan Konflik Orangutan di Adiankoting Sumatera Utara.

http://ksdae.menlhk.go.id/info/7489/penanganan-konflik-orangutan-di-adiankoting-

sumatera-utara.html. Ministry of Environment and Forestry. 2020;16 January 20202.

76.   Ancrenaz M, Oram F, Nardiyono, Silmi M, Jopony MEM, Voigt M, et al. The importance

of orangutans in small fragments for maintaining metapopulation dynamics. bioRxiv.

2020:2020.05.17.100842. doi:10.1101/2020.05.17.100842.

77.   Bruford MW, Ancrenaz M, Chikhi L, Lackman-Ancrenaz I, Andau M, Ambu L, et al.

Projecting genetic diversity and population viability for the fragmented orang-utan

population in the Kinabatangan floodplain, Sabah, Malaysia. Endangered Species

Research. 2010;12:249-61. doi:10.3354/esr00295.

78.   Campbell-Smith G, Campbell-Smith M, Singleton I, Linkie M. Raiders of the Lost Bark:

673   Orangutan Foraging Strategies in a Degraded Landscape. PloSONE. 2011;6

674   (6):e20962. doi:10.1371/journal.pone.0020962.

675   79.   Campbell-Smith G, Sembiring R, Linkie M. Evaluating the effectiveness of human-

676   orangutan conflict mitigation strategies in Sumatra. Journal of Applied Ecology.

677   2012;49:367-75. doi:10.1111/j.1365-2664.2012.02109.x.

678   80.   Karokaro AS, Hanafiah J. Indonesia rescues captive orangutans, but leaves their owners

679   untouched. Mongabay. 2019 February 8 2019.

680   81.   Schoneveld-de Lange N, Meijaard E, Löhr A. South to south learning in great ape

681   conservation. American Journal of Primatology. 2016;78(6):669-78.

682   doi:10.1002/ajp.22531.

683   82.   Gehrke V. Insights from Indonesia: The Social Impact of Environmental Conservation.

684   https://savegporangutans.org/insights-from-indonesia-the-social-impact-of-

685   environmental-conservation/. GCOCP. 2020;3 August 2020.