Individual heterogeneity and its importance for metapopulation
dynamics

Keywords:

Demography; Experimental Ecology; Arthropoda; Spatial Network; Dispersal

N. of Words in the main text: 5293

Elements of the manuscript: main text; figures 1-4; Supplementary Materials 1-7.

Article Type: Major Article

Abstract

Landscape connectedness shapes the exchange of individuals among patches, and hence
metapopulation connectivity and dynamics. Connectedness, and its resulting effects on
connectivity are therefore rightfully central in conservation biology. However, besides determining
demographic fluxes of individuals between patches, connectedness also generates phenotypic
sorting and thus impacts local and regional eco-evolutionary dynamics. Despite the central role of
connectedness, its effects on individual phenotypic heterogeneity and spatial organization are so
far neglected in theory and applications.
Through experimental metapopulations of *Tetranychus urticae* (two-spotted spider mite) with
three levels of landscape connectedness and by regularly removing phenotypic structure in a
subset of these populations, we tested how regional and local population dynamics are
determined both by network connectedness and phenotypic spatial organization.
We find that the self-organizing phenotypic spatial structure increases local equilibrium population
sizes and variability. It in contrast dampens the effects of imposed connectedness differences on
population sizes and is therefore anticipated to improve metapopulation persistence. Contrary to
theoretical expectations, the most locally connected patches within the network showed an
overall reduced local population size, possibly originating from a faster depletion of resources
from immigrants or transiting individuals.
This experiment shows how metapopulation dynamics can significantly deviate from theoretical
expectations due to individual heterogeneity. This potential rescue effect stemming from
phenotypical self-organization in space is a key point to consider for conservation actions,
especially based on translocations.
33

Introduction

Habitat fragmentation has a deep impact on the population structure and demography of almost every species at any trophic level (Ims and Andreassen 1999; Crooks 2002; Cushman 2006; McGuire et al. 2016). Even without any habitat loss, the physical disruption of a previously continuous habitat into a series of patches separated by a more hostile matrix can result in connectivity loss. Connectivity can be defined as the realized dispersal of individuals and/or propagules among patches of suitable habitat. It therefore depends on one end on the connectedness of the landscape, a measure of the average physical proximity among patches that results from the distance between them (Turner 1989; Cushman, McGarigal, and Neel 2008; Wang, Blanchet, and Koper 2014). Connectivity depends on how connectedness interacts with the dispersal capacities of the species involved (Tischendorf and Fahrig 2000). Barring indirect effects of evolution, loss of connectedness is expected to decrease connectivity, reducing the probability that an empty patch will be colonized, or that patches at the brink of extinction will be rescued; overall, it tends to lower the persistence and stability of a metapopulation (Fuller, Doyle, and Strayer 2015; Thompson, Rayfield, and Gonzalez 2017) and is therefore of prime importance for conservation biology.

Connectedness impacts local and regional fluctuations in population sizes, as well as the synchronicity of these changes within the entire metapopulation (De Roissart, Wang, and Bonte 2015; Wang, Haegeman, and Loreau 2015). Theory predicts that lower effective dispersal between disconnected patches raises both local and regional temporal variabilities in population size thereby increasing local patch extinctions while decreasing rescue from immigration (Abbott 2011; Wang, Haegeman, and Loreau 2015; Zhang, DeAngelis, and Ni 2021). On the other

hand, a more connected network will also show a higher level of spatial synchronicity (Gouhier,

Guichard, and Gonzalez 2010), which makes the whole system also more vulnerable to global

extinction because of declines in rescuing potential when all local populations decrease at the

same time (Lloyd and May 1999; Sabelis et al. 2005). This holds especially when local patches

are small, as small local populations are vulnerable to demographic stochasticity (Fahrig 2017).

Recent theory shows, however, a more complicated picture with local variability and

synchronicity in these local population fluctuations to depend on the pace of population growth

and recovery after perturbance (Luo et al. 2021). Connectivity can thus be considered a double-

edged sword for metapopulation conservation (Hudson and Cattadori 1999), but it is clear that

more insights are needed on the conditions causing perils for metapopulation persistence.

All these theoretical developments ignore evolutionary dynamics and the potential for eco-

evolutionary feedbacks. Landscape connectedness directly affects the costs of dispersal (Bonte

et al. 2012), imposing selection pressures on dispersal and life history. The resulting life history

and dispersal evolution will in turn feedback on connectivity and demography (Bonte, Masier,

and Mortier 2018; Govaert et al. 2019). The evolutionary feedbacks can occur at both the

regional (metapopulation) and local (patch) level. Connectivity is an environmental pressure

affecting all individuals in a suite of patches which may impact the demography and therefore

competition in single patches. Metapopulation dynamics are consequently expected to be

influenced by metapopulation-level selection on dispersal and associated traits that may equally

evolve at the local patch level. This can include, but is not limited to, feeding efficiency, offspring

investment and energy allocation (Masier and Bonte 2020). Indeed, the position and local

connectedness of each patch within the spatial network will determine the number of

immigrants (Tischendorf and Fahrig 2000; Fahrig 2017; Bonte and Bafort 2019; Karisto and Kisdi

2019). Patches with higher levels of local connectedness are for instance expected to have an

increased immigration, that should theoretically increase population densities (Ives et al. 2004;

Masier and Bonte 2020). Alternatively, as less connected patches will receive lower numbers of

immigrants, they should become genetically more isolated, kin-structured and inbred.

Connectivity not only affects fluxes of individuals among patches, and therefore local and

regional population sizes, it is also expected to determine the phenotypic structure within the

metapopulation. Changes in local densities affect competition and therefore directly impact the

phenotypic signature of the developing individuals (Lion 2018). This phenotypic plasticity is

usually expressed in body size, morphology, physiology, and behavior (Parsons et al. 2011;

Koeller, Mohn, and Etter 2000; Kamioka and Iwasa 2017; Le Galliard, Paquet, and Mugabo 2015;

Woodworth et al. 2017). Since dispersal is performed by a non-random subset of individuals

from the population that share specific life history traits (Bonte et al. 2012; Clobert et al. 2012;

Dahirel et al. 2019) and physiology (Goossens et al. 2020), connectivity changes will equally

affect the sorting of phenotypes, and therefore phenotypic variation (*kind structure*) among

patches. The spatial organization of phenotypes among and within patches then again has the

potential to affect population dynamics. More-over, because of intimate links to gene flow and

thus spatial genetic structure, connectivity will directly impact interactions among relatives (*kin*

structure). In absence of realized dispersal, competitive interactions among kin are elevated

(Perrin and Mazalov 2000). Kin competition is known to be an important driver of dispersal

(Gandon 1999), thus potentially imposing negative feedbacks on the metapopulation dynamics

(Poethke, Pfenning, and Hovestadt 2007). At the same time, local interactions between

individuals will impose an internal self-organization, with phenotypes exclusively emerging from

local demographic interactions (Lowe, Muhlfeld, and Allendorf 2015). This emergent complexity,

resulting from a balance between order (phenotypical sorting) and disorder (stochastic

distribution of phenotypes) determines the demographic self-organization of the entire

metapopulation (Parrott 2010; Urban et al. 2020).

Individual heterogeneity, phenotypic self-organization, selection and the resulting population

dynamics are anticipated to impact metapopulation viability (Thompson, Rayfield, and Gonzalez

2017). We used the two-spotted spider mite *Tetranychus urticae* as a model to demonstrate

how the spatial self-organization of phenotypes (i.e., the emerging kin and kind structure)

equalizes metapopulation dynamics across connectedness levels (Urban et al. 2020). The

prerequisites for such a phenotypic self-organization are present: phenotypic variation in

relation to local densities, stage and sex structure (Dahirel et al. 2019; De Roissart, Wang, and

Bonte 2015), density and kin-competition related dispersal (Van Petegem et al. 2018; Fronhofer,

Poethke, and Dieckmann 2015). Moreover, previous studies showed that this species'

phenotype-dependent dispersal maximizes fitness (Dahirel et al. 2019; Bonte et al. 2014). We

earlier demonstrated that changes in connectedness impose genetic and epigenetic changes in

dispersal timing, dispersal costs and fecundity, but not dispersal propensity (Masier and Bonte,

2020).

We initialized replicated experimental metapopulations according to three levels of

metapopulation-level connectedness and implemented a randomization treatment by

reshuffling individuals across the patches in the metapopulation, removing the spatial geno- and

phenotypic structure while leaving demographic attributes like densities, stage, sex, and age-

structure untouched. This treatment thus removed most of the emerging spatial kin and kind

structure (Masier and Bonte, 2020). We quantified and compared average local and

metapopulation sizes, as well as the level of demographic fluctuations (local and

metapopulation variability) and synchrony in local population size changes between treatments

(Wang and Loreau 2014). As variation in local connectedness may also affect local dynamics, we

additionally compared responses from patches within the same spatial network that varied in

their degree of local connectedness (i.e., core vs edge patches within a single metapopulation).

We generated some prior predictions on the role of connectedness versus connectivity on

metapopulation demography by means of an individual based model, and we later compared

them to our experimental results.

Materials and methods

<u>Study species</u>

To perform the experiment, we used the two-spotted spider mite *Tetranychus urticae* Koch

1836 (Acari: Tetranychidae) as a model. *T. urticae* is a parasite on a wide diversity of host plants

(Migeon, Nouguier, and Dorkeld 2010), and has short developmental time (7 to 10 days at

experimental conditions (Hance and Van Impe 1999)). We used the "LS-VL" lab strain

established in October 2000 (Van Leeuwen et al. 2008; Cazaux et al. 2014) as the base

population of spider mites. This population has been maintained on bean (*Phaseolus vulgaris* L.,

variety 'Prelude') in climate-controlled rooms and under a constant L:D 16:8 light regime since

then. As many experiments showed through the years (e.g., Van Petegem *et al.*, 2018), this

strain shows a high level standing genetic variation and is therefore of perfect use in

experimental evolution (Kawecki et al. 2012).

<u>Experimental setup</u>

To understand the effect of spatial self-organization and spatial connectedness on ecological

dynamics, we studied spider mites in 24 independent experimental metapopulations varying in

connectedness levels and the application or not of a randomization treatment (see below).

These metapopulations are the ones studied in (Masier and Bonte, 2020) for phenotypic

evolution; we present here a summary of the experimental setup. Further practical details about

the setup and maintenance of the experimental metapopulations are provided in (Masier and

Bonte, 2020) and as supplementary material (Supplementary Material S01).

Each experimental metapopulation consisted of a 3-by-3 square grid of nine patches, with

connections between patches following Moore neighborhood rules (each patch connected to its

"horizontal", "vertical" and diagonal neighbors). Patches were all the same size (5×5 cm²) and

were each cut from one of the two main leaves of a two-weeks-old bean plant. Patches were

connected using Parafilm bridges; each bridge was about 0.5 cm wide, and the length of the

bridges defined the "metapopulation-level connectedness" treatment: 8 metapopulations were

closely connected (4cm long bridges), 8 averagely connected (8cm) and 8 loosely connected

(16cm). Lengths in the previous sentence refer to all horizontal and vertical bridges; diagonal

bridges were $\approx \sqrt{2}$ times longer. Experimental trials showed that mortality during movement

increased with bridge length (Supplementary Material S02).

Within each replicate metapopulation, patches could be distinguished by their level of "local

connectedness": central patches were connected to all other patches (8 links), while side and

corner patches were less connected (5 and 3 links, respectively, (Masier and Bonte, 2020)).

Every week, we counted and collected all adult females using a thin paintbrush, put them on

fresh bean leaves to reduce stress as much as possible, after which they were collectively put

back in the metapopulation. In the standard treatment used as a control for the reshuffling

treatment (see lower), they were put back on the same leaf as the one they were collected

from. To tease out the consequences of phenotypic and genotypic heterogeneity (kind and kin

structure), we subjected three replicates out of eight in each metapopulation connectedness

level to a randomization treatment. In these, collected females were then shuffled and moved

back on a random patch of the same metapopulation. Randomization was done so the number

of adult females per patch remained unchanged pre- and post-manipulation. As a result, both

the absolute population numbers and the stage/sex structure of each patch remained unaltered

by randomization itself. The treatment therefore specifically destroys, or at least reduces, kin

and kind structure (genetic relatedness and phenotypic similarity; (Van Petegem et al. 2018)) as

well as any local adaptation caused i.e. by local density fluctuations, while maintaining the direct

ecological effect of connectedness on population densities themselves.

We decided to reshuffle only three out of eight replicates to ensure feasibility over the course of

the full experiment (10 months) while optimizing statistical power for both these demographic

and earlier evolutionary analyses (Masier and Bonte, 2020).

Each patch was initialized at the start of the experiment using 5 adult females and 2 adult males

taken randomly from our stock population (63 individuals per metapopulation, 1512 individuals

overall). After a 15-days (≈1.5 generations) "burn-in" period during which no count or

randomization was performed to allow (meta-)populations to reach an equilibrium (which was

reached out in 21 out of 24 replicates, Supplementary Material S03), the metapopulations were

monitored for 23 weeks (≈18 generations at the described conditions). This resulted in a total of

4941 abundance records at the patch level (9 patches × 24 replicates × 23 weeks, minus three

replicates missing one week each due to data transcription errors).

<u>Individual-based model</u>

To understand the importance of connectedness for the measured metapopulation metric, we

developed an individual-based version of a metapopulation model that resembled our

experimental systems (3×3 patch networks). The model did not incorporate evolutionary

dynamics or context (e.g. density-)dependent dispersal. Changes in connectedness were

implemented by simulating higher dispersal mortality (as from the experimental result

presented in material [Supplementary Material S02]). An extensive model description is

provided in the supplementary material (Supplementary Material S04). We present results from

this simulation model prior to those observed in the effective experiment.

<u>Metapopulation dynamics: statistical analysis</u>

Following Wang and Loreau (2014), we can describe the temporal dynamics of each

metapopulation by the patch-level average population sizes and a series of four metrics

summarizing variability at multiple scales (Table 1). To calculate these metrics and compare

them across treatments, one needs to gather estimates of mean population sizes for each patch and temporal covariances between patches for each metapopulation, as well as information about their uncertainties. Wang and Loreau (2014) do not provide ready-made methods to estimate these uncertainties; however, one can readily obtain both point estimates and uncertainties for each of these metrics using the output from a multivariate mixed/multilevel model where each patch is its separate response and patch responses are correlated (Supplementary Material S05).

Using numbers of adult females per local patch as our metric, we analyzed mite abundance data by fitting a Poisson multivariate multilevel model (Supplementary Material S05). We fitted this model in a Bayesian framework using the Stan language (Carpenter et al. 2017). The use of a MCMC-Bayesian framework is especially valuable here as, among other things, posterior samples greatly facilitate error propagation when combining multiple model parameters to estimate the metrics (described in Table 1). We wrote this model and analyzed it using R, version 4.0.5 (Team R Core 2021) and the *brms* package version 2.14.4 (Bürkner 2018) as frontends. For simplicity, we only present a conceptual summary below: a complete and formal description, including rationale for prior choices and how the metrics described in Table 1 can be derived from model parameters (fixed effects, random effects and (co)variances), is included in the archived code (see "Data and code availability") and in supplementary material (Supplementary Material S05).

This model included fixed effects of metapopulation-level connectedness, local connectedness, randomization, and the two randomization × connectedness interactions, as well as random effects of metapopulation and patch identity (nested in metapopulation). We converted fixed effects to centered dummy variables to facilitate the interpretation of interaction coefficients

(Schielzeth 2010) and the setting of priors. The model also included patch-specific observation-

level random effects (Harrison 2014), to account for within-patch temporal variability. These

random effects were correlated between patches, to allow for temporal covariance among

patches (the w_{ij} in Table 1). Regarding this within-patch (co)variation, it is important to note that

because landscapes can be freely rotated, x,y encodings for patch coordinates are somewhat

arbitrary and only valid within a given replicate metapopulation. This has important

consequences for model specification: indeed, the $x = 1$, $y = 1$ corner patches in any two

metapopulations do not intrinsically "match" each other more than, say, the $x = 1$, $y = 1$ corner

patch in one landscape and the $x = 1$, $y = 3$ corner patch in another. This prevents explicit in-

model partitioning of covariance matrices into mean treatment-level and residual components.

As a result, we fitted one within-patch covariance matrix per replicate, and posterior averaging

among replicates from a same treatment was only done once all within-replicate operations

were complete.

In a multilevel/mixed Poisson model as here, mean population sizes depend on both latent

intercepts and within-patch variability (de Villemereuil et al. 2016). Importantly, if the latter

random effect variance is allowed to vary between treatments, as is the case here, then we

need to account for it to accurately predict average differences due to connectedness and/or

reshuffling, as differences based on fixed effects only may not show the whole picture (Table 1).

We therefore used formulas in (de Villemereuil et al. 2016) to estimate posterior means on the

observed scale from latent-scale intercepts and covariance matrices, and compare them

between treatments. We additionally used methods in (de Villemereuil et al. 2016) to convert

posterior temporal covariance matrices from our model, which are on the latent log scale, to the

observed data scale, which is the scale relevant for the variability metrics (Table 1).

We compared mean local population sizes between replicates as a function of connectedness

(local and metapopulation-level) and randomization treatment, and connectedness ×

randomization interaction. For the variability metrics (α, β_2, $\square$, γ), due to a reduced number of

experimental points at the metapopulation scale, we focused solely on the additive effects of

connectedness and randomization.

In a secondary model, we analyzed mean metapopulation-level abundances between

treatments, i.e., the response variable was here the sum of adult females in one replicate.

Because there was no internal covariation to worry about at that scale, we used a negative

binomial model, in which both the mean abundance and the within-metapopulation temporal

variability (through the shape parameter) depended on fixed effects of landscape

connectedness, randomization and their interaction, and random effects of metapopulation

identity. We compared posterior of treatment effects on one side, and mean replicate-level

metapopulation size on the other side, between this model and the main one (in the latter

multiplying local predictions by the number of patches) to get metapopulation sizes.

Priors choices for both models combined general-purpose "weakly informative" priors based on

(McElreath 2020), with more informative priors for parameters where pre-existing knowledge

was strong (based on De Roissart, Wang, and Bonte 2015), or where constraints on model

parameters were needed (Gelman et al. 2020). We based the more informative priors (fixed

effect intercepts and random effect variance) on the empirical distribution of population

densities in (De Roissart, Wang, and Bonte 2015). For each model, we ran four chains for 4000

iterations, with 2000 iterations post-warmup per chain; this ensured both convergence (based

on graphical checks and the modified $\hat{R}$ statistic (Vehtari et al. 2020)), and that bulk and tail

effective sample sizes were satisfactory for all parameters ((Vehtari et al. 2020); at least 400,

and at least 1000 for most parameters). We relied mainly on the *tidybayes*, *bayesplot*,

patchwork packages as well as the *tidyverse* suite of packages for data handling, model post-

processing and plotting (Gabry et al. 2019; Pedersen 2019; Wickham et al. 2019; Kay and Mastny

2020). All posterior summaries given in text and figures are given as means [95% Higher

posterior Density Intervals (HDIs)].

Results

1. local and regional population sizes

From the simulation model, we expect population sizes to consistently decrease with a decreased regional (metapopulation) connectedness (Supplementary Material S04). Patches with higher local connectedness (central location) harbored consistently larger populations than less connected patches (corner location) (Supplementary Material S04).

Randomization, local and metapopulation connectedness impacted the populations sizes in the experimental metapopulations. The directions of effect were however different. For the additive effects, less connected metapopulations had higher mean population sizes (Fig. 1; predicted ratio 16 cm / 4cm = 117.2% [102.9%, 132.9%]). While evidence is weaker (95% interval overlaps 0), the direction of response is the same in response to local connectedness (Fig. 1, predicted ratio 3 links / 8 links = 113.5% [98.3%, 127.2%]). Local populations were on average smaller in reshuffled metapopulations compared to controls (Fig. 1, predicted ratio reshuffled/control = 87.0% [78.0%, 95.9%]).

Looking separately at the randomized versus non-randomized metapopulations, possible interactions became clear (Fig 2). The effect of metapopulation-level connectedness on population size was mostly apparent in the randomized metapopulations (predicted ratio 16cm /4 cm = 131.8% [106.4%, 157.8%]) but strongly reduced when looking only at control replicates (predicted ratio 16cm/4 cm = 110.1% [93.1%, 127.4%]). Conversely, the effect of the randomization treatment on population sizes was mostly detected at high levels of connectedness, but not at the lowest (randomization/ control ratios in 4 cm landscapes: 84.7%

[70.3%, 99.5%], in 8 cm landscapes: 74.6% [61.5%, 88.5%], in 16 cm landscapes: 101.1% [84.4%,

119.9%]). For a more detailed comparison between posterior distributions, please refer to the

Supplementary material (Supplementary Material S06)

Finally, the analysis of mean metapopulation sizes in a separate model shows no clear effects of

any treatment (Supplementary Material S07). However, although all effect size posteriors

overlap 0, possibly due to the much lower sample size at that scale, the directions of effect are

nonetheless the same as those implied by the patch-level model (Supplementary Material S07),

and mean replicate-level predictions are congruent with those that would be obtained by

summing local estimates from the patch-level model (Supplementary Material S07).

2. Metapopulation variability

Our simulations showed that both local (α) and regional (γ) variability are expected to increase

as a direct result of higher dispersal mortality under connectedness loss (Fig.3). Synchrony (ϕ),

by contrast, remained unaffected (Fig.3).

In our experiments, reducing metapopulation-level connectedness reduced spatial synchrony

(predicted ratio 16 cm/4cm = 85.1% [71.7%, 99.0%]) and increased additive spatial variability

(115.2% [99.7%, 131.4%]), but had no effect on α or γ variability (Fig. 4). By contrast,

randomization had a possible dampening effect on both local (α) variability (predicted ratio

reshuffled/control= 89.2% [78.3%, 99.8%]), and additive spatial variation (89.3% [79.0%,

100.3%]), but no detectable effect on synchrony or metapopulation-scale (γ) variability (Fig. 4).

Discussion

Metapopulation dynamics, as summarized by average population size, population size

variability, spatial unevenness and spatial synchrony are known to be impacted by the level of

connectedness (Wang and Loreau 2014; De Roissart, Wang, and Bonte 2015; Wang, Haegeman,

and Loreau 2015), dispersal (Jacob et al. 2019; Mortier et al. 2019) and the interaction of

dispersal with local environmental and demographic variation (Fronhofer et al. 2018). We earlier

quantified trait evolution in response to changes in connectedness (Masier and Bonte, 2020)

and now show that phenotypic sorting and spatial organization can counter the expected

decrease of population sizes loss, thereby potentially safeguarding metapopulation stability.

Dispersal in general is phenotype-dependent (Clobert et al. 2009; Bonte and Dahirel 2017),

leading to the sorting of phenotypes in metapopulations (Bonte et al. 2014; Dahirel et al. 2019).

We provide experimental evidence that such a phenotypic self-organization is an additional and

powerful driver of metapopulation dynamics. The exact underlying suites of trait changes that

drive this pattern remain, however, unknown.

Theory predicts that lower connectedness and dispersal intensify local and regional variabilities

in population sizes (Singh et al. 2004) and increase spatial asynchrony within the

metapopulation (Wang, Haegeman, and Loreau 2015). The former is also supported by the

simple individual-based model inspired by our experimental setup (Fig.3). We did, however, not

observe such a response in our experiments, and if anything, the opposite was found with

higher levels of synchrony emerging in the least connected metapopulations (Fig. 4).

Our experimental results can be reconciled with the abundant theory on the synchronizing

effects of dispersal in several ways. First, connectedness loss can in some cases drive the

evolution of higher dispersal (Cote et al. 2017) which then compensates the desynchronizing

effects of connectedness loss. Dispersal is known to often be positive-density dependent, both

in general (Harman et al. 2020) and in spider mites specifically (Bitume et al. 2013). It is worth

remembering that, as local density is not a fixed characteristic of patches, purely density-

dependent habitat choice fails to promote local evolutionary differences (Berner and Thibert-

Plante 2015). We expect a stronger local density-dependency when dispersal costs are higher

(as in our least connected treatments, Supplementary Material S04) (Travis, Murrell, and

Dytham 1999; Rodrigues and Johnstone 2014; Govindan et al. 2015). Our results here suggest

that because of dispersal variability, landscapes with low connectedness could thus counter-

intuitively lead to higher, and more homogenizing movement between patches than expected.

Second, if changes in connectedness lead to sufficiently different ecological conditions through

feedbacks, then individuals may plastically adjust their responses and eventually show higher

dispersal, but also on average higher fecundity and growth rates (Dahirel et al. 2019) that buffer

local population fluctuations. Finally, synchrony and variability may simply show opposite

patterns at long *versus* short time scales, depending on population growth responses to

perturbations (Luo et al. 2021). As such, our findings do match theoretical predictions of

dispersal (in our case connectivity) destabilizing populations and desynchronizing population

dynamics across patches at short time intervals for populations exhibiting under-compensatory

growth (i.e. slow recovery after being perturbed). At longer time intervals, these dynamics shift

towards the anticipated synchronizing and stabilizing dynamics as found in our theoretical

model. Evidently, such long-term experiments are challenging using model species with

generation times of larger magnitudes than protists and other microbial models.

Spatial heterogeneity in demographic rates may alter and even reverse population-level

predictions from mean-field theory (Holt 1985; Van Dyken and Zhang 2019; Deangelis et al.

2020). When resource conditions are homogeneous among patches, as in our experiment, such

a spatial variation in demography can only result from phenotypic structuring and self-sorting.

This spatial heterogeneity can be generated by for instance local differences in competition, or

any associated phenotypic and genotypic sorting in the network (Masier and Bonte, 2020). Our

randomization treatment, where population densities and stage structure were maintained but

not genetic (kin) and phenotypic (kind) structure (Van Petegem et al. 2018) thus provides

further insights on the relevance of phenotypic heterogeneity.

Disrupting phenotypic self-organization by randomization seems to have negative effects on

local populations, based on population sizes (Fig. 1). A closer look reveals that this effect is

actually connectedness-dependent (Fig. 2), or conversely that the effect of connectedness is

randomization-dependent: randomization has little to no effect in the least connected

metapopulations, while it does decrease population size in the others. Phenotypic self-

organization thus increases population size above the baseline expected with global random

dispersal when connectedness is high. The balance between the loss of group-living benefits

(Oku, Magalhães, and Dicke 2009; Yano 2012; Strodl and Schausberger 2013) and the spread of

favorable genotypes provides a tentative explanation. When costs are low (high connectedness),

randomization is expected to destroy favorable local structure without clear benefits in terms of

costs avoided, leading to reduced population sizes. When connectedness is low however, the

balance would be positive or neutral, as lost benefits are compensated by reduced costs of

dispersal, randomization acting as a no-costs source of global dispersal.

Besides the earlier discussed effect on population sizes, randomization also decreased local (α)

and to a lower extent regional (γ) variability (Fig. 4). This shows that (meta-)population

variability shifts in the directions anticipated from ecological theory when spatial individual

heterogeneity in metapopulation dynamics is removed (Vindenes, Engen, and Sæther 2008;

Gangloff, Sparkman, and Bronikowski 2018; Hamel et al. 2018; Smallegange, Fernandes, and

Croll 2018). Disrupting spatial phenotypic heterogeneity through randomization decreases in

consequence both population sizes and their variability across the connectedness gradient. Or,

put alternatively, the maintenance of this phenotypic organization leads to more (asynchronous)

fluctuating metapopulations of larger size, conditions known to rescue metapopulations at

intermediate levels (Wang and Loreau 2014; Wang, Haegeman, and Loreau 2015), but that can

also potentially lead the metapopulations to be more vulnerable towards local extinctions if said

fluctuations become too pronounced or erratic (Abbott 2011; Zhang, DeAngelis, and Ni 2021).

Spatial synchrony ($\square$) remains unaffected by the randomization treatment. In conjunction with

the effects on spatial variability (additive β), we propose the following explanation for this

paradoxical result. Spatial synchrony $\square$ *sensu* Wang & Loreau (2014) represents roughly how the

regional variability γ is partitioned between local variability α and inter-patches spatial

variability β (see also Table 1). As such, if local variability and spatial variability vary with similar

dynamics, the resulting spatial synchrony will remain unchanged. This is visible in of simulation

model (fig.3): local variability (α) and spatial variability (β) have an almost-identical shape,

resulting in a constant spatial synchrony (φ). In the same regard, the effect of randomization on

spatial variability is similar to the one for local variability (fig.4). To summarize, when phenotypic

self-sorting is removed from a (meta-)population, local population sizes decrease, while the

spatial synchronicity (φ) remains unaffected because both local variability (α) and spatial

variability (additive β) change in the same direction. Self-sorting thus increases both local

variability and spatial variability. They respectively impose a destabilizing and stabilizing effect,

with the net balance seeming overall positive (at least in terms of population sizes). Further

studies are recommended to confirm or falsify our finding, but also to test the consistency of

these dynamics on longer time periods, or under disturbances (Luo et al. 2021).

Our experiments additionally show how differences in connectedness within the spatial network

affect local population sizes. As the central patches showed the highest level of connectedness

in all metapopulations, increased population sizes due to positive immigration-emigration

balances were expected (Masier and Bonte, 2020). Surprisingly however, opposite patterns

were found with central patches systematically showing lower densities. This suggests that local

demographic changes are not captured by the metapopulation-level rates of dispersal. As we

did not record the total number of individuals, but only the adult females, it is possible that

juveniles development was slowed down in central patches due to resource limitation, which

was possibly caused by a large influx of immigrants that since then died or dispersed again. It is

known that *T. urticae* exhibits longer development times when faced with suboptimal food

sources, while longevity and egg-laying performances are not affected (Aucejo-Romero, Gómez-

Cadenas, and Jacas-Miret 2004). Likely such delays caused a reduced number of adults during

the counts of central patches relative to the others. Highly-connected patches are commonly

considered as the most resilient in a network precisely due to their high number of connections,

and thus the most valuable and cost-effective populations for conservation (Calabrese and

Fagan 2004; Gonzalez, Thompson, and Loreau 2017; Liao, Shen, and Liao 2020). Our experiment

shows how this is not necessarily the case.

The implemented randomization treatment can be considered as an extreme case of

hypermobility within the spatial network. Our experiment thus pinpoints the applied

consequences of hypermobility, for instance through human-mediated dispersal (McRae et al.

2012; Bullock et al. 2018; Rudnick et al. 2012). Apart from the potential to impose

maladaptation, or to hinder adaptation (Brady, Bolnick, Barrett, et al. 2019; Brady, Bolnick,

Angert, et al. 2019), hypermobility can lead to changes in the local and regional population

dynamics. While phenotypic variation has been shown to explain patterns in biodiversity

(Bolnick et al. 2011; Dahirel et al. 2019), we here demonstrate its importance for

metapopulation demography, and in extension metapopulation conservation. Removing spatial

phenotypic organization by for instance assisted migration needs to be done with care, as it

could have so far unforeseen detrimental effects for the persistence of the metapopulation.

References

Abbott, Karen C. 2011. "A Dispersal-Induced Paradox: Synchrony and Stability in Stochastic Metapopulations." *Ecology Letters* 14 (11): 1158–69. https://doi.org/10.1111/j.1461-0248.2011.01670.x.

Aucejo-Romero, Sílvia, Aurelio Gómez-Cadenas, and Josep Anton Jacas-Miret. 2004. "Effects of NaCl-Stressed Citrus Plants on Life-History Parameters of Tetranychus Urticae (Acari: Tetranychidae)." *Experimental and Applied Acarology* 33 (1–2): 55–67. https://doi.org/10.1023/B:APPA.0000030026.77800.0c.

Berner, D., and X. Thibert-Plante. 2015. "How Mechanisms of Habitat Preference Evolve and Promote Divergence with Gene Flow." *Journal of Evolutionary Biology* 28 (9): 1641–55. https://doi.org/10.1111/jeb.12683.

Bitume, E. V., D. Bonte, O. Ronce, F. Bach, E. Flaven, I. Olivieri, and C. M. Nieberding. 2013. "Density and Genetic Relatedness Increase Dispersal Distance in a Subsocial Organism." Edited by John Fryxell. *Ecology Letters* 16 (4): 430–37. https://doi.org/10.1111/ele.12057.

Bolnick, Daniel I., Priyanga Amarasekare, Márcio S. Araújo, Reinhard Bürger, Jonathan M. Levine, Mark Novak, Volker H.W. Rudolf, Sebastian J. Schreiber, Mark C. Urban, and David A. Vasseur. 2011. "Why Intraspecific Trait Variation Matters in Community Ecology." *Trends in Ecology and Evolution* 26 (4): 183–92. https://doi.org/10.1016/j.tree.2011.01.009.

Bonte, D., A. De Roissart, N. Wybouw, and T. Van Leeuwen. 2014. "Fitness Maximization by Dispersal: Evidence from an Invasion Experiment." *Ecology* 95 (11): 3104–11. https://doi.org/10.1890/13-2269.1.

Bonte, Dries, and Quinten Bafort. 2019. "The Importance and Adaptive Value of Life-History Evolution for Metapopulation Dynamics." *Journal of Animal Ecology* 88 (1): 24–34. https://doi.org/10.1111/1365-2656.12928.

Bonte, Dries, and Maxime Dahirel. 2017. "Dispersal: A Central and Independent Trait in Life History." *Oikos* 126 (4): 472–79. https://doi.org/10.1111/oik.03801.

Bonte, Dries, Hans Van Dyck, James M. Bullock, Aurélie Coulon, Maria Delgado, Melanie Gibbs, Valerie Lehouck, et al. 2012. "Costs of Dispersal." *Biological Reviews* 87 (2): 290–312. https://doi.org/10.1111/j.1469-185X.2011.00201.x.

Bonte, Dries, Stefano Masier, and Frederik Mortier. 2018. "Eco-Evolutionary Feedbacks Following Changes in Spatial Connectedness." *Current Opinion in Insect Science* 29 (October): 64–70. https://doi.org/10.1016/j.cois.2018.06.003.

Brady, Steven P., Daniel I. Bolnick, Amy L. Angert, Andrew Gonzalez, Rowan D.H. Barrett, Erika Crispo, Alison M. Derry, et al. 2019. "Causes of Maladaptation." *Evolutionary Applications* 12 (7): 1229–42. https://doi.org/10.1111/eva.12844.

Brady, Steven P., Daniel I. Bolnick, Rowan D.H. Barrett, Lauren Chapman, Erika Crispo, Alison M. Derry, Christopher G. Eckert, et al. 2019. "Understanding Maladaptation by Uniting Ecological and Evolutionary Perspectives." *American Naturalist* 194 (4): 495–515. https://doi.org/10.1086/705020.

Bullock, James M., Dries Bonte, Gesine Pufal, Carolina da Silva Carvalho, Daniel S. Chapman, Cristina García, Daniel García, Erik Matthysen, and Maria Mar Delgado. 2018. "Human-Mediated Dispersal and the Rewiring of Spatial Networks." *Trends in Ecology & Evolution* 33 (12): 958–70. https://doi.org/10.1016/j.tree.2018.09.008.

Bürkner, Paul Christian. 2018. "Advanced Bayesian Multilevel Modeling with the R Package Brms." *R Journal* 10 (1): 395–411. https://doi.org/10.32614/rj-2018-017.

Calabrese, Justin M, and William F Fagan. 2004. "A Comparison-Shopper 's Guide to Connectivity Metrics." *Frontiers in Ecology and the Environment* 2 (10): 529–36. https://doi.org/10.1890/1540-9295(2004)002[0529:ACGTCM]2.0.CO;2.

Carpenter, Bob, Andrew Gelman, Matthew D. Hoffman, Daniel Lee, Ben Goodrich, Michael Betancourt, Marcus A. Brubaker, Jiqiang Guo, Peter Li, and Allen Riddell. 2017. "Stan: A Probabilistic Programming Language." *Journal of Statistical Software* 76 (1). https://doi.org/10.18637/jss.v076.i01.

Cazaux, Marc, Marie Navarro, Kristie A. Bruinsma, Vladimir Zhurov, Tara Negrave, Thomas Van

525   Leeuwen, Vojislava Grbic, and Miodrag Grbic. 2014. "Application of Two-Spotted Spider
526   Mite <em>Tetranychus Urticae</Em> for Plant-Pest Interaction Studies." *Journal of*
527   *Visualized Experiments*, no. 89 (July): 1–7. https://doi.org/10.3791/51738.
528   Clobert, Jean, Michel Baguette, Tim G. Benton, and James M. Bullock. 2012. *Dispersal Ecology*
529   *and Evolution.*
530   Clobert, Jean, Jean François Le Galliard, Julien Cote, Sandrine Meylan, and Manuel Massot.
531   2009. "Informed Dispersal, Heterogeneity in Animal Dispersal Syndromes and the Dynamics
532   of Spatially Structured Populations." *Ecology Letters* 12 (3): 197–209.
533   https://doi.org/10.1111/j.1461-0248.2008.01267.x.
534   Cote, Julien, Elvire Bestion, Staffan Jacob, Justin Travis, Delphine Legrand, and Michel Baguette.
535   2017. "Evolution of Dispersal Strategies and Dispersal Syndromes in Fragmented
536   Landscapes." *Ecography* 40: 56–73. https://doi.org/doi: 10.1111/ecog.02538.
537   Crooks, Kevin R. 2002. "Relative Sensitivities of Mammalian Carnivores to Habitat
538   Fragmentation." *Conservation Biology* 16 (2): 488–502. https://doi.org/10.1046/j.1523-
539   1739.2002.00386.x.
540   Cushman, Samuel A. 2006. "Effects of Habitat Loss and Fragmentation on Amphibians: A Review
541   and Prospectus." *Biological Conservation* 128 (2): 231–40.
542   https://doi.org/10.1016/j.biocon.2005.09.031.
543   Cushman, Samuel A., Kevin McGarigal, and Maile C. Neel. 2008. "Parsimony in Landscape
544   Metrics: Strength, Universality, and Consistency." *Ecological Indicators* 8 (5): 691–703.
545   https://doi.org/10.1016/j.ecolind.2007.12.002.
546   Dahirel, Maxime, Stefano Masier, David Renault, and Dries Bonte. 2019. "The Distinct
547   Phenotypic Signatures of Dispersal and Stress in an Arthropod Model: From Physiology to
548   Life History." *Journal of Experimental Biology* 222 (August): jeb.203596.
549   https://doi.org/10.1242/jeb.203596.
550   Deangelis, D. L., Bo Zhang, Wei Ming Ni, and Yuanshi Wang. 2020. "Carrying Capacity of a
551   Population Diffusing in a Heterogeneous Environment." *Mathematics* 8 (1).
552   https://doi.org/10.3390/math8010049.
553   Dochtermann, Ned A., and Raphaël Royauté. 2019. "The Mean Matters: Going beyond
554   Repeatability to Interpret Behavioural Variation." *Animal Behaviour* 153: 147–50.
555   https://doi.org/10.1016/j.anbehav.2019.05.012.
556   Dyken, J. David Van, and Bo Zhang. 2019. "Carrying Capacity of a Spatially-Structured
557   Population: Disentangling the Effects of Dispersal, Growth Parameters, Habitat
558   Heterogeneity and Habitat Clustering." *Journal of Theoretical Biology* 460: 115–24.
559   https://doi.org/10.1016/j.jtbi.2018.09.015.
560   Fahrig, Lenore. 2017. "Ecological Responses to Habitat Fragmentation Per Se." *Annual Review*
561   *of Ecology, Evolution, and Systematics* 48 (1): 1–23. https://doi.org/10.1146/annurev-
562   ecolsys-110316-022612.
563   Fronhofer, Emanuel A., Delphine Legrand, Florian Altermatt, Armelle Ansart, Simon Blanchet,
564   Dries Bonte, Alexis Chaine, et al. 2018. "Bottom-up and Top-down Control of Dispersal
565   across Major Organismal Groups." *Nature Ecology & Evolution* 2 (12): 1859–63.
566   https://doi.org/10.1038/s41559-018-0686-0.
567   Fronhofer, Emanuel A., Hans Joachim Poethke, and Ulf Dieckmann. 2015. "Evolution of
568   Dispersal Distance: Maternal Investment Leads to Bimodal Dispersal Kernels." *Journal of*
569   *Theoretical Biology* 365: 270–79. https://doi.org/10.1016/j.jtbi.2014.10.024.
570   Fuller, Matthew R., Martin W. Doyle, and David L. Strayer. 2015. "Causes and Consequences of
571   Habitat Fragmentation in River Networks." *Annals of the New York Academy of Sciences*
572   1355 (1): 31–51. https://doi.org/10.1111/nyas.12853.
573   Gabry, Jonah, Daniel Simpson, Aki Vehtari, Michael Betancourt, and Andrew Gelman. 2019.
574   "Visualization in Bayesian Workflow." *Journal of the Royal Statistical Society. Series A:*
575   *Statistics in Society* 182 (2): 389–402. https://doi.org/10.1111/rssa.12378.
576   Galliard, J. F. Le, M. Paquet, and M. Mugabo. 2015. "An Experimental Test of Density-Dependent
577   Selection on Temperament Traits of Activity, Boldness and Sociability." *Journal of*
578   *Evolutionary Biology* 28 (5): 1144–55. https://doi.org/10.1111/jeb.12641.

Gandon, Sylvain. 1999. "Kin Competition, the Cost of Inbreeding and the Evolution of Dispersal."
Journal of Theoretical Biology 200 (4): 345–64. https://doi.org/10.1006/jtbi.1999.0994.
Gangloff, Eric J., Amanda M. Sparkman, and Anne M. Bronikowski. 2018. "Among-Individual
Heterogeneity in Maternal Behaviour and Physiology Affects Reproductive Allocation and
Offspring Life-History Traits in the Garter Snake Thamnophis Elegans." *Oikos* 127 (5): 612–
25.
Gelman, Andrew, Aki Vehtari, Daniel Simpson, Charles C. Margossian, Bob Carpenter, Yuling
Yao, Lauren Kennedy, Jonah Gabry, Paul-Christian Bürkner, and Martin Modrák. 2020.
"Bayesian Workflow," November. http://arxiv.org/abs/2011.01808.
Gonzalez, Andrew, Patrick Thompson, and Michel Loreau. 2017. "Spatial Ecological Networks:
Planning for Sustainability in the Long-Term." *Current Opinion in Environmental*
Sustainability 29 (December): 187–97. https://doi.org/10.1016/j.cosust.2018.03.012.
Goossens, Steven, Nicky Wybouw, Thomas Van Leeuwen, and Dries Bonte. 2020. "The
Physiology of Movement." *Movement Ecology* 8 (5).
https://doi.org/https://doi.org/10.1186/s40462-020-0192-2.
Gouhier, Tarik C., Frédéric Guichard, and Andrew Gonzalez. 2010. "Synchrony and Stability of
Food Webs in Metacommunities." *The American Naturalist* 175 (2): E16–34.
https://doi.org/10.1086/649579.
Govaert, Lynn, Emanuel A. Fronhofer, Sébastien Lion, Christophe Eizaguirre, Dries Bonte,
Martijn Egas, Andrew P. Hendry, et al. 2019. "Eco-Evolutionary Feedbacks—Theoretical
Models and Perspectives." *Functional Ecology* 33 (1): 13–30. https://doi.org/10.1111/1365-
2435.13241.
Govindan, Byju N., Zhilan Feng, Yssa D. DeWoody, and Robert K. Swihart. 2015. "Intermediate
Disturbance in Experimental Landscapes Improves Persistence of Beetle Metapopulations."
Ecology 96 (3): 728–36. https://doi.org/10.1890/14-0044.1.
Hamel, S., J.-M. Gaillard, M. Douhard, M. Festa-Bianchet, F. Pelletier, and N.G. Yoccoz. 2018.
"Quantifying Individual Heterogeneity and Its Influence on Life-History Trajectories: Different
Methods for Different Questions and Contexts." *Oikos* 127 (5).
Hance, Th, and G. Van Impe. 1999. "The Influence of Initial Age Structure on Predator – Prey
Interaction." *Ecological Modelling* 114 (2–3): 195–211. https://doi.org/10.1016/S0304-
3800(98)00148-3.
Harman, Rachel R., Jerome Goddard, Ratnasingham Shivaji, and James T. Cronin. 2020.
"Frequency of Occurrence and Population-Dynamic Consequences of Different Forms of
Density-Dependent Emigration." *The American Naturalist* 195 (5): 851–67.
https://doi.org/10.1086/708156.
Harrison, Xavier A. 2014. "Using Observation-Level Randomeffects to Model Overdispersion in
Count Data in Ecology and Evolution." *PeerJ* 2014 (1). https://doi.org/10.7717/peerj.616.
Holt, Robert D. 1985. "Population Dynamics in Two-Patch Environments: Some Anomalous
Consequences of an Optimal Habitat Distribution." *Theoretical Population Biology* 28 (2):
181–208. https://doi.org/10.1016/0040-5809(85)90027-9.
Hudson, P. J., and I. M. Cattadori. 1999. "The Moran Effect: A Cause of Population Synchrony."
Trends in Ecology and Evolution 14 (1): 1–2. https://doi.org/10.1016/S0169-5347(98)01498-
0.
Ims, Rolf A., and Harry P. Andreassen. 1999. "Effects of Experimental Habitat Fragmentation and
Connectivity on Root Vole Demography." *Journal of Animal Ecology* 68 (5): 839–52.
https://doi.org/10.1046/j.1365-2656.1999.00336.x.
Ives, Anthony R., Scott T. Woody, Erik V. Nordheim, Chris Nelson, and John H. Andrews. 2004.
"The Synergistic Effects of Stochasticity and Dispersal on Population Densities." *The*
American Naturalist 163 (3): 375–87. https://doi.org/10.1086/381942.
Jacob, Staffan, Alexis S. Chaine, Michèle Huet, Jean Clobert, and Delphine Legrand. 2019.
"Variability in Dispersal Syndromes Is a Key Driver of Metapopulation Dynamics in
Experimental Microcosms." *The American Naturalist* 194 (5): 613–26.
https://doi.org/10.1086/705410.
Kamioka, Takahiro, and Yoh Iwasa. 2017. "Evolution of Density-Dependent Wing Polymorphism

in Insects." *Evolutionary Ecology Research* 18 (1): 335–48.

Karisto, Petteri, and Éva Kisdi. 2019. "Joint Evolution of Dispersal and Connectivity." *Evolution* 73 (12): 2529–37. https://doi.org/10.1111/evo.13866.

Kawecki, Tadeusz J., Richard E. Lenski, Dieter Ebert, Brian Hollis, Isabelle Olivieri, and Michael C. Whitlock. 2012. "Experimental Evolution." *Trends in Ecology and Evolution* 27 (10): 547–60. https://doi.org/10.1016/j.tree.2012.06.001.

Kay, M, and T Mastny. 2020. "Tidybayes: Tidy Data and Geoms for Bayesian Models, 2020." *R Package Version* 2 (1.9000).

Koeller, P., R. Mohn, and M. Etter. 2000. "Density Dependant Sex Change in Northern Shrimp, Pandalus Borealis, on the Scotian Shelf." *Journal of Northwest Atlantic Fishery Science* 27 (1978): 107–18. https://doi.org/10.2960/J.v27.a10.

Leeuwen, Thomas Van, Bartel Vanholme, Steven Van Pottelberge, Pieter Van Nieuwenhuyse, Ralf Nauen, Luc Tirry, and Ian Denholm. 2008. "Mitochondrial Heteroplasmy and the Evolution of Insecticide Resistance: Non-Mendelian Inheritance in Action." *Proceedings of the National Academy of Sciences of the United States of America* 105 (16): 5980–85. https://doi.org/10.1073/pnas.0802224105.

Liao, Limei, Yang Shen, and Jinbao Liao. 2020. "Robustness of Dispersal Network Structure to Patch Loss." *Ecological Modelling* 424 (December 2019): 109036. https://doi.org/10.1016/j.ecolmodel.2020.109036.

Lion, Sébastien. 2018. "Theoretical Approaches in Evolutionary Ecology: Environmental Feedback as a Unifying Perspective." *American Naturalist* 191 (1): 21–44. https://doi.org/10.1086/694865.

Lloyd, Alun L., and R. M. May. 1999. "Synchronicity , Chaos and Population Cycles :" *Trends in Ecology & Evolution* 14 (11): 417–18.

Lowe, Winsor H., Clint C. Muhlfeld, and Fred W. Allendorf. 2015. "Spatial Sorting Promotes the Spread of Maladaptive Hybridization." *Trends in Ecology and Evolution* 30 (8): 456–62. https://doi.org/10.1016/j.tree.2015.05.008.

Luo, Mingyu, Daniel C. Reuman, Lauren M. Hallett, Lauren Shoemaker, Lei Zhao, Max C.N. Castorani, Joan C. Dudney, et al. 2021. "The Effects of Dispersal on Spatial Synchrony in Metapopulations Differ by Timescale." *Oikos*, no. July: 1–11. https://doi.org/10.1111/oik.08298.

Masier, Stefano, and Dries Bonte. 2020. "Spatial Connectedness Imposes Local- and Metapopulation-level Selection on Life History through Feedbacks on Demography." Edited by Tim Coulson. *Ecology Letters* 23 (2): 242–53. https://doi.org/10.1111/ele.13421.

McElreath, Richard. 2020. *Statistical Rethinking: A Bayesian Course with Examples in R and Stan*. Second. Chapman and Hall/CRC press.

McGuire, Jenny L., Joshua J. Lawler, Brad H. McRae, Tristan A. Nuñez, and David M. Theobald. 2016. "Achieving Climate Connectivity in a Fragmented Landscape." *Proceedings of the National Academy of Sciences* 113 (26): 7195–7200. https://doi.org/10.1166/jctn.2009.1107.

McRae, Brad H., Sonia A. Hall, Paul Beier, and David M. Theobald. 2012. "Where to Restore Ecological Connectivity? Detecting Barriers and Quantifying Restoration Benefits." *PLoS ONE* 7 (12). https://doi.org/10.1371/journal.pone.0052604.

Migeon, Alain, Elodie Nouguier, and Franck Dorkeld. 2010. "Spider Mites Web: A Comprehensive Database for the Tetranychidae." In *Trends in Acarology*, edited by Maurice W. Sabelis and Jan Bruin, 557–60. Dordrecht: Springer Netherlands. https://doi.org/10.1007/978-90-481-9837-5_96.

Mortier, Frederik, Staffan Jacob, Martijn L. Vandegehuchte, and Dries Bonte. 2019. "Habitat Choice Stabilizes Metapopulation Dynamics through Increased Ecological Specialisation." *Oikos* 128 (4): 529–39. https://doi.org/10.1101/267575.

Oku, Keiko, Sara Magalhães, and Marcel Dicke. 2009. "The Presence of Webbing Affects the Oviposition Rate of Two-Spotted Spider Mites, Tetranychus Urticae (Acari: Tetranychidae)." *Experimental and Applied Acarology* 49 (3): 167–72. https://doi.org/10.1007/s10493-009-9252-4.

Parrott, Lael. 2010. "Measuring Ecological Complexity." *Ecological Indicators* 10 (6): 1069–76.

https://doi.org/10.1016/j.ecolind.2010.03.014.

Parsons, Kevin J., H. D. Sheets, S. Skúlason, and M. M. Ferguson. 2011. "Phenotypic Plasticity,
Heterochrony and Ontogenetic Repatterning during Juvenile Development of Divergent
Arctic Charr (Salvelinus Alpinus)." *Journal of Evolutionary Biology* 24 (8): 1640–52.
https://doi.org/10.1111/j.1420-9101.2011.02301.x.

Pedersen, Thomas Lin. 2019. "Patchwork: The Composer of Plots." *R Package Version* 1 (0).

Perrin, Nicolas, and Vladimir Mazalov. 2000. "Local Competition, Inbreeding, and the Evolution of
Sex‐Biased Dispersal." *The American Naturalist* 155 (1): 116–27.
https://doi.org/10.1086/303296.

Petegem, Katrien Hilde Petra Van, Felix Moerman, Maxime Dahirel, Emanuel A. Fronhofer,
Martijn L. Vandegehuchte, Thomas Van Leeuwen, Nicky Wybouw, Robby Stoks, and Dries
Bonte. 2018. "Kin Competition Accelerates Experimental Range Expansion in an Arthropod
Herbivore." *Ecology Letters* 21 (2): 225–34. https://doi.org/10.1111/ele.12887.

Poethke, H.J., B. Pfenning, and T. Hovestadt. 2007. "The Relative Contribution of Individual and
Kin Selection to the Evolution of Density-Dependent Dispersal Rates." *Evolutionary Ecology*
Research 9 (1): 41–50.

Rodrigues, António M. M., and Rufus A. Johnstone. 2014. "Evolution of Positive and Negative
Density-Dependent Dispersal." *Proceedings of the Royal Society B: Biological Sciences* 281
(1791): 20141226. https://doi.org/10.1098/rspb.2014.1226.

Roissart, Annelies De, Shaopeng Wang, and Dries Bonte. 2015. "Spatial and Spatiotemporal
Variation in Metapopulation Structure Affects Population Dynamics in a Passively
Dispersing Arthropod." *Journal of Animal Ecology* 84 (6): 1565–74.
https://doi.org/10.1111/1365-2656.12400.

Rudnick, Deborah A., Michael G Ryan, Mark E Harmon, Richard a Birdsey, Christian P Giardina,
Linda S Heath, Richard a Houghton, et al. 2012. "The Role of Landscape Connectivity in
Planning and Implementating Conservation and Restoration Priorities." *Issue in Ecology* 13
(16): 1–16.

Sabelis, Maurice W., Arne Janssen, Odo Diekmann, Vincent A.A. Jansen, Erik van Gool, and
Minus van Baalen. 2005. "Global Persistence Despite Local Extinction in Acarine Predator-
Prey Systems: Lessons From Experimental and Mathematical Exercises." *Advances in*
Ecological Research 37 (04): 183–220. https://doi.org/10.1016/S0065-2504(04)37006-6.

Schielzeth, Holger. 2010. "Simple Means to Improve the Interpretability of Regression
Coefficients." *Methods in Ecology and Evolution* 1 (2): 103–13.
https://doi.org/10.1111/j.2041-210x.2010.00012.x.

Singh, Brajendra K., J. Subba Rao, R. Ramaswamy, and Somdatta Sinha. 2004. "The Role of
Heterogeneity on the Spatiotemporal Dynamics of Host-Parasite Metapopulation."
Ecological Modelling 180 (2–3): 435–43. https://doi.org/10.1016/j.ecolmodel.2004.04.031.

Smallegange, Isabel M., Rianne E. Fernandes, and Jasper C. Croll. 2018. "Population
Consequences of Individual Heterogeneity in Life Histories: Overcompensation in Response
to Harvesting of Alternative Reproductive Tactics." *Oikos* 127 (5): 738–49.
https://doi.org/10.1111/oik.04130.

Strodl, Markus A., and Peter Schausberger. 2013. "Social Familiarity Relaxes the Constraints of
Limited Attention and Enhances Reproduction of Group-Living Predatory Mites." *Oikos* 122
(8): 1217–26. https://doi.org/10.1111/j.1600-0706.2012.20833.x.

Team R Core. 2021. "R: A Language and Environment for Statistical Computing (4.0.5)."

Thompson, Patrick L., Bronwyn Rayfield, and Andrew Gonzalez. 2017. "Loss of Habitat and
Connectivity Erodes Species Diversity, Ecosystem Functioning, and Stability in
Metacommunity Networks." *Ecography* 40 (1): 98–108. https://doi.org/10.1111/ecog.02558.

Tischendorf, Lutz, and Lenore Fahrig. 2000. "On the Usage and Measurement of Landscape
Connectivity." *Oikos* 90 (1): 7–19. https://doi.org/10.1034/j.1600-0706.2000.900102.x.

Travis, Justin M J, David J Murrell, and Calvin Dytham. 1999. "The Evolution of Density–
Dependent Dispersal." *Proceedings of the Royal Society of London. Series B: Biological*
Sciences 266 (1431): 1837–42. https://doi.org/10.1098/rspb.1999.0854.

Turner, M G. 1989. "Landscape Ecology: The Effect of Pattern on Process." *Annual Review of*

Ecology and Systematics 20 (1): 171–97.
https://doi.org/10.1146/annurev.es.20.110189.001131.
Urban, Mark C., Sharon Y. Strauss, Fanie Pelletier, Eric P. Palkovacs, Mathew A. Leibold,
Andrew P. Hendry, Luc de Meester, Stephanie M. Carlson, Amy L. Angert, and Sean T.
Giery. 2020. "Evolutionary Origins for Ecological Patterns in Space." *Proceedings of the*
National Academy of Sciences of the United States of America 117 (30): 17482–90.
https://doi.org/10.1073/pnas.1918960117.
Vehtari, Aki, Andrew Gelman, Daniel Simpson, Bob Carpenter, and Paul-Christian Bürkner. 2020.
"Rank-Normalization, Folding, and Localization: An Improved $\widehat{R}$ for Assessing
Convergence of MCMC." *Bayesian Analysis*, 1–26. https://doi.org/10.1214/20-ba1221.
Villemereuil, Pierre de, Holger Schielzeth, Shinichi Nakagawa, and Michael Morrissey. 2016.
"General Methods for Evolutionary Quantitative Genetic Inference from Generalized Mixed
Models." *Genetics* 204 (3): 1281–94. https://doi.org/10.1534/genetics.115.186536.
Vindenes, Yngvild, Steinar Engen, and Bernt Erik Sæther. 2008. "Individual Heterogeneity in Vital
Demographic Stochasticity." *American Naturalist* 171 (4): 455–67.
https://doi.org/10.1086/528965.
Wang, Shaopeng, Bart Haegeman, and Michel Loreau. 2015. "Dispersal and Metapopulation
Stability." *PeerJ* 3 (10): 1–16. https://doi.org/10.7717/peerj.1295.
Wang, Shaopeng, and Michel Loreau. 2014. "Ecosystem Stability in Space: α, β and γ
Variability." Edited by Andrew Liebhold. *Ecology Letters* 17 (8): 891–901.
https://doi.org/10.1111/ele.12292.
Wang, Xianli, F. Guillaume Blanchet, and Nicola Koper. 2014. "Measuring Habitat Fragmentation:
An Evaluation of Landscape Pattern Metrics." *Methods in Ecology and Evolution* 5 (7): 634–
46. https://doi.org/10.1111/2041-210X.12198.
Wickham, Hadley, Mara Averick, Jennifer Bryan, Winston Chang, Lucy McGowan, Romain
François, Garrett Grolemund, et al. 2019. "Welcome to the Tidyverse." *Journal of Open*
Source Software 4 (43): 1686. https://doi.org/10.21105/joss.01686.
Woodworth, Bradley K., Nathaniel T. Wheelwright, Amy E.M. Newman, and D. Ryan Norris. 2017.
"Local Density Regulates Migratory Songbird Reproductive Success through Effects on
Double-Brooding and Nest Predation." *Ecology* 98 (8): 2039–48.
https://doi.org/10.1002/ecy.1911.
Yano, Shuichi. 2012. "Cooperative Web Sharing against Predators Promotes Group Living in
Spider Mites." *Behavioral Ecology and Sociobiology* 66 (6): 845–53.
https://doi.org/10.1007/s00265-012-1332-5.
Zhang, Bo, Donald L. DeAngelis, and Wei-Ming Ni. 2021. "Carrying Capacity of Spatially
Distributed Metapopulations." *Trends in Ecology & Evolution* 36 (2): 164–73.
https://doi.org/10.1016/j.tree.2020.10.007.
778
779

 Figures legends

Figure 1. Posterior additive effects of treatments on mean local population size. For each

treatment variable, posteriors (with mean and 95% CI in black) are displayed after averaging

over the other two. Grey dots are empirical patch-level means, transparency of the dots is

proportional to relative variability, with more precise estimates being more opaque (using the

inverse of the mean-scaled variance as a measure of precision; see e.g. (Dochtermann and

Royauté 2019)).

Figure 2. Posterior effect of metapopulation connectedness on mean local population size as a

function of randomization. For each connectedness level, posteriors (with mean and 95% CI in

black) are displayed after averaging over the other one. Grey dots are empirical patch-level

means, transparency of the dots is proportional to relative variability, with more precise

estimates being more opaque (using the inverse of the mean-scaled variance as a measure of

precision).

Figure 3. Landscape variability metrics calculated from our simulation model (no evolution

included, dispersal mortality used as a proxy for connectedness): variability is expected to

increase with lower connectedness, both at local (first panel) and metapopulation level (fourth

panel); spatial variability is expected to increase similarly (second panel). Conversely, spatial

synchrony is expected to be largely unaffected (third panel).

Figure 4. Effect of landscape-level variables of measures of landscape variability based on (Wang

and Loreau, 2014). For each treatment variable, posteriors (with mean and 95% CI in black) are

displayed after averaging over the other two. Grey "shadows" in the panels showing the effect

of connectedness denote the posteriors that would be obtained using only control, non-

randomized replicates.

Description	Notation
Temporal variance of local population size in patch i [a]	w_{ii}
Temporal mean local population size of patch i [a,b]	μ_i
Temporal (co)variance between patches i and j from the same metapopulation [a]	w_{ij}
Temporal mean metapopulation size [c]	$\mu_M = \sum_i \mu_i / m$, where m is the number of patches
Temporal variability at the local patch scale α: weighted average of patch-level coefficients of temporal variation	$\alpha = \left(\dfrac{\sum_i \sqrt{w_{ii}}}{\mu_M} \right)^2$
Temporal variability at the metapopulation scale γ	$\gamma = \left(\dfrac{\sum_{i,j} \sqrt{w_{ij}}}{\mu_M} \right)^2$
Additive spatial variability β_2	$\beta_2 = \alpha - \gamma$
Spatial synchrony $\varphi = 1/\beta_1$ [d]	$\varphi = \gamma/\alpha$

Table 1: Summary of the key variables estimated from the multivariate Poisson model of local
population sizes, based on Wang and Loreau (2014) (See Supplementary material S05 for how
each metric relates to combination of fixed and random effects).

[a] see Supplementary Material S05 and de Villemereuil et al. 2016 for how to obtain this parameter on the data scale
from latent-scale mixed model coefficients

[b] It is important to note that in a Poisson model with observation-level random effects, differences in patch-level
intercepts reflect differences in (log-transformed) medians; if present, as in here, differences in within-patch
variances should be accounted for to estimate differences in means (see de Villemereuil et al. 2016)

[c] Can also be estimated from a model with metapopulation size as a response (see Supplementary Material S07 for a
comparison)

[d] Spatial synchrony among patches from the same populations; connects local temporal variability and additive spatial
variability to metapopulation temporal variability; $\alpha \times \varphi = \gamma$, $\alpha \times (1 - \varphi) = \beta_2$.