Genomically Complex Human Angiosarcoma and Canine Hemangiosarcoma Establish

Convergent Angiogenic Transcriptional Programs Driven by Novel Gene Fusions

Jong Hyuk Kim[1,2,3,4*], Kate Megquier[5], Rachael Thomas[6], Aaron L. Sarver[1,3,7], Jung Min Song[3],

Yoon Tae Kim[8], Nuojin Cheng[9,a], Ashley J. Schulte[1,2,3], Michael A. Linden[1,3,10], Paari

Murugan[1,3,10], LeAnn Oseth[3], Colleen L. Forster[11], Ingegerd Elvers[5,12], Ross Swofford[5], Jason

Turner-Maier[5], Elinor K. Karlsson[5,13], Matthew Breen[6,14], Kerstin Lindblad-Toh[5,12], Jaime F.

Modiano[1,2,3,4,10,15,16]

[1]Animal Cancer Care and Research Program, University of Minnesota, St Paul, MN, USA

[2]Department of Veterinary Clinical Sciences, College of Veterinary Medicine, University of

Minnesota, St Paul, MN, USA

[3]Masonic Cancer Center, University of Minnesota, Minneapolis, MN, USA

[4]Institute for Engineering in Medicine, University of Minnesota, Minneapolis, MN, USA

[5]Broad Institute of Harvard and MIT, Cambridge, MA, USA

[6]Department of Molecular Biomedical Sciences, College of Veterinary Medicine & Comparative

Medicine Institute, North Carolina State University, Raleigh, NC, USA

[7]Institute for Health Informatics, University of Minnesota, Minneapolis, MN, USA

[8]Department of Electrical Engineering and Computer Science, York University, Toronto,

Ontario, Canada

[9]School of Mathematics, College of Science and Engineering, University of Minnesota,

Minneapolis, MN, USA

[10]Department of Laboratory Medicine and Pathology, School of Medicine, University of

Minnesota, Minneapolis, MN, USA

[11]The University of Minnesota Biological Materials Procurement Network (BioNet), University

of Minnesota, Minneapolis, MN, USA

[12]Science for Life Laboratory, Department of Medical Biochemistry and Microbiology, Uppsala

University, Uppsala, Sweden

[13]University of Massachusetts Medical School, Worcester, MA, USA

[14]Cancer Genetics Program, University of North Carolina Lineberger Comprehensive Cancer

Center, Raleigh, NC, USA

[15]Center for Immunology, University of Minnesota, Minneapolis, MN, USA

[16]Stem Cell Institute, University of Minnesota, Minneapolis, MN, USA

[a]Current address: Applied Mathematics, University of Colorado Boulder, Boulder, CO, USA

Running Title: Angiogenic transcription programs in angiosarcoma

Keywords: Angiosarcoma, chromosome translocation, fusion gene, hemangiosarcoma, *TP53*

Corresponding Author: Jong Hyuk Kim (jhkim@umn.edu), MCRB560A, Masonic Cancer

Center, 420 Delaware St. SE, Minneapolis, MN 55455. Phone: 612-624-3612, Email:

jhkim@umn.edu

Abstract

Sporadic angiosarcomas (ASs) are aggressive vascular sarcomas whose rarity and genomic complexity present significant obstacles in deciphering the pathogenic significance of individual genetic alterations. Numerous fusion genes have been identified across multiple types of cancers, but their existence and significance remain unclear in sporadic ASs. In this study, we leveraged RNA sequencing data from thirteen human ASs and 76 spontaneous canine hemangiosarcomas (HSAs) to identify fusion genes associated with spontaneous vascular malignancies. Ten novel protein-coding fusion genes, including *TEX2-PECAM1* and *ATP8A2-FLT1*, were identified in seven of the thirteen human tumors, with two tumors showing mutations of *TP53*. *HRAS* and *NRAS* mutations were found in ASs without fusions or *TP53* mutations. We found fifteen novel protein-coding fusion genes including *MYO16-PTK2*, *GABRA3-FLT1*, and *AKT3-XPNPEP1* in eleven of the 76 canine HSAs; these fusion genes were seen exclusively in tumors of the angiogenic molecular subtype that contained recurrent mutations in *TP53*, *PIK3CA*, *PIK3R1*, and *NRAS*. In particular, fusion genes and mutations of *TP53* co-occurred in tumors with higher frequency than expected by random chance, and they enriched gene signatures predicting activation of angiogenic pathways. Comparative transcriptomic analysis of human ASs and canine HSAs identified shared molecular signatures associated with activation of PI3K/AKT/mTOR pathways. Our data show that, while driver events of malignant vasoformative tumors of humans and dogs include diverse mutations and stochastic rearrangements that create novel fusion genes, convergent transcriptional programs govern the highly conserved morphological organization and biological behavior of these tumors in both species.

Introduction

Sarcomas are diverse tumors that arise from cells of mesenchymal origin in soft tissues such as blood and lymphatic vessels, fat, bone, cartilage, muscle, and connective tissues. The heterogeneity of sarcomas has provided an impetus for developing molecular approaches to classify these tumors (1,2), leading to their categorization into genomically simple and genomically complex sarcomas (1,3). Angiosarcomas (ASs) are rare, highly aggressive, genomically complex sarcomas of blood vessel-forming cells (3,4). The five-year survival rate of AS is approximately 40% (5-7), but half of patients have metastatic or unresectable disease with a median overall survival of less than 6 months (8). The events that drive progression are incompletely understood; previous studies have identified recurrent mutations of *RAS*, *PTPRB*, *PLCG1*, *KDR* (kinase insert domain receptor, also known as *VEGFR2*), *TP53*, *PIK3CA*, and *FLT4* (*VEGFR3*) in human ASs (9-12). *MYC* gene amplification and alterations in the TP53, CDKN2, NF-κB/IL-6, PIK3CA/AKT/mTOR pathways have also been reported (13); however, these studies represent a small case series, precluding definitive conclusions regarding pathogenic mechanisms that contribute to the genetic cause and to the progression of the disease.

Hemangiosarcoma (HSA) is a malignant vascular tumor that is common in dogs with an estimated tens of thousands of cases diagnosed each year (14-16). Canine HSA shares clinical and morphological features with human AS, as well as aspects of its mutational landscape (17-20). We previously documented three molecular subtypes of HSA, characterized by angiogenic, inflammatory, and adipogenic transcriptomic signatures (21). These gene expression signatures are conserved in HSA progenitor cells that show multipotency and self-renewal (21). Nevertheless, the transcriptional state of these HSA progenitor cells seems to be somewhat malleable, regulated by immune and metabolic reprogramming (22). Mutations in genes that

regulate genomic integrity, such as *TP53,* can alter the intrinsic transcriptional program of tumor

cells; however, genomic instability in the tumor can create even more dramatic changes by

modulating transcriptional programs of heterotypic stromal cells in the tumor tissue, as well as in

the composition of the niche (23,24).

Chromosome translocations and the resulting fusion genes are important contributors to

the pathogenesis of cancer, particularly in sarcomas and hematopoietic malignancies (25).

However, the nature and frequency of these events in canine HSA and human AS remains

unclear. Here, we used next generation RNA sequencing (RNA-Seq) data to identify fusion

genes in thirteen human ASs and 76 visceral HSAs originating from 74 dogs, and we

investigated the relationship of these fusions to the mutational landscape of the tumor. We

identified ten novel protein-coding fusion genes including *TEX2-PECAM1* and *ATP8A2-FLT1* in

seven of thirteen human ASs, and two of the fusion-detected tumors showed mutations of *TP53*

(R248Q and P250L). In canine HSAs, we found novel protein-coding fusion genes in a subset of

the tumors of the angiogenic subtype. These fusion genes co-occurred with *TP53* mutations and

were associated with gene enrichment for activated angiogenic pathways in the tumors. Our data

suggest that genomic instability induced by mutations of *TP53* creates a permissive environment

for fusion genes, with selection for angiogenic molecular programs in malignant vasoformative

tumors. Our data also demonstrate that human AS and canine HSA maintain molecular programs

that activate convergent signaling pathways to establish angiogenic phenotypes despite their

genomic complexity.

Materials and Methods

Human tissue samples

Snap frozen and formalin fixed paraffin embedded (FFPE) tissues for human biospecimens were

obtained from the University of Minnesota Biological Materials Procurement Network (UMN

BioNet) and from the Cooperative Human Tissue Network (CHTN) under their standardized

patient consent protocols. The demographic characteristics of human patients from whom we

obtained ASs ($n = 13$) and normal tissue samples ($n = 6$) are summarized in **Supplementary**

Table S1.

Dog tissue samples

Seventy-six snap frozen and FFPE tissue samples were obtained from 74 dogs with HSAs.

Frozen and FFPE tissues samples from 10 dogs with splenic hematomas, which are benign

lesions with enlarged vascular spaces lined by endothelial cells, were used as controls. Samples

were obtained as part of medically necessary diagnostic procedures and were used for research

with owner consent. The origin of these samples was reported previously (14,21,26-28), or they

were collected from dogs with HSA or with splenic hematomas at the Veterinary Medical

Center, University of Minnesota. Procedures involving animal use were done with approval and

under the supervision of the University of Minnesota Animal Care and Use Committee

(protocols 1110A06186, 1507-32804A, 0802A27363, 1101A94713, 1312-31131A, and 1702-

34548A). The demographic characteristics of dogs ($n = 74$) from whom we acquired HSA and

non-malignant splenic hematomas ($n = 10$) are summarized in **Supplementary Table S2**.

Histological assessment

FFPE sections (4 μm) were stained with hematoxylin and eosin (H&E) and examined by

veterinary pathologists to assign a histological diagnosis of canine HSA. Solid, capillary,

cavernous, or mixed histological subtypes were assigned using accepted criteria (29); mitotic

index (MI) was calculated per 1,000 cells in 5-10 random fields under 400X magnification (30).

H&E slides were further reviewed for tumor content by two board-certified medical pathologists

(ML and PM) (31), with the percent of sample containing viable nucleated cells corresponding to

tumor recorded in a range of 0 to > 90% based on the planar surface of the sections. Diagnostic

and histopathology reports of human tissues were provided by the specimen providers, the UMN

BioNet and the CHTN.

RNA isolation and generation of RNA-Seq libraries

Total RNA was isolated from tissue samples using the TriPure Isolation Reagent (Roche Applied

Science, Indianapolis, IN, USA). The RNeasy Mini Kit (Qiagen, Valencia, CA, USA) was used

for clean-up according to the manufacturer's instructions. RNA-Seq from 74 canine HSA tissues

is published (17,21,32,33) and an additional data set was generated from two canine HSA tissues

and from 10 non-malignant splenic hematoma tissues. Total RNA was also extracted from 13

human AS tissues and from six normal tissues. Two μg of total RNA from each sample were

quantified and assessed for quality; RNA-Seq libraries were generated as described (21) using

the TruSeq RNA sample preparation kit (Illumina Inc., San Diego, CA). Sequencing was

performed using HiSeq 2000 or 2500 systems (Illumina Inc.). Each sample was sequenced to a

targeted depth of 20 – 80 million paired-end reads with mate-pair distance of 50 bp. Primary

analysis and demultiplexing were performed using CASAVA software version 1.8.2 (Illumina

Inc.) to verify the quality of the sequence data. The end result of the CASAVA workflow was

demultiplexed into FASTQ files for analysis. Bioanalyzer quality control and RNA-Seq were

performed at the University of Minnesota Genomics Center (UMGC) or at the Broad Institute.

Bioinformatics analysis

The original FASTQ files prepared from thirteen human ASs and six non-malignant tissues were

mapped to the human reference genome (GRCh38). The FASTQ files generated from 76 canine

HSAs and ten non-malignant splenic hematomas were mapped to the dog reference genome

(Canfam3.1). Sequencing quality was assessed by FastQC. The deFuse algorithm (34) was used

to identify putative fusion events. To discriminate true fusion candidates from artifacts, we

included fusion events with exon boundaries in both fusion partners and excluded events created

from adjacent genes that showed breakpoint homology (>1). We also filtered highly recurrent

fusion events that were found at implausible frequencies across tumor and non-malignant tissue

samples (35) and transcription-induced chimeras. The split sequences of the fusion genes were

validated by *de novo* assembly using Trinity (36). TranscriptsToOrfs and deFuse-Trinity tools

verified the deFuse fusion predictions with Trinity-assembled transcripts and open reading

frames. TopHat2 was used to generate BAM files, and the Integrative Genomics Viewer (IGV

2.3; Broad Institute, Cambridge, MA) was used to visualize the mate pair sequences of fusion

genes. A protein translation tool in Expert Protein Analysis System (ExPASy; SIB Swiss

Institute of Bioinformatics, Lausanne, Switzerland) was used to determine in-frame fusion

proteins. Tumor purity and microenvironment scores were assessed using the bioinformatics

tools ESTIMATE (37) and *xCell* (38).

Reverse transcription polymerase chain reaction (RT-PCR) and Sanger sequencing

RT-PCR was performed to validate fusion transcripts identified by deFuse (39). Briefly, cDNA

was synthesized using SuperScript® VILO cDNA Synthesis Kit and Master Mix (Invitrogen).

PCR amplification was performed using a conventional thermocycler with HotStarTaq DNA

polymerase (Qiagen) or using a LightCycler® 96 (Roche Applied Science, Indianapolis, IN,

USA) with FastStart SYBR Green Master Mix (Roche Applied Science) for quantitative real-

time RT-PCR (40). PCR primer pairs used for fusion gene amplification are presented in the

Results section. *GAPDH* was used as a control for RNA integrity and for the RT-PCR reactions.

The forward and reverse primer sequences for *GAPDH* were 5'-GGA GTC CAC TGG CGT

CTT CAC-3' and 5'-GAG GCA TTG CTG ATG ATC TTG AGG-3', respectively. Relative

mRNA values were expressed as delta-Ct values normalized to *GAPDH*. Sanger sequencing was

performed at the UMGC.

Fluorescence *in situ* hybridization (FISH)

FISH was performed to detect *MYO16-PTK2* and *GABRA3-FLT1* fusion genes by designing

FISH probes derived from the genome-anchored canine CHORI-82 bacterial artificial

chromosome (BAC) library (41). Single locus probes were used for proximal *MYO16* at dog

chromosome (CFA) 22:57,565,917-57,750,789 (clone 183H20), distal *MYO16* at CFA

22:57,750,801-57,967,880 (clone 385H13), and *PTK2* at CFA 13:35,302,679-35,483,060 (clone

451H13) with distinct fluorescent tags. For *GABRA3-FLT1* fusion, break-apart FISH probes

were used for proximal *FLT1* at CFA 25:11,057,892-11,263,935 (clone 363B20) and distal *FLT1*

at CFA 25:11,274,078-11,471,538 (clone 235H9). The PureLink® HiPure Plasmid Maxiprep Kit

(Invitrogen) was used for BAC DNA extraction. For preparation and hybridization of FISH

probes, BAC DNA probes were labeled by Nick Translation Kit (Abbott Molecular) using

Green-500 dUTP, Orange-552 dUTP and Aqua-431 dUTP (Enzo Life Science). Labeled DNA

was precipitated in COT-1 DNA, salmon sperm DNA, sodium acetate and 95% ethanol, then

dried and resuspended in 50% formamide hybridization buffer. The Red-proximal *MYO16*,

Green-distal *MYO16* and Aqua-*PTK2* probes were combined into one 3-color FISH probe for

MYO16-PTK2 fusion. The Red-proximal *FLT1* and Green-distal *FLT1* break-apart probes were

applied for the split *FLT1* gene.

212   FFPE sections (4 μm) were processed according to the Dako IQFISH protocol; probes

were applied to the slide and hybridized for 24 hours at 37°C in a humidified chamber. After

hybridization, slides were washed and counterstained with DAPI. Fluorescent signals were

visualized on an Olympus BX61 microscope workstation (Applied Spectral Imaging, Vista, CA)

with DAPI, FITC, Texas Red and Aqua filter sets. FISH images were captured using an

interferometer-based CCD cooled camera (ASI) and FISHView ASI software. A total of 200

interphase cells were examined for each sample. Non-malignant canine spleen tissues were used

as controls for the FISH experiment.

Validation of somatic mutations using RNA-Seq data

A pipeline was developed to identify the bases present at locations defined as somatic mutations

in the Tumor-Normal exome calls (17). Briefly, RNA-Seq data were mapped using the STAR-

Mapper (42) with STAR-FUSION mapping settings (43) to the Canfam3.1 or GRCh38 genome.

BAM files generated by STAR were sorted and indexed using Samtools (44). Starting from a file

containing somatic mutation locations and a file containing a list of BAM file locations, the

pipeline uses Samtools (44) functions to identify the bases present at each location at each file

and then reports whether a variant is found at that location. For inclusion within the variant file,

at least one sample must have at least 3 reads that support the variant and that represent greater

than 10% of all reads at that location.

Gene expression profiling

CLC Bio Genomics Workbench 10 (CLC Bio, Aarhus, Denmark) and DESeq2 R packages were

used to quantify gene expression and differential gene expression analysis as described (45).

Briefly, paired-end RNA-Seq data with mate-pair distance of 50 bp in FASTQ format were

imported, and sequencing quality was determined. Transcriptomics analysis was then performed

to generate the expression level of each gene presented as total reads by mapping the sequencing

reads to Canfam3.1 or GRCh38. Heatmaps and hierarchical clustering based on average linkage

were visualized using Cluster 3.0, Morpheus (https://clue.io/morpheus), or R packages. GO

Enrichment Analysis (46-48) or Ingenuity® Pathway Analysis software version 8.6 (Qiagen,

Redwood City, CA) were used to define biological functions, canonical pathways, and upstream

regulators associated with differently expressed genes (DEGs) between groups using Benjamini-

Hochberg multiple testing corrections to evaluate significance. For gene expression profiling,

unsupervised PCA and hierarchical clustering were performed to define subtypes of canine

HSAs as described previously (21). Gene expression data of human sarcomas in The Cancer

Genome Atlas (TCGA) database were also compared with our data sets.

TMA generation and immunohistochemistry (IHC)

Canine TMA blocks were generated from 45 HSA tissues, including 32 tumors used for RNA-

Seq and eight non-tumor tissues (six splenic hematomas and two non-malignant liver samples).

Tissue cores of 1-mm diameter in quadruplicate from each sample were assembled in random

order in four TMA blocks. One TMA block with mouse tissues was generated for staining

controls. Immunostaining with CD31, Vimentin and Pan-Cytokeratin antibodies was evaluated to

support tumor content estimates. A human TMA block was generated from ten AS tissues and

six non-malignant tissues (submandibular gland, skin, breast adipose tissue, thigh skeletal

muscle, spleen, and lung).

Unstained TMA sections (4 μm) were de-paraffinized and rehydrated using standard

methods for IHC. All of the immunohistochemical assays, including validation for antibodies,

were performed and optimized at the UMN BioNet Histology Laboratory or the Veterinary

Diagnostic Laboratory at the University of Minnesota. Antibodies used for IHC are summarized

in **Supplementary Table S3**. The immunostaining score assigned to each case was a

semiquantitative assessment derived from the product of two integers, ranging from 0 to 3 and

from 1 to 3, that respectively reflect the percentage of positive cells in a sample and the intensity

of staining at high power magnification (400X) as described previously with some modifications

(49). The percentage of positive cells was scored from 0 to 3+, where 0 reflected specific

staining in <1% of the cells, 1+ reflected specific staining in >1% and <25% of the cells, 2+

reflected specific staining in 25–75% of the cells, and 3+ reflected specific staining in >75% of

the cells. The intensity was assessed as weak (intensity score 1), moderate (intensity score 2), or

strong (intensity score 3). Immunostaining results were scored (ranging from 0 to 9) by

multiplying the percentage of positive cells (score 0-3) by the intensity (score 1-3).

Statistical analysis

Chi-square or Fisher's exact test, was performed for contingency tables analysis. Continuous

values were analyzed by Welch's (Heteroscedastic) T-test or Mann-Whitney U test. The

statistical tests were two-tailed. Statistical analysis was performed using GraphPad Prism 6

(GraphPad Software, Inc., San Diego, CA). *P*-values are reported without inference of

significance, consistent with the American Statistical Association's Statement on Statistical

Significance and P-Values (50).

Results

Novel protein-coding fusion genes are identified in human ASs and canine HSAs

Putative fusion gene events were identified from RNA-Seq data as paired-end sequence reads

that mapped connecting two distant genes (**Supplementary Fig. S1A**). We identified novel in-

frame protein-coding fusion transcripts for ten fusion events in 7 of 13 (53.8%) human ASs **(Fig.**

1A; **Table 1)**. Two of the fusions were inter-chromosomal events and eight were intra-

chromosomal events. The fusions included *TEX2-PECAM1*, which contained the gene that

encodes CD31, and *ATP8A2-FLT1*, a kinase fusion gene that encodes the vascular endothelial

growth factor receptor 1 (VEGFR1). None of the ten fusion events was seen in more than one

tumor, and three of the seven fusion-positive tumors contained two distinct fusion events each.

In canine HSA, we found fifteen novel protein-coding fusion genes in eleven of 76

tumors (14.5%) (**Fig. 1B**; **Table 2**). Ten of the fusions were inter-chromosomal events and five

were intra-chromosomal events. None of the fifteen fusion events was seen in more than one

tumor, and four of the eleven fusion-positive tumors involved two distinct fusion genes each.

One fusion partner in four of the translocations encoded either a protein kinase or a protein

phosphatase associated with angiogenic signaling (*MYO16-PTK2, GABRA3-FLT1*, *AKT3-*

XPNPEP1, and *PTPRB-NOL10*). Eight of the fusion genes were associated with kinase signaling

or kinase binding activity, such as PI3 kinase signaling, a MAP kinase, a receptor tyrosine

kinase, a protein serine/threonine kinase, and an NAD+ kinase. Gene ontology annotations of the

fusion partners for every translocation are described in **Supplementary Table S4** for human AS

and **Supplementary Table S5** for canine HSA. Fusion genes were not present in any of the

human non-malignant tissues ($n = 6$) or canine hematomas ($n = 10$) examined. Conserved driver

translocations such as *BCR-ABL* and *MYC-IGH* that are present in both human and canine

chronic myelogenous leukemia and Burkitt lymphoma, respectively (51), were not identified in

human AS and canine HSA. All the fusion events identified in the seven human ASs and eleven

canine HSAs involved different gene pairs, with the exception of the *FLT1* gene, which created

fusions with a different partner gene in one case from each species.

To determine whether the non-tumor components in the tumor tissue affected the

detection of fusion genes, we quantified tumor content histologically and bioinformatically in

canine HSAs (**Supplementary Fig. S2A - E**). Seventy of the 76 HSA samples were

histologically evaluated, and the tumor content was not different between HSA samples with

fusion events ($n = 11$) and those without fusion events ($n = 59$). We used two independent

bioinformatic tools, *xCell* and ESTIMATE, to predict stromal and immune cell components, and

these algorithms generated consistent output scores (Pearson's $R = 0.84$; $R^2 = 0.71$; $P < 0.00001$)

that showed the presence of fusion genes was not associated with tumor purity. The detection of

fusion genes was also independent of sequencing depth (**Supplementary Fig. S2F**). To rule out

artifacts from the computational process, we validated the presence of the inter-chromosomal

fusion gene, *SCLT1-NIPBL,* in the original human AS sample where it was identified, in two

additional samples where it was undetectable based on sequencing data, and in a non-malignant

tissue sample. The fusion transcript was detectable by quantitative real time RT-PCR

amplification; PCR primer pairs were designed to amplify putative split sequences (up to 200

base pairs) involving the breakpoints identified by deFuse (**Supplementary Fig. S3A**). We

confirmed that the junction sequences between the two genes producing the new fusion event

were amplified by PCR (**Supplementary Fig. S3B**). Four representative fusion transcripts found

in canine HSAs (*MYO16-PTK2*; *AKT3-XPNPEP1*; *AP4E1-BAIAP2*; *NOL10-PTPRB*) were also

detected, but only in the respective cases where they were identified in the sequencing data

(**Supplementary Fig. S3C - E**). Each PCR amplification product was verified by Sanger

sequencing. We then used RT-PCR to evaluate RNA-Seq data from 63 canine tissue samples (53

HSAs and 10 hematomas) for the presence of these four fusion transcripts. The results were

consistent between RNA-Seq and PCR, as we found neither false-positive nor false-negative

events in the samples tested (**Supplementary Table S6**).

Fusion genes are associated with DNA copy number variations

We then determined if any of the fusion partner genes identified in our analysis were associated

with DNA copy number alterations. Publicly available whole Exome-sequencing data generated

from an independent data set of 36 human patients with ASs was used (12). Copy number

variations were found in twelve of the twenty (60%) fusion partner genes: nine genes were

amplified, and five genes were deleted (**Fig. 1C**; **Supplementary Fig. S4**). *TEX2* (39%),

STEAP1B (25%) and *PECAM1* (25%) were the top three genes where copy number gains

occurred most frequently. For canine HSA, we used oligonucleotide array comparative genomic

hybridization (oaCGH) in a larger HSA dataset (*n* = 123) (19). Copy number gains were

observed in 29 of the 30 (96.7%) fusion partners, and copy number losses were observed in 27 of

the 30 (90.0%) fusion partners. Protein kinase-encoding genes, *PTK2*, *FLT1*, and *AKT*3 revealed

a higher frequency of copy number gain: 15.5% gain vs 0.8% loss for *PTK2*; 4.9% gain vs 0.8%

loss for *FLT1*; and 4.1% gain vs 0.8% loss for *AKT3*, suggesting that copy number alterations

leading to dysregulation of downstream kinase signaling contribute at least partly to the

angiogenic program in a subset of canine HSAs (**Fig. 1D**).

Chromosomal translocations resulting in fusion genes are detectable in canine HSAs

We performed FISH to confirm that fusion genes were generated by chromosome translocations.

We chose representative inter-chromosomal fusions, *MYO16-PTK2* and *GABRA3-FLT1* for

cytogenetic validation because PTK2 and VEGFR1 are key molecules that regulate pathogenic

signaling in vascular cancers, including canine HSA (52). **Fig. 2A** illustrates the predicted

structure of *MYO16-PTK2* inter-chromosomal fusion between CFA 22 and CFA 13, based on

deFuse and Sanger sequencing data (**Fig. 2B**). The predicted fusion gene comprises exons 1-32

of *MYO16* (CFA 22) and exons 12-31 of *PTK2* (CFA 13), with the putative junction joining

MYO16 exon 32 and *PTK2* exon 12. Breakage occurs between exons 32 and 33 of *MYO16* at

CFA 22:57,750,807 bp, and between exons 11 and 12 of *PTK2* at CFA 13:35,397,284 bp.

Independent FISH probes identifying the association between proximal and distal *MYO16* and

the breakpoint of *PTK2* (**Fig. 2C**) confirmed the presence of the *MYO16-PTK2* fusion gene

between CFA 22 and 13 in archival FFPE samples from the same dog tumor. The *MYO16-PTK2*

fusion was identified by deFuse and RT-PCR (**Fig. 2D**). The t(CFA 13;CFA 22) translocation

was present in interphase nuclei of 16.8% of the tumor cells, with a smaller subpopulation

showing amplification of the fusion (**Fig. 2E**).

We used break-apart FISH to validate the presence of the *GABRA3-FLT1* fusion gene

(**Fig. 2F**). Split *FLT1* probes were found in 36.7% of tumor cells in archival FFPE samples from

the dog tumor in which the *GABRA3-FLT1* fusion was identified by deFuse and RT-PCR.

Interestingly, in this tumor the intact *FLT1* gene showed consistent amplification (up to four

copies), suggesting Flt-1 (also known as VEGFR1) activation in this tumor might have occurred

through multiple mechanisms (**Fig. 2G**). We next used FISH analysis to assess recurrence of the

MYO16-PTK2 fusion in a tissue microarray (TMA) comprised of 45 visceral HSAs and eight

non-malignant tissues (six spleens; two livers). The *MYO16-PTK2* fusion was once again present

in the sample from the canine tumor in which it was discovered, but it was not seen in any other

sample on the TMA. We also performed FISH to detect the *ATP8A2-FLT1* fusion in human AS

using a break-apart *FLT1* probe, but the fusion was undetectable in our FFPE sample. This might

have been due to the small number of tumor cells that were likely to contain the fusion event in a

heterogeneous clonal population. Since none of the fusion transcripts identified in our cohorts of

human and canine tumors were recurrent, we sought to determine if the fusion events were

associated with other genetic and molecular programs.

Fusion genes and somatic variants in human ASs enrich angiogenic gene signatures

To examine genomic aberrations associated with the fusion genes, we determined somatic

variations and gene expression profiles using RNA-Seq data. In human ASs, *TP53* mutations

(R248Q and P250L) were observed in two of thirteen human ASs, which also had fusion genes

(*SMURF1-TMEM139* and *AGO2-TRAPPC9* in one tumor, *IRF9-THTPA* in the other tumor).

NRAS (Q61L; $n = 1$) or *HRAS* (Q61L; $n = 1$) mutations were also detected, and both were present

in tumors that did not have fusion genes or *TP53* mutations (**Fig. 3A**; **Table 1**). The RNA-Seq

data did not provide evidence of mutations in *PIK3CA, PTEN*, or *KRAS* in this group of thirteen

ASs.

390   We established transcriptomic profiles of human ASs to identify molecular traits that

regulate global gene expression. We identified 1,237 differentially expressed genes between ASs

(n = 13) and non-malignant controls (N = 6) (FDR P-value < 0.05): 490 genes were upregulated

and 747 genes were downregulated in ASs. Biological functions and pathway analysis revealed

that upregulated genes in ASs were associated with cancer, angiogenesis, vasculogenesis, and

development of vasculature (P < 0.00001) (**Supplementary Table S7**). Additionally, we

performed cell type enrichment analysis using the *xCell* tool (38) to predict relative populations

of cellular components that comprised the AS tissues (**Supplementary Fig. S5**). We compared

the cell type signature of AS with that of sarcomas (n = 263) from the TCGA database, which

did not include AS. The results showed that gene signatures associated with endothelial cells and

activated dendritic cells were highly enriched in ASs, while other sarcomas in the TCGA

revealed gene enrichment of smooth muscle cells. Gene expression profiles of non-malignant

samples indicated distinct tissue-specific patterns of submandibular gland, skin, breast adipose

tissue, thigh skeletal muscle, spleen, and lung. The ASs showed upregulation of key angiogenic

genes such as *PECAM1* (*CD31*), *FLT1* (*VEGFR1*), *KDR* (*VEGFR2*), and *FLT4* (*VEGFR3)*

compared to non-malignant tissues (**Fig. 3B**). In addition, 6 of 20 (30.0%) fusion partners

(including *PECAM1*) showed a higher level of expression in the tumors compared to non-

malignant tissues (P < 0.05; fold change in a range from 1.9 to 35.0) (**Fig. 3C**). Collectively, our

data showed enriched angiogenic molecular programs were present in human ASs. Furthermore,

both tumors with *TP53* mutations also harbored fusion genes.

Fusion genes that co-occur with mutations of *TP53* are present exclusively in angiogenic

canine HSAs

In canine HSAs, *TP53* (n = 24/74; 32.4%), *PIK3CA* (n = 16/74; 21.6%), *PIK3R1* (n = 5/74;

6.8%), and *NRAS* (n = 4/74; 5.4%) transcripts showed recurrent mutations that were consistent

with those identified using tumor:normal Exome-sequencing (17). We found associations

between mutations of *TP53* (*TP53*^mt) and *PIK3CA* (*PIK3CA*^mt) and fusion genes (Fusion$^+$)

(**Supplementary Table S8**). Specifically, *TP53*^mt commonly co-occurred with *PIK3CA*^mt (P =

0.004) and with fusion genes (P = 0.004); and Fusion$^+$ tumors were seen in the tumors with

TP53^mt or *PIK3CA*^mt (P = 0.005) more frequently than would be expected by random chance.

When fusion genes co-occurred with *PIK3CA*^mt, they invariably co-occurred with mutations of

TP53 (P = 0.037), and they were not associated with *PIK3CA*^mt alone (P = 0.324).

Next, we sought to determine if fusion genes were preferentially associated with specific

molecular subtypes of canine HSA. We previously defined distinct angiogenic, inflammatory,

and adipogenic molecular subtypes of canine HSA (21). To further validate this classification,

we applied unsupervised principal component analysis (PCA) and hierarchical clustering to

identify distinct groups in the sample cohort from this study (**Supplementary Fig. S6**). Our

results show that the three molecular subtypes (58 angiogenic, 14 inflammatory, and 4

adipogenic) were reproducibly identified in the current dataset, as illustrated in the heatmap of

1,477 DEGs (FDR P < 0.001; fold change > |3|) shown in **Fig. 3D**. Interestingly, fusion genes

were present only in tumors of the angiogenic HSA subtype (P = 0.046). Likewise, *TP53*

mutations were identified in 23 of 56 (41.1%) angiogenic HSAs, and in one of 18 tumors from

the two other molecular subtypes (P = 0.008). The somatic variants of *TP53*, *PIK3CA*, *PIK3R1*,

and *NRAS* were found in 33 of 56 (58.9%) angiogenic HSAs, and in 3 of 18 (16.7%) tumors

from the two other subtypes (P = 0.002). The angiogenic subtype of HSA also showed

upregulation of *PECAM1* (*CD31*), *FLT1* (*VEGFR1*), *KDR* (*VEGFR2)*, and *FLT4* (*VEGFR3)*

compared to the other two HSA subtypes and to non-malignant hematomas (**Fig. 3E**). We found

that five of 32 dogs (16%) with angiogenic HSA lived longer than five months, while four of 10

dogs (40%) with inflammatory HSA survived longer than that (**Supplementary Fig. S7**).

Eighteen of 30 (60.0%) fusion partners, including protein kinase-encoding genes such as *FLT1*,

PTK2, and *AKT3,* showed higher levels of expression in HSAs compared to non-malignant

controls ($P < 0.05$; fold change in a range from 1.3 to 6.0) (**Fig. 3F**). Neither breed, sex, neuter

status, age, nor affected organs were associated with the presence of fusion genes

(**Supplementary Fig. S8**). There was also no association between the fusion events and

histological subtype or mitotic index (**Supplementary Table S9**).

Next, we analyzed DEGs (FDR $P < 0.05$; fold change > |2|) and gene pathways to

examine gene signatures enriched in HSAs that had both fusion genes and *TP53* mutations. We

classified tumors according to their mutations as summarized in **Supplementary Table S10**. Co-

occurrence of fusion genes with *TP53* mutations (i.e., *TP53*^mt/Fusion^+/*PIK3CA*^wt or PF tumors)

was associated with angiogenic and vascular signaling with enrichment of genes in pathways

such as PI3K, VEGF, and PDGF (**Supplementary Table S11 - S13**). Thirteen genes that were

commonly enriched in PF tumors were associated with activation of WNT3A as an upstream

regulator (**Supplementary Fig. S9**). **Fig. 4** illustrates a model integrating the data from these

findings to highlight potential pathogenetic contributions of fusion genes and recurrent mutations

in canine HSA.

Human ASs and canine HSAs establish molecular programs that activate convergent

signaling pathways

To determine if the genetic and molecular features of human AS and canine HSAs contributed to

the activation of functional pathways, we performed IHC in eleven human AS tissues and in 44

canine HSAs (**Fig. 5**; **Supplementary Tables S14** and **S15**). First, we evaluated the effects of

TP53 mutation on the presence and location of p53, phospho-p53 (Ser15), and phospho-p53

(Ser20) (**Fig. 5A** and **B**). In human ASs, nuclear expression of p53 was found in all of eleven

(100%) tumors, showing various levels of expression. Nuclear expression of phospo-p53

(Ser15) was seen in seven of eleven (63.6%) tumors, showing low expression in five of seven

(71%) tumors (IHC score $\leq$ 3). Nuclear and cytoplasmic expression of phospho-p53 (Ser20)

protein was detected in all eleven (100%) tumors, and seven of those showed high expression

(IHC score $\geq$ 7). In canine HSAs, p53 protein was localized to the nucleus in 34 of 44 (77%)

tumors with various levels of expression. Immunoreactivity of phospho-p53 (Ser15) was

observed in the nuclei of tumor cells in 38 of 40 (95%) cases, with 34 (90%) showing low or

medium expression (IHC score $\leq$ 6). Nuclear and cytoplasmic expression of phospho-p53

(Ser20) was seen in all of 40 HSAs (100%), with 32 tumors (80%) showing high levels of

expression. These data revealed that patterns of p53 and activated p53 were comparable in

human ASs and canine HSAs; especially, p53 was strongly phosphorylated at residue Ser20 in

tumors from both species. We found no association between phosphorylated p53 and *TP53*

mutations or fusion genes (**Supplementary Fig. S10**), suggesting that DNA damage and cellular

stress are widespread among these tumors, and they are likely to activate p53-mediated repair

mechanisms independent of these genetic alterations.

The PI3K/AKT/mTOR signaling pathway is important for regulation of angiogenic,

vascular, and energetic functions. To assess whether PI3K mutations resulted in higher levels of

downstream pathway activation, we evaluated expression of AKT and phospho-AKT proteins in

human ASs and canine HSAs (**Fig. 5C** and **D**). Expression of nuclear and cytoplasmic AKT

protein was observed in all eleven human ASs (100%) with eight of eleven (73%) tumors

showing medium or high expression. Expression of phospho-AKT (Thr308) was evaluated in

eight tumors; all of them (100%) showed weak or medium levels of expression. Similarly, AKT

protein was detectable in the nucleus and cytoplasm of all forty (100%) canine HSAs with 37

(93%) showing medium or high expression. Phospho-AKT (Thr308) was detected in the nuclei

and cytoplasm of all evaluable 39 (100%) canine HSAs, with 34 (87%) expressing medium or

high level of the protein. Strong AKT immunoreactivity was also seen in scattered stromal cells

in both human AS and canine HSA. Neither mutations of *TP53*, *PIK3CA*, *PIK3R1* nor the

presence of fusion genes were associated with expression of AKT and phospho-AKT in human

or canine tumors (**Supplementary Fig. S11**). Since *PIK3CA* and *PIK3R1* mutations were

undetectable in this set of eleven human ASs, we examined expression of mTOR and phospho-

mTOR (Ser2448) proteins as surrogates to confirm activation of their downstream pathways in

these tumors. mTOR protein was observed in the nuclei and cytoplasm of all eleven tumors (IHC

score $\geq$ 6), and it was not associated with the presence of *TP53* mutations or fusions

(**Supplementary Fig. S12A** and **B**). However, nuclear and cytoplasmic expression of phospho-

mTOR was higher in ASs that had *TP53* mutations or that had fusion genes than it was in tumors

without one of these genetic changes (**Supplementary Fig. S12C** and **D**). In summary, the

immunostaining data suggest that human AS and canine HSA have comparable activation of the

p53 and PI3K/AKT/mTOR pathways, and these events are largely independent of their

mutational states. Our results further suggest that these vasoformative tumors from both species

activate convergent signaling pathways that contribute to their final architecture and organization

with predictable enrichment of angiogenic gene signatures.

Discussion

For this study, our objective was to identify novel fusion genes in human ASs and spontaneous

canine HSAs. We showed that novel protein-coding fusion genes were identified in

approximately 50% of human ASs of which two had *TP53* mutations. In canine HSAs, protein-

coding fusion genes were detectable in ~15% of tumors, and those were associated with p53

deficiency and enrichment of angiogenic gene signatures. Our data suggest that convergent

molecular mechanisms associated with p53 inactivation and enhanced PI3K/AKT/mTOR

signaling pathways are operational in genomically complex human ASs and canine HSAs.

In the past decade, advances in next-generation sequencing and bioinformatics have

enabled genome-wide identification of unbiased cancer-associated fusion transcripts in a variety

of tumor types. Previous studies have reported 7,887 fusion transcripts identified across thirteen

tumor types in TCGA datasets and 9,928 fusion genes with a 3% recurrence rate in the Mitelman

Database of Chromosome Aberrations and Gene Fusions in Cancer (53-55). These findings

illustrate the complexity of the cancer-associated fusion gene landscape, showing a relatively

high rate of fusions with low recurrence, possibly arising from catastrophic chromosome

rearrangements by chromothripsis (56) and chromoplexy (57). Despite this relatively high

frequency of fusion genes, a solution to define their pathogenic significance remains elusive.

One key finding from this work was that protein-coding fusion genes co-occurred with mutations

of *TP53* in the angiogenic molecular subtype of canine HSA, suggesting that genomic instability

might create a predisposition for translocations and the resultant fusion genes and that, in turn,

these fusion genes create unique transcriptional programs that promote angiogenic phenotypes in

these p53-deficient backgrounds.

Specifically, kinase fusion genes involving *FLT1*, *PTK2*, and *AKT3* can activate key

convergent gene pathways associated with blood vessel formation and remodeling (58), and they

represent potential therapeutic targets for kinase inhibitors (35,59). When we consider that

sarcomas have the highest frequency of kinase fusions in TCGA datasets (35), but that they also

have extremely low recurrence, a more rational approach might be to develop agents that target

these convergent angiogenic pathways instead of the products from the individual fusion genes

themselves (60).

The two fusion genes that we confirmed by genomic structural evaluation in canine

HSAs were present in approximately 20 - 40% of cells in the tumor, both genes showing chaotic

amplification. Several explanations could account for these observations. One is that histology

and bioinformatics assays overestimated tumor content and tumor purity. Another is that fusions

are epiphenomena arising from chaotic genomes with no influence on selection. A third, which

we believe is most likely, is that translocation and the resulting fusion events occur stochastically

in genomically unstable cells late in the course of tumor evolution. However, the enrichment of

fusion genes and angiogenic transcriptional programs suggests that these traits endow tumor cells

with selective growth and/or survival advantages that contribute to tumor progression by

promoting proangiogenic environments. It is worth noting that the selective pressures in the AS

milieu favor not only fusion-positive clones, but also fusion-negative clones, as the establishment

of a proangiogenic environment could improve survival of all the subpopulations within the

tumor. Indeed, a similar mechanism might be operative in alveolar rhabdomyosarcomas, where

PAX3-FOXO1A fusions are necessary for tumor initiation but have no effect on tumor recurrence

(61,62). Further work will be necessary to distinguish which among these non-mutually

exclusive possibilities are operative, and to better understand the role of fusion genes in tumor

evolution of human AS and canine HSA and, potentially, in promoting clonal heterogeneity

through the creation of a permissive niche.

Fusion genes have been reported in human ASs (10,63-65). For instance, one study found

a *CIC-LEUTX* fusion in one of 120 (0.8%) FFPE ASs examined (10); another found a *CEP85L-*

ROS1 fusion in one of 34 (3.0%) ASs examined (63); and a third found an *EWSR1-ATF1* fusion

in one case of AS (65). A *NUP160-SLC43A3* fusion has also been reported in the ISO-HAS AS

cell line (64). However, none of these fusion genes has been identified recurrently in subsequent

studies of AS samples. While these observations are consistent with our stochastic hypothesis,

we cannot completely exclude the possibility that fusion genes in human ASs, or for that matter

in canine HSAs, are non-pathogenic passenger aberrations.

A larger case series will be required to define the fusion gene landscape in human AS,

but canine HSA provides potential insights for what might be expected. Mutations of *TP53* are

largely mutually exclusive of mutations in *KDR*, *PIK3CA*, and *RAS* gene family in human AS,

and the mutational patterns seem to be associated with the location of the primary tumor

(10,12,17,66,67). We see a similar pattern emerge in a subset of canine HSA, and from our data

we propose a model that can be used as a foundation to test mechanistic links between the

mutational and transcriptional landscapes in malignant vascular tumors (**Fig. 4**) and determine

their roles in tumor progression. In this model, inflammatory HSAs harbor no mutations of

PIK3CA, and only rarely of *TP53*, maintaining sufficient genomic stability that disfavors

formation of fusion genes. Furthermore, the transcriptional programs in these inflammatory

HSAs are weakly angiogenic, and their permissive inflammatory environments restrain growth

and metastasis. Conversely, angiogenic HSAs harbor frequent mutations of *PIK3CA* and *TP53*,

and fusion events. Mutations of *PIK3CA* in p53-proficient backgrounds promote pro-angiogenic

environments, while in p53-deficient backgrounds, these mutations promote altered chromatin

regulation and immunomodulatory transcriptional programs. Finally, mutations of *TP53* enable

genomic instability with formation of fusion genes. These events are stochastic, but fusion genes

that promote pro-angiogenic transcriptional programs can enhance or even supplant the effects of

PIK3CA mutations and create environments that accelerate tumor growth and metastatic

propensity.

Molecular distinctions among human ASs could be driven by their clinical phenotype and

potential therapeutic responses (9,68,69). For instance, a subset of ASs harbor gene

amplifications of *MYC* and *FLT4* which frequently co-occur in tumors associated with ultraviolet

(UV) irradiation- or therapeutic radiation. Mutational signatures associated with UV exposure

and high mutational burden might predict more favorable immunotherapeutic responses in AS

patients, as they do in patients diagnosed with malignant melanoma; however, supportive clinical

trials to test this premise are limited, and the use of immune checkpoint inhibitors in human AS

patients thus far has yielded mixed results (70,71). The mutational signatures in canine HSA are

largely confined to the "aging" (cellular replication) signature (17), and total mutational burden

is relatively low (17,72), so this condition is unlikely to provide a model to address the utility of

immunotherapy in this context. However, canine HSA could provide a suitable model to address

other treatments, whether pharmacologic or immunologic, directed at the molecular programs

that drive progression and maintenance of the tumors in both species (33). Such validation

studies could alter the paradigms for diagnosis and treatment of human AS and canine HSA, as

well as of other aggressive, genomically complex sarcomas that affect humans and dogs alike.

Data availability

RNA-Seq gene expression data generated from human sarcomas are available from the TCGA

Research Network (https://www.cancer.gov/tcga). Exome sequencing data from human

angiosarcomas are available from The Angiosarcoma Project (https://ascproject.org), a project of

Count Me In (https://joincountmein.org/). RNA-Seq data from human AS tissues are available

through the Gene Expression Omnibus (GEO; http://www.ncbi.nlm. nih.gov/geo; accession

number GSE163359). RNA-Seq data from canine HSA tissues are published (17,21,32,33) and

available through the GEO (accession number GSE95183) and the NCBI Sequence Read

Archive (accession number PRJNA562916). All other data generated from this study are

available upon request to the corresponding author.

Acknowledgements

The authors would like to acknowledge Dr. Corrie Painter for reviewing the manuscript and

providing feedback. The authors acknowledge Mitzi Lewellen for assistance with inventory,

database management, and editorial assistance. The authors would also like to thank Lauren

Mills for processing of the next generation sequencing data and Dr. Douglas Yee, Director of

Masonic Cancer Center, for assisting with the collection of human tissues. Human biospecimens

were obtained from the UMN BioNet and from the CHTN. Tissue samples were provided by the

CHTN which is funded by the National Cancer Institute (NCI). Other investigators may have

received specimens form the same subjects. This work was partially supported by grants

1R03CA191713-01 (J.F. Modiano, A.L. Sarver, J.H. Kim) and R37CA218570 (E.K. Karlsson)

from the NCI of the National Institutes of Health (NIH), grants #422 (J.F. Modiano) and 1889-G

(J.F. Modiano, M. Breen, K. Lindblad-Toh) from the AKC Canine Health Foundation, grant

JHK15MN-004 (J.H. Kim) from the National Canine Cancer Foundation, grant D10-501 (J.F.

Modiano, M. Breen, K. Lindblad-Toh) from Morris Animal Foundation, and a grant from

Swedish Cancerfonden (K. Lindblad-Toh). This work was also supported by an NIH NCI R50

grant, CA211249 (A.L. Sarver). The NIH Comprehensive Cancer Center Support Grant to the

Masonic Cancer Center, University of Minnesota (P30 CA077598) provided support for the

cytogenetic analyses performed in the Cytogenomics Shared Resource. K. Megquier is supported

by the NCI of the NIH under Award Number F32CA247088. The content is solely the

responsibility of the authors and does not necessarily represent the official views of the NIH. M.

Breen is supported in part by the Oscar J. Fletcher Distinguished Professorship in Comparative

Oncology Genetics at North Carolina State University. K. Lindblad-Toh is supported by a

Distinguished Professor award from the Swedish Research Council. J.F. Modiano is supported

by the Alvin and June Perlman Chair in Animal Oncology. The UMGC

(http://genomics.umn.edu) supported for generation of genomic sequencing data libraries, and

the Minnesota Supercomputing Institute (MSI) at the University of Minnesota

(http://www.msi.umn.edu) provided computational resources that contributed to the results in

this study. The authors gratefully acknowledge donations to the Animal Cancer Care and

Research Program of the University of Minnesota that helped support this project.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Authors' Contributions

Conception and design: J.H. Kim, J.F. Modiano

Development of methodology: J.H. Kim, K. Megquier, A.L. Sarver, R. Thomas, J.F. Modiano

Acquisition of data (provided animals, acquired and managed patients, provided facilities,

etc.): J.H. Kim, K. Megquier, R. Thomas, A.L. Sarver, N. Cheng, M.A. Linden, P. Murugan, L.

Oseth, C.L. Foster

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational

analysis): J.H. Kim, K. Megquier, R. Thomas, A.L. Sarver, J.M. Song, Y.T. Kim, N. Cheng,

M.A. Linden, P. Murugan, I. Elvers, R. Swofford, J. Turner-Maier, E.K. Karlsson, M. Breen, K.

Lindblad-Toh, J.F. Modiano

Writing, review, and/or revision of the manuscript: J.H. Kim, K. Megquier, R. Thomas, A.J.

Graef, K. Lindblad-Toh, J.F. Modiano with help from all authors

Administrative, technical, or material support: A.J. Graef

Study supervision: J.H. Kim, M. Breen, K. Lindblad-Toh, J.F. Modiano

References

1.   Taylor BS, Barretina J, Maki RG, Antonescu CR, Singer S, Ladanyi M. Advances in

sarcoma genomics and new therapeutic targets. Nat Rev Cancer **2011**;11(8):541-57 doi

10.1038/nrc3087.

2.   Neuville A, Ranchere-Vince D, Dei Tos AP, Montesco MC, Hostein I, Toffolatti L*, et al.*

Impact of molecular analysis on the final sarcoma diagnosis: a study on 763 cases

collected during a European epidemiological study. Am J Surg Pathol **2013**;37(8):1259-

68 doi 10.1097/PAS.0b013e31828f51b9.

3.   Helman LJ, Meltzer P. Mechanisms of sarcoma development. Nat Rev Cancer

2003;3(9):685-94 doi 10.1038/nrc1168.

4.   Italiano A, Chen CL, Thomas R, Breen M, Bonnet F, Sevenet N*, et al.* Alterations of the

p53 and PIK3CA/AKT/mTOR pathways in angiosarcomas: a pattern distinct from other

sarcomas with complex genomics. Cancer **2012**;118(23):5878-87 doi

10.1002/cncr.27614.

5.   Buehler D, Rice SR, Moody JS, Rush P, Hafez GR, Attia S*, et al.* Angiosarcoma

outcomes and prognostic factors: a 25-year single institution experience. Am J Clin

Oncol **2014**;37(5):473-9 doi 10.1097/COC.0b013e31827e4e7b.

6.   Lahat G, Dhuka AR, Hallevi H, Xiao L, Zou C, Smith KD*, et al.* Angiosarcoma: clinical

and molecular insights. Ann Surg **2010**;251(6):1098-106 doi

10.1097/SLA.0b013e3181dbb75a.

7.   Fayette J, Martin E, Piperno-Neumann S, Le Cesne A, Robert C, Bonvalot S*, et al.*

Angiosarcomas, a heterogeneous group of sarcomas with specific behavior depending on

primary site: a retrospective study of 161 cases. Ann Oncol **2007**;18(12):2030-6 doi

10.1093/annonc/mdm381.

8.   Smrke A, Hamm J, Karvat A, Simmons C, Srikanthan A. A retrospective review of 145

patients with angiosarcoma: Radiation therapy, extent of resection and chemotherapy are

important predictors of survival. Molecular and Clinical Oncology. Molecular and

Clinical Oncology **2020**;13:179-85 doi https://doi.org/10.3892/mco.2020.2055.

9.   Behjati S, Tarpey PS, Sheldon H, Martincorena I, Van Loo P, Gundem G*, et al.*

Recurrent PTPRB and PLCG1 mutations in angiosarcoma. Nat Genet **2014**;46(4):376-9

doi 10.1038/ng.2921.

10.   Huang SC, Zhang L, Sung YS, Chen CL, Kao YC, Agaram NP*, et al.* Recurrent CIC

Gene Abnormalities in Angiosarcomas: A Molecular Study of 120 Cases With

Concurrent Investigation of PLCG1, KDR, MYC, and FLT4 Gene Alterations. Am J

Surg Pathol **2016**;40(5):645-55 doi 10.1097/PAS.0000000000000582.

11.   Antonescu CR, Yoshida A, Guo T, Chang NE, Zhang L, Agaram NP*, et al.* KDR

activating mutations in human angiosarcomas are sensitive to specific kinase inhibitors.

Cancer Res **2009**;69(18):7175-9 doi 10.1158/0008-5472.CAN-09-2068.

12.   Painter CA, Jain E, Tomson BN, Dunphy M, Stoddard RE, Thomas BS*, et al.* The

Angiosarcoma Project: enabling genomic and clinical discoveries in a rare cancer through

patient-partnered research. Nat Med **2020**;26(2):181-7 doi 10.1038/s41591-019-0749-z.

13.   Antonescu C. Malignant vascular tumors--an update. Mod Pathol **2014**;27 Suppl 1:S30-8

doi 10.1038/modpathol.2013.176.

14.   Lamerato-Kozicki AR, Helm KM, Jubala CM, Cutter GR, Modiano JF. Canine

hemangiosarcoma originates from hematopoietic precursors with potential for endothelial

differentiation. Exp Hematol **2006**;34(7):870-8.

15.   Cohen SM, Storer RD, Criswell KA, Doerrer NG, Dellarco VL, Pegg DG*, et al.*

Hemangiosarcoma in rodents: mode-of-action evaluation and human relevance. Toxicol

Sci **2009**;111(1):4-18 doi 10.1093/toxsci/kfp131.

16.   Mullin C, Clifford CA. Histiocytic Sarcoma and Hemangiosarcoma Update. Vet Clin

North Am Small Anim Pract **2019**;49(5):855-79 doi 10.1016/j.cvsm.2019.04.009.

17.   Megquier K, Turner-Maier J, Swofford R, Kim JH, Sarver AL, Wang C*, et al.*

Comparative Genomics Reveals Shared Mutational Landscape in Canine

Hemangiosarcoma and Human Angiosarcoma. Mol Cancer Res **2019**;17(12):2410-21 doi

10.1158/1541-7786.MCR-19-0221.

18.   Wang G, Wu M, Durham AC, Radaelli E, Mason NJ, Xu X*, et al.* Molecular subtypes in

canine hemangiosarcoma reveal similarities with human angiosarcoma. PLoS One

2020;15(3):e0229728 doi 10.1371/journal.pone.0229728.

19.   Thomas R, Borst L, Rotroff D, Motsinger-Reif A, Lindblad-Toh K, Modiano JF*, et al.*

Genomic profiling reveals extensive heterogeneity in somatic DNA copy number

aberrations of canine hemangiosarcoma. Chromosome Res **2014**;22(3):305-19 doi

10.1007/s10577-014-9406-z.

20.   Andersen NJ, Nickoloff BJ, Dykema KJ, Boguslawski EA, Krivochenitser RI, Froman

RE*, et al.* Pharmacologic inhibition of MEK signaling prevents growth of canine

hemangiosarcoma. Mol Cancer Ther **2013**;12(9):1701-14 doi 10.1158/1535-7163.MCT-

12-0893.

21.   Gorden BH, Kim JH, Sarver AL, Frantz AM, Breen M, Lindblad-Toh K*, et al.*

Identification of three molecular and functional subtypes in canine hemangiosarcoma

through gene expression profiling and progenitor cell characterization. Am J Pathol

2014;184(4):985-95 doi 10.1016/j.ajpath.2013.12.025.

22.   Kim JH, Frantz AM, Sarver AL, Gorden Klukas BH, Lewellen M, O'Brien TD*, et al.*

Modulation of fatty acid metabolism and immune suppression are features of in vitro

733  tumour sphere formation in ontogenetically distinct dog cancers. Vet Comp Oncol

734  **2018**;16(1):E176-E84 doi 10.1111/vco.12368.

735 23. Hill R, Song Y, Cardiff RD, Van Dyke T. Selective evolution of stromal mesenchyme

736  with p53 loss in response to epithelial tumorigenesis. Cell **2005**;123(6):1001-11 doi

737  10.1016/j.cell.2005.09.030.

738 24. Goldstein I, Marcel V, Olivier M, Oren M, Rotter V, Hainaut P. Understanding wild-type

739  and mutant p53 activities in human cancer: new landmarks on the way to targeted

740  therapies. Cancer Gene Ther **2011**;18(1):2-11 doi 10.1038/cgt.2010.63.

741 25. Mitelman F, Johansson B, Mertens F. The impact of translocations and gene fusions on

742  cancer causation. Nat Rev Cancer **2007**;7(4):233-45 doi 10.1038/nrc2091.

743 26. Fosmire SP, Dickerson EB, Scott AM, Bianco SR, Pettengill MJ, Meylemans H*, et al.*

744  Canine malignant hemangiosarcoma as a model of primitive angiogenic endothelium.

745  Lab Invest **2004**;84(5):562-72.

746 27. Dickerson EB, Thomas R, Fosmire SP, Lamerato-Kozicki AR, Bianco SR, Wojcieszyn

747  JW*, et al.* Mutations of phosphatase and tensin homolog deleted from chromosome 10 in

748  canine hemangiosarcoma. Vet Pathol **2005**;42(5):618-32 doi 10.1354/vp.42-5-618.

749 28. Tamburini BA, Trapp S, Phang TL, Schappa JT, Hunter LE, Modiano JF. Gene

750  expression profiles of sporadic canine hemangiosarcoma are uniquely associated with

751  breed. PLoS One **2009**;4(5):e5549 doi 10.1371/journal.pone.0005549.

752 29. Kim J-H, Graef A, Dickerson E, Modiano J. Pathobiology of Hemangiosarcoma in Dogs:

753  Research Advances and Future Perspectives. Veterinary Sciences **2015**;2(4):388.

30.   Rudolph P, Peters J, Lorenz D, Schmidt D, Parwaresch R. Correlation between mitotic

and Ki-67 labeling indices in paraffin-embedded carcinoma specimens. Hum Pathol

1998;29(11):1216-22.

31.   Tschida BR, Temiz NA, Kuka TP, Lee LA, Riordan JD, Tierrablanca CA*, et al.* Sleeping

Beauty Insertional Mutagenesis in Mice Identifies Drivers of Steatosis-Associated

Hepatic Tumors. Cancer Res **2017**;77(23):6576-88 doi 10.1158/0008-5472.CAN-17-

2281.

32.   Tonomura N, Elvers I, Thomas R, Megquier K, Turner-Maier J, Howald C*, et al.*

Genome-wide association study identifies shared risk loci common to two malignancies

in golden retrievers. PLoS Genet **2015**;11(2):e1004922 doi

10.1371/journal.pgen.1004922.

33.   Borgatti A, Koopmeiners JS, Sarver AL, Winter AL, Stuebner K, Todhunter D*, et al.*

Safe and Effective Sarcoma Therapy through Bispecific Targeting of EGFR and uPAR.

Mol Cancer Ther **2017**;16(5):956-65 doi 10.1158/1535-7163.MCT-16-0637.

34.   McPherson A, Hormozdiari F, Zayed A, Giuliany R, Ha G, Sun MG*, et al.* deFuse: an

algorithm for gene fusion discovery in tumor RNA-Seq data. PLoS Comput Biol

2011;7(5):e1001138 doi 10.1371/journal.pcbi.1001138.

35.   Stransky N, Cerami E, Schalm S, Kim JL, Lengauer C. The landscape of kinase fusions

in cancer. Nat Commun **2014**;5:4846 doi 10.1038/ncomms5846.

36.   Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J*, et al.* De novo

transcript sequence reconstruction from RNA-seq using the Trinity platform for reference

generation and analysis. Nat Protoc **2013**;8(8):1494-512 doi 10.1038/nprot.2013.084.

37.   Yoshihara K, Shahmoradgoli M, Martinez E, Vegesna R, Kim H, Torres-Garcia W, *et al.*

777   Inferring tumour purity and stromal and immune cell admixture from expression data.

778   Nat Commun **2013**;4:2612 doi 10.1038/ncomms3612.

779   38.   Aran D, Hu Z, Butte AJ. xCell: digitally portraying the tissue cellular heterogeneity

780   landscape. Genome Biol **2017**;18(1):220 doi 10.1186/s13059-017-1349-1.

781   39.   Kim JH, Frantz AM, Anderson KL, Graef AJ, Scott MC, Robinson S, *et al.* Interleukin-8

782   promotes canine hemangiosarcoma growth by regulating the tumor microenvironment.

783   Experimental cell research **2014**;323(1):155-64 doi 10.1016/j.yexcr.2014.02.020.

784   40.   Anderson KL, Snyder KM, Ito D, Lins DC, Mills LJ, Weiskopf K, *et al.* Evolutionarily

785   conserved resistance to phagocytosis observed in melanoma cells is insensitive to

786   upregulation of pro-phagocytic signals and to CD47 blockade. Melanoma Res

787   **2020**;30(2):147-58 doi 10.1097/CMR.0000000000000629.

788   41.   Thomas R, Duke SE, Karlsson EK, Evans A, Ellis P, Lindblad-Toh K, *et al.* A genome

789   assembly-integrated dog 1 Mb BAC microarray: a cytogenetic resource for canine cancer

790   studies and comparative genomic analysis. Cytogenet Genome Res **2008**;122(2):110-21

791   doi 10.1159/000163088.

792   42.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, *et al.* STAR: ultrafast

793   universal RNA-seq aligner. Bioinformatics **2013**;29(1):15-21 doi

794   10.1093/bioinformatics/bts635.

795   43.   Haas B, Dobin A, Stransky N, Li B, Yang X, Tickle T, *et al.* 2017 STAR-Fusion: Fast

796   and Accurate Fusion Transcript Detection from RNA-Seq. bioRxiv

797   <https://www.biorxiv.org/content/early/2017/03/24/120295>.

798   44.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N*, et al.* The Sequence

799   Alignment/Map format and SAMtools. Bioinformatics **2009**;25(16):2078-9 doi

800   10.1093/bioinformatics/btp352.

801   45.   Qian X, Khammanivong A, Song JM, Teferi F, Upadhyaya P, Dickerson E*, et al.* RNA-

802   sequencing studies identify genes differentially regulated during inflammation-driven

803   lung tumorigenesis and targeted by chemopreventive agents. Inflamm Res

804   **2015**;64(5):343-61 doi 10.1007/s00011-015-0815-z.

805   46.   Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM*, et al.* Gene ontology:

806   tool for the unification of biology. The Gene Ontology Consortium. Nat Genet

807   **2000**;25(1):25-9 doi 10.1038/75556.

808   47.   The Gene Ontology C. The Gene Ontology Resource: 20 years and still GOing strong.

809   Nucleic Acids Res **2019**;47(D1):D330-D8 doi 10.1093/nar/gky1055.

810   48.   Mi H, Muruganujan A, Ebert D, Huang X, Thomas PD. PANTHER version 14: more

811   genomes, a new PANTHER GO-slim and improvements in enrichment analysis tools.

812   Nucleic Acids Res **2019**;47(D1):D419-D26 doi 10.1093/nar/gky1038.

813   49.   Rodriguez AM, Graef AJ, LeVine DN, Cohen IR, Modiano JF, Kim JH. Association of

814   Sphingosine-1-phosphate (S1P)/S1P Receptor-1 Pathway with Cell Proliferation and

815   Survival in Canine Hemangiosarcoma. J Vet Intern Med **2015**;29(4):1088-97 doi

816   10.1111/jvim.13570.

817   50.   Ronald L. Wasserstein, Lazar NA. The ASA's Statement on p-Values: Context, Process,

818   and Purpose. The American Statistician **2016**;70(2):129-33 doi

819   10.1080/00031305.2016.1154108.

820   51.   Breen M, Modiano JF. Evolutionarily conserved cytogenetic changes in hematological

821   malignancies of dogs and humans--man and his best friend share more than

822   companionship. Chromosome Res **2008**;16(1):145-54 doi 10.1007/s10577-007-1212-4.

823   52.   Marley K, Maier CS, Helfand SC. Phosphotyrosine enrichment identifies focal adhesion

824   kinase and other tyrosine kinases for targeting in canine hemangiosarcoma. Vet Comp

825   Oncol **2012**;10(3):214-22 doi 10.1111/j.1476-5829.2012.00325.x.

826   53.   Mertens F, Johansson B, Fioretos T, Mitelman F. The emerging complexity of gene

827   fusions in cancer. Nat Rev Cancer **2015**;15(6):371-81 doi 10.1038/nrc3947.

828   54.   Yoshihara K, Wang Q, Torres-Garcia W, Zheng S, Vegesna R, Kim H*, et al.* The

829   landscape and therapeutic relevance of cancer-associated transcript fusions. Oncogene

830   **2015**;34(37):4845-54 doi 10.1038/onc.2014.406.

831   55.   Mitelman F, Johansson B, Mertens F. 2020 Mitelman Database Chromosome Aberrations

832   and Gene Fusions in Cancer.   <https://mitelmandatabase.isb-cgc.org/>.

833   56.   Stephens PJ, Greenman CD, Fu B, Yang F, Bignell GR, Mudie LJ*, et al.* Massive

834   genomic rearrangement acquired in a single catastrophic event during cancer

835   development. Cell **2011**;144(1):27-40 doi 10.1016/j.cell.2010.11.055.

836   57.   Anderson ND, de Borja R, Young MD, Fuligni F, Rosic A, Roberts ND*, et al.*

837   Rearrangement bursts generate canonical gene fusions in bone and soft tissue tumors.

838   Science **2018**;361(6405) doi 10.1126/science.aam8419.

839   58.   Lee HK, Chauhan SK, Kay E, Dana R. Flt-1 regulates vascular endothelial cell migration

840   via a protein tyrosine kinase-7-dependent pathway. Blood **2011**;117(21):5762-71 doi

841   10.1182/blood-2010-09-306928.

59.   Kim P, Jia P, Zhao Z. Kinase impact assessment in the landscape of fusion genes that

retain kinase domains: a pan-cancer study. Brief Bioinform **2018**;19(3):450-60 doi

10.1093/bib/bbw127.

60.   Medves S, Demoulin JB. Tyrosine kinase gene fusions in cancer: translating mechanisms

into targeted therapies. J Cell Mol Med **2012**;16(2):237-48 doi 10.1111/j.1582-

4934.2011.01415.x.

61.   Pandey PR, Chatterjee B, Olanich ME, Khan J, Miettinen MM, Hewitt SM*, et al.* PAX3-

FOXO1 is essential for tumour initiation and maintenance but not recurrence in a human

myoblast model of rhabdomyosarcoma. J Pathol **2017**;241(5):626-37 doi

10.1002/path.4867.

62.   Selfe JL, Shipley J. Fusion gene addiction: can tumours be forced to give up the habit? J

Pathol **2017**;242(3):263-6 doi 10.1002/path.4902.

63.   Giacomini CP, Sun S, Varma S, Shain AH, Giacomini MM, Balagtas J*, et al.* Breakpoint

analysis of transcriptional and genomic profiles uncovers novel gene fusions spanning

multiple human cancer types. PLoS Genet **2013**;9(4):e1003464 doi

10.1371/journal.pgen.1003464.

64.   Shimozono N, Jinnin M, Masuzawa M, Masuzawa M, Wang Z, Hirano A*, et al.*

NUP160-SLC43A3 is a novel recurrent fusion oncogene in angiosarcoma. Cancer Res

2015;75(21):4458-65 doi 10.1158/0008-5472.CAN-15-0418.

65.   Gru AA, Becker N, Pfeifer JD. Angiosarcoma of the parotid gland with a t(12;22)

translocation creating a EWSR1-ATF1 fusion: a diagnostic dilemma. J Clin Pathol

2013;66(5):452-4 doi 10.1136/jclinpath-2012-201433.

66.   Guo T, Zhang L, Chang NE, Singer S, Maki RG, Antonescu CR. Consistent MYC and

865   FLT4 gene amplification in radiation-induced angiosarcoma but not in other radiation-

866   associated atypical vascular lesions. Genes Chromosomes Cancer **2011**;50(1):25-33 doi

867   10.1002/gcc.20827.

868   67.   Cornejo KM, Deng A, Wu H, Cosar EF, Khan A, St Cyr M, *et al.* The utility of MYC and

869   FLT4 in the diagnosis and treatment of postradiation atypical vascular lesion and

870   angiosarcoma of the breast. Hum Pathol **2015**;46(6):868-75 doi

871   10.1016/j.humpath.2015.02.014.

872   68.   Beca F, Krings G, Chen YY, Hosfield EM, Vohra P, Sibley RK, *et al.* Primary mammary

873   angiosarcomas harbor frequent mutations in KDR and PIK3CA and show evidence of

874   distinct pathogenesis. Mod Pathol **2020**;33(8):1518-26 doi 10.1038/s41379-020-0511-6.

875   69.   Chan JY, Lim JQ, Yeong J, Ravi V, Guan P, Boot A, *et al.* Multiomic analysis and

876   immunoprofiling reveal distinct subtypes of human angiosarcoma. J Clin Invest

877   **2020**;130(11):5833-46 doi 10.1172/JCI139080.

878   70.   Florou V, Rosenberg AE, Wieder E, Komanduri KV, Kolonias D, Uduman M, *et al.*

879   Angiosarcoma patients treated with immune checkpoint inhibitors: a case series of seven

880   patients from a single institution. J Immunother Cancer **2019**;7(1):213 doi

881   10.1186/s40425-019-0689-7.

882   71.   Dancsok AR, Gao D, Lee AF, Steigen SE, Blay JY, Thomas DM, *et al.* Tumor-associated

883   macrophages and macrophage-related immune checkpoint expression in sarcomas.

884   Oncoimmunology **2020**;9(1):1747340 doi 10.1080/2162402X.2020.1747340.

885   72.   Wang G, Wu M, Maloneyhuss MA, Wojcik J, Durham AC, Mason NJ*, et al.* Actionable

886   mutations in canine hemangiosarcoma. PLoS One **2017**;12(11):e0188667 doi

887   10.1371/journal.pone.0188667.

Table 1. Putative fusion genes identified in transcriptomic data of human ASs

Patient ID	Gene 1	Gene 2	Putative fusion gene	Gene 1 chromosome	Gene 2 chromosome	Gene 1 fusion location	Gene 2 fusion location	Fusion Type	Gene 1 Ensembl ID	Gene 2 Ensembl ID	Genomic break position in gene 1	Genomic break position in gene 2	Somatic variants
Patient 1	-	-	-	-	-	-	-	-	-	-	-	-	-
Patient 2	-	-	-	-	-	-	-	-	-	-	-	-	-
Patient 3	VKORC1L1	STEAP1B	VKORC1L1-STEAP1B	7	7	coding	coding	Intra-chromosomal	ENSG00000196715	ENSG00000105889	65873565	22419836	-
	PPP1R13B	ATP5MPL	PPP1R13B-ATP5MPL	14	14	coding	coding	Intra-chromosomal	ENSG00000088808	ENSG00000156411	103847299	103915189	
Patient 4	-	-	-	-	-	-	-	-	-	-	-	-	-
Patient 5	-	-	-	-	-	-	-	-	-	-	-	-	HRAS
Patient 6	-	-	-	-	-	-	-	-	-	-	-	-	-
Patient 7	SMURF1	TMEM139	SMURF1-TMEM139	7	7	coding	intron	Intra-chromosomal	ENSG00000198742	ENSG00000178826	99143726	143286424	TP53
	AGO2	TRAPPC9	AGO2-TRAPPC9	8	8	coding	coding	Intra-chromosomal	ENSG00000123908	ENSG00000167632	140635485	140451383	
Patient 8	-	-	-	-	-	-	-	-	-	-	-	-	NRAS
Patient 9	PEMT	ANKRD6	PEMT-ANKRD6	17	6	coding	utr5p	Inter-chromosomal	ENSG00000133027	ENSG00000135299	17577027	89433375	-
	CPD	NSRP1	CPD-NSRP1	17	17	coding	coding	Intra-chromosomal	ENSG00000108582	ENSG00000126653	30461311	30172542	
Patient 10	SCLT1	NIPBL	SCLT1-NIPBL	4	5	coding	coding	Inter-chromosomal	ENSG00000151466	ENSG00000164190	128888679	37057186	-
Patient 11	TEX2	PECAM1	TEX2-PECAM1	17	17	utr5p	utr5p	Intra-chromosomal	ENSG00000136478	ENSG00000261371	64263168	64390763	-
Patient 12	ATP8A2	FLT1	ATP8A2-FLT1	13	13	coding	coding	Intra-chromosomal	ENSG00000132932	ENSG00000102755	25968679	28357685	-
Patient 13	IRF9	THTPA	IRF9-THTPA	14	14	coding	coding	Intra-chromosomal	ENSG00000213928	ENSG00000259431	24164134	23558695	TP53

Table 2. Putative fusion genes identified in transcriptomic data of canine HSAs

Dog sample ID	Gene 1	Gene 2	Putative fusion gene	Gene 1 chromosome	Gene 2 chromosome	Gene 1 fusion location	Gene 2 fusion location	Fusion type	Gene 1 Ensembl ID	Gene 2 Ensembl ID	Genomic break position in gene 1	Genomic break position in gene 2	Somatic variants	
CHAD-B7	PREX2	LPCAT1	PREX2-LPCAT1	29	34	coding	coding	Inter-chromosomal	ENSCAFG00000007620	ENSCAFG00000010491	17653781	11166324	TP53	-
DHSA-1204	AP4E1	BAIAP2	AP4E1-BAIAP2	30	9	coding	coding	Inter-chromosomal	ENSCAFG00000015318	ENSCAFG00000005700	16748493	968509	TP53	-
	NOL10	PTPRB	NOL10-PTPRB	17	10	coding	coding	Inter-chromosomal	ENSCAFG00000003435	ENSCAFG00000000446	7515966	12372230		
DHSA-0906	MYO16	PTK2	MYO16-PTK2	22	13	coding	coding	Inter-chromosomal	ENSCAFG00000006050	ENSCAFG00000001217	57750807	35397284	TP53	-
	ATP9A	SNX5	ATP9A-SNX5	24	24	coding	coding	Intra-chromosomal	ENSCAFG00000011659	ENSCAFG00000005463	37856900	5096627		
DHSA1101	GABRA3	FLT1	GABRA3-FLT1	X	25	utr5p	coding	Inter-chromosomal	ENSCAFG00000019161	ENSCAFG00000006701	120379631	11232377	TP53	-
DHSA1407	ANKH	ATG16L1	ANKH-ATG16L1	4	25	coding	coding	Inter-chromosomal	ENSCAFG00000014290	ENSCAFG00000011752	88259666	44761299	-	-
DHSA1416	LAMB1	CBLB	LAMB1-CBLB	18	33	coding	coding	Inter-chromosomal	ENSCAFG00000025057	ENSCAFG00000009793	12675054	11259031	TP53	PIK3CA
DHSA1513	PIK3AP1	REV3L	PIK3AP1-REV3L	28	12	coding	coding	Inter-chromosomal	ENSCAFG00000008880	ENSCAFG00000003942	10085967	67953124	TP53	PIK3CA
	AKT3	XPNPEP1	AKT3-XPNPEP1	7	28	coding	coding	Inter-chromosomal	ENSCAFG00000015806	ENSCAFG00000010661	34778111	21321899		
DHSA-1015	MRPS35	CACNA1C	MRPS35-CACNA1C	27	27	coding	coding	Intra-chromosomal	ENSCAFG00000010963	ENSCAFG00000016051	20003244	44487698	-	PIK3CA
	CCDC172	ABLIM1	CCDC172-ABLIM1	28	28	coding	coding	Intra-chromosomal	ENSCAFG00000011803	ENSCAFG00000011513	26980198	25377079		
DHSA-0803	COPS5	NCOA2	COPS5-NCOA2	29	29	utr5p	utr5p	Intra-chromosomal	ENSCAFG00000007386	ENSCAFG00000007775	16578975	19448341	TP53	PIK3CA
DHSA-0805	MSN	LUC7L2	MSN-LUC7L2	X	16	coding	coding	Inter-chromosomal	ENSCAFG00000016607	ENSCAFG00000004069	50748329	9413652	TP53	PIK3CA
DHSA-1113	RANBP3L	ARL15	RANBP3L-ARL15	4	4	coding	coding	Intra-chromosomal	ENSCAFG00000018711	ENSCAFG00000018394	72325076	61484049	-	-

Figures

Figure 1. Identification of novel putative protein-coding fusion transcripts in human AS and

canine HSA. The *Circos* plots visualize fusion genes identified in human ASs (**A**, $n = 13$) and

canine HSA (**B**, $n = 76$). Bar graphs show DNA copy number alterations for each fusion partner

gene using publicly available Exome-sequencing data in an independent dataset of human AS

(**C**, $n = 36$; data retrieved from Ref (12)) and using oaCGH in a larger HSA dataset (**D**, $n = 123$;

data retrieved from Ref (19)).

Figure 2. Validation of fusion genes in canine HSA. **A**, *MYO16-PTK2* fusion gene track and

visualization of the breakpoint in UCSC Genome Browser (Canfam3.1). **B**, Sanger sequencing

result of the PCR product for *MYO16-PTK2* fusion gene. **C**, Schematic illustration of the

putative *MYO16-PTK2* fusion gene and designed FISH probes. **D**, Detection of the *MYO16-*

PTK2 fusion gene on primary canine HSA tissue by FISH. In wild type cells an association

between BAC clones 183H20 (red) and 385H13 (green) can be appreciated, both showing

independent localization from 451H13 (aqua). A portion of tumor cells shows a breakage

within 451H13, with one half of that signal associating with 183H20, and independent of the

localization of 385H13 indicating the existence of the *MYO16-PTK2* fusion at the genomic level.

E, Arrows indicate the amplification of the *MYO16-PTK2* fusion gene. **F**, Detection of the

GABRA3-FLT1 fusion gene by FISH. The *GABRA3-FLT1* fusion is identified by break-apart

FISH probes for proximal *FLT1* (clone 363B20; red) at CFA 25 and distal *FLT1* at CFA 25

(clone 235H9; green). Split *FLT1* genes indicate the fusion event identified by single color signal

(white arrows). Dual colors represent the intact *FLT1* gene (grey arrows). **G**, DNA amplification

of *FLT1* gene.

Figure 3. Fusion genes, mutations of somatic variants, and molecular subtypes of human AS (A-

C) and canine HSA (D-F). **A,** 1,237 differentially expressed genes were identified between

human ASs ($n = 13$) and non-malignant tissue samples ($n = 6$) (False Discovery Rate or FDR $P <$

0.05): 490 genes were upregulated and 747 genes were downregulated in ASs. Ten fusion genes

are marked as yellow bars in seven AS samples. Somatic variations in *TP53* ($n = 2$), *NRAS* ($n = $

1), and *HRAS* ($n = 1$) were found in four ASs. **B,** Box plots show gene expression of *PECAM1,*

FLT1, *KDR*, and *FLT4* representing vasculogenic and angiogenic functions in human ASs and

non-malignant tissues (two-tailed Mann-Whitney test). ***, $P < 0.001$. **C**, Bar graphs show the

relative expression of genes in human ASs normalized to the expression of non-malignant

tissues. Six of 20 genes where the *P*-value was less than 0.05 are displayed (two-tailed Welch's

T-test). **D,** Heatmap illustrates 1,477 significant differentially expressed genes among three

subtypes of canine HSA ($n = 76$) (FDR $P < 0.001$; Fold change > 3). Somatic variant analysis

identified mutations in *TP53* ($n = 24$), *PIK3CA* ($n = 16$), *PIK3R1* ($n = 5$), *NRAS* ($n = 4$), and

ARPC1A ($n = 1$). Grey bars indicate unavailable somatic variants data. Fifteen fusion genes are

marked as yellow bars in 11 HSA samples. **E,** Box plots display gene expression of *PECAM1,*

FLT1, *KDR*, and *FLT4* representing vasculogenic and angiogenic functions in subtypes of HSA

and non-malignant hematomas (two-tailed Mann-Whitney test). ****, $P < 0.0001$; ***, $P <$

0.001; **, $P < 0.01$; *, $P < 0.05$. **F,** Bar graphs show the relative expression of genes in canine

HSA normalized to the expression of hematomas. Eighteen of 30 genes where the P-value is less

than 0.05 are displayed (two-tailed Welch's T-test). Heatmaps (**A** and **D**) show unsupervised

hierarchical clustering (average linkage). Heatmap colors display mean-centered fold change

expression following $\log_2$ transformation. Upregulated genes are presented in red and

downregulated genes are shown in green.

Figure 4. Hypothetical model for angiogenic pathogenesis of canine HSA.

Figure 5. Immunohistochemical expression of p53 and AKT proteins in human AS and canine

HSA. **A,** Bar graphs show the number of cases (y-axis) plotted as a function of IHC scores (x-

axis) for staining with anti-p53, anti-phosphorylated (p)-p53 (Ser15), and anti-p-p53 (Ser20)

antibodies in human ASs (Left panel) and canine HSAs (Right panel). **B,** Representative

photomicrographs show IHC staining of p53, p-p53 (Ser15), and p-p53 (Ser20) in human AS

(Left panel) and canine HSA tissues (Right panel). Bar graphs (**C**) for IHC scores of AKT and p-

AKT (Thr308) proteins and representative photomicrographs (**D**) are also displayed for human

ASs and canine HSAs. H&E = hematoxylin and eosin stain. IHC staining (Horseradish

peroxidase and hematoxylin counterstain). 200X magnification.

Figure 1

A   Human Angiosarcoma

Sample ID	Fusion gene
Patient 3	*VKORC1L1-STEAP1B / PPP1R13B-ATP5MPL*
Patient 7	*SMURF1-TMEM139 / AGO2-TRAPPC9*
Patient 9	*PEMT-ANKRD6 / CPD-NSRP1*
Patient 10	*SCLT1-NIPBL*
Patient 11	*TEX2-PECAM1*
Patient 12	*ATP8A2-FLT1*
Patient 13	*IRF9-THTPA*

B   Canine Hemangiosarcoma

Sample ID	Fusion gene
CHAD-B7	*PREX2-LPCAT1*
DHSA-1204	*AP4E1-BAIAP2 / NOL10-PTPRB*
DHSA-0906	*MYO16-PTK2 / ATP9A-SNX5*
DHSA-1101	*GABRA3-FLT1*
DHSA-1407	*ANKH-ATG16L1*
DHSA-1416	*LAMB1-CBLB*
DHSA-1513	*PIK3AP1-REV3L / AKT3-XPNPEP1*
DHSA-1015	*MRPS35-CACNA1C / CCDC172-ABLIM1*
DHSA-0803	*COPS5-NCOA2*
DHSA-0805	*MSN-LUC7L2*
DHSA-1113	*RANBP3L-ARL15*

C

D

Figure 2

GTGAAGGTTTTGAAACTAACATGAACTTAAGTGGCCGAGATGATCCTGGTAGATCAGAAATTGCATCTGAAACTCAAGATAG
AAATGCAAATAACCATGGAATTCAATTACCTAATTCACTATCGAGTGCTATCACTGCTGAAAATGGGAATTCCATCTCAAATG/
AAGGTGAACGGGCTTTGCCATCGATACCAAAGCTGGCCAACAGCGAGAAGCAAGGCGTGCGGACACACGCGGTCTCCGTG
TCAGAAACAGATGATTATGCTGAGATTATAGATGAAGAAGATACCTACACCATGCCGTCAACCAGG

Figure 3

Figure 4

Figure 5