The Cell Wall Integrity MAPK pathway controls actin cytoskeleton assembly during fungal somatic cell fusion.

Antonio Serrano, Hamzeh H. Hammadeh, Natalie Schwarz, Ulrike Brandt and André Fleißner[#]

Institut für Genetik, Technische Universität Braunschweig, Spielmannstraße 7, 38106 Braunschweig, Germany

21                    #Corresponding author

22                    André Fleißner

23                    Institut für Genetik

24            Technische Universität Braunschweig

25                    Spielmannstraße 7

26                    38106 Braunschweig

27                    Germany

28                  Tel: +49-531-3915795

29                  FAX: +49-531-3915765

30                  a.fleissner@tu-bs.de

33                    running title:

34            MAPK regulation of directed growth

36                    key words:

37  MAPK, cell fusion, directed growth, cell polarization, cell communication, *Neurospora*

38                    *crassa*

**Summary statement**

The CWI MAPK MAK-1 pathway controls actin cytoskeleton assembly at the cell tips

through activation of the Rho-GTPase RAC-1 exclusively on somatic cell fusion.

**Abstract**

Somatic cell fusion is widely studied in the filamentous fungus *Neurospora crassa*. The

interaction of genetically identical germlings is mediated by a signaling mechanism in

which the cells take turns in signal-sending and receiving. The switch between these

physiological states is represented by the alternating membrane recruitment of the SO

protein and the MAPK MAK-2. This dialog-like behavior is observed until the cells

establish physical contact, when the cell-wall-integrity MAK-1 is recruited to the contact

area to control the final steps of the cell fusion process. This work revealed, for the first-

time, an additional MAK-1-function during the tropic growth phase. Specific inhibition of

MAK-1 during tropic-growth resulted in disassembly of the actin-aster, and mislocalization

of SO and MAK-2. Similar defects were observed after the inhibition of the Rho-GTPase

RAC-1, suggesting a functional link between them, being MAK-1 upstream of RAC-1. In

contrast, after inhibition of MAK-2, the actin-aster stayed intact, however, its subcellular

localization became instable within the cell-membrane. Together these observations led

to a new working model, in which MAK-1 promotes the formation and stability of the actin-

aster, while MAK-2 controls its positionning and cell growth directionality.

**Introduction**

Cell-cell fusion (also known as cell fusion) is fundamental for the development of Eukaryotic organisms. It plays an important role in the asexual development of organisms (so-called somatic cell fusion), in which the cells fuse to create organs (e.g. placenta), different cell types (e.g. osteoclasts) or networks (e.g. fungal mycelia) and in the sexual cycle of haploid cells. Both processes are studied in many different model organisms or cell types (e.g. monocytes, myoblasts, yeast etc) and, interestingly, all share very similar and defined stages. These are: 1) the induction of a competence for cell fusion, in which the cells become potential fusion partners by activating specific developmental programs, 2) the comminment of the two fusing cells, when cells actively express signals and cognate receptors to establish a stable communication that induce cell polarization of both partners and, finally, 3) cell fusion, at which the two cells control the correct merging of the plasma membranes and connection of both cytoplasms (Aguilar et al., 2013). In sexual reproduction this step will continue with the fusion of the genomes and the differentiation of a new organism, while in somatic cells new rounds of cell fusion will form, in some cases, a giant syncitia or, like in filamentous fungi, an interconnected cellular network.

In both types of cell fusion, cells need to respond to a chemoattractant (e.g. pheromones in sexual fusion) and either migrate (chemotaxis) or growth (chemotropism) toward the highest  gradient. The correct establishment of cell polarization is essential for the chemotropic response. In fungi, the polarized growth of hyphal cells to chemoattractants is mediated through a highly regulated machinery which involves the activation of mitogen-activated protein kinases (MAPKs) and spatial focalization of polarity core

factors, such as formins, or Rho-GTPases (Aguilar et al., 2013; Berepiki et al., 2010;

Lichius et al., 2014; Martin and Arkowitz, 2014). The concentration of formin-created actin

at the cell tip is also known as a formin-nucleated actin aster (Dudin et al., 2015; Dudin

et al., 2017). In fission yeast, formation of the actin aster is controlled by the formin Fus1

and type V myosins. The formin is needed for nucleation of the actin aster, and type V

myosins are essential for the release of cell wall hydrolases, which degrade the cell wall

between the interacting cells and allow the culmination of the cell fusion process (Dudin

et al., 2015). In budding yeast, actin aster and cable assembly are essential for

recruitment of the MAPK Fus3p (Qi and Elion, 2005). In addition, Fus3p positively

regulates actin cable assembly through phosphorylation of the formin bni1p (Matheos et

al., 2004), yielding a regulatory feedback loop that increases the phosphorylation of

Fus3p and therefore the response to the pheromone signal. In filamentous fungi, actin is

essential for all types of growth (vegetative growth or cell fusion). Disruption of actin

during hyphal extension leads to rapid tip swelling, indicating an important function of

actin into polarized growth (Berepiki et al., 2010). A highly dynamic actin aster is also

observed in *Neurospora crassa* cells undergoing cell fusion, and disruption of actin also

arrests the growth of the interacting cells (Berepiki et al., 2011; Berepiki et al., 2010).

Actin is, overall, the main component of the polarization during cell fusion although its

implications on the process had not been yet elucidated. Two of the main regulators of

actin assembly during fungal growth, the Rho-GTPase RAC-1 and CDC-42, had been

extensively studied in *N. crassa*, showing an independent and exclusive role of RAC-1

for cell fusion and CDC-42 for vegetative growth (Lichius et al., 2014).

The cell fusion process in *N. crassa* is characterized by the alternating oscillatory recruitment of the MAPK MAK-2 (homolog to Fus3p), and the protein SO, to the opposite tips of the interacting cells. The recruitment of each factors is hypothesized to represent two physiological states on the cells: signal sending and signal receiving, with two very different developmental programs (Fleissner et al., 2009; Goryachev et al., 2012; Serrano et al., 2017). During the cell fusion interaction, the cells switch between the two states, which relies on switching of the recruitment of MAK-2 and SO. After cells establish physical contact, the growth is arrested and cell wall and plasma membrane remodeled to allow the complete fusion and mixing of the cytoplasm. A second MAPK, the cell-wall-integrity (CWI) MAK-1, is only recruited after the cells establish physical contact, and the chemical inhibition of an analog-sensitive variant results in the disruption of the cell merger, suggesting that the kinase activity is essential for the process (Weichert et al., 2016). Interestingly, MAK-1 as well as MAK-2 and SO are essential for the process and the mutation of their respective genes results in the complete absent of cell fusion (Fleissner et al., 2005; Fu et al., 2011; Pandey et al., 2004). This suggests that MAK-1 (besides MAK-2 and SO) is essential for the establishing of a competence state on the cells for cell fusion. However, no conclusive data have shown whether MAK-1 is needed for the communication phase of the cell fusion process, which is featured by the oscillatory recruitment of MAK-2 and SO and the release and sensing of the so-far unknown fusion signal.

In this study, we will address the question on MAK-1 functionality during the communication phase of the cell fusion process and the implications of MAK-2 regulation. We will follow a chemical genetics strategy in order to answer these questions.

## Results

### *MAK-1 kinase-activity inhibition results in disruption of the cell fusion process*

The CWI MAPK MAK-1 plays important functions for the sucessful completion of the cell fusion process. This kinase plays an unknown essential role during the establishment of the cell fusion competence and, in addition, its activity is crucial for the growt arrest of the fusing cells after establishing physical contact (Fu et al., 2011; Weichert et al., 2016). While these data indicate that MAK-1 is required for the onset of the fusion process and after cell-cell contact, it remains unclear if this MAPK also plays a role during the tropic interaction of the fusion partners. Since the deletion of the *mak-1* gene results in the complete disruption of the cell fusion competence (Fu et al., 2011), we decided to turn into the creation of an analog-sensitive version, also known as *Shokat* allele (Bishop et al., 2000), of *mak-1* to manipulate its activity in a spatio-temporal manner. For that purpose, the point mutation that renders the encoded kinase analog-sensitive was integrated at the original gene locus. The gatekeeper amino acid was identified in a previous study (glutamine at position 104) and exchanged by glycine, resulting in the *mak-1$^{E104G}$* gene which encoded the analog-sensitive kinase (Weichert et al., 2016). The plasmid 738 containing the *mak-1$^{E104G}$-hph* construct (details on Mats and method section), was used as a template to amplify the full-length knock-in fragment by PCR (using primers 1304 and 1307), which was transformed into the *N. crassa* strain N1-06 (FGSC 9719) (Fig. S1A). This recipient strain possesses a mutation in the gene *mus52* (*mutagen sensitive 52*), whose gene product is required for DNA repair by nonhomologous end-joining .The mutation results in an increase in the frequency of homologous recombination up to almost 100% (Ninomiya et al., 2004). The resulting

transformants were first tested by PCR by amplifying the *mak-1* locus and the fragment

was sequenced to confirm the base pair exchange (E104G; CAG to GGC). The proper

integration of the transforming DNA and the absence of additional heterologous

integrations were successfully tested by Southern blot of homokaryotic isolates obtained

from a sexual cross of the primary transformants with the wild-type strain (Fig. S1A and

B). The resulting homokaryotic strain, 849 (similar observations were made in 848, 850

and 851), appeared macroscopically wild type-like, indicating that the mutated kinase is

functional in absence of the inhibitor (ATP-analog, 1-NM-PP1) and supplemented with

1% DMSO (solvent of the 1-NM-PP1) as a control (Fig. S1C). In contrast, addition of the

inhibitor (40 µM final concentration) to the growth medium resulted in a Δ*mak-1*-like

phenotype, whilst the wild-type control strain remained unaffected (Fig. S1C).

Microscopically, the presence of the inhibitor drastically reduced the cell fusion rate of the

MAK-1$^{E104G}$ strain, proving that it can be inhibited, and allowing us to analyze the functions

of MAK-1 at specific stages of the cell fusion process, such as the tropic growth phase

(Fig. S1D). Additionally and as previously shown in other publications, neither the solvent

DMSO nor the ATP-analog have any effect on wild-type cells, proving its specificity to

analog-sensitive variants (Fig. S1C and D).

***Inhibition of MAK-1 during the tropic growth phase results in disruption of MAK-***

***2/SO and actin cytoskeleton***

In order to test the consequences of MAK-1 inhibition during the cell-cell communication

steps, multiple strains were created. The most important proteins involved in the cell

fusion process are MAK-2 and SO, so we decided to first test the localization of these two

after MAK-1 inhibition. The strain carrying the *mak-1::mak-1$^{E104G}$-hph* (849) genotype was

independently crossed with the strains *his-3::Ptef-1-mak-2-gfp* (665) and *his-3::Ptef-1-so-*

*gfp* (714), resulting in strains 865 and 874, respectively. Both strains were tested by PCR

for integration of the *mak-1::mak-1$^{E104G}$-hph* locus and by fluorescence microscopy for

each GFP construct integration. Microscopically, all strains showed comparable MAK-1

inhibition phenotypes at the presence of 1-NM-PP1 (data not shown). When cell-cell

communication was observed with the microscope, the addition of DMSO did not affect

the normal oscillatory recruitment of MAK-2 (Fig. 1A). However, addition of the kinase

inhibitor completely disrupted the cell-cell communication process, and the usual dynamic

localization of MAK-2 was interrupted. The kinase disappered from the membrane and

no detachment was observed after almost 18 minutes (Fig. 1B). Similarly, addition of

DMSO did not affect the usual oscillatory recruitment of SO (Fig. 1C). However, 10

minutes after addition of the kinase inhibitor, SO was accumulated at the membrane in

both cell tips at the same time, and the GFP signals were more widely distributed around

the cell tips (Fig. 1D). Taken together these data indicate that MAK-1$^{E104G}$ activity is

essential for the tropic growth phase and for proper MAK-2 and SO dynamics. As

previously described, cell fusion is observed at the early stage of colony establishment

between germinated and ungerminated spores, but also later in development between

hyphae in the inner part of the fungal colony in a process known as hyphal fusion.

In previous work, we showed that the characteristic MAK-2/SO dynamics also occur

during hyphal fusion (Serrano et al., 2018). In order to test whether MAK-1 also plays a

role during hyphal fusion, we tested the consequences of MAK-1 inhibition by microscopic

analysis of hyphal fusion events. With DMSO, MAK-2 oscillated in a wild-type manner

(Fig. S2A). When MAK-1$^{E104G}$ was inhibited, MAK-2 showed a disruption of their dynamics

similar to the findings in germlings (Fig S2B). When looking to SO, DMSO did not affect

the usual oscillatory dynamics (Fig. S2C), but addition of the inhibitor completely disrupt

the process (Fig. S2D). While MAK-2 completely vanished from the hyphal tip, SO

remained as single dots distributed widely around the plasma membrane of both

interacting hyphae. These data indicate that the function played by MAK-1 during

germling fusion is also conserved in hyphal fusion.

In fungi, proper actin organization is crucial for polarized hyphal extension. In *N. crassa*,

Berepiki *et al* adapted a method for indirect actin visualization: the actin-reporter life-act

linked to GFP (Berepiki et al., 2010). Lifeact is a small peptide (17 amino-acid), which

represents the N-terminus of the actin-binding protein Abp140 from *S. cerevisiae* (Riedl

et al., 2008). In non-interacting cells of *N. crassa* expressing this reporter, actin is mainly

found at the tips of growing germ tubes, in the form of actin patches localized around the

cell tips and actin cables. During cell fusion, disruption of actin assembly results in failure

of polarized, directed growth of the fusion tips, while microtubules are dispensable for this

process (Roca et al., 2010). Additionally, in *S. pombe* mating fusion, actin also

accumulates at the Shmoo of the interacting cells in a structure termed actin fusion aster

complex. This accumulation has been shown to mediate the release of cell wall degrading

enzymes involved in fusion pore formation (Dudin et al., 2015). All these data indicate

that actin is essential for tropic growth and cell fusion, but its regulation remains mostly

unknown. In order to test whether MAK-1 influences actin assembly and dynamics, we

created a strain carrying the *mak-1::mak-1$^{E104G}$-hph* construct together with the *his-*

*3::Ptef-1-lifeact-gfp* expression cassette, by crossing of strains 849 and 754, respectively,

resulting in strain 869. This isolate behaved as wild type in the presence of DMSO, and

typical actin cables were present at the cell tips during the communication phase and

during cell merger comparable to wild-type cells. To our surprise, actin dynamics showed

a comparable oscillatory and alternating accumulation at the cell tips of the interacting

cells as MAK-2 and SO  (Fig. 1E). When MAK-1$^{E104G}$ was inhibited by addition of the

kinase inhibitor, actin accumulation was clearly affected. Interestingly, the accumulation

of actin cables from the tips disappeared. In addition, actin patches were not focused

anymore around the cell tips, but localized in a wider distribution around the cell

membrane of the entire cell body (Fig. 1F). As expected, actin oscillation was also

observed in hyphal fusion (Fig. S2E). Inhibition of MAK-1$^{E104G}$ also disrupted actin

accumulation at the hyphal tips as observed in germling fusion (Fig. S2F).

These data suggest for the first time that MAK-1 might be involved in the regulation of the

actin cytoskeleton during the cell fusion process. An alternative explanation might be that

the actin misregulation represents an indirect effect of the disruption of the cell fusion

process in the cells. This will be addressed in the following sections.

***Actin accumulation at the cell tips oscillates out-of-phase of SO/MAK-2 usual***

***dynamics***

Surprisingly, the observation of actin localization in the MAK-1 inhibitable strain in the

presence of DMSO suggested that accumulation of actin occurs at the cell tips of the

interacting cells and that this accumulation oscillates in an antiphase manner. To further

decipher whether it was a specific effect of this strain, we decided to observe the

localization of actin on wild-type cells. By using the strain 754 (Berepiki et al., 2010) in

which the construct *lifeact-gfp* is transformed into the *his-3* locus in the wild-type

background, we made similar observations on actin accumulation at the cell tips and its

oscillation in an antiphase manner (Fig. 2A). Based on these data, we hypothesized that

actin accumulation is likely observed together with SO, since the presence of SO at the

cell tips is associated with a signal sending state of the fusing cell (Fleißner and Serrano,

2016; Goryachev et al., 2012). We therefore created a strain (H99) expressing both genes

in the *his-3* locus, lifeact-GFP and SO-mCherry, to allow observation of actin and SO

within the same cell.  When interacting cell pairs were observed, a dynamic interferes co-

localization was found, in which SO was recruited to the membrane at the tip of the first

pair while actin was recruited in the second interacting cell. After a few minutes, both

proteins co-localize at the tip of the first partner. Lastly, actin disassembled form the

membrane of the upper cell and was recruited in the lower cell, while SO stayed at the tip

of the upper cell. (Fig. 2B). This indicates that actin polymerization at the cell tips is out-

of-phase of the usual dynamics of MAK-2/SO. To test whether actin oscillates during

vegetative growth, germinated spores of strain 754 (expressing lifeact-gfp in the WT

background) were observed under the fluorescence microscope. Actin was localized to

the cell tips of growing cells as previously described (Berepiki et al., 2010) and no

oscillation was observed (Fig 2C), indicating that oscillatory accumulation of actin is

exclusive of the cell fusion growth.

***Inhibition of MAK-2 results in disruption of the usual MAK-2/SO and actin***

***localization during cell fusion***

Based on the previous results, we hypothesized that MAK-1 may be directly involved in

actin polimerization or that actin misregulation may be an indirect effect of the disruption

of the cell fusion process caused by MAK-1 inhibition. To test this hypothesis, we decided

to follow a similar strategy by creating an analog-sensitive version of MAK-2 to decipher

SO and actin dynamics during the tropic growth phase.

Similar to the design of the *mak-1$^{E104G}$-hph* construct, a *mak-2$^{Q100A}$-hph* gene knock-in

cassette was created and integrated at the original gene locus of strain N1-06 (Δ*mus-52*).

However, this construct proved to be not functional and the respective transformants

exhibited a Δ*mak-2*-like phenotype (data not shown). In previous studies, we found that

the Δ*mak-2* phenotype of the knock-out mutant was only fully complemented when the

analog-sensitive version was overexpressed by using strong constitutive promoters such

as *Pccg-1* or *Ptef-1* (Fleissner et al., 2009; Serrano et al., 2018). We therefore decided to

replace the promotor at the original gene locus by the P*tef-1* sequence. The linear *Ptef-*

*1-mak-2$^{Q100A}$-hph* knock-in cassette (details on Mats and Methods section) amplified from

the plasmid 846 with primers 1487 and 1488 was transformed into N1-05 (Δ*mus-52* like

N1-06, but different mating type) resulting in strain 882 (*Pmak-2-mak-2::Ptef-1-mak-*

*2$^{Q100A}$-hph*) (Fig. S3A). The primary transformants were tested by PCR of the *mak-2* locus

and sequencing of the resulting fragment. In order to obtain a homokaryotic strain, a

sexual cross was performed between wild type *mat a* (N1-02) and one of the primary

transformants, 882 (*mat A*). Four of the resulting progenies were tested by Southern blot

analysis for single integration of our construct. All four isolates showed clear bands

corresponding to the *Ptef-1-mak-2$^{Q100A}$* radioactive-labelled fragment, while it was absent

in the recipient strain (N1-05) (Fig. S3A and B). The generated strains showed a

macroscopic phenotype like wild type in the absent of the kinase inhibitor (plus DMSO)

and a Δ*mak-2*-like phenotype when it was present (Fig. S3C). Microscopically, cell fusion

rates  were not fully comparable to the wild-type strain in the present of DMSO, but cell

fusion was still observed. In contrast, cell fusion rate was comparable to Δ*mak-2* when

the kinase inhibitor was added (Fig. S3D). These data indicate that the strain is functional

for cell fusion in the absent of the kinase inhibitor and allow us to analyze the cell

dynamics when MAK-2$^{Q100A}$ is inhibited in a spatio-temporal manner.

To test the effect of MAK-2$^{Q100A}$ inhibition on the localization of SO and actin, we crossed

the strain *Pmak-2-mak-2::Ptef-1-mak-2$^{Q100A}$-hph* (899) independently with *his-3::Ptef-1-*

*so-gfp* (714) and *his-3::Ptef-1-lifeact-gfp* (754), generating the strains *his-3::Ptef-1-so-gfp;*

*Pmak-2-mak-2::Ptef-1-mak-2$^{Q100A}$-hph* (912) and *his-3::Ptef-1-lifeact-gfp; Pmak-2-mak-*

*2::Ptef-1-mak-2$^{Q100A}$-hph* (915), respectively. When cell fusion was observed with DMSO,

the dynamic localization of SO was wild-type-like and no significant differences to the

reference strain were observed (Fig. 3A). When the kinase inhibitor was added, the

oscillation rapidly ceased (after around 4 minutes) and, in 10/15 observations, the

localization of SO remained widely distributed around the cell membrane (Fig. S4A).

Interestingly, in 5/15 observations, the localization of SO switched to another area of the

cell, suggesting that the complex formed at the cell tips may travel together around the

cell-membrane after MAK-2 inhibition (Fig. 3B). Regarding actin localization, fusing cell

pairs were observed under the microscope with DMSO, showing the already described

oscillatory accumulation of actin to the cell tips (Fig. 3C). In the presence of the kinase

inhibitor, the actin localization switched to a different spot within the cell, in some of the

cases (10/20) travelling to the opposite cell tip, in a similar manner than SO (Fig. 3B).

After 10 minutes, actin was still accumulated at the new spots although the tropic

interaction and directed growth were already terminated (Fig. 3D). We also observed in

some cases (10/20) than actin switched to a different position, but remaining close to the

**14**

cell tip (Fig. S4B). These data clearly indicate that MAK-2 inhibition leads to a slightly

different disruption of the actin localization, suggesting that MAK-1 and MAK-2 play

different roles on the cytoskeleton organization during cell fusion and excluding the

hypothesis of the indirect effect on actin cytoskeleton due to the cell fusion disruption.

***The actin cytoskeleton is essential for the dynamic localization of MAK-2 and SO***

MAK-1$^{E104G}$ inhibition disrupts the proper MAK-2/SO membrane recruitment. At the same

time, actin accumulation at the cell tips is also disturbed. Actin cables serve as tracks for

the transport of multiple cargoes, including potential secretory vesicles, which might be

involved in the cell fusion process (Berepiki et al., 2011; Dudin et al., 2015). Our data

support two alternative hypotheses: either MAK-1 activity is directly involved in both MAK-

2/SO recruitment and actin organization, or it is only involved in actin organization, but

actin disruption results in a disturbed MAK-2/SO localization.

In yeast, Fus3p recruitment is mainly mediated through actin cables organized at the

Shmoo of the mating cells (Qi and Elion, 2005). In order to test whether actin is needed

for proper MAK-2/SO recruitment, we employed the actin-disrupting drug latrunculin A.

This toxin originating from the Red Sea sponge *Latrunculia magnifica* binds to actin

monomers, thereby preventing the formation of F-actin (Coué et al., 1987). In *N. crassa*,

it has been used to demonstrate that actin is essential during the cell fusion process, but

without testing the effect on MAK-2/SO dynamics (Berepiki et al., 2010).

In a pretest we determined that a concentration of 10 µM resulted in the disassociation of

F-actin in germinating spores within 6 minutes (Fig. S4C). When MAK-2 localization was

tested, we observed that actin disruption due to latrunculin A resulted in the dissasimble

of MAK-2 after 8 minutes, and no further MAK-2 membrane recruitment was observed in

the cells (Fig. 4A). Similarly, the dynamic localization and membrane recruitment of SO

were also affected by actin inhibition (Fig. 4B), suggesting that the proper polymerization

of actin at the cell tips is essential for membrane recruitment of MAK-2 and SO. To further

understand the implications of MAK-1/MAK-2 in this process, we decided to investigate

the roles of the upstream regulators of actin polymerization, the Rho-GTPases.

Rho GTPases are signaling G proteins conserved in all eukaryotic organisms that

regulate intracellular actin dynamics. These proteins cycle between an active (GTP-

bound) and an inactive (GDP-bound) conformation (Martin and Arkowitz, 2014). Upon

activation, the Rho GTPases translocate from the cytosol to the plasma membrane,

where they interact with actin polymerization proteins, such as formins, that regulates the

formation of actin cables (Berepiki et al., 2011).

In *N. crassa*, the Rho-type GTPase RAC-1 is essential for cell fusion, while the Rho-type

GTPase CDC-42 plays a role in germination and normal polarized germ tube growth. Both

proteins localize to the tips of growing cells (Lichius et al., 2014). To differentiate the

functions of both Rho-type GTPases, the specific RAC-1 inhibitor NSC3766 was

employed in an earlier study. The drug only affects RAC-1 but not CDC42 and had already

been used and proved functional in *N. crassa* (Lichius et al., 2014). In our experiments,

we tested if the specific inhibition (with NSC3766) of RAC-1 would affect the dynamics of

actin and the recruitment of MAK-2, SO and actin assembly. In interacting cell pairs

expressing MAK-2-GFP, the protein was recruited to the membrane in a wild-type manner

before addition of NSC3766. The addition of NSC23766 at a concentration of 100 µM

resulted in defects in the recruitment of MAK-2 after 12 minutes. MAK-2 fully disappeared

from the membranes of both cells and the interaction ended (Fig. 4C). When SO dynamics

were tested, the usual membrane accumulation at the cell tips was observed. Like the

previous experiment, defects in the localization of SO were observed 12 minutes after the

addition of the RAC-1-inhibitor. However, in contrast to MAK-2, SO remained widely

distributed at the membranes of both cells although the interaction had ended (Fig. 4D).

This mislocalization of SO was comparable to the one observed after MAK-1 inhibition

(actin patches distributed all around the cell membrane), suggesting that both proteins

(MAK-1 & RAC-1) may play functions within the same pathway.

Actin localization was also tested in the presence of the RAC-1 inhibitor. When the

inhibitor was added, the number of actin cables declined and actin patches were

distributed more widely around the plasma membrane. After 12 minutes, actin cables fully

disappeared and directed cell growth halted (Fig. S4D). All these data indicate that the

proper polymerization of actin at the cell tips of the interacting cells, through activation of

the Rho GTPase RAC-1, is essential for the dynamic localization of MAK-2 and SO.

***MAK-1 activity is essential for activation of RAC-1 while MAK-2 activity is important***

***for its positioning***

All previous experiments suggest a potential connection between MAK-1 activity and

RAC-1 activity. Both proteins are essential for actin assembly during cell fusion, and

similar defects are observed when each one is inhibited. In contrast, MAK-2 inhibition

disrupts the cell communication process although the actin aster becomes unstable within

the cell. Together, these data support two potential hypotheses: MAK-1 and RAC-1

function in the same pathway (either upstream/downstream), which controls actin

**17**

polymerization, or RAC-1 and MAK-1 regulate, independently, a common target that is

essential for actin assembly (e.g. the formin BNI-1).

To test these hypotheses, we decided to analyze active RAC-1 dynamics after MAK-1

and MAK-2 inhibition through visualization by the Cdc42-Rac-interacting-binding (CRIB)

reporter (see below). If our first potential hypothesis is correct, the outcome of the

experiment will show that MAK-1 inhibition results in inactivation of RAC-1 and therefore

membrane detachment of the reporter since Rho-GTPases are membrane-recruited only

when are activated (Araujo-Palomares et al., 2011; Corvest et al., 2013). If the second

hypothesis is correct, we should observe that MAK-1 inhibition does not have any effect

on RAC-1 activation, and therefore the reporter should remain activated and attached at

the plasma membrane, indicating that MAK-1 and RAC-1 control independently the same

factor, that is involved in actin polymerization during cell fusion.

The use of CRIB reporters in *N. crassa* was established by Lichius and colleagues. They

created a GFP-labelled reporter that binds as a GEF protein to activated Rho-GTPases

without disrupting their function, thereby allowing to localize both activated RAC-1 and

CDC-42 at the same time (Lichius et al., 2014). The use of the RAC-1 inhibitor allowed

them to distinguish the functions of both proteins. The CLA-4 CRIB reporter  strain

(*Pccg1::crib^cla-4-gfp::bar*+, 927) was crossed with the MAK-1$^{E104G}$ strain (849), resulting

in strain 943 (*mak-1::mak-1$^{E104G}$-hph; Pal1-crib$^{cla-4}$-bar*). As a control we first tested the

already reported effects of RAC-1 inhibition on the CRIB reporter. As previously reported,

the addition of the RAC-1-specific inhibitor NSC23766 to interacting cells resulted in

vanishing of the reporter from the apical tip and cell growth arrest (Fig. S5A). The

complete disappearance of the reporter from the tips of the interacting cells indicates that

only RAC-1 and not CDC-42 is present at the growth cone of the fusing cells, since CDC-

42 is not affected by the RAC-1 inhibitor.

In our control, we observed as already shown that the CRIB reporter was present

permanently in both cell tips during fusion (Fig. 5A) (Lichius et al., 2014). When in the

next step, MAK-1 was inhibited, the CRIB reporter disappeared from the plasma

membrane and cell growth was arrested (Fig. 5B). Interestingly, both RAC-1 and MAK-1

inhibition showed a similar effect on the CRIB reporters, suggesting that likely the

inhibition of MAK-1 specifically affect RAC-1 activity, therefore disrupting the actin

assembly, membrane recruitment of MAK-2 and SO and the general fusion process.

Consistent with our hypothesis, MAK-2 inhibition should not affect the activity of RAC-1,

but rather its localization, as we have observed with the actin in previous experiments. To

test this, the CRIB reporter strain (927) was crossed with MAK-2$^{Q100A}$ (899), resulting in

strain 949 (*Pmak-2-mak-2::Ptef-1-mak-2$^{Q100A}$-hph;Pal1-crib$^{cla-4}$-bar*). Like previously, the

localization of the CRIB reporter stayed permanently at both cell tips during the whole

tropic growth phase in our control (Fig. 5C). When MAK-2 was inhibited, the CRIB reporter

remained recruited to the plasma membrane and its localization appeared to be widely

distributed around the cell tip (Fig. 5D). These data suggest that MAK-2 might be

controlling the localization, and therefore growth directionality, of the Rho-GTPase RAC-

1.

However, we already mentioned that the CRIB reporter allows the localization of the

activated forms of RAC-1 and CDC-42 at the same time (Lichius et al., 2014). To decipher

if RAC-1, and not CDC-42, is switching its position after MAK-2 inhibition, we used the

RAC-1 inhibitor after inhibition of MAK-2. Like the previous experiments, fusing cells were

observed under the microscope and the kinase inhibitor was added. After observing,

again, the switching in the position of the CRIB reporter provoked by the inhibition of MAK-

2, the RAC-1 inhibitor was added. Interestingly, the CRIB reporter fully disappeared from

the plasma membrane, indicating that MAK-2 inhibition of fusing cells affects exclusively

the position of RAC-1 at the plasma membrane (Fig. 5E). These data indicate that MAK-

2 inhibition in tropically growing cells specifically affects the localization of RAC-1 (that

stays active), which remains recruited to the plasma membrane but loses growth

directionality towards the cell fusion partner and changes its position to another area in

the cell.

Together, all these data support a model in which inhibition of MAK-1 specifically affects

RAC-1 membrane recruitment, which then results in disassembly of the actin focus

organized by the Rho-GTPase and arrest growth of the interacting cells.

***MAK-1 exclusively controls actin polymerization during cell fusion***

We have shown that MAK-1 activity is essential for the correct polimerization of actin at

the cell tips of the interacting cells and it raises the question whether MAK-1 functions are

specific for directed growth or if they are also involved in general polar growth. In order to

test this, we decided to observe the effect of the kinase inhibitor and the RAC-1 inhibitor

(as a negative control) independently in non-interacting cells. Under the microscope, we

quantified the length of the germ tube before and after incubation of either the kinase

inhibitor, the RAC-1 inhibitor or DMSO. To ensure that only non-interacting cells were

tested, only germlings with a distance of more than 30 μm to the nearest cell were

analyzed. As a result, no significant differences were observed between the DMSO, the

kinase inhibitor or the RAC-1 inhibitor, supporting the idea that MAK-1 exclusively

regulates actin polimerization and therefore tropic growth during cell fusion (Fig. 6A).

Additionally, we also tested whether MAK-2 had any effect on growth rate by performing

the same experiment and obtaining similar results (Fig. 6A).  Furthemore, we also tested

the localization of the CRIB reporter and actin before and after addition of DMSO, the

kinase inhibitor or the RAC-1 inhibitor. Observations were made as previously described

but using a fluorescence microscope to record the localization of the CRIB reporter or

actin. As expected, no consistent differences were observed in the localization of the

CRIB reporter when the cells were incubated with DMSO or the kinase inhibitor (Fig. 6B)

and with $H_2O$ (solvent of the RAC-1 inhibitor) or the RAC-1 inhibitor (Fig. 6C) in the MAK-

1-inhibitable strains. Similarly, no differences were found in the localization of actin when

cells were treated with DMSO or the kinase inhibitor (Fig. 6D), or with $H_2O$ or the RAC-1

inhibitor (Fig. 6E) in the MAK-2-inhibitable strains. In order to confirm this finding from the

chemical genetics approach, we also studied the actin organization in the *mak-1* knock-

out mutant. The knock-out strain was obtained from the *N. crassa* gene knock-out

collection (NCU09482). As previously reported, the mutant showed a cell fusion-deficient

phenotype (Fu et al., 2011). In order to obtain an actin-labelled Δ*mak-1* strain, a sexual

cross was set up between the strains Δ*mak-1* and 754 (*his-3::Ptef-1-lifeact-gfp*), resulting

in strain 891 (*mak-1::hph;his-3::Ptef-1-lifeact-gfp*). Spores of this isolate were incubated

in MM for 2 hours and actin organization was studied by fluorescence microscopy. In the

wild-type cells, actin is organized at the growing cell tips and actin cables are expanding

from the growing tip (Berepiki et al., 2011). In the Δ*mak-1* mutant cells, germ tube

elongation and actin assembly at the growing cell tips was comparable to wild type (Fig.

S5B).

Together, all these data indicate that MAK-1 exclusively function as a actin polymerization

regulator during the cell fusion process and is not involved in actin polimerization during

general vegetative growth. In other words, MAK-1 is not involved within the same actin-

regulatory pathway than CDC-42.

***MAK-1 functions upstream of RAC-1***

As we have already shown, MAK-1 is essential during the communication phase, when

MAK-2 and SO are recruited to the cell tips in an oscillatory manner. Additionally, MAK-1

is also important for the growth arrest phase, in which the tropic growth is arrested and

cell wall is remodeled to allow the completition of the fusion process. When an analog-

sensitive version of MAK-1 is partially inhibited, the growth arrest and cell merger is

disrupted and fusing cells exhibit an unique phenotype of corkscrew-like structures (also

known a twisting germ tubes) (Weichert et al., 2016). To additionally test whether MAK-1

and RAC-1 are within the same pathway, we decided to analyze the formation of the

twisting germ tubes phenotype after partial inhibition of RAC-1 in wild-type cells, as

indicated in the previous study (Weichert et al., 2016). Interestingly, the partial inhibition

of RAC-1 induced the formation of twisting germ tubes, again indicating that both proteins

function in the same pathway. Three different concentrations were tested (100, 50 and

25 µM), albeit only 25 µM allowed the induction of cell fusion events (Fig. 7A).

Interestingly, a large number of cells exhibit defects in polar growth, which may be related

to the effect of RAC-1 inhibition. To further investigate this, we decided to quantify the

percentage of cells that shows tropic defects (observed as growth-forming circles). First,

we quantify the number of cells that underwent this growth after RAC-1 inhibition in a

medium with $Ca^{2+}$, and about 12.5% of the cells exhitied this growth (compared to the

water control). To show whether this defect may be caused by inhibition of the general

germ tube elongation machinery (not related to cell fusion growth), we performed similar

quantifications on wild-type cells placed on media without $Ca^{2+}$ (its absent inhibits the

induction of cell fusion) and on a cell fusion mutant, such as Δ*so.* We observed that either

in absent of $Ca^{2+}$ or in the cell fusion mutant, the polarity defects were not observed (Fig

7B), suggesting that this phenotype may be caused by partial inhibition of RAC-1 only

when cells are undergoing cell fusion.

Although MAK-1 and RAC-1 functions within the same actin-regulatory pathway during

cell fusion, there are not strong evidences on which is upstream/downstream.

Interestingly, both proteins are localized at the cell tips during the cell growth arrest phase.

We therefore decided to test how localization of MAK-1 and RAC-1 was affected after

inhibition of RAC-1 or MAK-1, respectively. Cells that just established contact were

observed in the microscope and either RAC-1 or the kinase inhibitor added to visualize

the effect on the subcellular localization of MAK-1/RAC-1. When RAC-1 was under

observation, its localization stayed at the contact area of the fusing cells that just made

contact and the addition of DMSO did not have any effect on its localization (Fig. 8A).

However, the full inhibition of MAK-1 in fusing cells induced a rapid mislocalization of

RAC-1, that completely dissappears from the plasma membrane while cells presumibely

stop their cellular interaction (Fig. 8B). Addition of water (solvent of RAC-1 inhibitor) did

not affect the cell fusion event and MAK-1 localization during the last steps of the cell

fusion (Fig. 8C). Interestingly, addition of the RAC-1 inhibitor in interacting cells as well

did not have any significant effect on the localization of MAK-1, that stayed at the contact

area, although cells likely stop their interaction (Fig. 8C).

These data clearly indicate that both MAK-1 and RAC-1 play a role within the same actin-

regulatory pathway, showing for the first-time that the MAPK MAK-1 controls, and

therefore is upstream, activation and membrane-recruitment of RAC-1 during the cell

fusion process.

**Discussion**

The correct actin-cytoskeleton assembly is essential for the somatic cell fusion in *N.*

*crassa*, which provides on outstanding model for studying actin-dynamics and its

regulation. In this study, we show for the first time the molecular insight regulation of the

CWI MAPK MAK-1 on actin-assembly, that happens through the activation of the

downstream Rho-GTPase RAC-1, to control actin assembly during the communication

phase and cell merger. In addition, we described a function of the MAPK MAK-2 on actin

reorganization by controlling RAC-1 positioning likely to the higher gradient intensity of

the unknown fusion signal generated by the opposite fusing cell.

**MAK-1 plays different roles in CWI regulation and cell fusion**

The central and well-described role of the CWI-signaling pathway is to detect and respond

to cell wall stresses that arise in the vegetative life cycle of fungi. In yeast, activation of

this pathway is well understood and has been reviewed several times (Correia et al.,

2010; de Nobel et al., 2000; Levin, 2011).

In filamentous fungi, activation of the pathway is mediated by a group of sensor proteins

named as WSC (cell wall integrity and stress-response component). These glycosylated

transmembrane proteins sense morphological changes induced by cell growth or cell wall

stresses. The signal is transduced to the MAPK cascade by the small Rho-GTPase RhoA

and the protein kinase C (PkcA). This activation yields a fully phosphorylated MpkA MAPK

(homolog to MAK-1) that further activates transcriptional factors mediating the appropriate

cellular responses (Yoshimi et al., 2016). In *N. crassa*, WSC-1 and its homolog WSC-2

activate the cell wall integrity MAK-1 MAPK pathway when the cell wall is stressed.

Deletion of both genes (*wsc-1* and *wsc-2*) results in strong morphological defects during

vegetative growth and colony formation (Maddi et al., 2012), which is very similar to the

observations made in the different mutants of the CWI MAPK MAK-1 pathway (Maerz et

al., 2008; Park et al., 2008; Vogt and Seiler, 2008). In addition to these defects in cell wall

stress response, mutation of every component of the MAK-1 MAPK module also results

in a cell fusion-defective phenotype (Fu et al., 2011). However, while the kinases are

essential to undergo cell fusion, WSC-1 and WSC-2 are dispensable for this process,

indicating that for the cell fusion process MAK-1 is activated independently of the CWI

pathway (Maddi et al., 2012).

Another upstream activator of MAK-1 is the HAM-7 protein. HAM-7 is a GPI-anchored

cell wall protein that is required for activation of MAK-1 (Maddi et al., 2012). Together with

HAM-7, the proteins HAM-6 and HAM-8 are hypothesized to work together in a complex

as a sensor of signals to trigger MAK-1 phosphorylation (Fu et al., 2014). Mutation of

either *ham-6*, *ham-7* or *ham-8* results in a cell fusion defective phenotype, therefore

suggesting that its function might be related to activation of MAK-1 for cell fusion

**25**

purposes. All these clearly suggest and support the idea that MAK-1 is activated by two

different set of sensors: the WSC complex that mediates the signals related to the cell

wall stresses and the HAM-6/HAM-7/HAM-8 sensor which is more associated to the

signals coming from the cell fusion process. In future studies, it will be of interest to

investigate the functions of these three proteins by controlling its expression with inducible

promoters and observe the defects induced in the cell fusion (e.g. twisting germ tubes).

**MAK-1 functions in all of the three main cell fusion steps: competence, signaling**

**and growth arrest**

As shown before, cell fusion comprises several independent steps: first the cells need to

acquire a cell fusion competence. Second, the cells recognize the presence of other cells

and start the tropic growth including the membrane recruitment of MAK-2 and SO. Third,

the cells establish physical contact, their growth is arrested and the cell wall is remodeled

in order to allow the subsequent fusion of the plasma membranes of both interacting cells.

In previous studies, MAK-1 had been shown to play potential functions during

competence and growth arrest. It is known that the mutation of *mak-1* results in a cell

fusion defective phenotype, in which cells fail to initiate interactions (Fu et al., 2011),

indicating that the cells need MAK-1 in order to start the cell fusion process. This also

suggests that MAK-1 plays a role in the activation of different factors that allow the

acquisition of the competence state by the cell.

One of these potential factors might be the Rho-GTPase RAC-1. As it has been previously

shown, mutation of *rac-1* also leads to a fusion defective phenotype and its inhibition

results in the loss of the ability of the cells to undergo cell fusion (Araujo-Palomares et al.,

2011; Lichius et al., 2014). Our experiments revealed that inhibition of MAK-1 results in

inactivation of RAC-1 and the failure of the cells to initiate interactions. We therefore

hypothesize that an active version of RAC-1 is needed to establish the cell fusion

competence in the cell and this activation is probably mediated by MAK-1. One way to

test if a mutant is blocked in the cell fusion competence state is to analyze its ability to

form CATs. For example, the mutation of *so* results in spores that are fusion defective,

but are still able to form CATs in the presence of wild-type cells, suggesting that the cells

are still fusion competent but unable to undergo the subsequent communication process

and directed growth (Fleissner et al., 2005). To analyze whether RAC-1 is also affected

in this state, a mix of Δ*rac-1* and wild-type spores should be analyzed by light microscopy

to observe whether CATs are formed. RAC-1 inhibition studies could also be performed,

by mixing of inhibited and uninhibited wild-type cells. These experiments would elucidate

whether RAC-1 is essential for cell fusion competence and would also help to solve the

implications of MAK-1 to this process.

As previously mentioned, MAK-1 also plays a role after the cells have established

physical contact, specifically in growth arrest and fusion pore formation. During the

communication phase, when MAK-2 and SO are dynamically recruited to the tips of the

interacting cells, MAK-1 remains cytoplasmic. The protein is only localized to the contact

area of the cells after establishment of cell-cell contact, once cell growth is arrested and

fusion pore is being formed (Weichert et al., 2016). Interestingly, partial inhibition of MAK-

1 which still allows fusion competence and the tropic cell-cell interaction, results in a

unique phenotype, in which the interaction partners fail to arrest growth after cell-cell

contact and twist around each other.

Together these observations suggest a role of MAK-1 in the recognition of cell-cell contact

and subsequent growth arrest. A similar failure in cell growth arrest is observed in the

Δ*erg-2* strain. The *erg-2* (*ergosterol-2*) gene encodes a sterol reductase, which mediates

the last step of the ergosterol biosynthesis in *N. crassa*. In this mutant, MAK-1 is not

recruited to the contact area and the twisting germ tubes phenotype is also observed

(Weichert et al., 2016). Interestingly, a potential connection between MAK-1 partial

inhibition and membrane sterol composition is the Rho GTPase RAC-1. The membrane

recruitment of Rho GTPases is dependent of the proper accumulation and synthesis of

sterol-rich membrane (SRM) domains which are defined by the sterol composition of the

membrane (Makushok et al., 2016). This suggest that the defects reported in the sterol-

biosynthesis mutant (Δ*erg-2*) (Weichert et al., 2016), could be caused by defects in the

composition and formation of the SRM domains, which may partially affect the proper

localization and likely activation of the Rho-GTPase RAC-1. It will also explain why partial

inhibition of MAK-1, which we can assume also cause partial inhibition of RAC-1,

produced a similar phenotype. In future studies, it will be of great interest to observe

whether the localization of activated RAC-1 (CRIB reporter) is affected in the Δ*erg-2*

strain.

**MAK-1 is essential for maintenance of the directed growth machinery while MAK-2**

**controls the direction of the growth**

As discussed above, MAK-1 possesses functions during two of the cell communication

steps: fusion competence and growth arrest after cell-cell contact. In this study we show

for the first time that, in addition, MAK-1 also plays a role during the phase of cell-cell

signaling and directed growth.

Specific inhibition of MAK-1 during this stage of the fusion process revealed that the

kinase is essential for the maintenance of an actin cluster formed by actin filaments at the

growing cell tips. This function seems to be again linked to the small Rho-type GTPase

RAC-1, whose inhibition results in a comparable disintegration of the actin aster,

suggesting that both factors function in the same pathway. Localization of activated RAC-

1 (visualized by the CRIB reporter) at the cell tip also required MAK-1 activity, suggesting

that MAK-1 functions upstream of RAC-1, meanwhile inhibition of RAC-1 does not disrupt

MAK-1 localization at the contact area of the fusing cells.

Our testing of the specific function of the Fus3p homolog MAK-2 in *N. crassa* cell fusion

revealed that the kinase is essential for the correct positioning of the actin cluster within

the cell. Consistent with our findings that MAK-1 and RAC-1 function in the same pathway

organizing the structure of this aster (see above) also the position of RAC-1 becomes

instable after MAK-2 inhibition. Future experiments should co-localize actin, the only

known formin in *N.* crassa, BNI-1, and RAC-1 after MAK-2 inhibition. If our hypothesis is

correct, all factors should co-localize in the same instable fashion after MAK-2 inhibition

(meaning they should all move around together in the inhibited cell). A major unanswered

question is how MAK-2 is positioning the actin cluster at the proper position in the cell,

thereby controlling growth directionality. We hypothesize that MAK-2 is directly involved

in the translocation of the cell-cell communication signal and is therefore marking the

position of highest signal intensity. Unfortunately, so far this signal and its cognate

receptor remain unknown. Common candidates, such as G-Protein coupled receptors or

histidine kinases could be excluded (Fleißner and Serrano, 2016). Future experiments

should therefore aim to identify the signal, the upstream activators of MAK-2, and MAK-

2 targets involved in positioning the actin cluster.

All together, the results obtained in this study support a model in which MAK-1 functions

as a key activator of actin reorganization during cell fusion. This activation occurs through

the Rho-GTPase RAC-1 which is the main organizer and also activator likely of the only

known formin in *N. crassa*, BNI-1. This actin reorganization is essential for the growth of

the fusing cells and therefore for somatic cell fusion. In contrast, MAK-2 plays the role of

positioning the actin cluster around the tip in order to respond and grow towards the

highest produced gradient of fusion signals. This way the cells are able to recognize the

presence of fusing partners in their surroundings and grow towards each other in order

to complete the fusion event.

**Material and methods**

*N. crassa strains and media*

The strains generated and used in this study are listed in the supplementary information

(Table S1). Strains were generally grown in Vogel's Minimal Medium (VMM) (Vogel,

1956) supplemented with 2% sucrose as the carbon source and with 1.5% agar for solid

media. Generally, all strains were incubated in solid VMM (supplemented with histidine

0.5 mg/ml when needed) for 3-4 days in dark at 30ºC and an additional day with natural

light at room temperature. Homokariotic strains were purified by either single spore

isolation (SSPi) or by crossing with the wild-type strain. For selection during SSPi, VMM

with hygromycin was employed. Crosses were performed on Westergaard's medium as

described earlier (Westergaard and Mitchell, 1947). The ascospores obtained from the

crosses were germinated in VMM with hygromycin and confirmed by polymerase chain

reaction (PCR).

*Plasmid construction*

The primers used for the generation of the plasmids are listed in table S2. In order to

construct the *mak-1^{E104G}* (analog-sensitive) knock-in cassette, a fragment containing a 5´

upstream region together with the *mak-1* open reading frame was PCR-amplified using

the primers 1304 and 1305 and using the previously generated *mak-1^{E104G}-gfp* plasmid

as a template (Weichert et al., 2016). By yeast recombinational cloning, the resulting

fragment was fused to the hygromycin resistance cassette (amplified with primers 82 and

83) and a 1 kb fragment homologous to the 3' downstream region of *mak-1* (generated

with primers 1306 and 1307), resulting in the plasmid 738.

For the generation of the *Ptef-1-mak-2^{Q100A}* strain, the fragments required for the

construction of the gene knock-in cassette were amplified by PCR. A 1 kb genomic region

positioned 2 kb upstream of the *mak-2* promoter (considering it to be the 1 kb region

upstream of the coding read frame) was amplified with primers 1489 and 1490. The *Ptef-*

*1-mak-2^{Q100A}-gfp* plasmid used in a previous study (Serrano et al., 2018) served as a

template for the amplification of the *Ptef-1-mak-2^{Q100A}* PCR fragment with primers 1488

and 1487. The resistance cassette used was *Hyg^{R}*, which was amplified with primers 82

and 83. Finally, the 1 kb genomic region downstream of the *mak-2* Stop-codon was

amplified with primers 1489 and 1492. The four fragments were fused by yeast

recombinational cloning generating the plasmid 846.. The plasmid 723 (*Ptef-1-dsRed-so*)

was generated by cloning of the *so* ORF amplified with primers 1419 and 1420 and ligated

into the plasmid 722 (pMF334-Tef-dsRed) with enzymes AscI/XbaI. The resulting plasmid

was transformed into N1-03 generating the strain 843 (*his-3::Ptef-1-dsRed-so*). Yeast

recombinational cloning was performed as described elsewhere (Da et al., 2000).

*Strain generation*

All strains were either generated by transformation as described in early studies (Margolin

et al., 1997) or by crossing (Westergaard and Mitchell, 1947). When indicated, strains

were purchased to the Fungal Genetics Stock Center. A helpful list of protocols and media

description is available at their website (www.fgsc.net).

*Live-cell imaging*

Sample preparation was performed as previously described (Schürg et al., 2012). For

observation of hyphal fusion in *N. crassa*, 5 µl of a $10^7$-spore suspension were inoculated

on the side of a MM agar plate and incubated for 15 hours at 30ºC. The preparation of

the sample was performed in a similar way than for the spores. The microscope used and

the settings for software analyses were described in an early study (Serrano et al., 2018).

All fluorescence images shown in this study were processed by the widely used

commercial deconvolution software Huygens (by Scientific Volume Imaging), more

specifically the Huygens Essential version. Z-stacks (usually n=10) up to 100 nm of all

fluorescence images were captured and assembled as a single tiff file by using ImageJ

(image-stacks-image to stack). The composite images were processed through the

deconvolution software with the following parameters (40-100 iterations, 12 signal/noise

ratio and 0.01% of quality change thresh).

*Quantitative analyses*

Quantification of the cell fusion and germination rate was performed as described

elsewhere (Fleissner et al., 2009; Schürg et al., 2012). For all cell fusion quantifications

at least 100 spores were quantified per sample/condition, and at least three independent

repetitions were studied.

*Chemical inhibition*

Three different chemicals were used for the corresponding experiments. Latrunculin A

(10 µM), dissolved in water, was used to test the consequences of actin-polymerization

inhibition, 1-NM-PP1 (40 µm) dissolved in DMSO was used to test the inhibition of the

analog-sensitive kinases, and NSC23766 (100 µM/25 µM), dissolved in water, was used

for specific inhibition of the Rho-GTPase RAC-1. As a control, either water or DMSO 1%

was used. For all inhibitor experiments, fresh spores were harvested and incubated for 2

hours at 30ºC in MM plates. A 1 cm$^2$ piece was cut from the agar plate and inverted onto

a coverslip. The chemical was added on the sample prior inversion onto the coverslip or

on one side of the agar block.

*Statistics applied to the data*

All quantitative data were statistically analyzed by performing a two-tailed paired t-test

using Excel. The P values were calculated for 0.001, 0.05 or 0.01, and indicated as ***,

** or * respectively. To analyze whether the variances were paired or unpaired, F-test

analyses were performed.

**Achnowledgements**

We greatly acknowledge use of materials generated by the project 'Functional analysis

of a model filamentous fungus', which was supported by a National Institutes of Health

grant (PO1 GM068087).

**Funding**

This work has been supported by funding from the Deutsche Forschungsgemeinschaft

(FL706-2) and from the European Commission (PITN-GA-2013-607963).

**References**

**Aguilar, P. S., Baylies, M. K., Fleissner, A., Helming, L., Inoue, N.,**
**Podbilewicz, B., Wang, H. and Wong, M.** (2013). Genetic basis of cell-cell fusion
mechanisms. *Trends Genet* **29**, 427-37.

**Araujo-Palomares, C. L., Richthammer, C., Seiler, S. and Castro, L.** (2011).
Functional Characterization and Cellular Dynamics of the CDC-42 - RAC - CDC-24
Module in *Neurospora crassa*. *PloS one* **6**.

**Berepiki, A., Lichius, A. and Read, N. D.** (2011). Actin Organization and
Dynamics in Filamentous Fungi. *Nature reviews. Microbiology* **9**.

**Berepiki, A., Lichius, A., Shoji, J. Y., Tilsner, J. and Read, N. D.** (2010). F-actin
dynamics in *Neurospora crassa*. *Eukaryot Cell* **9**, 547-57.

**Bishop, A. C., Ubersax, J. A., Petsch, D. T., Matheos, D. P., Gray, N. S.,**
**Blethrow, J., Shimizu, E., Tsien, J. Z., Schultz, P. G., Rose, M. D. et al.** (2000). A
Chemical Switch for Inhibitor-Sensitive Alleles of Any Protein Kinase. *Nature* **407**.

**Correia, I., Alonso-Monge, R. and Pla, J.** (2010). MAPK Cell-Cycle Regulation
in *Saccharomyces cerevisiae* and *Candida albicans*. *Future microbiology* **5**.

**Corvest, V., Bogliolo, S., Follette, P., Arkowitz, R. A. and Bassilana.** (2013).
Spatiotemporal Regulation of Rho1 and Cdc42 Activity During *Candida albicans*
Filamentous Growth. *Molecular microbiology* **89**.

**Coué, M., Brenner, S. L., Spector, I. and Korn, E.** (1987). Inhibition of Actin
Polymerization by Latrunculin A. *FEBS letters* **213**.

**Da, B., Dawson, D. and Stearns, T.** (2000). Methods In Yeast Genetics: A Cold
Spring Harbor Laboratory Course Manual.

**de Nobel, H., van Den Ende, H. and Klis, F. M.** (2000). Cell Wall Maintenance in
Fungi. *Trends in microbiology* **8**.

**Dudin, O., Bendezú, F. O., Groux, R., Laroche, T., Seitz, A. and Martin.** (2015).
A Formin-Nucleated Actin Aster Concentrates Cell Wall Hydrolases for Cell Fusion in
Fission Yeast. *The Journal of cell biology* **208**.

**Dudin, O., Merlini, L., Bendezú, F. O., Groux, R., Vincenzetti, V. and Martin.**
(2017). A Systematic Screen for Morphological Abnormalities During Fission Yeast
Sexual Reproduction Identifies a Mechanism of Actin Aster Formation for Cell Fusion.
*PLoS genetics* **13**.

**Fleissner, A., Leeder, A. C., Roca, M. G., Read, N. D. and Glass.** (2009).
Oscillatory Recruitment of Signaling Proteins to Cell Tips Promotes Coordinated Behavior
During Cell Fusion. *Proceedings of the National Academy of Sciences of the United*
*States of America* **106**.

**Fleissner, A., Sarkar, S., Jacobson, D. J., Roca, M. G., Read, N. D. and Glass.**
(2005). The So Locus Is Required for Vegetative Cell Fusion and Postfertilization Events
in *Neurospora crassa*. *Eukaryotic cell* **4**.

**Fleißner, A. and Serrano, A.** (2016). 7 The Art of Networking: Vegetative Hyphal
Fusion in Filamentous Ascomycete Fungi. In *Growth, Differentiation and Sexuality*, (ed.
J. Wendland), pp. 133-153. Cham: Springer International Publishing.

**Fu, C., Ao, J., Dettmann, A., Seiler, S. and Free, S. J.** (2014). Characterization
of the *Neurospora crassa* Cell Fusion Proteins, HAM-6, HAM-7, HAM-8, HAM-9, HAM-
10, AMPH-1 and WHI-2. *PloS one* **9**.

**Fu, C., Priyadarshini, I., Amrita, H., Julia, A., Angela, S. and Stephen, J.**
(2011). Identification and Characterization of Genes Required for Cell-to-Cell Fusion in
*Neurospora crassa*.

**Goryachev, A. B., Lichius, A., Wright, G. D. and Read.** (2012). Excitable
Behavior Can Explain the "Ping-Pong" Mode of Communication Between Cells Using the
Same Chemoattractant. *BioEssays : news and reviews in molecular, cellular and*
*developmental biology* **34**.

**Levin, D. E.** (2011). Regulation of Cell Wall Biogenesis in *Saccharomyces*
*cerevisiae*: The Cell Wall Integrity Signaling Pathway. *Genetics* **189**.

**Lichius, A., Goryachev, A. B., Fricker, M. D., Obara, B., Castro-Longoria, E.**
**and Read.** (2014). CDC-42 and RAC-1 Regulate Opposite Chemotropisms in
*Neurospora crassa*. *Journal of cell science* **127**.

**Maddi, A., Dettman, A., Fu, C., Seiler, S. and Free, S. J.** (2012). WSC-1 and
HAM-7 Are MAK-1 MAP Kinase Pathway Sensors Required for Cell Wall Integrity and
Hyphal Fusion in *Neurospora crassa*. *PloS one* **7**.

**Maerz, S., Ziv, C., Vogt, N., Helmstaedt, K., Cohen, N., Gorovits, R., Yarden,**
**O. and Seiler, S.** (2008). The Nuclear Dbf2-related Kinase COT1 and the Mitogen-
Activated Protein Kinases MAK1 and MAK2 Genetically Interact to Regulate Filamentous
Growth, Hyphal Fusion and Sexual Development in *Neurospora crassa*. *Genetics* **179**.

**Makushok, T., Alves, P., Huisman, S. M., Kijowski, A. R. and Brunner, D.**
(2016). Sterol-Rich Membrane Domains Define Fission Yeast Cell Polarity. *Cell* **165**.

**Margolin, B. S., M. Freitag and Selker., E. U.** (1997). Improved plasmids for gene
targeting at the his-3 locus of Neurospora crassa by electroporation,. *Fungal Genetics*
*Reports* **44**.

**Martin, S. G. and Arkowitz, R. A.** (2014). Cell polarization in budding and fission
yeasts. *FEMS Microbiol Rev* **38**, 228-53.

**Matheos, D., Metodiev, M., Muller, E., Stone, D. and Rose, M. D.** (2004).
Pheromone-induced Polarization Is Dependent on the Fus3p MAPK Acting Through the
Formin Bni1p. *The Journal of cell biology* **165**.

**Ninomiya, Y., Suzuki, K., Ishii, C. and Inoue.** (2004). Highly Efficient Gene
Replacements in *Neurospora* Strains Deficient for Nonhomologous End-Joining.
*Proceedings of the National Academy of Sciences of the United States of America* **101**.

**Pandey, A., Roca, M. G., Read, N. D. and Glass.** (2004). Role of a Mitogen-
Activated Protein Kinase Pathway During Conidial Germination and Hyphal Fusion in
Neurospora Crassa. *Eukaryotic cell* **3**.

**Park, G., Pan, S. and Borkovich, K. A.** (2008). Mitogen-activated Protein Kinase
Cascade Required for Regulation of Development and Secondary Metabolism in
*Neurospora crassa*. *Eukaryotic cell* **7**.

**Qi, M. and Elion, E.** (2005). Formin-induced Actin Cables Are Required for
Polarized Recruitment of the Ste5 Scaffold and High Level Activation of MAPK Fus3.
*Journal of cell science* **118**.

**Riedl, J., Crevenna, A. H., Kessenbrock, K., Yu, J. H., Neukirchen, D., Bista,**
**M., Bradke, F., Jenne, D., Holak, T. A., Werb, Z. et al.** (2008). Lifeact: A Versatile Marker
to Visualize F-actin. *Nature methods* **5**.

**Roca, M. G., Kuo, H. C., Lichius, A., Freitag, M. and Read, N.** (2010). Nuclear
Dynamics, Mitosis, and the Cytoskeleton During the Early Stages of Colony Initiation in
*Neurospora crassa*. *Eukaryotic cell* **9**.

**Schürg, T., Brandt, U., Adis, C. and Fleissner, A.** (2012). The Saccharomyces
Cerevisiae BEM1 Homologue in *Neurospora crassa* Promotes Co-Ordinated Cell
Behaviour Resulting in Cell Fusion. *Molecular microbiology* **86**.

**Serrano, A., Hammadeh, H. H., Herzog, S., Illgen, J., Schumann, M. R.,**
**Weichert, M. and Fleißner.** (2017). The Dynamics of Signal Complex Formation
Mediating Germling Fusion in *Neurospora crassa*. *Fungal genetics and biology : FG & B*
**101**.

**Serrano, A., Illgen, J., Brandt, U., Thieme, N., Letz, A., Lichius, A., Read, N. D.**
**and Fleissner, A.** (2018). Spatio-temporal MAPK dynamics mediate cell behavior
coordination during fungal somatic cell fusion. *J Cell Sci* **131**.

**Vogel, J. H.** (1956). A convenient growth medium for *Neurospora* (medium N).
*Microbial Genetics Bulletin* **13**, 42-43.

**Vogt, N. and Seiler, S.** (2008). The RHO1-specific GTPase-activating Protein
LRG1 Regulates Polar Tip Growth in Parallel to Ndr Kinase Signaling in *Neurospora*.
*Molecular biology of the cell* **19**.

**Weichert, M., Lichius, A., Priegnitz, B. E., Brandt, U., Gottschalk, J., Nawrath,**
**T., Groenhagen, U., Read, N. D., Schulz, S. and Fleissner, A.** (2016). Accumulation of
specific sterol precursors targets a MAP kinase cascade mediating cell-cell recognition
and fusion. *Proc Natl Acad Sci U S A* **113**, 11877-11882.

**Westergaard, M. and Mitchell, H. K.** (1947). *Neurospora* V. A Synthetic Medium
Favoring Sexual Reproduction. *Am. J. Bot.* **34**, 573-577.

**Yoshimi, A., Miyazawa, K. and Abe, K.** (2016). Cell Wall Structure and
Biogenesis in *Aspergillus* Species. *Bioscience, biotechnology, and biochemistry* **80**.

**Figures**

**Figure 1. Activity inhibition of MAK-1 disrupts the oscillatory localization of MAK-**
**2/SO and the formation of the fusion actin aster at the cell tips**. **(A)** Oscillatory
localization of MAK-2-GFP when DMSO (1%) is added in the strain 865 (*mak-1::mak-*
*1E104G-hph*; *his-3::Ptef-1-mak-2-gfp*). **(B)** Addition of 1-NM-PP1 (40 µM) disrupts the
oscillatory dynamic localization of MAK-2 in strain 865. **(C)** Oscillatory dynamic
localization of SO when DMSO (1%) is added in the strain 874 (*mak-1::mak-1E104G-hph*;
*his-3::Ptef-1-so-gfp*). **(D)** Addition of 1-NM-PP1 (40 µM) disrupts the oscillatory dynamic
localization of SO in strain 874. **(E)** Localization of actin (through lifeact-gfp) when DMSO
(1%) is added in the strain 869 (*mak-1::mak-1E104G-hph*; *his-3::Ptef-1-lifeact-gfp*) **(F)**
Addition of 1-NM-PP1 (40 µM) disrupts the localization of actin at the cell tips of the
interacting cells, in strain 869. All observations were made multiple times (n≥15). Scale
bars 5 µm.

**(A)**

**Figure 2. Actin focalize to the tips of the interactings cells in an oscillatory manner.**
**(A)** Oscillatory dynamic localization of actin fusion aster at the cell tips of the interacting cells in strain 754 (*his-3::Ptef-1-lifeact-gfp*). **(B)** Actin (Lifeact-GFP) co-localization with dsRed-SO during the interaction of fusing cells in the heterokaryon created with strains 754 and N1-30. Similar observation were made three times. **(C)** Actin localization in non-interacting cell vegetative growth, strain 754. Similar observations for A and C were made multiple times (n=10). Scale bars 5 µm

**Figure 3. Activity inhibition of MAK-2 disrupts the SO dynamic localization and mislocalizes actin around the cell membrane**. **(A)** Oscillatory localization of SO-GFP when DMSO (1%) is added in the strain 912 (*Pmak-2-mak-2::Ptef-1-mak-2$^{Q100A}$-hph*; *his-3::Ptef-1-so-gfp*). **(B)** Addition of 1-NM-PP1 (40 µM) disrupts the oscillatory dynamic localization of SO in strain 912. **(C)** Oscillatory localization of actin when DMSO (1%) is added in the strain 915 (*Pmak-2-mak-2::Ptef-1-mak-2$^{Q100A}$-hph*; *his-3::Ptef-1-lifeact-gfp*). **(D)** Addition of 1-NM-PP1 (40 µM) disrupts the oscillatory dynamic localization of actin in strain 915. All observations were made multiple times (n≥10). Scale bars 5 µm.

**(A)**

**(B)**

**(C)**

**(D)**

**Figure 4. Inhibition of actin assembly disrupts the MAK-2/SO dynamic localization.**
**(A)** Inhibition of actin with latrunculin A (10 μM) disrupts the localization of MAK-2 in strain
865 (*mak-1::mak-1$^{E104G}$-hph*; *his-3::Ptef-1-mak-2-gfp*) **(B)** Inhibition of actin with
latrunculin A (10 μM) disrupts the localization of SO in strain 874 (*mak-1::mak-1$^{E104G}$-hph*;
*his-3::Ptef-1-so-gfp*). **(C)** Inhibition of RAC-1 with NSC23766 (100 μM) disrupts the
localization of MAK-2 in strain 865 (*mak-1::mak-1$^{E104G}$-hph*; *his-3::Ptef-1-mak-2-gfp*)
(Next page)

**(D)** Inhibition of RAC-1 with NSC23766 (100 µM) disrupts the localization of SO in strain

874 (*mak-1::mak-1$^{E104G}$-hph*; *his-3::Ptef-1-so-gfp*). All observations were made multiple

times (n≥10). Scale bars 5 µm.

*(Figure 5 in next page)*

**Figure 5. Activity inhibition of MAK-1 results in RAC-1 inactivation while MAK-2 inhibition results in RAC-1 mislocalization. (A)** Localization of activated RAC-1/CDC-42 in both cell tips through the CRIB reporter (CRIB-GFP) during cell fusion when DMSO (1%) is added in strain 943 (*mak-1::mak-1^{E104G}-hph; Pal1-crib^{cla-4}-bar*). **(B)** Disruption in CRIB-GFP localization when 1-NM-PP1 (40 µM) is added in strain 943. **(C)** Localization of activated RAC-1/CDC-42 in both cell tips through the CRIB reporter (CRIB-GFP) during cell fusion when DMSO (1%) is added in strain 949 (*Pmak-2-mak-2::Ptef-1-mak-2^{Q100A}-hph;Pal1-crib^{cla-4}-bar*). **(D)** Mislocalization of CRIB-GFP localization when 1-NM-PP1 (40 µM) is added in strain 949. **(E)** Localization of the CRIB-GFP when 1-NM-PP1 (40 µM) is added in strain 949, followed by inhibition of RAC-1 with NSC23766 (100 µM). Comparable observations were made multiple times (n≥10). Scale bars 5 µm.

**Figure 6. Inhibition of either MAK-1 or RAC-1 does not affect actin localization in vegetative growth. (A)** Growth quantification in cells treated as control (DMSO 1%), 1-NM-PP1 40 µM (MAK-1/MAK-2 inhibition) and NSC23766 100 µM (RAC-1 inhibition). No significant differences are observed between all tests (n≥25). **(B)** Upper panel: localization of the CRIB reporter in strain 943 (*mak-1::mak-1$^{E104G}$-hph; Pal1-crib$^{cla-4}$-bar*) in cells treated with DMSO (1%) before and 30 minutes after addition. Bottom panel: localization of the CRIB reporter in strain 943 in cells treated with 1-NM-PP1 (40 µM) before and 30 minutes after addition. **(C)** Upper panel: localization of the CRIB reporter in strain 949 (*Pmak-2-mak-2::Ptef-1-mak-2$^{Q100A}$-hph;Pal1-crib$^{cla-4}$-bar*) in cells treated with DMSO (1%) before and 30 minutes after addition. Bottom panel: localization of the CRIB reporter in strain 949 in cells treated with 1-NM-PP1 (40 µM) before and 30 minutes after addition. **(D)** Upper panel: localization of actin (Lifeact-GFP) in strain 869 (*mak-1::mak-1$^{E104G}$-hph*; *his-3::Ptef-1-lifeact-gfp*) in cells treated with DMSO (1%) before and 30 minutes after addition. Bottom panel: localization of actin (Lifeact-GFP) in strain 869 in cells treated with 1-NM-PP1 (40 µM) before and 30 minutes after addition. **(E)** Upper panel: localization of actin (Lifeact-GFP) in strain 915 (*Pmak-2-mak-2::Ptef-1-mak-2$^{Q100A}$-hph*; *his-3::Ptef-1-lifeact-gfp*) in cells treated with DMSO (1%) before and 30 minutes after addition. Bottom panel: localization of actin (Lifeact-GFP) in strain 915 in cells treated with 1-NM-PP1 (40 µM) before and 30 minutes after addition. Similar observations were made multiple times (n≥5). Scale bars 5 µm.

**Figure 7. RAC-1 and MAK-1 acts within the same pathway. (A)** Quantification of the observation of twisting cell fusion growth during partial inhibition of RAC-1. Control ($H_2O$) and 25 µM (NSC23766). Significant differences were observed between the number of twisting growth observations made in the presence of the inhibitor compare to the control (n≥100) **(B)** Quantification of polarity growth defects when RAC-1 is partially inhibited during cell fusion. Control ($H_2O$) and 25 µM (NSC23766), in media with $Ca^{2+}$ or without. Significant differences were observed between the partial inhibition and the control only in the present of $Ca^{2+}$.

**Figure 8. MAK-1 is upstream of RAC-1. (A)** Localization of the CRIB-GFP during cell fusion when cells establish physical contact in strain 943 (*mak-1::mak-1$^{E104G}$-hph; Pal1-crib$^{cla-4}$-bar*) in cells treated with DMSO (1%). **(B)** Localization of the CRIB-GFP in strain 943 when MAK-1 is fully inhibited (with 1-NM-PP-1, 100 µM) in cells that just established physical contact. (*) to point the cell fusion event and signal dissapearence. **(C)** Localization of MAK-1-GFP in strain MW513 (*mak-1::hph; his-3::Pccg-1-mak-1-gfp*) with $H_2O$. **(D)** Localization of MAK-1-GFP when RAC-1 is fully inhibited with NSC23766 (100 µM). Similar observations were made multiple times (n=10) Scale bars 5 µm.