The biophysical basis underlying the maintenance of early phase long-term potentiation
Short title: AMPAR trafficking during early phase LTP

Moritz F. P. Becker *,[1] and Christian Tetzlaff[1,2]

[1]III. Institute of Physics – Biophysics, Georg-August University, Göttingen, Germany
[2]Bernstein Center for Computational Neuroscience, Göttingen, Germany

*Correspondence to:
Moritz F. P. Becker
moritz.becker@phys.uni-goettingen.de

Conflict of Interest Statement: The authors declare no competing financial interests.

Acknowledgments: We would like to thank Silvio Rizzoli for various fruitful discussions and helpful comments. The research was funded by the German Research Foundation (CRC1286, project C1 and project #419866478).

Abstract

The maintenance of synaptic changes resulting from long-term potentiation (LTP) is essential for brain function such as memory and learning. Different LTP phases have been associated with diverse molecular processes and pathways, and the molecular underpinnings of LTP on the short, as well as long time scales, are well established. However, the principles on the intermediate time scale of 1-6 hours that mediate the early phase of LTP (E-LTP) remain elusive. We hypothesize that the interplay between specific features of postsynaptic receptor trafficking is responsible for sustaining synaptic changes during this LTP phase. We test this hypothesis by formalizing a biophysical model that integrates several experimentally-motivated mechanisms. The model captures a wide range of experimental findings and predicts that synaptic changes are preserved for hours when the receptor dynamics are shaped by the interplay of structural changes of the spine in conjunction with increased trafficking from recycling endosomes and the cooperative binding of receptors. Furthermore, our model provides several predictions to verify our findings experimentally.

Author summary

The cognitive ability of learning is associated with plasticity-induced changes in synaptic transmission efficacy mediated by AMPA receptors. Synaptic changes depend on a multitude of molecular and physiological mechanisms, building complex interaction networks. By formalizing and employing a biophysical model of AMPAR trafficking, we unravel and evaluate the interplay between key mechanisms such as receptor binding, exocytosis, morphological changes, and cooperative receptor binding. Our findings indicate that cooperative receptor binding in conjunction with morphological changes of the spine and increased trafficking from recycling endosomes leads to the maintenance of synaptic changes on behaviorally relevant time spans. Characterizing the principles underlying synaptic changes will provide insight into the role of synaptic dynamics in neurodegenerative diseases.

 # Introduction

Fig 1. Temporal course of E-LTP and related processes. **(A)** E-LTP decays after 1-6 hours (squares; data taken from [1]) , while L-LTP remains stable for hours to days ([2]; not depicted here). When exocytosis is blocked at the postsynaptic spine, E-LTP decays after 10-20 min (triangles; data taken from [3]). **(B)** The exocytosis event rate of GluR1 at spines is increased for a few minutes after LTP-induction (data taken from [4]). **(C)** Functional LTP is accompanied by structural changes of the spine, which also depends on exocytosis (data taken from [5, 6]). **(D)** The activity of CaMKII is also increased for a few minutes after LTP-induction (data taken from [7]). The figures have been redrawn from [1] Fig 3C, [3] Fig 2C, [4] Fig 4A, [5] Fig 2C, [6] Fig 4C, [7] Fig 2B.

Synaptic plasticity regulates synaptic transmission strength and is associated with cognitive processes such as learning and memory. For excitatory synapses, the transmission strength is mainly determined by the number of α-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptors (AMPARs) accumulating at the postsynaptic density (PSD) [8, 9, 10, 11]. Long-term potentiation of synaptic strength (LTP) can be divided into different phases such as early-phase LTP (E-LTP), which decays by 1-6 hrs after induction, or late-phase LTP (L-LTP), which lasts for several hours or even days and depends on the synthesis of new proteins [12, 2, 13]. While reports have identified various

molecular processes and pathways underlying the different phases of LTP, especially for L-LTP [14, 2, 15], the biological and biophysical mechanisms maintaining E-LTP remain unclear. To address this issue, here we focus on protein synthesis independent E-LTP and investigate the influence of different experimentally motivated candidate mechanisms on AMPAR dynamics at the PSD [1].

AMPARs are continuously inserted and removed from the spine membrane by exocytosis and endocytosis [16, 4, 17, 18]. Experimental studies indicate that E-LTP decays after several minutes if the exocytosis of additional AMPARs is blocked ([3, 17, 19, 20]; Fig 1A, triangles). Although these results suggest that exocytosis could be the main mechanism underlying the maintenance of E-LTP, the rate of exocytosis events at dendritic spines is only increased for a few minutes after LTP-induction and exocytosis of receptors is slow compared to diffusional exchange ([4, 21, 16]; Fig 1B).

AMPARs are regularly exchanged between the postsynaptic and the dendritic membrane via lateral diffusion but can become bound and thereby functional at the PSD [22, 23, 24, 18]. Several studies indicate that the phosphorylation and interaction of PSD scaffolding proteins such as PSD95 and transmembrane AMPAR regulatory proteins (TARPs) such as Stargazin play a key role in binding AMPARs at the synapse [10, 25, 26]. However, the activity of kinase CaMKII which targets these PSD proteins is only upregulated for a few minutes after LTP-induction ([7]; Fig 1C). While the phosphorylated state of the TARPS probably outlasts CaMKII activity, the fast decay of LTP in the case of blocked exocytosis indicates that AMPAR binding rates are not increased for much longer [3, 7]. Also, concentrations of PSD proteins only increase in the presence of protein synthesis and persistent spine enlargement, indicating that the number of possible binding sites stays approximately constant during E-LTP [27, 28]. Taken together, these results demonstrate that the principles underlying the maintenance of E-LTP remain unknown as neither the dynamics of the exocytosis rate nor the CaMKII activity seems to match the slow decay of E-LTP (Fig 1A, squares).

However, additional biological and biophysical mechanisms influence the AMPAR dynamics at the dendritic spine. The induction of LTP for example also yields structural modifications of the spine (sLTP), which is expressed by an increase in spine volume

and the translocation of AMPAR-containing recycling endosomes (REs) into the spine [29, 30, 31, 32, 5, 33, 34, 6, 27] (Fig 1C). Both could influence AMPAR trafficking: RE translocation by increasing receptor recycling to the surface via AMPAR exocytosis [29, 30, 4] and spine growth by regulating diffusion between dendrite and spine [35, 36, 37]. Exocytosis from REs not only delivers AMPARs to the spine membrane but also stabilizes the spine after LTP induction and has been suggested to be involved in the early maintenance of LTP [30, 29, 31, 6, 38]. Importantly, changes in the spine volume and RE allocation last much longer than changes in the rate of exocytosis events at the spine [27, 29, 4, 21].

Furthermore, AMPARs are not homogeneously distributed in the PSD but cluster on PSD nanodomains formed by PSD95 and other scaffolding proteins [39, 40]. The emergence of such clusters can be related to receptor-receptor and receptor-scaffolding protein interactions, which in turn influence the binding of AMPARs [41, 42, 43, 44, 45]. The relationship between structural plasticity, AMPAR clustering, and the induction and maintenance of E-LTP remains unexplained.

In this study, based on well-established theoretical studies [46, 45] we derive a mathematical model of AMPAR trafficking in a dendritic spine that integrates diverse experimental data [4, 47, 5, 6, 29, 48, 49, 50, 51, 24, 7]. We use this model to systematically investigate the impact of the above-mentioned biological mechanisms on the maintenance of E-LTP by comparing the model with the experimental data (summary of experimental findings shown in Fig 1, data taken from [1, 7, 4, 3]).

We find that morphological changes in conjunction with increased RE trafficking and cooperative binding can complement each other and together provide a possible explanation for the decay time of E-LTP as well as the dependence of LTP on exocytosis. This combined model provides a new hypothesis about the principles underlying the maintenance of E-LTP and leads to specific, new experimental predictions that could verify our findings.

Results

Basic model of AMPAR-trafficking

The basic model integrates the core processes of AMPAR trafficking in the dendritic spine: lateral diffusion to and from the dendritic membrane, exo- and endocytosis within the spine, and binding to and disassociation from scaffolding proteins in the PSD (Fig 2A). We formalize the coaction of these processes by considering the number of AMPARs that freely move on the membrane of the spine denoted by U and of AMPARs that are bound in the PSD labeled by B. The surface area of the spine is denoted by A_{spine}. Please note that we do not explicitly model the number of AMPARs in the intracellular space of the spine or on the dendritic membrane. As the rate of exocytosis is slow [16, 4] and diffusion of surface receptors is fast [23] we assume that the dendritic and intracellular pools are rapidly replenished. On the synaptic membrane, U is increased by exocytosis at a rate $k_{exo} S_{exo}$ where S_{exo} denotes the size of an exocytic event and k_{exo} the rate at which exocytosis events occur. U is further increased by lateral diffusion from the dendritic to the spine membrane at a rate k_{in}, and by the release of AMPARs from the PSD described by $k_{BU} B$. U is decreased by endocytosis at a rate $k_{endo} \frac{U}{A_{spine}}$ and lateral diffusion to the dendrite at a rate $k_{out} \frac{U}{A_{spine}}$, as well as by AMPARs binding to the PSD scaffold described by $k_{UB}(P-B) \frac{U}{A_{spine}}$, where P denotes the number of slots formed by scaffolding proteins. Thus, the change in the number of unbound AMPARs can be formalized by

$$\frac{dU}{dt} = k_{exo} S_{exo} + k_{in} + k_{BU} B - (k_{endo} + k_{out} + k_{UB}(P - B)) \frac{U}{A_{spine}}. \tag{1}$$

We do not consider direct endo- and exocytosis of bound AMPARs such that the number is only regulated by the interchange with freely moving AMPARs.

$$\frac{dB}{dt} = k_{UB}(P - B) \frac{U}{A_{spine}} - k_{BU}B. \tag{2}$$

Equations (1) and (2) describe the AMPAR-dynamics on the spine's membrane under basal condition. We derived the corresponding set of parameter values of an average spine by estimating the values from several experimental studies that are mainly conducted in

hippocampal cultures and slices (see Table 1) and are described in more detail in the methods section.

Table 1: Parameters of AMPAR trafficking.

Parameter	Meaning	Value	Experiments
k_{exo}^0	Rate of exocytosis events at the spine under basal condition	$0.0018 \ s^{-}1$	[4]
k_{endo}	Rate of receptor endocytosis in the spine	$0.0021 \ \mu m^2/s$	This study, equation (4)
k_{out}	Rate at which AMPARs exit the spine via lateral diffusion	$0.018 \ \mu m^2/s$	[51, 22], this study
k_{in}	Rate at which AMPARs enter the spine via lateral diffusion	$0.2 \ \#/s$	[35, 50, 49, 4]
k_{BU}	Unbinding rate of receptors from PSD slots	$0.1 \ s^{-1}$	[24]
k_{UB}^0	Binding rate of AMPARs at the PSD	$0.0036 \ \mu m^2/(\#s)$	This study, equation (5)
k_{UB}^0 (Coop.)	Binding rate of AMPARs at the PSD in the cooperative binding model	$0.0005 \ \mu m^2/(\#s)$	This study, equation, 24
U^*, B^*	Steady state number of mobile and bound receptors at the average spine	$10, 20 \ \#$	[48, 52]
P	Number of slots at the PSD	$70 \ \#$	This study
V_{spine}	Spine volume	$0.08 \ \mu m^3$	[5, 37]
S_{exo}	Size of an AMPAR exocytosis event	$13 \ \#$	[47]

Exocytosis event and binding rate increase alone cannot account for the maintenance of E-LTP

To examine whether a brief increase in exocytosis event and binding rate can account for the maintenance of E-LTP for several hours as well as the fast decay in the absence of exocytosis, we first study a basic model of AMPAR trafficking.

We model the LTP-induced variation of the exocytosis event and binding rate (Fig 1B and D). Parameter values under basal conditions, i.e. before LTP induction, will be denoted by a superscript index 0 (k_{exo}^0, k_{UB}^0). The time-dependence of k_{exo} and k_{UB} is described by an exponential rise (τ_{exo}^1, τ_{UB}^1) and decay (τ_{exo}^2, τ_{UB}^2) with maximum amplitudes of A_{exo} and A_{UB} (see Table 2 and Equations (6)-(8)). Please note that the

Fig 2. Different model versions and resulting AMPAR dynamics compared to experimental data.

Fig 2. **(A)** *Basic model*: AMPA receptors are trafficked across the spine via several pathways. During LTP-induction, exocytosis (k_{exo}) and binding rate (k_{UB}) are briefly increased. The LTP-induced dynamics of the basic model (blue line) is compared to experimental data of E-LTP (black squares; data taken from [1]) and also to LTP with the blockage of exocytosis (orange line compared to black triangles; data taken from [3]). **(B)** *sLTP model*: Increase in spine volume and RE trafficking in addition to k_{exo} and k_{UB} influence the LTP-induced AMPAR dynamics. Note that B only slowly decays back to the baseline if exocytosis occurs but rapidly decays otherwise. **(C)** *Cooperative model*: Cooperative receptor binding slows down the decay of LTP induced increase in bound AMPARs and increases the sensitivity to exocytosis. **(D)** *sLTP+Cooperative Model*: sLTP and cooperative receptor binding in concert with increased exocytosis and binding rate sustain E-LTP . **(A-D)** LTP-induction is triggered at time point $0s$.

153 extensions of the basic model to also include structural plasticity or cooperative receptor

154 binding are described in the corresponding sections below.

Table 2: Parameters of LTP-induced changes in exocytosis event and binding rate.

Parameter	Meaning	Value	Experiments
A_{UB}	Factor of LTP-induced increase in binding rate in the basic model	30	this study
A_{UB} (*Coop.*)	Factor by which the binding rate increases upon LTP-induction	5	this study
τ_{UB}^1	Time constant of binding rate increase during LTP induction	5 s	[7, 3]
τ_{UB}^2	Decay time constant of binding rate after LTP induction	60 s	[7, 3]
A_{exo}	Factor by which the exocytosis event rate increases upon LTP-induction	5	[4, 17]
τ_{exo}^1	Time constant of exocytosis event rate increase during LTP induction	25 s	[4]
τ_{exo}^2	Decay time constant of exocytosis event rate after LTP induction	60 s	[4]

155 Under basal conditions, the different processes of AMPAR-dynamics described before

156 balance each other such that the number of mobile and bound AMPARs in the PSD and

157 on the synaptic membrane reach an equilibrium state of constant temporal averages with

158 28.5% of slots being occupied (parameters have been chosen explicitly to result in a low

159 occupancy of the PSD; see Methods).

The induction of LTP and the related increase in receptor exocytosis and binding lead to a fast increase in the number of mobile and bound AMPARs, which results in a stronger synaptic transmission strength, assuming that AMPAR EPSCs are proportional to the number of bound AMPARs at the PSD (Fig 2A, blue line). After a few minutes, the exocytosis event and binding rate return to basal levels. Given these dynamics, the increased level of bound AMPARs in the PSD declines such that it reaches the pre-LTP level after about $10 - 15$ minutes. The time course of bound AMPARs is similar in the case of blocked exocytosis (Fig 2A, orange line). This result indicates that some core mechanisms are missing, which maintain bound AMPARs in the PSD for a longer duration.

Structural plasticity is not sufficient to explain the maintenance of E-LTP

To account for the effects of sLTP we first model the change in spine volume V_{spine} (Equations (11) and (12)). Second, we assume that the magnitude of exocytosis events S_{exo} increases proportionally with the spine volume due to the translocation of AMPAR-containing REs (Fig 3C, Equation (13)) (for details see Materials and Methods).

All parameters of sLTP are based on previous studies ([27, 5, 37, 6]; see Materials and Methods). Similar to the basic model, the induction of LTP leads to a strong and fast increase in the number of bound AMPARs; however, the level of bound AMPARs falls significantly slower with sLTP than in the basic model (Fig 2B). This slower decay can be attributed to the increase in mobile receptor concentration that is mediated by the change in S_{exo} (Figs 3C and D) and disappears if exocytosis is blocked (Fig 2B, orange line). Given normal conditions, the time span until the AMPAR level falls back to the basal level is similar to experiments ($\approx 100min$). However, between 10 and 100 minutes the overall level of bound AMPARs is lower than in experimental data. Thus, the influence of the sLTP on the AMPAR-dynamics is not sufficient to explain the maintenance of E-LTP.

Fig 3. Basic principles and results of cooperative receptor binding and sLTP.
(A) Sketch of the cooperative model mechanisms: Slots of the PSD are spatially ordered into a grid with each slot having a binding $\hat{k}_{UB}^{coop}$ and unbinding $\hat{k}_{BU}^{coop}$ rate dependent on the number of nearest neighbours, indicated by different colors (white: occupied slot). **(B)** The mean binding $\langle\hat{k}_{UB}^{coop}\rangle$ and unbinding rate $\langle\hat{k}_{BU}^{coop}\rangle$ (Equations (17), (18)) vary with the grid size (slots P) and the total number of bound receptors. The simulation results are fitted by Equations (20) and (19) (dashed lines). The dynamics of the resulting mean-field model match well with the stochastic simulations (**A**, inset). **(C)** The enlargement of the spine during sLTP is accompanied by an increase in the size of exocytosis events S_{exo} (Equations (13) and (12)). **(D)** The concentration of mobile receptors U/A_{spine} increases after LTP-induction but is only sustained if accompanied by sLTP.

Cooperative receptor binding is not sufficient to explain the maintenance of E-LTP

Our model of cooperative AMPAR-dynamics, based on [45], enables us to investigate the influence of cooperative binding effects on the maintenance of E-LTP. In this model, the probability of receptors to bind to or unbind from a slot depends on the number of near-

est neighbours on a grid that represents the PSD (Fig 3A). We developed a mean-field approximation of this stochastic grid model (Equations (19), (20)), which is integrated into the basic model (Equations (1), (2)). In the mean-field model, cooperativity results in effective binding and unbinding rates k_{UB}^{coop}, k_{BU}^{coop} that depend on the number of already bound receptors B as well as the size of a PSD domain P (Fig 3B). Like before, LTP-related changes of the exocytosis event and binding rate are modeled by temporal variations of k_{exo} and k_{UB}. Similar to the basic model, LTP-induction yields a fast increase in the number of bound AMPARs followed by a fast decay (Fig 2C, blue line). Compared to the basic model, the decay is slower such that the level of bound AMPARs reaches the basal condition after $\approx 30 - 35$ minutes; however, it is significantly faster than in experiments. The blockage of exocytosis, after LTP-induction, reduces the level of bound AMPARs to $\approx 80\%$ (orange line), which is different from the basic or sLTP model. This effect of reduced transmission strength is also found in experiments (Fig 2C, triangles). Thus, the cooperative effect between receptors is not sufficient to explain the maintenance of E-LTP.

Exocytosis is the limiting factor in E-LTP maintenance

Although based on experimental data, the parameters considered before could embrace an imprecision in their exact values as exocytosis might be underestimated in experiments [53] and the amplitude and decay of the binding rate during LTP is uncertain. We, therefore, utilize a random sampling method (see Materials and Methods) to find the parameter set that results in the best match of the AMPAR-dynamics of the model with the experimentally measured temporal course of transmission strength (Fig 1A). For this, we randomly drew 2000 different parameter values for the trafficking rates $k_{BU}, k_{UB}^0, k_{exo}^0, k_{endo}$ and the LTP-induced temporal development of exocytosis (τ_{exo}^2, A_{exo}) and receptor binding rates (τ_{UB}^2, A_{UB}) from a reasonable regime (see Table 3). We simulated the resulting AMPAR-dynamics for the basic model and its extensions by sLTP or cooperative binding discussed before. To find the best match, for each trial we use a distance measure (Equation (9)) between the temporal development of the number of bound AMPARs in the model and the experimentally measured development of the EPSP with and without blockage of

Fig 4. Variation of parameter values yields parameter sets for all three model versions such that E-LTP is maintained for several hours. (A) The experimentally measured temporal course of E-LTP with and without the blockage of exocytosis is matched by all three model versions (basic, sLTP, cooperative) with corresponding parameter sets. Please note that lines overlie each other. **(B,C)** For the parameter sets used in (A), the LTP-induced increase of the binding rate of the models matches experimental data of CaMKII (B), whereas the exocytosis event rate is too high for the basic and sLTP model and decays too slow for all three model versions (C). **(D)** The concentration of mobile receptors U increases to $\approx 300\%$ of the baseline level for the basic and the sLTP model but only to $\approx 130\%$ for cooperative model. (A-D) Shown are the upper 0.5% that match best with the experimental data in (A). For the matching we used the cost function described by equation (9). Shades indicate standard deviation across parameter sets.

₂₂₀ exocytosis (see Materials and Methods).

₂₂₁   Indeed, for all three model versions, several parameter sets yield dynamics that are

₂₂₂ in agreement with experimental findings with and without the blockage of exocytosis

₂₂₃ (Fig 4A, colored lines; the 0.5% best matching parameter sets are shown). Furthermore,

₂₂₄ the temporal development of the binding rate is consistent with the CaMKII activity

₂₂₅ during LTP-induction (Fig 4B). However, for the basic model and the sLTP-model, the

₂₂₆ exocytosis event rate becomes much too high compared to experiments and the decrease

₂₂₇ becomes too slow (Fig 4C). Thereby, mobile receptor surface concentration U/A_{spine} in-

₂₂₈ creases by about 200%, which is far beyond experimental observations ($\approx 20\%$ for GluR1,

Table 3: Random parameter sampling.

Parameter	Meaning	Range
k_{exo}^0	Rate of exocytosis events at the spine under basal conditions.	$0...0.01\ s^{-1}$
τ_{exo}^2	Decay time constant of exocytosis event rate after LTP induction.	$0...67\ min.$
A_{exo}	Factor by which the exocytosis event rate increases upon LTP-induction.	$0...20$
k_{BU}	Unbinding rate of AMPARs from slots at the PSD.	$0...1\ s^{-1}$
τ_{UB}^2	Decay time constant of binding rate after LTP induction.	$0...8\ min.$
A_{UB}	Factor by which the binding rate increases upon LTP-induction.	$0...50$
$A_{UB}(Coop.)$	Factor by which the binding rate increases upon LTP-induction in the cooperative binding model.	$0...20$

[27]). For the model of AMPAR-dynamics with cooperative binding, the exocytosis event rate stays in a reasonable regime but also decays too slowly (lime green). However, the latter result indicates that the increase in exocytosis due to sLTP might be sufficient to maintain E-LTP in the cooperative binding model. This is also apparent from the increase in the mobile receptor concentration necessary to mediate E-LTP in the cooperative model (Fig 4D), which is close to the increase during sLTP ($\approx 25\%$, Fig 3D). We repeated this procedure for different values of the diffusion parameter k_{in} and k_{out} and the steady state U^*, B^*, yielding to similar results (S1 Fig).

sLTP and cooperative binding complement each other to maintain synaptic potentiation

We have shown that neither the basic AMPAR dynamics nor its extension by sLTP or cooperative binding seems to explain the maintenance of E-LTP. Next, we test whether a combination of cooperative binding and sLTP complement each other to mediate E-LTP as observed experimentally. All model parameters and the temporal developments of exocytosis event rate, binding rate, and spine volume are the same as before. In this combined model with experimentally derived parameters, the number of bound AMPARs increases rapidly after LTP induction (Fig 2D). If exocytosis is blocked, this increase is followed by a rapid decay below the baseline level (orange line). In presence of exocytosis,

Fig 5. The decay of E-LTP is determined by the size of spine and PSD. (A) In the cooperative model, the number of bound AMPARs in the fixed point B^* strongly depends on the amount of exocytosis from REs (S_{exo}) compared to the basic model. Thereby, small changes in the amount of exocytosed AMPARs ΔS_{exo} are sufficient for LTP expression (ΔB). **(B)** The decay time and relative amplitude of LTP change with the size of the PSD P. Whereas bound AMPARs B show fast dynamics for small PSDs (dark blue line), B only slowly decays back to the steady state baseline for large PSDs (light blue line). **(C)** E-LTP for different spine sizes where the PSD area scales linear with the spine surface area ($P \propto A_{spine}$) but the amount of baseline AMPAR exocytosis is kept constant ($S_{exo}^0 = 13$). Top: Although large spines lack pronounces sLTP, the slower dynamics due to the larger PSD result in E-LTP with a time scale similar to that of smaller spines. Bottom: The change in AMPAR abundance at the PSD 1 and 20 min. after LTP induction. **(D)** Same as in **(E)** but S_{exo}^0 depends on the initial spine volume V_{spine}^0 according to equation (13). Top: E-LTP decays rapidly for large and very small spines. Bottom: Same as in **(E)**.

₂₄₇ however, the combined dynamics of sLTP and cooperative binding sustain the level of

₂₄₈ bound AMPARs at a level that matches experimental data (blue line). But why are both

₂₄₉ mechanisms required to maintain E-LTP?

₂₅₀   To provide an answer, we assume that the trafficking dynamics of AMPARs are much

₂₅₁ faster than the changes triggered by LTP. This allows us to consider the fixed point

₂₅₂ of bound AMPARs B^* as a function of the spine volume V_{spine} and respectively the

size of exocytosis events S_{exo} (Fig 5A). This function illustrates the different influences cooperative binding, morphological changes, and RE trafficking have on the number of bound AMPARs:

- *Cooperative binding* influences the shape of this function. If the AMPAR dynamics are determined by cooperative binding (Fig 5A, blue line), the function becomes steep and sigmoidal-shaped. This is significantly different compared to the function of the basic model, which has a shallow slope (Fig 5A, black line).

- *Morphological changes* influence the argument of the function. A variation of the spine volume and the exocytosis event size shifts the system to a new state with a different number of bound AMPARs.

Thus, AMPAR-trafficking dynamics only extended by sLTP (black line) would imply that an increase by ΔS_{exo} results in a small change of the number of bound AMPARs. By contrast, if we additionally consider cooperative binding between AMPARs (blue line), a shift in V_{spine} and therefore in S_{exo} results in a significant increase of the number of bound receptors (here, ΔB).

Decay of E-LTP depends on PSD and spine size

For the results discussed before, we considered an average spine with a spine volume of $V_{spine}^0 = 0.08 \mu m^3$ before LTP-induction and a constant PSD size with 70 binding sites. Next, we evaluate to what degree E-LTP maintenance depends on the initial volume V_{spine}^0 with the corresponding number of slots P. We assume that the number of slots P changes linearly with the initial surface area of the spine A_{spine}^0:

$$P = P^{basal} \cdot \frac{A_{spine}^0}{A_{spine}^{basal}}, \tag{3}$$

where $P^{basal} = 70$ and $A_{spine}^{basal} = 0.898 \mu m^2$ such that the parameter values used before still hold (Table 1). The surface area is calculated from the volume assuming spherical spines. In addition, experimental studies have shown that sLTP mainly occurs for small to intermediate spines but not for large spines with volumes above about $0.3 \mu m^3$ [5, 34]

(Fig 7H). We integrated this dependence of ΔV_{spine} on V_{spine}^0 into our model using data

from several studies [5, 34, 6] (Equation (11)).

First, we assume that the size of exocytosis events under basal conditions does not

depend on the initial spine volume but is constant ($S_{exo}^0 = 13$). Surprisingly, in this case,

E-LTP is sustained even for large spines where sLTP is not as pronounced (Fig 5D). Simi-

larly, we find that the system takes longer to return to equilibrium after LTP-induction as

we increase the PSD size P while keeping V_{spine} constant and independent of P (Fig 5B).

This effect is also present in the non-cooperative model but much less pronounced (not

shown).

If the initial size of exocytosis events S_{exo}^0 scales with the initial spine volume (Equation

(13)), we find that E-LTP is sustained for intermediate initial spine sizes $V_{spine}^0 \approx 0.1 \mu m^3$

(Fig 5C). For very small ($V_{spine}^0 < 0.03 \mu m^3$) and large spines ($V_{spine}^0 > 0.35 \mu m^3$), however,

E-LTP decays after about 25 minutes, matching experimental insights indicating that

large spines do not experience long-lasting LTP [5, 34]. We argue that if exocytosis

increases with the spine size, the PSD will saturate due to the larger influx of receptors.

Thereby, receptors that are added to the spine during LTP induction only have a low

chance of getting bound such that B quickly returns to the baseline in the absence of

sLTP.

Model predictions

This study yields some experimentally verifiable predictions. Our main prediction is the

requirement of the interplay between cooperative binding of receptors and morphological

spine changes accompanied by an increase in AMPAR trafficking to the spine surface

from REs during E-LTP (Fig 2D). By contrast, each mechanism alone does not lead to

the required accumulation of bound receptors for a time span of several hours (Figs 2A-

C). We predict that due to cooperative effects, increased exocytic trafficking of AMPARs

is already sufficient for synaptic potentiation without the need for additional factors that

stabilize AMPARs at the PSD (Fig 5A). Some effects of cooperative receptor binding

should be readily visible in experiments.

1. Cooperativity results in a sigmoidal dependence between PSD occupancy and the

Fig 6. Receptor clustering influences the statistical properties and mobility of receptors at the PSD. (A) Characteristic for cooperative processes the steady state value of B shows a sigmoidal dependence on the concentration of mobile receptors U/A_{spine}. The mean-field model (dotted lines) matches the stochastic model (solid lines). Bars indicate standard deviations. **(B)** In the case of cooperative receptor binding, the CV steeply changes with the size and occupancy of the PSD (colored lines) in contrast to the case in which binding rates are constant (black dotted line). **(C)** The average dwell time $\langle \tau_{dwell}^{PSD} \rangle$ of receptors is not uniformly distributed across the PSD nanodomain. Despite the small size of the nanodomain, as the concentration of AMPARs increases, the receptor dwell times can range from seconds up to tens of minutes at the cluster core (left to right).

Fig 6. **(D)** The different dynamics of receptors in the cooperative model are visible in FRAP simulations. Top, left and right: As the PSD size P increases, the recovery of receptors is slowed down. The model matches experimental data of GluR1 recovery in hippocampal cultures [49]. Bottom, left: As the concentration of mobile receptors U/A_{spine} increases, recovery is slowed down considerably in the cooperative model. At a concentration $U/A_{spine} = 33.4\,\#/\mu m^2$ the model results match well with experimental data from rat visual cortex [54]. This effect is absent in the non-cooperative model (Bottom, right). Shown are the mean values over 100 trials of the stochastic simulation. The standard error of the mean is in the order of the line thickness.

concentration of mobile receptors in the spine membrane (Figs 6A and 4A).

2. Whether receptor binding is cooperative could be inferred from the statistics of re-
ceptor fluctuations. In the stochastic clustering model (see methods), the coefficient
of variation (CV) of the number of bound receptors B lies within the range of that
of EPSC peak amplitudes (Fig 6B, [55]: $CV \approx 20\% - 70\%$ with an average of
$\approx 39\%$). However, if binding is cooperative, the CV depends on the PSD size and
steeply on the occupancy (Fig 6B, colored lines), whereas in the non-cooperative
case the CV follows that of a binomially distributed random variable and is inde-
pendent of the PSD size (black, dotted line). Therefore, steep changes in the CV
upon changes in the occupancy of the PSD or changes in the concentration of mobile
receptors respectively could indicate cooperativity. It is however difficult to relate
fluctuations in e.g. the EPSC amplitude to specific processes such as presynaptic
neurotransmitter release or receptor properties.

3. Receptors are not uniformly distributed on nanodomains formed by PSD proteins
but cluster. This effect has been found in experiments [39].

4. Rigid receptor cluster formation due too cooperative binding results in a non-
uniform distribution of receptor dwell times τ_{dwell}^{PSD} within PSD nanodomains (Fig 6C),
where the average dwell time increases with the PSD occupancy ($k_{BU}^{coop} \sim 1/B$; Fig
3B, bottom). Indeed, the time that receptors spend within nanodomains ranges from
seconds to a few minutes [39]. Besides, fluorescence recovery after photo-bleaching
(FRAP) studies found pools of highly immobile receptors at synapses, which might

account for receptors at large cluster cores [49] while receptors within a small cluster or at the border of clusters may account for the very transient immobilization found in single receptor tracking studies [24].

5. Model simulations of FRAP show that, in the cooperative model, the AMPAR recovery is shaped considerably by the concentration of mobile receptors U/A_{spine} compared to the non-cooperative model (Fig 6D, bottom). As before, we assumed a linear relationship between spine area A_{spine} and PSD size P (Equation (3)). At low receptor concentrations $U/A_{spine} = 11.2\,\#/\mu m^2$, AMPAR recovery and its dependence on the spine size and PSD size P are almost indistinguishable between the two models (Fig 6D, top). Fast recovery found in hippocampal cultures [49] is matched at PSD size $P = 81$ and mobile receptor concentration $U/A_{spine} = 11.2\,\#/\mu m^2$. Slow recovery found in rat visual cortex [54] is matched by the cooperative model at PSD size $P = 144$ and a concentration of $U/A_{spine} = 33.4\,\#/\mu m^2$ (Fig 6D, bottom left), which is consistent with larger synaptic weights in the visual cortex [56] . However, these results should be interpreted with care when comparing to experimental data as similar results could be achieved by changes in the diffusion coefficient or the receptor unbinding rate. Still, this shows that cooperativity affects recovery and therefore could be inferred from FRAP experiments.

6. We assumed ongoing, rapid exchange of receptors observed experimentally [24]. If receptors are not stabilized otherwise, this implies the need for a sustained pool of extrasynaptic receptors that constantly replenishes receptors unbinding from the PSD scaffold. We, therefore, predict that cutting off the PSD from the mobile pool would result in a rapid decay of excitatory synaptic transmission.

7. We find that there exist three alternative explanations for the dependence of E-LTP on exocytosis: E-LTP is maintained by 1) stabilization of receptors due to an exocytic factor [53], 2) an exocytic factor that raises the probability of receptor binding or 3) a rapid increase in the number of slots after LTP induction [57, 58]. Most apparent, neither the generation of additional binding sites nor the exocytosis of factors that stabilize AMPARs require an increase in the concentration of the

spine's mobile receptor pool to maintain E-LTP, whereas in our model the mobile pool concentration needs to grow by $\approx 25\%$ (Fig 3D). We predict that this increase could be mediated by a sLTP dependent change in receptor exocytosis at the spine (Fig 3C,D).

Discussion

The question of how E-LTP is maintained has not been answered in previous studies. Also, the role of receptor exocytosis is still ambiguous. In this study, we focused on two major experimental observations: First, E-LTP can be long-lasting with decay times on the order of hours [12, 59]. Second, there exists a strong dependency between LTP and exocytosis, although the exocytosis of AMPARs is slow compared to the rapid exchange of receptors via lateral diffusion [3, 19, 60, 16, 4]. Our model indicates that the experimentally observed increase in the rate of AMPAR exocytosis events and their binding to the PSD scaffold are not sufficient to maintain E-LTP (Figs 2, 4, and S1 Fig). Furthermore, our results indicate that sLTP accompanied by increased AMPAR trafficking from REs during E-LTP [38, 29, 61] and cooperative receptor binding [45, 11] individually also can not account for the maintenance of E-LTP. However, we demonstrated and predict that cooperative receptor binding complemented by increased trafficking from REs is sufficient to sustain E-LTP. Thus, our study provides several insights about the functional roles of diverse mechanisms during LTP and yields predictions highlighted above.

Bound AMPARs as the determinant factor of synaptic strength We assumed that the EPSP amplitude measured in LTP experiments is determined by the number of bound AMPA receptors. This assumption is supported by several experimental studies demonstrating that LTP is mostly expressed at the postsynaptic site and that excitatory synaptic transmission depends on the number of AMPARs [62, 63, 64, 65, 3]. However, the relationship between receptor number and EPSP amplitude may be nonlinear. For example, the alignment of receptors with the presynaptic release site is an important factor that influences the postsynaptic response [66, 39, 67]. However, whether or to what degree enhanced alignment might occur during E-LTP is unknown. Other important factors that

shapes the postsynaptic response are the presynaptic release probability and the amount of glutamate that is released. Depending on the experimental setup and the conditions of LTP induction, long-lasting presynaptic changes sometimes contribute to LTP [68, 69]. However, several studies have shown that LTP is often independent of presynaptic changes and thus mainly driven postsynaptically [3, 62, 69]. Therefore, we target the question of whether changes in AMPAR trafficking alone could be sufficient for E-LTP expression. In addition, several types of AMPARs have been identified that may differently contribute to LTP and synaptic transmission [70, 71, 72, 73, 74, 49, 75, 76]. The overall conductance could be adapted by the insertion of different AMPAR types, each having a different conductivity [77]. The calcium-permeable GluR1 homomer (CP-AMPAR) for example has a conductance that is approximately four times higher than that of the GluR1/2 heteromer [77]. However, CP-AMPARs are not required for E-LTP expression [78] such that we limited our model to one type of receptor. Also, AMPARs already present at the PSD before induction might change their conductivity [79]. Stargazin, for example, has been shown to enhance the glutamate affinity and conductance of AMPARs [80, 81]. By contrast, the results from [3] indicate that the fast increase after LTP-induction is associated with the binding of additional AMPARs and possibly not with changes of receptor conductivity. Thus, in this study, we do not consider conductance changes of individual AMPARs. However, different levels of conductivity can be integrated into our model, for instance by weighting the contribution of an AMPAR to the synaptic strength, and by this, the influence of temporal variations in the conductivity by the possible exchange of AMPAR types on LTP could be investigated.

AMPAR mobility We assumed that AMPA receptors are transiently immobilized to become functional at the PSD and otherwise rapidly diffuse between spine and dendrite. This assumption is supported by several experimental studies [10, 11, 82]. However, the experimental literature is not clear regarding the dwell time of immobilized receptors. Studies blocking exocytosis [3, 19, 60] show that synaptic potentiation is very transient, which implies that newly bound AMPARs may unbind from binding sites within seconds or a few minutes and diffuse out of the PSD. Also, single-receptor-tracking studies revealed AMPAR dwell times in immobilized or confined states on the order of seconds [24, 83,

51, 39]. However, the estimates from these studies are biased by the fact that receptors are tracked for only $60 - 200s$. In contrast, studies investigating the fluorescence recovery after photo-bleaching (FRAP) suggest that immobile receptors are trapped for 30 minutes and longer [49, 50, 35]. However, it is not clear whether the pool of immobilized receptors measured in FRAP experiments is affected by LTP-induction. While one study found that the number of immobile GluR1, but not GluR2 increases after 2-photon glutamate uncaging [49], another study did not find a significant increase [4]. Furthermore, single-receptor-tracking as well as FRAP experiments use probes, which could physically interact with the system and, thus, impede the evaluation of receptor dwell times. While the role of Stargazin and MAGUKs such as PSD95 in receptor accumulation has been extensively studied [84, 83, 85, 86, 39], the entirety of mechanisms that drive receptor immobilization during synaptic plasticity is still unclear. In addition to changes in phosphorylation states of PSD proteins, different states of immobilization could be mediated by clustering, as shown in this study, but also by adaptation in steric hindrance/obstacle density via other proteins [87] or some, possibly exocytic, factors that alter the affinity between receptors and binding sites. To match the experimental observation that E-LTP rapidly decays in the absence of exocytosis, we set the time scale for the change in receptor binding k_{UB} to be small, which matches well with the decay of CaMKII activity after LTP induction. However, one should be careful when interpreting these results. Whereas this could be seen as an indicator for rapid dephosphorylation of synaptic proteins such as TARPs, the phosphorylated state may as well outlast CaMKII activity. The rapid decay in EPSP amplitude could also be attributed to the need for an exocytic factor or other processes that may require exocytosis dependent structural plasticity such as translocation of CaMKII or PSD reorganization. More experimental work is required.

PSD growth during LTP   We assumed no change in the number of binding sites during E-LTP, which is consistent with experimental observations showing that an increase in PSD size is protein synthesis dependent and only occurs about 50 minutes after LTP induction [27]. Also the number of PSD95 increases very gradually over the time course of 3 hours after LTP induction and only if the spine enlargement is stable [28]. However, it has been hypothesized that new binding sites may become available due to the un-

binding of the protein SynGAP from PSD95 upon phosphorylation [11, 57]. In addition, the CaMKII-NMDAR complex might mediate the composition of new binding sites [88]. However, more data is required to conclusively quantify the influence of PSD changes on the maintenance of E-LTP.

Correlation between RE trafficking, structural and functional LTP   How sLTP, RE trafficking, and functional LTP are interlinked is still ambiguous. Several experimental studies identified correlations between spine volume and synaptic transmission [5, 6, 33]. For instance, [5] have shown that AMPAR-mediated currents and spine volume increase during LTP are correlated in the very early phase of LTP $(1-5min)$ as well as at later stages $(>30min)$. These results indicate that the enlargement of the spine head might be an essential condition for the expression of LTP. In addition, exocytosis from AMPAR-containing REs is required for spine size maintenance [29, 6], and these are mobilized into the spine upon LTP-induction [38, 29, 30, 31]. Also, exocytosis is required for sustained E-LTP [3]. In this study, we, therefore, assumed that the spine volume is the determinant factor for the number of AMPARs exocytosed from REs during an exocytosis event. However, whether and how exocytosis scales with spine size is unclear and additional experimental data is needed. Also, it is unclear whether exocytosis of AMPARs or other cargo is the major factor shaping receptor accumulation [53] and it has been suggested that REs are required for the regulation of synaptic PSD-95 levels rather than AMPAR recycling [89]. Besides, there have been conflicting results regarding the impact of REs during the initial phase of sLTP [89, 29]. LTP not only influences exocytosis at the spine but also at the dendrite [4, 21]. It is however unclear whether receptors that are exocytosed outside the spine contribute to E-LTP expression. Theoretical studies have shown that the locus of exocytosis close to the PSD is important for synaptic capturing of AMPARs [36, 90]. A sustained increase in the exocytosis rate of GluR2 and GluR3 after LTP induction has been observed experimentally at the dendritic shaft [21]. Such a sustained local increase in dendritic receptor concentration close to a spine might increase the receptor flux (k_{in}) sufficiently to supplant or support RE mediated receptor recycling at spines and thereby contribute to E-LTP expression.

Conclusion

In line with several other studies [42, 43, 45, 39, 36, 31, 30], our results support the importance of cooperative binding, spine morphology, and RE trafficking for LTP. Beyond this, our model predicts that cooperative binding and spine morphology complement each other to sustain E-LTP such that other phases or processes of (late-phase) LTP can set in. As discussed above, the processes involved in the control of AMPAR mobility and trapping, in particular concerning morphological spine changes and exocytosis, remain to be fully elucidated. The model derived in this study provides a foundation for further investigations of these processes and could be applied to other types of synaptic plasticity such as LTD and synaptic scaling or the role of different AMPAR types. Proper functioning of various types of synaptic plasticity is important for learning and memory and dysregulation of processes involved in synaptic plasticity such as AMPAR trafficking have been implicated in different diseases such as Alzheimer's and epilepsy [91]. To understand how deficits in receptor trafficking can evolve into neurodegenerative diseases we need to obtain a profound understanding of how different trafficking mechanisms can influence synaptic transmission.

Materials and Methods

Parameter derivation

All model parameters are listed in Tables 1 and 2. In the following, we will motivate the different parameter choices. Parameter values under basal conditions, i.e. before LTP induction, will be denoted by a superscript index 0.

The *number of AMPARs* in the PSD has been estimated at around $15 - 30$ receptors [52, 92, 48, 39] (the maximum number could be up to 140 receptors [33, 48]). The exact fraction of mobile receptors in the spine is unclear. While some studies report around 50% mobile receptors at spines [23, 83, 35, 93] more recent studies show that $\approx 65 - 90\%$ of the AMPARs are confined in PSD nanodomains [94, 39]. The question remains how many of these confined AMPARs are also contributing to the excitatory transmission or

whether they functionally belong to the mobile pool U in our model. Considering the above here we assume a total number of 30 AMPARs in the steady state with 2/3 bound receptors ($B^* = 20$) and 1/3 mobile receptors ($U^* = 10$); $*$ marks the steady state.

The *exocytosis event rate* within spines has been measured by Patterson et al. [4] who found ≈ 0.11 exocytosis events per minute under basal conditions (k_{exo}^0). It has been estimated that about 51 ± 36 GluR1 subunits are released into the membrane upon an exocytosis event at the dendritic shaft [47]. However, exocytosis events at spines are on average smaller than at the dendritic shaft [4]. Another study by [21] only estimated $10 - 15$ subunits per event. AMPARs are tetrameric, containing either one or two types of subunits. Based on the above findings we estimated the exocytosis event at spines to have a size of about $S_{exo}^0 = 13$ receptors.

The *diffusion-dependent parameters* k_{out} and k_{in} are estimated from FRAP and receptor tracking experiments, assuming that the contribution of exocytosis to the fluorescence recovery is small [4, 49, 16]. The recovery half time in these FRAP experiments ranges from tens of seconds up to 100 seconds depending on the spine and AMPAR subunit type [50, 49, 4, 35, 54]. We, therefore, estimate that diffusing AMPARs enter the spine's membrane from the dendrite at a rate of $k_{in}' \approx 0.02 s^{-1}$. Given that receptor exchange occurs mainly due to diffusion [4, 49], in- and efflux need to be properly balanced in accordance with the steady state value U^*. Therefore, we set $k_{in} = k_{in}' \cdot U^*$. Single receptor tracking and theoretical studies reveal receptor dwell times in confined states at spines that are also on the order of tens of seconds [22, 51, 39, 36]. Therefore $k_{out} = k_{in}' A_{spine}$ is a convenient choice. Alternatively, we will sometimes use the term receptor dwell time $\tau_{dwell} = A_{spine}/k_{out}$.

The *unbinding rate* of AMPARs k_{BU} is estimated based on the dwell time of receptors being in an immobile state at the PSD. Given [24], we set $k_{BU} = 1/10 \, s^{-1}$. We assume that functional AMPARs need to be in an immobilized state, considering that precise alignment of AMPARs with the presynaptic active zone is important for excita-

tory synaptic transmission [95].

The number of binding sites or *slots* P contained in a PSD is not clear. However, the receptor number in spines is much smaller than the number of most scaffolding proteins being in the PSD [52]. Also, the EPSP amplitude increases by about $50 - 200\%$ during LTP-induction, which is consistent with the increase in GluR1 fluorescence [27]. Considering that $\approx 60 - 70\%$ of the receptors are located within PSD Nanodomains and from the assumption that the PSD does not change during E-LTP follows that the PSD is not saturated before induction. Therefore, to account for the 200% increase in AMPARs, we set $P = 3.5 \cdot B^*$.

For the average *spine volume* under basal conditions we assume $V^0_{spine} = 0.08 \mu m^3$ [5, 37]. We calculate the surface area A^0_{spine} from the volume considering spherical spines heads.

Given that we set the number of AMPARs under basal conditions, the *endocytosis rate* k_{endo} and the *binding rate* k^0_{UB} (before LTP induction) can be calculated analytically from the fixed point equation of the basic model:

$$k_{endo} = \frac{A^0_{spine} \left(k^0_{exo} S_{exo} + k_{in} \right) - k_{out} U^*}{U^*},$$ (4)

$$k^0_{UB} = \frac{A^0_{spine} \, k_{BU} \, B^*}{(P - B^*) \, U^*}.$$ (5)

Parameter changes during LTP-induction

LTP-induction triggers a change in the exocytosis event rate k_{exo} ($Events/s$) and the binding rate k_{UB} [4, 3, 7, 84] (see Introduction and Fig 1). The change in both parameters $k_x, x \in \{exo, UB\}$ during LTP-induction is modeled by the following functions:

$$k_x(t) = k^0_x \cdot f(t, C_x = 1),$$ (6)

with

$$f(t) = \left(C_x + A_x \frac{e^{-t/\tau_x^2} - e^{-t/\tau_x^1}}{A_{norm}} \right) \qquad (7)$$

where

$$A_{norm} = \left(\frac{\tau_x^2}{\tau_x^1} \right)^{\frac{\tau_x^1}{\tau_x^1 - \tau_x^2}} - \left(\frac{\tau_x^2}{\tau_x^1} \right)^{\frac{\tau_x^2}{\tau_x^1 - \tau_x^2}} \qquad (8)$$

is a normalizing factor, such that $max((exp(-t/\tau_x^2) - exp(-t/\tau_x^1))/A_{norm}) = 1$. A_x is the factor by which the parameter changes during LTP-induction. τ_x^2 and τ_x^1 are time constants. Upon LTP-induction the exocytosis event rate increases 5-fold for the duration of the stimulus ($\approx 1min.$), as shown in 2-photon-glutamate-uncaging experiments [4]. We therefore assume a rapid increase in the exocytosis event rate k_{exo} with time constants $\tau_{exo}^1 = 25s$ and $\tau_{exo}^2 = 60s$ during E-LTP-induction. Based on [4, 17], we set $A_{exo} = 5$.

During LTP induction, AMPARs are transiently trapped at the synapse [3], indicating that the binding rate k_{UB} is transiently increased. Therefore, we set $\tau_{UB}^1 = 5s$ and $\tau_{UB}^2 = 60s$. Consistent with this, CaMKII activity rapidly increases but decreases within $2\,min$ upon 2-photon glutamate uncaging [7, 96].

There exists no clear experimental data from which the change in amplitude of k_{UB} during LTP induction could be inferred. Thus, we assume that that the increase is large enough to nearly saturate the available slots, which is the case for $A_{UB} = 30$ in the non-cooperative model and $A_{UB} = 5$ in the cooperative model.

Computer simulation and random sampling of the parameter space

For each model version, we used the classical Runge-Kutta method to solve the corresponding set of differential equations.

In addition to the parameter estimates from above, we randomly sampled parameters (Figs 4 and S1 Fig) to evaluate within which parameter regime the model can be matched to the experimental data of E-LTP with and without exocytosis. In this method, parameters of the model are chosen randomly within a predefined regime (Table 3). The outcome of a simulation is compared to the reference experimental data using the following cost

function:

$$\Delta x = \frac{1}{T} \left(\int_0^T |B_{ref}^{E-LTP} - B_{sim}^{E-LTP}| dt + \int_0^T |B_{ref}^{k_{exo}=0} - B_{sim}^{k_{exo}=0}| dt \right), \qquad (9)$$

where B_{ref}^{E-LTP} (with exocytosis), $B_{ref}^{k_{exo}=0}$ (without exocytosis) are the reference data to which we match the model and T is the duration of measurements. The cost function can be interpreted as the time-averaged difference in the number of bound receptors between modeling and experimental data. We considered 2000 trials (different sets of parameter values) and evaluated the best 1%, according to the cost function.

We used as reference the experimental data from [1] and [3] (E-LTP, LTP with blocked exocytosis; Fig 1A), which is given in percentage of the baseline EPSPs. We fitted the EPSP data by the following function:

$$\begin{aligned} EPSP_x(t) = {} & f(t, C_{x,short}, A_{x,short}, T_{x,short}^2, T_{x,short}^1) \\ & + f(t, C_{x,long}, A_{x,long}, T_{x,long}^2, T_{x,long}^1), \end{aligned} \qquad (10)$$

where $f(\cdot)$ is defined by Equation (7) and $x \in \{E - LTP, k_{exo} = 0\}$. The function $EPSP_x(t)$ accounts for a short initial phase as well as for a subsequent long lasting phase. We transferred this data from % of baseline EPSPs into units of receptor numbers by assuming the same baseline number of receptors for the reference as we did for the model ($B_{ref}^{\,x} = \frac{EPSP^x(t)}{100\%} \cdot B^*$).

Model of sLTP

Similar to functional LTP, sLTP can be divided into different phases [5]. Directly after the induction of sLTP the spine volume rapidly increases and subsequently decays (V_{spine}^{short}) but settles at a level above the original baseline V_{spine}^{long}. sLTP and functional LTP are closely correlated and in the absence of protein synthesis the late phase of sLTP (V_{spine}^{long}) decays with a time constants similar to E-LTP [6, 27]. In addition, sLTP depends on exocytosis and rapidly decays after application of Botox [6, 29] (Fig 1C). Furthermore, several experiments suggest that the relative increase in spine volume triggered by LTP-induction ΔV_{spine}^{long} depends on the initial spine volume V_{spine}^0 [5, 34, 6]). We extracted

the data from these studies and did an exponential fit to be able to account for this dependence in our model (Fig 7H):

$$\Delta V_{spine}^{long} = \frac{a \cdot e^{V_{spine}^0/b} + c}{100\%} \tag{11}$$

with parameter values given in the legend of Fig 7H. Based on the findings discussed above, we formalized a model of the spine volume change:

$$
\begin{aligned}
V_{spine}(t) = V_{spine}^0 \cdot [&f(t, 1, 1 + 1.75 \cdot \Delta V_{spine}^{long}, T_1^2, T_2^1) \\
&+ f(t, 0, V_{spine}^{long}, T_3^2, T_4^1)],
\end{aligned}
\tag{12}
$$

where $f(\cdot)$ is defined by Equation (7). The first term on the right hand side describes the initial, transient phase of structural plasticity V_{spine}^{short} that decays within $10 - 20min$ ($T_2^1 = 5s$, $T_1^2 = 250s$). The amplitude depends on ΔV_{spine}^{long}, but it has a lower limit in the absence of a late phase ($1 + 1.75 \cdot \Delta V_{spine}^{long}$) [5]. The second term on the right hand side describes the long lasting second phase V_{spine}^{long} ($T_4^1 = 500s$). When L-LTP is triggered, T_3^2 is very large, i.e. the spine volume does not decay over the time course of hours. In the absence of L-LTP, we set $T_3 = 50min$, which is the decay time constant of E-LTP. If exocytosis is blocked, ΔV_{spine}^{long} is set to zero. Thereby, our model closely matches the time course of sLTP for different spine sizes and conditions (Fig 7G). To account for endosomal trafficking during sLTP we describe the dynamics of the exocytosis event size S_{exo} inside the spine by

$$\frac{dS_{exo}}{dt} = k_{in}^{RE} - k_{out}^{RE} \frac{S_{exo}}{V_{spine}(t)}, \tag{13}$$

where we set $k_{in}^{RE} = 0.1s^{-1}$. From the fixed point equation of (13) we derive $k_{out}^{RE} = k_{in}^{RE} V_{spine}^0 / S_{exo}^* = 0.000615 \, \mu m^3/(\#s)$ (where $S_{exo}^* = S_{exo}^0 = 13$, see methods section on parameter derivation).

Cooperative binding model

Thanks to super-resolution microscopy techniques like STORM and STED, in recent years, several studies revealed that synaptic proteins in the PSD are not homogeneously distributed but often clustered in so-called nanodomains [26, 39, 97, 40]. Several theo-

Fig 7. Functional relations in the stochastic grid model of cooperative binding and spine size dependence of sLTP. (A) The slope $m(P)$ of the relation between the mean cooperative binding rate $\langle \hat{k}_{UB} \rangle$ and the number of bound receptors B depends on P, as on small grids the mean number of nearest neighbours increases faster than on large grids. **(B,C)** The mean unbinding rate $\langle \hat{k}_{BU} \rangle$ decreases with B with functions $\lambda(P)$ (B) and $\beta(P)$ (C). **(D)** Example traces of the number of bound AMPARs B for different concentrations U/A_{spine} and PSD sizes P. **(E)** The steady state value of B depends linear on the PSD size P. The mean-field model (dotted lines) matches the stochastic model (solid lines). Bars indicate standard deviations. **(F)** Example of a cluster realization in the stochastic model where white squares denote occupied slots. **(G)** Spine volume increase during sLTP in the model (colored lines) compared to experimental data taken from [5, 6]. **(H)** The relative increase in spine volume depends on the initial spine size. The data has been fitted by an exponential function (Equation 11) with parameters shown in the legend. The shade indicates the standard deviation across fits to the different data sets taken from [6, 5, 34].

625 retical studies identified that clusters can emerge due to cooperative binding of proteins

626 [42, 43, 44, 98, 41, 45]. PSD scaffolding proteins such as PSD95 form nanodomains on

627 which AMPARs are not uniformly distributed but form smaller clusters themselves [39],

indicating cooperative binding of receptors.

To extend the basic model by cooperative binding of AMPARs in the PSD we utilize a recent theoretical model capturing the essence of cooperativity given a few general assumptions [45]. In our adaptation of this model the scaffolding proteins or slots are spatially organized on a grid (Fig 3A) where each mobile receptor U binds to a free slot with a certain rate, dependent on the number of occupied nearest-neighbor slots (Equation (14)). Similarly, the rate of unbinding decreases with more bound receptors B being in the surrounding (Equation (15)). Note that in this model the grid size ($N \times N$) is analog to P in the basic model (Equations (1) and (2)).

$$\hat{k}_{UB}^{coop} = k_{UB}(\alpha\chi + 1) \tag{14}$$

$$\hat{k}_{BU}^{coop} = k_{BU}(1 - \chi). \tag{15}$$

Here, χ is the fraction of occupied nearest neighbours on the grid (i.e. $n/8$ with $n = 0, 1, ..., 8$) and α is a measure for the cooperativity, where a big value coincides with a stronger influence of cooperativity. Taking into account the dependence on the mobile receptors concentration the total binding flux is $\hat{k}_{UB}^{coop} \frac{U}{A_{spine}^0}$. As apparent from Equations (14) and (15), the cooperative binding and unbinding rate $\hat{k}_{UB}^{coop}$, $\hat{k}_{BU}^{coop}$ change depending on the number of bound receptors on the grid and depending on the size of the grid P. We did a stochastic simulation of this model (as in [45]) with the binding and unbinding probabilities being $p_{UB}^{coop} = \hat{k}_{UB}^{coop} \cdot \Delta t$ and $p_{BU} = \hat{k}_{BU}^{coop} \cdot \Delta t$ with the integration time step $\Delta t = 0.5\,s$. In addition, we describe U by

$$dU = \left(k_{in} + k_{exo}\,S_{exo} - (k_{out} + k_{endo})\,\frac{U}{A_{spine}^0}\right)dt + dB_{off} - dB_{on}, \tag{16}$$

where dB_{off} and dB_{on} are the receptors unbinding/binding from/to the grid respectively within one time step of the stochastic simulation. To control the number of mobile receptors, and thereby, also the number of bound receptors on the grid, we vary the influx k_{in} between $0.1 - 1.2\,\#/s$. The binding parameter were set to $k_{UB} = 0.0005(\#s)^{-1}$, $\alpha = 16$ and $k_{BU} = 0.1s^{-1}$.

To integrate the key principles of the cooperative model into the basic model of AMPAR-dynamics (Equations (1) and (2)), we evaluated several iterations of the cooperative model for different numbers of available receptors U and slots P (grid size). From these, we obtained the mean binding $\langle \hat{k}_{UB}^{coop} \rangle$ and unbinding rate $\langle \hat{k}_{BU}^{coop} \rangle$ per mobile/bound receptor averaged across the grid and and across many realizations and their dependency on number of slots P and bound receptors B (Fig 3B).

$$\langle \hat{k}_{UB}^{coop} \rangle = \frac{1}{N} \sum_{i=0}^{N} \frac{1}{P - B_i} \sum_{grid=0} \hat{k}_{UB,i}^{coop} \ , \tag{17}$$

$$\langle \hat{k}_{BU}^{coop} \rangle = \frac{1}{N} \sum_{i=0}^{N} \frac{1}{B_i} \sum_{grid=1} \hat{k}_{BU,i}^{coop} \ , \tag{18}$$

where $grid = 0$ and $grid = 1$ denote the elements of the grid that are empty or occupied respectively and N the number of realizations. Note that the choices for k_{UB} and k_{BU} do not influence the dependence of $\langle \hat{k}_{UB}^{coop} \rangle$ or $\langle \hat{k}_{BU}^{coop} \rangle$ on P or B.

We then formalized the dependency of binding and unbinding rate on P and B by fitting $\langle \hat{k}_{UB}^{coop} \rangle$ and $\langle \hat{k}_{BU}^{coop} \rangle$ by the following functions (Fig 3B dashed lines):

$$\frac{k_{BU}^{coop}}{k_{BU}} = \frac{\lambda(P)}{\beta(P) + B} - 0.5, \tag{19}$$

$$\frac{k_{UB}^{coop}}{k_{UB}} = m(P)B^{0.8} + 1 \tag{20}$$

with functions

$$\beta = m_\beta P + c_\beta, \tag{21}$$

$$\lambda = m_\lambda P + c_\lambda, \tag{22}$$

$$m = a_m/(P + b_m). \tag{23}$$

$\beta(P)$, $\lambda(P)$ and $m(P)$ are shown in Figs 7A-C. By this, we eliminate the spatial component of the cooperative binding model and obtain a mean-field description that we can integrate

into our basic model by substituting k_{UB} and k_{BU} in Equations (1) and (2) by k_{UB}^{coop} and k_{UB}^{coop} (Fig 3B). The mean-field model closely matches the stochastic spatial clustering model's static and dynamic properties (Figs 6A, 7E, and 3A inset).

Steady state binding rate in the cooperative model

The dynamics of cooperative binding changes the steady state of the system. Therefore, the binding rate has to be derived from the the adapted fixed point equation resulting to:

$$k_{UB}^0 = \frac{A_{spine}^0 \, B^* \, k_{BU} \left(\frac{c_\lambda + m_\lambda P}{B^* + c_\beta + m_\beta P} - 0.5 \right)}{(P - B^*) \left(1 + \frac{a_m \, (B^*)^{0.8}}{bm + P} \right) U^*}. \tag{24}$$

Code Accessibility

Code and documentation are available on GitHub `https://github.com/MoritzB90/AMPAR-Trafficking-Model`.

References

[1] Barco A, Alarcon JM, Kandel ER. Expression of Constitutively Active CREB Protein Facilitates the Late Phase of Long-Term Potentiation by Enhancing Synaptic Capture. Cell. 2002;108(5):689–703. doi:10.1016/s0092-8674(02)00657-8.

[2] Abraham WC. How long will long-term potentiation last? Philosophical Transactions of the Royal Society of London Series B: Biological Sciences. 2003;358(1432):735–744. doi:10.1098/rstb.2002.1222.

[3] Penn AC, Zhang CL, Georges F, Royer L, Breillat C, Hosy E, et al. Hippocampal LTP and contextual learning require surface diffusion of AMPA receptors. Nature. 2017;549(7672):384–388. doi:10.1038/nature23658.

[4] Patterson MA, Szatmari EM, Yasuda R. AMPA receptors are exocytosed in stimulated spines and adjacent dendrites in a Ras-ERK-dependent manner during long-term potentiation. Proceedings of the National Academy of Sciences. 2010;107(36):15951–15956. doi:10.1073/pnas.0913875107.

[5] Matsuzaki M, Honkura N, Ellis-Davies GCR, Kasai H. Structural basis of long-term potentiation in single dendritic spines. Nature. 2004;429(6993):761–766. doi:10.1038/nature02617.

[6] Yang Y, b Wang X, Frerking M, Zhou Q. Spine Expansion and Stabilization Associated with Long-Term Potentiation. Journal of Neuroscience. 2008;28(22):5740–5751. doi:10.1523/jneurosci.3998-07.2008.

[7] Lee SJR, Escobedo-Lozoya Y, Szatmari EM, Yasuda R. Activation of CaMKII in single dendritic spines during long-term potentiation. Nature. 2009;458(7236):299–304. doi:10.1038/nature07842.

[8] Anggono V, Huganir RL. Regulation of AMPA receptor trafficking and synaptic plasticity. Current Opinion in Neurobiology. 2012;22(3):461–469. doi:10.1016/j.conb.2011.12.006.

[9] Lisman J, Yasuda R, Raghavachari S. Mechanisms of CaMKII action in long-term potentiation. Nature Reviews Neuroscience. 2012;13(3):169–182. doi:10.1038/nrn3192.

[10] Opazo P, Sainlos M, Choquet D. Regulation of AMPA receptor surface diffusion by PSD-95 slots. Current Opinion in Neurobiology. 2012;22(3):453–460. doi:10.1016/j.conb.2011.10.010.

[11] Choquet D. Linking Nanoscale Dynamics of AMPA Receptor Organization to Plasticity of Excitatory Synapses and Learning. The Journal of Neuroscience. 2018;38(44):9318–9329. doi:10.1523/jneurosci.2119-18.2018.

[12] Frey U, Morris RGM. Synaptic tagging and long-term potentiation. Nature. 1997;385(6616):533–536. doi:10.1038/385533a0.

[13] Bliss TVP, Collingridge GL, Morris RGM, Reymann KG. Long-term potentiation in the hippocampus: discovery, mechanisms and function. Neuroforum. 2018;24(3):A103–A120. doi:10.1515/nf-2017-a059.

[14] Malinow R, Malenka RC. AMPA Receptor Trafficking and Synap-

tic Plasticity. Annual Review of Neuroscience. 2002;25(1):103–126.
doi:10.1146/annurev.neuro.25.112701.142758.

[15] Gallimore AR, Kim T, Tanaka-Yamamoto K, Schutter ED. Switching On De-
pression and Potentiation in the Cerebellum. Cell Reports. 2018;22(3):722–733.
doi:10.1016/j.celrep.2017.12.084.

[16] Passafaro M, Piëch V, Sheng M. Subunit-specific temporal and spatial pat-
terns of AMPA receptor exocytosis in hippocampal neurons. Nature Neuroscience.
2001;4(9):917–926. doi:10.1038/nn0901-917.

[17] Kennedy MJ, Davison IG, Robinson CG, Ehlers MD. Syntaxin-4 Defines a Domain
for Activity-Dependent Exocytosis in Dendritic Spines. Cell. 2010;141(3):524–535.
doi:10.1016/j.cell.2010.02.042.

[18] Choquet D, Triller A. The Dynamic Synapse. Neuron. 2013;80(3):691–703.
doi:10.1016/j.neuron.2013.10.013.

[19] Wu D, Bacaj T, Morishita W, Goswami D, Arendt KL, Xu W, et al. Postsy-
naptic synaptotagmins mediate AMPA receptor exocytosis during LTP. Nature.
2017;544(7650):316–321. doi:10.1038/nature21720.

[20] Lledo PM, Zhang X, Südhof TC, Malenka RC, Nicoll RA. Postsynaptic Mem-
brane Fusion and Long-Term Potentiation. Science. 1998;279(5349):399–403.
doi:10.1126/science.279.5349.399.

[21] Tanaka H, Hirano T. Visualization of Subunit-Specific Delivery of Glutamate Recep-
tors to Postsynaptic Membrane during Hippocampal Long-Term Potentiation. Cell
Reports. 2012;1(4):291–298. doi:10.1016/j.celrep.2012.02.004.

[22] Borgdorff AJ, Choquet D. Regulation of AMPA receptor lateral movements. Nature.
2002;417(6889):649–653. doi:10.1038/nature00780.

[23] Tardin C, Cognet L, Bats C, Lounis B, Choquet D. Direct imaging of lateral move-
ments of AMPA receptors inside synapses. The EMBO Journal. 2003;22(18):4656–
4665. doi:10.1093/emboj/cdg463.

[24] Heine M, Groc L, Frischknecht R, que JCB, Lounis B, Rumbaugh G, et al. Surface Mobility of Postsynaptic AMPARs Tunes Synaptic Transmission. Science. 2008;320(5873):201–205. doi:10.1126/science.1152089.

[25] Huganir RL, Nicoll RA. AMPARs and Synaptic Plasticity: The Last 25 Years. Neuron. 2013;80(3):704–717. doi:10.1016/j.neuron.2013.10.025.

[26] MacGillavry HD, Song Y, Raghavachari S, Blanpied TA. Nanoscale Scaffolding Domains within the Postsynaptic Density Concentrate Synaptic AMPA Receptors. Neuron. 2013;78(4):615–622. doi:10.1016/j.neuron.2013.03.009.

[27] Bosch M, Castro J, Saneyoshi T, Matsuno H, Sur M, Hayashi Y. Structural and Molecular Remodeling of Dendritic Spine Substructures during Long-Term Potentiation. Neuron. 2014;82(2):444–459. doi:10.1016/j.neuron.2014.03.021.

[28] Meyer D, Bonhoeffer T, Scheuss V. Balance and Stability of Synaptic Structures during Synaptic Plasticity. Neuron. 2014;82(2):430–443. doi:10.1016/j.neuron.2014.02.031.

[29] Park M, Salgado JM, Ostroff L, Helton TD, Robinson CG, Harris KM, et al. Plasticity-Induced Growth of Dendritic Spines by Exocytic Trafficking from Recycling Endosomes. Neuron. 2006;52(5):817–830. doi:10.1016/j.neuron.2006.09.040.

[30] Park M, Penick EC, Edwards JG, Kauer JA, Ehlers MD. Recycling Endosomes Supply AMPA Receptors for LTP. Science. 2004;305(5692):1972–1975. doi:10.1126/science.1102026.

[31] Wang Z, Edwards JG, Riley N, Provance DW, Karcher R, dong Li X, et al. Myosin Vb Mobilizes Recycling Endosomes and AMPA Receptors for Postsynaptic Plasticity. Cell. 2008;135(3):535–548. doi:10.1016/j.cell.2008.09.057.

[32] Zheng N, Jeyifous O, Munro C, Montgomery JM, Green WN. Synaptic activity regulates AMPA receptor trafficking through different recycling pathways. eLife. 2015;4. doi:10.7554/elife.06878.

[33] Matsuzaki M, Ellis-Davies GCR, Nemoto T, Miyashita Y, Iino M, Kasai H. Dendritic spine geometry is critical for AMPA receptor expression in hippocampal CA1 pyramidal neurons. Nature Neuroscience. 2001;4(11):1086–1092. doi:10.1038/nn736.

[34] Tanaka J, Horiike Y, Matsuzaki M, Miyazaki T, Ellis-Davies GCR, Kasai H. Protein Synthesis and Neurotrophin-Dependent Structural Plasticity of Single Dendritic Spines. Science. 2008;319(5870):1683–1687. doi:10.1126/science.1152864.

[35] Ashby MC, Maier SR, Nishimune A, Henley JM. Lateral Diffusion Drives Constitutive Exchange of AMPA Receptors at Dendritic Spines and Is Regulated by Spine Morphology. Journal of Neuroscience. 2006;26(26):7046–7055. doi:10.1523/jneurosci.1235-06.2006.

[36] Kusters R, Kapitein LC, Hoogenraad CC, Storm C. Shape-Induced Asymmetric Diffusion in Dendritic Spines Allows Efficient Synaptic AMPA Receptor Trapping. Biophysical Journal. 2013;105(12):2743–2750. doi:10.1016/j.bpj.2013.11.016.

[37] Adrian M, Kusters R, Storm C, Hoogenraad CC, Kapitein LC. Probing the Interplay between Dendritic Spine Morphology and Membrane-Bound Diffusion. Biophysical Journal. 2017;113(10):2261–2270. doi:10.1016/j.bpj.2017.06.048.

[38] Kneussel M, Hausrat TJ. Postsynaptic Neurotransmitter Receptor Reserve Pools for Synaptic Potentiation. Trends in Neurosciences. 2016;39(3):170–182. doi:10.1016/j.tins.2016.01.002.

[39] Nair D, Hosy E, Petersen JD, Constals A, Giannone G, Choquet D, et al. Super-Resolution Imaging Reveals That AMPA Receptors Inside Synapses Are Dynamically Organized in Nanodomains Regulated by PSD95. Journal of Neuroscience. 2013;33(32):13204–13224. doi:10.1523/jneurosci.2381-12.2013.

[40] Hruska M, Henderson N, Marchand SJL, Jafri H, Dalva MB. Synaptic nanomodules underlie the organization and plasticity of spine synapses. Nature Neuroscience. 2018;21(5):671–682. doi:10.1038/s41593-018-0138-9.

[41] Ranft J, Almeida LG, Rodriguez PC, Triller A, Hakim V. An aggregation-removal model for the formation and size determination of post-synaptic

scaffold domains. PLOS Computational Biology. 2017;13(4):e1005516. doi:10.1371/journal.pcbi.1005516.

[42] Shouval HZ. Clusters of interacting receptors can stabilize synaptic efficacies. Proceedings of the National Academy of Sciences. 2005;102(40):14440–14445. doi:10.1073/pnas.0506934102.

[43] Burlakov VM, Emptage N, Goriely A, Bressloff PC. Synaptic Bistability Due to Nucleation and Evaporation of Receptor Clusters. Physical Review Letters. 2012;108(2). doi:10.1103/physrevlett.108.028101.

[44] Sekimoto K, Triller A. Compatibility between itinerant synaptic receptors and stable postsynaptic structure. Physical Review E. 2009;79(3). doi:10.1103/physreve.79.031905.

[45] Shomar A, Geyrhofer L, Ziv NE, Brenner N. Cooperative stochastic binding and unbinding explain synaptic size dynamics and statistics. PLOS Computational Biology. 2017;13(7):e1005668. doi:10.1371/journal.pcbi.1005668.

[46] Earnshaw BA, Bressloff PC. Biophysical Model of AMPA Receptor Trafficking and Its Regulation during Long-Term Potentiation/Long-Term Depression. Journal of Neuroscience. 2006;26(47):12362–12373. doi:10.1523/jneurosci.3601-06.2006.

[47] Yudowski GA, Puthenveedu MA, Leonoudakis D, Panicker S, Thorn KS, Beattie EC, et al. Real-Time Imaging of Discrete Exocytic Events Mediating Surface Delivery of AMPA Receptors. Journal of Neuroscience. 2007;27(41):11112–11121. doi:10.1523/jneurosci.2465-07.2007.

[48] Nusser Z, Lujan R, Laube G, Roberts JDB, Molnar E, Somogyi P. Cell Type and Pathway Dependence of Synaptic AMPA Receptor Number and Variability in the Hippocampus. Neuron. 1998;21(3):545–559. doi:10.1016/s0896-6273(00)80565-6.

[49] Makino H, Malinow R. AMPA Receptor Incorporation into Synapses during LTP: The Role of Lateral Movement and Exocytosis. Neuron. 2009;64(3):381–390. doi:10.1016/j.neuron.2009.08.035.

[50] Frischknecht R, Heine M, Perrais D, Seidenbecher CI, Choquet D, Gundelfinger ED. Brain extracellular matrix affects AMPA receptor lateral mobility and short-term synaptic plasticity. Nature Neuroscience. 2009;12(7):897–904. doi:10.1038/nn.2338.

[51] Ehlers MD, Heine M, Groc L, Lee MC, Choquet D. Diffusional Trapping of GluR1 AMPA Receptors by Input-Specific Synaptic Activity. Neuron. 2007;54(3):447–460. doi:10.1016/j.neuron.2007.04.010.

[52] Sheng M, Hoogenraad CC. The Postsynaptic Architecture of Excitatory Synapses: A More Quantitative View. Annual Review of Biochemistry. 2007;76(1):823–847. doi:10.1146/annurev.biochem.76.060805.160029.

[53] Hiester BG, Becker MI, Bowen AB, Schwartz SL, Kennedy MJ. Mechanisms and Role of Dendritic Membrane Trafficking for Long-Term Potentiation. Frontiers in Cellular Neuroscience. 2018;12. doi:10.3389/fncel.2018.00391.

[54] Tatavarty V, Sun Q, Turrigiano GG. How to Scale Down Postsynaptic Strength. Journal of Neuroscience. 2013;33(32):13179–13189. doi:10.1523/jneurosci.1676-13.2013.

[55] Hanse E, Gustafsson B. Quantal variability at glutamatergic synapses in area CA1 of the rat neonatal hippocampus. The Journal of Physiology. 2001;531(2):467–480. doi:10.1111/j.1469-7793.2001.0467i.x.

[56] Barbour B, Brunel N, Hakim V, Nadal JP. What can we learn from synaptic weight distributions? Trends in Neurosciences. 2007;30(12):622–629. doi:10.1016/j.tins.2007.09.005.

[57] Nicoll RA. A Brief History of Long-Term Potentiation. Neuron. 2017;93(2):281–290. doi:10.1016/j.neuron.2016.12.015.

[58] Herring BE, Nicoll RA. Long-Term Potentiation: From CaMKII to AMPA Receptor Trafficking. Annual Review of Physiology. 2016;78(1):351–365. doi:10.1146/annurev-physiol-021014-071753.

[59] Redondo RL, Morris RGM. Making memories last: the synaptic tagging and capture hypothesis. Nature Reviews Neuroscience. 2010;12(1):17–30. doi:10.1038/nrn2963.

[60] Lüscher C, Xia H, Beattie EC, Carroll RC, von Zastrow M, Malenka RC, et al. Role of AMPA Receptor Cycling in Synaptic Transmission and Plasticity. Neuron. 1999;24(3):649–658. doi:10.1016/s0896-6273(00)81119-8.

[61] Lamprecht R, LeDoux J. Structural plasticity and memory. Nat Rev Neurosci. 2004;5:45–54. doi:10.1038/nrn1301.

[62] Luscher C, Malenka RC. NMDA Receptor-Dependent Long-Term Potentiation and Long-Term Depression (LTP/LTD). Cold Spring Harbor Perspectives in Biology. 2012;4(6):a005710–a005710. doi:10.1101/cshperspect.a005710.

[63] Andrásfalvy BK, Magee JC. Changes in AMPA receptor currents following LTP induction on rat CA1 pyramidal neurones. The Journal of Physiology. 2004;559(2):543–554. doi:10.1113/jphysiol.2004.065219.

[64] Collingridge GL, Isaac JTR, Wang YT. Receptor trafficking and synaptic plasticity. Nature Reviews Neuroscience. 2004;5(12):952–962. doi:10.1038/nrn1556.

[65] Isaac JTR, Nicoll RA, Malenka RC. Evidence for silent synapses: Implications for the expression of LTP. Neuron. 1995;15(2):427–434. doi:10.1016/0896-6273(95)90046-2.

[66] Franks KM, Stevens CF, Sejnowski TJ. Independent Sources of Quantal Variability at Single Glutamatergic Synapses. The Journal of Neuroscience. 2003;23(8):3186–3195. doi:10.1523/jneurosci.23-08-03186.2003.

[67] Tang AH, Chen H, Li TP, Metzbower SR, MacGillavry HD, Blanpied TA. A trans-synaptic nanocolumn aligns neurotransmitter release to receptors. Nature. 2016;536(7615):210–214. doi:10.1038/nature19058.

[68] Padamsey Z, Emptage N. Two sides to long-term potentiation: a view towards reconciliation. Philosophical Transactions of the Royal Society B: Biological Sciences. 2014;369(1633):20130154. doi:10.1098/rstb.2013.0154.

[69] Bliss TV, Collingridge GL. Expression of NMDA receptor-dependent LTP in the hippocampus: bridging the divide. Molecular Brain. 2013;6(1):5. doi:10.1186/1756-6606-6-5.

[70] Zamanillo D, Sprengel R, Hvalby O, Jensen V, Burnashev N, Rozov A, et al. Importance of AMPA Receptors for Hippocampal Synaptic Plasticity But Not for Spatial Learning. Science. 1999;284(5421):1805–1811. doi:10.1126/science.284.5421.1805.

[71] Hayashi Y, Shi SH, Esteban JA, Piccini A, Poncer JC, Malinow R. Driving AMPA Receptors into Synapses by LTP and CaMKII: Requirement for GluR1 and PDZ Domain Interaction. Science. 2000;287(5461):2262–2267. doi:10.1126/science.287.5461.2262.

[72] Shi SH, Hayashi Y, Esteban JA, Malinow R. Subunit-Specific Rules Governing AMPA Receptor Trafficking to Synapses in Hippocampal Pyramidal Neurons. Cell. 2001;105(3):331–343. doi:10.1016/s0092-8674(01)00321-x.

[73] Meng Y, Zhang Y, Jia Z. Synaptic Transmission and Plasticity in the Absence of AMPA Glutamate Receptor GluR2 and GluR3. Neuron. 2003;39(1):163–176. doi:10.1016/s0896-6273(03)00368-4.

[74] Plant K, Pelkey KA, Bortolotto ZA, Morita D, Terashima A, McBain CJ, et al. Transient incorporation of native GluR2-lacking AMPA receptors during hippocampal long-term potentiation. Nature Neuroscience. 2006;9(5):602–604. doi:10.1038/nn1678.

[75] Isaac JTR, Ashby MC, McBain CJ. The Role of the GluR2 Subunit in AMPA Receptor Function and Synaptic Plasticity. Neuron. 2007;54(6):859–871. doi:10.1016/j.neuron.2007.06.001.

[76] Park P, Sanderson TM, Amici M, Choi SL, Bortolotto ZA, Zhuo M, et al. Calcium-Permeable AMPA Receptors Mediate the Induction of the Protein Kinase A-Dependent Component of Long-Term Potentiation in the Hippocampus. Journal of Neuroscience. 2016;36(2):622–631. doi:10.1523/jneurosci.3625-15.2016.

[77] Benke T, Traynelis SF. AMPA-Type Glutamate Receptor Conductance Changes and Plasticity: Still a Lot of Noise. Neurochemical Research. 2018;44(3):539–548. doi:10.1007/s11064-018-2491-1.

[78] Park P, Kang H, Sanderson TM, Bortolotto ZA, Georgiou J, Zhuo M, et al. The Role of Calcium-Permeable AMPARs in Long-Term Potentiation at Principal Neurons in the Rodent Hippocampus. Frontiers in Synaptic Neuroscience. 2018;10. doi:10.3389/fnsyn.2018.00042.

[79] Benke TA, Lüthi A, Isaac JTR, Collingridge GL. Modulation of AMPA receptor unitary conductance by synaptic activity. Nature. 1998;393(6687):793–797. doi:10.1038/31709.

[80] Tomita S, Adesnik H, Sekiguchi M, Zhang W, Wada K, Howe JR, et al. Stargazin modulates AMPA receptor gating and trafficking by distinct domains. Nature. 2005;435(7045):1052–1058. doi:10.1038/nature03624.

[81] Turetsky D, Garringer E, Patneau DK. Stargazin Modulates Native AMPA Receptor Functional Properties by Two Distinct Mechanisms. Journal of Neuroscience. 2005;25(32):7438–7448. doi:10.1523/jneurosci.1108-05.2005.

[82] Hastings MH, Man HY. Synaptic Capture of Laterally Diffusing AMPA Receptors – An Idea That Stuck. Trends in Neurosciences. 2018;41(6):330–332. doi:10.1016/j.tins.2018.03.016.

[83] Bats C, Groc L, Choquet D. The Interaction between Stargazin and PSD-95 Regulates AMPA Receptor Surface Trafficking. Neuron. 2007;53(5):719–734. doi:10.1016/j.neuron.2007.01.030.

[84] Opazo P, Labrecque S, Tigaret CM, Frouin A, Wiseman PW, Koninck PD, et al. CaMKII Triggers the Diffusional Trapping of Surface AMPARs through Phosphorylation of Stargazin. Neuron. 2010;67(2):239–252. doi:10.1016/j.neuron.2010.06.007.

[85] Schnell E, Sizemore M, Karimzadegan S, Chen L, Bredt DS, Nicoll RA. Direct interactions between PSD-95 and stargazin control synaptic AMPA receptor number. Proceedings of the National Academy of Sciences. 2002;99(21):13902–13907. doi:10.1073/pnas.172511199.

[86] Tomita S, Stein V, Stocker TJ, Nicoll RA, Bredt DS. Bidirectional Synaptic Plasticity

Regulated by Phosphorylation of Stargazin-like TARPs. Neuron. 2005;45(2):269–277. doi:10.1016/j.neuron.2005.01.009.

[87] Kokolaki ML, Fauquier A, Renner M. Molecular Crowding and Diffusion-Capture in Synapses. iScience. 2020;23(8):101382. doi:10.1016/j.isci.2020.101382.

[88] Sanhueza M, Lisman J. The CaMKII/NMDAR complex as a molecular memory. Molecular Brain. 2013;6(1):10. doi:10.1186/1756-6606-6-10.

[89] da Silva ME, Adrian M, Schätzle P, Lipka J, Watanabe T, Cho S, et al. Positioning of AMPA Receptor-Containing Endosomes Regulates Synapse Architecture. Cell Reports. 2015;13(5):933–943. doi:10.1016/j.celrep.2015.09.062.

[90] Czondor K, Mondin M, Garcia M, Heine M, Frischknecht R, Choquet D, et al. Unified quantitative model of AMPA receptor trafficking at synapses. Proceedings of the National Academy of Sciences. 2012;109(9):3522–3527. doi:10.1073/pnas.1109818109.

[91] Chater TE, Goda Y. The role of AMPA receptors in postsynaptic mechanisms of synaptic plasticity. Frontiers in Cellular Neuroscience. 2014;8. doi:10.3389/fncel.2014.00401.

[92] Cheng D, Hoogenraad CC, Rush J, Ramm E, Schlager MA, Duong DM, et al. Relative and Absolute Quantification of Postsynaptic Density Proteome Isolated from Rat Forebrain and Cerebellum. Molecular & Cellular Proteomics. 2006;5(6):1158–1170. doi:10.1074/mcp.d500009-mcp200.

[93] Petrini EM, Lu J, Cognet L, Lounis B, Ehlers MD, Choquet D. Endocytic Trafficking and Recycling Maintain a Pool of Mobile Surface AMPA Receptors Required for Synaptic Potentiation. Neuron. 2009;63(1):92–105. doi:10.1016/j.neuron.2009.05.025.

[94] Lee SH, Jin C, Cai E, Ge P, Ishitsuka Y, Teng KW, et al. Super-resolution imaging of synaptic and Extra-synaptic AMPA receptors with different-sized fluorescent probes. eLife. 2017;6. doi:10.7554/elife.27744.

[95] Haas KT, Compans B, Letellier M, Bartol TM, Grillo-Bosch D, Sejnowski TJ, et al.

Pre-post synaptic alignment through neuroligin-1 tunes synaptic transmission efficiency. eLife. 2018;7. doi:10.7554/elife.31755.

[96] Chang JY, Parra-Bueno P, Laviv T, Szatmari EM, Lee SJR, Yasuda R. CaMKII Autophosphorylation Is Necessary for Optimal Integration of Ca 2+ Signals during LTP Induction, but Not Maintenance. Neuron. 2017;94(4):800–808.e4. doi:10.1016/j.neuron.2017.04.041.

[97] Compans B, Choquet D, Hosy E. Review on the role of AMPA receptor nano-organization and dynamic in the properties of synaptic transmission. Neurophotonics. 2016;3(4):041811. doi:10.1117/1.nph.3.4.041811.

[98] Haselwandter CA, Calamai M, Kardar M, Triller A, da Silveira RA. Formation and Stability of Synaptic Receptor Domains. Physical Review Letters. 2011;106(23). doi:10.1103/physrevlett.106.238104.

Supporting information

S1 Fig. The parameter values of exocytosis for the best 0.5% parameter sets depend on the number of bound AMPARs under basal condition and the dwell time of mobile receptors at the spine. For the three models of AMPAR-dynamics (first column: basic model; second column: sLTP model; third column: cooperative binding model), the exocytosis event rate under basal condition (k_{exo}^0; first row), the maximal amplitude of LTP-induced exocytosis event rate ($k_{exo}^A = k_{exo}^0(1 + A_{exo})$; second row), and the decay time of the exocytosis event rate after LTP-induction (τ_{exo}; third row) are shown. Black lines depict standard deviations. The red areas indicate the biologically plausible regime of the corresponding indicators based on experimental studies [16, 47, 4].