Construction of a complete set of *Neisseria meningitidis* defined mutants – the NeMeSys 2.0 collection – and its use for the phenotypic profiling of the genome of an important human pathogen

Alastair Muir[1], Ishwori Gurung[1], Ana Cehovin[1], Adelme Bazin[2], David Vallenet[2], Vladimir Pelicic[1,*]

[1]MRC Centre for Molecular Bacteriology and Infection, Imperial College London, London, United Kingdom.

[2]LABGeM, Génomique Métabolique, CEA, Genoscope, Institut François Jacob, Université d'Evry, Université Paris-Saclay, CNRS, Evry, France.

*Corresponding author

email: v.pelicic@imperial.ac.uk

Abstract

One of the great challenges in biology is to determine the function of millions of genes of unknown function. Even in model bacterial species, there is a sizeable proportion of such genes, which has fundamental and practical consequences. Here, we constructed a complete collection of defined mutants in protein-coding genes – named NeMeSys 2.0 – in the human pathogen *Neisseria meningitidis*, consisting of individual mutants in 1,584 non-essential genes. This effort identified 391 essential genes – broadly conserved in other bacteria – leading to a full overview of the essential meningococcal genome, associated with just four underlying basic biological functions: 1) expression of genome information, 2) preservation of genome information, 3) cell membrane structure/function, and 4) cytosolic metabolism. Subsequently, we illustrated the utility of the NeMeSys 2.0 collection for determining gene function by identifying 1) a novel and conserved family of histidinol-phosphatase, 2) 20 genes, including three new ones, involved in the biology of type IV pili, a widespread virulence factor, and 3) several conditionally essential genes found in regions of genome plasticity encoding antitoxins and/or immunity proteins, which become dispensable when the gene encoding the cognate toxin is deleted. These findings have implications beyond the meningococcus. The NeMeSys 2.0 collection is an invaluable resource paving the way for a global phenotypic landscape in a major human bacterial pathogen.

Introduction

Low-cost massively parallel sequencing methods have transformed biology, with availability of complete genome sequences increasing exponentially. In bacteria, genome sequences can be determined for hundreds of isolates in parallel in just a matter of days. For bacterial pathogens, genome sequences are often available for thousands of clinical isolates, and represent a key resource for epidemiology and comparative genomics[1]. This explosion has also led to the identification of millions of new genes, with the important insight that many are of unknown function. Even in a model species such as *Escherichia coli* K-12, perhaps the most thoroughly studied biological entity, only 60 % of 4,623 protein-coding genes have experimentally verified functions[2], while 35 % have either no predicted function, or a predicted function that is yet to be verified[2]. In less well studied species, this partition is dramatically skewed towards the latter class, with an overwhelming majority of genes having predicted but not experimentally verified functions, or no predicted function at all. This indicates that there are still major gaps in our understanding of basic bacterial physiology, including for well-studied metabolic pathways[3]. This has fundamental consequences for systems and synthetic biology, by hindering holistic understanding of the life-supporting functions and processes allowing bacteria to grow and replicate. It also has practical consequences by slowing down the identification of new means for controlling bacterial pathogens, which are needed more than ever in this era of antimicrobial resistance. Therefore, developing methods for systematically elucidating gene function is a critical biological endeavour.

Molecular genetics, which involves the creation and phenotypic analysis of large collections of mutants, is the most direct path to determining gene function on a genome scale by providing a crucial link between phenotype and genotype. In many bacteria, genome-wide mutant collections in protein-coding genes have been constructed using either targeted mutagenesis or random transposon (Tn) mutagenesis. Because of its time and cost-effectiveness, Tn mutagenesis is more

widely used[4-6], especially since it has been coupled with massively parallel sequencing of Tn insertion sites (Tn-Seq)[7], which allows simultaneous monitoring of the fitness in pools of hundreds of thousands of mutants[8,9]. However, Tn-Seq is not without drawbacks. 1) Not all intragenic Tn insertions lead to a loss of function. 2) Some properties cannot be assessed in a pooled assay format. 3) Specific mutants cannot be recovered from pools for further individual study. 4) Libraries of Tn mutant are never complete, no matter how large. In contrast, complete collections of mutants constructed by targeted mutagenesis do not suffer from these limitations, but their construction is laborious and expensive. For this reason, such collections are available for only a handful of bacterial species: the model Gram-negative *E. coli* K-12 (Gammaproteobacteria)[10], the model Gram-positive *Bacillus subtilis* (Firmicute)[11], *Acinetobacter baylyi* (Gammaproteobacteria) studied for its metabolic properties and possible biotechnological applications[12], and *Streptococcus sanguinis* (Firmicute) an opportunistic pathogen causing infective endocarditis[13]. The utility of such resources is best exemplified by a large-scale study using the Keio collection of approx. 4,000 mutants in *E. coli* K-12, which identified mutants presenting altered colony size upon growth in more than 300 different conditions[14]. More than 10,000 phenotypes were identified for approx. 2,000 mutants, including many in genes of unknown function[14]. Another major advantage of mutant collections constructed by targeted mutagenesis is that they also reveal complete lists of essential genes, those that cannot be mutated because their loss is lethal. These genes that encode proteins essential for cellular life, many of which are expected to be common to all living organisms[15], have attracted particular interest because they are expected to shed light on the origin of life. Additionally, essential genes specific to bacteria are also interesting because they are prime targets for the design of new drugs to tackle antimicrobial resistance.

Since the few species in which complete collections of mutants have been constructed represent only a tiny fraction of bacterial diversity, additional complete collections of mutants in diverse species are needed to tackle more effectively the

challenge of genes of unknown function. Human bacterial pathogens are particularly attractive candidates for this venture because of the potential practical implications for human health. Among these, *Neisseria meningitidis* – a Gram-negative Betaproteobacteria causing life-threatening meningitis and sepsis – stands out as an ideal candidate for several reasons. 1) It has a small genome with approx. 2,000 protein-coding genes, and yet displays a robust metabolism allowing rapid growth (doubling time of 40 min), even on minimal medium. 2) It has been the subject of intensive investigation for decades and there are several thousand publicly available meningococcal genome sequences. 3) It is naturally competent, which makes it a model for molecular genetics. Previously, we have used these advantages to design a modular toolbox for <u>Ne</u>isseria <u>me</u>ningitidis <u>sys</u>tematic analysis of gene function, named NeMeSys[16]. The central module of NeMeSys was an ordered library of approx. 4,500 Tn mutants in which the Tn insertion sites were sequence-defined[16,17], revealing insertions in more than 900 genes[16].

In the present study, we have 1) used the original library of Tn mutants as a starting point to generate a complete collection of defined mutants in *N. meningitidis* – named NeMeSys 2.0 – comprising one mutant in each non-essential protein-coding gene, 2) defined and analysed in depth the essential meningococcal genome, and 3) illustrated the potential of NeMeSys 2.0 for the phenotypic profiling of the meningococcal genome by identifying the function of multiple genes of unknown function.

Results and Discussion

Construction of the complete NeMeSys 2.0 collection of meningococcal mutants

The present study gave new impetus to a systematic re-annotation of the genome of *N. meningitidis* 8013 published in 2009[16]. We took into account new information in the public databases, new publications, an RNA-Seq analysis performed in 8013[18], and findings in the present study. In brief, we dropped one previously annotated gene, updated the gene product annotations of 521 genes, added 69 non-coding RNAs, and changed 176 gene start sites. The genome of *N. meningitidis* 8013 contains 2,060 protein-coding genes (labelled NMV_). In parallel, we re-sequenced the genome of the 8013 wild-type (WT) strain by Illumina whole genome sequencing (WGS), which confirmed the accuracy of the original genome sequence determined by Sanger sequencing more than 12 years ago[16], with an error rate of approx. 1 per 134,000 bases. We identified 17 differences with the published sequence, which are likely to be genuine since they were also found in the few mutants that were verified by WGS (see below). Of these differences, 11 correspond to single-nucleotide polymorphisms, five are single-nucleotide indels, and one is the deletion of a GT dinucleotide within a repeated GT tract in NMV_0677, which therefore corresponds to a phase variation event[19].

In keeping with previous studies describing the construction of complete collections of mutants[10-13], 85 genes (4.1 %) were not considered to be meaningful targets for mutagenesis (Fig. 1) because they encode transposases for repeated insertion sequences, or they correspond to short remnants of truncated genes and/or non-expressed cassettes (Supplementary Table 1). We therefore set out to assemble a complete collection of mutants in the remaining 1,975 genes (95.9 %) following a two-step approach described below. In brief, we first selected a subset of suitable Tn

mutants from the NeMeSys library, and then systematically mutagenised the

remaining target genes by allelic exchange (Supplementary Fig. 1).

First, since we previously constructed an arrayed library of Tn mutants in strain

8013 in which Tn insertion sites were sequence-defined[16,17], we selected a subset of

suitable mutants from this library. Specifically, mutations with a Tn insertion closer to

the centre of the genes (excluding the first and last 15 % that are often not gene-

inactivating) (Supplementary Fig. 1) were first re-transformed into strain 8013, which

is naturally competent. After extraction of the genomic DNA of the transformants, we

PCR-verified that they contained a Tn inserted in the expected target gene using

suitable pairs of flanking primers (Supplementary Table 2). We thus selected a

subset of 801 Tn mutants (40.6 % of target genes), harbouring Tn insertions

expected to be gene-inactivating (Fig. 1) (Supplementary Table 3). As a control,

Illumina WGS of one mutant, in *nlaB*, confirmed that it contained the expected

mutation, a Tn inserted after a TA dinucleotide at position 2,157,750 in the genome.

Next, we set out to systematically mutagenise the remaining 1,174 target

genes (59.4 %) by allelic exchange. For this, we used a previously validated no-

cloning mutagenesis method[20], based on splicing PCR (sPCR), to construct non-

polar mutants in which a central portion of the target gene (between the first and last

30 %) would be deleted and replaced by the same kanamycin (Km) resistance

cassette present in the above Tn (Supplementary Fig. 1). In brief, three PCR

products were first amplified, corresponding to the Km cassette and regions

upstream and downstream target genes. For failed reactions, we undertook as many

rounds of primer design as needed, until all PCRs were successful. The three PCR

products were then combined and spliced together. The final sPCR product was

transformed directly into *N. meningitidis* 8013. Typically, for non-essential genes, we

obtained hundreds of colonies resistant to kanamycin (Km^R). For each successful

transformation, two Km^R colonies were isolated and PCR-verified to contain the

expected mutation. To minimise false positive identification of essential genes, each

transformation that yielded no transformants was repeated at least three times with

different sPCR products. Only when all transformations failed to yield transformants,

was the target gene deemed to be essential. In summary (Fig. 1), out of the 1,174

genes that were targeted, we could disrupt 783 (Supplementary Table 3), while 391

could not be mutated and are thus essential for growth on rich medium

(Supplementary Table 4). As above, WGS of a few mutants constructed by sPCR, in

Int, *tatB* and *secB*, confirmed that each contained the expected mutation.

Taken together, using the above two-step approach, we constructed a

comprehensive set of 1,584 defined mutants in strain 8013, named the NeMeSys 2.0

collection. In addition, we also identified 391 candidate essential genes (19 % of all

protein-coding genes), which are required for growth of the meningococcus on rich

medium. For each gene that was successfully mutated, one mutant was stored at -

80ºC in glycerol, while the corresponding genomic DNA was stored at -20ºC. This

allows for easy distribution of mutants to the community and/or re-transformation of

the corresponding mutations in 8013, other meningococcal strains, or even

genetically closely related species like the gonococcus, which share most of their

genome with the meningococcus.

In depth analysis of the meningococcal essential genome

Considering their fundamental and practical importance, we first focused on the 391

genes essential for meningococcal life. Since corresponding proteins are expected to

be broadly conserved, we determined the partition of the 391 essential genes

between conserved and variable genomes in meningococci. To do this, we first used

the recently described PPanGGOLiN software[21] – an expectation-maximisation

algorithm based on multivariate Bernoulli mixture model coupled with a Markov

random field – to compute the pangenome of *N. meningitidis*, based on complete

genome sequences of 108 meningococcal isolates available in RefSeq

(Supplementary Table 5). We thereby classified genes in three categories[21],

persistent (gene families present in almost all genomes), shell (present at

intermediate frequencies), or cloud (present at low frequency). We then determined

how the 2,060 protein-coding genes of *N. meningitidis* 8013 partitioned between

these three classes (Supplementary Table 5). Only 1,664 genes (80.8 %) belong to

the persistent genome, while 396 genes (19.2 %) correspond to gene families

present at intermediate/low frequency in meningococci since they are part of the shell

(241 genes) and cloud genomes (155 genes) (Fig. 2a). Critically, besides highlighting

the well-known genomic plasticity of this naturally competent species, this analysis

confirmed our original prediction by revealing that essential genes are

overwhelmingly conserved in meningococcal isolates. Indeed, 382 essential genes

(97.4 %) are part of the persistent meningococcal genome (Fig. 2b) (Supplementary

Table 5). Of the remaining nine essential genes, six are part of the shell genome,

while three belong to the cloud genome (Supplementary Table 5).

Furthermore, many meningococcal essential genes (30.2 %) are conserved

and essential in other bacteria where systematic mutagenesis efforts have been

performed (Fig. 3a), including in phylogenetically distant species such as *S.*

sanguinis (Supplementary Table 6)[10,12,13]. When conservation was assessed with

Proteobacteria only (*E. coli* and *A. baylyi*), which are more closely related to *N.*

meningitidis, half of the 391 essential meningococcal genes (195 in total) are

conserved and essential in these three Gram-negative species (Fig. 3a).

Interestingly, a similar proportion of meningococcal essential genes (175 in total) was

conserved in JCVI-syn3.0 (Fig. 3b) (Supplementary Table 7), a synthetic

Mycoplasma mycoides bacterium engineered with a minimal genome[22] smaller than

that of any naturally occurring autonomously replicating cell. Taken together, these

observations are consistent with the notion that a sizeable portion of the genome

essential for cellular life is widely conserved in bacteria.

Next, to have a more global understanding of the biological functions that are

essential in the meningococcus, we analysed the distribution of the 391 essential

genes in functional categories (Supplementary Table 8). Genes were classified in specific functional categories using the bioinformatic tools embedded in the MicroScope platform[23], which hosts NeMeSys 2.0. In particular, we used predictions from MultiFun[24], MetaCyc[25], eggNOG[26], COG[27], FIGfam[28], and/or InterProScan[29]. As can be seen in Table 1, the essential meningococcal genes could be distributed in a surprisingly limited number of pathways (Table 1). In keeping with expectations, many essential genes are involved in key cellular processes such as 1) transcription (11 genes encoding subunits of the RNA polymerase, a series of factors modulating transcription, and two transcriptional regulators), 2) RNA modification/degradation (19 genes), and 3) protein biosynthesis (120 genes) (Table 1). The large class of genes involved in protein biosynthesis encode 1) all 24 proteins involved in tRNA charging, 2) virtually all ribosomal proteins (51/56), 3) five proteins involved in ribosome biogenesis/maturation, 4) 10 factors modulating translation, 5) two enzymes introducing post-translational modifications in proteins, 6) seven factors facilitating protein folding, and 7) 21 factors involved in protein export (signal peptidase, Lol system involved in lipoprotein export, general secretion Sec system, Tat translocation system, and TAM translocation and assembly module specific for autotransporters) (Table 1). Furthermore, many other essential genes encode proteins involved in additional key processes such as 1) genome replication and maintenance (20 proteins including subunits of the DNA polymerase, topoisomerases, gyrase, ligase *etc.*), 2) cell division (13 proteins), 3) peptidoglycan cell wall biogenesis (17 proteins), and 4) membrane biogenesis (37 proteins) (Table 1). The latter class includes all 12 proteins required for fatty acid biosynthesis, eight proteins required for the biosynthesis of phospholipids (phosphatidylethanolamine (PE) and phosphatidylglycerol (PG)), and 17 proteins involved in the biosynthesis and export of the lipo-oligosaccharide (LOS), which constitutes the external leaflet of the outer membrane. Finally, most of the remaining genes of known function (Table 1) are involved in energy generation, production of key metabolic intermediates (such

as dihydroxyacetone phosphate (DHAP), or 5-phosphoribosyl diphosphate (PRPP))

and a variety of metabolic pathways leading to the biosynthesis of 1) vitamins (B1,

B2, B6, and B9), 2) nucleotides, 3) amino acids (notably *meso*-diaminopimelate (M-

DAP) a component of the peptidoglycan), 4) isoprenoid, 5) heme, 6) CoA/ac-CoA, 7)

NAD/NADP, 8) iron-sulfur (Fe-S) clusters, 9) ubiquinol, 10) lipoate, and 11) S-

adenosyl-methionine (SAM). Many of these compounds are cofactors and/or

coenzymes known to be essential for the activity of many bacterial enzymes.

When essential genes were further integrated into networks and metabolic

pathways – using primarily MetaCyc[25] – the above picture became dramatically

simpler. As many as 91.3 % of the meningococcal essential genes partitioned into

just four major functional categories (Fig. 4). Namely, 1) gene/protein expression, 2)

genome/cell replication, 3) membrane/cell wall biogenesis, and 4) cytosolic

metabolism. Strikingly, for most multi-step enzymatic pathways, most, and often all,

the corresponding genes were identified as essential (Fig. 5), although they are most

often scattered throughout the genome. We view this as an important quality control

of our library. Only 34 essential genes (8.7 %) could not be clearly assigned to these

four categories. Critically, this functional partition is coherent with the one previously

determined for the minimal synthetic bacterium JCVI-syn3.0[22]. As a result, we were

able to generate a concise overview of the meningococcal essential genome in the

context of the cell (Fig. 5), in which many of the above pathways are detailed and

often linked by critical metabolic intermediates. This overview provides a useful

blueprint for systems and synthetic biology and a global understanding of

meningococcal biology.

Essential genes that are part of the variable genome are conditionally essential

A puzzling finding was that some essential genes, which could not be assigned to the

above four functional groups, are part of shell/cloud genomes present at

intermediate/low frequency in the meningococcus (Supplementary Table 5).

Interestingly, many of the corresponding genes are located in regions of genome

plasticity (RGP) – often thought to have been acquired by horizontal gene transfer –

as confirmed by the novel predictive method panRGP[30] that identifies RGP using

pangenome data generated by PPanGGOLiN[21]. A panRGP analysis identified 32

RGP in the genome of strain 8013, encompassing a total of 348,885 bp (15.3 % of

the genome) (Supplementary Table 9). Interestingly, one of these genes offered a

plausible explanation to the apparent paradox of essential genes that are not part of

the persistent meningococcal genome. NMV_2289 is predicted to encode a DNA-

methyltransferase, based on the presence of a DNA methylase N-4/N-6 domain

(InterPro IPR002941). Since the neighbouring NMV_2288 is predicted to encode a

restriction enzyme, the probable role of the NMV_2289 methyltransferase is to

protect meningococcal DNA against degradation by this restriction enzyme. This

would explain the lethal phenotype of the ΔNMV_2289 mutant. A closer examination

of the other essential genes that are part of shell/cloud genomes and often found in

RGP suggests that a similar scenario might be often applicable. We noticed that

NMV_1478, which encodes a protein of unknown function in RGP_2, is likely to be

co-transcribed with NMV_1479 that is predicted to encode the toxin of a toxin-

antitoxin (TA) system[31] (Fig. 6a). Therefore, since a toxin and its cognate antitoxin

are often encoded by closely linked genes, we hypothesized that NMV_1478 might

be the antitoxin for the neighbouring NMV_1479 toxin. This would explain why

NMV_1478 is essential, *i.e.* in its absence the NMV_1479 toxin would kill the cell. If

this is true, we reasoned that it should be possible to delete NMV_1478 together with

NMV_1479, which was attempted. As predicted, in contrast to NMV_1478 which

could not be deleted individually despite repeated attempts, a double deletion mutant

ΔNMV_1478/1479 could be readily obtained (Fig. 6a). This confirms that the

NMV_1478 gene is essential only when the toxin product of NMV_1479 is also

present. Such a scenario was also tested for NMV_0559, which is found in a large

RGP of 86.8 kbp (RGP_0), encompassing genes NMV_0527 to NMV_0618/0619

(Fig. 6b). We named this RGP *tps* because it contains multiple *tpsA* genes predicted

to encode large haemagglutinin/haemolysin-related proteins of two partner secretion

systems[32]. Since NMV_0559 is the only essential gene in the *tps* RGP, we tested

whether it might be conditionally essential by attempting to delete most of RGP_0. As

predicted, while NMV_0559 could not be deleted on its own despite repeated

attempts, an 80 kbp Δ*tps* deletion in RGP_0 encompassing NMV_0559 could be

readily obtained (Fig. 6b). This suggests that NMV_0559 is essential only in the

presence of another gene in RGP_0, which remains to be identified. We predict that

several other essential genes of unknown function that are not part of the persistent

meningococcal genome may be similarly conditionally essential, including 1)

NMV_1305 and NMV_1310 part of a putative prophage (RGP_1), 2) NMV_1541 the

putative antitoxin for neighbouring NMV_1539 toxin, 3) NMV_1544 putative antitoxin

for neighbouring NMV_1543 toxin, 4) NMV_1761 putative immunity protein against

neighbouring MafB toxins (RGP_4), which represent a newly identified family of

secreted toxins in pathogenic *Neisseria* species[33], 5) NMV_1918 part of a second *tps*

RGP (RGP_7), 6) NMV_2010 the putative immunity protein against neighbouring

bacteriocins (RGP_3), and 7) NMV_2333 and NMV_2335 putative immunity proteins

in a second *maf* RGP (RGP_11).

Filling holes in metabolic pathways: identification of a novel and widespread

histidinol-phosphatase

Another major utility of the NeMeSys 2.0 collection of mutants is to determine the

function of genes of unknown function by reverse genetics. We therefore wished to

illustrate this aspect. We noticed that the 10-step metabolic pathway leading to

histidine biosynthesis presents an apparent "hole" (Fig. 7a). The "missing" enzyme in

8013 corresponds to histidinol-phosphatase (EC 3.1.3.15), which catalyses the

dephosphorylation of histidinol-P into histidinol (Fig. 7a). The meningococcal genome

does not have homologs of known histidinol-phosphatases[34], a feature it shares with

many other bacterial species according to MetaCyc[25]. First, we excluded the

possibility that the meningococcus might be auxotrophic for histidine by showing that

8013 can grow on M9 minimal medium without added histidine (Fig. 7b). Next, we

sought to identify the unknown histidinol-phosphatase using NeMeSys 2.0, by

specifically testing mutants in genes encoding putative phosphatases of unknown

function, in search of a mutant that would not grow on M9 without added histidine.

Using this approach, we identified ΔNMV_1317 as a histidine auxotroph, growing on

M9 plates only when histidine was added (Fig. 7b). The product of this gene was

annotated as a putative hydrolase of unknown function, belonging to the IB subfamily

of the haloacid dehalogenase superfamily of aspartate-nucleophile hydrolases

(IPR006385). The corresponding sequences include a variety of phosphatases none

of which is annotated as a histidinol-phosphatase. Complementation of ΔNMV_1317

mutant with NMV_1317 restored growth on M9 without added histidine (Fig. 7b),

confirming that the auxotrophic phenotype was due to the mutation in NMV_1317. To

confirm that NMV_1317 encodes the missing enzyme in histidine biosynthesis, we

performed a cross-species complementation assay with the *hisB* gene from *E. coli*

DH5α (*hisB$_{EC}$*), which encodes a histidinol-phosphatase unrelated to NMV_1317.

Complementation of ΔNMV_1317 with *hisB$_{EC}$* restored growth on M9 without added

histidine (Fig. 7b). Complementation was gene specific, since *hisB$_{EC}$* could not

complement the growth deficiency of a different histidine auxotrophic mutant in

NMV_1778 (*hisH*) (Fig. 7b). Together, these results show that NMV_1317 defines a

previously unrecognised class of histidinol-phosphatases, allowing us to fill a hole in

the histidine biosynthesis pathway in the meningococcus. This finding, which

illustrates the utility of NeMeSys 2.0 for annotating genes of unknown function by

reverse genetics, has implications for other species lacking known histidinol-

phosphatases. Indeed, NMV_1317 homologs are widespread in Betaproteobacteria

and Gammaproteobacteria in which no histidinol-phosphatase has been identified

according to MetaCyc[25]. We expect that NMV_1317 homologs will encode the

elusive histidinol-phosphatase in a variety of Burkholderiales and Pseudomonadales,

including many pathogenic species of *Bordetella*, *Pseudomonas*, and *Burkholderia*

(there is some evidence for this in *Burkholderia phytofirmans*[35]).

Identification of a comprehensive set of genes involved in type IV pilus biology

in *N. meningitidis*, including three new ones

Another utility of NeMeSys 2.0 we wished to illustrate is the possibility to perform

whole-genome phenotypic screens to identify all the genes responsible for a

phenotype of interest. We chose to focus on type IV pili (T4P), which are pivotal

virulence factors in the meningococcus[36], for two reasons. First, T4P, are ubiquitous

in prokaryotes, which has made them a hot topic for research for the past 40 years[37].

Second, although many aspects of T4P biology remain incompletely understood,

most genes composing the multi-protein machinery involved in the assembly of these

filaments and/or their multiple functions have been identified, including in the

meningococcus that is one of the mainstream T4P models[36]. Hence, we could readily

benchmark the results of our screen against previous mutational analyses, especially

for the recovery rate of expected mutants. The two T4P-linked phenotypes we

decided to study are the formation of bacterial aggregates and twitching motility. Both

phenotypes can be simultaneously assessed – allowing thus a dual screen – by

observing the mutants growing in liquid medium by phase-contrast microscopy[38]. We

thus scored the 1,589 mutants for the presence of round aggregates, and for the

continuous and vigorous jerky movements of cells within these aggregates, which

corresponds to twitching motility[38] (Supplementary Table 10). This analysis revealed

that 20 mutants (1.2 %) present phenotypic defects in these two T4P-linked

phenotypes (Table 2). Significantly, we identified 100 % of the expected single

mutants in 17 *pil* genes known to affect these phenotypes in the meningococcus[38-40].

This is another quality control of the NeMeSys 2.0 collection of mutants, which

confirms excellent correlation between phenotype and genotype, and shows that

when a robust screening method is used all the genes involved in a given phenotype

can be identified. Furthermore, we could readily identify mutants such as Δ*pilT* able

to form (irregular) aggregates, but showing no twitching motility[38]. PilT is known to

encode the motor powering pilus retraction, which is directly responsible for twitching

motility[41]. No other mutant in the library exhibited a similar phenotype. Strikingly, we

also identified mutants in three genes not previously associated with T4P biology in

N. meningitidis (Table 2), which was unexpected considering that T4P have been

studied for decades in this species. TsaP has been shown in the closely related

pathogen *Neisseria gonorrhoeae* to interact with the secretin PilQ[42], which forms a

pore in the outer membrane through which T4P translocate onto the cell surface.

TsaP plays a poorly understood role in T4P biology in *N. gonorrhoeae* and

Myxococcus xanthus[42], but has apparently no role in *Pseudomonas aeruginosa*[43]. In

addition, we identified NMV_1205 and NMV_2228, which to the best of our

knowledge have never been previously linked to T4P biology. NMV_1205 is

predicted to encode a periplasmic protein of unknown function and is found mainly in

Neisseriales, where it is widespread. NMV_2228 is predicted to encode a carbonic

anhydrase, an enzyme catalysing the reversible hydration of carbon dioxide[44].

Next, we analysed the three new mutants in detail using an approach

previously validated in the meningococcus with many *pil* genes[38,45]. We first tested

whether piliation was affected in the mutants by purifying T4P using a procedure in

which filaments sheared by vortexing are precipitated using ammonium sulfate[38].

Pilus preparations, obtained from equivalent numbers of cells, were separated by

SDS-PAGE and either stained using Coomassie blue, or tested by immunoblotting

using an anti-PilE antibody. This revealed the major pilin PilE as a 17 kDa species

(Fig. 8a) and confirmed that all three mutants were piliated. However, when

compared to the WT, pilus yields differed between the mutants. While piliation in

ΔNMV_2228 was apparently normal, it was decreased in Δ*tsaP* and ΔNMV_1205.

The decrease was dramatic in ΔNMV_1205, where filaments could be detected only

by immunoblotting (Fig. 8a). Quantification of piliation using a whole-cell ELISA

procedure[38] confirmed these findings (Fig. 8b) and showed that piliation levels were

40 %, 31 % and 154 % of WT, in ΔtsaP, ΔNMV_1205 and ΔNMV_2228, respectively.

Therefore, none of the corresponding proteins is required for T4P biogenesis.

We then further characterised the new mutants for T4P-linked functions. Since

it is known that in some meningococcal mutants piliation and/or aggregation defects

can be restored when filament retraction is abolished by a concurrent mutation in

pilT[38,45], we tested if that might be the case for the new mutants. First, we showed

that aggregation, which is abolished in ΔtsaP and ΔNMV_1205 and dramatically

affected in ΔNMV_2228, could be restored when mutants were complemented with

the corresponding WT alleles (Fig. 9a). This confirmed that the phenotypic defects in

these mutants were indeed due to the mutations in the above three genes. Similarly,

some aggregation was restored in ΔtsaP/ΔpilT and ΔNMV_1205/ΔpilT

(ΔNMV_2228/ΔpilT could not be constructed), which harboured a concurrent

mutation in pilT that encodes the retraction motor PilT. However, aggregates were

morphologically distinct from the round aggregates formed by the WT or the irregular

aggregates formed by ΔpilT. These findings suggest that tsaP and NMV_1205

participate in the formation of aggregates by counterbalancing PilT-mediated pilus

retraction. Lastly, we checked whether the mutants were affected for another T4P-

linked phenotype, competence for DNA uptake which makes the meningococcus

naturally transformable. We quantified competence in the three mutants as

previously described[46] by transformation to rifampicin resistance (Fig. 9b). We found

that ΔtsaP was almost as transformable as the WT, ΔNMV_1205 showed a 26-fold

decrease in transformation, while competence was completely abolished in

ΔNMV_2228 (explaining why the ΔNMV_2228/ΔpilT double mutant could not be

constructed). These findings show that tsaP, NMV_1205 and NMV_2228 contribute

to the fine-tuning of multiple functions mediated by T4P in N. meningitidis.

Taken together, these findings are of significance in two ways. The

identification of three new genes playing a role in T4P biology (how exactly remains

to be understood), will contribute to a better understanding of these filaments. This

has broad implications because T4P and T4P-related filamentous nanomachines are

ubiquitous in Bacteria and Archaea[47]. In addition, the identification of new genes

contributing to phenotypes that have been extensively studied in several species for

the past 30 years is unambiguous evidence of the potential of NeMeSys 2.0 to lead

to a global phenotypic profiling of the meningococcal genome.

Concluding remarks

Here, we described the construction of a complete collection of defined mutants in *N.*

meningitidis, which has been fully integrated in our modular NeMeSys toolbox[16],

accessible online in MicroScope[23]. Furthermore, we illustrated NeMeSys 2.0 utility for

tackling the challenge of genes of unknown function, following a variety of

approaches. Although they were limited at this stage to a few different biological

properties, these experiments have provided new information on meningococcal

biology with possible implications for many other species. They have also clearly

demonstrated the potential of NeMeSys 2.0 to make a significant contribution to

ongoing efforts directed towards a comprehensive understanding of a bacterial cell.

Such potential is further amplified by several useful properties of *N. meningitidis* such

as 1) its small genome with limited functional redundancy, which in species with

larger genomes can obscure the link between phenotype and genotype, 2) its hardy

nature with a robust metabolism allowing it to grow on minimal medium, 3) its

taxonomy making it the first Betaproteobacteria in which a complete library of

mutants is available, and 4) the fact that it is a major human pathogen, which allows

addressing virulence properties absent in non-pathogenic model species. This latter

point makes the meningococcus a prime model for the identification of new means

for controlling bacterial pathogens, which would have practical implications for human

health.

The other major achievement in the present study is the identification of the

essential meningococcal genome, comprising 391 genes that could not be disrupted.

The finding that more than 90 % of these genes are associated with just four basic

biological functions, helped us generate a coherent concise cellular overview of the

meningococcal essential metabolism. We surmise that most, if not all, of the

remaining essential genes of unknown function, which are conserved in other

bacterial species, will be involved in either expression of genome information,

preservation of genome information, cell membrane structure/function, or cytosolic

metabolism (in particular the production of key metabolic intermediates, vitamins and

crucial cofactors/coenzymes). Our cellular overview provides a useful starting point

for systems biology, which together with previous genome-scale metabolic network

models[48,49], could help us progress towards a global understanding of meningococcal

biology. This, in turn, would have widespread implications, because many of the

meningococcal genes, especially the essential ones, are widely conserved.

Materials and methods

Strains and growth conditions

All the *N. meningitidis* strains that were generated and used in this study were derived from a highly adhesive variant – sometimes called clone 12 or 2C43 – of the clinical isolate 8013[50]. This serogroup C strain, which belongs to the sequence type 177 and clonal complex ST-18, has been previously sequenced[16]. Meningococci were routinely grown on plates with GC agar Base (GCB) (Difco) containing 5 g/l agar (all chemicals were from Sigma unless stated otherwise) and Kellogg's supplements (4 g/l glucose, 0.59 µM thiamine hydrochloride, 12.37 µM $Fe(NO_3)_3 \cdot 9H_2O$, 68.4 µM L-glutamine). Plates were incubated overnight (O/N) at 37°C in a moist atmosphere containing 5 % CO_2. Alternatively, we used agar plates with M9 minimal medium (4 g/l glucose, 4.78 mM Na_2HPO_4, 2.2 mM KH_2PO_4, 1.87 mM NH_4Cl, 8.56 mM NaCl, 2 mM $MgSO_4$, 0.1 mM $CaCl_2$). When required, plates contained 100 µg/ml kanamycin, 3 µg/ml erythromycin, 5 µg/ml rifampicin, 25 µg/ml L-histidine, and/or 0.4 mM isopropyl-β-D-thiogalactopyranoside (IPTG) (Merck). Strains were stored at -80ºC in 10 % glycerol in liquid GC (15 g/l protease peptone No. 3, 23 mM K_2HPO_4, 7.34 mM KH_2PO_4, 85.6 mM NaCl). *E. coli* TOP10 and DH5α were grown at 37°C in liquid or solid lysogeny broth (LB), which contained 100 µg/ml spectinomycin or 50 µg/ml kanamycin, when appropriate.

Construction of strains

Genomic DNA from *N. meningitidis* and *E. coli* strains was prepared using the Wizard Genomic DNA Purification kit (Promega) following the manufacturer's instructions. Plasmid DNA from *E. coli* strains was purified using the QIAprep Spin Miniprep Kit (Qiagen) following the manufacturer's instructions. Bacteria were transformed as follows. *N. meningitidis* is naturally competent for transformation. A loopful of bacteria grown on GCB plates was resuspended in liquid GC containing 5 mM $MgCl_2$ (GC

transfo), 200 μl was aliquoted in the well of a 24-well plate, and DNA was added.

After incubation for 30 min at 37°C on an orbital shaker, 0.8 ml GC transfo was

added to the wells and the plates were further incubated for 3 h at 37°C, without

shaking. Transformants were selected on GCB plates containing suitable antibiotics.

For transformation of *E. coli*, ultra-competent cells were prepared as described

elsewhere[51] and transformed by a standard heat shock procedure[52]. Transformants

were selected on LB plates containing the suitable antibiotic.

The construction of the NeMeSys 2.0 library of meningococcal mutants

followed a two-step procedure explained in the Results section (Supplementary Fig.

1). First, we selected a subset of potentially suitable mutants from an archived library

of Tn mutants with sequence-defined Tn insertion sites, which was described

previously[16,17]. The corresponding genomic DNAs were PCR-verified using primers

flanking the Tn insertion sites (Supplementary Table 2). When the mutants were

confirmed to be correct, the corresponding mutations were re-transformed in 8013,

the mutants were PCR-verified once again and stored at -80ºC. The remaining target

genes, for which no Tn mutants were available, were then systematically submitted

to non-polar targeted mutagenesis using a validated no-cloning method[20]. In brief,

using high-fidelity PfuUltra II Fusion HS DNA Polymerase (Agilent) and two sets of

specific primers (F1/R1 and F2/R2), we amplified 500-750 bp PCR products

upstream and downstream from each target gene, respectively. The R1 and F2

primers were consecutive, non-overlapping and chosen within the target gene

(excluding the first and last 30 %) (Supplementary Fig. 1). These primers contained

20-mer overhangs complementary to the F3/R3 primers used to amplify the 1,518 bp

kanamycin resistance cassette present in the Tn (Supplementary Table 2), which

contains a DNA uptake sequence necessary for efficient transformation in the

meningococcus[53] (Supplementary Fig. 1). In the first step, three PCR products were

amplified separately using F1/R1, F2/R2 and F3/R3 pairs of primers. The first two

PCRs contained a 4-fold excess of outer primers F1 and R2[54]. The three products

were then combined (3.3 μl of each) and spliced together by PCR using the excess

F1/R2 primers added in the first reaction[54]. The sPCR products were directly

transformed into *N. meningitidis*. For each successful transformation, two colonies

were isolated and verified by PCR using F1/R2. When transformations yielded no

transformants, they were repeated at least three times with different sPCR products.

This method was also used to construct several polymutants in which we deleted the

80 kbp RGP_0, the TA system NMV_1478/1479, and three repeated prophages

(NMV_1286/1294, NMV_1387/1398, and NMV_1412/1418). For each gene that was

successfully mutated, one mutant was individually stored at -80ºC in glycerol, and the

tubes were ordered according to the gene NMV_ label. Likewise, corresponding

genomic DNAs that allow easy re-transformation of these mutations (even in other

Neisseria strains), were individually ordered in Eppendorf tubes and stored at -20ºC.

Several mutants, as well as the WT strain, were verified by WGS. Sequencing was

performed by MicrobesNG on an Illumina sequencer using standard Nextera

protocols. An average of 122 Mb of read sequences (between 80 and 179 Mb) were

obtained for each sequencing project, representing an average 53-fold genome

coverage. The corresponding Illumina reads have been deposited in the European

Nucleotide Archive.

The Δ*tsaP*/Δ*pilT* and ΔNMV_1205/Δ*pilT* double mutants were constructed by

transforming Δ*tsaP* and ΔNMV_1205 with genomic DNA extracted from a *pilT* mutant

disrupted by a cloned *ermAM* cassette conferring resistance to erythromycin[55].

ΔNMV_2228/Δ*pilT* could not be constructed because both mutations abolish

competence. For complementation assays, complementing genes were amplified

using specific indF/indR primers (Supplementary Table 2), with overhangs

introducing flanking *Pac*I sites for cloning, and a ribosome binding site in front of the

gene. The corresponding PCR products were cloned into pCR8/GW/TOPO

(Invitrogen), verified by Sanger sequencing and subcloned into pGCC4[56]. This placed

the genes under the transcriptional control of an IPTG-inducible promoter, within a

DNA an intragenic region of the gonococcal chromosome conserved in *N.*

meningitidis[56]. The resulting plasmids were transformed in the desired

meningococcal mutant, leading to ectopical insertion of the complementing gene. We

thus constructed ΔNMV_1317::NMV_1317, ΔNMV_1317::*hisB$_{EC}$*,

ΔNMV_1718::*hisB$_{EC}$*, Δ*tsaP*::*tsaP*, ΔNMV_1205::NMV_1205 and

ΔNMV_2228::NMV_2228. Since ΔNMV_2228 is not competent, the last strain was

constructed by first transforming the corresponding pGCC4 derivative into the WT

strain, before transforming the ΔNMV_2228 mutation. Expression of the

complementing genes in *N. meningitidis* was induced by growing the strains on GCB

plates containing 0.5 mM IPTG.

Dual screen for mutants affected for aggregation and/or twitching motility

N. meningitidis mutants grown O/N on GCB plates were resuspended individually

with a 200 µl pipette tip in the wells of 24-well plates containing 500 µl pre-warmed

RPMI 1640 with L-glutamine (PAA Laboratories), supplemented with 10 % heat-

inactivated fetal bovine serum Gold (PAA Laboratories). Plates were incubated for

approx. 2 h at 37ºC. Aggregates forming on the bottom of the wells were visualised

by phase-contrast microscopy using a Nikon TS100F microscope[38]. Digital images of

aggregates were recorded using a Sony HDR-CX11 camcorder mounted onto the

microscope. In parallel, we scored twitching motility by observing whether bacteria in

aggregates exhibited continuous and vigorous jerky movement[38]. This movement is

abolished in a Δ*pilT* mutant, in which the T4P retraction motor that powers twitching

motility is not produced anymore.

Detection and quantification of T4P

Pilus purification by ammonium sulfate precipitation was carried out as follows.

Bacteria grown O/N on GCB plates were first resuspended in 1.5 ml ethanolamine

buffer (150 mM ethanolamine, 65 mM NaCl) at pH 10.5. Filaments were sheared by

vortexing at maximum speed for 1 min, before OD_{600} was adjusted to 9-12 using

ethanolamine buffer (in 1.5 ml final volume). Bacteria were then pelleted by

centrifugation at 17,000 g at 4°C during 10 min. The supernatant (1.35 ml) was

recovered, topped to 1.5 ml with ethanolamine buffer. This step was repeated once,

and filaments were precipitated for approx. 1 h at room temperature (RT) by adding

150 µl ethanolamine buffer saturated with ammonium sulphate. Filaments were then

pelleted by centrifugation at 17,000 g at 4°C during 15 min. Pellets were rinsed once

with 100 µl Tris-buffered saline at pH 8, and finally resuspended in 100 µl of Laemmli

buffer (BioRad) containing β-mercaptoethanol.

Piliation was quantified by performing whole-cell ELISA[38] using the 20D9

mouse monoclonal antibody that is specific for the T4P in strain 8013[57]. Bacteria

grown O/N on GCB plates were resuspended in PBS, adjusted to OD_{600} 0.1 and

heat-killed during 1 h at 56ºC. Serial 2-fold dilutions were then aliquoted (100 µl) in

the wells of 96-well plates. Each well was made in triplicate. The plates were dried

O/N at RT in a running safety cabinet. The next day, wells were washed seven times

with washing solution (0.1 % Tween 80 in PBS), before adding 100 µl/well of 20D9

antibody (diluted 1/1,000 in washing solution containing 5 % skimmed milk). Plates

were incubated 1 h at RT, and then washed seven times with washing solution. Next,

we added 100 µl/well of Amersham ECL-HRP linked mouse IgG antibody (GE

Healthcare) diluted 1/10,000. Plates were incubated 1 h at RT, and then washed

seven times with washing solution. We then added 100 µl/well of TMB solution

(Thermo Scientific) and incubated the plates during 20 min at RT in the dark. Finally,

we stopped the reaction by adding 100 µl/well of 0.18 M sulfuric acid, before reading

the plates at 450 nm using a plate reader. Statistical analyses were performed with

Prism (GraphPad Software) using appropriate tests as described in the legends to

figures.

SDS-PAGE, Coomassie staining and immunoblotting

Purified T4P were separated by SDS-PAGE using 15 % polyacrylamide gels. Gels

were stained using Bio-Safe Coomassie stain (BioRad) or blotted to Amersham

Hybond ECL membranes (GE Healthcare) using standard molecular biology

techniques[52]. Blocking, incubation with primary/secondary antibodies and detection

using Amersham ECL Plus reagents (GE Healthcare) were done following the

manufacturer's instructions. The primary antibody was a previously described rabbit

anti-PilE serum (1/2,500)[58], while the secondary antibody (1/10,000) was a

commercial Amersham ECL-HRP linked rabbit IgG (GE Healthcare).

Quantifying transformation in the meningococcus

Since the tested mutants are Km^R, we tested their competence by transforming them

to Rif^R using DNA from a mutant of 8013 spontaneously resistant to rifampicin[59]. We

first amplified by PCR, using rpoF/rpoR primers, a 1,172 bp internal portion of *rpoB*,

which usually contains point mutations[60] leading to Rif^R. This PCR product was

cloned in pCR8/GW/TOPO and sequenced, which revealed a single point mutation,

leading to a His $\rightarrow$ Tyr substitution at position 553 in RpoB. This pCR8/GW/TOPO

derivative was used to amplify the PCR product used for quantifying competence,

which was done as follows. Bacteria grown O/N on GCB plates were resuspended in

pre-warmed liquid GC transfo at an OD_{600} of 0.1. The number of bacterial cells in this

suspension was quantified by performing colony-forming unit (CFU) counts. We

mixed 200 µl of bacterial suspension with 100 ng of transforming DNA in the wells of

a 24-well plate. After incubating for 30 min at 37ºC on an orbital shaker, 0.8 ml GC

transfo was added to the wells and the plates were further incubated during 3 h at

37ºC without shaking. Transformants were selected by plating appropriate dilutions

on plates containing rifampicin and by counting, the next day, the number of Rif^R

CFU. Transformation frequencies are expressed as % of transformed recipient cells.

Bioinformatics

The Qiagen CLC Genomics Workbench software was used for WGS analysis. In

brief, Illumina reads from each strain were mapped onto the published sequence of

8013[16], and good quality and high frequency (>70 %) base changes and small

deletions/insertions were identified. Large deletions/insertions were identified using

annotated assemblies (N50 between 38,959 and 45,097 bp), which were visualised

in Artemis[61], by performing pairwise sequence alignments using DNA Strider[62].

The genome of strain 8013 and a total of 108 complete genomes from NCBI

RefSeq (last accessed June 8[th], 2020) (Supplementary Table 5) were used to

compute the *N. meningitidis* pangenome using the PPanGGOLiN software[21] (version

1.1.85). The original annotations of the genomes have been kept in order to compute

gene families. In brief, PPanGGOLiN uses a statistical model to infer the pangenome

classes (persistent, shell and cloud) based on both the presence/absence of gene

families and genomic neighbourhood information[21]. The options "--use pseudo" and

"--defrag" were used to consider pseudogenes in the gene family computation, and to

associate fragmented genes with their original gene family, respectively. The other

parameters have been used with default values to compute the persistent, shell and

cloud partitions. The PPanGGOLiN results were then used to predict regions of

genomic plasticity using the panRGP method with default options[30].

Datasets and annotations generated during this study have been stored within

MicroScope[23]. This publicly accessible web interface can be used to visualise

genomes (simultaneously with synteny maps in other microbial genomes), perform

queries (by BLAST or keyword searches) and download datasets/annotations in a

variety of formats (including EMBL and GenBank).

Data availability

Whole genome sequencing data that support the findings in this study have been

deposited in the European Nucleotide Archive under accession number

PRJEB39197. All the datasets generated during this study are either included in this

published article (and its Supplementary Information files), or available in MicroScope

(http://mage.genoscope.cns.fr/microscope/mage/viewer.php?O_id=99).

References

1. Jolley, K.A. & Maiden, M.C. BIGSdb: Scalable analysis of bacterial genome
variation at the population level. *BMC Bioinfo.* **11**, 595 (2010).

2. Ghatak, S., King, Z.A., Sastry, A. & Palsson, B.O. The y-ome defines the 35% of
Escherichia coli genes that lack experimental evidence of function. *Nucleic Acids*
Res. **47**, 2446-2454 (2019).

3. Galperin, M.Y. & Koonin, E.V. From complete genome sequence to 'complete'
understanding? *Trends Biotechnol.* **28**, 398-406 (2010).

4. Akerley, B.J. *et al.* A genome-scale analysis for identification of genes required for
growth or survival of *Haemophilus influenzae*. *Proc. Natl. Acad. Sci. USA* **99**, 966-
971 (2002).

5. Liberati, N.T. *et al.* An ordered, nonredundant library of *Pseudomonas aeruginosa*
strain PA14 transposon insertion mutants. *Proc. Natl. Acad. Sci. USA* **103**, 2833-
2838 (2006).

6. Deutschbauer, A. *et al.* Evidence-based annotation of gene function in *Shewanella*
oneidensis MR-1 using genome-wide fitness profiling across 121 conditions. *PLoS*
Genet. **7**, e1002385 (2011).

7. van Opijnen, T., Lazinski, D.W. & Camilli, A. Genome-wide fitness and genetic
interactions determined by Tn-seq, a high-throughput massively parallel
sequencing method for microorganisms. *Curr. Protoc. Mol. Biol.* **106**, 7.16.1-
7.16.24 (2014).

8. Langridge, G.C. *et al.* Simultaneous assay of every *Salmonella* Typhi gene using
one million transposon mutants. *Genome Res.* **19**, 2308-2316 (2009).

9. Christen, B. *et al.* The essential genome of a bacterium. *Mol. Syst. Biol.* **7**, 528
(2011).

10. Baba, T. *et al.* Construction of *Escherichia coli* K-12 in-frame, single-gene
knockout mutants: the Keio collection. *Mol. Syst. Biol.* **2**, 2006.0008 (2006).

11. Kobayashi, K. *et al.* Essential *Bacillus subtilis* genes. *Proc. Natl. Acad. Sci. USA*
100, 4678-4683 (2003).

12. de Berardinis, V. *et al.* A complete collection of single-gene deletion mutants of
Acinetobacter baylyi ADP1. *Mol. Syst. Biol.* **4**, 174 (2008).

13. Xu, P. *et al.* Genome-wide essential gene identification in *Streptococcus*
sanguinis. *Sci. Rep.* **1**, 125 (2011).

14. Nichols, R.J. *et al.* Phenotypic landscape of a bacterial cell. *Cell* **144**, 143-156
(2011).

15. Mushegian, A.R. & Koonin, E.V. A minimal gene set for cellular life derived by
comparison of complete bacterial genomes. *Proc. Natl. Acad. Sci. USA* **93**, 10268-
10273 (1996).

16. Rusniok, C. *et al.* NeMeSys: a resource for narrowing the gap between sequence
and function in the human pathogen *Neisseria meningitidis*. *Genome Biol.* **10**,
R110 (2009).

17. Geoffroy, M., Floquet, S., Métais, A., Nassif, X. & Pelicic, V. Large-scale analysis
of the meningococcus genome by gene disruption: resistance to complement-
mediated lysis. *Genome Res.* **13**, 391-398 (2003).

18. Heidrich, N. *et al.* The primary transcriptome of *Neisseria meningitidis* and its
interaction with the RNA chaperone Hfq. *Nucleic Acids Res.* **45**, 6147-6167
(2017).

19. Martin, P. *et al.* Experimentally revised repertoire of putative contingency loci in
Neisseria meningitidis strain MC58: evidence for a novel mechanism of phase
variation. *Mol. Microbiol.* **50**, 245-257 (2003).

20. Georgiadou, M., Castagnini, M., Karimova, G., Ladant, D. & Pelicic, V. Large-
scale study of the interactions between proteins involved in type IV pilus biology in
Neisseria meningitidis: characterization of a subcomplex involved in pilus
assembly. *Mol. Microbiol.* **84**, 857-873 (2012).

21. Gautreau, G. *et al.* PPanGGOLiN: Depicting microbial diversity via a partitioned

pangenome graph. *PLoS Comput. Biol.* **16**, e1007732 (2020).

22. Hutchison, C.A. *et al.* Design and synthesis of a minimal bacterial genome.

Science **351**, aad6253 (2016).

23. Vallenet, D. *et al.* MicroScope: an integrated platform for the annotation and

exploration of microbial gene functions through genomic, pangenomic and

metabolic comparative analysis. *Nucleic Acids Resesarch* **48**, D579-D589 (2020).

24. Serres, M.H. & Riley, M. MultiFun, a multifunctional classification scheme for

Escherichia coli K-12 gene products. *Microb. Comp. Genomics* **5**, 205-222 (2000).

25. Caspi, R. *et al.* The MetaCyc database of metabolic pathways and enzymes - a

2019 update. *Nucleic Acids Res.* **48**, D445-D453 (2020).

26. Huerta-Cepas, J. *et al.* eggNOG 5.0: a hierarchical, functionally and

phylogenetically annotated orthology resource based on 5090 organisms and

2502 viruses. *Nucleic Acids Res.* **47**, D309-D314 (2019).

27. Galperin, M.Y., Makarova, K.S., Wolf, Y.I. & Koonin, E.V. Expanded microbial

genome coverage and improved protein family annotation in the COG database.

Nucleic Acids Res. **43**, D261-D269 (2015).

28. Meyer, F., Overbeek, R. & Rodriguez, A. FIGfams: yet another set of protein

families. *Nucleic Acids Res.* **37**, 6643-6654 (2009).

29. Jones, P. *et al.* InterProScan 5: genome-scale protein function classification.

Bioinformatics **30**, 1236-1240 (2014).

30. Bazin, A., Gautreau, G., Médigue, C., Vallenet, D. & Calteau, A. panRGP: a

pangenome-based method to predict genomic islands and explore their diversity.

Preprint at http://www.biorxiv.org/content/10.1101/2020.03.26.007484v1 (2020).

31. Yamaguchi, Y., Park, J.H. & Inouye, M. Toxin-antitoxin systems in bacteria and

archaea. *Annu. Rev. Genet.* **45**, 61-79 (2011).

32. van Ulsen, P., Rutten, L., Feller, M., Tommassen, J. & van der Ende, A. Two-

partner secretion systems of *Neisseria meningitidis* associated with invasive clonal

complexes. *Infect. Immun.* **76**, 4649-4658 (2008).

33. Jamet, A. *et al.* A new family of secreted toxins in pathogenic *Neisseria* species.

PLoS Pathog. **11**, e1004592 (2015).

34. Kulis-Horn, R.K., Ruckert, C., Kalinowski, J. & Persicke, M. Sequence-based

identification of inositol monophosphatase-like histidinol-phosphate phosphatases

(HisN) in *Corynebacterium glutamicum*, Actinobacteria, and beyond. *BMC*

Microbiol. **17**, 161 (2017).

35. Price, M.N. *et al.* Mutant phenotypes for thousands of bacterial genes of unknown

function. *Nature* **557**, 503-509(2018).

36. Pelicic, V. Type IV pili: *e pluribus unum*? *Mol. Microbiol.* **68**, 827-837 (2008).

37. Berry, J.L. & Pelicic, V. Exceptionally widespread nano-machines composed of

type IV pilins: the prokaryotic Swiss Army knives. *FEMS Microbiol. Rev.* **39**, 134-

154 (2015).

38. Brown, D., Helaine, S., Carbonnelle, E. & Pelicic, V. Systematic functional

analysis reveals that a set of 7 genes is involved in fine tuning of the multiple

functions mediated by type IV pili in *Neisseria meningitidis*. *Infect. Immun.* **78**,

3053-3063 (2010).

39. Carbonnelle, E., Helaine, S., Prouvensier, L., Nassif, X. & Pelicic, V. Type IV

pilus biogenesis in *Neisseria meningitidis*: PilW is involved in a step occuring after

pilus assembly, essential for fiber stability and function. *Mol. Microbiol.* **55**, 54-64

(2005).

40. Helaine, S. *et al.* PilX, a pilus-associated protein essential for bacterial

aggregation, is a key to pilus-facilitated attachment of *Neisseria meningitidis* to

human cells. *Mol. Microbiol.* **55**, 65-77 (2005).

41. Merz, A.J., So, M. & Sheetz, M.P. Pilus retraction powers bacterial twitching

motility. *Nature* **407**, 98-102 (2000).

42. Siewering, K. *et al.* Peptidoglycan-binding protein TsaP functions in surface

assembly of type IV pili. *Proc. Natl. Acad. Sci. USA* **111**, E953-E961 (2014).

43. Koo, J., Lamers, R.P., Rubinstein, J.L., Burrows, L.L. & Howell, P.L. Structure of

the *Pseudomonas aeruginosa* type IVa pilus secretin at 7.4 Å. *Structure* **24**, 1778-

1787 (2016).

44. Smith, K.S. & Ferry, J.G. Prokaryotic carbonic anhydrases. *FEMS Microbiol. Rev.*

24, 335-366 (2000).

45. Carbonnelle, E., Helaine, S., Nassif, X. & Pelicic, V. A systematic genetic

analysis in *Neisseria meningitidis* defines the Pil proteins required for assembly,

functionality, stabilization and export of type IV pili. *Mol. Microbiol.* **61**, 1510-1522

(2006).

46. Berry, J.L., Cehovin, A., McDowell, M.A., Lea, S.M. & Pelicic, V. Functional

analysis of the interdependence between DNA uptake sequence and its cognate

ComP receptor during natural transformation in *Neisseria* species. *PLoS Genet.* **9**,

e1004014 (2013).

47. Denise, R., Abby, S.S. & Rocha, E.P.C. Diversification of the type IV filament

superfamily into machines for adhesion, protein secretion, DNA uptake, and

motility. *PLoS Biol.* **17**, e3000390 (2019).

48. Baart, G.J. *et al.* Modeling *Neisseria meningitidis* metabolism: from genome to

metabolic fluxes. *Genome Biol.* **8**, R136 (2007).

49. Mendum, T.A., Newcombe, J., Mannan, A.A., Kierzek, A.M. & McFadden, J.

Interrogation of global mutagenesis data with a genome scale model of *Neisseria*

meningitidis to assess gene fitness *in vitro* and in sera. *Genome Biol.* **12**, R127

(2011).

50. Nassif, X. *et al.* Antigenic variation of pilin regulates adhesion of *Neisseria*

meningitidis to human epithelial cells. *Mol. Microbiol.* **8**, 719-725 (1993).

51. Inoue, H., Nojima, H. & Okayama, H. High efficiency transformation of

Escherichia coli with plasmids. *Gene* **96**, 23-28 (1990).

52. Sambrook, J. & Russell, D.W. *Molecular cloning. A laboratory manual*, (Cold

Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 2001).

53. Pelicic, V., Morelle, S., Lampe, D. & Nassif, X. Mutagenesis of *Neisseria*

meningitidis by *in vitro* transposition of *Himar1 mariner*. *J. Bacteriol.* **182**, 5391-

5398 (2000).

54. Metzgar, D. *et al. Acinetobacter* sp. ADP1: an ideal model organism for genetic

analysis and genome engineering. *Nucleic Acids Res.* **32**, 5780-5790 (2004).

55. Pujol, C., Eugène, E., Marceau, M. & Nassif, X. The meningococcal PilT protein

is required for induction of intimate attachment to epithelial cells following pilus-

mediated adhesion. *Proc. Natl. Acad. Sci. USA* **96**, 4017-4022 (1999).

56. Mehr, I.J., Long, C.D., Serkin, C.D. & Seifert, H.S. A homologue of the

recombination-dependent growth gene, *rdgC*, is involved in gonococcal pilin

antigenic variation. *Genetics* **154**, 523-532 (2000).

57. Pujol, C., Eugène, E., de Saint Martin, L. & Nassif, X. Interaction of *Neisseria*

meningitidis with a polarized monolayer of epithelial cells. *Infect. Immun.* **65**, 4836-

4842 (1997).

58. Morand, P.C. *et al.* Type IV pilus retraction in pathogenic *Neisseria* is regulated

by the PilC proteins. *EMBO J.* **23**, 2009-2017 (2004).

59. Cehovin, A. *et al.* Specific DNA recognition mediated by a type IV pilin. *Proc.*

Natl. Acad. Sci. USA **110**, 3065-3070 (2013).

60. Carter, P.E., Abadi, F.J., Yakubu, D.E. & Pennington, T.H. Molecular

characterization of rifampin-resistant *Neisseria meningitidis*. *Antimicrob. Agents*

Chemother. **38**, 1256-1261 (1994).

61. Carver, T., Harris, S.R., Berriman, M., Parkhill, J. & McQuillan, J.A. Artemis: an

integrated platform for visualization and analysis of high-throughput sequence-

based experimental data. *Bioinformatics* **28**, 464-469 (2012).

62. Marck, C. 'DNA Strider': a 'C' program for the fast analysis of DNA and protein

sequences on the Apple Macintosh family of computers. *Nucleic Acids Res.* **16**,

1829-1836 (1988).

63. Luo, H., Lin, Y., Gao, F., Zhang, C.T. & Zhang, R. DEG 10, an update of the

database of essential genes that includes both protein-coding genes and

noncoding genomic elements. *Nucleic Acids Res.* **42**, D574-D580 (2014).

Acknowledgements

This work was supported by funding from the Wellcome Trust (092290/Z/10/Z) to VP. The France Génomique and French Bioinformatics Institute national infrastructures – funded as part of Investissement d'Avenir program managed by the Agence Nationale pour la Recherche (contracts ANR-10-INBS-09 and ANR-11-INBS-0013) – are acknowledged for support of the MicroScope annotation platform. We thank Angelika Gründling (Imperial College London) and Christoph Tang (University of Oxford) for critical reading of the manuscript.

Author contributions

V. P. was responsible for conception and supervision of the work, interpretation of data, and writing of manuscript. A. M., I. G., A. C. and V. P performed the experimental studies. A. B. and D. V. performed the computational studies.

Competing interests

The authors declare no competing interests.

Table 1. **Essential meningococcal genes listed by functional category/sub-category.** For this classification, we used predictions from MultiFun[24], MetaCyc[25], eggNOG[26], COG[27], FIGfam[28], and/or InterProScan[29]. The corresponding datasets are listed in Supplementary Table 8.

Functional category/sub-category	Number of genes	% of total
gene/protein expression	**150**	**38.4**
ribosomal protein	51	13
aminoacyl-tRNA synthesis	24	6.1
protein export	21	5.4
RNA modification/degradation	19	4.9
DNA transcription	11	2.8
translation factors	10	2.6
protein folding	7	1.8
ribosome biogenesis/maturation	5	1.3
protein post-translational modification	2	0.5
genome/cell replication	**33**	**8.4**
chromosome replication/maintenance	20	5.1
cell division	13	3.3
cell membrane/wall biogenesis	**54**	**13.8**
peptidoglycan biosynthesis/recycling	17	4.3
LOS biosynthesis	14	3.6
fatty acid biosynthesis	12	3.1
phospholipids biosynthesis	8	2
UDP-GlcNAc biosynthesis	3	0.8
cytosolic metabolism	**120**	**30.7**
vitamins biosynthesis	24	6.1
nucleotide biosynthesis	15	3.8
aerobic respiration/cytochrome c	12	3.1
amino acid metabolism	9	2.3
isoprenoid biosynthesis	9	2.3
electron transport	9	2.3
heme biosynthesis	9	2.3
sugar metabolism	9	2.3
Fe-S cluster biosynthesis	6	1.5
CoA/ac-CoA biosynthesis	6	1.5
ubiquinol biosynthesis	4	1
NAD/NADP biosynthesis	3	0.8
lipoate biosynthesis	2	0.5
SAM biosynthesis	1	0.3
phosphate metabolism	1	0.3
sulfur metabolism	1	0.3
unassigned	**34**	**8.7**
Total	**391**	**100**

Table 2. **Mutants in the complete NeMeSys 2.0 library of meningococcal mutants affected for two functions mediated by T4P: aggregation and twitching motility.** The other 1,569 mutants can form aggregates and exhibit twitching motility. N/A, not assayable because we could assess twitching motility only in mutants forming aggregates. The corresponding datasets are listed in Supplementary Table 10.

Mutated gene	Product	Aggregation	Twitching
Genes previously known to be involved in T4P biology in _N. meningitidis_			
pilE	major pilin PilE	-	N/A
pilT	type IV pilus retraction ATPase PilT	+ (irregular)	-
pilF	type IV pilus extension ATPase PilF	-	N/A
pilD	leader peptidase / N-methyltransferase PilD	-	N/A
pilG	type IV pilus biogenesis protein PilG	-	N/A
pilW	type IV pilus biogenesis lipoprotein PilW	-	N/A
pilX	minor pilin PilX	-	N/A
pilK	type IV pilus biogenesis protein PilK	-	N/A
pilJ	type IV pilus biogenesis protein PilJ	-	N/A
pilI	type IV pilus biogenesis protein PilI	-	N/A
pilH	type IV pilus biogenesis protein PilH	-	N/A
pilZ	PilZ protein	-	N/A
pilM	type IV pilus biogenesis protein PilM	-	N/A
pilN	type IV pilus biogenesis protein PilN	-	N/A
pilO	type IV pilus biogenesis protein PilO	-	N/A
pilP	type IV pilus biogenesis lipoprotein PilP	-	N/A
pilQ	type IV pilus secretin PilQ	-	N/A
Genes not previously known to be involved in T4P biology in _N. meningitidis_			
tsaP	secretin-associated protein TsaP	-	N/A
NMV_1205	conserved hypothetical periplasmic protein	-	N/A
NMV_2228	putative carbonic anhydrase	+/-	N/A

 Legends to Figures

Fig. 1. Flowchart of the construction of the NeMeSys 2.0 complete collection of

mutants in *N. meningitidis* 8013. As for similar efforts in other bacteria[10,12,13], we

first selected protein-coding genes to be targeted by systematic mutagenesis,

excluding 85 genes (4.1 %) because they encode transposases of repeated insertion

sequences, or correspond to short remnants of truncated genes or cassettes

(Supplementary Table 1). We then followed a two-step mutagenesis approach

explained in the text and in Supplementary Fig. 1. In brief, we first selected a subset

of sequence-verified Tn mutants from a previously constructed arrayed library[16,17].

Mutations were re-transformed in strain 8013 and PCR-verified. We thus selected

801 Tn mutants with a disrupting transposon in the corresponding target genes. Next,

we systematically mutagenised the remaining 1,174 target genes using a validated

no-cloning mutagenesis method relying on sPCR[20]. For each successful

transformation, two colonies were isolated and PCR-verified. To minimise false

positive identification of essential genes, each transformation that yielded no

transformants was repeated at least three times. In total, we could construct an

additional 783 mutants, generating an ordered library of defined mutants in 1,584

meningococcal genes (Supplementary Table 3). This effort also identified 391

candidate essential genes, which could not be disrupted, encoding proteins required

for *N. meningitidis* growth on rich medium (Supplementary Table 4).

Fig. 2. Partition of essential meningococcal genes into persistent, shell, and

cloud genomes. To perform this analysis, we used the PPanGGOLiN[21] method as

explained in the text. **a**) Partition of the 2,060 genes in the genome of *N. meningitidis*

8013. The corresponding datasets are listed in Supplementary Table 5. **b**) Partition of

the subset of 391 essential genes identified in this study. The corresponding datasets

are listed in Supplementary Table 5.

Fig. 3. Comparison of *N. meningitidis* essential genome to that of other bacteria. a) Edwards-Venn diagram displaying overlaps between essential genes in bacteria in which complete libraries of mutants have been constructed: *N. meningitidis* (black line), *E. coli* (green line)[10], *A. baylyi* (red line)[12] and *S. sanguinis* (blue line)[13]. To perform this analysis, we queried the DEG database[63] of essential genes using our set of essential genes. The corresponding datasets are listed in Supplementary Table 6. **b**) Comparison of *N. meningitidis* essential genome (black line) to JCVI-syn3.0 (red line), a synthetic *M. mycoides* designed with a minimal genome[22]. The corresponding datasets are listed in Supplementary Table 7.

Fig. 4. Partition of the meningococcal essential genes into four major functional groups. Gene/protein expression (38.4 %, orange), genome/cell replication (8.4 %, green), cell membrane/wall biogenesis (13.8 %, blue), and cytosolic metabolism (30.7 %, yellow). Only 34 essential genes (8.3 %) could not be clearly assigned to one of these four categories. The corresponding datasets are listed in Supplementary Table 8.

Fig. 5. Concise cellular overview of the essential meningococcal genome. Essential genes were integrated into networks and metabolic pathways using primarily MetaCyc[25]. The four basic functional groups are highlighted using the same colour as in Fig. 4, *i.e.* orange (gene/protein expression), green (genome/cell replication), blue (cell membrane/wall biogenesis), and yellow (cytosolic metabolism). The 34 essential genes that could not be clearly assigned to one of these four categories are not represented on the figure. Genes involved in the different reactions are indicated by their name or NMV_ label, in red when essential, in black when dispensable. *Genes involved in more than one pathway. Key compounds/proteins are abbreviated as follows. ACP, acyl carrier protein; AIR,

aminoimidazole ribotide; CDP-DAG, CDP-diacylglycerol; CMP-Kdo, CMP-

ketodeoxyoctonate; DHAP, dihydroxyacetone phosphate; DPP, dimethylallyl

diphosphate; DXP, 1-deoxyxylulose-5P; Fe-S, iron-sulfur; FMN, flavin

mononucleotide; G3P, glyceraldehyde-3P Glu-6P, glucose-6P; IMP, inosine

monophosphate; IPP, isopentenyl diphosphate; LOS, lipo-oligosaccharide; M-DAP,

meso-diaminopimelate; NaMN, nicotinate D-ribonucleotide; OPP, all-*trans*-octaprenyl

diphosphate; PE, phosphatidylethanolamine; PG, phosphatidylglycerol; PHBA, *p*-

hydroxybenzoate; PRPP, 5-phosphoribosyl diphosphate; Rib-5P, ribulose-5P; SAM,

S-adenosyl-methionine; UDP-GlcNAc, UDP-*N*-acetyl-glucosamine; UP, di-*trans*-poly-

cis-undecaprenyl monophosphate; UPP, di-*trans*-poly-*cis*-undecaprenyl diphosphate.

The corresponding datasets are listed in Supplementary Table 8.

Fig. 6. Essential genes in RGP – NMV_1479 and NMV-0559 – are conditionally

essential. a) Gene organisation of the putative antitoxin NMV_1478 (highlighted in

red) with its neighbouring NMV_1479 toxin. Results of the mutagenesis (✂) are

shown. ⬭, viable mutant; ☠, lethal phenotype. **b**) Gene organisation of the *tps*

RGP (RGP_0) to which NMV_0559 (highlighted in red) belongs (Supplementary

Table 9). Genes on the + strand are in white, genes on the - strand are in black.

Results of the mutagenesis (✂) are shown. ⬭, viable mutant; ☠, lethal phenotype.

In contrast to NMV_0559, each of the other target genes in the *tps* RGP could be

mutated individually (not shown for readability).

Fig. 7. NMV_1317 encodes a novel histidinol-phosphatase. a) Histidine

biosynthesis pathway in the meningococcus. PRPP, 5-phosphoribosyl diphosphate;

PRFAR, phosphoribulosylformimino-AICAR-P; IGP, erythro-imidazole-glycerol-P. **b**)

Growth on M9 minimal medium, with or without added histidine (His). The plates also

contained 0.5 mM IPTG for inducing expression of the complementing genes. WT,

strain 8013; Δ1317, ΔNMV_1317 mutant; Δ1317::1317, ΔNMV_1317 complemented

with NMV_1317; Δ1317::$hisB_{EC}$, ΔNMV_1317 cross-complemented with $hisB_{EC}$ from

E. coli, which encodes the unrelated histidinol-phosphatase present in this species;

Δ1718, ΔNMV_1718 mutant; Δ1718::$hisB_{EC}$, control showing that ΔNMV_1718 (*hisH*)

cannot be cross-complemented by $hisB_{EC}$.

Fig. 8. Assaying piliation in the mutants in genes not previously associated

with T4P biology in *N. meningitidis*. The WT strain and a non-piliated *ΔpilD* mutant

were included as positive and negative controls, respectively. *ΔtsaP*, *ΔtsaP* mutant;

Δ1205, ΔNMV_1205 mutant; Δ2228, ΔNMV_2228 mutant. **a**) T4P purified using a

shearing/precipitation method were separated by SDS-PAGE and either stained with

Coomassie blue (upper panel) or analysed by immunoblotting using an antibody

against the major pilin PilE (lower panel)[58]. Samples were prepared from equivalent

numbers of cells and identical volumes were loaded in each lane. MW, molecular

weight marker lane, with values in kDa. **b**) T4P were quantified by whole-cell ELISA

using a monoclonal antibody specific for the filaments of strain 8013[57]. Equivalent

numbers of cells, based on OD_{600} readings, were applied to the wells of microtiter

plates. Results are expressed in % piliation (ratio to WT) and are the average ±

standard deviations from five independent experiments. For statistical analysis, one-

way ANOVAs followed by Dunnett's multiple comparison tests were performed (*** *p*

< 0.0001).

Fig 9. Functional analysis of the mutants in genes not previously associated

with T4P biology in *N. meningitidis*. The WT strain and a non-piliated *pilD* mutant

were included as positive and negative controls, respectively. **a**) Aggregation in liquid

culture as assessed by phase-contrast microscopy[38]. *ΔtsaP::tsaP*, *ΔtsaP*

complemented with *tsaP*; *ΔtsaP/ΔpilT*, double mutant in *tsaP* and *pilT*; Δ1025::1205,

ΔNMV_1205 complemented with NMV_1205; Δ1205/*ΔpilT*, double mutant in

NMV_1205 and *pilT*; Δ2228::2228, ΔNMV_2228 complemented with NMV_2228. **b**)

Quantification of the competence for DNA transformation. Equivalent numbers of

recipient cells were transformed using a *rpoB* PCR product containing a point

mutation leading to rifampicin resistance. Results are expressed as transformation

frequencies and are the average ± standard deviations from four independent

experiments. For statistical analysis, one-way ANOVAs followed by Dunnett's

multiple comparison tests were performed (*** $p < 0.0001$).

Figure 1

Figure 2

a

S. sanguinis *N. meningitidis* 199 *A. bayli*

85 53

40 10 26 7

10 118

5 2

12 77

55 17

E. coli

b

216 175 263

N. meningitidis JCVI-syn3.0

Figure 3

Figure 4

Figure 5

Figure 6

a

PRPP
↓ *hisG + hisZ*
phosphoribosyl-ATP
↓ *hisE*
phosphoribosyl-AMP
↓ *hisI*
phosphoribosyl-formimino-aicar-P
↓ *hisA*
PRFAR
↓ *hisH + hisF*
IGP
↓ *hisB*
imidazole acetol-P
↓ *hisC*
histidinol-P
↓ NMV_1317
histidinol
↓ *hisD*
histidinal
↓ *hisD*
L-histidine

b

WT · D1778::*hisB$_{EC}$* · D1317 · D1778 · D1317::1317 · D1317::*hisB$_{EC}$*

M9 (no His)

WT · D1778::*hisB$_{EC}$* · D1317 · D1778 · D1317::1317 · D1317::*hisB$_{EC}$*

M9 (+ His)

Figure 7

Figure 8

Figure 9