1 **Bacterial gene essentiality under modeled microgravity**
2
3

4 Emanuel Burgos[1], Madeline M. Vroom[2], Ella Rotman[3§], Megan Murphy-Belcaster[3], Jamie S.
5 Foster[2], Mark J. Mandel[1,3,*]
6
7
8 [1] Department of Medical Microbiology and Immunology, University of Wisconsin-Madison,
9 Madison, WI USA
10
11 [2] Department of Microbiology and Cell Science, Space Life Science Lab, University of Florida,
12 Merritt Island, FL USA
13
14 [3] Department of Microbiology-Immunology, Northwestern University Feinberg School of
15 Medicine, Chicago, IL USA
16
17
18 [§] Current Address: Department of Microbiology, University of Chicago, Chicago, IL USA
19
20
21

22 Short title: *Vibrio fischeri* modeled microgravity mutant analysis
23
24 Keywords: modeled microgravity, essential genes, symbiosis, *Vibrio fischeri*, *Aliivibrio fischeri*,
25 *Euprymna scolopes*, INSeq, Tn-seq
26
27

28 * Correspondence to:
29
Mark J. Mandel
University of Wisconsin-Madison
Department of Medical Microbiology and Immunology
1550 Linden Drive
Madison, WI 53706
Phone: (608) 261-1170
Fax: (608) 262-8418
Email: mmandel@wisc.edu
ORCID: https://orcid.org/0000-0001-6506-6711

39
40
41

ABSTRACT

To support long-duration spaceflight, identifying genes that enable microbial survival and

stability under space stressors is crucial, as these factors can affect both host health and in-

space biomanufacturing efforts. Our objective was to determine what bacterial genes are

required for growth in culture under modeled, or simulated, microgravity conditions compared to

normal gravity controls. We focused on the marine bacterium *Vibrio fischeri*, which forms a

monospecific symbiosis with the Hawaiian bobtail squid, *Euprymna scolopes*. The symbiosis

has been studied during spaceflight and in ground-based modeled microgravity conditions.

Using transposon insertion sequencing (INSeq), we identified dozens of genes that exhibited

fitness defects under both conditions, yet we identified relatively few genes with differential

effects under modeled microgravity or gravity specifically. We additionally compared RNA-seq

and INSeq data and determined that expression under microgravity was not predictive of the

essentiality of a given gene. In summary, empirical determination of conditional gene

essentiality identifies few microgravity-specific genes for environmental growth of *V. fischeri*,

suggesting that the condition of microgravity has a minimal impact on symbiont gene

requirement during growth in media. These findings suggest that maintaining beneficial

microbial communities during spaceflight, and enabling their use in biomanufacturing

applications, may require minimal genetic adaptation, easing the challenges of supporting

healthy symbiotic relationships on long-duration missions.

INTRODUCTION

Spaceflight alters microbial physiology and their interactions with eukaryotes, and as the

number and duration of manned space missions expands beyond low Earth orbit, elucidating

the underlying mechanisms of these changes is essential[1–10]. To address these issues,

experiments are needed to examine bacterial growth, bacterial-host interactions, and host

health under spaceflight-like conditions. Although previous studies using both natural and

modeled microgravity conditions have resulted in a wide range of physiological and genetic

responses, many initial studies primarily targeted pathogenic strains[11–13]. More recently, studies

have begun to examine the effects of microgravity on beneficial microbes that associate with

animals and how spaceflight impacts these interactions[4–6,10,14,15]. Given the importance of

beneficial microbes to host health, we sought here to ask a fundamental question of what genes

are required for bacterial growth in microgravity.

Due to the number of logistical constraints in conducting spaceflight experiments, several

ground-based platforms exist to simulate the low-shear environment of microgravity[16]. Rotating

cell culture systems with High Aspect-Ratio Vessels (HARVs) represent one such platform as

they mimic the low-shear fluid conditions that occur in low Earth orbit and have been used for

decades to model microgravity environments[17]. As cells grow in the HARVs, the hydrodynamic

forces within each vessel offset the effects of gravity and the cells are essentially in "freefall"

and maintained in a constant suspension (**Fig. 1**). Bacterial strains grown in these low-shear

modeled microgravity (LSMMG) conditions can experience a wide range of different

physiological responses, including increases in growth rates, biofilm formation, secondary

metabolite production, environmental stress responses, and antibiotic resistance[6,18–23].

Fig. 1. Experimental setup for bacterial mutant enrichment in HARV vessels to model
microgravity (LSMMG) and to have the control gravity condition. INSeq libraries were
grown in HARV vessels that can simulate microgravity when placed perpendicular or gravity
when horizontal. Mutants exhibited either no defects in either condition or exhibited (1) a growth
defect in both conditions, (2) LSMMG defect, (3) or a gravity-specific defect.

The symbiosis of *Vibrio fischeri* colonization of the light organ of the Hawaiian bobtail squid,

Euprymna scolopes, has emerged as a valuable model to study the effects of microgravity on

host-microbe dynamics and function[1,10,24,25]. The squid hatches each generation without its

bacterial partner and then proceeds to harvest *V. fischeri* from the seawater[26]. The bacteria

colonize the dedicated symbiotic light organ of the host, where bioluminescence from the

bacteria is projected downward and camouflages the host in the moonlight[27]. This system has

been especially valuable to understand the molecular basis by which animals acquire specific

microbes from the environment[28]. The bacteria are genetically manipulatable, and the site of

colonization can be imaged directly in live animals, enabling studies that have revealed much of

the molecular dialogue between the two partners[29–32]. The specificity of the relationship, in

which only certain strains of *V. fischeri* can colonize the squid host, provides a strong model

system to study animal-microbiome formation, development, and evolution[26]. Furthermore, the

small size of the host animal and simplicity of the symbiosis has contributed to the value of this

system for studying host-microbe interactions during spaceflight[1,3,10,33].

In this study, we take a complementary approach to previous work on bacterial gene expression

under simulated microgravity. Previous studies using the squid-vibrio system have focused

extensively on the impact of simulated and actual spaceflight conditions on the host squid

physiology in the presence and absence of *V. fischeri*[1,3,10,33,34]. Results have shown that under

both simulated and actual spaceflight, the colonization of the host by *V. fischeri* results in the

down-regulation of several host genes associated with oxidative and heat stress, suggesting

that being in a symbiotic state is beneficial to the host during spaceflight. These studies do not,

however, examine the impact of the space-like environment on the symbiont, specifically, what

genes are required for survival under microgravity and modeled microgravity conditions. Other

studies in the symbiont have applied a genome-scale genetic approach technology to globally

identify *V. fischeri* genes that are essential, and conditionally essential, under specific

environmental conditions and during growth in media under laboratory conditions[35]. Here, we

apply a similar global approach to determine what genes are required for bacterial growth under

simulated microgravity conditions to ascertain the overall impact that microgravity-like conditions

have on symbiont health and physiology.

RESULTS

Global analysis of bacterial genes required for growth in low-shear modeled

microgravity. A number of studies have examined differential gene expression in beneficial

microbes under microgravity or simulated microgravity conditions[36–39], including in the marine

bacterium *V. fischeri*[39,40]. The current study asks a distinct, yet related, question of what genes

are differentially required for growth under simulated microgravity conditions compared to a

gravity control condition. To identify mutants with a competitive growth defect in simulated

microgravity, we applied the transposon insertion sequencing (INSeq/Tn-seq) approach to *V.*

fischeri strains cultivated under HARV conditions in both the low-shear modeled microgravity

(LSMMG) and gravity positions (**Fig. 1**)[40,41]. We started with a characterized mutant library of

over 40,000 transposon mutants in *V. fischeri* strain ES114[35]. The resulting "input" library was

introduced into HARV vessels under either LSMMG or normal gravity conditions (1 x *g*). We

simultaneously examined the mutant library grown in LSMMG and gravity control conditions

after growth for approximately 15 generations, and each resulting "output" pool of mutants was

frozen. A total of six LSMMG biological replicates and three gravity control replicates were

compared to the input library. From each sample, DNA was isolated, from which INSeq libraries

were constructed[41]. The libraries were sequenced on an Illumina HiSeq 2000 instrument, and

the location of each transposon insertion was mapped using the pyinseq Python package.

For each sample, we obtained between 6.65 - 8.18 x 10^6 Illumina reads (**Table S1**). We

observed similar numbers and patterns of unique transposon hits across the samples tested,

suggesting that the library did not undergo significant bottlenecks during the experiment (**Table**

S1). Next, for each sample, we examined the normalized transposon insertion counts (CPM;

counts-per-million Illumina reads) in each gene. We compared the similarity of these gene-level

counts across the samples in the analysis using pairwise correlation analysis. All pairwise

comparisons had a high level of similarity (Spearman R^2 > 0.95), and the HARV-grown samples

were clearly distinguishable from the input libraries (**Fig. 2**). Examination of the heat map

suggested little overall differentiation between the LSMMG and gravity samples that were

otherwise grown similarly. We proceeded to compare individual genes that were depleted under

LSMMG, gravity, or both conditions.

Fig. 2. Pairwise correlation between samples revealed a clear difference between input
samples and output samples from the HARV conditions. Spearman correlations between
samples, in which the normalized transposon insertion counts in each gene were compared.
Input replicates were the original INSeq library grown for inoculation into the HARV, and low-
shear modeled microgravity (LSMMG) and normal gravity replicates were outputs from the
respective HARV-grown experiments.

To identify genes that had significant differential depletion under modeled microgravity we used

DESeq2 to calculate the median representation of each mutant in the output LSMMG or gravity

pools, and plotted those values compared to the input pool. We focused our analysis on genes

that were depleted at least 2-fold in the different conditions. Genes that did not meet these

criteria, genes that were poorly represented in the input pool, and genes for which a previous

study suggested that mutants impaired bacterial growth were excluded from this analysis (**Fig.**

3; open black circles). A total of 109 genes exhibited depletion under both LSMMG and gravity

conditions (**Fig. 3**; filled circles). Of these genes, most were similarly depleted under both

conditions. However, there were genes in this group that had a ≥ 2-fold difference in one

condition (LSMMG or gravity) relative to the other, including 10 genes more depleted under

gravity conditions and one gene (*rodA*) more depleted under LSMMG. Furthermore, there were

two genes in the analysis that were significantly depleted ($p < 0.05$ from DESeq2 analysis)

under gravity conditions and not under LSMMG conditions (*flgD, rfaD*). There were no genes

that were only significantly depleted under LSMMG conditions.

Fig. 3. Mutant behavior under LSMMG or gravity revealed fitness in each HARV condition,
normalized to the input transposon sequencing library. A) Comparison scatter plot of
microgravity (x-axis) and gravity (y-axis) Log(Fold-Change) values. Counts for each gene for
each replicate were normalized and used to calculate Log_{10}(Fold-Change) values as described
in Materials and Methods. Genes that were poorly represented in the input pool, did not exhibit
significant depletion, or were previously determined to be growth deficient in LBS medium were
designated as open black circles. Filled black circles are genes that were significantly (p-value <
0.05) depleted 2-fold change under both simulated microgravity and gravity conditions. Genes
that are filled red were significant in both conditions but more depleted under gravity, whereas
open red circles are exclusively significant in gravity. Filled blue genes are significant in both,
but more depleted under simulated microgravity. Remaining genes are shown as open black
circles.

Validation of INSeq results through defined mutant competitions. The above results were

derived from an analysis of complex mutant pools with > 40,000 mutants. We therefore sought

to determine whether we would observe the same behavior using one-versus-one competitions

between defined mutant strains and the parental strain. For this analysis, we considered the

entire data set (i.e., Table S3), and we proceeded to isolate mutant strains that had a depletion

from the INSeq analysis, as well as mutants in two control genes that were not depleted under

either condition (*brnQ* and *nhaR*). We examined mutants that had a range of fitness defects and

that were represented in a mutant collection in the the laboratory. Each defined mutant strain

was grown in culture, then competed in the HARVs against the parental strain that carries the

LacZ-expressing plasmid pVSV103[42]. The input and output pools from each experiment were

plated onto LBS-Xgal medium, and the ratios of blue:white colonies in the samples were

calculated. A competitive index was calculated to determine the fitness of each mutant under

each condition. The results of this analysis are plotted in **Fig. 4**. There was strong overall

concordance between the INSeq and defined competition results. All the genes for which

mutants were significantly depleted under INSeq were also substantially depleted in the defined

competitions, and the control mutants exhibited no substantial depletion. Therefore, we

conclude that the INSeq analysis can reliably predict conditional gene requirements in the

simulated microgravity environment.

Examination of the relationship between bacterial gene requirement and gene induction

in modeled microgravity. We recently examined bacterial genes for which mRNA levels are

induced in the modeled microgravity condition, compared to gravity controls[42]. We analyzed our

current study of gene requirement (i.e., INSeq) with the previous dataset on gene expression

218   **Fig. 4. Competition of defined mutants provided validation of mutant fitness from within**
219   **the complex INSeq library.** Mutants were competed against wild-type *V. fischeri* under
220   simulated microgravity and gravity using the HARV vessels. Dots represent fold-change for 1:1
221   competition of mutant versus marked parental strain (MJM1575).

222
223

Fig. 5. Gene expression (RNA-Seq) does not predict conditional gene requirement
(INSeq) under LSMMG conditions. INSeq (y-axis) and RNA-Seq (x-axis) were compared by
using log-normalized fold-change values for different experimental conditions. A) INSeq
(LSMMG/Gravity) was compared to a previously published RNA-Seq (LSMMG/Gravity) under
HARV conditions and had an R^2 of 0.08. We also calculated fold-changes for RNA-Seq HARV
conditions using a dataset from SWT medium. B) INSeq (Gravity/Input) were compared to RNA-
Seq (Gravity/SWT) and had an R^2 of 0.18. C) INSeq (LSMMG/Input) was compared to RNA-
Seq (LSMMG/SWT) and had an R^2 0.11.(i.e., RNA-Seq) to ask whether there is a correlation
between gene requirement and gene expression. As shown in **Fig. 5A**, there was no significant
correlation between gene requirement and transcript induction (Pearson R^2 = 0.08). Our data
above suggest that the HARV platform may play a more significant role in shaping mutant
communities than the specific LSMMG or gravity conditions as has been determined for other
bacterial strains[43]. Therefore, we explored whether genes required in the HARV are also
induced in the HARV when normalized to non-HARV samples. For our INSeq data, we
normalized to the input sample that did not experience the HARV condition, and for RNA-Seq,
we normalized to published data on culture-grown *V. fischeri*[44]. As shown in **Fig. 5B-C**, there
was a lack of overall correlation between the gene expression and gene requirement (Pearson
R^2 = 0.18, 0.11 for **Fig. 5B, 5C**, respectively). Therefore, we conclude that gene requirement in
the HARV cannot be predicted from transcriptome induction data, emphasizing the need for
empirical determination of gene essentiality.

DISCUSSION

This study provides a global view of *V. fischeri* genes that are required for growth in conditions

that simulate microgravity. Given the prominence of modeled microgravity platforms, such as

HARVs, in examining the impact of spaceflight on animal-microbe interactions, a motivation for

this work was to understand how the basic requirements for bacterial survival and growth are

altered during growth in the microgravity analog platforms. A surprising finding was that there

was little difference in gene requirement under these conditions, despite dramatic differences in

gene expression. These findings, and their implications, are discussed in greater detail below.

The global data provided by INSeq, along with the comparison of the INSeq data with results

from defined mutants, provided sensitive internal controls to examine growth in multiple HARV

samples. Overall, the data we obtained from HARV samples were highly consistent. As shown

in the heat map in **Fig. 2**, the high correlation of INSeq replicates within a treatment (e.g., for

LSMMG samples) suggests that there was no substantial bottleneck during inoculation or

growth in the HARVs. Furthermore, a strong correlation between INSeq results and the results

with defined mutants (**Fig. 4**) provides support that the global data are representative of gene-

level data. In fact, the consistency we observed between vessel replicates was also observed

between LSMMG and gravity samples in the HARVs. The heat map in **Fig. 2**, coupled with the

tight correlation of the samples in **Fig. 3** (Pearson R^2 = 0.96), illustrates that there was little

variation observed between the LSMMG and gravity conditions. A striking consistency between

genes required for bacterial growth under gravity and those required for growth under modeled

microgravity is the major finding from this study. As the *V. fischeri* system is used for more

extensive research on microgravity and is emerging as a tool for space biomanufacturing[45], this

result indicates there will not be a confounding effect of genes that are simply required for

bacterial growth under microgravity. Put another way, in the future, if genes are identified that

are required for colonization under microgravity, they are likely to be required specifically for

interaction with the animal host, and not simply for growth under this altered gravity condition.

A common follow-up experiment for INSeq experiments is to validate the global data with

individual gene mutants. We conducted such an experiment in **Fig. 4** and found that the results

were consistent across over two orders of magnitude. Another common follow-up experiment is

to complement back the wild-type allele for mutants under study to then demonstrate that the

wild-type phenotype is restored. Given the negative result that we observed—i.e., that mutants

did not display different effects in simulated microgravity compared to the gravity control—there

was not the impetus to conduct this experiment. However, as described above, we argue that

this negative finding is an important one given the expectation of the experiment, and given the

rigor with which we validated the results with the single-gene mutants in **Fig. 4**.

Despite the overall patterns of concordance between LSMMG and gravity, there were genes for

which we observed differential effects between the two conditions, as plotted in **Fig. 3** and

detailed in **Table S2**. For example, depletion in *rodA (mrdB)* under both LSMMG and gravity

was observed, yet the gene was more significantly depleted under LSMMG. RodA is a SEDS-

family peptidoglycan polymerase that has multiple effects on bacterial cell shape and division [46].

This result suggests that differences in bacterial morphology under the two different conditions

may impact the genetic requirement for *rodA*, though we note that its absence does affect

growth under both regimes.

We additionally identified several genes with mutants depleted under both conditions but that

were more significantly depleted under gravity (**Fig. 3**, filled red dots). Notably, multiple genes

for the F0F1-ATPase were depleted under both conditions, but more so under gravity.

Interestingly, prior work in *Escherichia coli* demonstrated that a similar set of genes was

essential for aerobic growth in minimal glycerol media, even though the metabolic model used

predicted that they would not be required[47]. Together with our results, this suggests that this

subset of genes (*atpA, atpB, atpC, atpF*) may perform a function separate from their role in ATP

synthesis. Finally, there were two genes for which mutants exhibited significant depletion under

gravity but not LSMMG: *flgD* (encoding the hook capping protein FlgD) and *rfaD* (encoding the

LPS biosynthesis enzyme ADP-L-glycero-D-mannoheptose 6-epimerase). Both genes affect the

outer surface of the bacteria, and consistent with the *rodA* results above, support the idea that

the bacterial envelope is most susceptible to differential effects of gravity.

It is also important to note that we observed dozens of genes for which mutants exhibited similar

depletion under both LSMMG and gravity conditions (**Fig. 3**; black filled dots). We note that

significant growth defects for mutants in these genes were not observed during 15 generations

of growth in LBS medium under aerobic conditions[35]. Therefore, these genes are likely required

for robust growth in the HARV, but not for growth in LSMMG versus gravity. Given the

anaerobic environment of the HARV, it seems likely that many of these genes may be required

for optimal growth under anaerobic conditions. This hypothesis is supported by the presence of

four genes of the Na^+-translocating NADH:quinone oxidoreductase (*nqrA*, *nqrB*, *nqrD*, *nqrE*),

which, in the related species *V. cholerae*, conducts 90% of the membrane NADH

dehydrogenase activity under anaerobic conditions[48,49]. We also note the presence of some

genes that are required for robust symbiosis in this category, including *degS*, *dnaJ*, and

ompU[35,50]. Given that mutants in these genes exhibited growth defects in both LSMMG and

gravity conditions in the HARVs, our results suggest that colonization in the HARV may proceed

differently than under standard laboratory conditions. Furthermore, we speculate that these

genes may play a role in *V. fischeri* growth under anaerobic conditions.

A possible limitation of the current study is that the transposon library used was built on agar

plates in a standard microbiology laboratory (i.e., under normal gravity conditions). However,

were this a major limitation, then in **Fig. 3**, we would have observed many mutants with no

defect in gravity but with a defect in LSMMG (e.g., in the top left of the figure). Not only did we

not observe a substantial number of such genes, but they were outnumbered by the genes that

fell in the bottom right. Therefore, we can conclude that the origination of the library under unit

gravity conditions did not impair our ability to investigate this question.

We provide a comparison of our INSeq data with previously published RNA-Seq data. As shown

in **Fig. 5**, there is little correlation between gene requirement (INSeq) and gene expression

(RNA-Seq). It is important to consider both gene requirement and gene expression, as genes

that are not induced in a condition may nonetheless be required for growth and/or survival. For

example, a gene that is expressed at a constant level under two conditions may be required for

survival in only one of those conditions. A broader analysis comparing gene requirement and

expression has demonstrated that these categories are often unlinked[51]. Our results argue that

genes that are induced in simulated microgravity are not preferentially required for bacterial

growth under these conditions.

Overall, this work identifies genes required under exposure to LSMMG, unit gravity, or the

general conditions within the analog HARV environment. Importantly, these results demonstrate

that microgravity-induced transcripts do not predict which genes are conditionally essential for

survival in this environment. Finally, we demonstrated that the HARV environment is amenable

to high-throughput genetic experiments and was reproducible within treatments. By laying this

groundwork, our findings enable deeper exploration of the genetic mechanisms underlying how

microbes adapt and thrive during spaceflight. Such insights are crucial for maintaining healthy

host-microbe associations and advancing space biomanufacturing, since enabling reliable

biological production processes in reduced gravity is key to optimizing resource utilization and

supporting long-term human missions beyond low Earth orbit.

MATERIALS AND METHODS

Media and growth conditions. *V. fischeri* strains (**Table 1**) were grown at 25°C in Luria-Bertani

salt (LBS) medium (per liter, 10 g Bacto-tryptone, 5 g yeast extract and 20 g NaCl, 50 ml 1 M

Tris buffer, pH 7.5, in distilled water). When appropriate, antibiotics or supplements were added

to media at the following concentrations: erythromycin, 5 µg/ml; chloramphenicol, 5 µg/ml; X-gal

(5-bromo-4-chloro-3-indolyl-β-D-galactopyranoside), 80 µg/ml. Growth media were solidified

with 1.5% agar as needed.

Table 1. *V. fischeri* strains used in this study.

Strain	Genotype	Reference/Source
MJM1100	Wild-type ES114	[52,53]
MJM1538	MJM1100/pLostfoX	[54]
MJM1575	MJM1100/pVSV103	[42,55]
MJM3801	MJM1100 *sspA*::Tn*erm* (transformed from VFS025D03)	This work
MJM3802	MJM1100 *mukB*::Tn*erm* (transformed from VFS001A10)	This work
MJM3804	MJM1100 *ruvB*::Tn*erm* (transformed from VFS020B09)	This work
MJM3805	MJM1100 *brnQ*::Tn*erm* (transformed from VFS022E03)	This work
MJM3807	MJM1100 *nhaR*::Tn*erm* (transformed from VFS022G01)	This work
MJM3808	MJM1100 *ubiH*::Tn*erm* (transformed from MJM1976)	This work
MJM3809	MJM1100 *hslO*::Tn*erm* (transformed from MJM1628)	This work

Preparation of input and output samples from HARV growth. The Lib04 library of pMarVF1

mariner transposon insertions in MJM1100 (ES114) was described previously[35]. Twenty-five

microliters of the library (approximately 2×10^8 CFU) were inoculated into 5 ml LBS, and grown

at 26 °C for approximately 5 h, to an OD_{600} of approximately 0.3. Aliquots (552 µl) of culture was

transferred to each of eight flasks containing 100 ml LBS (i.e., $2^{7.5}$-fold dilution) and mixed well;

an additional 500 µl of input culture was frozen in 17% glycerol (v/v) as the input sample. From

each flask, 50 ml was inoculated into each HARV: LSMMG vessels A, B, C, and D; and gravity

vessels A, B, C, and D. Each vessel D was used to monitor growth rate. Samples were grown

for approximately 5.25 h to an OD_{600} ≤ 0.3; diluted another $2^{7.5}$-fold; grown to a final OD_{600} of

approximately 0.3, then frozen as the 15-generation output samples. In some cases, HARV

vessels leaked or had significantly reduced growth rate, and those samples were excluded from

further analysis.

INSeq sample preparation. DNA from the input and output samples was prepared using the

MoBio Biofilm Isolation Kit (Carlsbad, CA). Samples were prepared for INSeq analysis using the

protocol of Goodman *et al.*[41], with double the BioSamA primer concentration. Samples were

submitted to the Tufts University Core Facility (TUCF) for sequencing on the Illumina HiSeq

2500 (single-end 50 bp reads). The resulting reads were deposited at NCBI SRA under

accession number SRR12394639.

Bioinformatic Analysis of INSeq data. Each sample was processed using the bioinformatic

software pyinseq (https://github.com/mjmlab/pyinseq, v0.2.0) to quantify transposon insertions

and then analyzed using Python visualization and statistical modules. Pyinseq starts by

demultiplexing the raw reads using a barcode index and then maps them to the reference

genome (ES114v2: CP000020.2, CP000021.2, CP000022.1) using the short-read aligner

software Bowtie[56] with parameters that allow a single base-pair mismatch. Reads with multiple

alignments (e.g., to the 12 semi-redundant rRNA operons and in tRNA genes) were excluded.

The output alignment files were used to quantify the frequency of reads at each transposon

insertion site (TA dinucleotides, for the mariner transposon). Transposon-insertion sites were

analyzed if they contain a minimum of three reads and have reads from both the left and right

flanking sequence (with a maximum difference of 10-fold abundance for one side over the

other). For each sample, a T50 value was calculated, which is defined as the minimum number

of transposon insertion sites that account for 50% of the reads in that sample. Gene-level

analysis consolidates the site-level data for insertions that fall in the 5'-most 90% of each gene

(-d parameter of 0.9). The pyinseq summary gene table (**Table S3**) was further analyzed using

pandas, a python module for manipulating large datasets [57,58]. In addition, pandas was used to

calculate spearman correlation values of each pairwise sample comparison. The technical

averages for each sample was further analyzed using DESeq2, an R package that normalizes

the dataset and performs appropriate statistical tests on high-throughput count data that does

not follow a normal distribution such as INSeq[59]. DESeq2 was used to estimate the library size,

normalize sequencing depth of samples, and calculate variation of each gene for statistical

testing.

Transformation of VFS into MJM1100 background. The VFS mutant collection was

assembled by sequencing random pEVS170 transposon insertions into ES114[35,60]. Mutant

alleles were moved into the MJM1100 background using transformation under *tfox* induction[61].

Selected VFS strains grown on LBS plates were verified with PCR amplification using locus-

specific primers and transposon-anchored primers MJM-127 (pEVS170 transposon) or SamA

(pMarVF1 transposon) (**Table 2**). Recipient strain MJM1538 (MJM1100 carrying *pLostfoX)* was

grown overnight in 3 ml of LBS containing 2.5 µg/ml chloramphenicol and then subcultured

1:100 into 3 ml of Tris Minimal with *N*-acetylglucosamine (GlcNac) for overnight growth. The

recipient was then subcultured 1:50 into fresh Tris-Minimal-GlcNac-Cam with aeration until the

OD_{600} reached 0.2-0.3, when 500 µl of recipient culture was incubated with 2.4 µg of VFS donor

DNA (prepared with the Qiagen Blood and Tissue kit, Gram-negative bacteria protocol),

followed by a brief vortex, and then static incubation at room temperature for 30 min. One ml of

LBS was added to culture, transferred to a glass culture tube, and incubated overnight with

aeration. The culture was spun down (8000 x *g*, 1 min), 900 µl of supernatant was removed, and

the pellet was resuspended in the remaining ~100 µl of LBS. Aliquots (50 µl) of each sample

were plated on LBS-erythromycin (LBS-Erm; 5 µg/ml) and three candidates were selected.

Colonies were restreaked on LBS-Erm plates and then patched on selective media to check for

absence of *pLostfoX* (LBS-Cam[S]) and the presence of the transformed DNA (LBS-Erm[R]).

Transformation was verified with PCR amplification using primers that target the transposon

junction and the gene target (**Table 2**).

431 **Table 2. Oligonucleotide primers**

Name	Sequence (5`-3`)	Source
MJM-127	ACAAGCATAAAGCTTGCTCAATCAATCACC	[62]
SamA	AAGCAGAAGACGGCATACGAAGACC	[41]
EB01 (*sspA*)	AACGGAGCTAAGTAACAATCCACT	This work
EB02 (*mukB*)	CACCTAGTGTATCAGCAAGACCTT	This work
EB03 (*ruvB*)	TCACTCGCTTTCAATTAACTCAGC	This work
EB06 (*brnQ*)	CAGCAACTTTCGCAGCATCTAATA	This work
EB08 (*nhaR*)	TTGTAAGTCCATATAGACGCCCAC	This work
EB10 (*hslO*)	AACATGGTTGATGGAGGTAAAGTT	This work
EB11 (*ubiH*)	CAAATTCAATCTTAGCAAAGCTGTC	This work

Competitions of individual gene mutants with a marked parental strain in the HARVs.

The competitive fitness of seven transposon mutants (**Table 1**) was evaluated in one-on-one

competition assays against the LacZ-expressing wild type, MJM1575, in the HARV vessels.

Overnight cultures of each strain were prepared in LBS with shaking at 25 °C and diluted 1:100

in fresh medium the following morning. After 2 h of growth at 25 °C with shaking, the mutant and

wild-type subcultures were normalized by OD_{600} and combined at a 1:1 ratio in fresh LBS. The

resultant starting culture was subsequently loaded into the HARV vessel. At this time, two

samples of the mixed input culture were collected: the first was preserved in 33% glycerol (v/v)

and stored at -80 °C for later analysis, whereas the second was serially diluted in PBS (pH 7.0)

and plated on LBS-Xgal agar (2 mg/ml) to reveal the starting ratio of mutant:control, based on

the proportion of white-to-blue colonies, as previously described [35]. In the HARVs, the mutant

and marked wild type were grown in competition with each other under LSMMG or gravity

conditions for approximately 10 generations at 25 °C and 13 rpm. After 10 doublings, which was

determined by OD_{600}, samples of the output cultures were preserved in 33% glycerol (v/v) and

stored at -80 °C. The input and output samples were plated on LBS-Xgal agar to calculate the

competitive index for each sample. The competitive index is equal to the Log_{10} value of the

mutant/wild type ratio after competition normalized to its measured ratio at the beginning of the

competition.

DATA AVAILABILITY

Illumina data for the INSeq reads are available at NCBI SRA, Accession number

SRR12394639.

ACKNOWLEDGMENTS

We acknowledge Cheryl Whistler and Randi Foxall for VFS Library mutants and A. Murat Eren

and his laboratory for technical advice and mentorship. Funding was provided by NASA's Illinois

Space Grant Consortium Research Seed Grant (M.J.M.), NIGMS R35 GM119627 (M.J.M.), an

American Society for Microbiology Undergraduate Research Fellowship (E.B.), and the McNair

Scholars Program (E.B.). The project was also supported in part by a NASA Space Biology

award 80NSSC18K1465 (J.S.F.). The funders played no role in study design, data collection,

analysis and interpretation of data, or the writing of this manuscript.

AUTHOR CONTRIBUTIONS

EB analyzed the data and wrote the paper. MMV conducted the HARV experiments. ER and

MMB conducted the INSeq library preparation. MJM and JSF conceived of the study, obtained

funding for the study, and wrote and edited the manuscript.

COMPETING INTERESTS

All authors declare no financial or non-financial competing interests.

SUPPLEMENTAL DATA

The file Supplemental_Tables.xlsx includes the following tables:

Table S1: INSeq sample details

Table S2: List of genes depleted as shown in Figure 3

Table S3: Summary gene table output from pyinseq

REFERENCES

1.   Foster, J. S., Khodadad, C. L. M., Ahrendt, S. R. & Parrish, M. L. Impact of simulated

microgravity on the normal developmental time line of an animal-bacteria symbiosis. *Sci.*

Rep. **3**, 1340 (2013).

2.   Voorhies, A. A. & Lorenzi, H. A. The challenge of maintaining a healthy microbiome during

long-duration space missions. *Frontiers in Astronomy and Space Sciences* **3**, 23 (2016).

3.   Casaburi, G., Goncharenko-Foster, I., Duscher, A. A. & Foster, J. S. Transcriptomic

changes in an animal-bacterial symbiosis under modeled microgravity conditions. *Sci. Rep.*

7, 46318 (2017).

4.   Castro-Wallace, S., Stahl, S., Voorhies, A., Lorenzi, H. & Douglas, G. L. Response of

Lactobacillus acidophilus ATCC 4356 to low-shear modeled microgravity. *Acta Astronaut.*

139, 463–468 (2017).

5.   Shao, D. *et al.* Simulated microgravity affects some biological characteristics of

Lactobacillus acidophilus. Appl. Microbiol. Biotechnol. **101**, 3439–3449 (2017).

6.   Duscher, A. A. *et al.* Transcriptional profiling of the mutualistic bacterium *Vibrio fischeri* and

an *hfq* mutant under modeled microgravity. *npj Microgravity* **4**, 25 (2018).

7.   Garrett-Bakelman, F. E. *et al.* The NASA Twins Study: A multidimensional analysis of a

year-long human spaceflight. *Science* **364**, eaau8650 (2019).

8.   Nickerson, C. A. *et al.* Microbiology of human spaceflight: microbial responses to

mechanical forces that impact health and habitat sustainability. *Microbiol. Mol. Biol. Rev.*

88, e0014423 (2024).

9.   Winer, D. A. *et al.* Astroimmunology: the effects of spaceflight and its associated stressors

on the immune system. *Nat. Rev. Immunol.* **26**, 189–212 (2025).

10.   Koch, E. J. *et al.* Beneficial microbes mitigate molecular stress responses and accelerate

developmental pathways in host animals during spaceflight. *Front. Space Technol.* **7**,

1791484 (2026).

11. Chang, D. *et al.* A multi-omic analysis of an *Enterococcus faecium* mutant reveals specific

genetic mutations and dramatic changes in mRNA and protein expression. *BMC Microbiol.*

13, 304 (2013).

12. Wilson, J. W. *et al.* Media ion composition controls regulatory and virulence response of

Salmonella in spaceflight. *PLoS One* **3**, e3923 (2008).

13. Simões, M. F. & Antunes, A. Microbial pathogenicity in space. *Pathogens* **10**, 450 (2021).

14. Tesei, D., Jewczynko, A., Lynch, A. M. & Urbaniak, C. Understanding the complexities and

changes of the astronaut microbiome for successful long-duration space missions. *Life*

(Basel) **12**, 495 (2022).

15. Gonzalez, E. *et al.* Spaceflight alters host-gut microbiota interactions. *NPJ Biofilms*

Microbiomes **10**, 71 (2024).

16. Nickerson, C. A., Ott, C. M., Wilson, J. W., Ramamurthy, R. & Pierson, D. L. Microbial

responses to microgravity and other low-shear environments. *Microbiol. Mol. Biol. Rev.* **68**,

345–361 (2004).

17. Schwarz, R. P., Goodwin, T. J. & Wolf, D. A. Cell culture for three-dimensional modeling in

rotating-wall vessels: an application of simulated microgravity. *J. Tissue Cult. Methods* **14**,

51–57 (1992).

18. Fang, A., Pierson, D. L., Koenig, D. W., Mishra, S. K. & Demain, A. L. Effect of simulated

microgravity and shear stress on microcin B17 production by *Escherichia coli* and on its

excretion into the medium. *Appl. Environ. Microbiol.* **63**, 4090–4092 (1997).

19. Demain, A. L. & Fang, A. Secondary metabolism in simulated microgravity. *Chem. Rec.* **1**,

333–346 (2001).

20. Castro, S. L., Nelman-Gonzalez, M., Nickerson, C. A. & Mark Ott, C. Induction of

attachment-independent biofilm formation and repression of *hfq* expression by low-fluid-

shear culture of *Staphylococcus aureus*. *Appl. Environ. Microbiol.* **77**, 6368–6378 (2011).

21. Wang, H. *et al.* Increased biofilm formation ability in *Klebsiella pneumoniae* after short-term

exposure to a simulated microgravity environment. *Microbiologyopen* **5**, 793–801 (2016).

22. Abshire, C. F. *et al.* Exposure of *Mycobacterium marinum* to low-shear modeled

microgravity: effect on growth, the transcriptome and survival under stress. *NPJ*

Microgravity **2**, 16038 (2016).

23. Acres, J. M., Youngapelian, M. J. & Nadeau, J. The influence of spaceflight and simulated

microgravity on bacterial motility and chemotaxis. *NPJ Microgravity* **7**, 7 (2021).

24. Bongrand, C. & Foster, J. S. Modelled microgravity impacts *Vibrio fischeri* population

structure in a mutualistic association with an animal host. *Environ. Microbiol.* (2023)

doi:10.1111/1462-2920.16522.

25. Duscher, A. A., Vroom, M. M. & Foster, J. S. Impact of modeled microgravity stress on

innate immunity in a beneficial animal-microbe symbiosis. *Sci. Rep.* **14**, 2912 (2024).

26. Mandel, M. J. Models and approaches to dissect host-symbiont specificity. *Trends*

Microbiol. **18**, 504–511 (2010).

27. Jones, B. W. & Nishiguchi, M. K. Counterillumination in the Hawaiian bobtail squid,

Euprymna scolopes Berry (Mollusca: Cephalopoda). *Mar. Biol.* **144**, 1151–1155 (2004).

28. Mandel, M. J. & Dunn, A. K. Impact and influence of the natural *Vibrio*-squid symbiosis in

understanding bacterial-animal interactions. *Front. Microbiol.* **7**, 1982 (2016).

29. Koropatnick, T. A. *et al.* Microbial factor-mediated development in a host-bacterial

mutualism. *Science* **306**, 1186–1188 (2004).

30. Mandel, M. J., Wollenberg, M. S., Stabb, E. V., Visick, K. L. & Ruby, E. G. A single

regulatory gene is sufficient to alter bacterial host range. *Nature* **458**, 215–218 (2009).

31. Mandel, M. J. *et al.* Squid-derived chitin oligosaccharides are a chemotactic signal during

colonization by *Vibrio fischeri*. *Appl. Environ. Microbiol.* **78**, 4620–4626 (2012).

32. Brooks, J. F., 2nd & Mandel, M. J. The histidine kinase BinK Is a negative regulator of

biofilm formation and squid colonization. *J. Bacteriol.* **198**, 2596–2607 (2016).

33. Foster, J. S., Kerney, K. R., Parrish, M. L., Khodadad, C. L. M. & Ahrendt, S. R. Potential of

the *Euprymna/Vibrio* symbiosis as a model to assess the impact of microgravity on

bacteria-induced animal development. *Gravitational and Space Research* **25**, 44–47

(2011).

34. Vroom, M. M., Troncoso-Garcia, A., Duscher, A. A. & Foster, J. S. Modeled microgravity

alters apoptotic gene expression and caspase activity in the squid-vibrio symbiosis. *BMC*

Microbiol. **22**, 202 (2022).

35. Brooks, J. F., 2nd *et al.* Global discovery of colonization determinants in the squid symbiont

Vibrio fischeri. Proc. Natl. Acad. Sci. U. S. A. **111**, 17284–17289 (2014).

36. Orsini, S. S., Lewis, A. M. & Rice, K. C. Investigation of simulated microgravity effects on

Streptococcus mutans physiology and global gene expression. *NPJ Microgravity* **3**, 4

(2017).

37. Salavatifar, M., Ahmadi, S. M., Todorov, S. D., Khosravi-Darani, K. & Tripathy, A. Impact of

microgravity on virulence, antibiotic resistance and gene expression in beneficial and

pathogenic microorganisms. *Mini Rev. Med. Chem.* **23**, 1608–1622 (2023).

38. Lozzi, B. *et al.* Simulated microgravity triggers a membrane adaptation to stress in *E. coli*

REL606. *BMC Microbiol.* **25**, 362 (2025).

39. Wang, X. *et al.* Phenotypic, transcriptomic and metabolomic changes in *Klebsiella*

pneumoniae after long term exposure to simulated microgravity. *NPJ Microgravity* **11**, 35

(2025).

40. Goodman, A. L. *et al.* Identifying genetic determinants needed to establish a human gut

symbiont in its habitat. *Cell Host Microbe* **6**, 279–289 (2009).

41. Goodman, A. L., Wu, M. & Gordon, J. I. Identifying microbial fitness determinants by

insertion sequencing using genome-wide transposon mutant libraries. *Nat. Protoc.* **6**, 1969–

1980 (2011).

42. Dunn, A. K., Millikan, D. S., Adin, D. M., Bose, J. L. & Stabb, E. V. New *rfp-* and pES213-

derived tools for analyzing symbiotic *Vibrio fischeri* reveal patterns of infection and *lux*

expression in situ. *Appl. Environ. Microbiol.* **72**, 802–810 (2006).

43. Hauserman, M. R., Sullivan, L. E., James, K. L., Ferraro, M. J. & Rice, K. C. Response of

Staphylococcus aureus physiology and Agr quorum sensing to low-shear modeled

microgravity. *J. Bacteriol.* **206**, e0027224 (2024).

44. Thompson, L. R. *et al.* Transcriptional characterization of *Vibrio fischeri* during colonization

of juvenile *Euprymna scolopes*. *Environ. Microbiol.* **19**, 1845–1856 (2017).

45. Vroom, M. M., Rodriguez-Ocasio, Y., Lynch, J. B., Ruby, E. G. & Foster, J. S. Modeled

microgravity alters lipopolysaccharide and outer membrane vesicle production of the

beneficial symbiont *Vibrio fischeri*. *NPJ Microgravity* **7**, 8 (2021).

46. Meeske, A. J. *et al.* SEDS proteins are a widespread family of bacterial cell wall

polymerases. *Nature* **537**, 634–638 (2016).

47. Joyce, A. R. *et al.* Experimental and computational assessment of conditionally essential

genes in *Escherichia coli*. *J. Bacteriol.* **188**, 8259–8271 (2006).

48. Casutt, M. S., Schlosser, A., Buckel, W. & Steuber, J. The single NqrB and NqrC subunits

in the Na^+-translocating NADH: quinone oxidoreductase (Na^+-NQR) from *Vibrio cholerae*

each carry one covalently attached FMN. *Biochim. Biophys. Acta* **1817**, 1817–1822 (2012).

49. Steuber, J. *et al.* Central role of the Na^+-translocating NADH:quinone oxidoreductase (Na^+-

NQR) in sodium bioenergetics of *Vibrio cholerae*. *Biol. Chem.* **395**, 1389–1399 (2014).

50. Aeckersberg, F., Lupp, C., Feliciano, B. & Ruby, E. G. *Vibrio fischeri* outer membrane

protein OmpU plays a role in normal symbiotic colonization. *J. Bacteriol.* **183**, 6590–6597

(2001).

51. Turner, K. H., Everett, J., Trivedi, U., Rumbaugh, K. P. & Whiteley, M. Requirements for

Pseudomonas aeruginosa acute burn and chronic surgical wound infection. *PLoS Genet.*

10, e1004518 (2014).

52. Boettcher, K. J. & Ruby, E. G. Depressed light emission by symbiotic *Vibrio fischeri* of the

sepiolid squid *Euprymna scolopes*. *J. Bacteriol.* **172**, 3701–3706 (1990).

53. Mandel, M. J., Stabb, E. V. & Ruby, E. G. Comparative genomics-based investigation of

resequencing targets in *Vibrio fischeri*: focus on point miscalls and artefactual expansions.

BMC Genomics **9**, 138 (2008).

54. Brooks, J. F., 2nd, Gyllborg, M. C., Kocher, A. A., Markey, L. E. H. & Mandel, M. J. TfoX-

based genetic mapping identifies *Vibrio fischeri* strain-level differences and reveals a

common lineage of laboratory strains. *J. Bacteriol.* **197**, 1065–1074 (2015).

55. Post, D. M. B. *et al.* O-antigen and core carbohydrate of *Vibrio fischeri* lipopolysaccharide:

composition and analysis of their role in *Euprymna scolopes* light organ colonization. *J.*

Biol. Chem. **287**, 8515–8530 (2012).

56. Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. L. Ultrafast and memory-efficient

alignment of short DNA sequences to the human genome. *Genome Biol.* **10**, R25 (2009).

57. Reback, J. *et al. Pandas-Dev/Pandas: Pandas 1.1.0.* (2020). doi:10.5281/zenodo.3964380.

58. McKinney, W. & Others. Data structures for statistical computing in python. in *Proceedings*

of the 9th Python in Science Conference vol. 445 51–56 (Austin, TX, 2010).

59. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

60. Lyell, N. L., Dunn, A. K., Bose, J. L., Vescovi, S. L. & Stabb, E. V. Effective mutagenesis of

Vibrio fischeri by using hyperactive mini-Tn*5* derivatives. *Appl. Environ. Microbiol.* **74**,

7059–7063 (2008).

61. Pollack-Berti, A., Wollenberg, M. S. & Ruby, E. G. Natural transformation of *Vibrio fischeri*

requires *tfoX* and *tfoY*. *Environ. Microbiol.* **12**, 2302–2311 (2010).

62. Studer, S. V., Mandel, M. J. & Ruby, E. G. AinS quorum sensing regulates the *Vibrio*

fischeri acetate switch. *J. Bacteriol.* **190**, 5915–5923 (2008).