Bacterial gene essentiality under modeled microgravity

Emanuel Burgos[1], Madeline M. Vroom[2], Ella Rotman[3§], Megan Murphy-Belcaster[3†], Jamie S.
Foster[2], Mark J. Mandel[1,3,*]

[1] Department of Medical Microbiology and Immunology, University of Wisconsin-Madison,
Madison, WI USA

[2] Department of Microbiology and Cell Science, Space Life Science Lab, University of Florida,
Merritt Island, FL USA

[3] Department of Microbiology-Immunology, Northwestern University Feinberg School of
Medicine, Chicago, IL USA

[§] Current Address: Department of Microbiology, University of Chicago, Chicago, IL USA

[†] Current Address: University of Wisconsin-Madison School of Medicine and Public Health,
Madison, WI USA

Short title: *Vibrio fischeri* modeled microgravity mutant analysis

Keywords: modeled microgravity, essential genes, symbiosis, *Vibrio fischeri*, *Aliivibrio fischeri*,
Euprymna scolopes, INSeq, Tn-seq

* Correspondence to:

Mark J. Mandel
University of Wisconsin-Madison
Department of Medical Microbiology and Immunology
1550 Linden Drive
Madison, WI 53706
Phone: (608) 261-1170
Fax: (608) 262-8418
Email: mmandel@wisc.edu

ABSTRACT

The health of eukaryotic hosts is tightly connected to relationships with symbiotic

microorganisms, yet how these relationships develop and evolve during long-duration

spaceflight is not well understood. In this study, we asked what bacterial genes are required for

growth under modeled, or simulated, microgravity conditions compared to normal gravity

controls. To conduct this study, we focused on the marine bacterium *Vibrio fischeri*, which forms

a monospecific symbiosis with the Hawaiian bobtail squid, *Euprymna scolopes.* The symbiosis

has been studied during spaceflight and in ground-based modeled microgravity conditions. We

employed a library of over 40,000 *V. fischeri* transposon mutants and compared the fitness of

mutants in modeled microgravity compared to the gravity controls using transposon insertion

sequencing (INSeq). We identified dozens of genes that exhibited fitness defects under both

conditions, likely due to the controlled anaerobic environment, yet we identified relatively few

genes with differential effects under modeled microgravity or gravity specifically: only mutants in

rodA were more depleted under modeled microgravity, and mutants in 12 genes exhibited

greater depletion under gravity conditions. We additionally compared RNA-seq and INSeq data

and determined that expression under microgravity was not predictive of the essentiality of a

given gene. In summary, empirical determination of conditional gene essentiality identifies few

microgravity-specific genes for environmental growth of *V. fischeri*, suggesting that the condition

of microgravity has a minimal impact on symbiont gene requirement.

IMPORTANCE

There is substantial evidence that both the host immune system and microbial physiology are

altered during space travel. It is difficult to discern the molecular mechanisms of these

processes in a complex microbial consortium and during the short durations of experiments in

space. By using a model organism that is amenable to high-throughput genetic approaches, we

have determined that *V. fischeri* does not require a separate genetic repertoire for media growth

in modeled microgravity versus gravity conditions. Our results argue that future studies on how

this organism forms a specific and stable association with its animal host will not be confounded

by growth effects in the environment. The identification of similar genetic requirements under

modeled microgravity and gravity suggest that fitness pressures on microbiome growth in space

may be similar to those on Earth and may not negatively impact their animal hosts during long-

duration spaceflight.

INTRODUCTION

During spaceflight, microbes--and microbe-host relationships--are altered, and understanding

the mechanisms of these effects are critical as people, plants, and animals spend more time in

space (1–7). To address this issue, experiments are needed to examine bacterial growth,

bacterial-host interactions, and host health. Although previous studies using both natural and

modeled microgravity conditions have resulted in a wide range of physiological and genetic

responses, most have primarily targeted pathogenic strains (8, 9). While fewer studies have

examined the effects of microgravity on beneficial microbes, there are recent studies that have

begun to examine this issue (4–6). Given the importance of beneficial microbes to host health,

we sought here to ask a fundamental question of what genes are required for bacterial growth in

microgravity.

Due to the number of logistical constraints in conducting spaceflight experiments, several

ground-based platforms exist to mimic the low-shear environment of microgravity (10). Rotating

High Aspect-Ratio Vessels (HARVs) represent one such platform; they mimic the low-shear fluid

conditions that occur at low Earth orbit, and have been used for decades to simulate

microgravity environments (11). As cells grow in the HARVs, the hydrodynamic forces within

each vessel offset the effects of gravity and the cells are essentially in "freefall" and maintained

in a constant suspension (Fig. 1). Bacterial strains grown in these low-shear modeled

microgravity (LSMMG) conditions can experience a wide range of different physiological

responses, including increases in growth rates, biofilm formation, secondary metabolite

production, environmental stress responses, and antibiotic resistance (6, 12–16).

The model symbiosis of *Vibrio fischeri* colonization of the light organ of the Hawaiian bobtail

squid, *Euprymna scolopes*, has emerged as a valuable model to study the effects of

microgravity on microbiome assembly and function (1). The squid hatches each generation

without its bacterial partner, and then proceeds to harvest *V. fischeri* from the seawater (17).

The bacteria colonize the dedicated symbiotic light organ of the host, where bioluminescence

from the bacteria is projected downward and camouflages the host in the moonlight (18). This

system has been especially valuable to understand the molecular basis by which animals

acquire specific microbes from the environment. The bacteria are genetically manipulatable, and

the site of colonization can be imaged directly in live animals, enabling studies that have

revealed much of the molecular dialogue between the two partners (19–22). The specific

relationship--only *V. fischeri*, and only specific strains of *V. fischeri*, can colonize the squid host--

provides a strong model system to study animal-microbiome formation, development, and

evolution (17). Furthermore, the small size of the host animal and simplicity of the symbiosis has

contributed to the value of this system for studying microbiome formation during spaceflight (1,

23).

In this study, we take a complementary approach to previous work on bacterial gene expression

under simulated microgravity to ask what genes are required for survival under this condition.

Our lab has previously published a study to globally identify *V. fischeri* genes that are essential,

and that are conditionally essential under specific environmental conditions (24). Here we apply

a similar global approach to determine what genes are required for bacterial growth under

simulated microgravity conditions to ascertain the overall impact that microgravity-like conditions

have on symbiont health and physiology.

RESULTS

A number of studies have examined differential-gene expression in beneficial microbes under

microgravity or simulated microgravity conditions, including in the marine bacterium *V. fischeri*

(6). The current study asks a distinct, yet related, question of what genes are differentially

required for growth under simulated microgravity conditions compared to a gravity control

condition. To identify mutants with a competitive growth defect in simulated microgravity, we

applied the transposon insertion sequencing (INSeq/Tn-seq) approach to *V. fischeri* strains

cultivated under HARV conditions in both the LSMMG and gravity positions (Fig. 1) (25, 26). We

started with a characterized mutant library of over 40,000 transposon mutants in *V. fischeri*

strain ES114 (24). The resulting "input" library was introduced into HARV vessels under either

LSMMG or normal gravity conditions (1 x *g*). We simultaneously examined the mutant library

grown in LSMMG and gravity control conditions after growth for approximately 15 generations,

and each resulting "output" pool of mutants was frozen. A total of six LSMMG biological

replicates and three gravity control replicates were compared to the input library. From each

sample, DNA was isolated, from which INSeq libraries were constructed (26). The libraries were

sequenced on an Illumina HiSeq 2000 instrument, and the location of each transposon insertion

was mapped using the pyinseq Python package.

For each sample, we obtained between 6.65 - 8.18 x 10^6 Illumina reads (Table S1). We

observed similar numbers and patterns of unique transposon hits across the samples tested,

suggesting that the library did not undergo significant bottlenecks during the experiment (Table

S1). Next, for each sample, we examined the normalized transposon insertion counts (CPM;

counts-per-million Illumina reads) in each gene. We compared the similarity of these gene-level

counts across the samples in the analysis using pairwise correlation analysis. All pairwise

comparisons had a high level of similarity (Spearman R^2 > 0.95), and the HARV-grown samples

were clearly distinguishable from the input libraries (Fig. 2). Examination of the heat map

suggested little overall differentiation between the LSMMG and gravity samples that were

otherwise grown similarly. We proceeded to compare individual genes that were depleted under

LSMMG, gravity, or both conditions. To identify genes that had significant differential depletion

under modeled microgravity we used DESeq2 to calculate the median representation of each

mutant in the output LSMMG or gravity pools, and plotted those values compared to the input

pool. We focused our analysis on genes that were depleted at least 2-fold in the different

conditions. Genes that did not meet these criteria, genes that were poorly represented in the

input pool, and genes for which a previous study suggested that mutants impaired bacterial

growth were excluded from further analyses (Fig. 3; open black circles). A total of 109 genes

exhibited depletion under both LSMMG and gravity conditions (Fig. 3; filled circles). Of these

genes, most were similarly depleted under both conditions. However, there were genes in this

group that had a ≥ 2-fold difference in one condition (LSMMG or gravity) relative to the other:

this includes 10 genes more depleted under gravity and one gene (*rodA*) more depleted under

LSMMG. Furthermore, there were two genes in the analysis that were significantly depleted ($p <$

0.05 from DESeq2 analysis) under gravity conditions and not under LSMMG conditions (*flgD,*

rfaD). There were no genes that were only significantly depleted under LSMMG conditions.

The above results were derived from an analysis of complex mutant pools with > 40,000

mutants. We therefore sought to determine whether we would observe the same behavior using

one-versus-one competitions between defined mutant strains and the parental strain. We

proceeded to isolate mutant strains that had a depletion from the INSeq analysis, as well as

mutants in two control genes that were not depleted under either condition (*brnQ* and *nhaR*).

Each defined mutant strain was grown in culture, then competed in the HARVs against the

parental strain that carries the LacZ-expressing plasmid pVSV103 (27). The input and output

pools from each experiment were plated onto LBS-Xgal medium, and the ratio of blue:white

colonies in the samples were calculated. A competitive index was calculated to determine the

fitness of each mutant under each condition. Plotted in Figure 4 are the results of this analysis.

Although some differences were observed between the massive INSeq competition and the

defined one-vs-one competitions, there was strong concordance between the INSeq and

defined competition results. All of the genes for which mutants were significantly depleted under

INSeq were also substantially depleted in the defined competitions, and the control mutants

exhibited no substantial depletion. Therefore, we conclude that the INSeq analysis can reliably

predict conditional gene requirements in the HARV environment.

We recently examined bacterial genes for which mRNA levels are induced in the modeled

microgravity condition, compared to gravity controls (6). We analyzed our current study of gene

requirement (i.e., INSeq) with the previous dataset on gene expression (i.e., RNA-Seq) to ask

whether there is a correlation between gene requirement and gene expression. As shown in

Figure 5A, there was no significant correlation between gene requirement and transcript

induction (Pearson R^2 = 0.08). Our data above suggest that the HARV platform may play a

more significant role in shaping mutant communities than the specific LSMMG or gravity

conditions. Therefore, we asked whether genes required in the HARV are also induced in the

HARV, when normalized to non-HARV samples. For our INSeq data, we normalized to the input

sample that did not experience the HARV condition, and for RNA-Seq we normalized to

published data on culture-grown *V. fischeri* (28). As shown in Figure 5B-C, there is similarly a

lack of overall correlation between the gene expression and gene requirement (Pearson R^2 =

0.18, 0.11 for Fig. 5B, 5C, respectively). Therefore, we conclude that gene requirement in the

HARV cannot be predicted from transcriptome induction data, emphasizing the need for

empirical determination of gene essentiality.

DISCUSSION

This study provides a global view of *V. fischeri* genes that are required for growth in conditions that simulate microgravity. Given the prominence of modeled microgravity platforms, such as HARVs, in examining the impact of spaceflight on animal-microbe interactions, a motivation for this work was to understand how the basic requirements for bacterial survival and growth are altered during growth in the HARV platforms. A surprising finding was that there is little difference in gene requirement under these conditions, despite dramatic differences in gene expression. These findings, and their implications, are discussed in greater detail below.

The global data provided by INSeq, along with the comparison of the INSeq data with results from defined mutants, provided sensitive internal controls to examine growth in multiple HARV samples. Overall, the data we obtained from HARV samples was highly consistent. As shown in the heat map in Figure 2, the high correlation of INSeq replicates within a treatment (e.g., for LSMMG samples) argues that there was no substantial bottleneck during inoculation or growth in the HARVs. Furthermore, strong correlation between INSeq results and the results with defined mutants (Fig. 4) provides support that the global data are representative of gene-level data. In fact, the consistency we observed between vessel replicates was also observed between HARV LSMMG and gravity samples. The heat map in Fig. 2 coupled with the tight correlation of the samples in Figure 3 (Pearson R^2 = 0.96) illustrate that there was little variation observed between the LSMMG and gravity conditions. A striking consistency between genes

required for bacterial growth under gravity and those required for growth under modeled

microgravity is the major finding from this study. As the *V. fischeri* system is used for more

extensive research on microgravity, this result indicates there will not be a confounding effect of

genes that are simply required for bacterial growth under microgravity. Put another way, in the

future, if genes are identified that are required for colonization under microgravity, they are likely

to be required specifically for interaction with the animal host, and not simply for growth under

this altered gravity condition. Our findings, in concert with previous studies showing limited

changes in gene expression during spaceflight, suggest that spaceflight missions will have

minimal negative consequences on the microbiome.

Despite the overall patterns of concordance between LSMMG and gravity, there were genes for

which we observed differential effects between the two conditions, as plotted in Figure 3 and

detailed in Table S3. For example, depletion in *rodA (mrdB)* under both LSMMG and gravity

was observed, yet the gene was more significantly depleted under LSMMG. RodA is a SEDS-

family peptidoglycan polymerase that has multiple effects on bacterial cell shape and division

(29). This result suggests that differences in bacterial shape under the two different conditions

may impact the genetic requirement for *rodA*, though we note that its absence does affect

growth under both regimes. We additionally identified a number of genes with mutants depleted

under both conditions but that were more significantly depleted under gravity (Figure 3, filled red

dots). Notably, multiple genes for the F0F1-ATPase were depleted under both conditions, but

more so under gravity. Interestingly, prior work in *Escherichia coli* demonstrated that a similar

set of genes were essential for aerobic growth in minimal glycerol media, even though the

metabolic model used predicted that they would not be required (30). Together with our results,

this suggests that this subset of genes (*atpA, atpB, atpC, atpF*) may perform a function separate

from the role in ATP synthesis. Finally, there were two genes for which mutants exhibited

significant depletion under gravity but not LSMMG: *flgD* (encoding the hook capping protein

FlgD) and *rfaD* (encoding the LPS biosynthesis enzyme ADP-L-glycero-D-mannoheptose 6-

epimerase). Both genes affect the outer surface of the bacteria, and consistent with the *rodA*

results above, support the idea that the bacterial envelope is most susceptible to differential

effects of gravity.

It is also important to note that we observed dozens of genes for which mutants exhibited similar

depletion under both LSMMG and gravity conditions (Fig. 3; black filled dots). We note that

significant growth defects for mutants in these genes was not observed during 15 generations of

growth in LBS medium under aerobic conditions (24). Therefore, these genes are likely required

for robust growth in the HARV, but not for growth in LSMMG versus gravity. Given the

anaerobic environment of the HARV, it seems likely that many of these genes may be required

for optimal growth under anaerobic conditions. This hypothesis is supported by the presence of

four genes of the Na$^+$-translocating NADH:quinone oxidoreductase (*nqrA*, *nqrB*, *nqrD*, *nqrE*),

which, in the related species *V. cholerae*, conducts 90% of the membrane NADH

dehydrogenase activity under anaerobic conditions (31, 32). We also note the presence of some

genes that are required for robust symbiosis in this category, including *degS*, *dnaJ*, and *ompU*

(24, 33). Given that mutants in these genes exhibited growth defects in both LSMMG and

gravity conditions in the HARVs, our results suggest that colonization in the HARV may proceed

differently than under standard laboratory conditions. Furthermore, we speculate that these

genes may play a role in *V. fischeri* growth under anaerobic conditions.

A possible limitation of the current study is that the transposon library used was built on agar

plates in a standard microbiology laboratory; i.e., under normal gravity conditions. However,

were this a major limitation, then in Figure 3 we would have observed a large number of

mutants with no defect in gravity but with a defect in LSMMG (e.g., in the top left of the figure).

Not only did we not observe a substantial number of such genes, but they were outnumbered by

the genes that fell in the bottom right. Therefore, we can conclude that the origination of the

library in the gravity condition did not impair our ability to investigate this question.

We provide a comparison of our INSeq data with previously published RNA-Seq data. As shown

in Figure 5, there is little correlation between gene requirement (INSeq) and gene expression

(RNA-Seq). It is important to consider both gene requirement and gene expression, as genes

that are not induced in a condition may nonetheless be required for growth and/or survival. For

example, a gene that is expressed at a constant level under two conditions may be required for

survival in only one of those conditions. Broader analysis comparing gene requirement and

expression have demonstrated that these categories are often unlinked (34). Our results argue

that genes that are induced in simulated microgravity are not preferentially required for bacterial

growth under these conditions.

Overall, this work establishes which genes were impacted by growth in LSMMG, gravity, or the

HARV environment generally. We additionally determined that microgravity-induced transcripts

do not predict which genes are conditionally essential under this condition. Finally, we

demonstrated that the HARV environment is amenable to high-throughput genetic experiments

and was reproducible within treatments. This study, therefore, provides a solid foundation for

future studies that seek to identify the genetic basis by which bacteria and animals form specific,

robust interactions under conditions of reduced gravity.

MATERIALS AND METHODS

Media and growth conditions. *V. fischeri* strains were grown at 25°C in Luria-Bertani salt

(LBS) medium (per liter, 10 g Bacto-tryptone, 5 g yeast extract and 20 g NaCl, 50 ml 1 M Tris

buffer, pH 7.5, in distilled water). When appropriate, antibiotics or supplements were added to

media at the following concentrations: erythromycin, 5 µg/ml; chloramphenicol, 5 µg/ml; X-gal

(5-bromo-4-chloro-3-indolyl-β-D-galactopyranoside), 80 µg/ml. Growth media were solidified

with 1.5% agar as needed.

Preparation of input and output samples from HARV growth. The Lib04 library of pMarVF1

mariner transposon insertions in MJM1100 (ES114) was described previously (24). Twenty-five

microliters of the library (approximately 2 x 10^8 CFU) were inoculated into 5 ml LBS, and grown

at 26 °C for approximately 5 h, to an OD_{600} of approximately 0.3. Aliquots (552 µl) of culture was

transferred to each of eight flasks containing 100 ml LBS (i.e., $2^{7.5}$-fold dilution) and mixed well;

an additional 500 µl of input culture was frozen in 17% glycerol (v/v) as the input sample. From

each flask, 50 ml was inoculated into each HARV: LSMMG vessels A, B, C, and D; and gravity

vessels A, B, C, and D. Each vessel D was used to monitor growth rate. Samples were grown

for approximately 5.25 h to an OD_{600} ≤ 0.3; diluted another $2^{7.5}$-fold; grown to a final OD_{600} of

approximately 0.3, then frozen as the 15-generation output samples. In some cases, HARV

vessels leaked or had significantly reduced growth rate, and those samples were excluded from

further analysis.

INSeq sample preparation. DNA from the input and output samples was prepared using the

MoBio Biofilm Isolation Kit (Carlsbad, CA). Samples were prepared for INSeq analysis using the

protocol of Goodman et al. (26), with double the BioSamA primer concentration. Samples were

submitted to the Tufts University Core Facility (TUCF) for sequencing on the Illumina HiSeq

2500 (single-end 50 bp reads). The resulting reads were deposited at NCBI SRA under

accession number SRR12394639.

Bioinformatic Analysis of INSeq data. Each sample was processed using the bioinformatic

software pyinseq (https://github.com/mjmlab/pyinseq, v0.2.0) to quantify transposon insertions

and then analyzed using Python visualization and statistical modules. Pyinseq starts by

demultiplexing the raw reads using a barcode index and then maps them to the reference

genome (ES114v2: CP000020.2, CP000021.2, CP000022.1) using the short-read aligner

software Bowtie (35) with parameters that allow a single base-pair mismatch. Reads with

multiple alignments (e.g., to the 12 semi-redundant rRNA operons and in tRNA genes) were

excluded. The output alignment files were used to quantify the frequency of reads at each

transposon insertion site (TA dinucleotides, for the mariner transposon). Transposon-insertion

sites were analyzed if they contain a minimum of three reads and have reads from both the left

and right flanking sequence (with a maximum difference of 10-fold abundance for one side over

the other). For each sample, a T50 value was calculated, which is defined as the minimum

number of transposon insertion sites that account for 50% of the reads in that sample. Gene-

level analysis consolidates the site-level data for insertions that fall in the 5'-most 90% of each

gene (-d parameter of 0.9). The pyinseq summary gene table (Table S2) was further analyzed

using pandas, a python module for manipulating large datasets (36, 37). In addition, pandas

was used to calculate spearman correlation values of each pairwise sample comparison. The

technical averages for each sample was further analyzed using DESeq2, an R package that

normalizes the dataset and performs appropriate statistical tests on high-throughput count data

that does not follow a normal distribution such as INSeq (38). DESeq2 was used to estimate the

library size, normalize sequencing depth of samples, and calculate variation of each gene for

statistical testing.

Transformation of VFS into MJM1100 background. The VFS mutant collection was

assembled by sequencing random pEVS170 transposon insertions into ES114 (24, 39). Mutant

alleles were moved into the MJM1100 background using transformation under *tfox* induction

(40). Selected VFS strains grown on LBS plates were verified with PCR amplification using

locus-specific primers and transposon-anchored primers MJM-127 (pEVS170 transposon) or

SamA (pMarVF1 transposon) (Table 2). Recipient strain MJM1538 (MJM1100 carrying

pLostfoX) was grown overnight in 3 ml of LBS containing 2.5 µg/ml chloramphenicol and then

subcultured 1:100 into 3 ml of Tris Minimal with *N*-acetylglucosamine (GlcNac) for overnight

growth. The recipient was then subcultured 1:50 into fresh Tris-Minimal-GlcNac-Cam with

aeration until the OD_{600} reached 0.2-0.3, when 500 µl of recipient culture was incubated with 2.4

µg of VFS donor DNA (prepared with the Qiagen Blood and Tissue kit, Gram-negative bacteria

protocol), followed by a brief vortex, and then static incubation at room temperature for 30 min.

One ml of LBS was added to culture, transferred to a glass culture tube, and incubated

overnight with aeration. The culture was spun down (8000 x *g*, 1 min), 900 µl of supernatant

was removed, and the pellet was resuspended in the remaining ~100 µl of LBS. Aliquots (50 µl)

of each sample were plated on LBS-erythromycin (LBS-Erm; 5 µg/ml) and three candidates

were selected. Colonies were restreaked on LBS-Erm plates and then patched on selective

media to check for absence of *pLostfoX* (LBS-Cam[S]) and the presence of the transformed DNA

(LBS-Erm^R). Transformation was verified with PCR amplification using primers that target the

transposon junction and the gene target (Table 2).

Competitions of individual gene mutants with a marked parental strain in the HARVs.

The competitive fitness of seven transposon mutants (Table 1) was evaluated in one-

on-one competition assays against the LacZ-expressing wild type, MJM1575, in the

HARV vessels. Overnight cultures of each strain were prepared in LBS with shaking at

25 °C and diluted 1:100 in fresh medium the following morning. After 2 h of growth at 25

°C with shaking, the mutant and wild-type subcultures were normalized by OD$_{600}$ and

combined at a 1:1 ratio in fresh LBS. The resultant starting culture was subsequently

loaded into the HARV vessel. At this time, two samples of the mixed input culture were

collected: the first was preserved in 33% glycerol (v/v) and stored at -80 °C for later

analysis, whereas the second was serially diluted in PBS (pH 7.0) and plated on LBS-

Xgal agar (2 mg/ml) to reveal the starting ratio of mutant:control, based on the

proportion of white-to-blue colonies, as previously described (24). In the HARVs, the

mutant and marked wild type were grown in competition with each other under LSMMG

or gravity conditions for approximately 10 generations at 25 °C and 13 rpm. After 10

doublings, which was determined by OD$_{600}$, samples of the output cultures were

preserved in 33% glycerol (v/v) and stored at -80 °C. The input and output samples

were plated on LBS-Xgal agar to calculate the competitive index for each sample. The

competitive index is equal to the Log_{10} value of the mutant/wild type ratio after

competition normalized to its measured ratio at the beginning of the competition.

DATA AVAILABILITY

Illumina data for the INSeq reads are available at NCBI SRA, Accession number

SRR12394639.

TABLES

Table 1. *V. fischeri* strains used in this study.

Strain	Genotype	Reference/Source
MJM1100	Wild-type ES114	(41, 42)
MJM1538	MJM1100/pLostfoX	(43)
MJM1575	MJM1100/pVSV103	(27, 44)
MJM3801	MJM1100 *sspA*::Tn*erm* (transformed from VFS025D03)	This work
MJM3802	MJM1100 *mukB*::Tn*erm* (transformed from VFS001A10)	This work
MJM3804	MJM1100 *ruvB*::Tn*erm* (transformed from VFS020B09)	This work
MJM3805	MJM1100 *brnQ*::Tn*erm* (transformed from VFS022E03)	This work
MJM3807	MJM1100 *nhaR*::Tn*erm* (transformed from VFS022G01)	This work
MJM3808	MJM1100 *ubiH*::Tn*erm* (transformed from MJM1976)	This work
MJM3809	MJM1100 *hslO*::Tn*erm* (transformed from MJM1628)	This work

404   **Table 2. Oligonucleotide primers**

Name	Sequence (5`-3`)	Source
MJM-127	ACAAGCATAAAGCTTGCTCAATCAATCACC	(45)
SamA	AAGCAGAAGACGGCATACGAAGACC	(26)
EB01 (*sspA*)	AACGGAGCTAAGTAACAATCCACT	This work
EB02 (*mukB*)	CACCTAGTGTATCAGCAAGACCTT	This work
EB03 (*ruvB*)	TCACTCGCTTTCAATTAACTCAGC	This work
EB06 (*brnQ*)	CAGCAACTTTCGCAGCATCTAATA	This work
EB08 (*nhaR*)	TTGTAAGTCCATATAGACGCCCAC	This work
EB10 (*hslO*)	AACATGGTTGATGGAGGTAAAGTT	This work
EB11 (*ubiH*)	CAAATTCAATCTTAGCAAAGCTGTC	This work

     FIGURES

Figure 1. Experimental setup for bacterial mutant enrichment in HARV vessels to model

microgravity (LSMMG) and to have the control gravity condition. INSeq libraries were

grown in HARV vessels that can simulate microgravity when placed perpendicular or gravity

when horizontal. Mutants exhibited either no defects in either condition, or exhibited (1) a growth

defect in both conditions, (2) LSMMG defect, (3) or a gravity-specific defect.

Figure 2. Pairwise correlation between samples revealed a clear difference between input

samples and output samples from the HARV conditions . Spearman correlations between

samples, in which the normalized transposon insertion counts in each gene were compared.

Input replicates were the original INSeq library grown for inoculation into the HARV, and low-

shear modeled microgravity (LSMMG) and normal gravity replicates were outputs from the

respective HARV-grown experiments.

Figure 3. Mutant behavior under LSMMG or gravity revealed fitness in each HARV

condition, normalized to the input transposon sequencing library. A) Comparison scatter

plot of microgravity (x-axis) and gravity (y-axis) Log(Fold-Change) values. Counts for each gene

for each replicate were normalized and used to calculate Log10(Fold-Change) values as

described in Materials and Methods. Genes that were poorly represented in the input pool did

not exhibit significant depletion, or were previously determined to be growth deficient in LBS

medium were designated as open black circles. Filled black circles are genes that were

significantly (p-value < 0.05) depleted 2-fold change under both simulated microgravity and

gravity conditions. Genes that are filled red were significant in both conditions but more depleted

under gravity, whereas open red circles are exclusively significant in gravity. Filled blue genes

are significant in both but more depleted under simulated microgravity. Remaining genes are

shown as open black circles.

Figure 4. Competition of defined mutants provided validation of mutant fitness from

within the complex INSeq library. Mutants were competed against wild-type *V. fischeri* under

simulated microgravity and gravity using the HARV vessels. Dots represent fold-change for 1:1

competition of mutant vs. marked parental strain (MJM1575).

Figure 5.Gene expression (RNA-Seq) does not predict conditional gene requirement

(INSeq) under LSMMG conditions. INSeq (y-axis) and RNA-Seq (x-axis) were compared by

using log-normalized fold-change values for different experimental conditions. A) INSeq

(LSMMG/Gravity) was compared to a previously published RNA-Seq (LSMMG/Gravity) under

HARV conditions and had an R^2 of 0.08. We also calculated fold-changes for RNA-Seq HARV

conditions using a dataset from SWT medium. B) INSeq (Gravity/Input) were compared to RNA-

Seq (Gravity/SWT) and had an R^2 of 0.18. C) INSeq (LSMMG/Input) was compared to RNA-

Seq (LSMMG/SWT) and had an R^2 0.11.

SUPPLEMENTAL DATA

The file Supplemental_Tables.xlsx includes the following tables:

Table S1: INSeq sample details

Table S2: Summary gene table output from pyinseq

Table S3: List of genes depleted as shown in Figure 3

ACKNOWLEDGMENTS

We acknowledge Cheryl Whistler and Randi Foxall for VFS Library mutants and A. Murat Eren

and his laboratory for technical advice and mentorship. Funding was provided by NASA's Illinois

Space Grant Consortium Research Seed Grant (M.J.M.), NIGMS R35 GM119627 (M.J.M.), an

American Society for Microbiology Undergraduate Research Fellowship (E.B.), and the McNair

Scholars Program (E.B.). The project was also supported in part by a NASA Space Biology

award 80NSSC18K1465 (J.S.F.).

REFERENCES

1. Foster JS, Khodadad CLM, Ahrendt SR, Parrish ML. 2013. Impact of simulated

microgravity on the normal developmental time line of an animal-bacteria symbiosis. Sci

Rep 3:srep01340.

2. Voorhies AA, Lorenzi HA. 2016. The Challenge of Maintaining a Healthy Microbiome during

Long-Duration Space Missions. Frontiers in Astronomy and Space Sciences 3:23.

3. Casaburi G, Goncharenko-Foster I, Duscher AA, Foster JS. 2017. Transcriptomic changes

in an animal-bacterial symbiosis under modeled microgravity conditions. Sci Rep 7:46318.

4. Castro-Wallace S, Stahl S, Voorhies A, Lorenzi H, Douglas GL. 2017. Response of

Lactobacillus acidophilus ATCC 4356 to low-shear modeled microgravity. Acta Astronaut

139:463–468.

5. Shao D, Yao L, Riaz MS, Zhu J, Shi J, Jin M, Huang Q, Yang H. 2017. Simulated

microgravity affects some biological characteristics of *Lactobacillus acidophilus*. Appl

Microbiol Biotechnol 101:3439–3449.

6. Duscher AA, Conesa A, Bishop M, Vroom MM, Zubizarreta SD, Foster JS. 2018.

Transcriptional profiling of the mutualistic bacterium *Vibrio fischeri* and an *hfq* mutant under

modeled microgravity. npj Microgravity 4:25.

7. Garrett-Bakelman FE, Darshi M, Green SJ, Gur RC, Lin L, Macias BR, McKenna MJ,

Meydan C, Mishra T, Nasrini J, Piening BD, Rizzardi LF, Sharma K, Siamwala JH, Taylor L,

Vitaterna MH, Afkarian M, Afshinnekoo E, Ahadi S, Ambati A, Arya M, Bezdan D, Callahan

CM, Chen S, Choi AMK, Chlipala GE, Contrepois K, Covington M, Crucian BE, De Vivo I,

Dinges DF, Ebert DJ, Feinberg JI, Gandara JA, George KA, Goutsias J, Grills GS, Hargens

AR, Heer M, Hillary RP, Hoofnagle AN, Hook VYH, Jenkinson G, Jiang P, Keshavarzian A,

Laurie SS, Lee-McMullen B, Lumpkins SB, MacKay M, Maienschein-Cline MG, Melnick

AM, Moore TM, Nakahira K, Patel HH, Pietrzyk R, Rao V, Saito R, Salins DN, Schilling JM,

Sears DD, Sheridan CK, Stenger MB, Tryggvadottir R, Urban AE, Vaisar T, Van Espen B,

Zhang J, Ziegler MG, Zwart SR, Charles JB, Kundrot CE, Scott GBI, Bailey SM, Basner M,

Feinberg AP, Lee SMC, Mason CE, Mignot E, Rana BK, Smith SM, Snyder MP, Turek FW.

2019. The NASA Twins Study: A multidimensional analysis of a year-long human

spaceflight. Science 364.

8.   Chang D, Zhu Y, An L, Liu J, Su L, Guo Y, Chen Z, Wang Y, Wang L, Wang J, Li T, Fang

X, Fang C, Yang R, Liu C. 2013. A multi-omic analysis of an *Enterococcus faecium* mutant

reveals specific genetic mutations and dramatic changes in mRNA and protein expression.

BMC Microbiol 13:304.

9.   Wilson JW, Ott CM, Quick L, Davis R, Höner zu Bentrup K, Crabbé A, Richter E, Sarker S,

Barrila J, Porwollik S, Cheng P, McClelland M, Tsaprailis G, Radabaugh T, Hunt A, Shah

M, Nelman-Gonzalez M, Hing S, Parra M, Dumars P, Norwood K, Bober R, Devich J,

Ruggles A, CdeBaca A, Narayan S, Benjamin J, Goulart C, Rupert M, Catella L, Schurr MJ,

Buchanan K, Morici L, McCracken J, Porter MD, Pierson DL, Smith SM, Mergeay M, Leys

N, Stefanyshyn-Piper HM, Gorie D, Nickerson CA. 2008. Media ion composition controls

regulatory and virulence response of *Salmonella* in spaceflight. PLoS One 3:e3923.

10. Nickerson CA, Ott CM, Wilson JW, Ramamurthy R, Pierson DL. 2004. Microbial responses

to microgravity and other low-shear environments. Microbiol Mol Biol Rev 68:345–361.

11. Schwarz RP, Goodwin TJ, Wolf DA. 1992. Cell culture for three-dimensional modeling in

rotating-wall vessels: an application of simulated microgravity. J Tissue Cult Methods

14:51–57.

12. Fang A, Pierson DL, Koenig DW, Mishra SK, Demain AL. 1997. Effect of simulated

microgravity and shear stress on microcin B17 production by *Escherichia coli* and on its

excretion into the medium. Appl Environ Microbiol 63:4090–4092.

13. Demain AL, Fang A. 2001. Secondary metabolism in simulated microgravity. Chem Rec

1:333–346.

14. Castro SL, Nelman-Gonzalez M, Nickerson CA, Mark Ott C. 2011. Induction of Attachment-

Independent Biofilm Formation and Repression of *hfq* Expression by Low-Fluid-Shear

Culture of *Staphylococcus aureus*. Appl Environ Microbiol 77:6368–6378.

15. Wang H, Yan Y, Rong D, Wang J, Wang H, Liu Z, Wang J, Yang R, Han Y. 2016.

Increased biofilm formation ability in *Klebsiella pneumoniae* after short-term exposure to a

simulated microgravity environment. Microbiologyopen 5:793–801.

16. Abshire CF, Prasai K, Soto I, Shi R, Concha M, Baddoo M, Flemington EK, Ennis DG, Scott

RS, Harrison L. 2016. Exposure of *Mycobacterium marinum* to low-shear modeled

microgravity: effect on growth, the transcriptome and survival under stress. NPJ

Microgravity 2:16038.

17. Mandel MJ. 2010. Models and approaches to dissect host-symbiont specificity. Trends

Microbiol 18:504–511.

18. Jones BW, Nishiguchi MK. 2004. Counterillumination in the Hawaiian bobtail squid,

Euprymna scolopes Berry (Mollusca: Cephalopoda). Mar Biol 144:1151–1155.

19. Koropatnick TA, Engle JT, Apicella MA, Stabb EV, Goldman WE, McFall-Ngai MJ. 2004.

Microbial factor-mediated development in a host-bacterial mutualism. Science 306:1186–

1188.

20. Mandel MJ, Wollenberg MS, Stabb EV, Visick KL, Ruby EG. 2009. A single regulatory gene

is sufficient to alter bacterial host range. Nature 458:215–218.

21. Mandel MJ, Schaefer AL, Brennan CA, Heath-Heckman EAC, Deloney-Marino CR, McFall-

Ngai MJ, Ruby EG. 2012. Squid-derived chitin oligosaccharides are a chemotactic signal

during colonization by *Vibrio fischeri*. Appl Environ Microbiol 78:4620–4626.

22. Brooks JF 2nd, Mandel MJ. 2016. The histidine kinase BinK Is a negative regulator of

biofilm formation and squid colonization. J Bacteriol 198:2596–2607.

23. Foster JS, Kerney KR, Parrish ML, Khodadad CLM, Ahrendt SR. 2011. Potential of the

Euprymna/Vibrio symbiosis as a model to assess the impact of microgravity on bacteria-

induced animal development. Gravitational and Space Research 25:44–47.

24. Brooks JF 2nd, Gyllborg MC, Cronin DC, Quillin SJ, Mallama CA, Foxall R, Whistler C,

Goodman AL, Mandel MJ. 2014. Global discovery of colonization determinants in the squid

symbiont *Vibrio fischeri*. Proc Natl Acad Sci U S A 111:17284–17289.

25. Goodman AL, McNulty NP, Zhao Y, Leip D, Mitra RD, Lozupone CA, Knight R, Gordon JI.

2009. Identifying genetic determinants needed to establish a human gut symbiont in its

habitat. Cell Host Microbe 6:279–289.

26. Goodman AL, Wu M, Gordon JI. 2011. Identifying microbial fitness determinants by

insertion sequencing using genome-wide transposon mutant libraries. Nat Protoc 6:1969–

1980.

27. Dunn AK, Millikan DS, Adin DM, Bose JL, Stabb EV. 2006. New *rfp*- and pES213-derived

tools for analyzing symbiotic *Vibrio fischeri* reveal patterns of infection and *lux* expression in

situ. Appl Environ Microbiol 72:802–810.

28. Thompson LR, Nikolakakis K, Pan S, Reed J, Knight R, Ruby EG. 2017. Transcriptional

characterization of *Vibrio fischeri* during colonization of juvenile *Euprymna scolopes*.

Environ Microbiol 19:1845–1856.

29. Meeske AJ, Riley EP, Robins WP, Uehara T, Mekalanos JJ, Kahne D, Walker S, Kruse AC,

Bernhardt TG, Rudner DZ. 2016. SEDS proteins are a widespread family of bacterial cell

wall polymerases. Nature 537:634–638.

30. Joyce AR, Reed JL, White A, Edwards R, Osterman A, Baba T, Mori H, Lesely SA, Palsson

BØ, Agarwalla S. 2006. Experimental and computational assessment of conditionally

essential genes in *Escherichia coli*. J Bacteriol 188:8259–8271.

31. Casutt MS, Schlosser A, Buckel W, Steuber J. 2012. The single NqrB and NqrC subunits in

the Na^+-translocating NADH: quinone oxidoreductase (Na^+-NQR) from *Vibrio cholerae* each

carry one covalently attached FMN. Biochim Biophys Acta 1817:1817–1822.

32. Steuber J, Halang P, Vorburger T, Steffen W, Vohl G, Fritz G. 2014. Central role of the Na^+-

translocating NADH:quinone oxidoreductase (Na^+-NQR) in sodium bioenergetics of *Vibrio*

cholerae. Biol Chem 395:1389–1399.

33. Aeckersberg F, Lupp C, Feliciano B, Ruby EG. 2001. *Vibrio fischeri* outer membrane

protein OmpU plays a role in normal symbiotic colonization. J Bacteriol 183:6590–6597.

34. Turner KH, Everett J, Trivedi U, Rumbaugh KP, Whiteley M. 2014. Requirements for

Pseudomonas aeruginosa acute burn and chronic surgical wound infection. PLoS Genet

10:e1004518.

35. Langmead B, Trapnell C, Pop M, Salzberg SL. 2009. Ultrafast and memory-efficient

alignment of short DNA sequences to the human genome. Genome Biol 10:R25.

36. Reback J, McKinney W, jbrockmendel, Van den Bossche J, Augspurger T, Cloud P,

gfyoung, Sinhrks, Klein A, Hawkins S, Roeschke M, Tratner J, She C, Ayd W, Petersen T,

MomIsBestFriend, Garcia M, Schendel J, Hayden A, Jancauskas V, Battiston P, Saxton D,

Seabold S, McMaster A, chris-b, h-vetinari, Hoyer S, Dong K, Overmeire W, Winkel M.

2020. pandas-dev/pandas: Pandas 1.1.0.

37. McKinney W, Others. 2010. Data structures for statistical computing in python, p. 51–56. *In*

Proceedings of the 9th Python in Science Conference. Austin, TX.

38. Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. Genome Biol 15:550.

39. Lyell NL, Dunn AK, Bose JL, Vescovi SL, Stabb EV. 2008. Effective Mutagenesis of *Vibrio*

fischeri by Using Hyperactive Mini-Tn*5* Derivatives. Appl Environ Microbiol 74:7059–7063.

40. Pollack-Berti A, Wollenberg MS, Ruby EG. 2010. Natural transformation of *Vibrio fischeri*

requires *tfoX* and *tfoY*. Environ Microbiol 12:2302–2311.

41. Boettcher KJ, Ruby EG. 1990. Depressed light emission by symbiotic *Vibrio fischeri* of the

sepiolid squid *Euprymna scolopes*. J Bacteriol 172:3701–3706.

42. Mandel MJ, Stabb EV, Ruby EG. 2008. Comparative genomics-based investigation of

resequencing targets in *Vibrio fischeri*: focus on point miscalls and artefactual expansions.

BMC Genomics 9:138.

43. Brooks JF 2nd, Gyllborg MC, Kocher AA, Markey LEH, Mandel MJ. 2015. TfoX-based

genetic mapping identifies *Vibrio fischeri* strain-level differences and reveals a common

lineage of laboratory strains. J Bacteriol 197:1065–1074.

44. Post DMB, Yu L, Krasity BC, Choudhury B, Mandel MJ, Brennan CA, Ruby EG, McFall-

Ngai MJ, Gibson BW, Apicella MA. 2012. O-antigen and core carbohydrate of *Vibrio fischeri*

lipopolysaccharide: composition and analysis of their role in *Euprymna scolopes* light organ

colonization. J Biol Chem 287:8515–8530.

45. Studer SV, Mandel MJ, Ruby EG. 2008. AinS quorum sensing regulates the *Vibrio fischeri*

acetate switch. J Bacteriol 190:5915–5923.

INPUT POOL

LSMMG

Gravity