In plain sight: the role of alpha-1-antitrypsin in COVID-19 pathogenesis and

therapeutics.

Kasopefoluwa Y Oguntuyo[1#], Christian S Stevens[1#], Mohammed NA Siddiquey [2],

Robert M Schilke [2], Matthew D Woolard [2], Hongbo Zhang [2], Joshua A Acklin [1], Satoshi

Ikegame[1], Chuan-Tien Huang[1], Jean K Lim[1], Robert W Cross[3], Thomas W Geisbert[3],

Stanimir S Ivanov [2], Jeremy P Kamil[2], and Benhur Lee[1]

Affiliations:
1. Department of Microbiology, Icahn School of Medicine at Mount Sinai, New York,
NY 10029
2. Department of Microbiology and Immunology, Louisiana State University Health
Shreveport, Shreveport, LA 71103
3. Department of Microbiology and Immunology, University of Texas Medical
Branch at Galveston, Galveston, TX 77555
16

[#]These authors contributed equally to this work.

Author contributions:

KYO, CSS and BL conceived and designed the study. KYO, CSS, MNAS, RMS, MDW,

HZ, and JAA collected data. SI, CTH, JKL, RWC, TWG, SSI, and JPK contributed

valuable reagents, data and/or tools. KYO and CSS analyzed the data and wrote the

original drafts of the paper. BL reviewed the draft, supported data analysis, and

provided invaluable direction throughout the conceptualization and execution of the

project. All authors had the opportunity to review the manuscript prior to submission and

JAA, SI, JKL, RWC, TWG, SSI, JPK provided valuable feedback during the editing

process.

ABSTRACT

Entry of SARS-CoV-2 is facilitated by endogenous and exogenous proteases. These

proteases proteolytically activate the SARS-CoV-2 spike glycoprotein and are key

modulators of virus tropism. We show that SARS-CoV-2 naïve serum exhibits significant

inhibition of SARS-CoV-2 entry. We identify alpha-1-antitrypsin (AAT) as the major

serum protease inhibitor that potently restrict protease-mediated entry of SARS-CoV-2.

AAT inhibition of protease-mediated SARS-CoV-2 entry in vitro occurs at concentrations

far below what is present in serum and bronchoalveolar tissues, suggesting that AAT

effects are physiologically relevant. Moreover, AAT deficiency affects up to 20% of the

population and its symptomatic manifestations coincides with many risk factors

associated with severe COVID-19 disease. In addition to the effects that AAT may have

on viral entry itself, we argue that the anti-inflammatory and coagulation regulatory

activity of AAT have implications for coronavirus disease 2019 (COVID-19)

pathogenicity, SARS-CoV-2 tissue restriction, convalescent plasma therapies, and even

potentially AAT therapy.

KEYWORDS: COVID-19, SARS-CoV-2, Alpha-1-antitrypsin, SERPINA1, alpha-2-

macroglobulin, TMPRSS2, protease, convalescent plasma

The COVID-19 pandemic that began in December 2019 has resulted in tens of millions of cases with hundreds of thousands of deaths. The central conundrum of this pandemic is the heterogeneity of disease severity in SARS-CoV-2 infected individuals. The widespread disparities in outcomes has spurred global efforts to better understand SARS-CoV-2 pathogenesis, investigate the factors contributing to the clinical course of COVID-19, and develop viable therapeutics. One of the most promising therapeutic targets is the Spike (S) glycoprotein of SARS-CoV-2, which bears the fusion machinery necessary to mediate viral entry. Vaccines, monoclonal antibodies and convalescent plasma therapy are all premised upon neutralizing SARS-CoV-2 spike (CoV2-S) mediated entry. All three modalities are being developed at unprecedented speed. Unfortunately, the lack of standardized virus neutralization assays (VNAs) or reporting metrics have made it difficult to compare the efficacy across the proliferating number of vaccine platforms and CoV2-S targeted treatment modalities.

Standardizing a SARS-CoV-2 Viral Neutralization Assay

In an accompanying study, we worked to develop a scalable, standardized VNA that reflects the complex interplay between CoV2-S receptor interaction and proteolytic activation. To that end we generated and validated VSVΔG pseudotyped particles bearing SARS-CoV-2 spike (CoV2pp).[1] We initially optimized infection conditions in serum free media. Due to the role of proteolytic activation in SARS-CoV-2 entry, we utilized exogenous trypsin as well as soybean trypsin inhibitor to maximize entry of the CoV2pp while limiting cytotoxicity. These conditions were designed noting that proteolytic activation of CoV2-S is required for the receptor-induced conformational

changes that result in virus-cell membrane fusion and viral entry. Several endosomal,

cell surface, and exogenous proteases, including furin, select cathepsins and

TMPRSSs, and trypsin-like proteases have been implicated in mediating these

cleavage events for SARS-CoV-2 and enhancing cellular entry (Fig. 1A).[2–4] A similar

body of evidence also indicates that these proteases and others, such as elastase, play

critical roles in the productive processing of the S protein from SARS-CoV-1, MERS-

CoV, and other betacoronaviruses.[5–7] In the case of MERS-CoV, proteolytic processing

of spike is capable of dictating cell tropism and correlates well with virulence.[8] For

SARS-CoV-2, the role of cell surface protease-mediated entry is of sufficient importance

that TMPRSS2 overexpression can render some cell lines refractory to chloroquine

mediated inhibition of virus entry.[9]

To standardize our VNA, we initially utilized trypsin-treated CoV2pp for human serum

neutralization experiments. However, we observed that under these conditions, sera

from patients not exposed to SARS-CoV-2 was capable of neutralizing these

pseudoviruses, though, as expected, to a lesser extent than sera from patients that

tested positive for SARS-CoV-2 antibodies (Fig. 1B)[1].

Uncovering AAT: a heat-labile, CoV2pp neutralizing factor in SARS-CoV-2 naïve

serum

To pre-empt the variable neutralizing effect of seronegative serum, we diluted our

trypsin-treated CoV2pp in DMEM containing 10% fetal bovine serum (FBS). However,

while this provided an easy solution in standardizing our assay for "out-of-the-box" use

when sent to multiple labs, it left the question of "why" unanswered. This neutralizing

effect of SARS-CoV-2 naïve human sera was ameliorated through heat-inactivation by

incubation at 56⁰C for 1hr (Fig. 1B), and also appeared to be specific to CoV2-S

mediated entry as VSV-G pseudotyped particles (VSV-Gpp) were unaffected. Thus, the

serum neutralizing factor(s) was unlikely to be cross-reactive antibodies to seasonal

coronaviruses or complement *per se*. We therefore suspected a heat-labile serum

factor(s) capable of inhibiting trypsin.[10]

To validate these observations, we tested a panel of non-heat-inactivated human sera

for neutralization activity against CoV2pp and VSV-Gpp. SARS-CoV-2 seropositive and

seronegative sera were diluted in SFM (Supplemental Fig. 1A). Here, we observed

neutralization of trypsin-treated CoV2pp by both seronegative and seropositive sera

(Fig. 2A and Supplemental Fig. 1B). Importantly, VSV-Gpp was not inhibited by the

samples tested (Fig. 2B and Supplemental Fig. 1C), once again showing that this effect

was CoV2pp-specific. At the same time, an external group at Louisiana State University

Health Shreveport (LSUHS) independently observed neutralization by seronegative

sera under similar experimental conditions using our CoV2pp (Fig. 2C, Supplemental

Fig. 1D and 1E). Remarkably, in both groups, SARS-CoV-2 naïve sera inhibited

CoV2pp entry by 90-97% (Fig. 2A and 2C, left panels). Nonetheless, seropositive

patient sera showed inhibition orders of magnitude beyond this threshold, suggesting

antibody mediated inhibition of CoV2pp entry (Fig. 2A and C, right panels). Additionally,

using the identical serum samples in Fig. 2C, collaborators at the University of Texas

Medical Branch at Galveston (UTMB) observed modest, but significant, neutralization of

live virus by seronegative sera as assayed by a plaque reduction neutralization assay

(Fig. 2D and Supplemental Fig. 1F).

Upon making these repeated observations, we searched the literature for highly

abundant and heat labile products in serum and identified alpha-1-antitrypsin (AAT) and

alpha-2-macroglobulin (A2M) as potential candidates.[10] These blood products are

typically present in human serum at high concentrations (1.1-2.2 mg/mL for AAT and 2-

4 mg/mL for A2M) and have been described to inhibit both exogenous and endogenous

proteases.[11–13] Despite the careful characterizations of the role endogenous and

exogenous proteases play in SARS-CoV-2 entry, there have been limited

characterizations of the role *in vivo* protease inhibitors play in modulating SARS-CoV-2

entry. A2M and AAT alone are responsible for approximately 10% and 90% of serum

antiprotease capacity, respectively.[14]

A2M functions to inhibit a broad range of proteases, such as serine and cysteine

proteases. In addition to protease inhibitory functions, A2M also inhibits thrombin to

prevent coagulation and binds to growth factors and cytokines. No clinical conditions

have yet been associated with low plasma levels of A2M.[11] On the other hand, AAT is a

protease inhibitor that irreversibly binds serine proteases and plays additional roles in

the regulation of inflammation and coagulation.[15] Notably, decreased plasma

concentrations of or function of AAT have been associated with liver and lung disease,

particularly pulmonary emphysema due to unregulated neutrophil elastase activity.[12]

Mutations leading to these conditions are highly prevalent as nearly 20% of individuals

have non-wildtype AAT alleles.[13]

To assess whether AAT and/or A2M alone could inhibit trypsin-treated CoV2pp entry,

we added each at the time of infection and observed potent entry inhibition by AAT and

modest inhibition by A2M, with IC50s of 14.47μg/mL and 54.20μg/mL, respectively (Fig.

3A, left panel). Importantly, neither protein inhibited VSV-Gpp (Fig. 3A, right panel).

Albumin, the most abundant protein in blood, showed no significant reduction of entry of

either CoV2pp or VSV-Gpp (Fig. 3A), which underscores that the inhibitory effects of

AAT and A2M on CoV2-S mediated entry was specific.

While these findings suggest that AAT, and to a lesser extent A2M, can inhibit

exogenous trypsin-like proteases known to enhance SARS-CoV-2 entry, tissue

restriction of SARS-CoV-2 infection is also mediated by proteases at the cell surface.[2,3]

Therefore, we sought to investigate whether either protein could inhibit TMPRSS2, an

endogenous serine protease implicated in SARS-CoV-2 pathogenicity. We previously

engineered two ultra-permissive 293T clones stably expressing ACE2 (clone 5-7) or

ACE2+TMPRSS2 (clone F8-2). Each of these lines was capable of highly efficient

CoV2pp entry in the absence of trypsin pre-treatment. To assess entry inhibition by AAT

and A2M, we performed VNAs in both clones using CoV2pp without any trypsin pre-

treatment but diluted in standard media (DMEM+10% FBS). Here, we observed that

AAT inhibited CoV2pp entry into TMPRSS2 expressing F8-2 clones, but not the 5-7

clones (Fig. 3B). For both cell lines, A2M and albumin both displayed no entry inhibition

at the concentrations tested. As additional controls, we observe potent inhibition of

CoV2pp by sRBD in both cell lines and, as expected, Nafamostat mesylate, a serine

protease inhibitor, inhibited entry in only the TMPRSS2-expressing F8-2 clones

(Supplemental Fig. 2). Notably, AAT and Nafamostat inhibition of CoV2pp entry into

TMPRSS2-expressing F8-2 cells approached a maximal inhibition of ~80%, suggesting

that SARS-CoV-2 can enter via other pathways noted in Fig. 1A.

Together, these observations show that not only can SARS-CoV-2 naïve sera potently

inhibit protease-mediated entry of SARS-CoV-2, but AAT and A2M appear to be

potential regulators of protease-mediated entry by SARS-CoV-2. In concert with the

large body of literature about AAT and the proteolytic processing of coronavirus spike

proteins, we are led to three distinct, but important hypotheses. The first is that AAT and

A2M may play a biologically relevant role in tissue restriction in SARS-CoV-2 infection.

The second is that, if the first hypothesis is true, variant AAT genotypes could influence

the relative severity of COVID-19. And thirdly, treatment with the already FDA-approved

AAT, could play a role in simultaneously controlling viral burden as well as aberrant

immune responses.

The suspected roles of protease inhibitors as uncharacterized players in COVID-

19 pathogenesis and therapy.

In an accompanying paper, we reported that a factor, or factors, in SARS-CoV-2

seronegative sera is capable of inhibiting trypsin-mediated entry of CoV2pp. Here, we

pinpoint AAT and A2M as highly abundant serum factors that inhibit the effect of

exogenous proteases on CoV2pp entry. Moreover, at the concentrations tested, we

show that AAT, but not A2M, can inhibit TMPRSS2-mediated entry of CoV2pp at sub-

physiologic concentrations. Two recent pre-peer review reports support these findings.

Azouz et al utilized a fluorescent cleavage assay in a screen that identified four small

molecules and AAT as inhibitors of TMPRSS2 enzymatic activity.[16] Wettstein et al

report neutralization of SARS-CoV-2 by polypeptides extracted from pooled

bronchoalveolar lavage (BAL) fluids and, after fractionation and mass spectrometry,

identify AAT as a component driving this activity.[17] Although previously reported, the

identification of AAT in the BAL confirms that it can diffuse into lung tissues and

suggests that it can be present at the site of SAR-CoV-2 infection and replication. They

also show that while the fraction containing AAT inhibits >99% of infection, there is

another set of fractions that inhibit just under 90% of infection. We speculate that these

fractions may contain A2M which we show to be a less potent inhibitor of trypsin-

mediated CoV2pp entry, perhaps due to the requirement of a tetramer to trap two

proteases (Fig. 3A).[11] Interestingly, this lack of potency may be offset by its broader

protease inhibition potential, particularly for cysteine proteases such as Cathepsin B

and L, which have been reported to play a role in endosomal mediated SARS-CoV-2

entry. Additionally, elastase—a serine protease released by neutrophils—has been

previously reported to play a role in enhancing SARS-CoV-1 and MERS entry.[7]

Although the role of elastase in SARS-CoV-2 entry has not been elucidated, elevation of

neutrophil counts in BAL and serum have been consistently associated with severe

COVID-19 cases.[18–20] In spite of its name, AAT has a stronger binding affinity to

elastase than trypsin, and this is borne out by the clinically significant sequalae

associated with AAT deficiency.[15]

Though the findings presented here focus on protease inhibitors' ability to inhibit

protease-mediated SARS-CoV-2 entry, the inhibitors play significant additional roles.

While others have speculated that neutrophil elastase should be considered as a target

for potential COVID-19 prophylactics, AAT was not noted specifically.[21] AAT, as an

acute phase protein, has been characterized to play roles in modulating inflammation by

inhibiting elastase among other factors. Elastase is critical for the formation of neutrophil

extracellular traps (NETs) in acute pneumonia, which can amplify inflammatory

responses if not resolved by AAT. Runaway pulmonary inflammation and NETosis is an

emerging theme in COVID-19 pathogenesis.[22] AAT is also known to modulate activities

that result in downstream IL-6 inhibition, which is heavily implicated in COVID-19

pathogenicity.[12,23] Seeking to capitalize on these anti-inflammatory roles and the already

well established use of recombinant AAT to treat AAT deficiency, McElvaney et al

recently reported the initiation of a clinical trial for AAT treatment of COVID-19 based on

their published work.[24] Moreover, AAT has regulatory roles in the coagulation cascade[25]

and, via elastase inhibition, could inhibit NET-triggered immunothromboses.[26] Notably,

inflammatory dysregulation and coagulopathies have been reported to play a role in the

disparate COVID-19 severities between patients.[27]

The potent neutralization of protease-mediated cellular entry by SARS-CoV-2 along with

the wide range and prevalence of functionally different AAT genotypes implies a

potential role for AAT in variable COVID-19 severity. Of particular note, nearly 20% of

individuals are either heterozygous or homozygous for non-wildtype alleles, many of

which have been described to result in reduced levels of AAT in the blood or reduced

AAT function.[13] Although many mutations have been characterized, new mutations that

impair abundance or function are still being identified. For example, AAT variants such

as G373R or the R223C (F allele) have been identified and reported to have wildtype

levels in the blood, but impaired inhibitory activity against neutrophil elastase.[28,29] We

speculate that this CoV2pp VNA assay may be capable of serving as a scalable means

by which one could screen for deficient AAT functionality and not simply abundance.

Considering that widespread screening of AAT is rarely performed in the absence of

emphysema, it is reasonable to expect that there may be more unidentified mutations

that impair abundance or function of AAT, which may subsequently result in aberrant

response to SARS-CoV-2 infection. This undesired response by individuals with

functional AAT deficits may enable effective viral entry, dysregulated inflammation,

and/or coagulopathies (Fig. 4). It also raises the possibility that AAT may represent a

novel therapeutic approach in the fight against SARS-CoV-2.[25]

However, in the case of convalescent plasma therapies, the presence of AAT in blood

plasma from donors may play a beneficial role as non-immunoglobulin products are

currently not excluded from transfused plasma. In line with this, a recent meta-analysis

from Joyner et al suggests that convalescent plasma therapy is beneficial to its

recipients.[30] While the purported benefits are attributed to the presence of neutralizing

antibodies, the authors also acknowledge that "other biological mechanisms" may

contribute to these observations. Additionally, a recent bioRxiv report shows modest

benefits from the transfer of standard IVIG, suggesting that neutralizing antibodies are

not the only serum components that may play a role in alleviating the burden of COVID-

19 in patients.[31] In its myriad roles, the transfer of AAT may provide additional benefits

to convalescent plasma recipients by inhibiting SARS-CoV-2 entry, restraining

inflammation and/or moderating coagulation.

In sum, these findings highlight the importance of protease inhibitors in restricting

exogenous and/or endogenous protease-mediated enhancement of SARS-CoV-2 entry.

We also speculate that the diversity of AAT genotypes, the complex regulation of its

activity, and its myriad roles in inflammation and coagulation, implicates functional AAT

levels in COVID-19 pathogenicity. There is an urgent need to address:

1) The biologically relevant roles AAT, A2M, or other proteases play in tissue

265   restriction in SARS-CoV-2 infection.

266   2) The impact of variant AAT genotypes on the relative severity of COVID-19.

267   3) Whether treatment with, the already FDA-approved, AAT, may be able to play

268   a role in simultaneously controlling viral burden as well as aberrant immune

269   responses and if simultaneous treatment with A2M should be considered

270   given their ability to form inhibitor complexes.[32]

METHODS

<u>Maintenance and generation of isogenic cell lines:</u>

Vero-CCL81, parental 293T, and isogenic 293T cells were cultured in DMEM with 10% heat inactivated FBS at 37ºC in the presence of 5% CO2. Isogenic 293T cell clones 5-7 and F8-2 were generated by lentivirus transduction to stably express ACE2 only or ACE2 and TMPRSS2, respectively. ACE2 expression was under puromycin selection and TMPRSS2 was under blasticidin selection as previously described.

<u>Production of VSV∆G pseudotyped particles and neutralization studies:</u>

Detailed protocols for the production and use of standardized VSVpp (CoV2pp and VSV-Gpp) are given in Oguntuyo and Stevens et al.[1] Briefly, 293T producer cells were transfected to express the viral surface glycoprotein of interest, infected with VSV∆G-rLuc-G* reporter virus, then virus supernatant collected and clarified 2 days post infection prior to use. Trypsin treated CoV2pp were treated as previously described.[1] All pseudotyped viruses (PsV) were aliquoted prior to storage at -80ºC and tittered prior to usage for neutralization experiments. Neutralization experiments were performed by diluting the appropriate PsV with a 4-fold serial dilution of Albumin (Sigma-Aldrich, A8763), alpha-1-antitrypsin (Sigma-Aldrich, SRP6312), alpha-2-Macroglobulin (Sigma-Aldrich, SRP6314) or Nafamostat mesylate (Selleckchem, S1386). SARS-CoV-2 soluble RBD (sRBD) with human IgG-Fc was produced by the Lee Lab using a recombinant Sendai virus expression platform (manuscript in preparation). De-identified patient sera were obtained via institutional biobanks that allowed use for research

purposes. All infections were processed for detection of Renilla luciferase activity at

20hrs post-infection, and luminescence was read on the Cytation3 (BioTek).

<u>Plaque reduction neutralization titration (PRNT) by sera of SARS-CoV-2:</u>

Neutralization experiments with live virus were performed by incubating sera with 50-

100 PFU of SARS-CoV-2 for one hour at 37ºC. All sera were diluted in serum free

DMEM. Serial dilutions started at a four-fold dilution and went through seven three-fold

serial dilutions. The virus-serum mixture was then used to inoculate Vero E6 cells for

one hour at 37ºC and 5% CO2. Cells were overlaid with EMEM medium (no FBS) and

1.25% Avicel, incubated for 3 days, and plaques were counted after staining with 1%

crystal violet in formalin.

ACKNOWLEDGMENTS

The authors acknowledge the following funding: KYO and CS were supported by Viral-

Host Pathogenesis Training Grant T32 AI07647; KYO was additionally supported

by F31 AI154739. BL acknowledges flexible funding support from NIH grants R01

AI123449, R21 AI1498033, and the Department of Microbiology and the Ward-Coleman

estate for endowing the Ward-Coleman Chairs at the ISMMS. JPK and SSI

acknowledge funding from a LSUHS COVID-19 intramural grant. JPK and SSI

acknowledge additional funding from NIH grants AI116851 and AI143839, respectively.

BL wishes to dedicate this paper to Ernest L Robles-Levroney, the first graduate student

BL had the privilege to train. Ernie Robles-Levroney was dedicated teacher, role model

and trailblazer who passed away unexpectedly during the course of writing this

manuscript.

 FIGURES / FIGURE LEGENDS

Figure 1. Overview of SARS-CoV-2 entry and inhibition of trypsin treated CoV2pp

entry by COVID seronegative sera. (A) Overview of SARS-CoV-2 entry. Three modes

of entry are displayed: (1) Entry mediated by endosomal proteases, such as Cathepsin

B, (2) entry mediated by cell surface proteases, such as TMPRSS2, and (3) entry

mediated by exogenous proteases, such as trypsin. This model was created in

Biorender. **(B)** Inhibition of trypsin treated CoV2pp entry by COVID seronegative sera.

Presented is a schematized version of the results presented in supplemental Figure 3A

of our previous publication. For this experiment, sera samples were incubated with

trypsin treated CoV2pp for 30mins prior to addition to Vero-CCL81 cells. Both the

CoV2pp and the sera samples were diluted in serum free media. The grey lines

represent COVID seronegative sera, they purple lines are COVID seropositive sera, and

the dashed lines are samples that were heat inactivated (HI) for 1hr at 56ºC prior to use

for CoV2pp neutralizations.

Figure 2. Negative patient sera inhibit exogenous protease mediated

enhancement of CoV2pp. (A) SARS-CoV-2 seronegative sera inhibit trypsin treated

CoV2pp. Seronegative or seropositive samples were first identified based on IgG

antibodies against Spike (Supplemental Fig. 1). The indicated sera diluted in serum-free

DMEM were incubated with a pre-titered amount of CoV2pp prior to spinoculation on

Vero-CCL81 cells as described.[1] Sera were not heat inactivated before use in our

neutralization assays. Normalized infection data at the highest and lowest dilutions

tested are shown as % maximal infection (media only) with results from seronegative

(left) and seropositive (right) sera plotted on linear and log scale, respectively. Data

points represent the mean of neutralizations performed in triplicate. Dilutions shown

were compared using a paired t test (ns, not significant; **, $p < 0.01$; ***, $p < 0.005$, and

****, $p < 0.0001$). Full neutralization curves are shown in Supplemental Fig. 1. **(B)**

SARS-CoV-2 seronegative sera do not inhibit VSV-Gpp. Experiment performed and

presented as in Fig. 2A. **(C)** Inhibition of trypsin treated CoV2pp entry by SARS-CoV-2

seronegative sera independently observed. Collaborators in a different state

independently performed the identical experiment described in Fig. 1A with their own

cohort of seropositive and seronegative samples. Data shown are means from technical

quadruplicates/sample/dilution and presented exactly as for Fig. 2A. **(D)** Live SARS-

CoV-2 is modestly inhibited by seronegative sera. Sera samples presented in Fig. 2C

were utilized for plaque reduction neutralization experiments (PRNT) with live virus (left

panel) as described in the materials and methods. Presented here are the mean of one

experiment done in technical duplicates and error bars show SEM. Data presented as in

Fig. 2A.

Figure 3. Alpha-1-antitrypsin (AAT) and alpha-2-macroglobulin (A2M) inhibit

protease mediated enhancement of CoV2pp entry. (A) AAT and A2M inhibit trypsin-

mediated enhancement of CoV2pp entry. Trypsin treated CoV2pp (left panel) and

standard VSV-Gpp (right) were diluted in serum free media, then used to infect Vero-

CCL81 cells in the presence of the indicated concentrations of albumin, AAT, or A2M.

Data are from two independent experiments and are presented as percent relative

infection where each concentration was normalized to the lowest concentration of the

test reagent used. Data fit as described in Fig. 1A. **(B)** AAT inhibits TMPRSS2-mediated

enhancement of CoV2pp entry. CoV2pp not treated with trypsin were diluted in

DMEM+10% FBS and utilized to infect 293T-ACE2+TMPRSS2 clone F8-2 (left panel) or

293T-ACE2 clone (5-7) in the presence of the indicated concentrations of A2M, AAT, or

Albumin. Data points are means +/- SEM a representative experiment performed in

triplicates, but otherwise presented as described as in Fig. 3A.

Figure 4. Putative relationship between alpha-1-antitrypsin function and SARS-

CoV-2. Differential alpha-1-antitrypsin (AAT) abundance and/or function may result in a

differential response to infection by SARS-CoV-2. Due to our observations that AAT can

inhibit CoV2pp protease-mediated entry, we expect that in the presence of functional

AAT (light blue, right) there is only modest amounts of SARS-CoV-2 protease-mediated

entry relative to those with AAT functional deficiencies (orange, left). In addition to its

entry effects, we speculate that normal AAT abundance and/or function may reign in an

otherwise dysregulated immune response. Moreover, AAT's role in regulation of the

coagulation cascade may further prevent the development of coagulopathies. For the

latter, a normal immune response to infection is indicated by a single up arrow and the

absence of coagulopathy is indicated by a dash. Figure generated in Biorender.

Supplemental Figure 1. Spike ELISA data and full neutralization curves. (A) Spike

ectodomain ELISAs for JBA and JBB samples. Our seronegative and seropositive

samples were utilized. ELISAs performed as previously described[(ref)] and shown are the

OD490 values from the 1:100 sera dilution. Shown are the median and interquartile

range. **(B)** Full neutralization curves from trypsin treated CoV2pp. Data points are mean

+/- SEM from experiment done in triplicates. **(C)** Full neutralization curves from VSV-

Gpp. Presented are the means of an experiment in technical triplicate with error bars

showing SEM. **(D)** Spike ectodomain ELISAs from LSA and LSB samples. Twelve

seropositive and twelve seronegative samples were utilized. Shown are the OD450

values from the 1:100 sera dilution. **(E)** Full neutralization curves from LSU CoV2pp

neutralization are shown here. **(F)** Live virus full neutralization curves. Live virus

neutralizations performed as described in the Methods and the same samples as in

Supplemental Fig. 1E were used. Presented here are the means of one experiment

done in technical duplicate and error bars show SEM and data were fit using variable

slope, 4-parameter logistics regression curve (robust fitting method).

Supplemental Figure 2. Nafamostat mesylate inhibits CoV2pp entry into

TMPRSS2 expressing cells. CoV2pp were mixed with a serial dilution of either

Nafamostat or sRBD prior to infection of isogenic cells stably expressing

ACE2+TMPRSS2 (clone F8, left panel) or ACE2 (clone 5-7, right panel). Presented

here are the results of an experiment done in technical triplicates. Error bars show SEM

and data were fit using variable slope, 4-parameter logistics regression curve (robust

fitting method).

REFERENCES

1. Oguntuyo, K. Y. and Stevens, C. *et al.* Quantifying absolute neutralization titers against SARS-CoV-2 by a standardized virus neutralization assay allows for cross-cohort comparisons of COVID-19 sera. *medRxiv* (2020).

2. Hoffmann, M. *et al.* SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* **181**, 271-280.e8 (2020).

3. Zhu, F. C. *et al.* Proteolytic activation of the SARS-coronavirus spike protein: Cutting enzymes at the cutting edge of antiviral research. *J. Virol.* **395**, 379–397 (2020).

4. Shang, J. *et al.* Cell entry mechanisms of SARS-CoV-2. *Proc. Natl. Acad. Sci.* **117**, 11727–11734 (2020).

5. Shulla, A. *et al.* A Transmembrane Serine Protease Is Linked to the Severe Acute Respiratory Syndrome Coronavirus Receptor and Activates Virus Entry. *J. Virol.* **85**, 873–882 (2011).

6. Menachery, V. D. *et al.* Trypsin Treatment Unlocks Barrier for Zoonotic Bat Coronavirus Infection. *J. Virol.* **94**, (2019).

7. Belouzard, S., Madu, I. & Whittaker, G. R. Elastase-mediated Activation of the Severe Acute Respiratory Syndrome Coronavirus Spike Protein at Discrete Sites within the S2 Domain. *J. Biol. Chem.* **285**, 22758–22763 (2010).

8. Park, J. E. *et al.* Proteolytic processing of Middle East respiratory syndrome coronavirus spikes expands virus tropism. *Proc. Natl. Acad. Sci.* **113**, 12262–

12267 (2016).

9.   Hoffmann, M. *et al.* Chloroquine does not inhibit infection of human lung cells with

SARS-CoV-2. *Nature* (2020). doi:10.1038/s41586-020-2575-3

10. Sriram Vemuri, C. Tony Yu, and N. R. Formulation and Stability of Recombinant

Alpha-1-Antitrypsin. in *Stability and Characterization of Protein and Peptide Drugs*

(eds. Wang, Y. J. & Pearlman, R.) **5**, 263–285 (Springer US, 1993).

11. Rehman, A. A., Ahsan, H. & Khan, F. H. Alpha-2-macroglobulin: A physiological

guardian. *J. Cell. Physiol.* **228**, 1665–1675 (2013).

12. Blanco, I. *Alpha-1 Antitrypsin Biology. Blanco's Overview of Alpha-1 Antitrypsin*

Deficiency **1**, (Elsevier Inc., 2017).

13. Bornhorst, J. A., Greene, D. N., Ashwood, E. R. & Grenache, D. G. α1-Antitrypsin

phenotypes and associated serum protein concentrations in a large clinical

population. *Chest* **143**, 1000–1008 (2013).

14. de Serres, F. & Blanco, I. Role of alpha-1 antitrypsin in human health and

disease. *J. Intern. Med.* **276**, 311–335 (2014).

15. Strnad, P., McElvaney, N. G. & Lomas, D. A. Alpha 1 -Antitrypsin Deficiency. *N.*

Engl. J. Med. **382**, 1443–1455 (2020).

16. Azouz, N. P., Klingler, A. M. & Rothenberg, M. E. Alpha 1 Antitrypsin is an

Inhibitor of the SARS-CoV2–Priming Protease TMPRSS2. *bioRxiv* (2020).

doi:10.1101/2020.05.04.077826

17. Wettstein, L. *et al.* Alpha-1 antitrypsin inhibits SARS-CoV-2 infection. *bioRxiv*

(2020). doi:10.1101/2020.07.02.183764

18. Zhou, Z. *et al.* Heightened Innate Immune Responses in the Respiratory Tract of

COVID-19 Patients. *Cell Host Microbe* **27**, 883-890.e2 (2020).

19.   Lucas, C. *et al.* Longitudinal analyses reveal immunological misfiring in severe

COVID-19. *Nature* (2020). doi:10.1038/s41586-020-2588-y

20.   Liao, M. *et al.* Single-cell landscape of bronchoalveolar immune cells in patients

with COVID-19. *Nat. Med.* **26**, 842–844 (2020).

21.   Mohamed, M. M. A., El-Shimy, I. A. & Hadi, M. A. Neutrophil Elastase Inhibitors: A

potential prophylactic treatment option for SARS-CoV-2-induced respiratory

complications? *Crit. Care* **24**, 311 (2020).

22.   Barnes, B. J. *et al.* Targeting potential drivers of COVID-19: Neutrophil

extracellular traps. *J. Exp. Med.* **217**, (2020).

23.   Grifoni, E. *et al.* Interleukin-6 as prognosticator in patients with COVID-19. *J.*

Infect. (2020). doi:10.1016/j.jinf.2020.06.008

24.   McElvaney, O. J. *et al.* Characterization of the Inflammatory Response to Severe

COVID-19 Illness. *Am. J. Respir. Crit. Care Med.* (2020).

doi:10.1164/rccm.202005-1583OC

25.   Janciauskiene, S. & Welte, T. Well-Known and Less Well-Known Functions of

Alpha-1 Antitrypsin. Its Role in Chronic Obstructive Pulmonary Disease and Other

Disease Developments. *Ann. Am. Thorac. Soc.* **13**, S280–S288 (2016).

26.   Middleton, E. A. *et al.* Neutrophil Extracellular Traps (NETs) Contribute to

Immunothrombosis in COVID-19 Acute Respiratory Distress Syndrome. *Blood*

(2020). doi:10.1182/blood.2020007008

27.   Jose, R. J. & Manuel, A. COVID-19 cytokine storm: the interplay between

inflammation and coagulation. *Lancet Respir. Med.* **8**, e46–e47 (2020).

28. Cook, L., Burdon, J. G. W., Brenton, S., Knight, K. R. & Janus, E. D. Kinetic

characterisation of alpha-1-antitrypsin F as an inhibitor of human neutrophil

elastase. *Pathology* **28**, 242–247 (1996).

29. Laffranchi, M. *et al.* Characterisation of a type II functionally-deficient variant of

alpha-1-antitrypsin discovered in the general population. *PLoS One* **14**, e0206955

(2019).

30. Joyner, M. J. *et al.* Early Safety Indicators of COVID-19 Convalescent Plasma in

5,000 Patients. *medRxiv* (2020). doi:10.1101/2020.05.12.20099879

31. Sakoulas, G. *et al.* Intravenous Immunoglobulin (IVIG) Significantly Reduces

Respiratory Morbidity in COVID-19 Pneumonia: A Prospective Randomized Trial.

medRxiv (2020). doi:10.1101/2020.07.20.20157891

32. Dejgaard, S., Ortapamuk, O. & Özer, I. The Trypsin-Inhibitory Efficiency of Human

α_2-Macroglobulin in the Presence of α_1-Proteinase Inhibitor: Evidence for the

Formation of an α_2-Macroglobulin-α_1-Proteinase Inhibitor Complex. *J. Enzyme*

Inhib. **14**, 391–405 (1999).

Figure 1. Overview of SARS-CoV-2 entry and inhibition of trypsin treated CoV2pp entry by COVID-19 seronegative sera.

Figure 2. Negative patient sera inhibit exogenous protease mediated enhancement of CoV2pp.

Figure 3. Alpha-1-antitrypsin (AAT) and alpha-2-macroglobulin (A2M) inhibit protease mediated enhancement of CoV2pp entry.

Figure 4. Putative relationship between alpha-1-antitrypsin function and SARS-CoV-2.