Research Manuscript

Alpha-1-antitrypsin and its variant-dependent role in COVID-19 pathogenesis

Christian S Stevens[1][#], Kasopefoluwa Y Oguntuyo[1][#], Shreyas Kowdle[1], Luca Brambilla[1],

Griffin Haas[1], Aditya Gowlikar[1], Mohammed NA Siddiquey[2], Robert M Schilke[2], Matthew

D Woolard[2], Hongbo Zhang[2], Joshua A Acklin[1], Satoshi Ikegame[1], Chuan-Tien Huang[1],

Jean K Lim[1], Robert W Cross[3], Thomas W Geisbert[3], Stanimir S Ivanov[2], Jeremy P

Kamil[2], the Alpha-1 Foundation[4], and Benhur Lee[1]

Affiliations:

1. Department of Microbiology, Icahn School of Medicine at Mount Sinai, New York,

11   NY 10029

12   2. Department of Microbiology and Immunology, Louisiana State University Health

13   Shreveport, Shreveport, LA 71103

14   3. Department of Microbiology and Immunology, University of Texas Medical

15   Branch at Galveston, Galveston, TX 77555

4. The Alpha-1 Foundation, University of Florida, Coral Gables, FL 33134

[#]These authors contributed equally to this work.

Author contributions:

CSS, KYO, and BL conceived and designed the study. CSS, KYO, SK, LB, GH, AG,

MNAS, RMS, MDW, HZ, and JAA collected data. SI, CTH, JKL, JKL, RWC, TWG, SSI,

and JPK contributed valuable reagents, data, and/or tools. CSS and KYO analyzed the

data and wrote the original drafts of the paper. BL reviewed the draft, supported data

analysis, and provided invaluable direction throughout the conceptualization and

execution of the project. All authors had the opportunity to review the manuscript prior to

submission and JAA, SI, JKL, RWC, TWG, SSI, JPK provided valuable feedback during

the editing process.

Research Manuscript

ABSTRACT

Rationale: SARS-CoV-2 entry into host cells is facilitated by endogenous and

exogenous proteases that proteolytically activate the spike glycoprotein and

antiproteases inhibiting this process. Understanding the key actors in viral entry is

crucial for advancing knowledge of virus tropism, pathogenesis, and potential

therapeutic targets.

Objectives: We aimed to investigate the role of naïve serum and alpha-1-antitrypsin

(AAT) in inhibiting protease-mediated SARS-CoV-2 entry and explore the implications of

AAT deficiency on susceptibility to different SARS-CoV-2 variants.

Findings: Our study demonstrates that naïve serum exhibits significant inhibition of

SARS-CoV-2 entry, with AAT identified as the major serum protease inhibitor potently

restricting entry. Using pseudoparticles, replication-competent pseudoviruses, and

authentic SARS-CoV-2, we show that AAT inhibition occurs at low concentrations

compared with those in serum and bronchoalveolar tissues, suggesting physiological

relevance. Furthermore, sera from subjects with an AAT-deficient genotype show

reduced ability to inhibit entry of both Wuhan-Hu-1 (WT) and B.1.617.2 (Delta) but

exhibit no difference in inhibiting B.1.1.529 (Omicron) entry.

Conclusions: AAT may have a variant-dependent therapeutic potential against SARS-

CoV-2. Our findings highlight the importance of further investigating the complex

interplay between proteases, antiproteases, and spike glycoprotein activation in SARS-

CoV-2 and other respiratory viruses to identify potential therapeutic targets and improve

understanding of disease pathogenesis.

KEYWORDS: COVID-19, SARS-CoV-2, alpha-1-antitrypsin, SERPINA1, alpha-2-

macroglobulin, TMPRSS2, proteolytic activation, variants of concern, Omicron, Delta

INTRODUCTION

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), the causative agent

of coronavirus disease 2019 (COVID-19), depends on the complex interplay between

the virus and host for cellular entry (1–3). Understanding the various steps and factors

involved in viral entry is vital to our ability to successfully model the process, identify

potential therapeutics, and even predict genetic risk factors. For SARS-CoV-2, this

process is mediated by the spike glycoprotein (S). Entry involves not only the simple

binding interaction between spike and the entry receptor ACE2, but also the delicate

balance of proteases and antiproteases that contribute to proteolytic activation and

facilitate viral fusion (4–8). SARS-CoV-2 S undergoes two sequential proteolytic

activation steps: first it is cleaved at the S1/S2 polybasic site, and second it is then

cleaved at S2′ revealing the fusion peptide (9).

A variety of proteases are implicated in the proteolytic activation of SARS-CoV-2

including furin-like proteases, cathepsins, trypsin, neutrophil elastase, and TMPRSS2

(1, 9–16). This intricate process enables the virus to enter and fuse either at the plasma

membrane or within endosomes, engaging different proteases and pathways at each

stage. The complexities of viral entry for SARS-CoV-2 are particularly important to

understand, as they can be context-dependent, influenced by factors such as local

tissue-specific protease and antiprotease milieu, host genotype that impact these

milieux, and virus variants that alter susceptibility to cognate protease activation. As an

example of the way these relationships can develop and change over time, increasingly

efficient proteolytic processing and cell fusion were driving factors in the selective

pressure that gave rise to new variants in the first half of the pandemic. New SARS-

CoV-2 variants displayed more enhanced cell fusogenicity and proteolytic efficiency

relative to earlier strains, starting with the first dominant spike mutation, D614G. Alpha

and Beta both showed significantly increased fusogenicity and Delta even more so

relative to them (17–21). However, this pattern ended with the rise of the Omicron

sublineages as they rely predominantly on endosomal-mediated entry and have

significantly reduced fusogenicity (22–26).

Virus-specific mutations are not the only determinant of viral entry and pathogenicity, as

various host genotypes can affect virus-host interactions. SNPs in TMPRSS2 and ACE2

for example have been associated with differential COVID-19 risk and/or severity (27–

31). Extensive research has been conducted better elucidate the determinants of

SARS-CoV-2 pathogenicity. In this study, we investigated the cause of an unexpected

inhibition of viral entry by serum samples from patients not previously exposed to

SARS-CoV-2. We identify the likely causative factor and discuss its potential role within

a complex system where proteases, antiproteases, host genomics, and viral genomics

interact and influence each other.

RESULTS

Serum inhibition of protease-mediated entry of SARS-CoV-2 pseudoparticles and live virus

SARS-CoV-2 entry is efficiently mediated by a host of endogenous, exogenous, and cell surface proteases. Assays investigating viral entry and entry inhibition should faithfully recapitulate proteolytic activation (Figure 1A) for optimal physiological relevance. Using VSVΔG pseudotyped particles bearing SARS-CoV-2 spike (CoV2pp) we have shown that we can optimize entry efficiency by treatment with exogenous trypsin and followed by treatment with soybean trypsin inhibitor to limit cytotoxicity (32). We tested our trypsin-treated CoV2pp for validity using human serum samples, careful to only using pre-pandemic samples as negative controls (Supplementary Figure 1A-B) (32). As anticipated, sera from SARS-CoV-2 antibody-positive patients exhibit significantly stronger neutralization compared to seronegative sera (Figure 1B-C and Supplementary Figure 1C-F).

Unexpectedly, we found that sera from SARS-CoV-2 naïve patients were also capable of neutralizing these pseudoviruses despite negative Spike ELISA results (Supplementary Figure 1A). An external group at Louisiana State University Health Shreveport (LSUHS) confirmed these findings using the same CoV2pp assay and experimental conditions (Figure 1C and Supplementary Figure 1B, C-F). In both groups, SARS-CoV-2 naïve sera inhibited CoV2pp entry by 90-97% (Supplementary Figure 1E, F). However, seropositive patient sera exhibited inhibition orders of magnitude beyond

this threshold, suggesting antibody mediated inhibition of CoV2pp entry (<mark>Figure 1C and</mark>

<mark>Supplementary Figure 1C-F</mark>). Additionally, using identical serum samples as LSUHS,

collaborators at the University of Texas Medical Branch at Galveston (UTMB) also

observed significant neutralization of authentic SARS-CoV-2 by seronegative sera, as

assayed by a plaque reduction neutralization assay (PRNT) (<mark>Fig. 1E and Supplemental</mark>

<mark>Fig. 1G</mark>).

During the validation and optimization our CoV2pp system, we described that this

inhibition was abolished by heat-inactivation at 56ºC for 1hr (32). Here we show a

schematic representing entry inhibition by naïve sera from our previously published

findings and replicated by other published SARS-CoV-2 neutralization assays using

proteolytic activation without sufficient heat inactivation (33–35) (<mark>Figure 1B</mark>). These

findings imply the existence of a heat-labile serum factor or factors capable of inhibiting

protease-mediated entry of SARS-CoV-2.

A serum factor capable of inhibiting protease-mediated entry

Upon observing and verifying our results, we identified alpha-1-antitrypsin (AAT) and

alpha-2-macroglobulin (A2M) as abundant and heat labile products in serum that may

be responsible for inhibition of protease-mediated entry (36–39). These blood products

are typically present in human serum at high concentrations—1.1-2.2 mg/mL for AAT

and 2-4 mg/mL for A2M—and have been described to inhibit both exogenous and

endogenous proteases (40). A2M and AAT alone are responsible for approximately

10% and 90% of serum antiprotease capacity, respectively (41).

In spite of the name, the primary physiological target of AAT is neutrophil elastase (NE),

a protease released by neutrophils and found at high concentrations during acute

inflammation, especially in the context of acute respiratory distress syndrome (ARDS)

secondary to COVID-19 (42–44). In order to lend weight to the AAT hypothesis, we

therefore investigated whether NE, like trypsin or TMPRSS2, was also capable of

enhancing CoV2pp entry. We found that NE potently enhances cellular entry relative to

untreated particles (Figure 2A), further underlying the potential role of AAT in inhibiting

protease-mediate entry *in vivo*.

To assess whether AAT and/or A2M alone could inhibit protease-mediated CoV2pp

entry, we treated with each at the time of infection and observed dose-dependent entry

inhibition by both AAT and A2M, with IC50s of 14.47µg/mL and 54.20µg/mL,

respectively (Figure 2B), values that are 50-100-fold below their concentration in serum

and bronchoalveolar lavage (BAL) (36, 40, 45). Importantly, neither protein inhibited

VSV-Gpp, replicating the inhibitory effect of naïve serum (Figure 2C). Albumin, the most

abundant protein in blood, showed no significant reduction of entry of either CoV2pp or

VSV-Gpp (Figure 2B), underscoring that the inhibitory effects of AAT and A2M on

CoV2-S mediated entry was specific.

While these findings suggest that AAT—and to a lesser extent A2M—can inhibit

exogenous proteases known to enhance SARS-CoV-2 entry, SARS-CoV-2 infection is

also mediated by proteases at the cell surface as well as endosomal proteases (Figure

1A). TMPRSS2 and cathepsin L are well-characterized examples of cell surface and

endosomal proteases, respectively. AAT and A2M are secreted extracellular proteins

that can access the former but likely not the latter. Therefore, we sought to investigate

whether either protein could inhibit TMPRSS2-mediated SARS-CoV-2 entry. We first

showed that saturating amounts of nafamostat mesylate, a specific inhibitor of

TMPRSS2, maximally inhibited ~80% of CoV2pp entry in 293T-ACE2/TMPRSS2 cells

(Figure 3A) but had no effect on isogenic 293T-ACE2 cells (Figure 3B). This is

consistent with the use of endosomal proteases such as Cathepsin L in the absence of

TMPRSS2 or other exogenous proteases. Soluble Spike receptor binding domain

(sRBD) completely abolished CoV2pp entry in both cell lines confirming that entry was

still entirely ACE2-dependent (Figure 3A-B).

To examine inhibition of TMPRSS2-mediated entry by AAT and A2M, we infected both

cell lines with non-protease-treated CoV2pp. We observed that AAT inhibited CoV2pp

entry into 293T-ACE/TMPRSS2 cells, but not 293T-ACE2 cells (Figure 3C, D). A2M and

albumin both displayed no entry inhibition at the concentrations tested. AAT inhibition of

entry into the 293T ACE2-TMPRSS2 cells resulted in approximately a 70% drop in

relative infection (Figure 3C). This finding indicates that AAT inhibition accounts for

much of the entry attributed to TMPRSS2 enhancement, given that the use of

nafamostat mesylate resulted in a maximal inhibition of 80% in the 293T ACE2-

TMPRSS2 cells (Figure 3A). This observation suggests that AAT plays a significant role

in inhibiting TMPRSS2-mediated entry even at concentrations far below those seen in

serum and bronchioalveolar lavage (BAL) (1.1 – 2.2 milligrams/ml) (36, 40, 45).

In summary, our findings provide evidence that AAT, and to a lesser extent A2M, can

inhibit protease-mediated entry of SARS-CoV-2 in a cell culture model. AAT

demonstrated the ability to inhibit not only exogenous proteases like trypsin but also cell

surface protease TMPRSS2, which plays a crucial role in SARS-CoV-2 entry. These

results highlight the potential role of these serum factors in modulating viral entry *in*

vivo.

AAT inhibits protease-mediated entry of authentic SARS-CoV-2

To test whether these findings held for authentic SARS-CoV-2, we infected 293T-ACE2

cells and treated them with elastase, trypsin, or a cathepsin inhibitor (E64). We

measured the fraction of cells infected with SARS-CoV-2 at 6, 12, 24, and 36 hours

post-infection. Consistent with our CoV2pp observations, both elastase and trypsin

significantly enhanced entry, while E64 inhibited infection as expected, by inhibiting the

cathepsin-mediated pathway utilized when exogenous proteases and TMPRSS2 are

absent (Figure 4A).

Next, we performed serial dilutions of AAT in the presence of trypsin- and elastase-

treated SARS-CoV-2 at 24 hours post-infection (Figure 4B,C). We observed a dose-

dependent inhibition of protease-mediated entry, with potent inhibition at concentrations

far below those present in serum and bronchoalveolar lavage fluid. These results are

consistent with those obtained using the CoV2pp system, further demonstrating that

AAT can inhibit protease-mediated entry of live SARS-CoV-2.

In conclusion, our experiments with authentic SARS-CoV-2 validate the findings

obtained using the CoV2pp system, showing that AAT can effectively inhibit protease-

mediated entry of the virus. This highlights the potential physiological relevance of AAT

in modulating SARS-CoV-2 infection.

Serum-inhibition of rVSVCov2 entry is reduced in subjects with AAT deficient

genotypes

Recognizing AATs potential as an inhibitor of protease-mediated entry enhancement,

we aimed to better elucidate the potential physiological relevance of these findings to

individuals of various AAT genotypes. AAT levels in serum varies across individuals and

are determined by co-dominant alleles in the SERPINA1 gene. The four genotypes

investigated here are designated as PI*MM, PI*MS, PI*MZ, and PI*ZZ. Typically, PI*MM

individuals have normal AAT levels, while PI*MS and PI*MZ individuals have varying

amounts that range from near to sub-normal AAT levels. PI*ZZ individuals are

considered AAT deficient (AATD) (Figure 5A).

To better elucidate the physiological implications our findings we obtained pre-2020

serum samples from individuals with these four major AAT genotypes and tested their

ability to inhibit SARS-CoV-2 Spike-mediated entry. We used replication-competent

VSV containing Spike in place of G (rVSV-CoV-2-S) to model SARS-CoV-2 entry. We

first infected Vero cells with trypsin-treated WT rVSV-CoV-2-S in the presence of

serially diluted serum from the indicated genotypes (Fig. 5B and Supplemental Figure

2A). At 12 hours post-infection, we observed elevated GFP expression at higher serum

concentrations for PI*ZZ samples compared to PI*MM, PI*MS, and PI*MZ samples

(Figure 5B and Supplemental Figure 2A). This suggests that the most severe AATD

genotypes have reduced serum inhibitory potential against protease-mediated entry of

WT SARS-CoV-2 Spike, but this reduction is not observed for heterozygotes of the Z

and S alleles.

During the SARS-CoV-2 pandemic, several viral lineages have emerged and become

dominant. While many similarities exist between all variants of concern (VOCs), some

show distinct differences, particularly in viral entry and proteolytic processing of Spike

(4, 22, 25). Therefore, we investigated whether the effects of AAT would remain

consistent across all SARS-CoV-2 VOCs.

Using our rVSV-CoV-2-S system, we compared entry of isogenic viruses differing only

in Spike to model entry by different SARS-CoV-2 VOCs (1). The pattern observed for

WT rVSV-CoV-2-S was also seen with Delta rVSV-CoV-2-S containing the Spike

protein from B.1.617.2. This VOC became dominant in mid to late 2021, and AATD sera

were still less able to inhibit protease-mediated entry (Figure 5C).

Finally, we examined Omicron rVSV-CoV2 containing the Spike protein from B.1.1.529.

The Omicron lineage became dominant in 2022 and has remained the dominant lineage

since (46), overtaking Delta as the primary VOC in circulation. Notably, Omicron has

been shown to undergo less enhancement of entry by proteolytic activation of Spike

(22, 25). Using Omicron rVSV-CoV2, we observed no significant difference in inhibitory

potential between PI*ZZ and other AAT genotypes (Figure 5D), unlike what we observe

for Delta and WT (Figure 5B,C). These data suggest that AAT, and consequently AATD,

may have less impact on protease-mediated entry for the Omicron variant compared to

WT and Delta variants.

DISCUSSION

In this study, we identify alpha-1 antitrypsin as the primary inhibitor of protease-

mediated entry of SARS-CoV-2 in naïve serum. We demonstrated that AAT effectively

inhibits entry mediated by trypsin, elastase, or TMPRSS2. Furthermore, we found that

serum from individuals with at least one functional AAT allele (PI*MM, PI*MS, PI*MZ)

can inhibit protease-mediated entry by both WT and Delta SARS-CoV-2. Notably,

serum from subjects homozygous for AATD alleles (PI*ZZ) exhibited reduced ability to

inhibit WT and Delta entry. However, genotype-dependent inhibition of entry was not

observed for the Omicron variant, as all serum samples—regardless of subject

genotype—displayed similar inhibitory potential. This finding highlights the potential for

AAT's role to vary depending on the entry mechanisms predominantly relied on by the

virus.

Approximately 116 million individuals worldwide carry AATD alleles—PI*S or PI*Z—and

3.4 million people are homozygous for AATD—PI*SS, PI*SZ, and PI*ZZ (47). In the

early stages of the COVID-19 pandemic, studies reported correlations between mortality

rates and the geographic distribution of AAT deficiency (48–50). More recent

retrospective analyses indicate that heterozygotes for AATD alleles (e.g. PI*MS, PI*MZ)

do not appear to face an increased risk of severe COVID-19 (51). This observation

aligns with our findings regarding the inhibitory potential of PI*MM, PI*MS, and PI*MZ

sera. Unfortunately, determining a clear increased risk of COVID-19 severity for PI*ZZ

patients remains elusive, as large-scale population studies have not yet included a

sufficient number of these individuals (52–54). Moreover, disentangling the confounding

influence of patients' awareness of their AATD status presents a challenge. In Germany,

survey data from March 2021 revealed that individuals diagnosed with AATD were more

concerned with being infected, more likely to restrict their social groups in response,

and saw a 65% drop in the percent who had been infected relative to the general

population (55).

An additional contributing factor in elucidating the risk associated the alpha-1 antitrypsin

deficiency is the emergence of the Omicron lineage of SARS-CoV-2, which is less

dependent on the protease-mediated entry pathway for efficient infection (25). As

Omicron has become the dominant variant, it has affected the potential risk factors for

more severe COVID-19. New variants may alter the role of AAT in disease severity,

leading to a different risk profile for individuals with AATD. The COVID-AATD Study

from Spain, involving over 2000 patients, showed that both AATD mutations and AAT

serum concentration below 116 mg/dl were associated with severe COVID-19 (56). One

limitation of this study might be that it was conducted from 1 May 2021 to 1 September

2022, and Omicron (BA.1) overtook Delta as the dominant strain in Spain around 3

January 2022 (57). If AATs antiprotease activity is playing a significant role in their

findings of increased risk of COVID-19 severity, separating subjects by estimated date

of infection may uncover whether the relationship is lessened after January 2022.

AAT has been predominantly described in the SARS-CoV-2 literature for its role as an

acute phase protein in modulating the host immune response (58). AAT plays a critical

part in modulating inflammation by inhibiting elastase and other factors (59–62). It is

involved in the formation of neutrophil extracellular traps (NETs) in acute pneumonia

and can modulate activities resulting in downstream IL-6 inhibition, which is heavily

implicated in COVID-19 pathogenicity (41–44, 63). In particular, the IL-6:AAT ratio is

highly correlated with COVID-19 severity (64). AAT is also noted for its regulatory role in

the coagulation cascade which may have relevance in preventing COVID-19

thromboses (43, 65–67). Previous works have also noted the importance of AAT's

ability to inhibit TMPRSS2 (68). Our study further elucidates the antiprotease role of

AAT and suggests that the presence or absence of functional AAT could influence the

efficiency of viral entry in other respiratory pathogens.

Patients diagnosed with AATD can receive supplementation through an FDA approved

medication: either IV or aerosolized AAT. Four clinical trials for use of AAT in the

context of COVID-19 have been registered. One never proceeded beyond recruitment

(NCT04385836), one was completed but has not published results (NCT04495101),

and one was stopped for futility (NCT04547140). One focused on treating with AAT

following a diagnosis of Acute Respiratory Distress Syndrome (ARDS) and saw

moderate effects, mostly on modulating inflammation (EudraCT 2020-001391-15) (69).

AAT as a treatment for proteolytically activated respiratory viruses shows promise as

both an immune modulator as well as a protease-inhibitor and would be well served by

clinical trials investigating treatment of a larger patient population as well as treatment in

the earlier stages of infection.

Despite the modest results seen in clinical trials of AAT supplementation, the clinical

implications of these findings for patients with an AAT deficiency are significant. Typical

clinical guidance is that AATD patients are only at increased risk of respiratory

infections secondary to chronic obstructive pulmonary disease, not primarily due to the

lack of functional AAT except in the case of immune dysregulation (70, 71). Our work

suggests this may miss the AATs role as an inhibitor of serine proteases and how that

might affect respiratory pathogens reliant on proteolytic activation. For example, not

only is TMPRSS2 known to be an important protease for proteolytic activation of the

surface glycoprotein hemagglutinin in H1, H3, H7, and H10 influenza A viruses (IAV)

(72, 73) but AAT specifically has been shown to inhibit protease-mediated entry of

H3N2 IAV and influenza B virus (72). Additionally, multiple studies have implicated AAT

in HIV entry as well (62, 74, 75). AAT's role in viral entry may be specific to the reliance

on proteolytic processing at or outside the cell. We show that in the case of SARS-CoV-

2, new variants may alter that role, leading to a different risk profile for individuals with

AATD. It is crucial to continue investigating the relationship between AAT and new

variants to better understand the evolving landscape of risk factors for COVID-19.

Furthermore, our findings suggest that it may be worthwhile to invest more into

investigating the role of AAT in other respiratory viruses mediated by serine proteases

(76, 77). A better understanding of the interplay between AAT and these viruses could

help identify potential therapeutic targets, improve patient outcomes, and affect clinical

guidance for AATD patients in relation to other respiratory pathogens.

In conclusion, our study highlights the critical role of AAT in inhibiting protease-mediated

entry of SARS-CoV-2 and its potential implications for treatment, especially in patients

with AAT deficiency. As the COVID-19 pandemic evolves with the emergence of new

variants, understanding the role of AAT in the context of these variants is crucial for

assessing the shifting risk profiles of individuals with AATD. Our findings also suggest

that AAT may play a role in the pathogenesis of other respiratory viruses mediated by

serine proteases, opening avenues for future research into the therapeutic potential of

AAT in treating these infections. Investigating the role of AAT in other respiratory

viruses could lead to the identification of potential therapeutic targets, improved patient

outcomes, and updated clinical guidance for AATD patients in relation to other

respiratory pathogens. Lastly, further investigation is needed to expand our

understanding of AAT's clinical implications, including the efficacy of AAT

supplementation for treating respiratory infections and determining the risk factors for

AATD patients in the context of COVID-19 and other respiratory illnesses. By deepening

our understanding of AAT's role in viral pathogenicity and its potential as a therapeutic

target, we can better inform clinical practice and contribute to improved public health

outcomes.

METHODS

<u>Maintenance and generation of isogenic cell lines:</u>

Vero-CCL81, BHK-21, Bsr-T7 (78), 293T, 293T ACE2, and 293T ACE2-TMPRSS2 cells

were cultured in DMEM with 10% heat inactivated FBS at 37ºC in the presence of 5%

CO_2. Isogenic 293T ACE2, and 293T ACE2-TMPRSS2 cell clones were generated by

lentivirus transduction to stably express ACE2 only or ACE2 and TMPRSS2. ACE2

expression was under puromycin selection and TMPRSS2 was under blasticidin

selection as previously described (32).

<u>Production of VSV∆G pseudotyped particles and neutralization studies:</u>

Detailed protocols for the production and use of standardized VSVpp (CoV2pp and

VSV-Gpp) are given in Oguntuyo and Stevens et al (32). Briefly, 293T producer cells

were transfected to express the viral surface glycoprotein of interest, infected with

VSV∆G-rLuc-G* reporter virus, then virus supernatant collected and clarified 2 days

post infection prior to use. Trypsin-treated CoV2pp were treated as previously described

(32). All pseudotyped viruses were aliquoted prior to storage at -80ºC and tittered prior

to usage for neutralization experiments. Neutralization experiments were performed by

diluting the appropriate pseudotyped virus with a 4-fold serial dilution of Albumin

(Sigma-Aldrich, A8763), alpha-1-antitrypsin (Sigma-Aldrich, SRP6312), alpha-2-

macroglobulin (Sigma-Aldrich, SRP6314) or Nafamostat mesylate (Selleckchem,

S1386). SARS-CoV-2 soluble RBD (sRBD) with human IgG-Fc was produced using a

recombinant Sendai virus expression platform further described below. All infections

were processed for detection of Renilla luciferase activity at 20hrs post-infection, and

luminescence was read on the Cytation3 (BioTek).

<u>SARS-CoV-2 plaque reduction neutralization titration (PRNT) by sera:</u>

Neutralization experiments with live virus were performed by incubating sera with 50-

100 PFU of SARS-CoV-2, isolate USA-WA1/2020 P4, for one hour at 37ºC. All sera

were diluted in serum free DMEM. Serial dilutions started at a four-fold dilution and went

through seven three-fold serial dilutions. The virus-serum mixture was then used to

inoculate Vero E6 cells for one hour at 37ºC and 5% CO_2. Cells were overlaid with

EMEM medium (no FBS) and 1.25% Avicel, incubated for 3 days, and plaques were

counted after staining with 1% crystal violet in formalin.

<u>SARS-CoV-2 infection of 293T-ACE2 cells and protease treatment:</u>

SARS-CoV-2 isolate USA-WA1/2020 (NR-52281) was provided by the Center for

Disease Control and Prevention and obtained through BEI Resources, NIAID, NIH.

$1x10^6$ 293T cells stably transduced to express ACE2 were plated in a 6 well dish and

infected at an MOI of 0.01 for the time period indicated. Just before infection, virus was

treated with alpha-1-antitrypsin (Sigma-Aldrich, SRP6312) at the concentration noted,

elastase from human leukocytes (Sigma-Aldrich, E8140) at 0.167 mg/mL, TPCK-treated

trypsin (Sigma-Aldrich, T1426-1G) at 50 µg/mL, or E-64 (Sigma-Aldrich, E3132) at 100

µM. Cells were harvested into 4% PFA and were allowed to fix for 30 minutes prior to

staining for flow cytometry. Infection was determined with mouse anti-SARS-CoV

nucleoprotein antibody, directly conjugated to Alexa Fluor 594. Samples were collected

on through an Attune NxT Flow Cytometer and data was analyzed using FlowJo

software (v10.6.2). All SARS-CoV-2 work was performed in the CDC and USDA-

approved BSL-3 facility at the Icahn School of Medicine at Mount Sinai in accordance

with institutional biosafety requirements.

<u>Production of Replication competent VSV∆G with SARS-CoV-2 Spike Variants and</u>

<u>Neutralization Studies:</u>

VSV-eGFP was cloned into the pEMC vector containing an optimized T7 promotor and

hammerhead ribozyme. Original VSV-eGFP sequence was from pVSV-eGFP; a gift

from Dr. John Rose (79). pEMC-VSV-eGFP-CoV2 Spike was cloned as follows: human

codon optimized SARS-CoV-2 Spike variants with the 21 amino acid truncation of the

cytoplasmic tail were inserted into the VSV-G open reading frame (80) (rVSV-CoV2).

The Spike transcriptional unit is flanked by MluI and PacI restriction sites. Expression

plasmids containing VSV N, P, M, G, and L open reading frames were each cloned into

a pCI vector backbone to allow for efficient virus rescue, generating pCI-VSV-N, pCI-

VSV-P, pCI-VSV-M, pCI-VSV-G, and pCI-VSV-L.

rVSV-CoV2 was rescued in $4x10^5$ 293T ACE2-TMPRSS2 or BHK-21 ACE2 cells (32) in

each well of a 6-well plate. 2000 ng of pEMC-VSV-EGFP-CoV2 spike, 2500 ng of

pCAGGS-T7opt (81), 850 ng of pCI-VSV-N, 400 ng of pCI-VSV-P, 100 ug of pCI-VSV-

M, 100 ng of pCI-VSV-G, 100 ng of pCI-VSV-L were mixed with 4 ml of Plus reagent

and 6.6 ml of Lipofectamine LTX (Invitrogen). 30 minutes later, the transfection mixture

was applied to 293T ACE2-TMPRSS2 cells dropwise. Cells were maintained with

medium replacement every day for 4-5 days until GFP positive syncytia appeared.

Rescued viruses were amplified in VeroCCL81 TMPRSS2 cells (32), harvested after 6

days, stored at -80C. For titration, 5×10^4 293T ACE2-TMPRSS2 cells were seeded onto

each well of a 96-well plate and 24 hours later were infected with serially diluted rVSV-

CoV2 stock. Virus titer (IU/mL) was calculated 10 hours later by counting GFP positive

cells on the Celigo imaging cytometer (Nexcelom).

Human Sera Samples:

All patient sera were acquired after approval by the respective institutional review

boards (IRBs) and/or equivalent oversight bodies (Bioethics Committee, Independent

Ethics Committee), as follows: (i) the Mount Sinai Hospital IRB (New York, NY, USA),

(ii) the Louisiana State University Health Sciences Center—Shreveport (LSUHS, LA,

USA), and (iii) the Alpha-1 Foundation (University of Florida, Coral Gables, FL, USA).

Samples were deidentified at the source institutions or by the respective principal

investigators (PIs) of the IRB-approved protocols for sample collection before analyses

performed in this study. All necessary patient/participant consent has been obtained,

and the appropriate institutional forms have been archived. Specifically, SERPINA1

genotyped sera samples collected before 2019 were obtained from the Alpha-1

Foundation.

Enzyme-Linked Immunosorbent Assay:

Spike ELISAs for patient sera from the Krammer lab were performed in a clinical setting

using the two-step protocol previously published (32, 82). Briefly, this involves screening

patient sera (at a 1:50 dilution) with the sRBD; samples determined to be positive were

further screened at 5 dilutions for reactivity to the spike ectodomain. Background was

subtracted from the OD values, samples were determined to be positive if their ODs

were ≥3-fold over that of the negative control, and the AUC was calculated in PRISM.

ELISAs performed by the LSUHS group utilized the sRBD with a 1:50 dilution of patient

serum to screen all samples, followed by use of the spike ectodomain with patient sera

at a 1:100 dilution. Background-subtracted OD values are reported for both sets of

ELISAs.

Production of Soluble SARS-CoV-2 Spike Receptor Binding Domain (sRBD):

Sendai virus (SeV) Z strain (AB855655.1) was cloned into a pRS vector backbone with

an additional eGFP transcriptional unit upstream of N. The F transcriptional unit was

derived from the SeV Fushimi strain (KY295909.1). We then generated an additional

transcriptional unit between the P gene and M gene. SARS-CoV-2 Spike receptor

binding domain (sRBD), amino acids 319-541, was taken from human codon optimized

Spike (MN908947) in a pCAGGS backbone, a gift from Dr. Florian Krammer (82), and

was fused to human IgG1 Fc (amino acids 220-449 of P0DOX5.2) at the C-terminus

(SeV-Z-eGFP-sRBD).

$2x10^5$ Bsr-T7 cells, stably expressing T7-polymerase, were seeded in a 6-well plate. 24

hours later 4 μg of pRS-SeV-Z-eGFP-sRBD, 4 μg of pCAGGS-T7opt, 1.44 μg of SeV-N,

0.77 μg of SeV-P, 0.07 μg of SeV-L were mixed with 5.5 μl of Plus reagent and 8.9 μl of

Lipofectamine LTX (Invitrogen). 30 minutes later, the transfection mixture was applied to

Bsr-T7 cells dropwise. Medium was replaced with DMEM + 0.2 µg /ml of TPCK-trypsin

(Millipore Sigma, #T1426) at one day post transfection, followed by media replacement

every day until infection reached confluency. Supernatant was stored at -80C.

Amplification was performed by seeding 5×10^6 cells in a T175cm^2-flask one day before

infection. Cells were infected by SeV-Z-eGFP-sRBD at an MOI of 0.01 for one hour,

followed by a media replacement with 0.2 mg/ml of TPCK-trypsin-containing DMEM.

Cells were maintained with medium replacement by the same every day until infection

reached confluency. At maximal infection the medium was changed and replaced with

plain DMEM. Cells were incubated for additional 24 hours to allow for maximum protein

production. Supernatant was collected and centrifuged at 360 g for 5 minutes, then

filtered with 0.1 µm filter (Corning 500 mL Vacuum Filter/Storage Bottle System, 0.1 µm

Pore). The flow-through was then applied to a Protein G Sepharose (Millipore Sigma,

#GE17-0618-01) containing column (5ml polypropylene columns; ThermoFisher,

#29922), followed by wash and elution.

<u>Statistics and reproducibility:</u>

All statistical tests were performed using GraphPad Prism 9 software (La Jolla, CA).

For all figures, error bars represent standard deviation of the mean. Sample size and

replicates for each experiment are indicated in the figure legends. Technical replicates

were prepared in parallel within one experiment, and experimental replicates were

performed on separate days. Statistical comparisons as noted in figure legends.

ACKNOWLEDGMENTS

The authors acknowledge the following funding: KYO and CS were supported by Viral-

Host Pathogenesis Training Grant T32 AI07647; KYO was additionally supported

by F31 AI154739. BL acknowledges flexible funding support from NIH grants R01

AI123449, R21 AI1498033, and the Department of Microbiology and the Ward-Coleman

estate for endowing the Ward-Coleman Chairs at the ISMMS. JPK and SSI

acknowledge funding from a LSUHS COVID-19 intramural grant. JPK and SSI

acknowledge additional funding from NIH grants AI116851 and AI143839, respectively.

Figures created with BioRender.com. We thank Randy A. Albrecht for oversight of the

conventional BSL3 biocontainment facility. We would also like to acknowledge the

Alpha-1 Foundation and the University of Florida which kindly provided serum samples

from SERPINA1-genotyped patients. BL wishes to dedicate this paper to Ernest L

Robles-Levroney, the first graduate student BL had the privilege to train. Ernie Robles-

Levroney was dedicated teacher, role model and trailblazer who passed away

unexpectedly during the course of writing this manuscript.

ACKNOWLEDGMENTS

The authors acknowledge the following funding: KYO and CS were supported by Viral-

Host Pathogenesis Training Grant T32 AI07647; KYO was additionally supported

by F31 AI154739. BL acknowledges flexible funding support from NIH grants R01

AI123449, R21 AI1498033, and the Department of Microbiology and the Ward-Coleman

estate for endowing the Ward-Coleman Chairs at the ISMMS. JPK and SSI

acknowledge funding from a LSUHS COVID-19 intramural grant. JPK and SSI

acknowledge additional funding from NIH grants AI116851 and AI143839, respectively.

Figures created with BioRender.com. We thank Randy A. Albrecht for oversight of the

conventional BSL3 biocontainment facility. We would also like to acknowledge the

Alpha-1 Foundation and the University of Florida which kindly provided serum samples

from SERPINA1-genotyped patients. BL wishes to dedicate this paper to Ernest L

Robles-Levroney, the first graduate student BL had the privilege to train. Ernie Robles-

Levroney was dedicated teacher, role model and trailblazer who passed away

unexpectedly during the course of writing this manuscript.

FIGURES

Figure 1. Overview of SARS-CoV-2 entry and inhibition of trypsin treated CoV2pp

entry by COVID-19 seronegative sera. **(A)** Overview of SARS-CoV-2 entry. Two

modes of entry are displayed: 1) Entry mediated by endosomal proteases such as

Cathepsin B/L (late entry) and 2) protease-mediated entry (early entry), driven by cell-

surface proteases like TMPRSS2 and extracellular proteases such as trypsin and

elastase. Protease-mediated entry may be inhibited by the presence of antiproteases.

This model was created in Biorender. **(B)** A representative schematic of entry inhibition

of trypsin-treated SARS-CoV-2 pseudotyped particles (CoV2pp) by sera from COVID-19

recovered or naïve individuals (COVID sero (+) and COVID sero (-) sera, respectively).

This is a representation of results previously presented in supplemental Figure 3A of

Oguntuyo and Stevens et al, mBio 2021, as well as Figure 5B of Nie et al, 2020. Here,

sera samples were incubated with trypsin-treated CoV2pp prior to infection of Vero-

CCL81 cells. Grey lines represent seronegative sera and purple lines are COVID-19

seropositive sera. The dashed lines are samples that were heat inactivated (HI) prior to

use. **(C)** Seronegative and seropositive samples were first identified based on IgG

antibodies against Spike (Supplemental Figure 1B). Normalized infection data at the

highest and lowest dilutions tested are shown as % maximal infection (media only) with

results from seronegative plotted on log scale. Data points represent the mean of

neutralizations performed in quadruplicate with SEM bars, each line indicating a sample

from a unique donor. Maximal sera inhibition was compared using Welch's t test. **(D)**

SARS-CoV-2 seronegative sera do not inhibit VSV-Gpp. Four serum samples were

analyzed each in technical triplicate, means with SEM error bars shown. **(E)** Authentic

SARS-CoV-2 is modestly inhibited by seronegative sera. Sera samples also presented

in Supplementary Figure 1E were utilized for plaque reduction neutralization

experiments (PRNT) with live virus. Presented here are the mean of one experiment

done in technical duplicates with SEM error bars. Maximal sera inhibition was compared

using Welch's t test. (ns, not significant; **, $p < 0.01$; ***, $p < 0.005$, and ****, $p <$

0.0001).

Figure 2. CoV2pp is enhanced by elastase treatment and alpha-1-antitrypsin

(AAT) and alpha-2-macroglobulin (A2M) inhibit trypsin-mediated enhancement of

CoV2pp entry. (A) Treatment of both CoV2pp and VSV-Gpp with elastase in serum-

free media. All points are means with SEM bars for samples performed in technical

triplicate. Red dotted line marks normalized maximal infection level (100%). **(B)** AAT

and A2M inhibit trypsin-mediated enhancement of CoV2pp. Trypsin-treated

pseudoparticles were diluted in serum free media, then used to infect Vero-CCL81 cells

in the presence of the indicated concentrations of albumin, AAT, or A2M. Data are from

two independent experiments and are presented as percent relative infection where

each concentration was normalized to the lowest concentration of the test reagent used.

(C) Performed identical to (B), AAT, A2M, and albumin have no effect on VSV-Gpp

entry.

Figure 3. Alpha-1-antitrypsin (AAT) inhibits TMPRSS2 mediated enhancement of

CoV2pp entry. (A,B) CoV2pp were mixed with a serial dilution of either Nafamostat or

sRBD prior to infection of isogenic cells stably expressing (A) ACE2 and TMPRSS2 or

(B) ACE2. **(C)** AAT inhibits TMPRSS2-mediated enhancement of CoV2pp entry.

CoV2pp not treated with trypsin were diluted in DMEM+10% FBS and utilized to infect

293T-ACE2+TMPRSS2 clonal in the presence of the indicated concentrations of A2M,

AAT, or Albumin. **(D)** CoV2pp, treated exactly as in **(C)**, were used to infect a 293T-

ACE2 clonal cell line. Presented here are the results of an experiment done in technical

triplicates. Error bars show SEM and data were fit using variable slope, 4-parameter

logistics regression curve (robust fitting method). Significance calculated using a

Welch's T Test on the Area Under the Curve for each condition. (ns, not significant; **, p

< 0.01; ***, p < 0.005, and ****, p < 0.0001).

Figure 4. Alpha-1-antitrypsin (AAT) inhibits protease-mediated enhancement of

authentic SARS-CoV-2 entry. (A) Authentic SARS-CoV-2 entry in 293T-ACE2 clonal

cells over 36 hours under treatment of elastase, trypsin, E64, or untreated. Significance

calculated using a Welch's T Test on the Area Under the Curve for each condition. **(B)**

Authentic SARS-CoV-2 entry in 293T-ACE2 cells mediated by trypsin, treated by

increasing concentrations of AAT, collected at 0, 12, 24, or 36 hours post-infection. **(C)**

Authentic SARS-CoV-2 entry in 293T-ACE2 cells mediated by elastase, treated by

increasing concentrations of AAT. Data points are means +/- SEM from a representative

experiment performed in triplicate. (ns, not significant; **, p < 0.01; ***, p < 0.005, and

****, p < 0.0001).

Figure 5. rVSV-CoV2 entry inhibition by serum from subjects with variable AAT

genotypes. (A) Expected range of concentrations of AAT in serum relative to AAT

genotype (data from Lopes et al (83)). **(B)** WT rVSV-CoV2 entry inhibition by serum

from 4 PI*MM subjects, 2 PI*MS subjects, 2 PI*MZ subjects, and 4 PI*ZZ subjects

performed in technical triplicate. Subject serum samples are identical for all data shown

across all variants. Boxes span 25-75th percentiles and median is noted. Whiskers span

minimum to maximum. **(C)** Delta rVSV-CoV2 entry inhibition by serum as described in

3B except only 2 PI*MM and 2 PI*ZZ subjects are shown. **(D)** Omicron rVSV-CoV2

entry inhibition by serum as described in 3B. All data shown performed in technical

triplicate. All rVSV-CoV-2 is trypsin-treated. Significance calculated using Welch's t test

(ns, not significant; **, $p < 0.01$; ***, $p < 0.005$, and ****, $p < 0.0001$).

Supplemental Figure 1. Spike ELISA data and neutralization curves. Spike

ectodomain ELISAs for **(A)** ISMMS or **(B)** LSUHS samples. Four seronegative and

seropositive samples were utilized. Shown are the OD490 values from the 1:100 sera

dilution with the median and interquartile range. **(C)** Seronegative and seropositive

samples were first identified based on IgG antibodies against Spike (Supplemental

Figure 1A). Normalized infection data at the highest and lowest dilutions tested are

shown as % maximal infection (media only) with results from seronegative plotted on

log scale. Data points represent the mean of neutralizations performed in quadruplicate

with SEM bars, each line indicating a sample from a unique donor. Maximal sera

inhibition was compared using Welch's t test. **(D)** Sera inhibition comparison against

VSV-Gpp for 4 sera samples collected at ISMMS. Sera inhibition comparisons for **(E)**

ISMMS or **(F)** LSUHS. Inhibition of trypsin treated CoV2pp entry by SARS-CoV-2

seropositive and seronegative sera independently observed. Collaborators in a different

state independently performed the identical experiment described in Fig. 1A with their

own cohort of seropositive and seronegative samples. Data shown are means from

technical quadruplicates/sample/dilution. Experiment performed and presented as in

Fig. 1C. **(G)** Live SARS-CoV-2 is inhibited by seropositive and seronegative sera. Sera

samples presented in Fig. 1C and (F) above were utilized for plaque reduction

neutralization experiments (PRNT) with live virus as described in the materials and

methods. Presented here are the mean of one experiment done in technical duplicates

and error bars show SEM. (ns, not significant; **, $p < 0.01$; ***, $p < 0.005$, and ****, $p <$

0.0001).

Supplemental Figure 2. Individual curves of rVSV-CoV2 entry inhibition by serum from

subjects with variable AAT genotypes in spike variants: **(A)** WT (Wuhan-Hu-1) and **(B)**

Omicron (B.1.1.529). Error bars show SEM and data were fit using variable slope, 4-

parameter logistics regression curve (robust fitting method). **(C)** Estimation plots for

area under the curve shown for PI*MM compared to PI*ZZ in WT rVSV-CoV2 and **(D)**

PI*MM compared to PI*ZZ in Omicron rVSV-CoV2. Significance calculated using a

Welch's t test on the Area Under the Curve for each condition. (ns, not significant; **, p

< 0.01; ***, $p < 0.005$, and ****, $p < 0.0001$).

REFERENCES

1. Stevens CS, Oguntuyo KY, Lee B. Proteases and variants: context matters for SARS-CoV-2

entry assays. *Curr. Opin. Virol.* 2021;50:49–58.

2. Hoffmann M et al. SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked

by a Clinically Proven Protease Inhibitor. *Cell* 2020;181(2):271-280.e8.

3. Zhu N et al. A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N. Engl. J.*

Med. 2020;382(8):727–733.

4. Letko M, Marzi A, Munster V. Functional assessment of cell entry and receptor usage for

SARS-CoV-2 and other lineage B betacoronaviruses. *Nat. Microbiol.* 2020;5(4):562–569.

5. Tang T, Bidon M, Jaimes JA, Whittaker GR, Daniel S. Coronavirus membrane fusion

mechanism offers a potential target for antiviral development. *Antiviral Res.*

2020;178(March):104792.

6. Wrapp D et al. Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation.

Science 2020;367(6483):1260–1263.

7. Zhu FC et al. Proteolytic activation of the SARS-coronavirus spike protein: Cutting enzymes

at the cutting edge of antiviral research. *J. Virol.* 2020;395(2):379–397.

8. Shang J et al. Cell entry mechanisms of SARS-CoV-2. *Proc. Natl. Acad. Sci.*

2020;117(21):11727–11734.

9. Tang T et al. Proteolytic Activation of SARS-CoV-2 Spike at the S1/S2 Boundary: Potential

Role of Proteases beyond Furin. *ACS Infect. Dis.* 2021;7(2):264–272.

10. Bhattacharyya C et al. SARS-CoV-2 mutation 614G creates an elastase cleavage site

enhancing its spread in high AAT-deficient regions. *Infect. Genet. Evol.* 2021;90:104760.

11. Belouzard S, Madu I, Whittaker GR. Elastase-mediated Activation of the Severe Acute

Respiratory Syndrome Coronavirus Spike Protein at Discrete Sites within the S2 Domain. *J.*

Biol. Chem. 2010;285(30):22758–22763.

12. Fuentes-Prior P. Priming of SARS-CoV-2 S protein by several membrane-bound serine

proteinases could explain enhanced viral infectivity and systemic COVID-19 infection. *J. Biol.*

Chem. 2021;296:100135.

13. Bollavaram K et al. Multiple sites on SARS-CoV-2 spike protein are susceptible to

proteolysis by cathepsins B, K, L, S, and V. *Protein Sci.* 2021;30(6):1131–1143.

14. B C et al. The spike glycoprotein of the new coronavirus 2019-nCoV contains a furin-like

cleavage site absent in CoV of the same clade [Internet]. *Antiviral Res.* 2020;176.

doi:10.1016/j.antiviral.2020.104742

15. G P et al. Furin cleavage of SARS-CoV-2 Spike promotes but is not essential for infection

and cell-cell fusion [Internet]. *PLoS Pathog.* 2021;17(1). doi:10.1371/journal.ppat.1009246

16. Peacock TP et al. The furin cleavage site in the SARS-CoV-2 spike protein is required for

transmission in ferrets. *Nat. Microbiol.* 2021;6(7):899–909.

17. Mlcochova P et al. SARS-CoV-2 B.1.617.2 Delta variant replication and immune evasion.

Nature 2021;599(7883):114–119.

18. Gobeil SM-C et al. D614G Mutation Alters SARS-CoV-2 Spike Conformation and Enhances

Protease Cleavage at the S1/S2 Junction. *Cell Rep.* 2021;34(2):108630.

19. Lubinski B et al. Spike Protein Cleavage-Activation in the Context of the SARS-CoV-2

P681R Mutation: an Analysis from Its First Appearance in Lineage A.23.1 Identified in Uganda.

Microbiol. Spectr. 2022;10(4):e01514-22.

20. Plante JA et al. Spike mutation D614G alters SARS-CoV-2 fitness. *Nature*

2021;592(7852):116–121.

21. Rajah MM et al. SARS-CoV-2 Alpha, Beta, and Delta variants display enhanced Spike-

mediated syncytia formation. *EMBO J.* 2021;40(24):e108944.

22. Zhao H et al. SARS-CoV-2 Omicron variant shows less efficient replication and fusion

activity when compared with Delta variant in TMPRSS2-expressed cells. *Emerg. Microbes*

Infect. 2022;11(1):277–283.

23. Hui KPY et al. SARS-CoV-2 Omicron variant replication in human bronchus and lung ex

vivo. *Nature* 2022;603(7902):715–720.

24. Meng B et al. Altered TMPRSS2 usage by SARS-CoV-2 Omicron impacts infectivity and

fusogenicity. *Nature* 2022;603(7902):706–714.

25. Willett BJ et al. SARS-CoV-2 Omicron is an immune escape variant with an altered cell

entry pathway. *Nat. Microbiol.* 2022;7(8):1161–1179.

26. Kumar S, Thambiraja TS, Karuppanan K, Subramaniam G. Omicron and Delta variant of

SARS-CoV-2: A comparative computational study of spike protein. *J. Med. Virol.*

2022;94(4):1641–1649.

27. Paniri A, Hosseini MM, Akhavan-Niaki H. First comprehensive computational analysis of

functional consequences of TMPRSS2 SNPs in susceptibility to SARS-CoV-2 among different

populations. *J. Biomol. Struct. Dyn.* 2021;39(10):3576–3593.

28. Jeon S et al. Regional TMPRSS2 V197M Allele Frequencies Are Correlated with COVID-19

Case Fatality Rates. *Mol. Cells* 2021;44(9):680–687.

29. Monticelli M et al. Protective Role of a TMPRSS2 Variant on Severe COVID-19 Outcome in

Young Males and Elderly Women. *Genes* 2021;12(4):596.

30. Li J et al. Polymorphisms and mutations of ACE2 and TMPRSS2 genes are associated with

COVID-19: a systematic review. *Eur. J. Med. Res.* 2022;27(1):26.

31. Schulert GS, Blum SA, Cron RQ. Host genetics of pediatric SARS-CoV-2 COVID-19 and

multisystem inflammatory syndrome in children. *Curr. Opin. Pediatr.* 2021;33(6):549.

32. Oguntuyo KY et al. Quantifying Absolute Neutralization Titers against SARS-CoV-2 by a

Standardized Virus Neutralization Assay Allows for Cross-Cohort Comparisons of COVID-19

Sera. *mBio* 2021;12(1):e02492-20.

33. Nie J et al. Establishment and validation of a pseudovirus neutralization assay for SARS-

CoV-2. *Emerg. Microbes Infect.* 2020;9(1):680–686.

34. Cruz-Cardenas JA et al. A pseudovirus-based platform to measure neutralizing antibodies in

Mexico using SARS-CoV-2 as proof-of-concept. *Sci. Rep.* 2022;12(1):17966.

35. Lam JH et al. Polymersomes as Stable Nanocarriers for a Highly Immunogenic and Durable

SARS-CoV-2 Spike Protein Subunit Vaccine. *ACS Nano* 2021;15(10):15754–15770.

36. Rehman AA, Ahsan H, Khan FH. Alpha-2-macroglobulin: A physiological guardian. *J. Cell.*

Physiol. 2013;228(8):1665–1675.

37. Blanco I. *Alpha-1 Antitrypsin Biology [Internet]*. Elsevier Inc.; 2017:

38. Bornhorst JA, Greene DN, Ashwood ER, Grenache DG. α1-Antitrypsin phenotypes and

associated serum protein concentrations in a large clinical population. *Chest* 2013;143(4):1000–

1008.

39. Sriram Vemuri, C. Tony Yu and NR. Formulation and Stability of Recombinant Alpha-1-

Antitrypsin [Internet]. In: Wang YJ, Pearlman R eds. *Stability and Characterization of Protein*

and Peptide Drugs. Boston, MA: Springer US; 1993:263–285

40. Strnad P, McElvaney NG, Lomas DA. Alpha 1 -Antitrypsin Deficiency. *N. Engl. J. Med.*

2020;382(15):1443–1455.

41. de Serres F, Blanco I. Role of alpha-1 antitrypsin in human health and disease. *J. Intern.*

Med. 2014;276(4):311–335.

42. Mohamed MMA, El-Shimy IA, Hadi MA. Neutrophil Elastase Inhibitors: A potential

prophylactic treatment option for SARS-CoV-2-induced respiratory complications?. *Crit. Care*

2020;24(1):311.

43. Middleton EA et al. Neutrophil Extracellular Traps (NETs) Contribute to Immunothrombosis

in COVID-19 Acute Respiratory Distress Syndrome [Internet]. *Blood* [published online ahead of

print: June 29, 2020]; doi:10.1182/blood.2020007008

44. Barnes BJ et al. Targeting potential drivers of COVID-19: Neutrophil extracellular traps

[Internet]. *J. Exp. Med.* 2020;217(6). doi:10.1084/jem.20200652

45. Olsen GN, Harris JO, Castle JR, Waldman RH, Karmgard HJ. Alpha-1-antitrypsin content in

the serum, alveolar macrophages, and alveolar lavage fluid of smoking and nonsmoking normal

subjects.. *J. Clin. Invest.* 1975;55(2):427–430.

46. Kirsebom FCM et al. COVID-19 vaccine effectiveness against the omicron (BA.2) variant in

England. *Lancet Infect. Dis.* 2022;22(7):931–933.

47. Greene CM et al. α1-Antitrypsin deficiency. *Nat. Rev. Dis. Primer* 2016;2(1):1–17.

48. Yoshikura H. Epidemiological correlation between COVID-19 epidemic and prevalence of

α-1 antitrypsin deficiency in the world. *Glob. Health Med.* 2021;3(2):73–81.

49. Shapira G, Shomron N, Gurwitz D. Ethnic differences in alpha-1 antitrypsin deficiency allele

frequencies may partially explain national differences in COVID-19 fatality rates. *FASEB J.*

2020;34(11):14160–14165.

50. Vianello A et al. Correlation between α1-Antitrypsin Deficiency and SARS-CoV-2 Infection:

Epidemiological Data and Pathogenetic Hypotheses. *J. Clin. Med.* 2021;10(19):4493.

51. Nygren D et al. Low Prevalence of Mild Alpha-1-Antitrypsin Deficiency in Hospitalized

COVID-19-Patients. *Int. J. Gen. Med.* 2022;Volume 15:5843–5848.

52. Faria N, Costa MI, Gomes J, Sucena M. Alpha-1 antitrypsin deficiency severity and the risk

of COVID-19: A Portuguese cohort [Internet]. *Respir. Med.* 2021;181.

doi:10.1016/j.rmed.2021.106387

53. Rodríguez-García C et al. Is SARS-COV-2 associated with alpha-1 antitrypsin deficiency?

[Internet]. *J. Thorac. Dis.* 2023;15(2). doi:10.21037/jtd-22-1062

54. Schneider CV, Strnad P. SARS-CoV-2 infection in alpha1-antitrypsin deficiency [Internet].

Respir. Med. 2021;184. doi:10.1016/j.rmed.2021.106466

55. Köhnlein T, Wilkens M, Eydt K. Auswirkungen der COVID-19-Pandemie auf das

Informationsmanagement und die Therapieadhärenz von substituierten Patienten mit Alpha-1-

Antitrypsin-Mangel (AATM). *Pneumologie* 2022;76(07):494–498.

56. Rodríguez Hermosa JL et al. Severe COVID-19 Illness and α1-Antitrypsin Deficiency:

COVID-AATD Study. *Biomedicines* 2023;11(2):516.

57. SARS-CoV-2 sequences by variant [Internet]. *Our World Data*

https://ourworldindata.org/grapher/covid-variants-bar. cited April 1, 2023

58. de Loyola MB et al. Alpha-1-antitrypsin: A possible host protective factor against Covid-19.

Rev. Med. Virol. 2021;31(2):e2157.

59. Zhou Z et al. Heightened Innate Immune Responses in the Respiratory Tract of COVID-19

Patients. *Cell Host Microbe* 2020;27(6):883-890.e2.

60. Lucas C et al. Longitudinal analyses reveal immunological misfiring in severe COVID-19

[Internet]. *Nature* [published online ahead of print: July 27, 2020]; doi:10.1038/s41586-020-

2588-y

61. Liao M et al. Single-cell landscape of bronchoalveolar immune cells in patients with

COVID-19. *Nat. Med.* 2020;26(6):842–844.

62. Bryan CL et al. HIV infection is associated with reduced serum alpha-1-antitrypsin

concentrations. *Clin. Investig. Med. Med. Clin. Exp.* 2010;33(6):E384-389.

63. Grifoni E et al. Interleukin-6 as prognosticator in patients with COVID-19 [Internet]. *J.*

Infect. [published online ahead of print: June 2020]; doi:10.1016/j.jinf.2020.06.008

64. McElvaney OJ et al. Characterization of the Inflammatory Response to Severe COVID-19

Illness. *Am. J. Respir. Crit. Care Med.* 2020;202(6):812–821.

65. Janciauskiene S, Welte T. Well-Known and Less Well-Known Functions of Alpha-1

Antitrypsin. Its Role in Chronic Obstructive Pulmonary Disease and Other Disease

Developments. *Ann. Am. Thorac. Soc.* 2016;13(Supplement_4):S280–S288.

66. Jose RJ, Manuel A. COVID-19 cytokine storm: the interplay between inflammation and

coagulation. *Lancet Respir. Med.* 2020;8(6):e46–e47.

67. Giannis D, Ziogas IA, Gianni P. Coagulation disorders in coronavirus infected patients:

COVID-19, SARS-CoV-1, MERS-CoV and lessons from the past [Internet]. *J. Clin. Virol.*

2020;127. doi:10.1016/j.jcv.2020.104362

68. Azouz NP et al. Alpha 1 Antitrypsin is an Inhibitor of the SARS-CoV-2–Priming Protease

TMPRSS2. *Pathog. Immun.* 2021;6(1):55–74.

69. McElvaney OJ et al. A randomized, double-blind, placebo-controlled trial of intravenous

alpha-1 antitrypsin for ARDS secondary to COVID-19. *Med* 2022;3(4):233-248.e6.

70. Meseeha M, Attia M. Alpha 1 Antitrypsin Deficiency [Internet]. In: *StatPearls*. Treasure

Island (FL): StatPearls Publishing; 2023:

71. Black L, Kueppers F. alpha1-Antitrypsin deficiency in nonsmokers.. *Am. Rev. Respir. Dis.*

1978;117(3):421–428.

72. Harbig A et al. Transcriptome profiling and protease inhibition experiments identify

proteases that activate H3N2 influenza A and influenza B viruses in murine airways. *J. Biol.*

Chem. 2020;295(33):11388–11407.

73. Böttcher-Friebertshäuser E et al. Cleavage of Influenza Virus Hemagglutinin by Airway

Proteases TMPRSS2 and HAT Differs in Subcellular Localization and Susceptibility to Protease

Inhibitors. *J. Virol.* 2010;84(11):5605–5614.

74. Potthoff AV, Münch J, Kirchhoff F, Brockmeyer NH. HIV infection in a patient with alpha-1

antitrypsin deficiency: a detrimental combination?. *AIDS Lond. Engl.* 2007;21(15):2115–2116.

75. Zhou X et al. Alpha-1-antitrypsin interacts with gp41 to block HIV-1 entry into CD4+ T

lymphocytes. *BMC Microbiol.* 2016;16(1):172.

76. Peitsch C, Klenk H-D, Garten W, Böttcher-Friebertshäuser E. Activation of Influenza A

Viruses by Host Proteases from Swine Airway Epithelium. *J. Virol.* 2014;88(1):282–291.

77. Hatesuer B et al. Tmprss2 Is Essential for Influenza H1N1 Virus Pathogenesis in Mice.

PLOS Pathog. 2013;9(12):e1003774.

78. Buchholz UJ, Finke S, Conzelmann KK. Generation of bovine respiratory syncytial virus

(BRSV) from cDNA: BRSV NS2 is not essential for virus replication in tissue culture, and the

human RSV leader region acts as a functional BRSV genome promoter. *J. Virol.*

1999;73(1):251–259.

79. Ramsburg E et al. A Vesicular Stomatitis Virus Recombinant Expressing Granulocyte-

Macrophage Colony-Stimulating Factor Induces Enhanced T-Cell Responses and Is Highly

Attenuated for Replication in Animals. *J. Virol.* 2005;79(24):15043–15053.

80. Case JB et al. Neutralizing Antibody and Soluble ACE2 Inhibition of a Replication-

Competent VSV-SARS-CoV-2 and a Clinical Isolate of SARS-CoV-2. *Cell Host Microbe*

2020;28(3):475-485.e5.

81. Beaty SM et al. Efficient and Robust Paramyxoviridae Reverse Genetics Systems. *mSphere*

2017;2(2):e00376-16.

82. Amanat F et al. A serological assay to detect SARS-CoV-2 seroconversion in humans. *Nat.*

Med. 2020;26(7):1033–1036.

83. Lopes AP et al. Portuguese consensus document for the management of alpha-1-antitrypsin

deficiency. *Pulmonology* 2018;24 Suppl 1:1–21.

Alpha-1-antitrypsin and its variant-dependent role in COVID-19 pathogenesis

Christian S Stevens[1#], Kasopefoluwa Y Oguntuyo[1#], Shreyas Kowdle[1], Luca Brambilla[1], Griffin Haas[1], Aditya Gowlikar[1], Mohammed NA Siddiquey[2], Robert M Schilke[2], Matthew D Woolard[2], Hongbo Zhang[2], Joshua A Acklin[1], Satoshi Ikegame[1], Chuan-Tien Huang[1], Jean K Lim[1], Robert W Cross[3], Thomas W Geisbert[3], Stanimir S Ivanov[2], Jeremy P Kamil[2], the Alpha-1 Foundation[4], and Benhur Lee[1]

A.

B.

C.

D.