The specific features of the developing T cell compartment of the neonatal lung

are a determinant of respiratory syncytial virus immunopathogenesis

Thomas Démoulins[1,2], Melanie Brügger[1,2,3], Beatrice Zumkehr[1,2], Blandina I. Oliveira

Esteves[1,2], Kemal Mehinagic[1,2,3,4], Amal Fahmi[1,2,3], Loïc Borcard[1,2,3], Adriano Taddeo[1,2], Horst

Posthaus[4,5], Charaf Benarafa[1,2], Nicolas Ruggli[1,2], and Marco P. Alves[1,2,*]

[1]The Institute of Virology and Immunology, Mittelhäusern and Bern, Switzerland

[2]Department of Infectious Diseases and Pathobiology, Vetsuisse Faculty, University of Bern,

Bern, Switzerland

[3]Graduate School for Cellular and Biomedical Sciences, University of Bern, Bern, Switzerland

[4]Institute of Animal Pathology, Department of Infectious Diseases and Pathobiology,

Vetsuisse Faculty, University of Bern, Bern, Switzerland

[5]COMPATH, Vetsuisse Faculty & Faculty of Medicine, University of Bern, Bern, Switzerland

[*]Corresponding Author: marco.alves@vetsuisse.unibe.ch

Short Title: Atypical neonatal immune response to RSV infection

Descriptor number: 10.09 Pathogen/Host Cell Interactions

Word count abstract: 204

Manuscript word count: 5671

References: 61

Figures: 6

Supplementary material: 5 figures, 2 tables

ABSTRACT

The human respiratory syncytial virus (RSV) is a major cause of severe lower respiratory tract

infections in infants, possibly due to the specific features of the immature neonatal pulmonary

immune system. Using the newborn lamb, a classical model of human lung development and

a state-of-the-art model of RSV infection, we aimed to explore the role of cell-mediated

immunity in RSV disease during early life. Remarkably, in healthy conditions, the developing

T cell compartment of the neonatal lung showed major differences to that seen in the mature

adult lung. The most striking observation being a high baseline frequency of bronchoalveolar

IL-4-producing CD4 and CD8 T cells, which declined progressively over developmental age.

RSV infection exacerbated this pro-type 2 environment in the bronchoalveolar space, rather

than inducing a type 2 response *per se*. Moreover, regulatory T cell suppressive functions

occurred very early to dampen this pro-type 2 environment, rather than shutting them down

afterwards, while $\gamma\delta$ T cells dropped and failed to produce IL-17. Importantly, RSV disease

severity was related to the magnitude of those unconventional bronchoalveolar T cell

responses. These findings provide novel insights in the mechanisms of RSV

immunopathogenesis in early life, and constitute a major step for the understanding of RSV

disease severity.

Keywords: Respiratory syncytial virus; neonatal immunity; bronchiolitis; interstitial

pneumonia, immunopathogenesis; Cell-mediated immunity, T cells.

AUTHOR SUMMARY

By using a state-of-the-art translational model with full accessibility to the small airways at

defined early life periods, we provide an unpreceded characterization of the developing T cell

compartment in the distal lungs of healthy and RSV-infected neonates. This process is highly

dynamic and tightly regulated, characterized by colonizing T-cell subsets that synergize

towards a narrow pro-tolerogenic immunological window. We believe our work constitutes a

solid basis to clarify the age dependency of RSV immunopathogenesis, and should be

considered in vaccine design, which remains challenging after five decades of effort.

INTRODUCTION

The human respiratory syncytial virus (RSV) is a seasonal virus, known as the major cause of lower respiratory tract infection during early childhood (1, 2). Worldwide, RSV leads to around 160'000 deaths each year (3). Although there are risk factors for RSV severity such as prematurity and congenital heart disease, the majority of hospitalized infants are previously healthy (4-7), suggesting that the disease is partially linked to inherent properties of early life immunity. The precise understanding of the immune-driven susceptibility of neonates is essential, considering there is no licensed vaccine and available prophylactic treatments are mostly based on palivizumab, a neutralizing antibody used in high-risk group infants (8).

The first exposure to pathogens occurs during the early postnatal period and is critical for lung colonization by immune cells. There is increasing evidence that the overall neonatal T-cell compartment is distinctive during this temporal window (9). Upon stimulation neonatal T-cells tend to differentiate into regulatory T-cells (Tregs), to facilitate self-tolerance to developing organs (10). Also, the developing lung is characterized by a type 2 immune phenotype, and the pulmonary T-cell responses present a T helper type 2 (Th2) bias (11, 12). Remarkably, neonatal $\gamma\delta$ T-cells, a T-cell subset enriched at mucosal barrier sites, have an impaired capacity to respond to stimulation (13-16).

Due to the aforementioned reasons, young infants do not mount a classical pulmonary immune response to pathogens such as RSV, which may contribute to high morbidity and mortality (17, 18). A likely explanation could be a combined contribution of viral load (19) and an inappropriate and/or dysfunctional immune response, through an over-exuberant inflammatory response or a biased T-cell response (9, 20, 21). Neonatal T-cells may have both protective and harmful effects in responses to RSV infection. While children with T-cell deficiencies cannot efficiently clear the virus (22, 23), it remains controversial whether the severity of RSV disease is associated with a bias towards type 2 cytokines such as IL-4 (24-26).

Knowledge of the course of RSV disease is still limited due to ethical and technical difficulties

associated with the investigation of the immune response in human neonates. The neonatal

lamb model of human RSV infection recapitulates most of the features of the paediatric

disease. Furthermore, due to similarities in development and pulmonary immune system, the

ovine lung is a classical model of the human respiratory tract (27). Human RSV-infected

animals clinically present rhinitis, coughing and respiratory distress. Typical pulmonary lesions

are bronchiolitis and interstitial pneumonia associated with inflammatory infiltrates (28-30).

Herein, this model, immunologically related to human, was exploited to have full access to the

small airways and follow the development of the pulmonary immune cell features in early life.

Moreover, by peribronchial lymph nodes (LNs), lung, and bronchoalveolar space sampling

following RSV infection, we provided an unprecedented delineation of the contributors of the

host neonatal T cell compartment over the course of RSV disease.

RESULTS

RSV infection is causing lower respiratory tract disease and leading to a potentially

protective humoral response

To evaluate the immune responses during the acute, recovery and convalescence phases of

RSV disease, time-points were Selected as follow: 3 and 6 days post-infection (p.i.), to

evaluate the innate and early adaptive responses; 14 days p.i. to assess the adaptive

mechanisms and virus clearance; 42 days p.i. to investigate the impact of RSV infection on

the immune cell colonization of the developing lung. As controls, non-infected newborns and

adults were used (Figure 1A). We quantified RSV in BALs (cellular and fluid fractions) and in

the lung tissue. Although RSV clearance was slower in the cellular fraction of BALs, it became

undetectable in any sample at day 42 p.i., demonstrating an efficient clearance (Figure 1B).

Immunohistochemistry of lung tissue sections revealed the presence of RSV-antigens in type

I pneumocytes and macrophages (Figure 1C). Clinically, RSV infected neonates showed signs

of rhinitis with clear nasal discharge, occasional coughing and wheezing. Macroscopical

lesions were visible at days 3-6 p.i, including failure of pulmonary collapse and focal, dark-red areas of subpleural pulmonary consolidation and atelectasis. Histological lesions, present at days 3-6 p.i consisted of multifocal-coalescing areas of lymphocytic and histiocytic infiltrates within interalveolar septae (interstitial pneumonia) and accumulation of inflammatory cells within alveolar spaces and bronchioles (bronchiolitis) (Figure 1D). Longitudinal measurements of white blood counts and neutrophil counts in the circulation showed no difference between mock and infected neonates (Figure S1). Next, we monitored the presence of RSV neutralizing antibodies (NAbs) (Figures 1E, S2). In the serum of healthy adults and mock neonates, no RSV NAbs were detected. In contrast, RSV-infected neonates have naturally acquired NAbs at day 13-14 p.i., except one animal. Notably, two siblings had RSV NAbs preceding infection; this was passively transmitted by their mother whose serum also had a strong RSV-neutralization (Figure S3). We extended the assay to later time points to test the persistence of RSV neutralization. No loss of humoral memory was measured up to 42 days p.i (Figure 1F). Altogether, these results indicate that neonates can mount an efficient humoral response suggesting potential protection for a subsequent RSV infection. When quantifying the cytokine profiles in BALs from mock control and RSV-infected animals, the most noticeable induction upon infection were IL-10 (day 6 p.i.), IL-8 and CXCL10/IP-10 (days 3 and 6 p.i.). The latter being a potent lymphocyte chemoattractant (Figure 1G).

Neonatal RSV infection leads to a transient depletion of $\gamma\delta$ T cells and a deficiency of IL-17-mediated immunity

We first assessed central components of the pulmonary immune responses bridging innate and adaptive responses, namely plasmacytoid DCs (pDCs) and $\gamma\delta$ T cells. In RSV disease, pDCs are known to be involved during the early stages of the pulmonary immune response and have a protective function by modulating the local immune microenvironment (31). When quantified in the BALs, pDC numbers of mock neonates were similar to those of healthy adults. Upon RSV infection, we observed a trend to pDC recruitment as early as 3 days p.i., and this

was enhanced at day 6 p.i. (p<0.05). This recruitment was specific of bronchoalveolar space,

since we failed to detect any increase in peribronchial LN and lung tissue. However, the pDC

recruitment in the bronchoalveolar space was transitional and tightly regulated, since day 14

p.i corresponded to a return to steady-state levels (Figure S4A, S5A). To verify whether the

extent of pDC recruitment was linked to the magnitude of RSV shedding, pDC counts were

plotted as a function of RSV copies in the fluid fraction of BALs. A significant association was

found (Figure S5B). We next evaluated the maturation and activation phenotype of recruited

pDCs. A significant CD86 upregulation was found at days 3-6 p.i. (Figure S5C, D). This

upregulation was transitional and back to steady-state levels at day 14 p.i. In line with the early

pDC recruitment and activation, we observed a significant upregulation of IFN-β during the

acute phase of neonatal RSV infection, suggesting the induction of an antiviral state in the

respiratory tract (Figure S5E).

Similar to pDCs, $\gamma\delta$ T cells link the innate and adaptive arms of the immune system, particularly

at mucosal sites including the respiratory tract. Thus, we wondered whether they were

involved in the early immune response to RSV infection (32). $\gamma\delta$ T cell numbers in the

peribronchial LNs, lung tissue and BALs of mock neonates were comparable to those

calculated in healthy adults. However, neonatal RSV infection led to a noticeable decrease in

$\gamma\delta$ T cell frequency in the BALs at day 14 p.i., whereas this remained stable in other lung

compartments. Nevertheless, this $\gamma\delta$ T cell depletion was entirely reversed at day 42 p.i.

(Figure 2A, S4A). To exclude the possibility that the $\gamma\delta$ T cell reduction could be due to

migration or relocation, we calculated their counts as a function of the percentage of

Live/Dead-positive cells. A negative association was found, suggesting that neonatal $\gamma\delta$ T cells

death is linked to the presence of RSV in the bronchoalveolar space (Figure 2B). We next

ascertained whether neonatal $\gamma\delta$ T cells in BALs remained immature during the recovery

phase. Remarkably, in healthy adults, half of the $\gamma\delta$ T cell pool was composed of mature CD25$^+$

cells, whereas the latter were absent in mock neonates at day 3. The proportion of mature $\gamma\delta$

T cells progressively strengthened over time, without reaching the level detected in healthy

adults. RSV infection exerts an effect on this maturation of neonatal $\gamma\delta$ T cells. Not only the

observed depletion was rapidly followed by a complete replenishment at day 42 p.i., but also

the restoration was accompanied by a marked increase in the percentage of CD25-positive

cells, compared to mock neonates (Figure 2C, D). Finally, we demonstrated the inability of

neonatal $\gamma\delta$ T cells to produce detectable levels of IL-17 (Figure 2E).

Neonatal RSV infection induces a fast and massive Tc2 influx in the bronchoalveolar

space

Assessment moved on to adaptive immune response, and more particularly to IL-4 *versus*

IFN-γ production by CD4$^+$ or CD8$^+$ T cells, a determining factor in the outcome of protective

versus harmful immune response to RSV infection (25, 33) (Figure 3A, S4C). In BALs, mock

neonates had a very low number of CD4$^+$ and CD8$^+$ T-cells compared to healthy adults,

whereas numbers were comparable in draining LNs. RSV infection led to their strong

recruitment in the bronchoalveolar space, albeit with distinct kinetics. An unexpected reduction

of CD4$^+$ T-cell recruitment was seen as soon as 14 days p.i. This contradicted the values

obtained for CD8$^+$ T-cells, where the significant enhancement following RSV infection was

maintained. Interestingly, this influx of CD8$^+$ T cells was preceded by their significant increase

in the lung, indicating that they transit through this compartment. When we considered both

CD4$^+$ and CD8$^+$ fractions at a later time-point, the numbers were comparable to those of mock

neonates (Figure 3B). Surprisingly, mock neonates had high baseline frequencies of Th2 cells

in BALs, which declined progressively over time. Elevated amounts of spontaneous produced

IL-4 were a signature of the developing CD4$^+$ fraction, since it was no longer detectable at day

42, or in healthy adults. Upon infection, this analysis revealed a trend towards an increase in

the frequency of those Th2 cells, whereas there was no consistent influence on IFN-γ induction

(Figures 3C, D). Consequently, RSV infection in early life exacerbates the overall lung pro-

Th2 cytokine environment, rather than inducing a Th2 immune response *per se*.

High baseline frequencies of IL-4-expressing CD8$^+$ T-cells (Tc2) were also found in BALs of

in mock neonates. Again, a progressive decline of Tc2 cells occurred up to day 42 p.i.,

comparable to negligible levels measured in healthy adults. RSV disease induced a greater

overall Tc2-derived cytokine environment in BALs. This exacerbation was tightly regulated,

reduced to a level similar to healthy adults at day 14 p.i. Again, no consistent influence of RSV

infection was seen on IFN-γ induction (Figure 3E, F). Altogether, these results highlight the

role of Tc2 cells as an important source of IL-4 production during neonatal RSV infection.

Finally, we then tested the activation and differentiation of naïve T lymphocytes into memory

subsets. The most striking observation was achieved at day 42 p.i.; the percent contributions

of different subsets seen in RSV-infected neonates was highly superimposable to that found

in healthy adults. This was not the case for mock neonates who had lower frequencies of

activated memory T-cells, particularly for the CD8$^+$ fraction (Figure 3G). Consequently, RSV

infection in early life modulates the activation and the maturation of the T-cell pool.

Nonetheless, it is unknown whether this precipitated activation/maturation is RSV epitope-

specific and a predictive factor for severe outcomes in the context of a secondary RSV

infection (34).

Neonatal RSV infection rapidly induces Treg suppressive functions to dampen the Th2

and Tc2 responses

The neonatal immune system is tolerogenic, a mechanism that promotes self-tolerance to

developing organs; fetal CD4$^+$ T cells are more likely to differentiate into Tregs after stimulation

(10). The shortened kinetics observed for CD4$^+$ T-cells was elaborated by determining whether

this was due to an early Treg rather than a classical CD4$^+$ T-cell response (Figure 4A, S4D).

RSV infection elicited a rapid Treg recruitment in BALs that mirrored the one previously

observed for CD4$^+$ T-cells. This Treg dynamic is specific for the respiratory mucosa, since

Treg frequencies remained unchanged in the lung tissue (Figure 4A). We then noticed a trend

for Treg subset overrepresentation following RSV infection (Figure 4B). The proportion of

Tregs declined progressively over time, tending towards similar levels to those detected in

healthy adults. RSV infection promoted enhancement of TGF-β production by Tregs, however

not significant (Figure 4C). When Treg counts were calculated as a function of non-Treg CD4$^+$

counts, an association was found, suggesting that CD4$^+$ T-cell pool tends to differentiate into

Tregs. When Treg counts were calculated as a function of CD8$^+$ counts, an association was

still found, confirming that CD8$^+$ T-cells and Tregs colonized the alveolar space in a

coordinated manner. By plotting CD8$^+$ count as a function of non-Tregs CD4$^+$ counts, it

appeared that T-cell response was mainly driven by the CD8$^+$ fraction (high numbers at days

6 and 14 p.i.) (Figure 4D).

Neonates and adults are mounting a distinct pulmonary T cell response following RSV

infection

We followed the same approach to dissect the cellular immune response in RSV-infected

adults. While the viral loads and virus clearance was comparable between adults and

neonates (Figure 5A), respiratory lesions were absent in adults (Figure 5B); this suggests that

the increased rate of severe disease reported in infants is not due to a failure to control the

virus, reinforcing the hypothesis for immune-mediated RSV pathogenesis.

The rapid and robust pDC recruitment observed in infected neonates did not occur in infected

adults. However, lung-resident pDCs of the latter were prompt to react, displaying an even

more mature and activated phenotype Importantly, this mature pDC response was also tightly

regulated (Figure 5C, D). The strong decrease in γδ T-cell counts observed in infected

neonates was absent in adults. Moreover, adult γδ T-cells produced high amounts of IL-17,

suggesting that IL-17-mediated immunity is deficient in early life (Figure 5E, F) (15). This IL-

17 response was accompanied by an activated phenotype, as indicated by the enhanced

proportion of CD25$^+$ cells (Figure 5G). From these results, we could ascertain that neonates

lack a mature and functional γδ T-cell pool in the lower airways. For the CD4$^+$ subset, no

significant recruitment was found (Figure 5H). The percentage of IL-4-producing cells was

very low, and RSV infection had no consistent influence on IFN-γ induction (Figure 5I). Another

clear difference was the contraction of effector memory pool (CD25$^+$CD45RO$^+$) at day 14 p.i.,

indicative of a specific immune response (Figure 5J). The recruitment of CD8$^+$ T-cells was

delayed compared to neonates, only detectable at day 14 p.i. (Figure 5K). Again, the

percentage of IL-4-producing cells was low (Figure 5L), whereas the precipitous contraction

of effector memory pool at day 14 p.i. suggested a RSV-specific CD8$^+$ T-cell response (Figure

5M). From these results, it became clear that the inaptitude of neonates to mount an effector

T-cell response is tentatively substituted by a fast and massive Th2/Tc2 cell recruitment.

Finally, we failed to detect any Treg recruitment. However, the rare tissue-resident Tregs were

able to produce TGF-β, suggesting that an immune regulation happened (Figure 5N, O).

The severity of RSV disease relates to the magnitude of the bronchoalveolar T cell

responses

The pulmonary T-cell response was evaluated in neonates infected with RSV-ON1-H1 strain,

which induced a milder disease (Figure 6A). Interestingly, the viral loads and virus clearance

was comparable between both strains, again reinforcing the hypothesis for immune-mediated

RSV pathogenesis (Figure 6B). Effectively, all T cell-related responses were attenuated. We

did not observe a γδ T-cell depletion, and the CD8$^+$ T-cell recruitment was delayed.

Furthermore, the Treg recruitment was lower compared to RSV A2 infection (Figure 6C). By

combining results obtained with both strains, we found a significant association between Treg

and pDC counts in BALs (Figure 6D). This indicates that, despite their skew, all distinct

subsets of neonatal T-cells cooperate tightly with the magnitude of the cellular immune

response against RSV infection. Altogether, our data clarified how neonates elicit an over-

exuberant pulmonary T cell response characterized by a bronchoalveolar influx of Treg, Th2,

and Tc2 cells, contrasting to a classical T cell response.

DISCUSSION

To properly assess the immunopathogenesis of RSV, it is essential to select a model that

recapitulates the features of human disease. We selected the neonatal lamb, as RSV is a

natural pathogen for this species, lung development is similar, and the key features mimic

human RSV infection in infants (28, 29). One strength of our experimental approach was the

ease sampling and the full accessibility to the lower airways throughout the disease when

sampling in infants is mainly restricted to the upper airways, the latter representing only an

approximate surrogate of the distal lung where RSV disease occurs. Moreover, we offer an

unprecedented view of the development of pulmonary neonatal immunity in healthy

conditions. This process is highly dynamic and tightly regulated, with a short and early stage

where colonizing T-cell subsets synergize towards a narrow pro-tolerogenic window, namely

a "Treg/Th2/Tc2" environment. This certainly contributes to the maturation of the immune

system and may constitute an unsuitable state to resolve RSV infection.

The mechanisms underlying the severity of disease are still unclear, although RSV itself was

shown to induce distal lung obstruction by shedding of necrotic airway epithelial cells (35).

Also, it is unclear yet if RSV disease severity is associated with viral loads (36, 37). Here we

demonstrate that while RSV disease was more severe in neonates compare to adult animals,

RSV loads and virus clearance was comparable for both age groups. In line with this, RSV

loads and virus clearance were similar for neonates infected with RSV strains inducing an

RSV disease with distinct severity. Clearly, our data strongly reinforce the hypothesis of an

inappropriate neonatal immune response as a determinant of RSV disease severity. We

observed a rapid and robust recruitment of neonatal pDCs, whose extent correlated with the

viral loads of infected neonates. Infected adults did not require this recruitment, since their

lung-resident pDCs reacted promptly by maturing to a higher level than that measured in

neonates. This partially corroborates recent studies reporting low counts and maturation levels

of pDCs in mice and infants (38, 39). This reduced capacity to respond against RSV might be

due to a direct disturbance of signaling by RSV itself (40), or an intrinsic deficiency of neonatal

pDCs to produce elevated amounts of cytokines (41-43). This neonatal defect would prevent

conventional DCs from adequate presentation of viral antigens, contributing to poor specific T

cell responses (44).

Controversial is whether γδ T-cells participate in viral clearance or immunopathology. Herein,

the γδ T-cell depletion conflicted with a previous report. Indeed, children with severe RSV

bronchiolitis had reduced frequencies of γδ T-cells in peripheral blood, which was explained

by the authors as a likely redistribution towards the lungs; unfortunately, they could not

evaluate their hypothesis (45). In line with our findings, mice depleted of γδ T-cells before RSV

infection display increased viral titers (13). An alternative hypothesis that would reconcile our

study with others, is that γδ T-cell tissue redistribution plays less of a role than its immature

state in early life. Peripheral blood γδ T-cells of infants with severe disease failed to produce

IFN-γ when restimulated (14), whereas γδ T-cells from neonatal mice had an impaired ability

to produce IL-17 (15). Finally, Tc17 and Th17 cells were recently associated with shorter

hospital stays and proposed to play a protective role (26). Altogether, our data demonstrate

that the protective role of γδ T-cells in RSV infection is largely inefficient in early life.

The BALs of healthy neonates displayed a high proportion of Tregs that declined rapidly to

levels comparable to healthy adults. As fetal CD4$^+$ T-cells tend to differentiate preferentially

into Tregs after stimulation (10), this tolerogenic mechanism is likely to promote self-tolerance

to the developing lung. We found that Treg response to RSV infection happened very early in

neonates, accompanying CD4$^+$ and CD8$^+$ T-cell expansion rather than shutting them down

afterwards. This strong Treg recruitment was absent in adults. We speculate that the neonate

immune system faces a dilemma between avoiding potential autoimmune disorders and failing

to mount a specific T-cell response to harmful pathogens. Studies conducted in young infants

with RSV disease support this statement. A selective depletion of peripheral Tregs was shown,

probably due to massive recruitment to the lungs (46). A higher level of TGF-β transcript was

measured in neonatal DCs compared to adults (47). In contrast, some reports emphasized

the protective role of Tregs, since Treg-depleted young mice had an enhanced RSV disease,

with abundant T-cells exhibiting an activated phenotype (48). Another remarkable observation

was the strong correlation between pDC and Treg counts. Despite the need for further

investigations, it is worth noting that pDC to Treg ratio was linked to the clearance *versus*

persistence of human papillomavirus (49).

Whether the cellular arm of the adaptive pulmonary response is detrimental or beneficial in

RSV disease is still debated. Particularly, the contribution of IL-4 to viral clearance versus

immunopathogenesis remains controversial (25, 50-52). Recently, a study performed in

hospitalized infants showed a possible involvement of Tc2 cells in disease severity. Although

convincing, this study was for obvious ethical reasons restricted to nasal aspirates (26).

Interestingly, Tc2 cells are increasingly proposed to play an important role in the etiology of

asthma (53). Herein, we provide a significant advancement in understanding this yet unclear

IL-4 contribution, by showing that type 2 cytokine environment in the lung is a footprint of early

life that decreases rapidly with time in healthy conditions. RSV infection circumvents this

decline by mediating the massive recruitment of $CD4^+$ and $CD8^+$ T-cells, exacerbating the

overall type 2 cytokine environment. We think that the link between IL-4 and RSV disease is

the direct consequence of a strong trafficking of endogenous Th2 and Tc2 lymphocytes

towards infected pulmonary tissues, rather than an RSV-specific type 2 immune response *per*

se.

The high proportion of pulmonary T-cells producing spontaneously IL-4 was undetectable in

the healthy adults. Neither was the RSV-specific immune response of adults oriented towards

a Th2 or Tc2 profile. This corroborates a previous report mentioning an endogenously poised

cytokine profile toward IL-4 in $CD4^+$ T-cells in cord blood of neonates and infants, establishing

a link between IL-4 and development (54). However, we provide new insights by

demonstrating that i) $CD8^+$ Tc2 cells are an equal and possibly superior source of IL-4 than

$CD4^+$ Th2 cells; ii) this immunological state occurs during a short period *in vivo*, raising the

likelihood that numerous animal trials failed to detect it due to the study design. Remarkably,

this transient type 2 cytokine environment was exacerbated upon RSV infection. So far, similar

observations were only made at the systemic level or in nasal aspirates, and authors could

only speculate on the relocation to the lower respiratory tract (26, 55). Herein, not only we

show that this happens, but we also unify the link between IL-4 and RSV disease and

endogenous IL-4 program in neonates. This finding constitutes an important step towards

deciphering why neonates and young infants are particularly at risk.

Conclusively, we provid novel insights into the identification of the host's contributing factors

to RSV immunopathology, strengthening the hypothesis of an inappropriate immune response

in the narrow temporal window of early life. This can now be characterized as a combined

bronchoalveolar influx of Treg, Th2 and Tc2 cells, associated with a strong depletion of $\gamma\delta$ T

cells and IL-17 deficiency. These findings constitute a novel basis for further exploration of

RSV immunopathogenesis in infants and should be considered in RSV vaccine design, which

remains challenging after five decades of effort.

MATERIAL AND METHODS

Virus

Human RSV-A2 strain was obtained from the American Type Culture Collection (ATCC, VR-

1540, GenBank accession number KT992094.1). The strain RSV-ON1-H1 was isolated from

nasopharyngeal aspirates of a child below the age of five years with confirmed RSV infection,

hospitalized at Hannover Medical School, Germany. The caregivers gave informed consent

for the donation of nasopharyngeal aspirates, and all steps were conducted in compliance with

good clinical and ethical practice and approved by the local ethical committee at Hannover

Medical School (permission number 63090/2012) (56). Both RSV strains were propagated on

HEp-2 cells (ATCC, CCL-23) cultured in DMEM (Gibco) supplemented with 5% FBS (Gibco),

at 37 °C in 5% CO_2 atmosphere. Briefly, viruses were incubated for 36-44 h in tissue culture

flasks at a multiplicity of infection (MOI) of 0.02 plaque-forming unit (PFU) per cell. When the

cytopathogenic effect (CPE) reached approximately 60%, the cultures were frozen overnight

at −70 °C, to disrupt the Hep-2 cells and release the virus. Then, the T-175 cm2 flasks were

thawed, HEp-2 cells were scraped and the complete contents of the flask were spun down at

1000 g, 10 min, 4 °C to collect cell debris. The clarified supernatants were layered over 20%

sucrose (w/v) solution prepared in sterile PBS and ultra-centrifuged for 3 h 30 min at 71'000 g at 4 °C. The resultant RSV stocks were aliquoted and stored at −150 °C until further use. Virus titers were determined on HEp-2 cells, using biotinylated anti-RSV antibody (goat polyclonal IgG, Bio-Rad) for 1 h at RT. The plaques were visualized using ExtrAvidin®-Peroxidase (Sigma), 30 min at RT, and revealed by DAB substrate (Sigma), left for up to 30 min at RT. When the plaques appeared with the desired coloration, the reaction was stopped with PBS. RSV titers were expressed as PFU per ml.

Animals

The experiments in lambs were performed in compliance with the Swiss animal protection law (TSchG SR 455; TSchV SR 455.1; TVV SR 455.163) under the authorization BE125/17. The experiments were reviewed by the cantonal committee on animal experiments of the canton of Bern, Switzerland, and approved by the cantonal veterinary authority (Amt für Landwirtschaft und Natur LANAT, Veterinärdienst VeD, Bern, Switzerland). For the duration of the experiment, the animals were housed in the high containment animal facility. *Postpartum*, the acclimatization lasted 3 to 6 days before RSV infection. In accordance with the principles of the 3Rs, suckling lambs were held together with their adult mothers and fed *ad libitum*. The animals were kept on straw with a local ambient temperature of 20-22 °C. In order to prevent secondary bacterial infection, all the animals received a prophylactic long-acting oxytetracycline treatment starting at 18 to 24 h before RSV infection by intramuscular injection of Cyclosol LA, 20 mg/kg (Dr. E. Graeub AG), repeated after 4 days. The developmental age of sheep does not follow a linear curve compared to humans but in terms of lung development, animals euthanized at day 3, 6, and 14 post-infection (p.i.) can be considered as infants and at day 42 as adolescents (57). The transtracheal RSV inoculation was performed under sedation and analgesia by intramuscular injection of a mixture of midazolam (Dormicum®, Roche) 0.2 mg/kg and butorphanol (Morphasol®, Dr. E. Graeub AG) 0.2 mg/kg. The injection site was shaved and disinfected with 70% ethanol. The trachea was punctured between the proximal tracheal rings with a 0.9 x 40 mm needle (20G, BD) and a

volume of 2 ml of virus (RSV-infected animals: 10^8 PFU for RSV-A2; 10^7 PFU for RSV-ON1-

H1) or PBS (mock-treated controls) were injected slowly. Before and after infection, we

monitored body temperature, and clinical status daily. The clinical status of the lambs was

assessed by a veterinarian, whenever possible always the same person to ensure unbiased

clinical assessment. The rectal body temperature was measured with a digital thermometer,

respiratory and heart rates with a stethoscope.

Histopathology and Immunohistochemistry

Samples of each lung lobe were fixed in 10% buffered formalin for 48 h, embedded in paraffin

and routinely processed for histology. Three μm Histological sections were stained with

hematoxylin and eosin. For Immunohistochemistry, unstained 3 μm sections were

deparaffinized with xylol for 5 minutes followed by rehydration in descending concentrations

of ethanol (100, 95, 80, and 75%). Endogenous peroxidase activity was inhibited by H_2O_2

(3.25% in methanol, 10 min at RT). Then, heat-induced antigen retrieval was carried out by

incubating the slides in boiling citrate buffer (pH 6.0) for 10 min. Nonspecific binding of

antibody was blocked with 1% BSA for 10 min. Slides were incubated overnight at 4° with

primary antibodies diluted 1:200 in PBS (clone 302, ThermoFisher). LSAB and AEC Kits

(DakoCytomation) were used as indicated by the manufacturer for secondary antibody

incubation and signal detection. Slides were counterstained with Ehrlich hematoxylin and

cover slips were mounted using Aquatex (Merck) (58).

Cell suspension preparation

The isolation of cells from peribronchial lymph nodes was done using a Collagenase D (Sigma)

and DNAse I (BioConcept) enzyme mix and the gentleMACS Octo Dissociator (Miltenyi

Biotec). For the isolation of immune cells from the lung tissue, we used a Collagenase I

(BioConcept), Collagenase II (BioConcept) and DNase I (BioConcept) enzyme mix. The

dissociated tissues were passed through a 100 μm cell strainer (Falcon) and centrifuged at

250 g for 10 min, 4 °C. Next, the cell pellets were resuspended, passed through a 70 μm cell

strainer (Falcon) to ensure the removal of clumps, and centrifuged 250 g for 10 min, 4 °C. For

the lung suspensions, the cells were further purified using the Ficoll-Paque approach (Sigma).

For the isolation of cells from bronchoalveolar lavages (BALs), the lungs were taken out with

the trachea, which was clamped before cutting, to prevent blood from entering the lungs. Then,

PBS was poured through the trachea using a sterile funnel. The single-cell suspensions were

applied to a sieve to remove any remaining larger particles. Depending on the BAL volume,

single-cell suspensions were distributed in the appropriate number of 50 ml Falcon tubes for

centrifugation at 350 g for 10min, 4°C. Then, the pellets were resuspended, passed through a

100 µm cell strainer (Falcon) and centrifuged at 350 g for 10 min, 4 °C. Next, the pellets were

resuspended, passed through a 70 µm cell strainer (Falcon) to ensure the removal of clumps,

and centrifuged at 350 g for 10 min, 4 °C. If needed, red blood cells were lysed by

resuspending the pellet with 2 ml of H_2O and washed immediately in cold PBS/EDTA, before

centrifugation at 350 g for 10 min, at 4 °C.

Human RSV neutralization assay

Serial 2-fold dilutions of heat-inactivated sera (30min, 56°C) were prepared in serum-free

DMEM medium, with a starting dilution of 1:4. Serial serum dilutions were incubated at 37 °C

for 60 min with an equal volume of RSV to provide 100 PFU per 100 µl. The serum-virus

mixtures were added in duplicate to monolayers of HEp-2 cells in 96-well flat-bottom tissue

culture-treated microtiter plates and incubated at 37 °C with 5% CO_2 in DMEM 5% FBS. After

48 h, cells were washed with 100 µl PBS per well and fixed for 20 min at RT with methanol

containing 2% H_2O_2. Cells were washed twice with 100 µl PBS per well, and biotinylated anti-

RSV antibody (goat polyclonal IgG, Bio-Rad) was added for 1 h at RT. Cells were washed

twice with 100 µl PBS 1% BSA per well and ExtrAvidin®-Peroxidase (Sigma) was added for

30 min at RT. Cells were washed twice with 100 µl PBS 1% BSA per well and DAB substrate

(Sigma) was added and left up to 30 min at RT. When the plaques appeared with the desired

coloration, the reaction was stopped with PBS. Serial two-fold dilutions of Palivizumab®

(Synagis), beginning at 100 µg/ml, and virus-free DMEM were used as controls. The plates

were scanned with an ImmunoSpot® analyzer (Cellular Technology Limited (CTL)).

RNA isolation and quantitative PCR

Total RNA was extracted from BAL fluids using the QiAamp® Viral RNA Mini Kit (QIAgen) and

Nucleospin® RNA Plus Kit (Macherey Nagel) for lung tissues and the BAL cellular fraction

according to the manufacturer's protocol. For quantitation of RNA expression levels,

quantitative PCR (qPCR) was performed using AgPath-ID™ One-Step RT-PCR Reagents

(ThermoFisher), according to the manufacturer's instructions. The primer and probe

sequences used to detect RSV-A2 and the housekeeping control 18S RNAs have been

previously described: RSV-A2 (FW: 5′-GAACTCAGTGTAGGTAGAATGTTTGCA-3′, RV: 5′-

TTCAGCTATCATTTTCTCTGCCAAT-3′,   Probe: 5′-FAM-

TTTGAACCTGTCTGAACATTCCCGGTT-TAMRA-3′)   (59);   18S   (FW: 5′-

CGCCGCTAGAGGTGAAATTCT-3′, RV: 5′-CATTCTTGGCAAATGCTTTCG-3′, Probe: 5′-

FAM-ACCGGCGCAAGACGGACCAGA-TAMRA-3′) (60). RNA expression levels of RSV and

18S were determined by absolute quantification by the serial dilution of plasmids containing

the sequence of interest. For the quantification of interferon-β (IFN-β), primers were designed

as   follow: FW: 5′-TGGTTCTCCTGCTGTGTTTCTC-3′;   RV: 5′-

CGTTGTTGGAATCGAAGCAA-3′; Probe: 5′-FAM-ACCACAGCTCTTTCCAGGAGCTACA-

BHQ1-3′ (61). For the amplification on ABI Fast 7500 Sequence Detection System (Applied

Biosystems), the following conditions were used: 95°C 20 sec, 40-45 cycles of the following:

95 °C 3 sec, 60 °C 30 sec, real-time data were analyzed using the SDS software (Applied

Biosystems).

Flow cytometry

The different ovine immune cell subtypes were identified by flow cytometry (FCM) using an

eight-step, seven-color staining protocols. All antibodies used for the procedure, as well as

their clones, host, working dilution, and references, are listed in Table E1. Combination

stainings analyzed plasmacytoid dendritic cell (pDC), $\gamma\delta$ T cells, Tregs, CD4 and CD8 T cell

subsets. For the acquisitions, 10^6 events were accumulated for each sample. The

experimental schedule is summarized in Table E2. For the counts of cell subtypes, we

calculated the percentage of total events: ratio (number of events in the gated cell subtype) to

(number of all events, excepting cell aggregates and debris). FCM acquisitions were

performed on a FACS Canto II (BD Bioscience) using the DIVA software and further analyzed

with FlowJo (TreeStar).

Multiplex immunoassay

For multiplex assay, the fluidic fraction of BALs from mock controls and RSV-A2 were

collected. Cytokine concentrations were assessed using the MILLIPLEX® Ovine

Cytokine/Chemokine Panel 1 (Life Science Research) according to the manufacturer's

protocol and read on a Luminex MAGPIX® System including a xPONENT Software version

4.2 software (Luminex).

Statistical Analysis

Statistical analysis was done using the GraphPad Prism 8 software (GraphPad software, La

Jolla, CA, USA). Data are presented as box and whisker plots and to determine differences

between two groups, non-parametric paired Wilcoxon tests or Mann–Whitney U-tests were

used as appropriate. Associations were tested using the Spearman rank correlation test. A p

value < 0.05 was considered statistically significant.

NOTES

Acknowledgements

We thank Artur Summerfield, Christoph Aebi and Sean R. R. Hall for helpful discussions; Sylvie Python, Aurélie Godel and Nathan Leborgne for technical assistance; Inês Berenguer Veiga for participation in the animal trial; Hans Peter Lüthi, Roman Troxler, Jan Salchli, Daniel Brechbühl and Katarzyna Sliz for animal care. The RSV isolate RSV-ON1-H1 was kindly provided by Sibylle Haid, Martin Wetzke, Gesine Hansen and Thomas Pietschmann. T.D. was supported in part by the EU FP7 Project UniVax (HEALTH-F3-2013-60173). This project was funded by the Swiss National Science Foundation (grant No. 310030_172895 to M.P.A.).

Data availability statement

The data generated and analysed during the current study are available from the corresponding author on reasonable request.

Authors contribution

All authors contributed to important elements to the work presented in this paper and reviewed and approved the final version of this manuscript. T.D. designed the various antibody combination staining, validated them *in vitro*, designed and performed the animal trials, performed the RSV quantifications in animal tissues and serum micro-neutralization assays, acquired, assembled, analyzed and interpreted the data, prepared the figures and wrote the manuscript. M.B. and B.Z established the RSV quantification and serum micro-neutralization assay protocols and designed and performed the animal trial. B.I.O.E. participated in the execution of the animal trial. K.M. and H.P. participated in the execution of the animal trial and evaluated the lung pathology. A.F. assisted in the animal trial and performed part of the immunohistochemistry. L.B. computed the data from WBC and PMN counts in the circulation with the R software and generated the related figures. A.T. helped with the multiplex assay. C.B. helped with clinical observations, interpretation of the data, and reviewed the manuscript.

N.R. participated in the design and execution of the animal trial, organized the partitioning of

animals in stables, and monitored the clinical scores. M.P.A. conceived the idea, overviewed,

designed, led and performed the animal trial, and edited the manuscript.

SUPPLEMENTARY MATERIAL

Table S1. List of antibodies validated for their use in the study.

Table S2. Combination immunostainings used for the identification of cell subsets.

Figure S1. Longitudinal evaluation of peripheral blood over the course of RSV disease.

Figure S2. Evaluation of serum RSV-specific neutralizing antibodies.

Figure S3. RSV-specific neutralizing activity present in one uninfected animal.

Figure S4. FCM gating strategy for immune cells identification.

Figure S5. Neonatal RSV infection induces bronchoalveolar recruitment of activated pDCs.

REFERENCES

1. Shi T, McAllister DA, O'Brien KL, Simoes EAF, Madhi SA, Gessner BD, et al. Global, regional, and national disease burden estimates of acute lower respiratory infections due to respiratory syncytial virus in young children in 2015: a systematic review and modelling study. Lancet. 2017;390(10098):946-58.

2. Pneumonia Etiology Research for Child Health Study G. Causes of severe pneumonia requiring hospital admission in children without HIV infection from Africa and Asia: the PERCH multi-country case-control study. Lancet. 2019;394(10200):757-79.

3. Nair H, Nokes DJ, Gessner BD, Dherani M, Madhi SA, Singleton RJ, et al. Global burden of acute lower respiratory infections due to respiratory syncytial virus in young children: a systematic review and meta-analysis. Lancet. 2010;375(9725):1545-55.

4. Hall CB, Weinberg GA, Iwane MK, Blumkin AK, Edwards KM, Staat MA, et al. The burden of respiratory syncytial virus infection in young children. N Engl J Med. 2009;360(6):588-98.

5. Murray J, Bottle A, Sharland M, Modi N, Aylin P, Majeed A, et al. Risk factors for hospital admission with RSV bronchiolitis in England: a population-based birth cohort study. PLoS One. 2014;9(2):e89186.

6. Garcia CG, Bhore R, Soriano-Fallas A, Trost M, Chason R, Ramilo O, et al. Risk factors in children hospitalized with RSV bronchiolitis versus non-RSV bronchiolitis. Pediatrics. 2010;126(6):e1453-60.

7. Hall CB, Weinberg GA, Blumkin AK, Edwards KM, Staat MA, Schultz AF, et al. Respiratory syncytial virus-associated hospitalizations among children less than 24 months of age. Pediatrics. 2013;132(2):e341-8.

8. Palivizumab, a Humanized Respiratory Syncytial Virus Monoclonal Antibody, Reduces Hospitalization From Respiratory Syncytial Virus Infection in High-risk Infants. Pediatrics. 1998;102(3):531-7.

9.   Openshaw PJM, Chiu C, Culley FJ, Johansson C. Protective and Harmful Immunity to

RSV Infection. Annu Rev Immunol. 2017;35:501-32.

10.   Mold JE, Venkatasubrahmanyam S, Burt TD, Michaelsson J, Rivera JM, Galkina SA,

et al. Fetal and adult hematopoietic stem cells give rise to distinct T cell lineages in humans.

Science. 2010;330(6011):1695-9.

11.   de Kleer IM, Kool M, de Bruijn MJ, Willart M, van Moorleghem J, Schuijs MJ, et al.

Perinatal Activation of the Interleukin-33 Pathway Promotes Type 2 Immunity in the

Developing Lung. Immunity. 2016;45(6):1285-98.

12.   PrabhuDas M, Adkins B, Gans H, King C, Levy O, Ramilo O, et al. Challenges in infant

immunity: implications for responses to infection and vaccines. Nat Immunol. 2011;12(3):189-

94.

13.   Dodd J, Riffault S, Kodituwakku JS, Hayday AC, Openshaw PJ. Pulmonary V gamma

4+ gamma delta T cells have proinflammatory and antiviral effects in viral lung disease. J

Immunol. 2009;182(2):1174-81.

14.   Aoyagi M, Shimojo N, Sekine K, Nishimuta T, Kohno Y. Respiratory syncytial virus

infection suppresses IFN-gamma production of gammadelta T cells. Clin Exp Immunol.

2003;131(2):312-7.

15.   Huang H, Saravia J, You D, Shaw AJ, Cormier SA. Impaired gamma delta T cell-

derived IL-17A and inflammasome activation during early respiratory syncytial virus infection

in infants. Immunol Cell Biol. 2015;93(2):126-35.

16.   McGill JL, Sacco RE. gammadelta T cells and the immune response to respiratory

syncytial virus infection. Vet Immunol Immunopathol. 2016;181:24-9.

17.   Restori KH, Srinivasa BT, Ward BJ, Fixman ED. Neonatal Immunity, Respiratory Virus

Infections, and the Development of Asthma. Front Immunol. 2018;9:1249.

18.   Levy O. Innate immunity of the newborn: basic mechanisms and clinical correlates.

Nat Rev Immunol. 2007;7(5):379-90.

19.   DeVincenzo JP, Wilkinson T, Vaishnaw A, Cehelsky J, Meyers R, Nochur S, et al. Viral

load drives disease in humans experimentally infected with respiratory syncytial virus. Am J

Respir Crit Care Med. 2010;182(10):1305-14.

20.   Pino M, Kelvin DJ, Bermejo-Martin JF, Alonso A, Matias V, Tenorio A, et al.

Nasopharyngeal aspirate cytokine levels 1 yr after severe respiratory syncytial virus infection.

Pediatr Allergy Immunol. 2009;20(8):791-5.

21.   Johnson JE, Gonzales RA, Olson SJ, Wright PF, Graham BS. The histopathology of

fatal untreated human respiratory syncytial virus infection. Mod Pathol. 2007;20(1):108-19.

22.   Piedra PA. Clinical experience with respiratory syncytial virus vaccines. Pediatr Infect

Dis J. 2003;22(2 Suppl):S94-9.

23.   Fishaut M, Tubergen D, McIntosh K. Cellular response to respiratory viruses with

particular reference to children with disorders of cell-mediated immunity. J Pediatr.

1980;96(2):179-86.

24.   Roman M, Calhoun WJ, Hinton KL, Avendano LF, Simon V, Escobar AM, et al.

Respiratory syncytial virus infection in infants is associated with predominant Th-2-like

response. Am J Respir Crit Care Med. 1997;156(1):190-5.

25.   Legg JP, Hussain IR, Warner JA, Johnston SL, Warner JO. Type 1 and type 2 cytokine

imbalance in acute respiratory syncytial virus bronchiolitis. Am J Respir Crit Care Med.

2003;168(6):633-9.

26.   Siefker DT, Vu L, You D, McBride A, Taylor R, Jones TL, et al. Respiratory Syncytial

Virus Disease Severity Is Associated with Distinct CD8(+) T-Cell Profiles. Am J Respir Crit

Care Med. 2020;201(3):325-34.

27.   Scheerlinck JP, Snibson KJ, Bowles VM, Sutton P. Biomedical applications of sheep

models: from asthma to vaccines. Trends Biotechnol. 2008;26(5):259-66.

28.   Olivier A, Gallup J, de Macedo MM, Varga SM, Ackermann M. Human respiratory

syncytial virus A2 strain replicates and induces innate immune responses by respiratory

epithelia of neonatal lambs. Int J Exp Pathol. 2009;90(4):431-8.

29.   Sitthicharoenchai P, Alnajjar S, Ackermann MR. A model of respiratory syncytial virus

(RSV) infection of infants in newborn lambs. Cell Tissue Res. 2020;380(2):313-24.

30.   Derscheid RJ, Ackermann MR. Perinatal lamb model of respiratory syncytial virus

(RSV) infection. Viruses. 2012;4(10):2359-78.

31.   Smit JJ, Rudd BD, Lukacs NW. Plasmacytoid dendritic cells inhibit pulmonary

immunopathology and promote clearance of respiratory syncytial virus. J Exp Med.

2006;203(5):1153-9.

32.   McCarthy NE, Eberl M. Human gammadelta T-Cell Control of Mucosal Immunity and

Inflammation. Front Immunol. 2018;9:985.

33.   Zhivaki D, Lemoine S, Lim A, Morva A, Vidalain PO, Schandene L, et al. Respiratory

Syncytial Virus Infects Regulatory B Cells in Human Neonates via Chemokine Receptor

CX3CR1 and Promotes Lung Disease Severity. Immunity. 2017;46(2):301-14.

34.   Schmidt ME, Knudson CJ, Hartwig SM, Pewe LL, Meyerholz DK, Langlois RA, et al.

Memory CD8 T cells mediate severe immunopathology following respiratory syncytial virus

infection. PLoS Pathog. 2018;14(1):e1006810.

35.   Liesman RM, Buchholz UJ, Luongo CL, Yang L, Proia AD, DeVincenzo JP, et al. RSV-

encoded NS2 promotes epithelial cell shedding and distal airway obstruction. J Clin Invest.

2014;124(5):2219-33.

36.   Garcia-Mauriño C, Moore-Clingenpeel M, Thomas J, Mertz S, Cohen DM, Ramilo O,

et al. Viral Load Dynamics and Clinical Disease Severity in Infants With Respiratory Syncytial

Virus Infection. The Journal of infectious diseases. 2019;219(8):1207-15.

37.   El Saleeby CM, Bush AJ, Harrison LM, Aitken JA, Devincenzo JP. Respiratory

syncytial virus load, viral dynamics, and disease severity in previously healthy naturally

infected children. The Journal of infectious diseases. 2011;204(7):996-1002.

38.   Kerrin A, Fitch P, Errington C, Kerr D, Waxman L, Riding K, et al. Differential lower

airway dendritic cell patterns may reveal distinct endotypes of RSV bronchiolitis. Thorax.

2017;72(7):620-7.

39.   Ruckwardt TJ, Malloy AM, Morabito KM, Graham BS. Quantitative and qualitative

deficits in neonatal lung-migratory dendritic cells impact the generation of the CD8+ T cell

response. PLoS Pathog. 2014;10(2):e1003934.

40.   Johnson TR, Johnson CN, Corbett KS, Edwards GC, Graham BS. Primary human

mDC1, mDC2, and pDC dendritic cells are differentially infected and activated by respiratory

syncytial virus. PLoS One. 2011;6(1):e16458.

41.   Kollmann TR, Crabtree J, Rein-Weston A, Blimkie D, Thommai F, Wang XY, et al.

Neonatal innate TLR-mediated responses are distinct from those of adults. J Immunol.

2009;183(11):7150-60.

42.   Marr N, Wang TI, Kam SH, Hu YS, Sharma AA, Lam A, et al. Attenuation of respiratory

syncytial virus-induced and RIG-I-dependent type I IFN responses in human neonates and

very young children. J Immunol. 2014;192(3):948-57.

43.   Danis B, George TC, Goriely S, Dutta B, Renneson J, Gatto L, et al. Interferon

regulatory factor 7-mediated responses are defective in cord blood plasmacytoid dendritic

cells. Eur J Immunol. 2008;38(2):507-17.

44.   Lau-Kilby AW, Turfkruyer M, Kehl M, Yang L, Buchholz UJ, Hickey K, et al. Type I IFN

ineffectively activates neonatal dendritic cells limiting respiratory antiviral T-cell responses.

Mucosal Immunol. 2020;13(2):371-80.

45.   De Weerd W, Twilhaar WN, Kimpen JL. T cell subset analysis in peripheral blood of

children with RSV bronchiolitis. Scand J Infect Dis. 1998;30(1):77-80.

46.   Raiden S, Pandolfi J, Payaslian F, Anderson M, Rivarola N, Ferrero F, et al. Depletion

of circulating regulatory T cells during severe respiratory syncytial virus infection in young

children. Am J Respir Crit Care Med. 2014;189(7):865-8.

47.   Thornburg NJ, Shepherd B, Crowe JE, Jr. Transforming growth factor beta is a major

regulator of human neonatal immune responses following respiratory syncytial virus infection.

J Virol. 2010;84(24):12895-902.

48. Durant LR, Makris S, Voorburg CM, Loebbermann J, Johansson C, Openshaw PJ.

Regulatory T cells prevent Th2 immune responses and pulmonary eosinophilia during

respiratory syncytial virus infection in mice. J Virol. 2013;87(20):10946-54.

49. Strickler HD, Martinson J, Desai S, Xie X, Burk RD, Anastos K, et al. The relation of

plasmacytoid dendritic cells (pDCs) and regulatory T-cells (Tregs) with HPV persistence in

HIV-infected and HIV-uninfected women. Viral Immunol. 2014;27(1):20-5.

50. Caballero MT, Serra ME, Acosta PL, Marzec J, Gibbons L, Salim M, et al. TLR4

genotype and environmental LPS mediate RSV bronchiolitis through Th2 polarization. J Clin

Invest. 2015;125(2):571-82.

51. Bermejo-Martin JF, Garcia-Arevalo MC, De Lejarazu RO, Ardura J, Eiros JM, Alonso

A, et al. Predominance of Th2 cytokines, CXC chemokines and innate immunity mediators at

the mucosal level during severe respiratory syncytial virus infection in children. Eur Cytokine

Netw. 2007;18(3):162-7.

52. Vu LD, Siefker D, Jones TL, You D, Taylor R, DeVincenzo J, et al. Elevated Levels of

Type 2 Respiratory Innate Lymphoid Cells in Human Infants with Severe Respiratory Syncytial

Virus Bronchiolitis. Am J Respir Crit Care Med. 2019;200(11):1414-23.

53. Hinks TSC, Hoyle RD, Gelfand EW. CD8(+) Tc2 cells: underappreciated contributors

to severe asthma. Eur Respir Rev. 2019;28(154).

54. Hebel K, Weinert S, Kuropka B, Knolle J, Kosak B, Jorch G, et al. CD4+ T cells from

human neonates and infants are poised spontaneously to run a nonclassical IL-4 program. J

Immunol. 2014;192(11):5160-70.

55. Bendelja K, Gagro A, Bace A, Lokar-Kolbas R, Krsulovic-Hresic V, Drazenovic V, et

al. Predominant type-2 response in infants with respiratory syncytial virus (RSV) infection

demonstrated by cytokine flow cytometry. Clin Exp Immunol. 2000;121(2):332-8.

56. Blockus S, Sake SM, Wetzke M, Grethe C, Graalmann T, Pils M, et al.

Labyrinthopeptins as virolytic inhibitors of respiratory syncytial virus cell entry. Antiviral Res.

2020;177:104774.

57.   Gnanalingham MG, Mostyn A, Gardner DS, Stephenson T, Symonds ME.

Developmental regulation of the lung in preparation for life after birth: hormonal and nutritional

manipulation of local glucocorticoid action and uncoupling protein-2. J Endocrinol.

2006;188(3):375-86.

58.   Mehinagic K, Pilo P, Vidondo B, Stokar-Regenscheit N. Coinfection of Swiss cattle with

bovine parainfluenza virus 3 and Mycoplasma bovis at acute and chronic stages of bovine

respiratory disease complex. J Vet Diagn Invest. 2019;31(5):674-80.

59.   Lee DC, Harker JA, Tregoning JS, Atabani SF, Johansson C, Schwarze J, et al. CD25+

natural regulatory T cells are critical in limiting innate and adaptive immunity and resolving

disease following respiratory syncytial virus infection. J Virol. 2010;84(17):8790-8.

60.   Vielle NJ, Zumkehr B, Garcia-Nicolas O, Blank F, Stojanov M, Musso D, et al. Silent

infection of human dendritic cells by African and Asian strains of Zika virus. Sci Rep.

2018;8(1):5440.

61.   Sow FB, Gallup JM, Olivier A, Krishnan S, Patera AC, Suzich J, et al. Respiratory

syncytial virus is associated with an inflammatory response in lungs and architectural

remodeling of lung-draining lymph nodes of newborn lambs. Am J Physiol Lung Cell Mol

Physiol. 2011;300(1):L12-24.

    FIGURE LEGENDS

    Figure 1. RSV infection is causing lower respiratory tract disease and leading to a

    potentially protective humoral response. (A) Illustration of the transtracheal RSV infection

    experimental design. (B) RSV was quantified in the respiratory tract (BAL, cell fraction; Lung;

    BAL, fluid) by qPCR. (C) RSV detection by immunohistochemistry in macrophages, and type

    I pneumocytes. (D) Representative histological lung tissue sections from neonates over the

    course of RSV infection (40X, HE). Lungs of RSV-infected animals (strain A2) show interstitial

    septae expanded by inflammatory cells (interstitial pneumonia) and accumulation of

    inflammatory cells within alveoli at days 3 and 6 p.i. No lesions at days 14 and 42 p.i. The

    lungs of mock neonates allow comparison with normal lung architecture within the first weeks

    of life. (E) Neonates infected with RSV can mount an antiviral neutralizing-antibody response.

    (F) Anti-RSV Nabs persist over a 42-day long period. (G) Multiplex immunoassay using BAL

    from the indicated animals. Heat map showing log_2-fold changes in concentration of 6

    measured cytokines/chemokines normalized by mean value of control group: Mock, 3 days

    p.i. Each symbol represents an individual animal (healthy adults, n=12; mock neonates, n=3

    per time point; neonates infected with RSV, n=6 per time point). Boxplots indicate median

    value (center line) and interquartile ranges (box edges), with whiskers extending to the lowest

    and the highest values. Stars indicate significance levels. *, $p < 0.05$.

    Figure 2. Neonatal RSV infection activates a transient depletion of $\gamma\delta$ T cells and

    deficiency of IL-17-mediated immunity. (A) Neonatal RSV infection induces a decrease in

    $\gamma\delta$ T cells number in the bronchoalveolar space, but not in the lung tissue. (B) Correlation

    coefficient (r) obtained with in $\gamma\delta$ T cell counts calculated as a function of % Live/Dead$^+$. (C-D)

    Replenishment and maturation (CD25 up-regulation) of $\gamma\delta$ T cell subset following the depletion

    after RSV infection (C, representative FACS contour plots; D, Pie charts displaying all

    individuals). (E) Inability of $\gamma\delta$ T cells to produce IL-17 (representative FCM contour plots).

    Each symbol represents an individual animal (healthy adults, n=12; mock neonates, n=3 per

time point; neonates infected with RSV, n=6 per time point). Boxplots indicate median value (center line) and interquartile ranges (box edges), with whiskers extending to the lowest and the highest values. Stars indicate significance levels. *, $p < 0.05$.

Figure 3. Neonatal RSV infection induces a fast and massive Tc2 influx in the bronchoalveolar space. (A) RSV infection in neonate accelerates the recruitment of CD4$^+$ in the bronchoalveolar space, but not in peribronchial LN or lung tissue. (B) RSV infection in neonate induces a strong recruitment of CD8$^+$ T cells in the bronchoalveolar space, and to a lesser extent in lung. (C-D) A high frequency of recruited CD4$^+$ T cells in bronchoalveolar space differentiate towards a Th2 cytokine profile (IL-4 production) (C, representative FACS contour plots; D, plot displaying all individuals). (E-F) A high frequency of recruited CD8$^+$ T cells in bronchoalveolar space differentiate towards a Tc2 cytokine profile (IL-4 production) (E, representative FCM contour plots; F, plot displaying all individuals). (G) RSV infection in neonates modulates the activation (cell surface expression of CD25) and the maturation (cell surface expression of T cell memory marker CD45RO) of CD4$^+$ and CD8$^+$ T cell pools in the lung. Each symbol represents an individual animal (healthy adults, n=12; mock neonates, n=3 per time point; neonates infected with RSV, n=6 per time point). Boxplots indicate median value (center line) and interquartile ranges (box edges), with whiskers extending to the lowest and the highest values. Stars indicate significance levels. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

Figure 4. Neonatal RSV infection rapidly induces Treg suppressive functions to dampen the Th2 and Tc2 responses. (A) RSV infection in neonates led to the recruitment of Tregs in the bronchoalveolar space, but not in peribronchial LN or lung tissue. (B) Progressive decrease with age of Treg subset among the CD4$^+$ T cell fraction (representative FCM contour plots). (C) Frequency of TGF-β-producing Tregs. (D) Correlation coefficient (r) obtained in BALs with Treg counts calculated as a function of non-Treg CD4$^+$ counts (left), Treg counts calculated as a function of CD8$^+$ counts (middle) and CD8$^+$ counts calculated as a function of

non-Treg CD4$^+$ counts (right). Significance was reached for the three correlations. Each

symbol represents an individual animal (healthy adults, n=12; mock neonates, n=3 per time

point; neonates infected with RSV, n=6 per time point). Boxplots indicate median value (center

line) and interquartile ranges (box edges), with whiskers extending to the lowest and the

highest values. Stars indicate significance levels. *, $p < 0.05$; **, $p < 0.01$.

Figure 5. Distinct cellular immune response in neonates compare to adults following

RSV infection. (A) RSV was quantified in the respiratory tract (BAL, cell fraction; lung; BAL,

fluid) of adults and neonates by qPCR. (B) Representative histological lung tissue sections

from RSV-infected neonates and adults at day 6 and 14 p.i. (40X, HE). Adults show no

histological lesions, whereas neonates show interstitial pneumonia at day 6.p.i. (C-D) Counts

of recruited pDCs and CD86 upregulation in bronchoalveolar space (C, plot displaying all

individuals; D, representative FCM contour plots). (E-G) $\gamma\delta$ T-cell counts, cytokine production,

and maturation in the bronchoalveolar space (E, plot displaying all individuals; F,

representative FCM contour plots; G, pie charts integrating all animals). (H-J) Recruitment,

activation, and maturation of the CD4$^+$ T-cell pool in the bronchoalveolar space (H, plot

displaying all individuals; I, representative FACS contour plots; J, pie charts integrating all

animals). (K-M) As in (H-J), but showing CD8$^+$ T-cell pool. (N-O) Recruitment of Tregs and

TGF-β production in the bronchoalveolar space (N, plot displaying all individuals; O,

representative FACS contour plots). Each symbol represents an individual animal (infected

adults, n=3 per time point; neonates infected with RSV, n=8 per time point). The grey zone

corresponds to the interval integrating the values of all corresponding mock controls. Boxplots

indicate median value (center line) and interquartile ranges (box edges), with whiskers

extending to the lowest and the highest values. Stars indicate significance levels. *, $p < 0.05$;

**, $p < 0.01$; ***, $p < 0.001$.

Figure 6. The severity of disease relates to the magnitude of the unconventional

bronchoalveolar T cell responses. (A) Representative histological lung tissue sections from

mock neonates and neonates infected with RSV-ON1-H1 and RSV A2 strain (40X, HE). At day 6 p.i. the RSV-ON1-H1 infected animals show mild interstitial pneumonia (upper right corner) with accumulation of inflammatory cells within alveoli. No lesions at day 14 p.i. or mock-infected animals. (B) RSV was quantified in the respiratory tract (BAL, cell fraction) of neonates infected with either RSV-ON1-H1 or RSV A2 strain by qPCR. (C) Comparison of immune cell subset trafficking into bronchoalveolar space after infection with RSV-ON1-H1 or RSV A2 strains. (D) Correlation coefficient (r) obtained in BALs with Treg counts calculated as a function of pDC counts. Each symbol represents an individual animal (neonates infected with RSV-ON1-H1, n=4 per time point; neonates infected with RSV-A2, n=8 per time point). The grey zone corresponds to the interval integrating the values of all corresponding mock controls. Boxplots indicate median value (center line) and interquartile ranges (box edges), with whiskers extending to the lowest and the highest values. Stars indicate significance levels. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

A

RSV

Birth
Acclimatization

-6   -3   0   3   6   9   12   15   19   22   25   31   34   37   40

* BALs
  Lung
  Draining LNs

▽ Clinical status
  Blood sampling

Healthy Adults | Mock | RSV — Day 3, p.i.
Healthy Adults | Mock | RSV — Day 6, p.i.
Healthy Adults | Mock | RSV — Day 14, p.i.
Healthy Adults | Mock | RSV — Day 42, p.i.

B

BAL, cell fraction

RSV copies/10¹² 18S copies

Adu.   3   6   14   42

Lung

Adu.   3   6   14   42
Days p.i.

p=0.083

BAL, fluid

RSV copies per ml

Adu.   3   6   14   42

p=0.071

C

6   6

Mock   RSV

D

3   6   14   42

Mock

RSV

E

Healthy adults   Mock   RSV

Serum neutralisation (%)

0   13/14   0   13/14   0   13/14

Days p.i.

***

F

0   13   22   33   42

Healthy adults

Mock

RSV

Serum neutralisation (%)

p=0.071   *   *   *

Adu. 0   13   22   33   42

Days p.i.

G

BAL, fluid

3   6   14   42

Mock   RSV   Mock   RSV   Mock   RSV   Mock   RSV

IP-10
IL-8
IL1-α
VEGF-A
IL-10
IL-36RA

Fold change (log$_2$)

-1

-2

6