The developing T-cell compartment of the neonatal lung orchestrates an atypical response to respiratory syncytial virus

Thomas Démoulins[1,2], Melanie Brügger[1,2,3], Beatrice Zumkehr[1,2], Blandina I. Oliveira Esteves[1,2], Kemal Mehinagic[1,2,3,4], Amal Fahmi[1,2,3], Loïc Borcard[1,2,3], Horst Posthaus[4,5], Charaf Benarafa[1,2], Nicolas Ruggli[1,2], and Marco P. Alves[1,2,*]

[1] *The Institute of Virology and Immunology, Mittelhäusern and Bern, Switzerland*

[2] *Department of Infectious Diseases and Pathobiology, Vetsuisse Faculty, University of Bern, Bern, Switzerland*

[3] *Graduate School for Cellular and Biomedical Sciences, University of Bern, Bern, Switzerland*

[4] *Institute of Animal Pathology, Department of Infectious Diseases and Pathobiology, Vetsuisse Faculty, University of Bern, Bern, Switzerland*

[5] *COMPATH, Vetsuisse Faculty & Faculty of Medicine, University of Bern, Bern, Switzerland*

[*] **Contact:**

marco.alves@vetsuisse.unibe.ch

Phone: +41 31 631 2483

Authors contribution

All authors contributed to important elements to the work presented in this paper and reviewed and approved the final version of this manuscript. T.D. designed the various antibody combination staining, validated them *in vitro*, designed and performed the

animal trials, performed the RSV quantifications in animal tissues and serum micro-neutralization assays, acquired, assembled, analyzed and interpreted the data, prepared the figures and wrote the manuscript. M.B. and B.Z established the RSV quantification and serum micro-neutralization assay protocols and designed and performed the animal trial. B.I.O.E. participated in the execution of the animal trial. K.M. and H.P. participated in the execution of the animal trial and evaluated the lung pathology. A.F. assisted in the animal trial and performed part of the immunohistochemistry. L.B. computed the data from WBC and PMN counts in the circulation with the R software and generated the related figures. C.B. helped with clinical observations, interpretation of the data, and reviewed the manuscript. N.R. participated in the design and execution of the animal trial, organized the partitioning of animals in stables, and monitored the clinical scores. M.P.A. conceived the idea, overviewed, designed, led and performed the animal trial, and edited the manuscript.

Sources of support

The RSV isolate RSV-ON1-H1 was kindly provided by Sibylle Haid, Martin Wetzke, Gesine Hansen and Thomas Pietschmann. T.D. was supported in part by the EU FP7 Project UniVax (HEALTH-F3-2013-60173). This project was funded by the Swiss National Science Foundation (grant No. 310030_172895 to M.P.A.).

Running Title: Atypical neonatal immune response to RSV infection

Abstract word count: 249

Total word count: 3863

Abstract

Rationale. Respiratory syncytial virus (RSV) is a major cause of lower respiratory tract infection in infancy, potentially aided by an inappropriate immune response. Sparse information is available for the distal lung, mostly because data arose from non-invasive samplings of peripheral blood and nasal aspirates.

Objectives. To determine the neonatal immune response to RSV in the bronchoalveolar space and better understand why neonates are at greater risk of developing severe disease.

Methods. We used the newborn lamb, a state-of-the-art translational model of human RSV infection, offering ease sampling and full accessibility to lower airways. Using a multiparameter flow cytometry assay, we evaluated the frequency and activation/maturation state of the major subsets of the developing T-cell compartment.

Measurements and Main Results. The T-cell compartment of the healthy developing lung was very distinct to that seen in adults. We observed a high frequency of type 2 CD4$^+$ (Th2) and CD8$^+$ (Tc2) T-cells, both being a large source of IL-4, which declined progressively over time. Remarkably, RSV infection exacerbated this pro-type 2 environment, rather than inducing a type 2 response *per se*. Neonatal regulatory T-cell (Treg) suppressive functions occurred very early to dampen those Th2 and Tc2 responses, while $\gamma\delta$ T-cells dropped and failed to produce IL-17. The disease severity was related to the magnitude of these T-cell responses.

26 **Conclusion.** The atypical neonatal immune response to RSV consists of distinct T-

27 cell subsets that tightly cooperate, namely a combined bronchoalveolar influx of Treg,

28 Th2 and Tc2 cells, associated with a depletion of $\gamma\delta$ T-cells.

30 **Keywords:** Respiratory syncytial virus; neonatal immunity; bronchiolitis;

31 immunopathogenesis; T cells immunity

Introduction

The human respiratory syncytial virus (RSV) is a seasonal virus, known as the major cause of lower respiratory tract infection during early childhood (1, 2). Worldwide, RSV leads to around 160'000 deaths each year (3). Although there are risk factors for RSV severity such as prematurity and congenital heart disease, the majority of hospitalized infants are previously healthy (4-7), suggesting that the disease is partially linked to inherent properties of early life immunity. The precise understanding of the immune-driven susceptibility of neonates is essential, considering there is no licensed vaccine and available prophylactic treatments are mostly based on palivizumab, a neutralizing antibody used in high-risk group infants (8).

The first exposure to pathogens occurs during the early postnatal period and is critical for lung colonization by immune cells. There is increasing evidence that the overall neonatal T-cell compartment is distinctive during this temporal window (9). Upon stimulation neonatal T-cells tend to differentiate into regulatory T-cells (Tregs), to facilitate self-tolerance to developing organs (10). Also, the developing lung is characterized by a type 2 immune phenotype, and the pulmonary T-cell responses present a T helper type 2 (Th2) bias (11, 12). Remarkably, neonatal $\gamma\delta$ T-cells, a T-cell subset enriched at mucosal barrier sites, have an impaired capacity to respond to stimulation (13-16).

Due to the aforementioned reasons, young infants do not mount a classical pulmonary immune response to pathogens such as RSV, which may contribute to high morbidity and mortality (17, 18). A likely explanation could be a combined contribution of viral load (19) and an inappropriate and/or dysfunctional immune response, through an over-exuberant inflammatory response or a biased T-cell response (9, 20, 21). Neonatal T-cells may have both protective and harmful effects in responses to RSV

infection. While children with T-cell deficiencies cannot efficiently clear the virus (22,

23), it remains controversial whether the severity of RSV disease is associated with a

bias towards type 2 cytokines such as IL-4 (24-26).

Knowledge of the course of RSV disease is still limited due to ethical and technical

difficulties associated with the investigation of the immune response in human

neonates. The neonatal lamb model of RSV infection recapitulates most of the

features of paediatric disease. Indeed, RSV is a natural pathogen of ruminants.

Experimentally-infected animals clinically present rhinitis, coughing and respiratory

distress. Typical pulmonary lesions are bronchiolitis and interstitial pneumonia

associated with inflammatory infiltrates (27-29). This model, immunologically related

to human, was exploited to have full access to the small airways and follow the

pulmonary immune cell features over the course of RSV disease.

Materials and methods

Virus

RSV-A2 strain was obtained from the American Type Culture Collection (ATCC® VR-

1540™). RSV-ON1-H1 strain was isolated from nasopharyngeal aspirates of a child

below the age of five years with confirmed RSV infection, hospitalized at Hannover

Medical School, Germany. The caregivers gave informed consent for sampling and all

steps were conducted in compliance with good clinical and ethical practice and

approved by the local ethical committee (permission number 63090/2012) (30). Virus

propagation and titre determination used standard methods, described in the online

supplement.

Animals

The experiments were approved by the cantonal veterinary authority, and performed

in compliance with the Swiss animal protection law under the authorization BE125/17.

Animals were housed in the high containment animal facility. *Postpartum*, the

acclimatization lasted 3-6 days before infection. In accordance with the principles of

the 3Rs, suckling lambs were held together with their mothers and fed *ad libitum*. More

details about the housing are given in the online supplement.

The transtracheal RSV inoculation was performed under sedation and analgesia by

intramuscular injection of a mixture of midazolam (Dormicum®, Roche) 0.2 mg/kg and

butorphanol (Morphasol®, Dr. E. Graeub AG) 0.2 mg/kg. The trachea was punctured

between the proximal tracheal rings with a 20G needle and a volume of 2 ml of virus

(10^8 PFU for RSV-A2; 10^7 PFU for RSV-ON1-H1) or PBS (mock-treated controls) were

injected slowly.

Histopathology and Immunohistochemistry

The methods used in the present work were described elsewhere (31).

Cell suspension preparation

Cell isolation used standardized methods, described in the online supplement.

Human RSV neutralization assay

Serum neutralization assay used standardized methods, described in the online

supplement.

RNA isolation and quantitative PCR

The primer and probe sequences used to detect RSV have been previously described

(32).

Flow cytometry

The different ovine immune cell subtypes were identified by flow cytometry (FCM)

using an eight-step, seven-color staining protocols. All antibodies used for the

procedure, as well as their clones, host, working dilution, and references, are listed in

Table E1. Combination stainings analyzed plasmacytoid dendritic cell (pDC), $\gamma\delta$ T-

cells, Tregs, CD4 and CD8 T-cell subsets. For the acquisitions, 10^6 events were

accumulated for each sample. The experimental schedule is summarized in Table E2.

For the counts of cell subtypes, we calculated the ratio (number of events in the gated

cell subtype) to (number of all events, excepting cell aggregates and debris). FCM

acquisitions were performed on a FACS Canto II (BD Bioscience) using the DIVA

software and further analyzed with FlowJo (TreeStar).

Statistical Analysis

Statistical analysis was done using GraphPad Prism 8 software. To determine

differences between two groups, non-parametric paired Wilcoxon tests or Mann–

Whitney U-tests were used as appropriate. Associations were tested using the

Spearman rank correlation test. A $p<0.05$ was considered significant.

Results

Respiratory disease following neonatal RSV infection

Selected time-points were as follow: 3 and 6 days post-infection (p.i.), to evaluate the innate and early adaptive responses; 14 days p.i. to assess the adaptive mechanisms and virus clearance; 42 days p.i. to investigate the impact of RSV infection on the immune cell colonization of the developing lung. As controls, non-infected newborns and adults were used (Figure 1A).

We quantified RSV in BALs (cellular and fluid fractions) and in the lung tissue. Although RSV clearance was slower in the cellular fraction of BALs, it became undetectable in any sample at day 42 p.i., demonstrating an efficient clearance (Figure 1B). Immunohistochemistry of lung tissue sections revealed the presence of RSV-antigens in type I pneumocytes and macrophages (Figure 1C).

Clinically, RSV infected neonates showed signs of rhinitis with clear nasal discharge, occasional coughing and wheezing. Macroscopical lesions were visible at days 3-6 p.i, including failure of pulmonary collapse and focal, dark-red areas of subpleural pulmonary consolidation and atelectasis. Histological lesions, present at days 3-6 p.i consisted of multifocal-coalescing areas of lymphocytic and histiocytic infiltrates within interalveolar septae (interstitial pneumonia) and accumulation of inflammatory cells within alveolar spaces and bronchioles (bronchiolitis) (Figure 1D). Longitudinal measurements of white blood counts and neutrophil counts in the circulation showed no difference between mock and infected neonates (Figure E1).

Neonatal RSV infection activates the initiators of the adaptive immune response

When quantified in the BALs, pDC numbers of mock neonates were similar to those of healthy adults. Upon RSV infection, we observed a trend to pDC recruitment as

early as 3 days p.i., and this was enhanced at day 6 p.i. (p<0.05). However, this pDC

recruitment was transitional and tightly regulated, since day 14 p.i corresponded to a

return to steady-state levels (Figure 2A,B). One out of six infected neonates failed to

display any pDC recruitment; this animal was the only one for which RSV was

undetectable in the fluid fraction of BAL (Figure 1B). To verify whether the extent of

pDC recruitment was linked to the magnitude of RSV shedding, pDC counts were

plotted as a function of RSV copies in the fluid fraction of BALs. A significant

association was found (Figure 2C). We next evaluated if the recruited pDCs displayed

a mature and activated phenotype. A significant CD86 upregulation was found at days

3-6 p.i. (Figure 2D,E). This upregulation was transitional and back to steady-state

levels at day 14 p.i.

Neonatal RSV infection leads to a transient depletion of $\gamma\delta$ T-cells

$\gamma\delta$ T-cell potential impairment in the bronchoalveolar space could contribute to RSV

immunopathogenesis (13-15). Their numbers in the BALs of mock neonates were

comparable to those of healthy adults. However, RSV infection led to a significant

decrease in $\gamma\delta$ T-cell frequency at day 14 p.i.. Nevertheless, this $\gamma\delta$ T-cell depletion

was entirely reversed at day 42 p.i. (Figure 2F,G). To exclude the possibility that the

$\gamma\delta$ T-cell reduction could be due to migration or relocation, we calculated their counts

as a function of the percentage of Live/Dead-positive cells. A negative association was

found, suggesting that neonatal $\gamma\delta$ T-cells death is linked to the presence of RSV in

the bronchoalveolar space (Figure 2H). Finally, BAL $\gamma\delta$ T-cells collected from infected

neonates failed to produce detectable levels of IL-17.

We next ascertained whether neonatal $\gamma\delta$ T-cells are characterized by an immature

phenotype. Remarkably, in BALs of healthy adults, half of the $\gamma\delta$ T-cell pool was

composed of mature CD25$^+$ cells, whereas the latter were absent in mock neonates

at day 3. The proportion of mature γδ T-cells progressively strengthened over time,

without reaching the level detected in healthy adults. RSV infection exerts an effect on

this maturation of neonatal γδ T-cells. Not only the observed depletion was rapidly

followed by a complete replenishment at day 42 p.i., but also the restoration was

accompanied by a marked increase in the percentage of CD25-positive cells,

compared to mock neonates (Figure 2I,J).

Neonatal RSV infection exacerbates the type 2 cytokine environment

Assessment moved on to adaptive immune response, and more particularly to IL-4

versus IFN-γ production by T-cells (Figure 3A). In BALs, mock neonates had a very

low number of CD4$^+$ and CD8$^+$ T-cells compared to healthy adults. RSV infection led

to their strong recruitment, albeit with distinct kinetics. An unexpected reduction of

CD4$^+$ T-cell recruitment was seen as soon as 14 days p.i. This contradicted the values

obtained for CD8$^+$ T-cells, where the significant enhancement following RSV infection

was maintained. When we considered both CD4$^+$ and CD8$^+$ fractions at a later time-

point, the numbers were comparable to those of mock neonates (Figure 3B).

Surprisingly, mock neonates had high frequencies of Th2 cells, which declined

progressively over time. Elevated amounts of spontaneous produced IL-4 were a

signature of the developing CD4$^+$ fraction, since it was no longer detectable at day 42,

or in healthy adults. Upon infection, this analysis revealed a trend towards an increase

in the frequency of those Th2 cells, whereas there was no consistent influence on IFN-

γ induction (Figures 3C,D). Consequently, RSV infection in early life exacerbates the

overall lung pro-Th2 cytokine environment, rather than inducing a Th2 immune

response *per se*.

High frequencies of IL-4-expressing CD8$^+$ T-cells (Tc2) were also found in mock

neonates. Again, a progressive decline of Tc2 cells occurred up to day 42 p.i.,

comparable to negligible levels measured in healthy adults. RSV disease induced a

greater overall Tc2-derived cytokine environment in BALs. This exacerbation was

tightly regulated, reduced to a level similar to healthy adults at day 14 p.i. Again, no

consistent influence of RSV infection was seen on IFN-γ induction (Figure 3E,F).

Altogether, these results highlight the role of Tc2 cells as an important source of IL-4

production during neonatal RSV infection.

We then tested the activation and differentiation of naïve T lymphocytes into memory

subsets. The most striking observation was achieved at day 42 p.i.; the percent

contributions of different subsets seen in RSV-infected neonates was highly

superimposable to that found in healthy adults. This was not the case for mock

neonates who had lower frequencies of activated memory T-cells, particularly for the

CD8$^+$ fraction (Figure 3G). Consequently, RSV infection in early life modulates the

activation and the maturation of the T-cell pool.

Next, we monitored the presence of RSV neutralizing antibodies (NAb) (Figures 3H,

E2). In the serum of healthy adults and mock neonates, no RSV neutralizing antibodies

were detected. In contrast, RSV-infected neonates have naturally acquired NAb at day

13-14 p.i., except one animal. Notably, two siblings had RSV NAb preceding infection;

this was passively transmitted by their mother whose serum also had a strong RSV-

neutralization (Figure E3). We extended the assay to later time points to test the

persistence of RSV neutralization. No loss of humoral memory was measured up to

42 days p.i (Figure 3I). Altogether, these results indicate that despite an inappropriate

cell-mediated immune response, neonates can clear RSV by mounting an efficient

humoral response.

Neonatal RSV infection rapidly induces Treg suppressive functions to dampen the Th2 and Tc2 responses

The shortened kinetics observed for CD4⁺ T-cells was elaborated by determining

whether this was due to an early Treg rather than a classical CD4⁺ T-cell response

(Figure 4A). RSV infection elicited a rapid Treg recruitment in BALs that mirrored the

one previously observed for CD4⁺ T-cells. This Treg dynamic is specific for the airway

mucosa, since Treg frequencies remained unchanged in the lung tissue (Figure 4B).

We then noticed a trend for Treg subset overrepresentation following RSV infection

(Figure 4C). The proportion of Tregs declined progressively over time, tending towards

similar levels to those detected in healthy adults. RSV infection promoted

enhancement of TGF-β production by Tregs, however not significant (Figure 4D).

When Treg counts were calculated as a function of non-Treg CD4⁺ counts, an

association was found, suggesting that CD4⁺ T-cell pool tends to differentiate into

Tregs. When Treg counts were calculated as a function of CD8⁺ counts, an association

was still found, confirming that CD8⁺ T-cells and Tregs colonized the alveolar space

in a coordinated manner. By plotting CD8⁺ count as a function of non-Tregs CD4⁺

counts, it appeared that T-cell response was mainly driven by the CD8⁺ fraction (high

numbers at days 6 and 14 p.i.) (Figure 4E). In peribronchial LNs following RSV

infection, CD8⁺ counts remained stable over the course of the disease, showing that

their accumulation is airway mucosa-specific (Figure 4F). Moreover, unlike the

situation described for the BAL compartment, RSV infection did not promote a further

increase of Tc2 cells (Figure 4G).

Distinct cellular immune response in neonates compare to adults following RSV infection

We followed the same approach to dissect the cellular immune response in RSV-infected adults. While the virus clearance was comparable between adults and neonates (Figure 5A), respiratory lesions were absent in adults (Figure 5B).

The rapid and robust pDC recruitment observed in infected neonates did not occur in infected adults. However, lung-resident pDCs of the latter were prompt to react, displaying an even more mature and activated phenotype Importantly, this mature pDC response was also tightly regulated (Figure 5C,D).

The strong decrease in $\gamma\delta$ T-cell counts observed in infected neonates was absent in adults. Moreover, adult $\gamma\delta$ T-cells produced high amounts of IL-17, suggesting that IL-17-mediated immunity is deficient in early life (Figure 5E,F) (15). This IL-17 response was accompanied by an activated phenotype, as indicated by the enhanced proportion of CD25$^+$ cells (Figure 5G). From these results, we could ascertain that neonates lack a mature and functional $\gamma\delta$ T-cell pool in the lower airways.

For the CD4$^+$ compartment, no significant recruitment was found (Figure 5H). The percentage of IL-4-producing cells was very low, and RSV infection had no consistent influence on IFN-γ induction (Figure 5I). Another clear difference was the contraction of effector memory pool (CD25$^+$CD45RO$^+$) at day 14 p.i., indicative of a specific immune response (Figure 5J).

The recruitment of CD8$^+$ T-cells was delayed compared to neonates, only detectable at day 14 p.i. (Figure 5K). Again, the percentage of IL-4-producing cells was low (Figure 5L), whereas the precipitous contraction of effector memory pool at day 14 p.i. suggested a RSV-specific CD8$^+$ T-cell response (Figure 5M). From these results, it

became clear that the inaptitude of neonates to mount an effector T-cell response is

tentatively substituted by a fast and massive Th2/Tc2 cell recruitment.

Finally, we failed to detect any Treg recruitment. However, the rare tissue-resident

Tregs were able to produce TGF-β, suggesting that an immune regulation happened

(Figure 5N,O).

The severity of disease relates to the magnitude of bronchoalveolar T-cell

responses

The pulmonary T-cell response was evaluated in neonates infected with RSV-ON1-

H1 strain, which induced a milder disease (Figure 6A). We did not observe a γδ T-cell

depletion, and the CD8$^+$ T-cell recruitment was delayed. Furthermore, the Treg

recruitment was lower compared to RSV A2 infection (Figure 6B). By combining

results obtained with both strains, we found a significant association between Treg

and pDC counts in BALs (Figure 6C). This indicates that, despite their skew, all distinct

subsets of neonatal T-cells cooperate tightly with the magnitude of the cellular immune

response against RSV infection.

Discussion

To properly assess the immunopathogenesis of RSV, it is essential to select a model

that recapitulates the features of human disease. We selected the neonatal lamb, as

RSV is a natural pathogen for this species, lung development is similar, and the key

features mimic human RSV infection in infants (27, 28). One strength of our

experimental approach was the ease sampling and the full accessibility to the lower

airways throughout the disease when sampling in infants is mainly restricted to the

upper airways, the latter representing only an approximate surrogate of the distal lung

where RSV disease occurs. Moreover, we offer an unprecedented view of the

development of pulmonary neonatal immunity in healthy conditions. This process is

highly dynamic and tightly regulated, with a short and early stage where colonizing T-

cell subsets synergize towards a narrow pro-tolerogenic window, namely a

"Treg/Th2/Tc2" environment. This certainly contributes to the maturation of the

immune system and may constitute an unsuitable state to resolve RSV infection.

The mechanisms underlying the severity of disease are still unclear. Herein, we

observed a comparable virus clearance between adults and neonates, reinforcing the

assumption of an inappropriate immune response. We observed a rapid and robust

recruitment of neonatal pDCs, whose extent correlated with the viral loads of infected

neonates. Infected adults did not require this recruitment, since their lung-resident

pDCs reacted promptly by maturing to a higher level than that measured in neonates.

This partially corroborates recent studies reporting low counts and maturation levels

of pDCs in mice and infants (33, 34). This reduced capacity to respond against RSV

might be due to a direct disturbance of signaling by RSV itself (35), or an intrinsic

deficiency of neonatal pDCs to produce elevated amounts of cytokines (36-38). This

neonatal defect would prevent conventional DCs from adequate presentation of viral

antigens, contributing to poor specific T-cell responses (39).

Controversial is whether $\gamma\delta$ T-cells participate in viral clearance or immunopathology.

Herein, the $\gamma\delta$ T-cell depletion conflicted with a previous report. Indeed, children with

severe RSV bronchiolitis had reduced frequencies of $\gamma\delta$ T-cells in peripheral blood,

which was explained by the authors as a likely redistribution towards the lungs;

unfortunately, they could not evaluate their hypothesis (40). In line with our findings,

mice depleted of $\gamma\delta$ T-cells before RSV infection display increased viral titers (13). An

alternative hypothesis that would reconcile our study with others, is that γδ T-cell tissue

redistribution plays less of a role than its immature state in early life. Peripheral blood

γδ T-cells of infants with severe disease failed to produce IFN-γ when restimulated

(14), whereas γδ T-cells from neonatal mice had an impaired ability to produce IL-17

(15). Finally, Tc17 and Th17 cells were recently associated with shorter hospital stays

and proposed to play a protective role (26). Altogether, our data demonstrate that the

protective role of γδ T-cells in RSV infection is largely inefficient in early life.

The BALs of neonates displayed a high proportion of Tregs that declined rapidly to

levels comparable to healthy adults. As fetal CD4$^+$ T-cells tend to differentiate

preferentially into Tregs after stimulation (10), this tolerogenic mechanism is likely to

promote self-tolerance to the developing lung. We found that Treg response to RSV

infection happened very early in neonates, accompanying CD4$^+$ and CD8$^+$ T-cell

expansion rather than shutting them down afterwards. This strong Treg recruitment

was absent in adults. We speculate that the neonate immune system faces a dilemma

between avoiding potential autoimmune disorders and failing to mount a specific T-

cell response to harmful pathogens. Studies conducted in young infants with RSV

disease support this statement. A selective depletion of peripheral Tregs was shown,

probably due to massive recruitment to the lungs (41). A higher level of TGF-β

transcript was measured in neonatal DCs compared to adults (42). In contrast, some

reports emphasized the protective role of Tregs, since Treg-depleted young mice had

an enhanced RSV disease, with abundant T-cells exhibiting an activated phenotype

(43). Another remarkable observation was the strong correlation between pDC and

Treg counts. Despite the need for further investigations, it is worth noting that pDC to

Treg ratio was linked to the clearance *versus* persistence of human papillomavirus

(44).

Whether the cellular arm of the adaptive pulmonary response is detrimental or beneficial in RSV disease is still debated. Particularly, the contribution of IL-4 to viral clearance versus immunopathogenesis remains controversial (25, 45-47). Recently, a study performed in hospitalized infants showed a possible involvement of Tc2 cells in disease severity. Although convincing, this study was for obvious ethical reasons restricted to nasal aspirates (26). Interestingly, Tc2 cells are increasingly proposed to play an important role in the etiology of asthma (48). Herein, we provide a significant advancement in understanding this yet unclear IL-4 contribution, by showing that type 2 cytokine environment in the lung is a footprint of early life that decreases rapidly with time in healthy conditions. RSV infection circumvents this decline by mediating the massive recruitment of CD4[+] and CD8[+] T-cells, exacerbating the overall type 2 cytokine environment. We think that the link between IL-4 and RSV disease is the direct consequence of a strong trafficking of endogenous Th2 and Tc2 lymphocytes towards infected pulmonary tissues, rather than an RSV-specific type 2 immune response *per se*.

The high proportion of pulmonary T-cells producing spontaneously IL-4 was undetectable in the healthy adults. Neither was the RSV-specific immune response of adults oriented towards a Th2 or Tc2 profile. This corroborates a previous report mentioning an endogenously poised cytokine profile toward IL-4 in CD4[+] T-cells in cord blood of neonates and infants, establishing a link between IL-4 and development (49). However, we provide new insights by demonstrating that i) CD8[+] Tc2 cells are an equal and possibly superior source of IL-4 than CD4[+] Th2 cells; ii) this immunological state occurs during a short period *in vivo*, raising the likelihood that numerous animal trials failed to detect it due to the study design. Remarkably, this transient type 2 cytokine environment was exacerbated upon RSV infection. So far,

similar observations were only made at the systemic level or in nasal aspirates, and

authors could only speculate on the relocation to the lower respiratory tract (26, 50).

Herein, not only we show that this happens, but we also unify the link between IL-4

and RSV disease and endogenous IL-4 program in neonates. This finding constitutes

an important step towards deciphering why neonates and young infants are

particularly at risk.

Conclusion

Our holistic approach provided novel insights into the identification of the host's

contributing factors to RSV immunopathology, strengthening the hypothesis of an

inappropriate immune response in the narrow temporal window of early life. This can

now be characterized as a combined bronchoalveolar influx of Treg, Th2 and Tc2 cells,

associated with a strong depletion of $\gamma\delta$ T-cells. These findings constitute a novel basis

for further exploration of RSV immunopathogenesis and should be considered in RSV

vaccine design, which remains challenging after five decades of effort.

Acknowledgements

We thank Artur Summerfield, Christoph Aebi and Sean R. R. Hall for helpful discussions; Sylvie Python and Aurélie Godel for technical assistance; Inês Berenguer Veiga for participation in the animal trial; Hans Peter Lüthi, Roman Troxler, Jan Salchli, Daniel Brechbühl and Katarzyna Sliz for animal care.

References

1. Shi T, McAllister DA, O'Brien KL, Simoes EAF, Madhi SA, Gessner BD, Polack FP, Balsells E, Acacio S, Aguayo C, Alassani I, Ali A, Antonio M, Awasthi S, Awori JO, Azziz-Baumgartner E, Baggett HC, Baillie VL, Balmaseda A, Barahona A, Basnet S, Bassat Q, Basualdo W, Bigogo G, Bont L, Breiman RF, Brooks WA, Broor S, Bruce N, Bruden D, Buchy P, Campbell S, Carosone-Link P, Chadha M, Chipeta J, Chou M, Clara W, Cohen C, de Cuellar E, Dang DA, Dash-Yandag B, Deloria-Knoll M, Dherani M, Eap T, Ebruke BE, Echavarria M, de Freitas Lazaro Emediato CC, Fasce RA, Feikin DR, Feng L, Gentile A, Gordon A, Goswami D, Goyet S, Groome M, Halasa N, Hirve S, Homaira N, Howie SRC, Jara J, Jroundi I, Kartasasmita CB, Khuri-Bulos N, Kotloff KL, Krishnan A, Libster R, Lopez O, Lucero MG, Lucion F, Lupisan SP, Marcone DN, McCracken JP, Mejia M, Moisi JC, Montgomery JM, Moore DP, Moraleda C, Moyes J, Munywoki P, Mutyara K, Nicol MP, Nokes DJ, Nymadawa P, da Costa Oliveira MT, Oshitani H, Pandey N, Paranhos-Baccala G, Phillips LN, Picot VS, Rahman M, Rakoto-Andrianarivelo M, Rasmussen ZA, Rath BA, Robinson A, Romero C, Russomando G, Salimi V, Sawatwong P, Scheltema N, Schweiger B, Scott JAG, Seidenberg P, Shen K, Singleton R, Sotomayor V, Strand TA, Sutanto A, Sylla M, Tapia MD, Thamthitiwat S, Thomas ED, Tokarz R, Turner C, Venter M, Waicharoen S, Wang J, Watthanaworawit W, Yoshida LM, Yu H, Zar HJ, Campbell H, Nair H, Network RSVGE. Global, regional, and national disease burden estimates of acute lower respiratory infections due to respiratory syncytial virus in young children in 2015: a systematic review and modelling study. *Lancet* 2017; 390: 946-958.

2. Pneumonia Etiology Research for Child Health Study G. Causes of severe pneumonia requiring hospital admission in children without HIV infection from Africa and Asia: the PERCH multi-country case-control study. *Lancet* 2019; 394: 757-779.

3. Nair H, Nokes DJ, Gessner BD, Dherani M, Madhi SA, Singleton RJ, O'Brien KL, Roca A,
Wright PF, Bruce N, Chandran A, Theodoratou E, Sutanto A, Sedyaningsih ER,
Ngama M, Munywoki PK, Kartasasmita C, Simoes EA, Rudan I, Weber MW, Campbell
H. Global burden of acute lower respiratory infections due to respiratory syncytial virus
in young children: a systematic review and meta-analysis. *Lancet* 2010; 375: 1545-
1555.

4. Hall CB, Weinberg GA, Iwane MK, Blumkin AK, Edwards KM, Staat MA, Auinger P, Griffin
MR, Poehling KA, Erdman D, Grijalva CG, Zhu Y, Szilagyi P. The burden of respiratory
syncytial virus infection in young children. *N Engl J Med* 2009; 360: 588-598.

5. Murray J, Bottle A, Sharland M, Modi N, Aylin P, Majeed A, Saxena S, Medicines for
Neonates Investigator G. Risk factors for hospital admission with RSV bronchiolitis in
England: a population-based birth cohort study. *PLoS One* 2014; 9: e89186.

6. Garcia CG, Bhore R, Soriano-Fallas A, Trost M, Chason R, Ramilo O, Mejias A. Risk factors
in children hospitalized with RSV bronchiolitis versus non-RSV bronchiolitis. *Pediatrics*
2010; 126: e1453-1460.

7. Hall CB, Weinberg GA, Blumkin AK, Edwards KM, Staat MA, Schultz AF, Poehling KA,
Szilagyi PG, Griffin MR, Williams JV, Zhu Y, Grijalva CG, Prill MM, Iwane MK.
Respiratory syncytial virus-associated hospitalizations among children less than 24
months of age. *Pediatrics* 2013; 132: e341-348.

8. Palivizumab, a Humanized Respiratory Syncytial Virus Monoclonal Antibody, Reduces
Hospitalization From Respiratory Syncytial Virus Infection in High-risk Infants.
Pediatrics 1998; 102: 531-537.

9. Openshaw PJM, Chiu C, Culley FJ, Johansson C. Protective and Harmful Immunity to RSV
Infection. *Annu Rev Immunol* 2017; 35: 501-532.

10. Mold JE, Venkatasubrahmanyam S, Burt TD, Michaelsson J, Rivera JM, Galkina SA,
Weinberg K, Stoddart CA, McCune JM. Fetal and adult hematopoietic stem cells give
rise to distinct T cell lineages in humans. *Science* 2010; 330: 1695-1699.

11. de Kleer IM, Kool M, de Bruijn MJ, Willart M, van Moorleghem J, Schuijs MJ, Plantinga M,

Beyaert R, Hams E, Fallon PG, Hammad H, Hendriks RW, Lambrecht BN. Perinatal

Activation of the Interleukin-33 Pathway Promotes Type 2 Immunity in the Developing

Lung. *Immunity* 2016; 45: 1285-1298.

12. PrabhuDas M, Adkins B, Gans H, King C, Levy O, Ramilo O, Siegrist CA. Challenges in

infant immunity: implications for responses to infection and vaccines. *Nat Immunol*

2011; 12: 189-194.

13. Dodd J, Riffault S, Kodituwakku JS, Hayday AC, Openshaw PJ. Pulmonary V gamma 4+

gamma delta T cells have proinflammatory and antiviral effects in viral lung disease. *J*

Immunol 2009; 182: 1174-1181.

14. Aoyagi M, Shimojo N, Sekine K, Nishimuta T, Kohno Y. Respiratory syncytial virus

infection suppresses IFN-gamma production of gammadelta T cells. *Clin Exp Immunol*

2003; 131: 312-317.

15. Huang H, Saravia J, You D, Shaw AJ, Cormier SA. Impaired gamma delta T cell-derived

IL-17A and inflammasome activation during early respiratory syncytial virus infection

in infants. *Immunol Cell Biol* 2015; 93: 126-135.

16. McGill JL, Sacco RE. gammadelta T cells and the immune response to respiratory

syncytial virus infection. *Vet Immunol Immunopathol* 2016; 181: 24-29.

17. Restori KH, Srinivasa BT, Ward BJ, Fixman ED. Neonatal Immunity, Respiratory Virus

Infections, and the Development of Asthma. *Front Immunol* 2018; 9: 1249.

18. Levy O. Innate immunity of the newborn: basic mechanisms and clinical correlates. *Nat*

Rev Immunol 2007; 7: 379-390.

19. DeVincenzo JP, Wilkinson T, Vaishnaw A, Cehelsky J, Meyers R, Nochur S, Harrison L,

Meeking P, Mann A, Moane E, Oxford J, Pareek R, Moore R, Walsh E, Studholme R,

Dorsett P, Alvarez R, Lambkin-Williams R. Viral load drives disease in humans

experimentally infected with respiratory syncytial virus. *Am J Respir Crit Care Med*

2010; 182: 1305-1314.

20. Pino M, Kelvin DJ, Bermejo-Martin JF, Alonso A, Matias V, Tenorio A, Rico L, Eiros JM,

Castrodeza J, Blanco-Quiros A, Ardura J, de Lejarazu RO. Nasopharyngeal aspirate

cytokine levels 1 yr after severe respiratory syncytial virus infection. *Pediatr Allergy*

Immunol 2009; 20: 791-795.

21. Johnson JE, Gonzales RA, Olson SJ, Wright PF, Graham BS. The histopathology of fatal

untreated human respiratory syncytial virus infection. *Mod Pathol* 2007; 20: 108-119.

22. Piedra PA. Clinical experience with respiratory syncytial virus vaccines. *Pediatr Infect Dis*

J 2003; 22: S94-99.

23. Fishaut M, Tubergen D, McIntosh K. Cellular response to respiratory viruses with particular

reference to children with disorders of cell-mediated immunity. *J Pediatr* 1980; 96: 179-

186.

24. Roman M, Calhoun WJ, Hinton KL, Avendano LF, Simon V, Escobar AM, Gaggero A, Diaz

PV. Respiratory syncytial virus infection in infants is associated with predominant Th-

2-like response. *Am J Respir Crit Care Med* 1997; 156: 190-195.

25. Legg JP, Hussain IR, Warner JA, Johnston SL, Warner JO. Type 1 and type 2 cytokine

imbalance in acute respiratory syncytial virus bronchiolitis. *Am J Respir Crit Care Med*

2003; 168: 633-639.

26. Siefker DT, Vu L, You D, McBride A, Taylor R, Jones TL, DeVincenzo J, Cormier SA.

Respiratory Syncytial Virus Disease Severity Is Associated with Distinct CD8(+) T-Cell

Profiles. *Am J Respir Crit Care Med* 2020; 201: 325-334.

27. Olivier A, Gallup J, de Macedo MM, Varga SM, Ackermann M. Human respiratory syncytial

virus A2 strain replicates and induces innate immune responses by respiratory

epithelia of neonatal lambs. *Int J Exp Pathol* 2009; 90: 431-438.

28. Sitthicharoenchai P, Alnajjar S, Ackermann MR. A model of respiratory syncytial virus

(RSV) infection of infants in newborn lambs. *Cell Tissue Res* 2020; 380: 313-324.

29. Derscheid RJ, Ackermann MR. Perinatal lamb model of respiratory syncytial virus (RSV)

infection. *Viruses* 2012; 4: 2359-2378.

30. Blockus S, Sake SM, Wetzke M, Grethe C, Graalmann T, Pils M, Le Goffic R, Galloux M,

Prochnow H, Rox K, Huttel S, Rupcic Z, Wiegmann B, Dijkman R, Rameix-Welti MA,

Eleouet JF, Duprex WP, Thiel V, Hansen G, Bronstrup M, Haid S, Pietschmann T.

Labyrinthopeptins as virolytic inhibitors of respiratory syncytial virus cell entry. *Antiviral*

Res 2020; 177: 104774.

31. Mehinagic K, Pilo P, Vidondo B, Stokar-Regenscheit N. Coinfection of Swiss cattle with

bovine parainfluenza virus 3 and Mycoplasma bovis at acute and chronic stages of

bovine respiratory disease complex. *J Vet Diagn Invest* 2019; 31: 674-680.

32. Lee DC, Harker JA, Tregoning JS, Atabani SF, Johansson C, Schwarze J, Openshaw PJ.

CD25+ natural regulatory T cells are critical in limiting innate and adaptive immunity

and resolving disease following respiratory syncytial virus infection. *J Virol* 2010; 84:

8790-8798.

33. Kerrin A, Fitch P, Errington C, Kerr D, Waxman L, Riding K, McCormack J, Mehendele F,

McSorley H, MacKenzie K, Wronski S, Braun A, Levin R, Theilen U, Schwarze J.

Differential lower airway dendritic cell patterns may reveal distinct endotypes of RSV

bronchiolitis. *Thorax* 2017; 72: 620-627.

34. Ruckwardt TJ, Malloy AM, Morabito KM, Graham BS. Quantitative and qualitative deficits

in neonatal lung-migratory dendritic cells impact the generation of the CD8+ T cell

response. *PLoS Pathog* 2014; 10: e1003934.

35. Johnson TR, Johnson CN, Corbett KS, Edwards GC, Graham BS. Primary human mDC1,

mDC2, and pDC dendritic cells are differentially infected and activated by respiratory

syncytial virus. *PLoS One* 2011; 6: e16458.

36. Kollmann TR, Crabtree J, Rein-Weston A, Blimkie D, Thommai F, Wang XY, Lavoie PM,

Furlong J, Fortuno ES, 3rd, Hajjar AM, Hawkins NR, Self SG, Wilson CB. Neonatal

innate TLR-mediated responses are distinct from those of adults. *J Immunol* 2009;

183: 7150-7160.

37. Marr N, Wang TI, Kam SH, Hu YS, Sharma AA, Lam A, Markowski J, Solimano A, Lavoie

PM, Turvey SE. Attenuation of respiratory syncytial virus-induced and RIG-I-

dependent type I IFN responses in human neonates and very young children. *J*

Immunol 2014; 192: 948-957.

38. Danis B, George TC, Goriely S, Dutta B, Renneson J, Gatto L, Fitzgerald-Bocarsly P,

Marchant A, Goldman M, Willems F, De Wit D. Interferon regulatory factor 7-mediated

responses are defective in cord blood plasmacytoid dendritic cells. *Eur J Immunol*

2008; 38: 507-517.

39. Lau-Kilby AW, Turfkruyer M, Kehl M, Yang L, Buchholz UJ, Hickey K, Malloy AMW. Type

I IFN ineffectively activates neonatal dendritic cells limiting respiratory antiviral T-cell

responses. *Mucosal Immunol* 2020; 13: 371-380.

40. De Weerd W, Twilhaar WN, Kimpen JL. T cell subset analysis in peripheral blood of

children with RSV bronchiolitis. *Scand J Infect Dis* 1998; 30: 77-80.

41. Raiden S, Pandolfi J, Payaslian F, Anderson M, Rivarola N, Ferrero F, Urtasun M,

Fainboim L, Geffner J, Arruvito L. Depletion of circulating regulatory T cells during

severe respiratory syncytial virus infection in young children. *Am J Respir Crit Care*

Med 2014; 189: 865-868.

42. Thornburg NJ, Shepherd B, Crowe JE, Jr. Transforming growth factor beta is a major

regulator of human neonatal immune responses following respiratory syncytial virus

infection. *J Virol* 2010; 84: 12895-12902.

43. Durant LR, Makris S, Voorburg CM, Loebbermann J, Johansson C, Openshaw PJ.

Regulatory T cells prevent Th2 immune responses and pulmonary eosinophilia during

respiratory syncytial virus infection in mice. *J Virol* 2013; 87: 10946-10954.

44. Strickler HD, Martinson J, Desai S, Xie X, Burk RD, Anastos K, Massad LS, Minkoff H,

Xue X, D'Souza G, Levine AM, Colie C, Watts DH, Palefsky JM, Landay A. The relation

of plasmacytoid dendritic cells (pDCs) and regulatory T-cells (Tregs) with HPV

persistence in HIV-infected and HIV-uninfected women. *Viral Immunol* 2014; 27: 20-

25.

45. Caballero MT, Serra ME, Acosta PL, Marzec J, Gibbons L, Salim M, Rodriguez A, Reynaldi

A, Garcia A, Bado D, Buchholz UJ, Hijano DR, Coviello S, Newcomb D, Bellabarba M,

Ferolla FM, Libster R, Berenstein A, Siniawaski S, Blumetti V, Echavarria M, Pinto L,

Lawrence A, Ossorio MF, Grosman A, Mateu CG, Bayle C, Dericco A, Pellegrini M,

Igarza I, Repetto HA, Grimaldi LA, Gudapati P, Polack NR, Althabe F, Shi M, Ferrero

F, Bergel E, Stein RT, Peebles RS, Boothby M, Kleeberger SR, Polack FP. TLR4

genotype and environmental LPS mediate RSV bronchiolitis through Th2 polarization.

J Clin Invest 2015; 125: 571-582.

46. Bermejo-Martin JF, Garcia-Arevalo MC, De Lejarazu RO, Ardura J, Eiros JM, Alonso A,

Matias V, Pino M, Bernardo D, Arranz E, Blanco-Quiros A. Predominance of Th2

cytokines, CXC chemokines and innate immunity mediators at the mucosal level during

severe respiratory syncytial virus infection in children. *Eur Cytokine Netw* 2007; 18:

162-167.

47. Vu LD, Siefker D, Jones TL, You D, Taylor R, DeVincenzo J, Cormier SA. Elevated Levels

of Type 2 Respiratory Innate Lymphoid Cells in Human Infants with Severe Respiratory

Syncytial Virus Bronchiolitis. *Am J Respir Crit Care Med* 2019; 200: 1414-1423.

48. Hinks TSC, Hoyle RD, Gelfand EW. CD8(+) Tc2 cells: underappreciated contributors to

severe asthma. *Eur Respir Rev* 2019; 28.

49. Hebel K, Weinert S, Kuropka B, Knolle J, Kosak B, Jorch G, Arens C, Krause E, Braun-

Dullaeus RC, Brunner-Weinzierl MC. CD4+ T cells from human neonates and infants

are poised spontaneously to run a nonclassical IL-4 program. *J Immunol* 2014; 192:

5160-5170.

50. Bendelja K, Gagro A, Bace A, Lokar-Kolbas R, Krsulovic-Hresic V, Drazenovic V, Mlinaric-

Galinovic G, Rabatic S. Predominant type-2 response in infants with respiratory

syncytial virus (RSV) infection demonstrated by cytokine flow cytometry. *Clin Exp*

Immunol 2000; 121: 332-338.

  Figure legends

Figure 1. Neonatal RSV infection causes bronchiolitis and interstitial

pneumonia. (A) Illustration of the transtracheal RSV infection experimental design.

(B) RSV was quantified in the respiratory tract (BAL, cell fraction; Lung; BAL, fluid) by

qPCR. (C) RSV detection by immunohistochemistry in macrophages, and type I

pneumocytes. (D) Representative histological lung tissue sections from neonates over

the course of RSV infection (40X, HE). Lungs of RSV-infected animals (strain A2)

show interstitial septae expanded by inflammatory cells (interstitial pneumonia) and

accumulation of inflammatory cells within alveoli at days 3 and 6 p.i. No lesions at days

14 and 42 p.i. The lungs of mock neonates allow comparison with normal lung

architecture within the first weeks of life. Each symbol represents an individual animal

(healthy adults, n=3; mock neonates, n=3; neonates infected with RSV, n=6). Boxplots

indicate median value (center line) and interquartile ranges (box edges), with whiskers

extending to the lowest and the highest values. Stars indicate significance levels. *, p

< 0.05.

Figure 2. Neonatal RSV infection activates the initiators of the adaptive immune

response and induces a transient depletion of $\gamma\delta$ T-cells. (A-E) Neonatal RSV

infection induces bronchoalveolar space recruitment of activated pDCs. (A) FCM

gating strategy for ovine pDCs. (B) Neonatal RSV infection induces an early and

transient bronchoalveolar space colonization by pDCs. (C) Correlation coefficient (r)

obtained with pDC counts calculated as a function of RSV copies/10^{12} 18S copies.

Significance was reached for BAL fluid but not for BAL, cell fraction. (D-E) A high

frequency of recruited pDCs in bronchoalveolar space upregulate CD86 (D,

representative FCM contour plots; E, plot displaying all individuals). (F-J) Neonatal

RSV infection leads to the transient depletion of γδ T-cell subset in the bronchoalveolar

space, followed by a full replenishment and an accelerated maturation. (F) FACS

gating strategy for ovine γδ T-cells. (G) Neonatal RSV infection induces a decrease in

γδ T-cells number in the bronchoalveolar space. (H) Correlation coefficient (r) obtained

with in γδ T-cell counts calculated as a function of % Live/Dead$^+$. (I-J) Replenishment

and maturation (CD25 up-regulation) of γδ T-cell subset following the depletion after

RSV infection (I, representative FACS contour plots; J, Pie charts displaying all

individuals). Each symbol represents an individual animal (healthy adults, n=3; mock

neonates, n=3; neonates infected with RSV, n=6). Boxplots indicate median value

(center line) and interquartile ranges (box edges), with whiskers extending to the

lowest and the highest values. Stars indicate significance levels. *, p < 0.05.

Figure 3. Neonatal RSV infection exacerbates the type 2 cytokine environment

in lung. (A) FCM gating strategy for ovine CD4$^+$ and CD8$^+$ T-cells. (B) RSV infection

in neonate accelerates the recruitment of CD4$^+$ and CD8$^+$ T-cells in the

bronchoalveolar space. (C-D) A high frequency of recruited CD4$^+$ T-cells in

bronchoalveolar space differentiate towards a Th2 cytokine profile (IL-4 production)

(C, representative FACS contour plots; D, plot displaying all individuals). (E-F) A high

frequency of recruited CD8$^+$ T-cells in bronchoalveolar space differentiate towards a

Tc2 cytokine profile (IL-4 production) (E, representative FCM contour plots; F, plot

displaying all individuals). (G) RSV infection in neonates modulates the activation (cell

surface expression of CD25) and the maturation (cell surface expression of T-cell

memory marker CD45RO) of CD4$^+$ and CD8$^+$ T-cell pools in the lung. (H) Neonates

infected with RSV can mount an antiviral neutralizing-antibody response. (I) Anti-RSV

Nabs persist over a 42-day long period. Each symbol represents an individual animal

(healthy adults, n=3; mock neonates, n=3; neonates infected with RSV, n=6). Boxplots

indicate median value (center line) and interquartile ranges (box edges), with whiskers

extending to the lowest and the highest values. Stars indicate significance levels. *, p

< 0.05; **, $p < 0.01$).

Figure 4. Neonatal RSV infection rapidly induces Treg suppressive functions to

dampen the Th2 and Tc2 responses. (A) FCM gating strategy for ovine CD4$^+$ and

CD8$^+$ T-cells. (B) RSV infection in neonates led to the recruitment of Tregs in the

bronchoalveolar space (left panel), but not in the lung (right panel). (C) Progressive

decrease with age of Treg subset among the CD4$^+$ T-cell fraction (representative FCM

contour plots). (D) Frequency of TGF-β-producing Tregs. (E) Correlation coefficient (r)

obtained in BALs with Treg counts calculated as a function of non-Treg CD4$^+$ counts

(left), Treg counts calculated as a function of CD8$^+$ counts (middle) and CD8$^+$ counts

calculated as a function of non-Treg CD4$^+$ counts (right). Significance was reached for

the three correlations. (F) CD8$^+$ T-cell counts remained stable in the peribronchial LNs

over the course of neonatal RSV infection. (G) Role of neonatal RSV infection on

cytokine production by CD8$^+$ T-cells in draining LNs. Upper panel: poor influence on

the frequency of IL-4$^+$ CD8$^+$ T-cells (representative FCM contour plots). Lower panel:

slight, but significant enhancement of IL-17$^+$ CD8$^+$ T-cells (representative FCM contour

plots). Each symbol represents an individual animal (healthy adults, n=3; mock

neonates, n=3; neonates infected with RSV, n=6). Boxplots indicate median value

(center line) and interquartile ranges (box edges), with whiskers extending to the

lowest and the highest values. Stars indicate significance levels. *, $p < 0.05$; **, $p <$

0.01.

Figure 5. Distinct cellular immune response in neonates compare to adults following RSV infection. (A) RSV was quantified in the respiratory tract (BAL, cell fraction; lung; BAL, fluid) of adults and neonates by qPCR. (B) Representative histological lung tissue sections from RSV-infected neonates and adults at day 6 and 14 p.i. (40X, HE). Adults show no histological lesions, whereas neonates show interstitial pneumonia at day 6.p.i. (C-D) Counts of recruited pDCs and CD86 upregulation in bronchoalveolar space (C, plot displaying all individuals; D, representative FCM contour plots). (E-G) $\gamma\delta$ T-cell counts, cytokine production, and maturation in the bronchoalveolar space (E, plot displaying all individuals; F, representative FCM contour plots; G, pie charts integrating all animals). (H-J) Recruitment, activation, and maturation of the $CD4^+$ T-cell pool in the bronchoalveolar space (H, plot displaying all individuals; I, representative FACS contour plots; J, pie charts integrating all animals). (K-M) As in (H-J), but showing $CD8^+$ T-cell pool. (N-O) Recruitment of Tregs and TGF-β production in the bronchoalveolar space (N, plot displaying all individuals; O, representative FACS contour plots). Each symbol represents an individual animal (infected adults, n=3; neonates infected with RSV, n=6). The grey zone corresponds to the interval integrating the values of all corresponding mock controls. Boxplots indicate median value (center line) and interquartile ranges (box edges), with whiskers extending to the lowest and the highest values. Stars indicate significance levels. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

Figure 6. The severity of disease relates to the magnitude of bronchoalveolar T-cell responses. (A) Representative histological lung tissue sections from mock neonates and neonates infected with RSV-ON1-H1 and RSV A2 strain (40X, HE). At day 6 p.i. the RSV-ON1-H1 infected animals show mild interstitial pneumonia (upper right corner) with accumulation of inflammatory cells within alveoli. No lesions at day 14 p.i. or mock-infected animals. (B) Comparison of immune cell subset trafficking into bronchoalveolar space after infection with RSV-ON1-H1 or RSV A2 strains. (C) Correlation coefficient (r) obtained in BALs with Treg counts calculated as a function of pDC counts. Each symbol represents an individual animal (infected adults, n=3; neonates infected with RSV, n=6). The grey zone corresponds to the interval integrating the values of all corresponding mock controls. Boxplots indicate median value (center line) and interquartile ranges (box edges), with whiskers extending to the lowest and the highest values. Stars indicate significance levels. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$.

Data availability statement

700   **Data availability statement**

701   The data generated and analysed during the current study are available from the

702   corresponding author on reasonable request.

Figure 1

705   **Figure 2**

Figure 3

709　**Figure 4**

Figure 5

714 **Figure 6**