1   **ELAV-like proteins bind and stabilize C/EBP mRNA in the induction of long-term**
2   **memory in *Aplysia***
3

4   Anastasios A. Mirisis[1], Ashley M. Kopec[2], and Thomas J. Carew[1]*
5
6   [1]Center for Neural Science, New York University, New York, NY, USA
7
8   [2]Department of Neuroscience and Experimental Therapeutics, Albany Medical College,
9   Albany, NY, USA
10
11   *Address correspondence to the Lead Contact: Thomas J. Carew, Ph.D.
12
Center for Neural Science

New York University

4 Washington Place

Room 621

New York, NY 10003

Email: tcarew@nyu.edu

Phone: (212) 998-7940

Running title: Post-transcriptional regulation by ELAV mediates LTM formation

Keywords: growth factor; long-term memory; Aplysia; ELAV; c/ebp; TGFβ; post-

transcriptional regulation;

Number of Figures: 5 + 1 Table

Number of Supplementary Figures: 3 + 1 Table

Conflict of Interest: The authors declare no competing financial interests.

1 **Summary**

Signaling through distinct growth factor pathways is required for long-term

memory (LTM) in *Aplysia*. In a two-trial paradigm, Trial 1 TrkB signaling is required for

expression of the immediate early gene *apc/ebp*, which is transient and necessary, but

insufficient, for LTM formation. A prolonged increase in *apc/ebp* gene expression, which

is both necessary and sufficient for LTM is achieved through TGFβ signaling during Trial

2. Here we demonstrate: (1) the prolonged increase in *apc/ebp* expression is not

dependent on *de novo* transcription but is dependent on p38 MAPK activity induced by

TGFβ; (2) Trial 2 increases the interaction between *apc/ebp* mRNA and the stabilizing

RNA-binding protein ApELAV; and (3) inhibition of the *apc/ebp* mRNA-ApELAV

interaction blocks induction of LTM. These results reveal a novel role for post-

transcriptional gene regulation: the stabilization of a specific transcript, *apc/ebp*, by

ELAV-like proteins, as a critical molecular step in LTM formation.

Introduction

Long-term memory (LTM) can be mechanistically distinguished from short-term and intermediate-term memory by its dependence on *de novo* transcription (Alberini, 2009; Kandel, 2012; Sutton et al., 2001). LTM for sensitization of the defensive tail and siphon withdrawal responses in *Aplysia californica* provides an experimentally advantageous system in which to study the molecular mechanisms underlying LTM formation. In this system, training trials (brief electrical shocks to the tail) lead to the release of the neuromodulator serotonin (5HT) in the CNS of *Aplysia* (Marinesco and Carew, 2002), which in turn enhances synaptic transmission in the reflex circuit (Brunelli et al., 1976). Studies focusing on the sensory neurons (SNs) mediating these reflexes have elucidated critical molecular events underlying LTM for sensitization. For example, initiation of CREB-mediated transcription of several target genes, including the immediate early gene *Aplysia c/ebp* (*apc/ebp*), is required for LTM formation (Alberini, 2009; Alberini et al., 1994; Kandel, 2012; Mirisis et al., 2016). The constitutive activation of a C/EBP complex by PKA alone can support long-term synaptic facilitation in the absence of CREB-mediated transcription (Lee et al., 2006). Moreover, C/EBP can regulate the transcription of a wide variety of genes by means of dimerization with different molecular partners and by generating a range of post-translational modifications (Tsukada et al., 2011).

As in many other species, the induction of LTM in *Aplysia* critically depends upon the number and pattern of training trials. For example, two training trials separated by 45 min (but not by 15 or 60 min), induces LTM formation (Philips et al., 2013), revealing a narrow time window during which a second trial interacts with a "molecular context"

produced by the first trial to give rise to LTM formation (Kopec et al., 2015; Philips et al.,

2013).

In this paper, we explore the role of growth factors (GFs) in LTM formation, with

an emphasis on regulation that occurs between transcription and translation within the

molecular context (Philips et al., 2013) established by Trial 1. In many systems, GF-

mediated regulation of RNA-binding proteins modulates the stability of gene transcripts

(Bai et al., 2012; Ge et al., 2016). Moreover, many of the canonical GF signaling

cascades engaged during development are re-engaged to support plasticity in the adult

(Kopec and Carew, 2013; Kopec et al., 2015; Lu et al., 2011). Because converging

signaling cascades and phenotypically similar functional outcomes are often induced by

different GFs, the relative contribution of each individual GF can be difficult to determine

(Kopec and Carew, 2013). To examine this question, Kopec et al. (2015) used the two-

trial training paradigm (Philips et al., 2013), and found that signaling through two distinct

GF families, TrkB and TGFβ, is required during different training trials for the induction

of the molecular events critical for LTM formation. Specifically, *apc/ebp* gene expression

induced by Trial 1 peaked at 45 mins, was dependent on TrkB signaling and returned to

baseline by 1 hr (Kopec et al., 2015; Philips et al., 2013). However, if Trial 2 was

delivered during peak *apc/ebp* mRNA levels at 45 min after Trial 1, *apc/ebp* mRNA level

remained elevated for more than 1 hr, and this elevation critically required TGFβ

signaling during Trial 2. In parallel behavioral experiments, Kopec et al. (2015) found

that the induction of LTM required TrkB signaling during Trial 1 and TGFβ signaling

during Trial 2.

Importantly, Trial 2-dependent *apc/ebp* gene expression (1 hr after Trial 1) was

directly dependent on the TrkB-mediated events set in motion by Trial 1 (Kopec et al.,

2015). This dependence could be explained in one of two ways: (i) TGFβ signaling in

Trial 2 also induces *apc/ebp* gene expression in Trial 2, or (ii) TGFβ signaling in Trial 2

engages post-transcriptional mechanisms which act to stabilize the *apc/ebp* mRNA that

was generated by TrkB signaling in Trial 1.

In considering the second alternative, a candidate mechanism for the

stabilization of the mRNA is the RNA-binding protein ELAV (embryonic lethal abnormal

vision), which confers stability to mRNAs containing AU-rich elements (AREs) in their 3'

UTR (Amadio et al., 2014; Brennan and Steitz, 2001; Pascale et al., 2005; Wang et al.,

2000). Since *apc/ebp* mRNA contains multiple AREs in its 3' UTR, it is a potential target

for stabilization by ApELAV (Yim et al., 2006). Two ELAV family members, ApELAV1

and ApELAV2, have been identified and cloned in *Aplysia*, and ApELAV1 has been

shown to bind to *apc/ebp* mRNA both *in vitro* and *in vivo* (Yim et al., 2006). Interestingly,

microinjection of the 3' UTR of *apc/ebp* mRNA, which contains the putative binding sites

(AREs) for ApELAV, into SNs, blocks the induction of long-term synaptic facilitation

(LTF) in sensorimotor neuron (SN-MN) co-culture (Yim et al., 2006). Moreover, ELAV

proteins have been implicated in memory formation in several mammalian model

systems, pointing to a possible evolutionarily conserved role for post-transcriptional

regulation in memory formation (Bolognani et al., 2004; Quattrone et al., 2001). In the

present paper, we first examined whether TGFβ signaling during Trial 2 actively induces

apc/ebp gene expression, or alternatively, whether Trial 2-dependent TGFβ induces the

prolonged expression of *apc/ebp* mRNA observed at 1 hr via engagement of ELAV-

dependent transcript stabilization, which would indicate a critical role for ApELAV during

Trial 2 in LTM formation.

We here report that: (i) The Trial 1-dependent increase in *apc/ebp* mRNA is

transcription-dependent, whereas the Trial 2-dependent increase in *apc/ebp* mRNA is

not dependent on *de novo* transcription; (ii) The Trial 2-dependent increase in *apc/ebp*

mRNA requires p38 MAPK activation downstream of TGFβ signaling; (iii) Treatment

with recombinant human TGFβ-1 at 45 min, even in the presence of transcriptional

inhibitors, is sufficient for increased *apc/ebp* mRNA at 1 hr after Trial 1; (iv) The

interaction between ApELAV and *apc/ebp* mRNA is increased after Trial 2, and this

increase critically depends on TGFβ signaling; (v) An inhibitor of ELAV-mRNA

interactions (CMLD-2) significantly reduces the level of Trial 2-dependent *apc/ebp*

mRNA at 1 hr; and (vi) CMLD-2 blocks the behavioral induction of 2-Trial LTM.

Collectively, these results reveal a novel role for a particular form of post-transcriptional

regulation: stabilization of a specific transcript (*apc/ebp*) by ELAV-like proteins, as a

necessary molecular step in LTM formation.

Results

The Trial 1-dependent increase in *apc/ebp* mRNA level is transcription-dependent

We have shown that *apc/ebp* gene expression is increased 45 min following Trial

1 (Kopec et al., 2015; Philips et al., 2013). Thus, we first asked whether Trial 1-

dependent increase in *apc/ebp* gene expression was due to (i) increased gene

transcription, or (ii) alternative mechanisms, such as an increase in transcript stability of

baseline levels of *apc/ebp* mRNA.

Figure 1

Figure 1. Trial 1 initiates a transcription-dependent increase in *apc/ebp* mRNA levels, whereas Trial 2 prolongs this gene expression through a transcription-independent mechanism.

(a) (Top) Experimental paradigm. Trial 1 is delivered at time=0:00 and SN somata are collected at 45 min following the onset of Trial 1. Dissected ganglia are treated with actinomycin D 15 min before Trial 1. Drug conditions are expressed as a ratio to vehicle conditions. **(Bottom)** Blocking transcription before Trial 1 significantly disrupts *apc/ebp* gene expression at 45 min. Histograms are labeled by their corresponding drug condition. In this and all subsequent figures, unless specified otherwise, data are displayed as aligned dot plots. Means are represented as red lines within each histogram. Within-group statistical significance is displayed with an asterisk above histograms, and between-group statistical significance is indicated above the histograms being compared. *p<0.05; n=7-8

(b) (Top) Experimental paradigm. SN somata are collected at 1 hr following the onset of Trial 1. Dissected ganglia are treated with actinomycin D 15 min before Trial 2. Drug conditions were expressed as a ratio to vehicle conditions. **(Bottom)** A second training trial (Trial 2) at 45 min following the onset of Trial 1 results in a significant transcription-independent significant increase in *apc/ebp* gene expression at 1 hr. *p<0.05, **p<0.01; n=7-11.

(c) (Top) Experimental paradigm. Trial 1 is delivered and SN somata are collected at 1 hr following the onset of Trial 1. Dissected ganglia are treated with actinomycin D or vehicle 15 min before washout (light yellow line) and TGFβ-1 or vehicle is applied at 45 min after Trial 1 until collection. Drug conditions were expressed as a ratio to vehicle conditions. **(Bottom)** Treatment with TGFβ-1 alone at 45 min results in a transcription-independent increase in *apc/ebp* mRNA level. *p<0.05; n=7-12

In order to distinguish between these possibilities, dissected pleural-pedal

ganglia were incubated with the irreversible transcription inhibitor Actinomycin D (40

µg/mL, Sigma) for 15 min, and contralateral within-animal control ganglia were

incubated with vehicle (an equivalent volume of ddH$_2$O). Following drug washout,

experimental ganglia were exposed to 5HT (50 µM in ASW, Sigma) for 5 min (Trial 1) to

simulate a tail shock training trial (Philips et al., 2013). 5HT was washed out with fresh

ASW, and pleural SN clusters were collected at 45 min and analyzed for *apc/ebp* gene

expression via quantitative PCR. In this and all following experiments, data from drug

conditions are expressed as a ratio to vehicle conditions. Consistent with previous

results (Kopec et al., 2015), in experimental ganglia which had not received inhibitor

pre-treatment, a within-group comparison revealed a significant increase in *apc/ebp*

gene expression in SN clusters collected 45 min after Trial 1 (Figure 1a; Table 1A). No

significant difference in experimental (+ActinoD) and within-animal control samples

(contralateral SN clusters) was observed (Figure 1a; Table 1B). A between-group

comparison revealed a significant difference between *apc/ebp* gene expression of SN

clusters which had received drug treatment compared to SN clusters which had not

received drug treatment (Figure 1a; Table 1C), demonstrating that Trial 1-dependent

apc/ebp gene expression at 45 min after Trial 1 is dependent on *de novo* gene

transcription.

The Trial 2-dependent increase in *apc/ebp* mRNA level is transcription-

independent

When Trial 2 is delivered 45 min after Trial 1, *apc/ebp* gene expression at 1 hr

after Trial 1 is dependent on (i) TrkB signaling during Trial 1, and (ii) TGFβ signaling

during Trial 2 (Kopec et al., 2015). This suggests that either (i) Trial 2 TGFβ-dependent

mechanisms cause a new wave of *apc/ebp* transcription, or (ii) Trial 2 induces a form of

transcription-independent stabilization of *apc/ebp* mRNA synthesized from Trial 1 (via

TrkB-dependent mechanisms). Ganglia were treated with Actinomycin D for 15 min

before Trial 2. Following drug washout, experimental ganglia were exposed to 5HT for 5

min (Trial 2). 5HT was washed out with fresh ASW, and pleural SN clusters were

collected at 1 hr after Trial 1. A within-group analysis revealed a significant increase in

apc/ebp mRNA levels in ganglia which had received Actinomycin D (Figure 1b; Table

1D). ANOVA revealed a significant difference among the three groups (Table 1E).

Subsequent planned t-tests revealed a significant increase in *apc/ebp* mRNA levels in

both the 2X 5HT -ActinoD and 2X 5HT +ActinoD groups, compared to ganglia that had

received only 1X 5HT, and no significant difference was noted between the 2X 5HT -

ActinoD and 2X 5HT +ActinoD groups (Figure 1b; Table 1F). These data demonstrate

that the Trial 2-dependent increase in *apc/ebp* mRNA level at 1 hr after Trial 1 is <u>not</u>

dependent on *de novo* gene transcription and supports the hypothesis that the

increased *apc/ebp* mRNA level may result from TGFβ-dependent mRNA stabilization.

TGFβ-1 treatment at 45 min is sufficient for Trial 2-dependent increase in *apc/ebp*

mRNA level

By sequestering endogenously released TGFβ-like ligands during Trial 2, we

have previously shown that intact TGFβ signaling is required during Trial 2 for *apc/ebp*

gene expression at 1 hr (Kopec et al., 2015). We here sought to determine whether

treatment with recombinant human TGFβ-1 alone could duplicate these effects.

To investigate this question, ganglia were treated with TGFβ-1 (100 ng/mL, R&D

Systems) starting at 45 min after Trial 1 until SN cluster collection and lysis at 1 hr after

Trial 1. Treatment with TGFβ-1 has previously been demonstrated to (i) recruit

persistent ERK signaling, a necessary step in LTM formation, and (ii) induce long-term

facilitation (LTF) in isolated *Aplysia* ganglia (Shobe et al., 2016; Zhang et al., 1997).

Contralateral control ganglia were treated with an equivalent volume of vehicle (0.1%

BSA in ASW). Treatment with TGFβ-1 resulted in a significant increase in *apc/ebp*

mRNA levels when compared to a within-animal control of SN clusters that had received

only vehicle (Figure 1c; Table 1G). Planned comparisons further revealed a significant

difference between the Trial 1 only and TGFβ-treated groups (Figure 1c; Table 1H).

Given that TGFβ-1 treatment (in the absence of 5HT) at 45 min after Trial 1 is

sufficient to increase *apc/ebp* mRNA level at 1 hr (Figure 1c), we further asked whether

the TGFβ signaling-dependent effect of increased *apc/ebp* mRNA level at 1 hr after Trial

1 required transcription. Ganglia were treated at 30 min after Trial 1 with Actinomycin D.

At 45 min after Trial 1, Actinomycin D was washed out and ganglia were immediately

treated with TGFβ-1 until SN cluster collection and lysis at 1 hr after Trial 1.

Contralateral within-animal control ganglia received vehicle (0.1% BSA in ASW and

ddH$_2$O). TGFβ-1 treatment concomitant with a transcriptional block still resulted in

significantly increased *apc/ebp* mRNA level when compared to within-animal control SN

clusters. Moreover, there were no significant differences between groups which

received TGFβ treatment alone and TGFβ treatment with ActinoD (Figure 1c: Table 1I).

Collectively, these data support the hypothesis that TGFβ signaling during Trial 2 is

responsible for the post-transcriptional stabilization effect on *apc/ebp* mRNA.

p38 MAPK is activated by two-trial training and is blocked by blocking TGFβ

signaling

Since ELAV is known to be an RNA-binding protein yielding a stabilizing effect on its target mRNAs (Brennan and Steitz, 2001; Yim et al., 2006), we explored the possibility that *apc/ebp* gene expression is modulated by ApELAV. How might ELAV be recruited to exert its stabilizing effects on target mRNAs? First, ELAV protein levels could be increased, or second, ELAV protein could be covalently modified, which would promote its ability to bind its target transcripts.

To distinguish between these possibilities, we first asked whether the protein levels of ApELAV were regulated by two-trial training. To assay changes in protein levels, we used an antibody against HuR, a mammalian member of the ELAV family. This antibody recognizes both *Aplysia* ELAV family members, ApELAV1 and ApELAV2 (Supplementary Figure 1a).

At the 1 hr timepoint (15 min after the onset of Trial 2), pleural SN clusters were collected and subjected to SDS-PAGE and Western blotting. Our results revealed no difference in ApELAV protein levels between the experimental and within-animal control ganglia, which had not received two-trial training (Supplementary Figure $1b_2$).

These findings, however, do not preclude the possibility of a post-translational modification of existing ApELAV proteins (such as phosphorylation) downstream of TGFβ signaling, which would result in an increase of ApELAV-*apc/ebp* mRNA association. In support of this possibility, a particular kinase, p38 MAPK, is both activated downstream of TGFβ signaling (Yamashita et al., 2008), and has previously been identified as a critical mediator of ELAV activation by phosphorylation(Bai et al.,

2012; Eberhardt et al., 2012; Lafarga et al., 2009; Pascale et al., 2005; Slone et al.,

2016). Our experiments revealed a significant increase in p38 MAPK activation

following two-trial training (Figure 2a: Table 1J). This increase was blocked either (i) by

eliminating Trial 2, or (ii) by delivering Trial 2 in the presence of TGFβsRII (5 µg/mL,

R&D Systems), which acts as an inhibitor of TGFβ signaling by sequestering

extracellular ligands (Figure 2a; Table 1K). These data demonstrate that p38 MAPK

activation at 1 hr is dependent on TGFβ signaling and may serve an intermediate role

between TGFβ signaling and ELAV activation.

Figure 2

a

b_1

2X ASW +DMSO   2X 5HT +DMSO   2X 5HT +SB

b_2

c_1

c_2

Figure 2. ELAV-mediated prolonged expression of *apc/ebp* mRNA requires TGFβ signaling initiated by p38 MAPK.
(a) (Top) Experimental paradigm. In all paradigms, SN somata are collected at 1 hr following the onset of Trial 1. Dissected ganglia are treated with CMLD-2, SB203580, or TGFβsRII from 15 min before Trial 2 until collection. Dissected ganglia are treated with TGFβ-1 at 45 min until collection. Dissected ganglia are treated with CMLD-2 from 15 min before TGFβ-1 application until collection. Drug conditions were expressed as a ratio to vehicle conditions. **(Bottom)** Following two-trial training, there is a significant increase in p38 MAPK activation which is blocked by treatment with TGFβsRII. *$p < 0.05$; n=6-7
(b_1) Representative maximum projection images of single isolated SNs.
(b_2) Puncta abundance in isolated SNs. Data are represented as a percentage of puncta abundance in SNs which received two-trial training in the presence of vehicle without 5HT. **$p < 0.01$; n=19-23
(c_1) Blocking the ApELAV-*apc/ebp* mRNA interaction significantly disrupts the increase in *apc/ebp* mRNA level at 1 hr after Trial 1. Inhibiting p38 MAPK activity similarly disrupts the increase in *apc/ebp* mRNA level. *$p < 0.05$, **$p < 0.01$; *n*=7-11.
(c_2) Blocking the ApELAV-mRNA interaction blocks the increase in *apc/ebp* mRNA level by TGFβ #$p < 0.3$, *$p < 0.05$; n=7-8

p38 MAPK activity is required for prolonged increase in *apc/ebp* mRNA levels

To further examine the role of p38 MAPK in the stabilization of *apc/ebp* mRNA, we directly measured *apc/ebp* mRNA levels, by performing single molecule fluorescence *in situ* hybridization (smFISH) on isolated *Aplysia* SNs (Figure 2b). This technique allows for fluorescent visualization of individual *apc/ebp* RNA molecules as puncta resulting from the combined fluorescence of 33 antisense DNA probes (LGC Biosearch Technology). ANOVA revealed a significant difference in puncta abundance between: (i) SNs that received two-trial training in the presence of vehicle, (ii) SNs which received two-trial training in the presence of SB203580, a p38 MAPK inhibitor previously used and characterized in *Aplysia* (Zhang et al., 2017) (10 µM, Tocris), and (iii) SNs that received two-trial training without 5HT and in the presence of vehicle (Figure 2b$_2$; Table 1L). Planned t-tests revealed a significant increase in puncta in SNs that were treated with two-trial training with 5HT in the presence of DMSO compared to each of the other groups (Figure 2b$_2$;Table 1M).

In order to examine whether p38 MAPK activation is important for the increase in *apc/ebp* mRNA level 1 hr after Trial 1, ganglia received Trial 2 training with 5HT in the presence of SB203580. 5HT was washed out following Trial 2, and SB203580 treatment proceeded until SN cluster collection and lysis at 1 hr. Within-animal control ganglia were treated with vehicle (0.1% DMSO) but did not receive two-trial training. Treatment with SB203580 indeed resulted in a significant block of the Trial 2-dependent increase in *apc/ebp* mRNA level when compared to two-trial training alone (Figure 2c$_1$; Table 1N).

ApELAV-mRNA interaction is required for prolonged increase in *apc/ebp* mRNA

levels

In order to directly manipulate the ability of ApELAV to interact with its target

transcripts, we used CMLD-2 (100 µM, Millipore), a potent inhibitor of HuR-ARE

interaction (Muralidharan et al., 2017; Slone et al., 2016; Wu et al., 2015). HuR is a

mammalian homolog of ELAV, and also contains three highly conserved RNA

recognition motifs which facilitate its binding to AREs of mRNAs. Since CMLD-2 directly

binds to the highly evolutionary conserved RNA-binding pocket of HuR, we

hypothesized that it would likewise bind and inhibit ApELAV binding. We found that

following treatment of ganglia with Actinomycin D to inhibit basal *de novo* transcription

of *apc/ebp*, addition of CMLD-2 accelerates *apc/ebp* mRNA degradation compared to

control (0.8% acetonitrile and ddH$_2$O) (Supplementary Figure 2a$_2$).

In order to examine whether the ApELAV-*apc/ebp* mRNA interaction plays a

critical role in increased *apc/ebp* mRNA levels during the two-trial training paradigm,

ganglia received Trial 2 in the presence of CMLD-2. 5HT was washed out following Trial

2, and CMLD-2 treatment proceeded until SN cluster collection and lysis at 1 hr. Within-

animal control ganglia were treated with vehicle but did not receive two-trial training. We

found that treatment with CMLD-2 significant blocked the increase of Trial 2-dependent

apc/ebp mRNA levels when compared to two-trial training alone (Figure 2c$_1$; Table 1O).

Treatment with vehicle alone did not affect the Trial 2-dependent increase in *apc/ebp*

mRNA level (data not shown). A similar effect on *apc/ebp* mRNA level was observed

when CMLD-2 was co-administered with TGFβ-1 treatment in lieu of 5HT. No significant

1 difference between experimental and within-animal control ganglia were observed

2 (Figure 2c$_2$; Table 1P).

3 Blocking TGFβ signaling, p38 MAPK activation, and the ELAV-ARE interaction

4 during Trial 2 blocks the ApELAV-*apc/ebp* mRNA interaction

Our hypothesis, that ApELAV is recruited by Trial 2 and binds to the *apc/ebp*

transcript, predicts that the ApELAV-*apc/ebp* mRNA interaction should be increased at

the 1 hr time point following Trial 1 in ganglia which receive two-trial training. To directly

test this hypothesis, we performed a cross-linking immunoprecipitation experiment

followed by qPCR (CLIP-qPCR), to compare the <u>physical interaction</u> of *apc/ebp* mRNA

with ApELAV between experimental (two-trial training) and within-animal control groups

(Figure 3a$_1$). Following two-trial training, we indeed found a significant increase in

apc/ebp mRNA association with ApELAV (Figure 3a$_2$; Table 1Q), which strengthens the

hypothesis that ELAV, by binding to *apc/ebp* transcript, promotes its stabilization at that

timepoint.

If ApELAV is responsible for stabilizing *apc/ebp* mRNA during Trial 2, thereby

prolonging its expression to 1 hr following Trial 1, then blocking TGFβ signaling should

block the ApELAV-*apc/ebp* mRNA interaction. We have previously shown that treatment

with TGFβsRII disrupted the increase in Trial 2-dependent *apc/ebp* mRNA level (Kopec

et al., 2015). Treatment with TGFβsRII resulted in the complete loss of the ApELAV-

apc/ebp mRNA interaction at the same time point when assayed by CLIP-qPCR and

compared to contralateral within-animal control ganglia which received only vehicle

(0.1% BSA in ASW) (Figure 3a$_2$; Table 1R), demonstrating that TGFβ signaling is

required for the increase in ApELAV-*apc/ebp* mRNA interaction at that time point.

Figure 3

Figure 3. Cross-linking immunoprecipitation (CLIP)-qPCR reveals increase in ApELAV-*apc/ebp* interaction following two-trial training.

(a) (Top) Experimental paradigm. SN somata are UV-fixed and collected at 1 hr following the onset of Trial 1. Dissected ganglia are treated with CMLD-2, SB203580, or TGFβsRII from 15 min before Trial 2 until collection.
(Bottom) CLIP-qPCR experimental flowthrough. See Methods for more details.
(b) Following two-trial training, there is a significant increase in ApELAV-*apc/ebp* mRNA interaction which is blocked by treatment with TGFβ-sRII, SB203580, or CMLD-2. Each data point represents the average of 3 full experiments of SN clusters pooled from 8 animals. *$p<0.05$, **$p<0.01$; n=3-5

Activation of p38 MAPK is dependent on intact TGFβ signaling during Trial 2

(Figure 2a). Thus, we predicted that blocking p38 MAPK should block the ability of

ApELAV to interact with *apc/ebp* mRNA. We found that blocking p38 MAPK with

SB203580 blocked the ApELAV-*apc/ebp* mRNA interaction compared to contralateral

within-animal control ganglia which received only vehicle (0.1% DMSO) (Figure $3a_2$:

Table 1S). These results demonstrate that TGFβ signaling and p38 MAPK activation

during Trial 2 are required for the interaction of ApELAV and *apc/ebp* mRNA.

To directly demonstrate if CMLD-2 blocks the ApELAV-*apc/ebp* mRNA

interaction, we treated ganglia with CMLD-2 during Trial 2 and performed CLIP-qPCR.

We found that treatment with CMLD-2 during Trial 2 significantly disrupted the ApELAV-

apc/ebp mRNA interaction compared to contralateral within-animal control ganglia

which had received only vehicle (Figure $3a_2$; Table 1T). These results demonstrate the

effectiveness of CMLD-2 as an inhibitor of the ApELAV-*apc/ebp* mRNA interaction in

our two-trial paradigm.

CMLD-2 disrupts the induction of LTM by Trial 2

The molecular observations described thus far provide clear predictions for the

role of TGFβ-mediated post-transcriptional regulation of *apc/ebp* mRNA by the RNA-

binding protein ApELAV in the behavioral induction of LTM. We have previously shown

that TGFβ signaling during Trial 2 is required for (i) the expression of *apc/ebp* mRNA at

the 1 hr time point and (ii) the induction of LTM for sensitization (Kopec et al., 2015). In

the present paper, our data show that TGFβ signaling during Trial 2 provides

stabilization of *apc/ebp* mRNA by means of increasing its association with ApELAV.

Thus collectively, our molecular data predict that the ApELAV-*apc/ebp* mRNA

Figure 4

a₁   a₂

Figure 4. ApELAV-*apc/ebp* mRNA interaction is required for LTM.
(a₁) (Top) Experimental paradigm. LTM is assessed by stimulating the test site before training, then 15-22 hr after training and measuring the T-SWR. (Bottom) Semi-intact preparation. Two-trial behavioral training is administered to the training site, and drug is applied to the isolated CNS chamber.
(a₂) Blocking the ApELAV-*apc/ebp* mRNA interaction during Trial 2 significantly disrupts LTM formation. *$p<0.05$; n=8.

interaction during Trial 2 is required for the prolonged increase of *apc/ebp* mRNA that is

necessary for the induction of LTM. We directly tested this prediction in a final set of

behavioral experiments examining LTM for sensitization of the tail-elicited siphon

withdrawal reflex (T-SWR). We used the T-SWR semi-intact preparation (Kopec et al.,

2015; Sutton et al., 2001) that permits the manipulation of the molecular environment of

the CNS while directly assaying withdrawal responses (see Figure 4a$_1$, Methods).

To test the hypothesis that the ApELAV-*apc/ebp* mRNA interaction is required

during Trial 2 for LTM formation, we exposed the CNS to 100 µM CMLD-2 or an

equivalent volume of vehicle during Trial 2 (Figure 4a$_1$). In the presence of vehicle, a

within-group comparison revealed significant LTM for sensitization of the T-SWR. In

contrast, the induction of LTM was significantly disrupted when the ApELAV-*apc/ebp*

mRNA interaction was blocked during Trial 2. Moreover, a between-group comparison

revealed a significant difference between vehicle and CMLD-2 groups (Figure 4a$_2$;

Table 1U). These behavioral data confirm the predictions derived from our molecular

observations and support a general model in which TGFβ signaling in Trial 2 mediates

RNA stabilization that is essential for LTM formation.

Discussion

Our results reveal a novel molecular mechanism underlying LTM formation: the

stabilization of previously synthesized mRNA. Specifically, we show that *apc/ebp* mRNA

is post-transcriptionally regulated by the RNA-binding protein ELAV. This regulation is

dependent upon TGFβ signaling, which is engaged by the second trial in a two-trial

paradigm and is required for the induction of LTM (Kopec et al., 2015)*.* Our data

supports a model (Figure 5) in which Trial 1 gives rise to TrkB signaling, resulting in an

Figure 5

Figure 5. Working model of molecular mechanisms underlying *apc/ebp* transcript stabilization in two-trial LTM formation.
Trial 1 gives rise to an increase in CREB mediated transcription and an increase in *apc/ebp* gene expression 45 min later. When Trial 2 occurs at that timepoint (45 min), TGFβ signaling gives rise to an increase in p38 MAPK activation, resulting in an increase in ELAV-*apc/ebp* interaction. This interaction increases the stability of *apc/ebp* transcript and prolongs its expression to 1 hr, which is permissive for the induction of LTM.

increase in CREB-mediated transcription and an increase in *apc/ebp* gene expression

45 min later. During Trial 2, TGFβ-like ligands act in SNs to activate ELAV through p38

MAPK causing an increase in its binding to *apc/ebp* mRNA, thereby stabilizing the

transcript, resulting in a transcription-independent prolonged increase in *apc/ebp* mRNA

level. Finally, the ApELAV-mRNA interaction during Trial 2 is a pre-requisite for LTM

formation. These experiments provide novel evidence for a GF-mediated process in

LTM formation resulting from post-transcriptional regulation that modulates the stability

of an immediate early gene transcript, which serves as a transcription factor required in

LTM.

C/EBP in Long-Term Memory Formation

The C/EBP family of transcription factors (having 6 isoforms) are immediate early

genes downstream of CREB (Alberini, 2009). C/EBPs, particularly the C/EBPβ and

C/EBPδ isoforms, are regulated by learning-related stimuli across a wide range of

species, as well as across diverse brain regions and learning tasks (Alberini, 2009;

Alberini et al., 1994). Together with these studies, our findings indicate that C/EBP

signaling is a significant molecular step mediating plasticity and memory formation.

Interestingly, C/EBP contributes to the induction of GF ligands and receptors in (i) IGF-II

in hippocampus (Chen et al., 2011), (ii) VEGF-C and its receptor in lymphatic

endothelial cells (Min et al., 2011), (iii) TGFβr-II in human embryonic stem cells

(Takayama et al., 2014), and (iv) NGF in cerebral cortex (McCauslin et al., 2006). This

raises the intriguing possibility of a GF-mediated positive feedback loop during memory

formation in which GF signaling upregulates C/EBP, which in turn increases GF ligand

and/or receptor expression.

Here we have observed two phases of *apc/ebp* gene expression during LTM

formation with two-trial training: First, following Trial 1, a *transcription-dependent*

increase in *apc/ebp* mRNA level, and second, following Trial 2, a *transcription-*

independent stabilization via the RNA-binding protein ApELAV, which prolongs the

increase in *apc/ebp* mRNA level. This is a critical point of regulation since the presence

of *cis*-acting regulatory elements on specific mRNAs, such as AREs, can modulate, or

even counteract, the effect of increased transcription (Lee et al., 2015; Ross, 1995).

The AU-rich element RNA binding proteins ApAUF1 and ApELAV can bi-

directionally regulate *apc/ebp* mRNA stability: ApAUF1 binding to the 3'UTR of *apc/ebp*

induces the degradation of the transcript, and overexpression of ApAUF1 inhibits 5HT-

induced LTF in SN-MN co-culture (Lee et al., 2012). Conversely, ApELAV binds to the

same AU rich domain but stabilizes the transcript (Yim et al., 2006). Using the small-

molecule inhibitor CMLD-2 to block the ApELAV-*apc/ebp* mRNA interaction during the

second trial of two-trial training, we demonstrated both the loss of the increase in

apc/ebp mRNA level and the disruption in the induction of LTM, underscoring the

importance of this stabilizing mechanism for prolonging mRNA level and its critical role

in LTM formation. Thus, the expression of ApAUF1 and ApELAV, or the ratio of these

proteins, has the capacity to favor either the degradation or stabilization of *apc/ebp*

mRNAs.

Finally, we should emphasize that, in addition to transcript stabilization, other

mechanisms of post-transcriptional regulation have previously been implicated in LTM

formation in *Aplysia* (Rajasethupathy et al., 2009).

TGFβ signaling and RNA-binding proteins

1  Signaling through distinct GF families is specifically engaged, both spatially and

2 temporally, during LTM formation (Kopec et al., 2015). In the first trial of a two-trial

3 training paradigm, TrkB signaling at the SN synapse is required for *apc/ebp* gene

4 expression 45 min following the onset of the trial, and TGFβ signaling is required at the

5 SN soma during the second trial (at 45 min) for prolonged increase in *apc/ebp* mRNA

6 level. Here, we demonstrate that: (i) direct activation of TGFβ signaling (in the absence

7 of 5HT at 45 min) results in a transcription-independent increase in *apc/ebp* mRNA

8 level, and (ii) this effect is generated by recruiting ApELAV to bind and stabilize the

9 *apc/ebp* transcript produced by TrkB signaling-dependent transcriptional mechanisms

10 initiated by Trial 1.

11  Interestingly, there is considerable evidence that TGFβ-1 initiates stabilization of

12 mRNAs via an RNA binding complex (Amara et al., 1995). In cardiac fibroblasts, TGFβ-

13 1 treatment induced HuR (a member of the mammalian ELAV family) translocation from

14 the nucleus to the cytoplasm where it bound the ARE region of the 3' UTR in TGFβ-1

15 mRNA thus stabilizing the mRNA and increasing protein level(Bai et al., 2012). These

16 data suggest the intriguing possibility that, in addition to TGFβr-II signaling initiating a

17 cascade to prolong the expression of critical learning-related genes like *apc/ebp*, other

18 GF ligands could be induced and stabilized in the same way (including a TGFβ-1-like

19 ligand itself). Indeed, GF ligands (Lim and Alkon, 2012) as well as their receptors

20 (Balmer et al., 2001) can be regulated by ARE-RNA binding proteins.

21  The mammalian ELAV-like protein HuR is phosphorylated by PKC and p38

22 MAPK (Doller et al., 2007; Doller et al., 2008; Doller et al., 2011; Eberhardt et al., 2012;

23 Lafarga et al., 2009). In our two-trial paradigm, we found that p38 MAPK activity was

engaged through TGFβ, and its activity was required both for the ApELAV-*apc/ebp*

mRNA interaction and for the prolonged increase in *apc/ebp* mRNA. Indeed,

phosphorylation of ELAV in many systems has been shown to (i) increase its binding

affinity for AREs (Doller et al., 2011; Eberhardt et al., 2012; Pascale et al., 2005), (ii)

modulate the binding affinity of ELAV to specific types of AREs (Doller et al., 2010;

Eberhardt et al., 2012), and (iii) regulate its nucleo-cytoplasmic shuttling ability (Bai et

al., 2012; Doller et al., 2015; Doller et al., 2008; Fan and Steitz, 1998; Wang et al.,

2002). Interestingly, p38 MAPK activation has additionally been shown to inhibit the

activity of destabilizing ARE-binding proteins, which would act to shift the balance

towards mRNA stabilization (Briata et al., 2005).

In conclusion, the present results, taken with previous work, illustrate the

importance of transcript regulation through GF signaling as a critical site of molecular

interaction underlying long-term memory formation.

Table 1

	Corresponding Figure	Test	Statistics
A	1A	Within-group paired t-test	-ActinoD: n=8, 1.456 ± 0.140, p<0.05
B	1A	Within-group paired t-test	+ActinoD: n=7, 0.8505 ± 0.1601
C	1A	Between-group unpaired t-test	Control versus Actino: t=2.862, df=13, p<0.05
D	1B	Within-group paired t-test	1X 5HT -ActinoD: 0.9600 ± 0.1118, n=7, p=0.73; 2X 5HT -ActinoD: 1.765 ± 0.1463, n=11, p<0.01; 2X 5HT +ActinoD: 1.527 ± 0.1830, n=8, p<0.05
E	1B	ANOVA	F=6.786; p<0.01
F	1B	Between-group unpaired t-test	1X 5HT -ActinoD versus 2X 5HT -ActinoD: t=3.926, df=16, p<0.01; 1X 5HT -ActinoD versus 2X 5HT +ActinoD: t=2.550, df=13, p<0.05; 2X 5HT -ActinoD versus 2X 5HT +ActinoD: t=1.028, df=17, p=0.32
G	1C	Within-group paired t-test	1X 5HT +TGFβ: 1.384 ± 0.1493, n=8, p<0.05
H	1C	Between-group unpaired t-test	1X 5HT -ActinoD versus 1X 5HT +TGFβ: t=2.219, df=13, p<0.05)
I	1C	Within-group paired t-test; Between-group unpaired t-test	1X 5HT +TGFβ +Actino: 1.230 ± 0.0954, n=12, p<0.05; 1X 5HT +TGFβ versus 1X 5HT +TGFβ +Actino: t=0.9160, df=18, p=0.37
J	2A	Within-group paired t-test	2X 5HT: 1.845 ± 0.232, n=6, p<0.05
K	2A	Within-group paired t-test; Between-group unpaired t-test	1X 5HT: 1.089 ± 0.184, n=7, p=0.99; 2X 5HT +TGFβsRII: 1.148 ± 0.086, n=7, p=0.25; 1X 5HT versus 2X 5HT: t=2.582, df=11, p<0.05; 2X 5HT versus 2X 5HT +TGFβsRII: t=2.996, df=11, p<0.05
L	2B	ANOVA	F=7.804, p<0.01
M	2B	Within-group paired t-test; Between-group unpaired t-test	2X 5HT +DMSO: 186.9% ± 23.85%, n=23, p<0.01, 1 outlier; 2X 5HT +SB: 94.82% ± 15.74%, n=15, p=0.747, 2 outliers; 2X ASW +DMSO: 100% ± 12.74%, n=23, 1 outlier; 2X 5HT +DMSO versus 2X 5HT +SB: t=2.855, p<0.01; 2X 5HT +DMSO versus 2X ASW +DMSO: t=3.213, p<0.01
N	2C	Within-group paired t-test; Between-group unpaired t-test	2X 5HT +SB: 0.734 ± 0.161, n=8, p=0.142; 2X 5HT versus 2X 5HT +SB: t=4.692, df=17, p<0.01
O	2C	Within-group paired t-test; Between-group unpaired t-test	2X 5HT +CMLD-2: 1.194 ± 0.201, n=9, p=0.62; 2X 5HT versus 2X 5HT +CMLD-2: t=2.350, df=18, p<0.05)
P	2C	Within-group paired t-test	1X 5HT +TGFβ +CMLD-2: 0.9825 ± 0.1928, n=7, p=0.93, 1 outlier: see Methods
Q	3A	Within-group paired t-test	2X 5HT: 201.6% ± 19.89, n=5, p<0.01
R	3A	Within-group paired t-test; Between-group unpaired t-test	2X 5HT +TGFβsRII: 119.0% ± 10.31, n=3, p=0.21; 2X 5HT versus 2X 5HT +TGFβsRII: t=2.995, df=6, p<0.05
S	3A	Within-group paired t-test; Between-group unpaired t-test	2X 5HT +SB: 124.0% ± 18.35, n=3, p=0.32; 2X 5HT versus 2X 5HT +SB: t=2.611, df=6, p<0.05
T	3A	Within-group paired t-test; Between-group unpaired t-test	2X 5HT +CMLD-2: 104.4% ± 7.91, n=3, p=0.63; 2X 5HT versus 2X 5HT +CMLD-2: t=3.579, p<0.05
U	4A	Within-group paired t-test; Between-group unpaired t-test	Vehicle: 128.5% ± 7.294, n=8, p<0.05; CMLD-2: 109.3% ± 6.008, n=8, p=0.33, Vehicle versus CMLD-2: t=2.032, df=14, p<0.05

Table 1

Statistical table. Refer to Methods for information on statistics.

Methods

CONTACT FOR REAGENT AND RESOURCE SHARING

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Thomas J. Carew (tcarew@nyu.edu).

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Aplysia californica were acquired from South Coast Bio-Marine and University of Miami National Resource for *Aplysia.* All animals were allowed to acclimate for at least 4 days in circulating tanks with artificial sea water (ASW) (Instant Ocean) at 15° C.

METHOD DETAILS

Ganglion Preparation

Pleural-pedal ganglia were dissected from anesthetized *Aplysia californica* (150-250 g). In a 1:1 solution of $MgCl_2$ and ASW, the pleural SN cluster and SN-MN neuropil were exposed. Ganglia were perfused with ASW for at least 1.5 hr to clear $MgCl_2$ prior to experimentation. Experimental and contralateral within-animal control ganglia both received GF/drug/vehicle treatment, while only the experimental ganglia received two-trial 5HT training.

Training and Drug Incubation for Gene Expression Analyses

After $MgCl_2$ washout, two-trial analog training was delivered by two pulses of 5HT (50 μM, Sigma, #H9523-25MG) in ASW (ITI = 45 min). As previously described (Kopec et al., 2015), ganglia in GF treatment experiments were blocked with BSA for at least 5 min pre-GF treatment, and GF/drug/vehicle was applied at the appropriate experimental time point. Drugs used were as follows: rhTGFβ-1 (R&D Systems, #240-B-002), rhTGFβ-sRII (R&D Systems, #241-R2-025), SB203580 (Tocris, #1202), Actinomycin D

(Sigma, #A1410-10MG), CMLD-2 (Millipore, #538339). Drug incubation was from 15

min before the training trial, during the training trial, and proceeded until SN cluster

collection. Within-animal control ganglia concurrently received appropriate vehicle but

did not receive two-trial training. Control and experimental SN clusters were collected at

the experimental time point and prepared for either Western Blotting or quantitative

PCR analysis.

ApELAV1 Protein Expression

The coding region for ApELAV1 was isolated by PCR from a previously

published plasmid kindly provided by the Bong-Kiun Kaang (Seoul, South Korea)(Yim et

al., 2006) and ligated into pGEX vector to make pGEX-ApELAV1-GST. Rosetta 2 DE3

cells (Millipore) were transformed with this plasmid, and cells were induced for 2 h at

37°C with 1 mM IPTG during log-phase growth. Cells were subsequently lysed and

subjected to Western blotting.

Western Blotting

Samples were lysed and snap-frozen in RIPA buffer (Sigma) with 1X Halt

protease and phosphatase inhibitor cocktail (Invitrogen), loaded onto 4-12% Bis-Tris

gels (Novex; Life Technologies) for electrophoresis and then transferred to

nitrocellulose membranes for incubation with primary antibodies and secondary

antibodies: anti-HuR (Santa Cruz, sc-5261), anti-β-Actin (Cell Signaling Technology,

#3700), anti-p-p38 MAPK (Cell Signaling Technology, #4511), anti-β-Tubulin (Cell

Signaling Technology, #2146), anti-mouse IgG IRDye 680LT (LICOR Biosciences,

#926-68020), anti-rabbit IgG IRDye 800CW (LICOR Biosciences, #926-32211). For

ApELAV and p-p38 MAPK experiments, protein levels within each sample was

assessed using the LICOR imaging system. Each protein of interest band was

normalized to the tubulin or actin within the same sample, and normalized protein of

interest in the experimental ganglia was compared to within-animal control ganglia. Data

are displayed as fold-expression of normalized protein of interest in experimental

sample relative to control sample.

RNA isolation, cDNA synthesis, quantitative PCR analysis

RNA was isolated and purified using Ambion RNAqueous Micro kits (Invitrogen),

and total RNA for cDNA synthesis was normalized between experimental and control

ganglia (~100ng RNA). cDNA was synthesized using Superscript IV cDNA synthesis

reagents (Invitrogen). Quantitative PCR was performed using a Roche LightCycler 480,

with 1 µL cDNA, 1 µL each of 5µM F' and R' primers, and 18 µL 1X SybrGreen (Roche).

Primer sequences are as follows: apc/ebp-F: caccacctcactcccatctc; apc/ebp-R:

ctgacgtctgcgagactttg; apgapdh-F: ctctgagggtgctttgaagg; apgapdh-R:

gttgtcgttgagggcaattc. The amplification program was 95°C for 3 mins, 30 cycles of 95°C

for 10 s, 60°C for 20 s, and 72°C for 30 s, and lastly a melting curve to confirm a single

PCR product. Data were analyzed by normalizing the quantity of each gene of interest

(measured by Ct) to *apgapdh* within the same sample (ΔCt), then comparing normalized

values between experimental and within-animal control groups ($\Delta\Delta$Ct). Data are

displayed as fold induction relative to the control group.

Single molecule fluorescence *in situ* hybridization (smFISH)

Aplysia SNs were isolated from the ventral SN cluster of the pleural ganglion of

80g animals according to (Zhao et al., 2009) in 50mm Mattek plates. After 4 days *in*

vitro, 7 plates of 10-20 SNs were chosen and separated into 4 different groups

randomly. 2 plates each were designated as "2X 5HT +DMSO", "2X 5HT +SB", and "2X

ASW +DMSO". 1 plate received no treatments but was used as a "no probe" control to

determine background fluorescence of SNs without DNA probes.

The smFISH protocol was adapted from (Eliscovich et al., 2017). Following

treatment according to the experimental paradigm (5HT treatment was 50 µM in ASW,

ASW treatment was an equivalent volume of ASW), the cells were fixed for 30 min at

room temperature in 4% formaldehyde in 30% sucrose / DEPC-treated PBS with

supplemented 1 mM magnesium chloride and 0.1 mM calcium chloride (DEPC-PBS-

MC). Cells were washed quickly 2X with DEPC-PBS-MC, and quenched with glycine in

PBS-MC. Quench buffer was removed and cells were incubated in permeabilization-

block buffer (0.2% Triton X-100, 0.5% Ultrapure BSA in DEPC-PBS-MC) for 30 min at

room temperature while shaking. Permabilization-block buffer was removed and cells

were incubated with pre-hybridization buffer (10% deionized formamide, 2X SSC, 0.5%

Ultrapure BSA in nuclease-free water) for 30 min at room temperature without shaking.

Pre-hybridization buffer was removed and cells were incubated with hybridization buffer

(10% deionized formamide, 2X SSC, 1 mg/mL *Escherichia coli* tRNA, 20 mg/mL BSA,

2X SSC, 2 mM Vanadyl Ribonucleoside Complex, 10% dextrane sulfate, 125 nM of

DNA probes in nuclease-free water) for 4 h at 37 °C. Thirty-three (33) DNA probes

conjugated to Quasar 670 dye (LGC Biosearch Technology) were designed to be

antisense to all regions on *apc/ebp* mRNA without significant complementarity to other

mRNAs in the *Aplysia* Refseq database (NCBI) and are listed in Supplementary Table

1. Cells were washed four times with 2X SSC and then once quickly with nuclease-free

water. Coverslips were detached from the plates and mounted using Prolong Gold

Antifade Reagent with DAPI (Life Technologies).

A Deltavision Elite imaging system (GE) equipped with Olympus ×60 / 1.42 NA

oil immersion lens and an InsightSSI Solid State Illumination module was used to image

samples. Individual full SN somata were imaged in a z-stack with 500 nm spacing. Raw

images were loaded into FISH-QUANT software (Mueller et al., 2013) and batch

processed after detecting nucleus size (by DAPI) and tracing the outline of each SN.

Initial analyses determined that, in the cells which received sham treatment (2X ASW),

the number of puncta counted in each cell was correlated strongly to the area of the cell

nucleus (Supplementary Figure 3), which corresponds to previous literature noting that

the absolute amount of RNA can be dependent on cell size (Marguerat and Bähler,

2012). Therefore, the number of puncta counted in each cell was normalized to the area

of the nucleus in each cell, as this value was calculated automatically by FISH-QUANT.

Cross-Linking Immunoprecipitation - qPCR

The protocol was adapted from (Yoon and Gorospe, 2016). Experimental and

contralateral within-animal control ganglia both received GF chimera/drug/vehicle

treatment, while only the experimental ganglia received two-trial 5HT training. At the

appropriate experimental timepoint, experimental and control ganglia were exposed

twice to 400 mJ UV (254nm) to crosslink ribonucleoprotein complexes. SN clusters of 8

animals were collected and lysed in ice-cold NP-40 lysis buffer (20 mM Tris-HCl pH 7.5,

100 mM KCl, 5 mM $MgCl_2$, 0.5% NP-40, 1 mM DTT) supplemented with 1X Halt

protease and phosphatase inhibitor cocktail and 40 U/mL RNaseOUT (Invitrogen), and

pooled for each experimental and contralateral control sample. Lysates were split into

two fractions each and immunoprecipitated with either antibodies against HuR (Santa-

Cruz, sc-5261) or normal mouse IgG_1 (Santa Cruz, sc-3877) using Protein G

Dynabeads (Invitrogen). Beads were washed with ice-cold NT-2 buffer (50 mM Tris-HCl

pH 7.5, 150 mM NaCl, 1 mM $MgCl_2$, 0.05% NP-40). Following consecutive digestions

with DNase I and Proteinase K (New England Biolabs), RNA was isolated with Trizol

extraction (Invitrogen) and cDNA was synthesized using Superscript IV cDNA synthesis

reagents (Invitrogen). Quantitative PCR was performed as described above.

Importantly, primers were designed to lie within the 3' UTR of *apc/ebp* mRNA, the

binding region of ELAV. The primer sequences were as follows: apc/ebpUTR-F:

cccaagcatgttgtgatagttgt; apc/ebpUTR-R: tgggaggagatagaagcagtg. Data were analyzed

by normalizing the difference in *apc/ebp* between ELAV- and IgG- immunoprecipitated

samples to the difference in *apgapdh* between the same samples, then comparing

these normalized values between experimental and within-animal control groups.

Semi-intact Behavioral Preparation

Preparations were completed as previously described (Kopec et al., 2015; Sutton

et al., 2001). Briefly, the ring ganglia (cerebral and paired pleural-pedal ganglia) from

anesthetized Aplysia (250-400 g; South Coast Bio-Marine) were surgically isolated,

leaving p9 and pleural-abdominal innervation to the tail and abdominal ganglia,

respectively, intact. The tail and mantle were surgically removed, and the siphon artery

was cannulated with Dow Corning silastic tubing (0.025 in I.D.; Fisher Scientific) and

perfused at ~5 ml/min, while the tail was perfused at ~0.5 ml/min via three 22-gauge

needles inserted into the tail. The tail and mantle were pinned to the chamber floor,

while the ring ganglia with both pleural-pedal ganglia de-sheathed were pinned in an

isolated CNS chamber. Both the chamber containing the peripheral tissues as well as

the CNS chamber were continuously perfused with seawater (Instant Ocean, 15°C).

The p9 and pleural-abdominal nerves exited the CNS chamber through small slits that

were sealed with Vaseline. Preparations were allowed at least 2 hrs to recover prior to

baseline measurements.

Training and Drug Incubation for LTM Analyses

An average of 3-4 baseline siphon-withdrawal measurements were recorded by

stimulating the medial posterior tip of the tail with a water jet (0.4 s, 45 psi; ITI 15 mins;

Teledyne Water Pik) and measuring the duration of time with which the siphon

withdrew. No significant differences in the pre-training baseline duration of T-SWRs

were observed in any experimental condition. The CNS was incubated with

Drug/Vehicle 15 mins prior to Trial 2, which was washed out 30 mins after Trial 2. Two-

Trial LTM training (100 mA, 1.5 s, ITI 45 mins) was delivered medially to the anterior

portion of the tail through a hand-held electrode. LTM for sensitization of the T-SWR

was assessed by 3-4 tests (ITI=15 mins) 15-22 hrs after training by an experimenter

blind to the experimental condition (Drug vs. Vehicle). The data are displayed as

duration of siphon withdrawal as a percent of pre-training baseline.

QUANTIFICATION AND STATISTICAL ANALYSIS

Data were analyzed with parametric statistics using GraphPad Prism. Statistical

details for all experiments can be found within the main text and/or in the figure legends.

Within-group analyses were performed using paired t-test. LTM for sensitization of the

T-SWR is compared relative to pre-training baseline T-SWR within the same

preparation. Between-group analyses (Drug vs. Vehicle) were performed using t-tests.

When appropriate, ANOVA analyses were used to determine if there was a difference among the groups and are indicated in the main text. Significant results were followed by planned parametric comparisons. Outliers greater than 2 standard deviations from the mean were excluded resulting in the removal of 5 data points (<5% of all data points; reported in Table 1). Due to normal distribution of data, data in all figures are depicted as means ± SEM. Significant within-group comparisons are displayed as asterisks above the summary data, and significant between-group comparisons are displayed as asterisks above a bar drawn between the summary data. Asterisks indicate statistical significance (*p<0.05, **p<0.01).

AUTHOR CONTRIBUTIONS

AAM conceived of, designed, performed, and analyzed experiments. AMK conceived of, designed, performed, and analyzed experiments. TJC conceived of and designed experiments. All authors wrote the paper.

ACKNOWLEDGEMENTS

We thank A. Alexandrescu, P.E. Miranda, and N.V. Kukushkin for their comments on the manuscript. We thank Bong-Kiun Kaang for sharing ApELAV plasmids. We thank the Hochwagen Laboratory for use of the Deltavision Elite microscope for smFISH imaging and the NYU Center for Genomics and Systems Biology Genomics Core for use of the Roche 480 LightCycler for qPCR experiments. Figures were created with BioRender.com. This work was supported by NIMH R01 MH 094792 to TJC and a Hellenic Medical Society of New York Leonidas Lantzounis Research Grant to AAM.

DECLARATION OF INTERESTS

The authors declare no competing interests.

REFERENCES

Alberini, C.M. (2009). Transcription factors in long-term memory and synaptic plasticity. Physiol Rev *89*, 121-145.

Alberini, C.M., Ghirardi, M., Metz, R., and Kandel, E.R. (1994). C/EBP is an immediate-early gene required for the consolidation of long-term facilitation in Aplysia. Cell *76*, 1099-1114.

Amadio, M., Scapagnini, G., Davinelli, S., Calabrese, V., Govoni, S., and Pascale, A. (2014). Involvement of ELAV RNA-binding proteins in the post-transcriptional regulation of HO-1. Front Cell Neurosci *8*, 459.

Amara, F.M., Chen, F.Y., and Wright, J.A. (1995). Defining a novel cis element in the 3'-untranslated region of mammalian ribonucleotide reductase component R2 mRNA: role in transforming growth factor-beta 1 induced mRNA stabilization. Nucleic Acids Res *23*, 1461-1467.

Bai, D., Gao, Q., Li, C., Ge, L., Gao, Y., and Wang, H. (2012). A conserved TGFbeta1/HuR feedback circuit regulates the fibrogenic response in fibroblasts. Cell Signal *24*, 1426-1432.

Balmer, L.A., Beveridge, D.J., Jazayeri, J.A., Thomson, A.M., Walker, C.E., and Leedman, P.J. (2001). Identification of a novel AU-Rich element in the 3' untranslated region of epidermal growth factor receptor mRNA that is the target for regulated RNA-binding proteins. Mol Cell Biol *21*, 2070-2084.

Bolognani, F., Merhege, M.A., Twiss, J., and Perrone-Bizzozero, N.I. (2004). Dendritic localization of the RNA-binding protein HuD in hippocampal neurons: association with polysomes and upregulation during contextual learning. Neurosci Lett *371*, 152-157.

Brennan, C.M., and Steitz, J.A. (2001). HuR and mRNA stability. Cell Mol Life Sci *58*, 266-277.

Briata, P., Forcales, S.V., Ponassi, M., Corte, G., Chen, C.Y., Karin, M., Puri, P.L., and Gherzi, R. (2005). p38-dependent phosphorylation of the mRNA decay-promoting factor KSRP controls the stability of select myogenic transcripts. Mol Cell *20*, 891-903.

Brunelli, M., Castellucci, V., and Kandel, E.R. (1976). Synaptic facilitation and behavioral sensitization in Aplysia: possible role of serotonin and cyclic AMP. Science *194*, 1178-1181.

Chen, D.Y., Stern, S.A., Garcia-Osta, A., Saunier-Rebori, B., Pollonini, G., Bambah-Mukku, D., Blitzer, R.D., and Alberini, C.M. (2011). A critical role for IGF-II in memory consolidation and enhancement. Nature *469*, 491-497.

Doller, A., Badawi, A., Schmid, T., Brauss, T., Pleli, T., zu Heringdorf, D.M., Piiper, A., Pfeilschifter, J., and Eberhardt, W. (2015). The cytoskeletal inhibitors latrunculin A and blebbistatin exert antitumorigenic properties in human hepatocellular carcinoma cells by interfering with intracellular HuR trafficking. Exp Cell Res *330*, 66-80.

Doller, A., Huwiler, A., Muller, R., Radeke, H.H., Pfeilschifter, J., and Eberhardt, W. (2007). Protein kinase C alpha-dependent phosphorylation of the mRNA-stabilizing factor HuR: implications for posttranscriptional regulation of cyclooxygenase-2. Mol Biol Cell *18*, 2137-2148.

Doller, A., Pfeilschifter, J., and Eberhardt, W. (2008). Signalling pathways regulating nucleo-cytoplasmic shuttling of the mRNA-binding protein HuR. Cell Signal *20*, 2165-2173.

Doller, A., Schlepckow, K., Schwalbe, H., Pfeilschifter, J., and Eberhardt, W. (2010).
Tandem phosphorylation of serines 221 and 318 by protein kinase Cdelta coordinates
mRNA binding and nucleocytoplasmic shuttling of HuR. Mol Cell Biol *30*, 1397-1410.
Doller, A., Winkler, C., Azrilian, I., Schulz, S., Hartmann, S., Pfeilschifter, J., and
Eberhardt, W. (2011). High-constitutive HuR phosphorylation at Ser 318 by PKC{delta}
propagates tumor relevant functions in colon carcinoma cells. Carcinogenesis *32*, 676-
685.
Eberhardt, W., Doller, A., and Pfeilschifter, J. (2012). Regulation of the mRNA-binding
protein HuR by posttranslational modification: spotlight on phosphorylation. Curr Protein
Pept Sci *13*, 380-390.
Eliscovich, C., Shenoy, S.M., and Singer, R.H. (2017). Imaging mRNA and protein
interactions within neurons. Proc Natl Acad Sci U S A *114*, E1875-E1884.
Fan, X.C., and Steitz, J.A. (1998). Overexpression of HuR, a nuclear-cytoplasmic
shuttling protein, increases the in vivo stability of ARE-containing mRNAs. EMBO J *17*,
3448-3460.
Ge, J., Chang, N., Zhao, Z., Tian, L., Duan, X., Yang, L., and Li, L. (2016). Essential
Roles of RNA-binding Protein HuR in Activation of Hepatic Stellate Cells Induced by
Transforming Growth Factor-beta1. Sci Rep *6*, 22141.
Kandel, E.R. (2012). The molecular biology of memory: cAMP, PKA, CRE, CREB-1,
CREB-2, and CPEB. Mol Brain *5*, 14.
Kopec, A.M., and Carew, T.J. (2013). Growth factor signaling and memory formation:
temporal and spatial integration of a molecular network. Learn Mem *20*, 531-539.
Kopec, A.M., Philips, G.T., and Carew, T.J. (2015). Distinct Growth Factor Families Are
Recruited in Unique Spatiotemporal Domains during Long-Term Memory Formation in
Aplysia californica. Neuron *86*, 1228-1239.
Lafarga, V., Cuadrado, A., Lopez de Silanes, I., Bengoechea, R., Fernandez-Capetillo,
O., and Nebreda, A.R. (2009). p38 Mitogen-activated protein kinase- and HuR-
dependent stabilization of p21(Cip1) mRNA mediates the G(1)/S checkpoint. Mol Cell
Biol *29*, 4341-4351.
Lee, J.A., Lee, S.H., Lee, C., Chang, D.J., Lee, Y., Kim, H., Cheang, Y.H., Ko, H.G.,
Lee, Y.S., Jun, H.*, et al.* (2006). PKA-activated ApAF-ApC/EBP heterodimer is a key
downstream effector of ApCREB and is necessary and sufficient for the consolidation of
long-term facilitation. J Cell Biol *174*, 827-838.
Lee, Y.S., Choi, S.L., Jun, H., Yim, S.J., Lee, J.A., Kim, H.F., Lee, S.H., Shim, J., Lee,
K., Jang, D.J.*, et al.* (2012). AU-rich element-binding protein negatively regulates
CCAAT enhancer-binding protein mRNA stability during long-term synaptic plasticity in
Aplysia. Proc Natl Acad Sci U S A *109*, 15520-15525.
Lee, Y.S., Lee, J.A., and Kaang, B.K. (2015). Regulation of mRNA stability by ARE-
binding proteins in synaptic plasticity and memory. Neurobiol Learn Mem *124*, 28-33.
Lim, C.S., and Alkon, D.L. (2012). Protein kinase C stimulates HuD-mediated mRNA
stability and protein expression of neurotrophic factors and enhances dendritic
maturation of hippocampal neurons in culture. Hippocampus *22*, 2303-2319.
Lu, Y., Ji, Y., Ganesan, S., Schloesser, R., Martinowich, K., Sun, M., Mei, F., Chao,
M.V., and Lu, B. (2011). TrkB as a potential synaptic and behavioral tag. J Neurosci *31*,
11762-11771.

Marguerat, S., and Bähler, J. (2012). Coordinating genome expression with cell size.
Trends Genet *28*, 560-565.
Marinesco, S., and Carew, T.J. (2002). Serotonin release evoked by tail nerve
stimulation in the CNS of aplysia: characterization and relationship to heterosynaptic
plasticity. J Neurosci *22*, 2299-2312.
McCauslin, C.S., Heath, V., Colangelo, A.M., Malik, R., Lee, S., Mallei, A., Mocchetti, I.,
and Johnson, P.F. (2006). CAAT/enhancer-binding protein delta and cAMP-response
element-binding protein mediate inducible expression of the nerve growth factor gene in
the central nervous system. J Biol Chem *281*, 17681-17688.
Min, Y., Ghose, S., Boelte, K., Li, J., Yang, L., and Lin, P.C. (2011). C/EBP-delta
regulates VEGF-C autocrine signaling in lymphangiogenesis and metastasis of lung
cancer through HIF-1alpha. Oncogene *30*, 4901-4909.
Mirisis, A.A., Alexandrescu, A., Carew, T.J., and Kopec, A.M. (2016). The Contribution
of Spatial and Temporal Molecular Networks in the Induction of Long-term Memory and
Its Underlying Synaptic Plasticity. AIMS Neurosci *3*, 356-384.
Mueller, F., Senecal, A., Tantale, K., Marie-Nelly, H., Ly, N., Collin, O., Basyuk, E.,
Bertrand, E., Darzacq, X., and Zimmer, C. (2013). FISH-quant: automatic counting of
transcripts in 3D FISH images. Nat Methods *10*, 277-278.
Muralidharan, R., Mehta, M., Ahmed, R., Roy, S., Xu, L., Aube, J., Chen, A., Zhao, Y.D.,
Herman, T., Ramesh, R*., et al.* (2017). HuR-targeted small molecule inhibitor exhibits
cytotoxicity towards human lung cancer cells. Sci Rep *7*, 9694.
Pascale, A., Amadio, M., Scapagnini, G., Lanni, C., Racchi, M., Provenzani, A., Govoni,
S., Alkon, D.L., and Quattrone, A. (2005). Neuronal ELAV proteins enhance mRNA
stability by a PKCalpha-dependent pathway. Proc Natl Acad Sci U S A *102*, 12065-
12070.
Philips, G.T., Ye, X., Kopec, A.M., and Carew, T.J. (2013). MAPK establishes a
molecular context that defines effective training patterns for long-term memory
formation. J Neurosci *33*, 7565-7573.
Quattrone, A., Pascale, A., Nogues, X., Zhao, W., Gusev, P., Pacini, A., and Alkon, D.L.
(2001). Posttranscriptional regulation of gene expression in learning by the neuronal
ELAV-like mRNA-stabilizing proteins. Proc Natl Acad Sci U S A *98*, 11668-11673.
Rajasethupathy, P., Fiumara, F., Sheridan, R., Betel, D., Puthanveettil, S.V., Russo,
J.J., Sander, C., Tuschl, T., and Kandel, E. (2009). Characterization of small RNAs in
Aplysia reveals a role for miR-124 in constraining synaptic plasticity through CREB.
Neuron *63*, 803-817.
Ross, J. (1995). mRNA stability in mammalian cells. Microbiol Rev *59*, 423-450.
Shobe, J., Philips, G.T., and Carew, T.J. (2016). Transforming growth factor beta
recruits persistent MAPK signaling to regulate long-term memory consolidation in
Aplysia californica. Learn Mem *23*, 182-188.
Slone, S., Anthony, S.R., Wu, X., Benoit, J.B., Aube, J., Xu, L., and Tranter, M. (2016).
Activation of HuR downstream of p38 MAPK promotes cardiomyocyte hypertrophy. Cell
Signal *28*, 1735-1741.
Sutton, M.A., Masters, S.E., Bagnall, M.W., and Carew, T.J. (2001). Molecular
mechanisms underlying a unique intermediate phase of memory in aplysia. Neuron *31*,
143-154.

Takayama, K., Kawabata, K., Nagamoto, Y., Inamura, M., Ohashi, K., Okuno, H.,
Yamaguchi, T., Tashiro, K., Sakurai, F., Hayakawa, T.*, et al.* (2014). CCAAT/enhancer
binding protein-mediated regulation of TGFbeta receptor 2 expression determines the
hepatoblast fate decision. Development *141*, 91-100.
Tsukada, J., Yoshida, Y., Kominato, Y., and Auron, P.E. (2011). The CCAAT/enhancer
(C/EBP) family of basic-leucine zipper (bZIP) transcription factors is a multifaceted
highly-regulated system for gene regulation. Cytokine *54*, 6-19.
Wang, W., Fan, J., Yang, X., Furer-Galban, S., Lopez de Silanes, I., von Kobbe, C.,
Guo, J., Georas, S.N., Foufelle, F., Hardie, D.G.*, et al.* (2002). AMP-activated kinase
regulates cytoplasmic HuR. Mol Cell Biol *22*, 3425-3436.
Wang, W., Furneaux, H., Cheng, H., Caldwell, M.C., Hutter, D., Liu, Y., Holbrook, N.,
and Gorospe, M. (2000). HuR regulates p21 mRNA stabilization by UV light. Mol Cell
Biol *20*, 760-769.
Wu, X., Lan, L., Wilson, D.M., Marquez, R.T., Tsao, W.C., Gao, P., Roy, A., Turner,
B.A., McDonald, P., Tunge, J.A.*, et al.* (2015). Identification and validation of novel
small molecule disruptors of HuR-mRNA interaction. ACS Chem Biol *10*, 1476-1484.
Yamashita, M., Fatyol, K., Jin, C., Wang, X., Liu, Z., and Zhang, Y.E. (2008). TRAF6
mediates Smad-independent activation of JNK and p38 by TGF-beta. Mol Cell *31*, 918-
924.
Yim, S.J., Lee, Y.S., Lee, J.A., Chang, D.J., Han, J.H., Kim, H., Park, H., Jun, H., Kim,
V.N., and Kaang, B.K. (2006). Regulation of ApC/EBP mRNA by the Aplysia AU-rich
element-binding protein, ApELAV, and its effects on 5-hydroxytryptamine-induced long-
term facilitation. J Neurochem *98*, 420-429.
Yoon, J.H., and Gorospe, M. (2016). Cross-Linking Immunoprecipitation and qPCR
(CLIP-qPCR) Analysis to Map Interactions Between Long Noncoding RNAs and RNA-
Binding Proteins. Methods Mol Biol *1402*, 11-17.
Zhang, F., Endo, S., Cleary, L.J., Eskin, A., and Byrne, J.H. (1997). Role of transforming
growth factor-beta in long-term synaptic facilitation in Aplysia. Science *275*, 1318-1320.
Zhang, Y., Smolen, P., Baxter, D.A., and Byrne, J.H. (2017). Biphasic Regulation of p38
MAPK by Serotonin Contributes to the Efficacy of Stimulus Protocols That Induce Long-
Term Synaptic Facilitation. eNeuro *4*.
Zhao, Y., Wang, D.O., and Martin, K.C. (2009). Preparation of Aplysia sensory-motor
neuronal cell cultures. J Vis Exp.
34
35

Supplementary Figure 1

a

b₁

b₂

Supplementary Figure 1.
(a) (Left) ApELAV1 and ApELAV2 are recognized by the antibody at their respective predicted molecular weights. ApELAV1 is ubiquitously expressed and ApELAV2 is neuron-specific. **(Right)** The antibody reacts strongly with recombinant ApELAV1 (expression induced with IPTG).
(b₁) Experimental paradigm. Experimental ganglia receive two-trial training with 5HT and contralateral control ganglia receive two-trial training without 5HT. Ganglia are collected and lysed at 1 hr.
(b₂) There is no significant change in ApELAV protein levels following two-trial training. Representative western blots are shown below the histograms (n=6).

Supplementary Figure 2

a₁

a₂

Minutes Following Onset of Treatment

Supplementary Figure 2.
(a₁) Experimental paradigm. Experimental ganglia are treated with actinomycin D and CMLD-2, or actinomycin D alone, and control ganglia are treated with vehicle. SN clusters are collected and lysed at 20 min or 40 min following treatment onset.
(a₂) CMLD-2 treatment results in significantly lower *apc/ebp* levels at 40 min post-treatment onset in experimental ganglia compared to actinomycin D alone.

1
2

Supplementary Figure 3

Supplementary Figure 3.
Distribution of smFISH data from 2X ASW +DMSO (control) group. There is a strong correlation between cell size (measured by area of nucleus) and number of puncta observed.

Supplementary Table 1

Probe Sequence

cggaaagctgaagtgaatcc
tgtgctgaagaaagtctgca
tagaagtcagctcacctttg
gcacaatcaattgactccac
gggatcgagaagatctaggg
agtctaaaactcgctccagg
ctgtttgtgggaatgtattc
aaataatgtggctgtccagg
tgggatctggggaaatgtga
aggctctgatgttgtagatc
tatttctcctcgtgcacgag
ctccaaaaagttgaccaccg
ctgtagtagctcgtcacgtc
atttgttgcgttgctgagtt
gagaggtcaactggctgaac
ttgaaaaccactggtgctgg
cacgtttcactaagtttggc
atctgtcctgactccgaaat
tgtggcatggagtaaatcca
gagtgtgaggatgtttagcg
ctatcgatgagggcacagat
gatgttggagatgggagtga
ccaggactagaagctgagta
tgcgagactttgtggagtag
cttatccacgtactcctgag
gcttttgcgaactgcaacat
ttcacgctgcttttcctttg
aagagacttcacacgaccct
gcaactggtcattttgatcg
actgagcaagtctactttct
gtcctttgagaacattcaat
gaagcaccaacgttcaagaa
cttgaggaattcatcaggta

Supplementary Table 1
List of DNA probes custom designed by LGC Biosearch Technology for detection of *apc/ebp* mRNA.