A neural signature of choice under sensory conflict in *Drosophila*

Preeti Sareen[1], Li Yan McCurdy[1,2], Michael N. Nitabach[1,3,4,*]

[1]Department of Cellular and Molecular Physiology, Yale University, New Haven, CT, USA

[2]Interdepartmental Neuroscience Program, Yale University, New Haven, CT, USA

[3]Department of Genetics, Yale University, New Haven, CT, USA

[4]Department of Neuroscience, Yale University, New Haven, CT, USA

* Lead contact: michael.nitabach@yale.edu

Summary

Feeding decisions are fundamental to survival, and decision making is often disrupted in disease, yet the neuronal and molecular mechanisms of adaptive decision making are not well understood. Here we show that neural activity in a small population of neurons projecting to the fan-shaped body higher-order central brain region of *Drosophila* represents final food choice during sensory conflict. We found that hungry flies made tradeoffs between appetitive and aversive values of food in a decision making task to choose bittersweet food with high sucrose concentration, but adulterated with bitter quinine, over sweet-only food with less sucrose. Using cell-specific optogenetics and receptor RNAi knockdown during the decision task, we identified an upstream neuropeptidergic and dopaminergic network that relays internal state and other decision-relevant information, such as valence and previous experience, to a specific subset of fan-shaped body neurons. Importantly, calcium imaging revealed that these neurons were strongly inhibited by the taste of the rejected food choice, suggesting that they encode final behavioral food choice. Our findings reveal that fan-shaped body taste responses to food choices are determined not only by taste quality, but also by previous experience (including choice outcome) and hunger state, which are integrated in the fan-shaped body to encode the decision before relay to downstream motor circuits for behavioral implementation. Our results uncover a novel role for the fan-shaped body in choice encoding, and reveal a neural substrate for sensory and internal state integration for decision making in a genetically tractable model organism to enable mechanistic dissection at circuit, cellular, and molecular levels.

Keywords: decision making, conflicting sensory context, value integration, internal state, central complex, fan-shaped body

Introduction

Animals integrate food-related sensory information from their external environment with their internal state in order to make adaptive decisions. Often food-related sensory information is conflicting in valence. For example, *Drosophila* flies forage on decomposing fruits and must balance obtaining essential nutrition with avoiding toxins, pathogens, etc. As flies forage, sweet and bitter taste receptors on their legs and wings signal the presence of sweet nutritive food and bitter potential toxins (Scott, 2018). Flies must adaptively weigh and integrate this conflicting information before consumption to enhance survival fitness. While some studies have shown integration of sweet and bitter tastes at the sensory neuron level (Chu et al., 2014; Jeong et al., 2013; Meunier et al., 2003), very little is known about how taste and other external sensory cues are integrated with internal state of the animal to form a feeding decision in the higher brain. We investigated how value-based decisions are made in the central brain of a hungry fly, using an experimental paradigm in which freely foraging flies sample and choose between different sweet and bittersweet foods (Fig. 1A). We quantified food choice and manipulated subsets of neurons while flies engaged in this decision task with conflicting taste information (Fig. 1A).

Results

Hungry flies make tradeoffs when faced with conflicting sensory information

We quantified the choices of wild-type flies deprived of food for different durations between a range of increasing concentration of sweet (sucrose-only) food option and a constant bittersweet (sucrose + quinine) option, with each option including either a red or blue food dye. When choosing between a low sucrose concentration sweet option and a high sucrose concentration bittersweet option, flies prefer higher sucrose bittersweet (Fig. 1B), as quantified by color of food ingested. As sucrose concentration of the sweet choice increased, flies

increasingly preferred it over bittersweet. This dose-dependent change in preference suggests

that when the sweet option reaches sufficient sucrose, the caloric advantage in choosing a less

palatable bittersweet food is outweighed by the danger-avoidance advantage of the sweet option

(Fig. 1B). In the absence of bitter, flies always chose the sweeter option (Fig. S1A), suggesting

that there was no saturation of sweet taste sensation for behavioral discrimination even at the

highest sucrose concentration tested (500 mM sucrose). Flies equally preferred the sweet and

bittersweet options at 10-fold sucrose concentration ratio (Fig. 1B, 50 mM vs. 500 mM sucrose +

1 mM quinine). This equal-preference point was identical at all of the tested food deprivation

durations (Fig. 1B), suggesting an external taste sensation and internal hunger state equilibrium

at this concentration ratio. The equal-preference point depends on the sucrose concentration ratio

between the two options and not absolute concentration (Fig. S1B). These results indicate that

hungry flies tradeoff the appetitive (sweet) and aversive (bitter) values of food in making feeding

decisions, and are consistent with known effects of bitter compounds on feeding, in which

increasing bitterness of food decreases feeding(Meunier et al., 2003).

To further characterize fly behavior in the decision task and determine whether sensory cues

other than taste could affect food choice, we also determined the spatial location of each fly at

the end of the decision task. As expected, position preference mirrored ingested food preference

(Fig. 1C). We also addressed whether size or sex ratio of groups of flies tested together

influenced the decision, since social interactions driven by group size (Ramdya et al., 2017;

Rooke et al., 2020) and sex ratio can affect fly behavior. There was no effect on ingested food

preference of male-to-female ratio within groups of flies tested together (Fig. 1D). At the equal-

preference condition (21h deprivation, 50 mM sucrose vs. 500 mM sucrose + 1 mM quinine),

neither food preference and group size (Fig. 1E), food preference and percent of flies that ate

(Fig. 1F), nor percent of flies that ate and group size (Fig. 1G) were correlated. There was also

no significant prediction of preference index by group size or percent of flies that ate in a

multiple regression model (Fig. 1H, Table S1), indicating no interaction between these variables

in the decision task. These results suggest that taste is the most salient sensory cue affecting food

choice in the decision assay.

A decision making neuronal ensemble converges on the fan-shaped body

During foraging, animals estimate values of internal state and external sensory parameters,

such as hunger level, valence of available foods, location of food, etc. These value estimates are

then integrated to form a decision. While neural substrates integrating internal state and external

stimuli in the *Drosophila* brain remain unclear, there are hints in the literature of cell-populations

and brain regions that could be involved. Various neuromodulators and the neurons that secrete

them regulate hunger dependent food intake (Cannell et al., 2016; Hentze et al., 2015; Lin et al.,

2019; Nassel and Zandawala, 2019; Wu et al., 2003; Yang et al., 2018; Zandawala et al., 2018),

reward (Huetteroth et al., 2015; Liu et al., 2012a; Lyutova et al., 2019; Yamagata et al., 2015) or

punishment (Riemensperger et al., 2005), and memory (Huetteroth et al., 2015; Masek et al.,

2015; Yamagata et al., 2015). The mushroom body is an insect central brain region involved in

gustatory learning and memory (Kirkhart and Scott, 2015; Masek and Scott, 2010) and valence

encoding (Aso et al., 2014), and is thought to be a major center controlling higher-order

behaviors, including associative learning (Cohn et al., 2015; Lewis et al., 2015; Tsao et al.,

2018). The insect central complex is an evolutionarily conserved central brain region whose

ellipsoid body and protocerebral bridge sub-regions have been implicated in navigation (Giraldo

et al., 2018; Green et al., 2019; Honkanen et al., 2019; Kim et al., 2017b; Mathejczyk and

Wernet, 2019; Seelig and Jayaraman, 2015; Sun et al., 2017; Turner-Evans et al., 2017) and

sleep (Guo et al., 2018; Liu et al., 2019; Liu et al., 2016). The central complex fan-shaped body,

a laminar sub-region, has been implicated in sleep (Donlea et al., 2014; Donlea et al., 2018; Liu

et al., 2012b; Pimentel et al., 2016) and ethanol preference (Azanchi et al., 2013; Scaplen et al.,

2020). Various neuromodulators (Kahsai and Winther, 2011), their receptors (Al-Anzi et al.,

2010; Cavey et al., 2016; Donlea et al., 2018), and dopaminergic inputs (Kim et al., 2020; Qian

et al., 2017) co-localize in the layers of fan-shaped body. We hypothesized that value estimates

of internal state and external environment encoded by neuromodulatory neurons will be required

inputs to higher-order brain regions for integration and generation of decisions. To test this

hypothesis, we genetically targeted acute optogenetic activation and inhibition to specific neural

subsets in neuromodulatory and higher-order brain centers using the GAL4-UAS binary

expression system (Brand and Perrimon, 1993), while flies actively sampled and consumed food

at the equal-preference condition (Fig. 1A, 1B, 21h food deprivation, 50 mM sucrose vs 500mM

sucrose + 1mM quinine). For optogenetic activation, we used red-light-sensitive CsChrimson

channelrhodopsin (Klapoetke et al., 2014), and for optogenetic inhibition we used green-light-

sensitive anion-conducting channelrhodopsin GtACR1 (Mohammad et al., 2017). This

optogenetic interrogation of modulatory neurons and higher order brain regions revealed

neuropeptidergic neurons (Leucokinin, Allatostatin A, NPF, DH44), subsets of dopaminergic

neurons (PPL1-$\gamma2\alpha$'1, PPL1-$\alpha3$, PAM-$\alpha1$), and a narrow subset of fan-shaped body layer 6

neurons (FBl6) whose activation or inhibition significantly shifted food choice in the equal-

preference condition (Fig. 2A).

The specific effects of activation or inhibition of restricted neuron populations on the sweet

versus bittersweet decision provides insights into the roles of such populations in value

estimation and integration. Optogenetic activation of Leucokinin (Lk) neurons suppressed

feeding in food deprived flies (Fig. 2A, 2B left panel, Fig. S2A-B), suggesting that Lk may

encode or relay metabolic state information. To confirm that Lk secreted by these neurons was

the molecular basis of this feeding suppression, we simultaneously knocked down Lk expression

with genetically encoded RNAi while optogenetically activating Lk neurons. While almost no

flies consumed food when Lk neurons were activated without Lk RNAi, simultaneous Lk

knockdown prevented this feeding suppression (Fig. 2B left panel, Fig. S2A-B). This indicates

that Lk secretion mediates feeding suppression by Lk neurons. Optogenetic silencing of Lk

neurons shifted preference towards bittersweet (Fig. 2A, 2B left panel, Fig. S2A-B). Feeding

suppression induced by Lk neuron activation implies a decrease in perceived hunger of food

deprived flies, consistent with the shift towards higher calorie bittersweet food by Lk neuron

inhibition reflecting increased perceived hunger. Optogenetic activation of Allatostatin A (AstA)

neurons shifted preference towards sweet, while inhibition shifted preference towards bittersweet

(Fig. 2A, 2C left panel). We confirmed that AstA was the molecular basis of this shift in

preference by simultaneous AstA RNAi knockdown and optogenetic activation of AstA neurons

(Fig. 2C left panel). Optogenetic activation of NPF neurons shifted preference towards sweet,

and this shift was abolished by simultaneous activation and NPF RNAi knockdown (Fig. 2A, 2D

left panel). Optogenetic activation of DH44 neurons had no significant effect, but inhibition

shifted preference towards bittersweet (Fig. 2A, 2E left). Dopaminergic subsets involved in

aversive memory (Fig. 2A PPL1-$\gamma2\alpha$'1) (Berry et al., 2012), taste conditioning (Fig. 2A PPL1-

$\alpha3$) (Masek et al., 2015), and long-term memory (Fig. 2A PAM-$\alpha1$) (Yamagata et al., 2015) also

affected food choice. Specifically, optogenetic activation of these dopaminergic subsets shifted

preference towards bittersweet (Fig. 2A). Optogenetic activation of various subsets of mushroom

body neurons, a brain region controlling higher-order behaviors, did not affect preference (Fig.

2A). However, optogenetic inhibition of a specific subset of FBl6 neurons shifted preference

towards bittersweet (Fig. 2A, 3A left panel). Value estimates of internal state and external

sensory cues, which are likely computed by identified modulatory neurons, are crucial for

decision making. Because many of these neuromodulators and their receptors (Al-Anzi et al.,

2010; Cavey et al., 2016; Donlea et al., 2018; Kahsai and Winther, 2011; Kim et al., 2020; Qian

et al., 2017) co-localize in the fan-shaped body, we thus hypothesize that neuromodulatory inputs

encoding value estimates are integrated in FBl6, where a decision is generated.

To test this hypothesis and determine whether the neurons we identified in our optogenetic

screen are connected in a behaviorally relevant ensemble, we employed a chemoconnectomics

approach (Deng et al., 2019) using cell-specific genetically encoded RNAi knockdown of

neuropeptide and dopamine receptors. By knocking down a specific receptor in a candidate

target neuron, we test whether that neuron receives direct input from a neuron secreting the

cognate ligand for that receptor. Knockdown of neuropeptide or dopamine receptors in Lk

neurons did not shift preference (Fig. 2B right panel). This indicates that Lk neurons do not

receive direct inputs from decision-relevant dopaminergic neurons or from neurons secreting the

ligands for tested neuropeptide receptors, and implies that Lk neurons must receive food

preference and hunger related information indirectly from other neurons. RNAi knockdown of

Dop1R1 dopamine receptor in AstA neurons shifted preference towards sweet (Fig. 2C right

panel), indicating that AstA neurons receive food-preference-relevant dopaminergic inputs.

RNAi knockdown of Lkr and Dop1R1 in NPF neurons shifted preference towards sweet (Fig. 2D

right panel), indicating that NPF neurons receive food-preference-relevant Lk and dopaminergic

inputs. RNAi knockdown of DopEcR dopamine receptor in DH44 neurons shifted preference

towards bittersweet (Fig. 2E right panel), indicating that DH44 neurons receive food-preference-

relevant dopaminergic inputs.

Importantly, RNAi knockdown of Lkr, AstA-R1, or DH44-R1 receptors in FBl6 neurons

shifted preference towards bittersweet (Fig. 3A right panel), indicating that FBl6 neurons are

directly modulated by these three neuropeptides to affect food choice. Furthermore, direction of

change in food preference induced by receptor RNAi in FBl6 mirrors direction of change in food

preference induced by optogenetic inhibition of the corresponding neuropeptide neurons. First,

both AstA-R1 receptor RNAi in FBl6 neurons (Fig. 3A right panel) and optogenetic inhibition of

AstA neurons (Fig. 2C left panel) shifted food preference towards bittersweet. Second, both

DH44-R1 receptor RNAi in FBl6 (Fig. 3A right panel) and optogenetic inhibition of DH44

neurons (Fig. 2D right panel) shifted preference towards bittersweet. Finally, both Lkr receptor

RNAi in FBl6 (Fig. 3A right panel) and optogenetic inhibition of Lk neurons (Fig. 2B right

panel) shifted preference towards bittersweet. RNAi knockdown of dopamine receptors in FBl6

had no effect (Fig. 3A right panel). This matrixed chemoconnectomics strategy mapped the

neuromodulatory connections between nodes in the ensemble to control choice, and uncovered

FBl6 as a previously unknown convergence node that is well positioned to integrate sensory,

metabolic, and experiential information for decision making.

Fan-shaped body neurons encode choice

Value estimates of internal state variables, such as hunger, external environmental cues,

appetitive or aversive value of food (i.e., valence), and past sensorimotor experience are

integrated with one another and transformed into behavioral choice. This raises the key question

whether FBl6 neurons compute value estimates or, instead, integrate these value estimates to

encode choice. If FBl6 neurons encode or estimate value of metabolic parameters such as hunger

or satiety, manipulating their activity would be expected to directly influence feeding behavior.

To test this, we first quantified the proportion of flies that consumed food during optogenetic

activation or inhibition of FBl6 neurons. During FBl6 neural activity manipulation, most food-

deprived flies consumed food while most fed flies did not (Fig. 3B, Table S1), revealing that

FBl6 activity manipulation neither prevented food consumption by food-deprived flies nor

induced food consumption by fed flies. These results demonstrate that hunger state is not

affected by FBl6 neural activity. Next, we quantified food consumption per fly during

optogenetic activation or inhibition of FBl6 neurons and found no effect on total amount of food

consumed per fly (Fig. 3C, Table S1). These manipulations also did not affect the quantity of

sweet versus bittersweet food consumed per fly within each condition (Fig. 3D, Table S1), again

demonstrating that FBl6 neural activity does not affect hunger state. Rather, the shift in food

preference induced by optogenetic inhibition of FBl6 (Fig. 3A left panel) was due to an increase

in the proportion of flies consuming bittersweet over sweet food, and not each fly consuming a

larger quantity of bittersweet food. Taken together, these results demonstrate that FBl6 does not

encode or affect metabolic signals of hunger or satiety.

If a neural population encodes reward or punishment, then manipulating the activity of that

population could directly shift a decision without integrating value estimates or encoding choice.

To test whether FBl6 activity is intrinsically rewarding or punishing, we illuminated half the

decision arena with either red (for optogenetic activation) or green (for optogenetic inhibition)

light in the absence of food and quantified preference for the illuminated versus dark sectors.

Flies expressing optogenetic actuators CsChrimson or GtACR1 in FBl6 did not alter their

preference for illuminated versus dark sectors compared to non-expressing controls (Fig. 3E),

demonstrating that FBl6 activation or inhibition is intrinsically neither rewarding nor punishing.

Animals accumulate information about past experience to inform future decisions. We

hypothesized that FBl6 integrates internal hunger state and external food-related value estimates

with experiential information to drive decisions. To understand how past experience affects FBl6

activity, we recorded FBl6 neural taste responses in flies that had different food-related

experiences. Equal-preference taste stimuli were presented to forelegs of the fly (Fig. 4A) while

measuring intracellular Ca^{2+} signals ratiometrically using GCaMP6f (Chen et al., 2013) and

tdTomato (Fig. 4B) expressed in FBl6 (Fig. 4B-C). First, we tested the effect of hunger on FBl6

taste responses in naïve flies, i.e., flies that had not experienced the decision task at all. FBl6

neurons of naïve food-deprived flies were strongly inhibited by the bittersweet stimulus, but not

sweet (Fig. 4D-E, "naïve deprived"). Flies often find bittersweet food aversive (Masek and Scott,

2010), and thus inhibitory taste responses in FBl6 to bittersweet food may represent rejection of

this option. In contrast to hungry food-deprived flies, fed flies reject both foods in the decision

task (Fig. 3B). If FBl6 neural activity represents rejected choice, then we expect FBl6 inhibition

by both sweet and bittersweet stimuli in naïve fed flies. Indeed, FBl6 neurons of naïve fed flies

were strongly inhibited by both bittersweet and sweet stimuli (Fig. 4D-E, "naïve fed").

Importantly, this is the first observation of gustatory responses in the *Drosophila* fan-shaped

body, or any other part of the central complex.

If, as in naïve flies, FBl6 neural activity also represents rejected choice in flies that have had

the opportunity to decide between sweet and bittersweet, then we expect decision outcomes to

affect FBl6 taste responses. Specifically, we expect FBl6 neurons of each fly to be strongly

inhibited by the rejected choice for that fly, based on its individual decision. Consistent with this,

when taste responses were subsequently measured in flies that chose sweet food in the decision

task, FBl6 neurons were strongly inhibited by the rejected bittersweet stimulus, but not by the

chosen sweet (Fig. 4D-E, "chose sweet"). Furthermore, FBl6 neurons of flies that chose

bittersweet food were strongly inhibited by the rejected sweet stimulus, but not by the chosen

bittersweet (Fig. 4D-E, "chose bittersweet"). Finally, FBl6 neurons of flies that chose neither

option, i.e., rejected both, were strongly inhibited by both bittersweet and sweet stimuli (Fig. 4D-

E, "chose neither"). In summary, FBl6 neural activity is strongly inhibited by the taste of food

that each individual fly rejects, suggesting that suppression of FBl6 activity is the neural

representation of behavioral food choice.

We then tested whether FBl6 neurons encode not only taste of rejected food, but also

behavioral choice associated with other sensory cues. Specifically, we determined whether FBl6

responds to water, and whether such responses are dependent on internal state and prior

experience of each fly. While FB16 neurons of food-deprived flies that chose sweet or neither in

the decision assay were inhibited by water applied to the forelegs, those of fed or food-deprived

naive flies were not (Fig. S4A-B). Furthermore, choice outcome in the decision assay affected

water responses, with water inhibiting FB16 neurons of flies choosing sweet or neither, but not

of flies choosing bittersweet (Fig. S4A-B). Individual flies that chose bittersweet higher-calorie

food over sweet are likely hungrier based on their individual behaviors during maintenance and

food deprivation than flies that chose lower-calorie sweet or neither (i.e., didn't consume), and

may correspondingly be thirstier. This could then suggest that the FBl6 inhibition by water we

observed specifically in flies that chose sweet or neither represents rejection of water by these

less-thirsty flies, and that lack of inhibition in flies that chose bittersweet represents acceptance.

Regardless of the specific interpretation (which requires further study), these water responses

indicate that, like for taste, FBl6 encodes not stimulus identity *per se*, but rather state- and

experience-dependent behavioral choice.

These state- and experience-dependent FBl6 responses do not simply encode taste or water

identity, but rather encode behavioral choice that arises from integration of internal state and

external sensory inputs. This neural representation of choice is modulated by taste (sweet vs.

bittersweet), previous experience (naïve vs. experience with the two-choice task, including

decision outcome), and hunger state (food-deprived vs. fed) (Fig. 4D-E, Fig. S4A-B). FBl6

neurons likely receive inputs encoding these parameters directly from AstA, DH44, and Lk

neurons, and indirectly from NPF and dopamine neurons. These signals within the decision

ensemble are then integrated by FBl6 to generate a representation of choice, which is relayed to

downstream motor circuits to implement the behavioral decision (Fig. 4F).

Discussion

Animals make decisions about which foods to consume by integrating internal physiological

state signals with external sensory cues. Here we delineated a neuronal ensemble in *Drosophila*

that underlies food-related decision making during sensory conflict between sweet and

bittersweet food choices (Fig. 1). Activating or silencing particular nodes in this ensemble shifts

the decision balance between sweet and bittersweet choices (Fig. 2, 3). Inputs encoding hunger

state, taste identity, and past experience converge on FBl6 neurons, which integrate these inputs

to generate a neural representation of food choice that drives motor outputs for foraging

decisions (Fig. 4).

Animals assess and assign value estimates to internal state and external environmental

parameters before integrating these estimates to guide adaptive decision making. Upstream

neuromodulatory subsets in the decision ensemble we identified have known roles in hunger

dependent food intake (Dus et al., 2015; Hentze et al., 2015; Yang et al., 2018; Zandawala et al.,

2018), reward (Shao et al., 2017), valence (Berry et al., 2012; Masek et al., 2015), and long-term

memory (Berry et al., 2012; Masek et al., 2015; Yamagata et al., 2015). These modulatory

neurons are well positioned to estimate value of features in the sensory environment and internal

hunger state. For example, AstA neuron activity influences relative carbohydrate and protein

preference (Hentze et al., 2015), while DH44 neurons sense sugars (Dus et al., 2015) and amino

acids (Yang et al., 2018). Thus, AstA and DH44 neurons could convey food identity information

to FBl6. NPF neuron activation is inherently rewarding (Shao et al., 2017), and thus could

convey food valence information. Lk neurons have been implicated in nutrient sensing

(Zandawala et al., 2018), and their activity regulates feeding in food deprived flies (Fig. 2A-B),

suggesting that internal metabolic state information could reach FBl6 through Lk/Lkr signaling.

Dopaminergic subsets involved in aversive memory (PPL1 $\gamma2\alpha$'1) (Berry et al., 2012), taste

conditioning (PPL1 α3) (Masek et al., 2015), and long-term memory (PAM α1) (Yamagata et al.,

2015) also affected food choice (Fig. 2A) and could provide signals for predicting and updating

value estimates in working memory, analogously to primate dopaminergic ventral tegmental area

(Sugrue et al., 2005). FBl6 neurons project their axons into the fan-shaped body (Kim et al.,

2020; Qian et al., 2017), have dense dendritic projections in the superior medial protocerebrum

(SMP), and have sparse dendritic projections in superior intermediate protocerebrum (SIP) and

superior lateral protocerebrum (SLP) (Kim et al., 2020; Qian et al., 2017) (Fig. 4G). In these

higher brain regions, FBl6 receives synaptic input from dopaminergic neurons (Liu et al., 2012b;

Pimentel et al., 2016) that regulate sleep (Liu et al., 2012b; Pimentel et al., 2016) and ethanol

preference (Azanchi et al., 2013). Interestingly, direct dopaminergic input to FBl6 through

dopamine receptors did not influence food choice (Fig. 3A). Instead, indirect dopaminergic

inputs likely conveyed by AstA, DH44, and NPF neurons regulated food choice (Fig. 2B-E).

Other neuromodulatory neurons, second-order sensory projection neurons, and interneurons

also likely interact with this decision ensemble to convey gustatory and other sensory inputs.

There are no known projections of primary gustatory sensory neurons or second-order taste

neurons (Bohra et al., 2018; Kain and Dahanukar, 2015; Kim et al., 2017a; Miyazaki et al., 2015)

to the fan-shaped body. Food preference was affected neither by acute optogenetic nor by

chronic inhibition (using tetanus toxin light chain to impair vesicle docking) of the only known

taste projection neurons (TPN3) that relay bitter taste information to SLP and are essential for

conditioned taste aversion (Kim et al., 2017a) (Fig. S5A). However, SMP, SLP, and SIP higher

brain regions, to which FBl6 neurons also densely project (Fig. 4G), have been implicated as

target areas for second-order taste neurons (Talay et al., 2017). It thus is possible that sweet and

bitter taste information reaches the fan-shaped body through second-order taste projection

neurons that terminate in these higher brain regions. Another possibility is that taste identity

information is conveyed to FBl6 by the dopaminergic neurons that respond to sweet (Huetteroth

et al., 2015) and bitter (Masek et al., 2015) tastes, are involved in taste conditioning (Huetteroth

et al., 2015; Masek et al., 2015), and shift preference in our decision assay (Fig. 2A, PAM-α1,

PPL1-α3).

A value integrator for food-related decision making requires as inputs estimates of taste

identity, previous experience, and hunger state. FBl6 neuron activity is modulated by these

parameters and strongly inhibited by the taste of rejected food choice. Lack of FBl6 inhibition in

response to a taste stimulus could represent acceptance of that option (Fig. 4D-E). The

mechanistic connection between optogenetic inhibition of FBl6 shifting food preference to

bittersweet (Fig. 3A), on the one hand, and rejected food choice strongly inhibiting FBl6 neural

activity (Fig. 4D-E), on the other hand, requires further investigation. It is possible that

optogenetic inhibition of FBl6 throughout the decision task disrupts value integration and

decision making resulting in the hungry fly defaulting to higher calorie bittersweet food.

Regardless, our findings indicate that the behavioral response of an individual fly can be

predicted by dynamic neural activity in FBl6.

Mammalian studies provide converging evidence for multiple interconnected networks in

frontal cortex and basal ganglia that compute and store value estimates of sensory environment

and motor events in that environment required for decision making (Lee et al., 2012; Sugrue et

al., 2005). The neural ensemble described in this study is an analogous framework of

interconnected networks for potentially storing, computing, and updating value estimates that are

likely integrated by FBl6 neurons. While decision making theories in mammals have focused on

how values are represented in the brain (Lee et al., 2012; Sugrue et al., 2005), mechanisms by

which the brain integrates value information to make decisions remain unclear (Shadlen and

Kiani, 2013; Tsetsos et al., 2012). Changes in neural spiking pattern are thought to underlie

sensory information accumulation and integration, with a decision being made when neural

firing rate reaches a threshold (Gold and Shadlen, 2007; Park et al., 2014; Wang, 2008).

Persistent neural activity and synaptic plasticity changes are thought to underlie storage of

sensory information and past choice history to guide adaptive decisions (Wang, 2008). Future

work is required to test the proposed hypotheses of specific roles for each node in the fly

decision ensemble, how upstream inputs are integrated in FBl6, how this integration is

transformed into the representation of choice, and how downstream motor circuits implement the

foraging decision.

Methods

Fly husbandry

Flies were cultured on standard cornmeal medium on 12:12 light:dark cycle at 25ºC. *w1118* lab

stock was used as wild type. All other genotypes and their sources are described in Table S2. 2-5

day old flies were wet starved for 2-48 h (based on experiment design) on a wet Kimwipe with

1.5 ml distilled water. For optogenetic experiments, flies were food deprived for 21 h before

testing on 0.4 mM all-*trans* Retinal (Cayman Chemicals) in 1% agar. Flies for RNAi knockdown

and their controls were moved to 28ºC for 21 h the day before testing, i.e., during food

deprivation, to induce strong RNAi. RNAi control was created for each GAL4 line by crossing

the respective GAL4 to UAS-Valium (see Table S2). All RNAi lines that we used were from

Harvard TRiP project (Ni et al., 2009; Perkins et al., 2015) and have been validated by

independent groups (see Table S2). Flies for simultaneous optogenetic and RNAi experiments

were created using the genotypes mentioned in Table S2. All experiments were conducted at

Zeitgeber Time 3-6.

Two-choice assay and optogenetics

Sweet foods were made with different concentrations of sucrose and bittersweet foods with

500mM sucrose (Sigma) and 1mM quinine (Alfa Aesar or Beantown Chemicals, CAS#207671-

44-1) dissolved in 1% agarose (AmericanBio) made in distilled water. 0.04% w/v red dye

(Sulforhodamine B, MP Biochemicals, CAS# 3520-42-1) and 0.02% w/v blue dye (Erioglaucine

A, Alfa Aesar, CAS# 3844-45-9) were used for food coloring. Dye colors were alternated

between sweet and bittersweet foods for each condition and there was no preference for one dye

over the other at the concentrations used. Fly arenas were prepared by pouring agarose based

foods in two-compartment petri-dishes (90-100 mm diameter) from Kord Valmark, EMS, or

Fisher Scientific. Because of a thin physical barrier between the two compartments in the arena

there was no diffusion between the two foods. Groups of 20-35 flies were aspirated and

introduced into the arena 5-10 sec before the start of the experiment. All experiments were

conducted in dark so that there was no effect of food color on preference. Arenas were placed on

a platform with IR backlight for video recording using a Flea Pointgrey camera (FL3-U3-

20E4C/M) at 15 fps. For optogenetics, we used high-power LEDs (Luxeon) placed adjacent to

backlight IR LEDs (based on Janelia ID&F design) of 627 nm (for CsChrimson) and 520 nm (for

GtACR1) that were controlled using Arduino Uno. For optogenetic screen, both red and green

lights were pulsed at 100% max intensity, 50Hz, 25% duty cycle. For follow up experiments,

CsChrimson experiments were conducted at 25% max intensity; GtACR1 follow up was done at

screen condition. Light was pulsed for the entire duration of the experiment. At the end of the

experiment, flies were anaesthetized using CO_2 and their belly color was recorded under a

dissection microscope. Preference index (PI) was calculated as (no. of sweet food flies + 0.5 no.

of both food flies) - (no. of bittersweet food flies + 0.5 no. of both food flies) / no. of total flies

that ate, where negative PI would mean that more number of flies ate bittersweet food.

Food intake quantification

Food intake was quantified using spectrophotometry as previously described (Deshpande et al.,

2014; Wong et al., 2009). After recording belly color flies were frozen in 1.5 ml Eppendorf tubes

at -20⁰C until intake quantification (1-2 days). Flies from each trial were separately homogenized

in distilled water (5 µl /fly) using a motorized pestle (BT Labsystems, BT703) for 1.5 min and

centrifuged at 13000 rpm for 5 min. Absorbance of the debris-cleared 2 µl supernatant was

measured on NanoDrop 2000 Spectrophotometer (Thermo Fisher Scientific) at 565 nm (for red

dye) and 630nm (for blue dye). Flies that ate uncolored 1% agarose with 50mM sucrose were

used as blank for baseline control. Red and blue dye concentrations were interpolated using their

respective standard curves (GraphPad Prism) acquired from serial dilutions of single dyes in

distilled water. Since we knew the number of flies that ate each color per trial, we could calculate

per fly blue and red concentrations in the same solution.

Calcium imaging and data analysis

3-5 day old flies (naïve or after two-choice assay) were aspirated and positioned in a custom

made fly holder in which they were glued using two-part transparent epoxy (Devcon). Only the

top of fly head (for imaging) and the forelegs (for taste delivery) were outside the holder, while

the rest of the fly, including proboscis were restrained in the fly holder. No anesthesia was used.

A small piece of head cuticle was dissected and air sacs removed using a 30-gauge syringe

needle and fine forceps, immediately followed by sealing the head capsule with a translucent

surgical silicone adhesive (Kwil-Sil, WPI). Dissected fly was then placed in a humidified

chamber for 15 min recovery before imaging.

Calcium imaging was performed on a Zeiss Axio Examiner upright microscope with 20x

air objective and a Colibri module for LED control. tdTomato was excited at 555 nm (80%

intensity) and GCaMP6 at 470 nm (100% intensity). An image splitter (Photometrics DV-2) was

used to split red and green channels and acquire simultaneous images for tdTomato and

GCaMP6 using a Hamamatsu ORCA-R2 C10600 camera. Images were acquired at 5 fps with

variable baseline (3 to 10 s required for stable tastant delivery) followed by tastant application to

the forelegs, using a syringe, for 3 s and 4 s of no tastant. Excess tastant was wicked from the

forelegs using absorbent tissue paper between each application. 10 s inter-trial interval was used

during which all lights were off. Water was always applied first, followed by either sweet or

bittersweet tastant. Sequence of sweet and bittersweet was alternated between flies. Sweet: 50

mM sucrose in distilled water; bittersweet: 500 mM sucrose + 1 mM quinine in distilled water.

For flies that chose sweet in the two-choice behavior assay only trials with sweet as the first

tastant were averaged and for flies that chose bittersweet, only trials with bittersweet as the first

tastant were averaged.

Pixel intensities were extracted in Fiji followed by data analysis in MATLAB, both using

custom written code. After background subtraction using Fiji's rolling-ball method (20 px), ROIs

were manually drawn and saved on the tdTomato image, and superimposed on the GCaMP

image (both reporters were expressed in the same neurons using the same driver) for mean ROI

pixel intensity extraction. The saved intensity signals were then analyzed in MATLAB.

tdTomato and GCaMP traces were individually corrected for photobleaching by fitting a single

exponential function. Corrected GCaMP trace was then divided by the corrected tdTomato trace

to obtain the ratiometric fluorescence trace (R). For relative fluorescence fold change ($\Delta R/R_0$)

determination, baseline fluorescence (R_0) was calculated by averaging R over 2 s preceding

tastant application. Peak $\Delta R/R_0$ was calculated during 4 s following tastant application.

EM reconstruction

Electron microscopy images were reconstructed from publically available Janelia FlyEM

hemibrain data using neuPRINT (Xu et al., 2020). Neuron identities were confirmed in

NeuronBridge (Jody et al., 2020) by cross-referencing EM traced FBl6 neurons matched with

light microscopy images of 84C10-GAL4 from FlyLight (Meissner et al., 2020). FBl6 mesh,

whole FB mesh, and SMP, SIP and SLP brain region meshes were used to depict brain regions

with neural projection areas.

Statistics

All data were plotted in either Python or MATLAB using custom written code. Statistics were

carried out in GraphPad Prism. If all data passed Kolmogorov-Smirnov normality test, ANOVA

was conducted, otherwise Kruskal-Wallis test was conducted, both followed by appropriate post-

hoc tests. Details of statistics for each figure are provided in Table S1. Sample sizes are reported

in parentheses next to dataset name in Table S1.

Acknowledgements

We thank Peter Niesman for help with calcium imaging data collection and Jason Braco for

helpful discussions. Yichen Luo from John Carlson's lab provided useful information for design

of the taste delivery imaging rig. Tanya Wolff provided insights into fan-shaped body

neuroanatomy. We also thank Gerry Rubin and Tanya Wolff for sharing unpublished fly lines

ss00208 and ss00225. These studies were supported in part by the National Institute of

Neurological Disorder and Stroke, National Institutes of Health (NIH) (R01NS091070) and the

National Institute of General Medical Sciences, NIH (R01GM098932).

Author contributions

P.S. conceptualized the study, designed and performed experiments, and analyzed data; L.Y.M.

contributed to experimental design and fly dissection; P.S. and M.N.N. interpreted data; P.S. and

M.N.N. prepared the manuscript with input from all authors.

Data and code availability

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the Lead Contact, Dr. Michael N. Nitabach (michael.nitabach@yale.edu). The data

and custom code that support the findings from this study are available from the Lead Contact

upon request.

Declaration of interests

Authors declare no competing interests.

Figure legends

Figure 1. Hungry flies make trade-offs between the appetitive and aversive value of food.

A, Schematic of the two-choice decision making assay. Sweet and bittersweet foods are prepared

in dissolved agarose, mixed with food dyes (e.g., sweet with blue and bittersweet with red), and

gelled in a circular arena. Dye colors are counterbalanced within each food tradeoff condition.

Flies are introduced into the dark food arena to walk, sample, and consume freely for 5 min,

while they are video recorded with infrared (IR) backlighting. Flies are then anaesthetized and

abdomen color is recorded under a dissecting microscope, indicating ingested food choice.

Preference index is calculated as (no. of sweet color flies + 0.5 purple flies) - (no. of bittersweet

color flies + 0.5 purple flies)/total no. of flies that ate, where positive index means flies preferred

sweet and negative index means flies preferred bittersweet. **B**, Preference index of wild-type

(*w1118*) flies deprived of food for increasing durations indicate the tradeoff between the sweet

and bittersweet food and reveal equal preference when the choice is between 50 mM sucrose-

only and 500 mM sucrose + 1mM quinine. **C**, Position preference index (i.e., sweet or

bittersweet sector preference as defined by the location of the flies at the end of the assay)

matches ingested food preference, with equal preference at 50 mM sucrose-only vs. 500 mM

sucrose + 1 mM quinine. **D,** Preferences of male and female flies within each group were

indistinguishable for all conditions tested. There was no significant correlation between

preference index and group size per trial (**E**), preference index and % of flies that ate per trial

(**F**), or % of flies that ate per trial and group size (**G**). **H,** Neither group size nor % flies that ate

significantly predicted preference index in a multiple regression model, indicating no interaction

between these measures. **B-D**, Plots show mean ± 95% CI, violins depict kernel density

estimation of the underlying data distribution with the width of each violin scaled by the number

of observations at that y-value. Each violin summarizes $10 \leq$ trials ≤ 30 with mode $= 10$. **E-H,**

Heatmaps depict bivariate distribution visualized using a kernel density estimation procedure;

darker regions indicate higher data density. r^2 is the square of Pearson's coefficient. See Table S1

for sample size and statistics.

Figure 2. A decision making neuronal ensemble revealed by combined optogenetic and

chemoconnectomic strategy. A, Cell-specific optogenetic activation and inhibition screen was

performed at the equal preference condition (50 mM sucrose vs. 500 mM sucrose + 1 mM

quinine) using flies deprived of food for 21 h. Neuronal subsets were genetically targeted using

the GAL4-UAS binary expression system. Red-light-sensitive CsChrimson (Chr) was used for

optogenetic activation and green-light-sensitive GtACR1 (Gt) for optogenetic silencing.

Optogenetic manipulation of AstA, DH44, Lk, and NPF neuropeptides; PPL1 γ2α'1, PPL1 α3,

and PAM α1 dopaminergic subsets; and fan-shaped body layer 6 neurons (FBl6) significantly

shifted preference away from equal preference compared to respective empty>Chr and empty>Gt

controls. Purple mean point = control, blue mean point = significant sweet preference, red mean

point = significant bittersweet preference, yellow mean point = simultaneous optogenetic

activation and RNAi. **B** (left), Optogenetic activation of Leucokinin (Lk) neurons suppressed

feeding in food deprived flies, while optogenetic inhibition shifted the preference towards

bittersweet. Simultaneous Lk RNAi in Lk neurons abolished the suppression of feeding induced

by optogenetic activation. **B** (right), RNAi in Lk neurons of analogous receptors for other

candidate neuromodulators had no effect. Adjacent schematic shows that Lk neurons convey

hunger state information to downstream neurons expressing Lk receptors. Lk neuron activation

suppresses feeding, conveying decreased hunger, while Lk inhibition leads to consumption of

higher caloric food, conveying increased hunger. **C** (left), Optogenetic activation of Allatostatin

A (AstA) neurons shifted preference towards sweet while optogenetic inhibition shifted it

towards bittersweet. Simultaneous AstA RNAi abolished the preference shift induced by

optogenetic activation. **C** (right), Dop1R1 RNAi in AstA neurons shifted preference towards

sweet. Adjacent schematic shows that AstA neurons receive decision-relevant direct

dopaminergic inputs through Dop1R1. Both AstA neuron activation and inhibition shift food

decision in opposite directions and send decision-relevant information to downstream neurons

expressing AstA receptors. **D** (left), Optogenetic activation of NPF neurons shifted preference

towards sweet. Simultaneous NPF RNAi abolished the preference shift induced by optogenetic

activation. **D** (right), Lkr or Dop1R1 RNAi in NPF neurons each shifted preference towards

sweet. Adjacent schematic shows that NPF neurons receive decision-relevant direct

dopaminergic inputs through Dop1R1 and Lk inputs through Lkr receptors. NPF neuron

activation shifts food decision to sweet and NPF neurons send decision-relevant information to

downstream neurons expressing NPF receptors. **E** (left), Optogenetic activation of DH44

neurons had no effect on preference, while optogenetic inhibition shifted preference towards

bittersweet. **E** (right), DopEcR RNAi in DH44 neurons shifted preference towards bittersweet.

Adjacent schematic shows that DH44 neurons receive decision-relevant direct dopaminergic

inputs through DopEcR. DH44 neuron inhibition shifts food decision to bittersweet and DH44

neurons send decision-relevant information to downstream neurons expressing DH44 receptors.

Plots show mean ± 95% CI, with violins depicting full data distribution; $5 \leq$ trials ≤ 30 per

violin, mode = 10. See Table S1 for sample size and statistics. p<0.00001=****, p<0.0001=***,

p<0.01=**, p<0.05=*.

Figure 3. Fan-shaped Body layer 6 is the convergence node of a decision making ensemble.

A (left), Optogenetic activation of FBl6 neurons did not affect preference. In contrast,

optogenetic inhibition shifted preference towards bittersweet. **A** (right), RNAi knockdown of

AstA-R1, DH44-R1, or Lkr receptors in FBl6 shifted preference towards bittersweet. **B**,

Optogenetic activation or inhibition of FBl6 did not induce feeding in fed flies or prevent feeding

in food-deprived flies. **C-D**, Optogenetic activation (**C**) or inhibition (**D**) of FBl6 did not change

total food consumption per fly compared to respective empty-GAL4 controls. Food consumption

was quantified by extracting red and blue food dyes from flies after the assay and measuring the

extract absorbance using UV-vis spectrometry. Concentrations of individual dyes were

calculated per trial by interpolating absorbance in the standard curve for each dye. Total volume

of agarose ingested was then calculated from red and blue dye concentrations and divided by

number of flies that ate red and blue food in that trial. Total food volume is the average of

consumption by each fly regardless of type of food consumed. Within each group, there was no

significant difference in sweet vs. bittersweet food consumption per fly. Optogenetic inhibition

of FBl6 shifted preference towards bittersweet (**A,** left), because more flies consumed bittersweet

than sweet. **E**, Optogenetic activation or inhibition of FBl6 did not significantly change

preference (place PI) of flies for illuminated vs. non-illuminated sectors of the arena, indicating

that neither activation nor inhibition of FBl6 is inherently rewarding or punishing. Plots show

mean $\pm$ 95% CI, with violins depicting full data distribution. Statistically different means are

shown in different color as in Fig. 2. See Table S1 for sample size and statistics.

$p < 0.00001 = ****, p < 0.0001 = ***, p < 0.01 = **, p < 0.05 = *$.

Figure 4. Neural activity in FBl6 encodes food choice. A, *In vivo* Ca^{2+} imaging of taste

responses in FBl6 neurons of flies in different hunger states making different sweet-bittersweet

decisions. Sweet or bittersweet taste stimuli are applied to the forelegs, and changes in

Ca^{2+} responses are measured ratiometrically in FBl6 using GCaMP6f and tdTomato. **B,** Neuronal

expression in FBl6 reported by tdTomato. Region of interest for fluorescence measurement is

outlined in cyan. 84C10-GAL4 driver line is used to specifically target strong fluorescent protein

expression in FBl6 neurons (FlyLight; Jenett et al., 2012) (Fig. S3B-D). **C,** EM reconstruction of

FBl6 neurons shows neural projections restricted to layer 6 of the fan-shaped body. EM

reconstruction of traced FBl6 neurons from the Janelia hemibrain data was performed using

neuPrint and NeuronBridge for neurons targeted by 84C10-GAL4 and embedded in the surface

representation of standardized FBl6 brain region. **D,** Ratiometric calcium responses, $\Delta R/R_0$, of

flies with different hunger state and decision outcomes. Sweet (50mM sucrose) and bittersweet

(500 mM sucrose + 1 mM quinine) taste stimuli from the equal-preference condition were

applied for 3 s, and neural response was quantified for 4 s post-stimulus application. Taste

stimulus application is indicated by gray background region. FBl6 neurons were strongly

inhibited by the behaviorally rejected taste stimulus. Specifically, FBl6 neurons of flies that

chose sweet were inhibited by bittersweet taste, while FBl6 neurons of flies that chose

bittersweet were inhibited by sweet taste. (**D-E**). Calcium activity trace depicts mean $\Delta R/R_0 \pm$

95% CI. **E**, Peak $\Delta R/R_0$ shows significant difference between responses to rejected vs. chosen

taste within each fly condition. $p < 0.05 = *$ (see Table S1 for details on statistics). Points on graphs

depict mean $\pm$ 95% CI, with violins depicting full data distribution. **F,** Schematic of the decision

making ensemble converging on FBl6. FBl6 activity is the neural representation of behavioral

food choice. This activity is modulated by taste, previous experience, and hunger state. FBl6

neurons likely receive inputs encoding these parameters directly through AstA, DH44, and Lk

receptor signaling, and indirectly through NPF and dopamine (DA) pathways. FBl6 integrates

the converging information to form a representation of choice, which is relayed to downstream

motor circuits for behavior implementation. **G**, EM reconstruction of FBl6 neurons targeted by

84C10-GAL4 was performed as previously and embedded in the surface representation of

standardized whole fan-shaped body (FB). FBl6 neural projections to superior medial

protocerebrum (SMP), superior intermediate protocerebrum (SIP), and superior lateral

protocerebrum (SLP) higher brain regions are shown embedded in the surface representation of

these standardized higher brain regions.

References

Al-Anzi, B., Armand, E., Nagamei, P., Olszewski, M., Sapin, V., Waters, C., Zinn, K., Wyman,

R.J., and Benzer, S. (2010). The leucokinin pathway and its neurons regulate meal size in

Drosophila. Curr Biol *20*, 969-978.

Aso, Y., Sitaraman, D., Ichinose, T., Kaun, K.R., Vogt, K., Belliart-Guerin, G., Placais, P.Y.,

Robie, A.A., Yamagata, N., Schnaitmann, C.*, et al.* (2014). Mushroom body output neurons

encode valence and guide memory-based action selection in Drosophila. Elife *3*, e04580.

Azanchi, R., Kaun, K.R., and Heberlein, U. (2013). Competing dopamine neurons drive

oviposition choice for ethanol in Drosophila. Proc Natl Acad Sci U S A *110*, 21153-21158.

Berry, J.A., Cervantes-Sandoval, I., Nicholas, E.P., and Davis, R.L. (2012). Dopamine is

required for learning and forgetting in Drosophila. Neuron *74*, 530-542.

Bohra, A.A., Kallman, B.R., Reichert, H., and VijayRaghavan, K. (2018). Identification of a

Single Pair of Interneurons for Bitter Taste Processing in the Drosophila Brain. Curr Biol *28*,

847-858 e843.

Brand, A.H., and Perrimon, N. (1993). Targeted gene expression as a means of altering cell fates

and generating dominant phenotypes. Development *118*, 401-415.

Cannell, E., Dornan, A.J., Halberg, K.A., Terhzaz, S., Dow, J.A.T., and Davies, S.A. (2016). The

corticotropin-releasing factor-like diuretic hormone 44 (DH44) and kinin neuropeptides

modulate desiccation and starvation tolerance in Drosophila melanogaster. Peptides *80*, 96-107.

Cavey, M., Collins, B., Bertet, C., and Blau, J. (2016). Circadian rhythms in neuronal activity

propagate through output circuits. Nat Neurosci *19*, 587-595.

Chen, T.W., Wardill, T.J., Sun, Y., Pulver, S.R., Renninger, S.L., Baohan, A., Schreiter, E.R.,

Kerr, R.A., Orger, M.B., Jayaraman, V.*, et al.* (2013). Ultrasensitive fluorescent proteins for

imaging neuronal activity. Nature *499*, 295-300.

Chu, B., Chui, V., Mann, K., and Gordon, M.D. (2014). Presynaptic gain control drives sweet

and bitter taste integration in Drosophila. Curr Biol *24*, 1978-1984.

Cohn, R., Morantte, I., and Ruta, V. (2015). Coordinated and Compartmentalized

Neuromodulation Shapes Sensory Processing in Drosophila. Cell *163*, 1742-1755.

Deng, B., Li, Q., Liu, X., Cao, Y., Li, B., Qian, Y., Xu, R., Mao, R., Zhou, E., Zhang, W.*, et al.*

(2019). Chemoconnectomics: Mapping Chemical Transmission in Drosophila. Neuron *101*, 876-

893 e874.

Deshpande, S.A., Carvalho, G.B., Amador, A., Phillips, A.M., Hoxha, S., Lizotte, K.J., and Ja,

W.W. (2014). Quantifying Drosophila food intake: comparative analysis of current methodology.

Nat Methods *11*, 535-540.

Donlea, J.M., Pimentel, D., and Miesenbock, G. (2014). Neuronal machinery of sleep

homeostasis in Drosophila. Neuron *81*, 860-872.

Donlea, J.M., Pimentel, D., Talbot, C.B., Kempf, A., Omoto, J.J., Hartenstein, V., and

Miesenbock, G. (2018). Recurrent Circuitry for Balancing Sleep Need and Sleep. Neuron *97*,

378-389 e374.

Dus, M., Lai, J.S., Gunapala, K.M., Min, S., Tayler, T.D., Hergarden, A.C., Geraud, E., Joseph,

C.M., and Suh, G.S. (2015). Nutrient Sensor in the Brain Directs the Action of the Brain-Gut

Axis in Drosophila. Neuron *87*, 139-151.

FlyLight, J. Janelia FlyLight.

Giraldo, Y.M., Leitch, K.J., Ros, I.G., Warren, T.L., Weir, P.T., and Dickinson, M.H. (2018).

Sun Navigation Requires Compass Neurons in Drosophila. Curr Biol *28*, 2845-2852 e2844.

Gold, J.I., and Shadlen, M.N. (2007). The neural basis of decision making. Annu Rev Neurosci

30, 535-574.

Green, J., Vijayan, V., Mussells Pires, P., Adachi, A., and Maimon, G. (2019). A neural heading

estimate is compared with an internal goal to guide oriented navigation. Nat Neurosci *22*, 1460-

1468.

Guo, F., Holla, M., Diaz, M.M., and Rosbash, M. (2018). A Circadian Output Circuit Controls

Sleep-Wake Arousal in Drosophila. Neuron *100*, 624-635 e624.

Hentze, J.L., Carlsson, M.A., Kondo, S., Nassel, D.R., and Rewitz, K.F. (2015). The

Neuropeptide Allatostatin A Regulates Metabolism and Feeding Decisions in Drosophila. Sci

Rep *5*, 11680.

Honkanen, A., Adden, A., da Silva Freitas, J., and Heinze, S. (2019). The insect central complex

and the neural basis of navigational strategies. J Exp Biol *222*.

Huetteroth, W., Perisse, E., Lin, S., Klappenbach, M., Burke, C., and Waddell, S. (2015). Sweet

taste and nutrient value subdivide rewarding dopaminergic neurons in Drosophila. Curr Biol *25*,

751-758.

Jenett, A., Rubin, G.M., Ngo, T.T., Shepherd, D., Murphy, C., Dionne, H., Pfeiffer, B.D.,

Cavallaro, A., Hall, D., Jeter, J.*, et al.* (2012). A GAL4-driver line resource for Drosophila

neurobiology. Cell Rep *2*, 991-1001.

Jeong, Y.T., Shim, J., Oh, S.R., Yoon, H.I., Kim, C.H., Moon, S.J., and Montell, C. (2013). An

odorant-binding protein required for suppression of sweet taste by bitter chemicals. Neuron *79*,

725-737.

Jody, C., Cristian, G., Antje, K., Hideo, O., Rob, S., and Konrad, R. (2020). NeuronBridge

Codebase.

669 Kahsai, L., and Winther, A.M. (2011). Chemical neuroanatomy of the Drosophila central

670 complex: distribution of multiple neuropeptides in relation to neurotransmitters. J Comp Neurol

671 *519*, 290-315.

672 Kain, P., and Dahanukar, A. (2015). Secondary taste neurons that convey sweet taste and

673 starvation in the Drosophila brain. Neuron *85*, 819-832.

674 Kim, H., Kirkhart, C., and Scott, K. (2017a). Long-range projection neurons in the taste circuit of

675 Drosophila. Elife *6*.

676 Kim, J.H., Ki, Y., Lee, H., Hur, M.S., Baik, B., Hur, J.H., Nam, D., and Lim, C. (2020). The

677 voltage-gated potassium channel Shaker promotes sleep via thermosensitive GABA

678 transmission. Commun Biol *3*, 174.

679 Kim, S.S., Rouault, H., Druckmann, S., and Jayaraman, V. (2017b). Ring attractor dynamics in

680 the Drosophila central brain. Science *356*, 849-853.

681 Kirkhart, C., and Scott, K. (2015). Gustatory learning and processing in the Drosophila

682 mushroom bodies. J Neurosci *35*, 5950-5958.

683 Klapoetke, N.C., Murata, Y., Kim, S.S., Pulver, S.R., Birdsey-Benson, A., Cho, Y.K., Morimoto,

684 T.K., Chuong, A.S., Carpenter, E.J., Tian, Z.*, et al.* (2014). Independent optical excitation of

685 distinct neural populations. Nat Methods *11*, 338-346.

686 Lee, D., Seo, H., and Jung, M.W. (2012). Neural basis of reinforcement learning and decision

687 making. Annu Rev Neurosci *35*, 287-308.

688 Lewis, L.P., Siju, K.P., Aso, Y., Friedrich, A.B., Bulteel, A.J., Rubin, G.M., and Grunwald

689 Kadow, I.C. (2015). A Higher Brain Circuit for Immediate Integration of Conflicting Sensory

690 Information in Drosophila. Curr Biol *25*, 2203-2214.

Lin, S., Senapati, B., and Tsao, C.H. (2019). Neural basis of hunger-driven behaviour in

Drosophila. Open Biol *9*, 180259.

Liu, C., Meng, Z., Wiggin, T.D., Yu, J., Reed, M.L., Guo, F., Zhang, Y., Rosbash, M., and

Griffith, L.C. (2019). A Serotonin-Modulated Circuit Controls Sleep Architecture to Regulate

Cognitive Function Independent of Total Sleep in Drosophila. Curr Biol *29*, 3635-3646 e3635.

Liu, C., Placais, P.Y., Yamagata, N., Pfeiffer, B.D., Aso, Y., Friedrich, A.B., Siwanowicz, I.,

Rubin, G.M., Preat, T., and Tanimoto, H. (2012a). A subset of dopamine neurons signals reward

for odour memory in Drosophila. Nature *488*, 512-516.

Liu, Q., Liu, S., Kodama, L., Driscoll, M.R., and Wu, M.N. (2012b). Two dopaminergic neurons

signal to the dorsal fan-shaped body to promote wakefulness in Drosophila. Curr Biol *22*, 2114-

2123.

Liu, S., Liu, Q., Tabuchi, M., and Wu, M.N. (2016). Sleep Drive Is Encoded by Neural Plastic

Changes in a Dedicated Circuit. Cell *165*, 1347-1360.

Lyutova, R., Selcho, M., Pfeuffer, M., Segebarth, D., Habenstein, J., Rohwedder, A.,

Frantzmann, F., Wegener, C., Thum, A.S., and Pauls, D. (2019). Reward signaling in a recurrent

circuit of dopaminergic neurons and peptidergic Kenyon cells. Nat Commun *10*, 3097.

Masek, P., and Scott, K. (2010). Limited taste discrimination in Drosophila. Proc Natl Acad Sci

U S A *107*, 14833-14838.

Masek, P., Worden, K., Aso, Y., Rubin, G.M., and Keene, A.C. (2015). A dopamine-modulated

neural circuit regulating aversive taste memory in Drosophila. Curr Biol *25*, 1535-1541.

Mathejczyk, T.F., and Wernet, M.F. (2019). Heading choices of flying Drosophila under

changing angles of polarized light. Sci Rep *9*, 16773.

Meissner, G.W., Dorman, Z., Nern, A., Forster, K., Gibney, T., Jeter, J., Johnson, L., He, Y.,

Lee, K., Melton, B.*, et al.* (2020). An image resource of subdivided Drosophila

GAL4-driver expression patterns for neuron-level searches. bioRxiv, 2020.2005.2029.080473.

Meunier, N., Marion-Poll, F., Rospars, J.P., and Tanimura, T. (2003). Peripheral coding of bitter

taste in Drosophila. J Neurobiol *56*, 139-152.

Miyazaki, T., Lin, T.Y., Ito, K., Lee, C.H., and Stopfer, M. (2015). A gustatory second-order

neuron that connects sucrose-sensitive primary neurons and a distinct region of the gnathal

ganglion in the Drosophila brain. J Neurogenet *29*, 144-155.

Mohammad, F., Stewart, J.C., Ott, S., Chlebikova, K., Chua, J.Y., Koh, T.W., Ho, J., and

Claridge-Chang, A. (2017). Optogenetic inhibition of behavior with anion channelrhodopsins.

Nat Methods *14*, 271-274.

Nassel, D.R., and Zandawala, M. (2019). Recent advances in neuropeptide signaling in

Drosophila, from genes to physiology and behavior. Prog Neurobiol *179*, 101607.

Ni, J.Q., Liu, L.P., Binari, R., Hardy, R., Shim, H.S., Cavallaro, A., Booker, M., Pfeiffer, B.D.,

Markstein, M., Wang, H.*, et al.* (2009). A Drosophila resource of transgenic RNAi lines for

neurogenetics. Genetics *182*, 1089-1100.

Park, I.M., Meister, M.L., Huk, A.C., and Pillow, J.W. (2014). Encoding and decoding in parietal

cortex during sensorimotor decision-making. Nat Neurosci *17*, 1395-1403.

Perkins, L.A., Holderbaum, L., Tao, R., Hu, Y., Sopko, R., McCall, K., Yang-Zhou, D.,

Flockhart, I., Binari, R., Shim, H.-S.*, et al.* (2015). The Transgenic RNAi Project at Harvard

Medical School: Resources and Validation. Genetics *201*, 843-852.

Pimentel, D., Donlea, J.M., Talbot, C.B., Song, S.M., Thurston, A.J.F., and Miesenbock, G.

(2016). Operation of a homeostatic sleep switch. Nature *536*, 333-337.

Qian, Y., Cao, Y., Deng, B., Yang, G., Li, J., Xu, R., Zhang, D., Huang, J., and Rao, Y. (2017).

Sleep homeostasis regulated by 5HT2b receptor in a small subset of neurons in the dorsal fan-

shaped body of drosophila. Elife *6*.

Ramdya, P., Schneider, J., and Levine, J.D. (2017). The neurogenetics of group behavior in

Drosophila melanogaster. J Exp Biol *220*, 35-41.

Riemensperger, T., Voller, T., Stock, P., Buchner, E., and Fiala, A. (2005). Punishment

prediction by dopaminergic neurons in Drosophila. Curr Biol *15*, 1953-1960.

Rooke, R., Rasool, A., Schneider, J., and Levine, J.D. (2020). Drosophila melanogaster

behaviour changes in different social environments based on group size and density. Commun

Biol *3*, 304.

Scaplen, K.M., Talay, M., Nunez, K.M., Salamon, S., Waterman, A.G., Gang, S., Song, S.L.,

Barnea, G., and Kaun, K.R. (2020). Circuits that encode and guide alcohol-associated preference.

Elife *9*.

Scott, K. (2018). Gustatory Processing in Drosophila melanogaster. Annu Rev Entomol *63*, 15-

30.

Seelig, J.D., and Jayaraman, V. (2015). Neural dynamics for landmark orientation and angular

path integration. Nature *521*, 186-191.

Shadlen, M.N., and Kiani, R. (2013). Decision making as a window on cognition. Neuron *80*,

791-806.

Shao, L., Saver, M., Chung, P., Ren, Q., Lee, T., Kent, C.F., and Heberlein, U. (2017).

Dissection of the Drosophila neuropeptide F circuit using a high-throughput two-choice assay.

Proc Natl Acad Sci U S A *114*, E8091-E8099.

Sugrue, L.P., Corrado, G.S., and Newsome, W.T. (2005). Choosing the greater of two goods:

neural currencies for valuation and decision making. Nat Rev Neurosci *6*, 363-375.

Sun, Y., Nern, A., Franconville, R., Dana, H., Schreiter, E.R., Looger, L.L., Svoboda, K., Kim,

D.S., Hermundstad, A.M., and Jayaraman, V. (2017). Neural signatures of dynamic stimulus

selection in Drosophila. Nat Neurosci *20*, 1104-1113.

Talay, M., Richman, E.B., Snell, N.J., Hartmann, G.G., Fisher, J.D., Sorkac, A., Santoyo, J.F.,

Chou-Freed, C., Nair, N., Johnson, M.*, et al.* (2017). Transsynaptic Mapping of Second-Order

Taste Neurons in Flies by trans-Tango. Neuron *96*, 783-795 e784.

Tsao, C.H., Chen, C.C., Lin, C.H., Yang, H.Y., and Lin, S. (2018). Drosophila mushroom bodies

integrate hunger and satiety signals to control innate food-seeking behavior. Elife *7*.

Tsetsos, K., Chater, N., and Usher, M. (2012). Salience driven value integration explains

decision biases and preference reversal. Proc Natl Acad Sci U S A *109*, 9659-9664.

Turner-Evans, D., Wegener, S., Rouault, H., Franconville, R., Wolff, T., Seelig, J.D.,

Druckmann, S., and Jayaraman, V. (2017). Angular velocity integration in a fly heading circuit.

Elife *6*.

Wang, X.J. (2008). Decision making in recurrent neuronal circuits. Neuron *60*, 215-234.

Wong, R., Piper, M.D., Wertheim, B., and Partridge, L. (2009). Quantification of food intake in

Drosophila. PLoS One *4*, e6063.

Wu, Q., Wen, T., Lee, G., Park, J.H., Cai, H.N., and Shen, P. (2003). Developmental control of

foraging and social behavior by the Drosophila neuropeptide Y-like system. Neuron *39*, 147-161.

Xu, C.S., Januszewski, M., Lu, Z., Takemura, S.-y., Hayworth, K.J., Huang, G., Shinomiya, K.,

Maitin-Shepard, J., Ackerman, D., Berg, S.*, et al.* (2020). A Connectome of the Adult

Drosophila Central Brain. bioRxiv, 2020.2001.2021.911859.

Yamagata, N., Ichinose, T., Aso, Y., Placais, P.Y., Friedrich, A.B., Sima, R.J., Preat, T., Rubin,

G.M., and Tanimoto, H. (2015). Distinct dopamine neurons mediate reward signals for short-

and long-term memories. Proc Natl Acad Sci U S A *112*, 578-583.

Yang, Z., Huang, R., Fu, X., Wang, G., Qi, W., Mao, D., Shi, Z., Shen, W.L., and Wang, L.

(2018). A post-ingestive amino acid sensor promotes food consumption in Drosophila. Cell Res

28, 1013-1025.

Zandawala, M., Yurgel, M.E., Texada, M.J., Liao, S., Rewitz, K.F., Keene, A.C., and Nassel,

D.R. (2018). Modulation of Drosophila post-feeding physiology and behavior by the

neuropeptide leucokinin. PLoS Genet *14*, e1007767.

Figure 1

Figure 2

Figure 3

Figure 4

A objective

cuticle window

taste

B FBI6 (20x)

84C10-GAL4 >
UAS-GCaMP6f; UAS-tdT

C FBI6 soma

D
sweet
bittersweet

naïve deprived   naïve fed   chose sweet   chose bittersweet   chose neither

0.5 ΔR/R₀

3 s   ▲taste

E

peak ΔR/R₀

1.0
0.0
-1.0

*   ns   *   *   ns

naïve deprived (n = 10)
naïve Fed (n = 12)
chose sweet (n = 5)
chose bittersweet (n = 5)
chose neither (n = 10)

F

DA   DA   DA
Dop1R1   DopEcR   Dop1R1

AstA   DH44   Lk —Lkr→ NPF
AstA-R1 DH44-R1 Lkr

metabolic state

FBI6 ⇨ ⇨choose

G

SMP   SLP

SIP

FB

A neural signature of choice under sensory conflict in *Drosophila*

Preeti Sareen[1], Li Yan McCurdy[1,2], Michael N. Nitabach[1,3,4,*]

[1]Department of Cellular and Molecular Physiology, Yale University, New Haven, CT, USA

[2]Interdepartmental Neuroscience Program, Yale University, New Haven, CT, USA

[3]Department of Genetics, Yale University, New Haven, CT, USA

[4]Department of Neuroscience, Yale University, New Haven, CT, USA

*Lead contact: michael.nitabach@yale.edu

Table S1

Detailed statistics and sample size for data in the main figures

Figure	Datasets compared	Statistics
Fig. 1D	w^{1118} male vs. female Preference index	Mixed-effects analysis, $F(9,118)=22.46$, $p<0.0001$
		Sidak's adjusted p:
	1mM male vs. female (n=10)	0.9968
	10mM male vs. female (n=10)	>0.9999
	50mM male vs. female (n=20)	0.1784
	100mM male vs. female (n=27)	0.5552
	500mM male vs. female (n=10)	0.8233
Fig. 1E	w^{1118} Preference index, Group size (n=77)	Pearson's $r^2=0.05655$, $p=0.0373$
Fig. 1F	w^{1118} Preference index, % ate (n=77)	Pearson's $r^2=0.225$, $p<0.0001$
Fig. 1G	w^{1118} % ate, Group size (n=77)	Pearson's $r^2=0.0006$, $p=0.8313$
Fig. 1H	w^{1118} Preference index, Group size, % ate	Multiple linear regression, $F(3,73)=9.393$, $p<0.0001$ $r^2=0.278$

Fig. 2A	Optogenetic Screen 20XUAS-Chrimson (Chr) empty>Chr (n=30)	One-way ANOVA, F(40,358)=5.397, p<0.0001
		Dunnett's adjusted p:
	empty>Chr vs. Akh>Chr (n=10)	0.9996
	empty>Chr vs. AstA>Chr (n=10)	**<0.0001**
	empty>Chr vs. Crz>Chr (n=10)	0.9990
	empty>Chr vs. DH44>Chr (n=10)	0.1302
	empty>Chr vs. Lk>Chr (n=10)	0.9997
	empty>Chr vs. NPF>Chr (n=10)	**<0.0001**
	empty>Chr vs. Proctolin>Chr (n=10)	0.9997
	empty>Chr vs. sNPF>Chr (n=10)	0.9983
	empty>Chr vs. Tk>Chr (n=10)	0.9993
	empty>Chr vs. TH>Chr (n=10)	0.9983
	empty>Chr vs. PPL1 (504B)>Chr (n=10)	0.9998
	empty>Chr vs. PPL1 (65B)>Chr (n=10)	0.9996
	empty>Chr vs. PAM (58E02)>Chr (n=10)	0.9993
	empty>Chr vs. OA/TA Tdc>Chr (n=10)	>0.9999
	empty>Chr vs. Ser/Trh>Chr (n=10)	0.9997
	empty>Chr vs. γ2α'1>Chr (n=10)	**<0.0001**
	empty>Chr vs. α3>Chr (n=10)	0.8644
	empty>Chr vs. γ1-pedc>Chr (n=10)	0.9986

	empty>Chr vs. α'2α2>Chr (n=10)	0.9997
	empty>Chr vs. α'2α2,γ2α'1>Chr (n=10)	0.9995
	empty>Chr vs. α1>Chr (n=10)	**0.0070**
	empty>Chr vs. β1>Chr (n=10)	0.1241
	empty>Chr vs. β1β2>Chr (n=10)	0.9924
	empty>Chr vs. γ5>Chr (n=10)	0.9997
	empty>Chr vs. β'2a>Chr (n=6)	0.9983
	empty>Chr vs. γ4,γ4<γ1γ2>Chr (n=10)	0.9982
	empty>Chr vs. γ3>Chr (n=10)	0.9990
	empty>Chr vs. allKC 10B>Chr (n=10)	0.9988
	empty>Chr vs. α/β 8B>Chr (n=8)	0.9988
	empty>Chr vs. α/β c739>Chr (n=10)	0.9988
	empty>Chr vs. α'/β' 5B>Chr (n=10)	0.9997
	empty>Chr vs. γ-m 131B>Chr (n=8)	0.3762
	empty>Chr vs. FBl4,6 ss20>Chr (n=15)	0.9997
	empty>Chr vs. FBl3,4,6 ss208>Chr (n=10)	0.9990
	empty>Chr vs. FBl3,4,6 ss225>Chr (n=10)	0.9993
	empty>Chr vs. FBl6 c205>Chr (n=10)	0.9777
	empty>Chr vs. FBl2,8,9 R89E07>Chr (n=10)	0.6555
	empty>Chr vs. FBl5,8,9 R38E07>Chr (n=10)	0.9994
	empty>Chr vs. ventral FB R58F03>Chr (n=10)	0.3548
	empty>Chr vs. FBl1,2 R52G12>Chr (n=10)	0.9987

Fig. 2A	Optogenetic Screen 20XUAS-GtACR1 (Gt)	One-way ANOVA, $F(10,129)=7.719$, $p<0.0001$
	empty>Gt (n=30)	
		Dunnett's adjusted p:
	empty>Gt vs. AstA>Gt (n=10)	**0.0004**
	empty>Gt vs. DH44>Gt (n=10)	**0.0023**
	empty>Gt vs. Lk>Gt (n=10)	**0.0081**
	empty>Gt vs. NPF>Gt (n=10)	0.4307
	empty>Gt vs. γ2α'1>Gt (n=10)	>0.9999
	empty>Gt vs. α3>Gt1 (n=10)	**<0.0001**
	empty>Gt vs. α1>Gt (n=10)	0.9628
	empty>Gt vs. β1>Gt (n=10)	0.9996
	empty>Gt vs. FBl6 c205>Gt (n=10)	**0.0002**
	empty>Gt vs. FBl2,8,9 89E07>Gt (n=20)	0.2042
Fig. 2B	Lk (left panel)	One-way ANOVA, $F(4,84)=8.136$, $p<0.0001$
		Sidak's adjusted p:
	empty>Chr (n=10) (n=10) vs. Lk>Chr (n=20)	0.2664
	empty>Chr (n=10) vs. Lk>UAS-Chr;UAS-DH44[RNAi] (n=30)	0.3550
		0.0005
	empty>Gt (n=10) **vs. Lk>Gt** (n=19)	

Fig. 2B	Lk (right panel) RNAi ctrl = Lk-GAL4>UAS-Valium (n=20)	One-way ANOVA, $F_{(7,106)}=1.973$, p=0.0655
	RNAi ctrl vs. AstA-R1^{RNAi} (n=14) RNAi ctrl vs. DH44-R1^{RNAi} (n=10) RNAi ctrl vs. NPFR^{RNAi} (n=10) RNAi ctrl vs. Dop1R1^{RNAi} (n=20) RNAi ctrl vs. Dop1R2^{RNAi} (n=10) RNAi ctrl vs. Dop2R^{RNAi} (n=20) RNAi ctrl vs. DopEcR^{RNAi} (n=10)	Multiple comparisons not carried out since ANOVA is not significant
Fig. 2C	AstA (left panel)	One-way ANOVA, $F_{(4,75)}=61.57$, p<0.0001
		Sidak's adjusted p:
	empty>Chr (n=10) **vs. Chr** (n=20) empty>Chr (n=10) vs. AstA>UAS-Chr;UAS-AstA^{RNAi} (n=20) **empty>Gt** (n=10) **vs. AstA>Gt** (n=20)	**<0.0001** 0.9814 **<0.0001**
Fig. 2C	AstA (right panel) RNAi ctrl = AstA-GAL4>UAS-Valium (n=20)	One-way ANOVA, $F_{(7,90)}=4.368$, p=0.0003
		Dunnett's adjusted p:
	RNAi ctrl vs. DH44-R1^{RNAi} (n=10)	0.9530

	RNAi ctrl vs. Lkr^{RNAi} (n=5)	0.2010
	RNAi ctrl vs. NPFR^{RNAi} (n=13)	0.5986
	RNAi ctrl vs. Dop1R1^{RNAi} (n=20)	**0.0005**
	RNAi ctrl vs. Dop1R2^{RNAi} (n=10)	0.9998
	RNAi ctrl vs. Dop2R^{RNAi} (n=10)	0.9979
	RNAi ctrl vs. DopEcR^{RNAi} (n=10)	0.9975
Fig. 2D	NPF (left panel)	One-way ANOVA, $F_{(5,89)}=11.81$, $p<0.0001$
		Sidak's adjusted p:
	empty>Chr (n=10) **vs. NPF>Chr** (n=20)	**0.0002**
	empty>Chr (n=10) vs. NPF>UAS-Chr;UAS-NPF^{RNAi} (n=25)	0.9855
		0.1928
	empty>Gt (n=10) vs. NPF>Gt (n=20)	
Fig. 2D	NPF (right panel) RNAi ctrl = NPF-GAL4>UAS-Valium (n=20)	One-way ANOVA, $F_{(7,127)}=3.657$, $p=0.0012$
		Dunnett's adjusted p:
	RNAi ctrl vs. AstA-R1^{RNAi} (n=15)	0.4148
	RNAi ctrl vs. DH44-R1^{RNAi} (n=10)	0.9972
	RNAi ctrl vs. Lkr ^{RNAi} (n=20)	**0.0188**
	RNAi ctrl vs. Dop1R1^{RNAi} (n=20)	**0.0026**
	RNAi ctrl vs. Dop1R2^{RNAi} (n=20)	0.1588

	RNAi ctrl vs. Dop2R^{RNAi} (n=20)	0.9212
	RNAi ctrl vs. DopEcR^{RNAi} (n=10)	0.9910
Fig. 2E	DH44 (left panel)	One-way ANOVA, F(4,75)=10.54, p<0.0001
		Sidak's adjusted p:
	empty>Chr (n=10) vs. DH44>Chr (n=20)	0.1591
	empty>Chr (n=10) vs. DH44>UAS-Chr;UAS-DH44^{RNAi} (n=20)	0.9807
		<0.0001
	empty>Gt (n=10) **vs. DH44>Gt** (n=20)	
Fig. 2E	DH44 (right panel) RNAi ctrl = DH44-GAL4>UAS-Valium (n=20)	One-way ANOVA, F(7,141)=5.56, p<0.0001
		Dunnett's adjusted p:
	RNAi ctrl vs. DH44>AstA-R1^{RNAi} (n=20)	0.7806
	RNAi ctrl vs. DH44>Lkr^{RNAi} (n=20)	0.6273
	RNAi ctrl vs. DH44>NPFR^{RNAi} (n=19)	0.9997
	RNAi ctrl vs. DH44>Dop1R1^{RNAi} (n=20)	0.9998
	RNAi ctrl vs. DH44>Dop1R2^{RNAi} (n=10)	0.9996
	RNAi ctrl vs. DH44>Dop2R^{RNAi} (n=20)	0.9952
	RNAi ctrl vs. DH44>DopEcR^{RNAi} (n=20)	**0.0001**

Fig. 3A	c205 (left panel)	One-way ANOVA, F(14,318)=3.315, p<0.0001
		Sidak's adjusted p:
	empty>Chr (n=26) vs. c205>Chr (n=45)	0.2150
	empty>Gt (n=20) **vs. c205>Gt** (n=20)	**<0.0001**
	c205 (right panel) RNAi ctrl = c205-GAL4>UAS-Valium (n=47)	Kruskal-Wallis stat=40.85, p<0.0001
		Dunn's adjusted p:
	RNAi ctrl vs. AstA^{RNAi} (n=20)	0.2550
	RNAi ctrl vs. AstA-R1^{RNAi} (n=20)	**0.0131**
	RNAi ctrl vs. DH44^{RNAi} (n=20)	0.1245
	RNAi ctrl vs. DH44-R1^{RNAi} (n=20)	**0.0001**
	RNAi ctrl vs. Lk^{RNAi} (n=20)	0.9999
	RNAi ctrl vs. Lkr^{RNAi} (n=20)	**0.0011**
	RNAi ctrl vs. NPF^{RNAi} (n=20)	0.9999
	RNAi ctrl vs. NPFR^{RNAi} (n=40)	0.9999
	RNAi ctrl vs. Dop1R1^{RNAi} (n=20)	0.9999
	RNAi ctrl vs. Dop1R2^{RNAi} (n=20)	0.9999
	RNAi ctrl vs. Dop2R^{RNAi} (n=20)	0.6954
	RNAi ctrl vs. DopEcR^{RNAi} (n=31)	0.9999
Fig. 3B	c205 % ate	Kruskal-Wallis stat=49.98, p<0.0001

		Dunn's adjusted p:
	c205>Chr deprived (n=29) **vs. fed** (n=10)	**<0.0001**
	c205>Gt deprived (n=20) **vs. fed** (n=10)	**<0.0001**
Fig. 3C	Food intake empty>Chr (n=10) c205>Chr (n=8)	Kruskal-Wallis stat=3.022, p=0.6966
Fig. 3D	Food intake empty>Gt (n=7), c205>Gt (n=7)	Kruskal-Wallis stat=4.189, p=0.5225
Fig. 3E	Place PI empty>Chr (n=8), c205>Chr (n=20) empty>Gt (n=7), c205>Gt (n=8)	One-way ANOVA, $F(3,39)=2.284$, p=0.094
Fig. 4E	Peak $\Delta R/R_0$	
		Wilcoxon matched pairs p:
	naïveDeprived sweet (n=10) vs. naïveDeprived bittersweet (n=10)	**0.0244**
	naïveFed sweet (n=12) vs. naïveFed bittersweet (n=12)	0.3110
	choseSweet sweet (n=5) vs. choseSweet bittersweet (n=5)	**0.0310**
	choseBittersweet sweet (n=5) vs. choseBittersweet bittersweet (n=5)	**0.0313**
	choseNeither sweet (n=10) vs. choseNeither bittersweet (n=10)	0.50

13 **Table S2**

14 Source for all the fly genotypes used

Figure	Genotype	Source
Fig. 2, 3	empty = Empty split-GAL4	FlyLight Robot ID: 3019156
Fig. 2, 3	Chr = 20XUAS-CsChrimson (X)	RRID:BDSC_55134
Fig. 2, 3	20XUAS-CsChrimson (II) for Chr;RNAi experiments	RRID:BDSC_55136
Fig. 2, 3	Gt = 20XUAS-GtACR1 (III)	Rebecca Yang (Duke), A. Claridge-Chang (Duke-NUS)
Fig. 2	Akh-GAL4	RRID:BDSC_25684
Fig. 2	AstA-GAL4	RRID:BDSC_51979
Fig. 2	Crz-GAL4	RRID:BDSC_51976
Fig. 2	DH44-GAL4	RRID:BDSC_51987
Fig. 2	Lk-GAL4	RRID:BDSC_51993
Fig. 2	NPF-GAL4	RRID:BDSC_25682
Fig. 2	Proctolin-GAL4	RRID:BDSC_51972
Fig. 2	sNPF-GAL4	RRID:BDSC_51991
Fig. 2	Tk-GAL4	RRID:BDSC_51973
Fig. 2	TH-GAL4 (ple-GAL4)	RRID:BDSC_8848
Fig. 2	(PPL1) MB504B-GAL4	RRID:BDSC_68329
Fig. 2	(PPL1) MB065B-GAL4	RRID:BDSC_68281
Fig. 2	(PAM) 58E02-GAL4	RRID:BDSC_41347
Fig. 2	Tdc-GAL4	RRID:BDSC_9313

Fig. 2	Trh-GAL4	RRID:BDSC_38388
Fig. 2	(PPL1-γ2α'1) MB296B-GAL4 (Aso et al., 2014; Aso and Rubin, 2016)	RRID:BDSC_68308
Fig. 2	(PPL1-α3) MB630B-GAL4 (Aso and Rubin, 2016)	RRID:BDSC_68334
Fig. 2	(PPL1-γ1-pedc) MB320C-GAL4 (Aso and Rubin, 2016)	RRID:BDSC_68253
Fig. 2	(PPL1-α'2α2) MB058B-GAL4 (Aso et al., 2014)	RRID:BDSC_68278
Fig. 2	(PPL1-α'2α2, γ2α'1) MB099C-GAL4 (Aso and Rubin, 2016)	RRID:BDSC_68290
Fig. 2	(PAM-α1) MB043C-GAL4 (Aso et al., 2014; Aso and Rubin, 2016)	RRID:BDSC_68363
Fig. 2	(PAM-β1) MB063B-GAL4 (Aso et al., 2014; Aso and Rubin, 2016)	RRID:BDSC_68248
Fig. 2	(PAM-β1β2) MB213B-GAL4 (Aso et al., 2014; Aso and Rubin, 2016)	RRID:BDSC_68273
Fig. 2	(PAM-γ5) MB315C-GAL4 (Aso et al., 2014; Aso and Rubin, 2016)	RRID:BDSC_68316
Fig. 2	(PAM-β'2a) MB109B-GAL4 (Aso et al., 2014; Aso and Rubin, 2016)	RRID:BDSC_68261
Fig. 2	(PAM-γ4,γ4< γ1γ2) MB312C-GAL4 (Aso et al., 2014)	RRID:BDSC_68252

Fig. 2	(PAM-γ3) MB441B-GAL4 (Aso et al., 2014)	RRID:BDSC_68251
Fig. 2	(all KC) MB010B-GAL4 (Aso et al., 2014)	FlyLight Robot ID: 2135061
Fig. 2	(α/β KC) MB008B-GAL4 (Aso et al., 2014)	FlyLight Robot ID: 2135059
Fig. 2	(α/β KC) c739-GAL4	RRID:BDSC_7362
Fig. 2	(α'/β' KC) MB005B-GAL4 (Aso et al., 2014)	FlyLight Robot ID: 2135056
Fig. 2	(γ-m KC) MB131B-GAL4 (Aso et al., 2014)	FlyLight Robot ID: 2135179
Fig. 2	(FBl4,6) ss20-GAL4 (III)	L. Shao, U. Heberlein, FlyLight
Fig. 2	(FBl4,6) ss208-GAL4 (III)	T. Wolff, A. Jenett, G. Rubin, FlyLight
Fig. 2	(FBl4,6) ss225-GAL4 (III)	T. Wolff, A. Jenett, G. Rubin, FlyLight
Fig. 2, 3	(FBl6) c205-GAL4	RRID:BDSC_30826
Fig. 2	(FBl2,8,9) 89E07-GAL4 (Hu et al., 2018)	RRID:BDSC_40553
Fig. 2	(FBl5,8,9) 38E07-GAL4 (Hu et al., 2018)	RRID:BDSC_50007
Fig. 2	(ventral FB) 58F03-GAL4	RRID:BDSC_39187
Fig. 2	(FBl1,2) 52G12-GAL4	RRID:BDSC_49581
Fig. 2, 3	UAS-Valium	RRID:BDSC_35786

Fig. 2, 3	UAS-Lk-RNAi (Cavey et al., 2016; Lee et al., 2014; Murphy et al., 2016; Senapati et al., 2019)	RRID:BDSC_25798
Fig. 2, 3	UAS-Lkr-RNAi (Cavey et al., 2016; Senapati et al., 2019; Zandawala et al., 2018)	RRID:BDSC_25936
Fig. 2, 3	UAS-AstA-RNAi (Lee et al., 2014; Schiemann et al., 2018)	RRID:BDSC_25866
Fig. 2, 3	UAS-AstA-R1-RNAi (Yu et al., 2016)	RRID:BDSC_27280
Fig. 2, 3	UAS-NPF-RNAi (Ameku et al., 2018; Guevara et al., 2018; Lee et al., 2014)	RRID:BDSC_27237
Fig. 2, 3	UAS-NPFR-RNAi (Ameku et al., 2018; Klose and Shaw, 2019; Senapati et al., 2019; Tsao et al., 2018)	RRID:BDSC_25939
Fig. 2, 3	UAS-DH44-RNAi (Cannell et al., 2016; King et al., 2017; Lee et al., 2014)	RRID:BDSC_25804
Fig. 2, 3	UAS-DH44-R1-RNAi (King et al., 2017)	RRID:BDSC_28780

Fig. 2, 3	UAS-Dop1R1-RNAi (Klose and Shaw, 2019)	RRID:BDSC_62193
Fig. 2, 3	UAS-Dop1R2-RNAi (Klose and Shaw, 2019)	RRID:BDSC_65997
Fig. 2, 3	UAS-Dop2R-RNAi (Klose and Shaw, 2019)	RRID:BDSC_26001
Fig. 2, 3	UAS-DopEcR-RNAi (Klose and Shaw, 2019)	RRID:BDSC_31981
Fig. 4	(FBl6) 84C10-GAL4	RRID:BDSC_48378
Fig. 4	UAS-GCaMP6f;UAS-tdTomato	D. Clark, Yale University

16 **Table S3**

17 Detailed statistics and sample size for data in the supplemental figures

Figure	Datasets compared	Statistics
Fig. S3B	84C10-GAL4 Preference index	One-way ANOVA, $F_{(3,50)}=7.823$, $p=0.0002$
		Sidak's adjusted p:
	84C10>Valium (n=17) vs. 84C10>Chr (n=17)	0.1463
	84C10>Valium (n=10) vs. 84C10>Gt (n=10)	**0.0034**
Fig. S3C	84C10-GAL4 % ate	One-way ANOVA, $F_{(3,55)}=186.1$, $p<0.0001$

		Sidak's adjusted p:
	deprived 84C10>Chr (n=17) vs. fed 84C10>Chr (n=15)	**<0.0001**
		<0.0001
	deprived 84C10>Gt (n=20) vs. fed 84C10>Gt (n=7)	

Supplementary Figure Legends

Figure S1. Wild-type fly behavior. A, w1118 flies always preferred higher sucrose

concentration when no quinine was present. **B**, Food preference was sucrose concentration ratio

dependent between two food options when quinine concentration was kept constant in the

bittersweet food. **C**, Most w1118 flies ate at 21 h food deprivation, with almost 100% eating at

the equal-preference 50 mM sucrose condition. Plots depict mean ± 95% CI; violins show data

distribution.

Figure S2. Percentage of flies that ate during decision assay. A, Percent of flies that ate

during the optogenetic screen for all the genotypes tested. **B**, Only ~4% of the flies ate when Lk

neurons were activated (Lk>Chr) and this effect was abolished (~57% ate) by knocking down Lk

in the same neurons during optogenetic activation (Lk>Chr;Lk-RNAi). Plots depict mean ± 95%

CI; violins show data distribution.

Figure S3. 84C10-GAL4 characterization. A, 84C10-GAL4 showed high baseline GCaMP6f

fluorescence. Images shown are raw florescence images from the same frame without

background subtraction. **B**, 84C10-GAL4 had the same behavioral phenotype as c205-GAL4

when optogenetically activated (84C10>Chr) and inhibited (84C10>Gt) compared to controls.

Flies prefered bittersweet food compared to control flies when FBl6 neurons were inhibited

using 84C10-GAL4. **C**, Feeding was not initiated in fed flies and not inhibited in food-deprived

flies on FBl6 activation or inhibition. **D**, Neither activation nor inhibition of FBl6 was inherently

rewarding or aversive since there was no significant difference in place preference without food

for illuminated vs. non-illuminated sectors of the arena. Plots depict mean ± 95% CI; violins

show data distribution. See Table S3 for statistics and sample size. $p < 0.00001 = ****$,

$p < 0.0001 = ***$, $p < 0.01 = **$, $p < 0.05 = *$.

Figure S4. FBl6 neurons respond to water choice. A, Ratiometric calcium responses to water

stimuli, $\Delta R/R_0$, of flies with different hunger state and decision outcomes. FBl6 water responses

were also modulated by hunger and past experience. FBl6 neurons of food-deprived flies that

experienced the decision assay and chose sweet ("chose sweet") or neither ("chose neither") food

were strongly inhibited by water, while FBl6 neurons of naïve flies were not ("naïve deprived",

"naïve fed"), even though FBl6 taste responses were similar in these flies (Fig. 4d-e). FBl6

neurons of food-deprived flies that chose bittersweet were not inhibited by water ("chose

bittersweet"). Gray background area represents water application. Calcium activity trace depicts

mean $\Delta R/R_0 \pm 95\%$ CI. **B**, Peak $\Delta R/R_0$ shows significant difference between response to rejected

vs. chosen water stimuli. $p < 0.01 = **$, $p < 0.05 = *$ (see Table S1 for details on statistics and sample

size). Points on graphs depict mean ± 95% CI, with violins depicting full data distribution.

Figure S5. Inhibition of TPN3 bitter taste projection neurons. A, Acute optogenetic

(TPN3>Gt) and chronic inhibition (TPN3>TNTe) of second-order taste projection neurons

(TPN3) did not change food preference at the equal-preference condition (50 mM) or when

sucrose concentration is the same in both sweet and bittersweet options (500 mM).

Supplementary References

Ameku, T., Yoshinari, Y., Texada, M.J., Kondo, S., Amezawa, K., Yoshizaki, G., Shimada-

Niwa, Y., and Niwa, R. (2018). Midgut-derived neuropeptide F controls germline stem cell

proliferation in a mating-dependent manner. PLoS Biol *16*, e2005004.

Aso, Y., Hattori, D., Yu, Y., Johnston, R.M., Iyer, N.A., Ngo, T.T., Dionne, H., Abbott, L.F.,

Axel, R., Tanimoto, H.*, et al.* (2014). The neuronal architecture of the mushroom body provides

a logic for associative learning. Elife *3*, e04577.

Aso, Y., and Rubin, G.M. (2016). Dopaminergic neurons write and update memories with cell-

type-specific rules. Elife *5*.

Cannell, E., Dornan, A.J., Halberg, K.A., Terhzaz, S., Dow, J.A.T., and Davies, S.A. (2016). The

corticotropin-releasing factor-like diuretic hormone 44 (DH44) and kinin neuropeptides

modulate desiccation and starvation tolerance in Drosophila melanogaster. Peptides *80*, 96-107.

Cavey, M., Collins, B., Bertet, C., and Blau, J. (2016). Circadian rhythms in neuronal activity

propagate through output circuits. Nat Neurosci *19*, 587-595.

Guevara, A., Gates, H., Urbina, B., and French, R. (2018). Developmental Ethanol Exposure

Causes Reduced Feeding and Reveals a Critical Role for Neuropeptide F in Survival. Front

Physiol *9*, 237.

Hu, W., Peng, Y., Sun, J., Zhang, F., Zhang, X., Wang, L., Li, Q., and Zhong, Y. (2018). Fan-

Shaped Body Neurons in the Drosophila Brain Regulate Both Innate and Conditioned

Nociceptive Avoidance. Cell Rep *24*, 1573-1584.

King, A.N., Barber, A.F., Smith, A.E., Dreyer, A.P., Sitaraman, D., Nitabach, M.N., Cavanaugh,

D.J., and Sehgal, A. (2017). A Peptidergic Circuit Links the Circadian Clock to Locomotor

Activity. Curr Biol *27*, 1915-1927 e1915.

Klose, M., and Shaw, P. (2019). Sleep-drive reprograms clock neuronal identity through CREB-

binding protein induced PDFR expression. bioRxiv, 2019.2012.2012.874628.

Lee, J.H., Bassel-Duby, R., and Olson, E.N. (2014). Heart- and muscle-derived signaling system

dependent on MED13 and Wingless controls obesity in Drosophila. Proc Natl Acad Sci U S A

111, 9491-9496.

Murphy, K.R., Deshpande, S.A., Yurgel, M.E., Quinn, J.P., Weissbach, J.L., Keene, A.C.,

Dawson-Scully, K., Huber, R., Tomchik, S.M., and Ja, W.W. (2016). Postprandial sleep

mechanics in Drosophila. Elife *5*.

Schiemann, R., Lammers, K., Janz, M., Lohmann, J., Paululat, A., and Meyer, H. (2018).

Identification and In Vivo Characterisation of Cardioactive Peptides in Drosophila melanogaster.

Int J Mol Sci *20*.

Senapati, B., Tsao, C.H., Juan, Y.A., Chiu, T.H., Wu, C.L., Waddell, S., and Lin, S. (2019). A

neural mechanism for deprivation state-specific expression of relevant memories in Drosophila.

Nat Neurosci *22*, 2029-2039.

Tsao, C.H., Chen, C.C., Lin, C.H., Yang, H.Y., and Lin, S. (2018). Drosophila mushroom bodies

integrate hunger and satiety signals to control innate food-seeking behavior. Elife *7*.

Yu, Y., Huang, R., Ye, J., Zhang, V., Wu, C., Cheng, G., Jia, J., and Wang, L. (2016).

Regulation of starvation-induced hyperactivity by insulin and glucagon signaling in adult

Drosophila. Elife *5*.

Zandawala, M., Yurgel, M.E., Texada, M.J., Liao, S., Rewitz, K.F., Keene, A.C., and Nassel,

D.R. (2018). Modulation of Drosophila post-feeding physiology and behavior by the

neuropeptide leucokinin. PLoS Genet *14*, e1007767.

Figure S1

A w1118 (21h)

B w1118 (21h)

C w1118 (21h)

Figure S2

Figure S3

A

84C10-GAL4>UAS-GCaMP6f

84C10-GAL4>UAS-tdT

B

Preference index

84C10>Valium   84C10>Chr   84C10>Valium   84C10>Gt

**

C

% ate

**** ****

84C10> Chr   Gt

○ food deprived
● fed

D

Place PI

1.0

0.0

-1.0

84C10>Chr   84C10>Gt

Figure S4

A

| naïve deprived | naïve fed | chose sweet | chose bittersweet | chose neither |

— response to water

B

Figure S5

A

TPN3 (R11H09-GAL4)

sucrose (mM): 50 500   50 500

Preference index

500 mM sucrose + 1 mM quinine