The yeast ISW1b ATP-dependent chromatin remodeler is critical for nucleosome spacing and dinucleosome resolution

Peter R. Eriksson and David J. Clark [1]

Division of Developmental Biology, *Eunice Kennedy Shriver* National Institute of Child Health and Human Development, National Institutes of Health, Bethesda, Maryland 20892, USA

[1] **For correspondence:** clarkda@mail.nih.gov. Tel. 301 496-6966.
Building 6A Room 2A02, 6 Center Drive, Bethesda MD 20892.

Competing interests: The authors declare that no competing interests exist.

Abstract

Isw1 and Chd1 are ATP-dependent nucleosome-spacing enzymes required to establish regular arrays of phased nucleosomes near transcription start sites of yeast genes. Cells lacking both Isw1 and Chd1 have extremely disrupted chromatin, with weak phasing, irregular spacing and a propensity to form closed-packed dinucleosomes. The Isw1 ATPase subunit occurs in two different remodeling complexes: ISW1a (composed of Isw1 and Ioc3) and ISW1b (Isw1, Ioc2 and Ioc4). We show that ISW1b is primarily responsible for setting nucleosome spacing and resolving some close-packed dinucleosomes, whereas ISW1a plays only a minor role. Chd1 also separates some dinucleosomes, but not all. Dinucleosome accumulation correlates with RNA polymerase II levels, except for very highly transcribed genes, which are partially depleted of dinucleosomes and mononucleosomes. Loss of the Set2 H3-K36 methyltransferase partly phenocopies loss of Ioc4, whereas loss of the Set1 H3-K4 methyltransferase has no global effect. In conclusion, ISW1b is critical for yeast chromatin organization.

Introduction

Eukaryotic DNA is packaged into the nucleus in the form of chromatin. The structural subunit of chromatin is the nucleosome, which is composed of an octamer of core histones (two molecules each of H3, H4, H2A and H2B), around which is wrapped ~146 bp of DNA in ~1.7 superhelical turns (**Luger et al., 1997**). Nucleosomes are regularly spaced along the DNA, like beads on a string, forming a fibre which spontaneously folds into higher-order chromatin structures (**Thoma et al., 1979**). Nucleosomes restrict access to DNA and are potent inhibitors of transcription and other DNA-dependent processes in vitro. Cells regulate access to their DNA in part by deploying ATP-dependent chromatin remodeling complexes that are capable of overcoming the nucleosome, either by removing it from the DNA or by sliding it along the DNA (**Chereji and Clark, 2018; Clapier et al., 2017; Paul and Bartholomew, 2018; Prajapati et al., 2020**).

The ISWI and CHD enzymes represent a major class of ATP-dependent chromatin remodelers conserved from yeast to mammals. They are primarily nucleosome sliding enzymes; many have nucleosome spacing activity in vitro (**Ito et al., 1997; Lieleg et al., 2015; Lusser et al., 2005; Pointner et al., 2012; Stockdale et al., 2006; Torigoe et al., 2013; Tsukiyama et al., 1999**). In vivo, ISWI enzymes are important for chromatin organization near promoters and other gene regulatory elements in vivo (**Barisic et al., 2019; Gkikopoulos et al., 2011; Ocampo et al., 2016; Wiechens et al., 2016**). ISWI complexes have additional functions in chromatin assembly (**Ito et al., 1997; Lusser et al., 2005; Vary et al., 2004**), stress-induced gene repression (**Lindstrom et al., 2006; Pinskaya et al., 2009**), transcript termination (**Morillon et al., 2003; Ocampo et al., 2019**) and quality control of mRNP biogenesis (**Babour et al., 2016**).

The budding yeast (*Saccharomyces cerevisiae*) possesses at least four ATP-dependent chromatin remodeling complexes capable of spacing nucleosomes in vitro: ISW1, ISW2, Chd1 and INO80 (**Krietenstein et al., 2016; Tsukiyama et al., 1999; Udugama et al., 2011; Vary et al., 2003**). In vivo, global chromatin organization in cells lacking Isw2 is very similar to wild type (**Gkikopoulos et al., 2011; Ocampo et al., 2016**), suggesting that ISW2 activity is more local than global. In contrast, cells lacking Ino80 (**Qiu et al., 2020; van Bakel et al., 2013; Yen et al., 2012**) or Isw1 (**Ocampo et al., 2016; Tirosh et al., 2010**) have shorter average nucleosome spacing than wild type cells. Cells lacking Chd1 have slightly shorter spacing and relatively poor nucleosome phasing (**Gkikopoulos et al., 2011; Ocampo et al., 2016**). Most impressively, cells lacking both Isw1 and Chd1 have extremely disrupted chromatin, indicating that both enzymes are required for proper chromatin organization (**Gkikopoulos et al., 2011; Ocampo et al., 2016**). Recently, we showed that an important contributory factor to chromatin disruption in the *chd1Δ isw1Δ* double mutant is a tendency for nucleosomes at the 5'-ends of yeast genes to be packed close together, resulting in dinucleosomes with little or no linker DNA (**Ocampo et al., 2019**). This observation suggests that Isw1 and/or Chd1 are critical for resolving dinucleosomes.

The yeast Isw1 ATPase subunit is found in two different complexes, termed ISW1a (containing Isw1 and Ioc3) and ISW1b (containing Isw1, Ioc2 and Ioc4) (**Mellor and Morillon, 2004; Vary et al., 2003**). There is genetic evidence for antagonistic interactions between ISW1a and ISW1b, suggesting that ISW1a has a negative role in transcription that is suppressed by ISW1b (**Lafon et al., 2012**). The Ioc subunits appear to have regulatory functions, since the isolated Isw1 subunit is inactive in vitro (**Vary et**

al., 2003), unless the AutoN inhibitory domain is mutated (*Yan et al., 2019*). The ISW1a complex is a
potent nucleosome spacing enzyme in vitro (*Gangaraju and Bartholomew, 2007; Richmond, 2012; Vary*
et al., 2003) and its structure has been determined, suggesting a mechanism involving separation of two
nucleosomes using a protein ruler (*Yamada et al., 2011*). The ISW1b complex is a weaker nucleosome
spacing enzyme than ISW1a in vitro (*Vary et al., 2003*). The Ioc2 subunit of ISW1b contains a PHD finger
(*Smolle et al., 2012*), which may bind H3-K4me3, suggesting that Isw1 binding to chromatin is mediated
by H3-K4 trimethylation (*Santos-Rosa et al., 2003*). The Ioc4 subunit of ISW1b has a PWWP domain
which binds to H3-K36me3, a histone modification associated with active transcription (*Maltby et al.,*
2012; Smolle et al., 2012). These observations suggest that H3-K4 and H3-K36 trimethylation may play a
critical role in ISW1b function.
Although ISW1a and ISW1b have been studied in depth in vitro, relatively little is known about
their contributions to ISW1 function in vivo. Here, we have assessed the contributions of ISW1a and
ISW1b to nucleosome spacing and separation of dinucleosomes. We find that ISW1b is the major
spacing enzyme, whereas ISW1a plays a very minor role in global genic chromatin organization that is
revealed only in the absence of ISW1b. ISW1b, together with Chd1, is required to resolve
dinucleosomes, whereas ISW1a makes little contribution.
89
90
91   **Results**
92
The ISW1b complex is the major nucleosome spacing enzyme in yeast
We have used MNase-seq to determine the chromatin organization at yeast genes. Briefly, this
technique involves micrococcal nuclease (MNase) digestion of the chromatin in purified nuclei to
predominantly mononucleosomes. Suitably digested DNA samples are then used to construct libraries
for paired-end sequencing. Alignment to the yeast genome indicates the location of each sequenced
nucleosome. The midpoint of each sequence is assumed to represent the midpoint of the nucleosome
(also called the dyad). An average plot for all 5770 yeast genes is obtained by aligning all genes on their
transcription start site (TSS) or on their major +1 nucleosome position, summing all of the nucleosome
dyads on every gene, and normalizing to the genomic average (set at 1). A typical plot for wild type cells
is shown in *Figure 1A*. We define nucleosome spacing as the average distance between the dyad peaks
for the +1 to +5 nucleosomes (i.e. the first five nucleosomes on the average gene), measured by the
slope of the regression line. However, in some mutants, the phasing is so poor that the spacing cannot
be measured accurately (see below). The degree of phasing is indicated by the height of the nucleosome
peaks: higher peaks indicate more coincident dyads and therefore better nucleosome positioning.
We have shown previously (*Ocampo et al., 2016*) that cells lacking Isw1 have weaker
nucleosome phasing and reduced spacing relative to wild type cells (*Figure 1B*). However, *isw1Δ* cells
lack both the ISW1a and the ISW1b remodeling complexes. To assess the relative contributions of ISW1a
and ISW1b to phasing and spacing, we examined the chromatin organization in various *ioc* mutants.
Cells lacking the ISW1a complex (*ioc3Δ*) show a slight increase in spacing (replicates: 166, 167 bp) with
no change in phasing relative to wild type (165, 165 bp) (*Figure 1C*). In contrast, cells lacking an ISW1b
subunit (*ioc2Δ* or *ioc4Δ*) have intermediate nucleosome spacing (161-163 bp) (*Figure 1D, E*): less than in

wild type (165 bp), but not as low as in *isw1Δ* cells (159, 160 bp). The spacing differences are mostly

small, but they are clear, because the nucleosome shift increases with the nucleosome number

(compare biological replicate experiments; ***Figure 1***). Phasing is weaker in the *ioc2Δ* mutant (***Figure 1D***),

but hardly affected in the *ioc4Δ* mutant (***Figure 1E***). Cells lacking both ISW1b ancillary subunits (the

ioc2Δ ioc4Δ double mutant) have similar chromatin organization to the *ioc2Δ* single mutant, exhibiting

the weaker phasing observed in *ioc2Δ* cells as well as the shorter spacing observed in both the *ioc2Δ* and

ioc4Δ single mutants (***Figure 1F***). The similar spacing in the *ioc2Δ* and *ioc4Δ* single mutants and the *ioc2Δ*

ioc4Δ double mutant (~162 bp) suggests that both Ioc2 and Ioc4 are required for ISW1b spacing activity.

Figure 1. The ISW1b complex (Isw1-Ioc2-Ioc4) is required for setting wild type nucleosome spacing, whereas the ISW1a complex (Isw1-Ioc3) sets intermediate spacing in the absence of ISW1b. Average nucleosome dyad density plots for all genes: (**A**) wild type, (**B**) *isw1Δ*, (**C**) *ioc3Δ*, (**D**) *ioc2Δ*, (**E**) *ioc4Δ*, (**F**) *ioc2Δ ioc4Δ*, (**G**) *ioc2Δ ioc3Δ ioc4Δ*. All yeast genes were aligned on the midpoints of their +1 nucleosomes. The dyad distribution was normalized to the global average (set at 1). For ease of comparison, wild type (WT) replicate A is shown as a black line with grey fill in all plots. Two biological replicate experiments (A and B) are shown for each strain in all of the plots (A: red line; B: blue line). The average spacing in bp is shown for replicates A and B in the bottom right corner (measured by regression analysis of the first 5 nucleosome peaks, beginning with the +1 nucleosome).

Chromatin organization in cells lacking all three ancillary subunits (the *ioc2Δ ioc3Δ ioc4Δ* triple
mutant) is very similar to that in *isw1Δ* cells (**Figure 1G**). This is expected, because both complexes
should be inactive in both mutants, given that the Isw1 subunit by itself has no remodeling activity in
vitro (**Vary et al., 2003**), but it also implies that ISW1a affects spacing since the triple mutant is not the
same as the *ioc2Δ ioc4Δ* double mutant. The shorter spacing in the *ioc2Δ ioc3Δ ioc4Δ* triple mutant is
suggests that ISW1a is responsible for the intermediate spacing observed in the absence of ISW1b. Since
the loss of Ioc3 has only a small effect on global spacing in the presence of ISW1b (the *ioc3Δ* single
mutant), ISW1b is more important for chromatin organization than ISW1a. That is, the contribution of
ISW1a is only clear at the global level if ISW1b is absent. We conclude that the ISW1b complex is
primarily responsible for wild type nucleosome spacing and that the ISW1a complex has a minor
underlying role, mediating intermediate spacing.
134
ISW1a and ISW1b activities are not restricted to target genes
Chromatin immunoprecipitation (ChIP-seq) experiments have shown that ISW1a (Ioc3), ISW1b (Ioc4)
and Isw1 are enriched on different sets of genes (**Yen et al., 2012**). We determined whether the
chromatin organization of these sets of target genes is differentially affected by loss of ISW1a or ISW1b,
as might be expected. In the case of ISW1a, we found that Ioc3-target genes show very slightly increased
spacing (166 vs. 165 bp) in wild type cells and in *ioc3Δ* cells (167 vs. 166 bp), but these differences are
probably negligible (**Figure S1A**). Thus, ISW1a has little or no differential effect on the chromatin
organization of its target genes. Moreover, the spacing on Ioc3-bound genes is affected in *ioc4Δ* cells,
indicating that ISW1a target genes are affected by ISW1b (Ioc4) (**Figure S1A**). In the case of ISW1b, Ioc4-
target and non-target genes have the same spacing and phasing in wild type cells, suggesting that they
are not differentially affected by ISW1b. In *ioc4Δ* cells, both sets of genes have shorter spacing than wild
type (**Figure S1B**). Ioc4-target genes may have slightly shorter spacing than non-target genes, but the
effect is subtle (**Figure S1B**). As expected, Ioc4-bound genes are not affected in *ioc3Δ* cells (**Figure S1B**).
Thus, ISW1b affects the chromatin organization of both target genes and non-target genes. Finally, Isw1-
target genes have the same spacing as non-target genes in wild type cells and, although the phasing is
slightly better on the Isw1-target genes, this effect is also observed in the absence of Isw1 (**Figure S1C**).
Isw1-target and non-target genes show similar changes in *isw1Δ* cells (short spacing and weaker
phasing), although there is a slight difference in spacing of 1-2 bp between target and non-target genes
(**Figure S1C**). Overall, the Isw1 complexes affect the chromatin organization of both target and non-
target genes similarly, with only subtle differences at most. It is unclear why we do not observe obvious
differences in chromatin organization for target and non-target genes defined by Ioc3, Ioc4 or Isw1
binding; perhaps all genes have some bound ISW1b, such that designated target genes are only
modestly enriched relative to non-target genes.
158
ISW1b and Chd1 account for the extreme chromatin disruption in cells lacking both Isw1 and Chd1
Cells lacking Chd1 have a mild chromatin organization defect (**Gkikopoulos et al., 2011; Ocampo et al.,**
2016), characterized by somewhat shorter spacing than wild type (164, 163 bp vs. 165, 165 bp) and
weaker phasing (**Figure 2A**). In contrast, cells lacking both Chd1 and Isw1 (the *chd1Δ isw1Δ* double
mutant) have extremely disrupted chromatin organization (**Gkikopoulos et al., 2011; Ocampo et al.,**
2016) (**Figure 2B**). We observed that the *chd1Δ ioc2Δ ioc4Δ* triple mutant (**Figure 2C**) also has extremely

disrupted chromatin, whereas the *chd1Δ ioc3Δ* double mutant is very similar to the *chd1Δ* single mutant
(**Figure 2D**). This observation is consistent with our conclusion that the ISW1b complex is much more
important than the ISW1a complex for global nucleosome spacing (**Figure 1**). However, chromatin
organization in the *chd1Δ ioc2Δ ioc4Δ* triple mutant is not quite as severely disrupted as in the *chd1Δ*
isw1Δ double mutant: the phasing in the triple mutant is very poor, but somewhat stronger than in the
double mutant (**Figure 2B, C**). Measurement of the spacing in these two mutants is not appropriate
because most of the peaks are too weak to measure accurately. The difference between the *chd1Δ*
ioc2Δ ioc4Δ triple mutant and the *chd1Δ isw1Δ* double mutant is that the ISW1a complex (Isw1-Ioc3) is
still present in the former mutant. This suggests that ISW1a contributes some residual order to genic
chromatin in the absence of both Chd1 and ISW1b.

Figure 2. ISW1b and Chd1 are the major remodelers required for normal chromatin organization. Average nucleosome dyad density plots for all genes: (**A**) *chd1Δ*, (**B**) *chd1Δ isw1Δ*, (**C**) *chd1Δ ioc2Δ ioc4Δ*, (**D**) *chd1Δ ioc3Δ*. All yeast genes were aligned on the midpoints of their +1 nucleosomes. The dyad distribution was normalized to the global average (set at 1). For ease of comparison, wild type replicate A is shown as a black line with grey fill in all plots. Two biological replicate experiments (A and B) are shown for each strain in all of the plots (A: red line; B: blue line). The average spacing in bp is shown for replicates A and B in the bottom right corner (measured by regression analysis of the first 5 nucleosome peaks, beginning with the +1 nucleosome); ND = not determined because the phasing is too weak for accurate measurement.

175
176

ISW1a and ISW1b have little effect on the chromatin of very active genes
Since ISW1b associates with transcriptionally active genes (**Maltby et al., 2012**), we examined whether
ISW1a or ISW1b mediate specific effects at highly active genes. We compared the average chromatin
structure of the 300 most active genes, defined by ChIP-seq for the Rpb3 subunit of Pol II (**Ocampo et**
al., 2016), with that of the remaining, less active, 5470 genes (**Figure S2**). In wild type cells, the
chromatin of the highly active genes is poorly organized, characterized by much reduced and irregular
spacing, weak phasing and a much wider nucleosome-depleted region (NDR) at the promoter, which
extends upstream (**Cole et al., 2014; Weiner et al., 2010**). Although the average spacing on the highly
active genes is difficult to measure accurately due to poor phasing, it is clearly shorter than the spacing

on the other genes (**Figure S2A**). The chromatin organization of the highly active genes is not strongly
differentially affected by any of the *iocΔ* mutations or *isw1Δ* (**Figure S2A**), whereas the other genes have
the spacing observed for all genes, as expected. We conclude that ISW1a and ISW1b have no differential
effect on the chromatin organization of highly active genes, with the caveat that their general state of
disruption might obscure subtle effects. On the other hand, nucleosome phasing on the top 300 active
genes is weaker in cells lacking Chd1 than in wild type or in any of the *ioc* mutants (**Figure S2B**),
suggesting that Chd1 is the most important spacing enzyme for highly active genes, and consistent with
a direct association of Chd1 with transcript elongation factors (**Simic et al., 2003**).
194
Both ISW1b and Chd1 are important for resolution of close-packed dinucleosomes
We reported previously that a major contributing factor to chromatin disruption in the *chd1Δ isw1Δ*
double mutant is the presence of close-packed dinucleosomes, primarily involving the +2 nucleosome
(i.e. dinucleosomes containing either the +1 and +2 nucleosomes or the +2 and +3 nucleosomes)
(**Ocampo et al., 2019**) (**Figure 3A**). These dinucleosomes are characterized by MNase-resistant DNA
fragments of 250 - 350 bp, presumably representing two nucleosomes (or perhaps sub-nucleosomes)
with little or no intervening linker DNA for MNase to cut. The dyad plots shown above (**Figures 1 and 2**)
include only mononucleosome data because the analysis was limited to DNA fragments of 120 - 180 bp.
A dyad plot is not appropriate for dinucleosomes, since the midpoint of a dinucleosome would be
located between the two nucleosomes. Instead, we used nucleosome occupancy (coverage) plots, in
which the number of times each genomic base pair appears in either mononucleosomes or
dinucleosomes is counted and normalized to the genomic average. In the *chd1Δ isw1Δ* double mutant,
there is a strong dinucleosome peak at the +2 position and a depressed +2 mononucleosome peak,
consistent with the presence of a significant fraction of all +2 nucleosomes in dinucleosomes (Ocampo
et al., 2019) (**Figure 3A; Figure S3**). We determined the relative contributions of Chd1 and Isw1 to
dinucleosome resolution. A dinucleosome peak is observed in both the *chd1Δ* and *isw1Δ* single mutants,
but it is weaker than in the *chd1Δ isw1Δ* double mutant (**Figure 3A**). The +2 mononucleosome peak is
also weaker in the single mutants relative to wild type. These data indicate that the Chd1 and Isw1
remodeling enzymes both contribute independently and make additive contributions to dinucleosome
resolution, such that neither enzyme is capable of resolving all dinucleosomes on its own.
215   Next, we determined the separate contributions of the ISW1a and ISW1b complexes to Isw1-
dependent dinucleosome resolution (**Figure 3B**). The *ioc2Δ* and *ioc4Δ* single mutants have an enhanced
dinucleosome peak and depressed mononucleosome peak, as observed for the *isw1Δ* single mutant,
suggesting that both ancillary subunits of the ISW1b complex are required for Isw1-dependent
dinucleosome resolution. Consistent with this conclusion, the *ioc2Δ ioc4Δ* double mutant has a
dinucleosome peak similar to that in the single mutants. In contrast, there are fewer dinucleosomes in
the *ioc3Δ* mutant, which has a slightly lower +2 dinucleosome peak than wild type and somewhat higher
levels of the +2 and +3 mononucleosomes (**Figure 3B; Figure S3**), indicating that ISW1a is not important
for dinucleosome resolution. These conclusions are supported by high levels of dinucleosomes in the
chd1Δ ioc2Δ ioc4Δ triple mutant versus wild type, resulting from loss of both ISW1b and Chd1 (**Figure
3C**). Similarly, the *chd1Δ ioc3Δ* double mutant has more dinucleosomes than wild type, but less than the
chd1Δ ioc2Δ ioc4Δ triple mutant, which can be accounted for by the absence of Chd1 with little

contribution from ISW1a (***Figure 3C***). Thus, the ISW1b complex accounts quite well for the role of Isw1
in dinucleosome resolution.
229

Figure 3. Both ISW1b and Chd1 are important for separating close-packed dinucleosomes. Average nucleosome occupancy (coverage) plots for all genes. All yeast genes were aligned on the midpoint of their average +1 nucleosome position. The occupancy was normalized to the global average (set at 1) for mononucleosomes (120 - 180 bp) or dinucleosomes (250 - 350 bp). Note different y-axis scales are used in different plots to separate the lines more clearly. Data for replicate A are shown (see ***Figure S3*** for comparison of biological replicate experiments). (**A**) Dinucleosomes involving the +2 nucleosome are enriched whereas +2 mononucleosomes are depleted in *chd1Δ isw1Δ* cells; smaller effects are observed in *chd1Δ* and *isw1Δ* cells. (**B**) ISW1b (Isw1-Ioc2-Ioc4) accounts for the increased level of dinucleosomes in *isw1Δ* cells. (**C**) ISW1b and Chd1 are both required to resolve all dinucleosomes.

230
231
Dinucleosome formation correlates with RNA polymerase II levels
Since highly active genes have much reduced nucleosome spacing (***Weiner et al., 2010***) (***Figure S2***), we
wondered whether transcription correlates with increased levels of dinucleosomes. We constructed
heat maps in which nucleosome occupancy data for all 5770 genes are aligned on the average +1
nucleosome position in wild type cells (set at 0 on the x-axis) and sorted by their Pol II density, as
measured by ChIP-seq for the Rpb3 subunit (Ocampo et al., 2016) (***Figure 4; Figure S4***). In wild type
cells, the +1 nucleosome is depleted in the dinucleosome heat map and well represented in the
mononucleosome heat map, indicating that there are relatively few +1/+2 dinucleosomes on all genes,
independent of Pol II level. However, in wild type cells, the prevalence of +2/+3 dinucleosomes does
correlate with gene activity, except for some of the top 300 highly active genes (***Figure 4A***; above the
horizontal dashed line). The latter are depleted of +2/+3 dinucleosomes, but somewhat enriched in
+4/+5 dinucleosomes instead, compared to other genes in the top 300 (***Figure 4A***). Mononucleosome
occupancy shows the opposite trend to dinucleosome occupancy: nucleosome occupancy on the 300

most active genes is relatively low, which we attribute to some loss of mononucleosomes to
dinucleosome formation and to some nucleosome loss associated with transcription (**Cole et al., 2014;**
Cui et al., 2012; Schwabish and Struhl, 2007, 2004). On all of the other genes, mononucleosome
occupancy increases as Pol II density decreases, correlating with decreasing dinucleosome levels (**Figure**
4A; below dashed line). The least active genes, at the bottom of the heat map, clearly have fewer
dinucleosomes and more mononucleosomes than the more active genes near the top of the heat map.

251
252   **Figure 4**. Dinucleosome formation correlates with RNA polymerase II levels, except for the most highly transcribed
253   genes. Heat map analysis of nucleosome occupancy (coverage) for all 5770 genes aligned on the midpoint of their
254   average +1 nucleosome position and sorted by Pol II density on the coding region (CDS), using ChIP-seq data for
       the Rpb3 subunit (**Ocampo et al., 2016**). Data for replicate A are shown (see **Figure S4** for replicate B). The dashed
       line separates the top 300 most active genes from the rest. The scale bar is the same for all plots, ranging from
       blue (low nucleosome occupancy) to red (high occupancy). (**A**) wild type, (**B**) isw1Δ, (**C**) chd1Δ isw1Δ, (**D**) ioc2Δ,
       (**E**) ioc3Δ, (**F**) ioc4Δ, (**G**) ioc2Δ ioc4Δ, (**H**) ioc2Δ ioc3Δ ioc4Δ, (**I**) chd1Δ, (**J**) chd1Δ ioc3Δ, (**K**) chd1Δ ioc2Δ ioc4Δ.

The trends observed in wild type cells are amplified in the *isw1Δ* single mutant (***Figure 4B***) and
very strongly amplified in the *chd1Δ isw1Δ* double mutant (***Figure 4C***); both have higher levels of
dinucleosomes than wild type (***Figure 3A***). In these two mutants, dinucleosomes involving the +2
nucleosome are much more prevalent relative to wild type, particularly in the bottom half of the heat
map where there is a greater increase in dinucleosomes relative to wild type; other dinucleosomes,
farther downstream, are much less prevalent relative to wild type (***Figure 4A-C***). In addition, in the
chd1Δ isw1Δ double mutant, the top 300 active genes are depleted of +4/+5 dinucleosomes. The least
active genes, right at the bottom of each heat map, remain low in dinucleosomes in both mutants
(***Figure 4A-C***).
The *ioc2Δ* single mutant, the *ioc2Δ ioc4Δ* double mutant and the *ioc2Δ ioc3Δ ioc4Δ* triple mutant
are all very similar to the *isw1Δ* mutant, whereas the *ioc4Δ* single mutant has higher levels of
downstream dinucleosomes than the other mutants and is more similar to wild type in this regard
(***Figure 4D-H***). In contrast, the *ioc3Δ* single mutant is similar to wild type (***Figure 4E***). This analysis
confirms that ISW1b is much more important than ISW1a for dinucleosome resolution. The *chd1Δ* single
mutant (***Figure 4I***) and the *chd1Δ ioc3Δ* double mutant (***Figure 4J***) are both quite similar to the *isw1Δ*
single mutant (***Figure 4B***). The *chd1Δ ioc2Δ ioc4Δ* triple mutant (***Figure 4K***) is very similar to the *chd1Δ*
isw1Δ double mutant (***Figure 4C***), again consistent with important roles for both Chd1 and ISW1b in
resolving dinucleosomes.
Overall, our data suggest that transcription is associated with higher levels of dinucleosomes,
perhaps formed by Pol II as it transcribes. The apparent anomaly of reduced dinucleosome prevalence
on the most active genes may be a consequence of transcription-associated nucleosome loss, which
strongly affects only the most active genes (***Cole et al., 2014; Schwabish and Struhl, 2007, 2004***).
277
Loss of Set2 has a subtle effect on chromatin organization
The Ioc4 PWWP domain binds preferentially to H3-K36me3 relative to unmethylated H3-K36 in vitro,
suggesting that ISW1b may be regulated by H3-K36me3 (***Maltby et al., 2012; Smolle et al., 2012***). The
only H3-K36 methyltransferase in yeast is encoded by *SET2*. Accordingly, we examined whether ISW1b-
mediated changes in nucleosome spacing and dinucleosome resolution depend on Set2 and H3-K36me3.
Nucleosome spacing in a *set2Δ* mutant (replicates: 165 and 163 bp) is slightly lower than wild type (165
and 165 bp), although this difference may not be significant (***Figure 5A***). Ioc4 has a clearer effect on
chromatin organization (***Figure 1E***; *ioc4Δ* replicates: 163 and 162 bp) than loss of Set2 (***Figure 5B***). Cells
lacking Set2 or Ioc4 show a similar correlation between dinucleosome levels and transcriptional activity
and show a somewhat higher prevalence of dinucleosomes than wild type (***Figure 5C and Figure S3***). We
conclude that loss of H3-K36me3 has some effect on ISW1b activity but does not fully nullify it, since
chromatin organization in the *set2Δ* mutant is not as defective as in the *ioc2Δ* or *ioc2Δ ioc4Δ* mutants
(***Figure 1D, F***).

291
292
Figure 5. Set2 has a subtle effect on chromatin organization whereas Set1 has no effect. (**A**) Average
nucleosome dyad density plot for all genes in *set2Δ* cells. Wild type replicate A is shown as a black line with
grey fill. Two biological replicate experiments are shown: A (red line) B (blue line). The average spacing (bp)
for replicates A and B are given (bottom right). (**B**) Comparison of chromatin organization in *set2Δ* cells
(blue line) and *ioc4Δ* cells (red line) (replicates A). (**C**) Heat map analysis of nucleosome occupancy
(coverage) in *set2Δ* cells for all genes aligned on the midpoint of the +1 nucleosome position and sorted by
Pol II density on the coding region (CDS). Replicates A are shown (see ***Figure S4*** for replicate B). Dashed line
separates the top 300 most active genes from the rest. The scale bar ranges from blue (low nucleosome
occupancy) to red (high). Equivalent analyses for cells lacking Set1: (**D**) Average nucleosome dyad density
plot for all genes in *set1Δ* cells. (**E**) Chromatin organization in *set1Δ* (blue line) and *ioc2Δ* cells (red line)
(replicates A). (**F**) Heat maps of dinucleosome and mononucleosome occupancies in *set1Δ* cells.

304
Loss of Set1 has no effect on global chromatin organization
The binding of ISW1 to chromatin is mediated through H3-K4 trimethylation by Set1, the only H3-K4
methyltransferase in yeast (***Santos-Rosa et al., 2003***). Since Ioc2 contains a PHD domain (***Smolle et al.,***
2012), it might mediate the binding of ISW1b to chromatin. However, we find that chromatin
organization in *set1Δ* cells is essentially identical to wild type at the global level (***Figure 5D***) and quite
different from that in *ioc2Δ* cells (***Figure 5E***). Dinucleosome distribution in *set1Δ* cells is also similar to
wild type cells (***Figure 5F; Figure S3***). Thus, H3-K4me3 is not required for ISW1-dependent nucleosome
spacing.
313
314

Deletion of the H4 N-terminal tail domain results in shorter nucleosome spacing but does not increase
dinucleosome levels
Nucleosome mobilisation in vitro by ISWI complexes isolated from different organisms requires the H4
N-terminal tail domain, specifically the "basic patch" (residues 16 to 19) (***Clapier et al., 2002, 2001; Yan***
et al., 2019). In addition, genetic interactions between *isw1* mutations and H4 mutations (point
mutations and a tail deletion) suggest involvement in a common pathway (Lindstrom et al., 2006). We
reasoned that deletion of the H4 N-tail might result in a chromatin organization similar to that observed
in the *isw1Δ* single mutant: reduced spacing, weaker phasing and increased dinucleosome formation
(***Figures 1B and 3A***). We constructed a yeast strain in which both H3/H4 gene loci (*HHT1-HHF1* and
HHT2-HHF2) were deleted and covered by a plasmid carrying wild type *HHT1-HHF1*, or by *HHT1-*
HHF1Δ21, in which the first 21 amino acid residues of the H4 N-tail had been deleted. This strain
displayed a clear growth phenotype, with a doubling time of ~2.6 h, compared with ~1.8 h for the wild
type strain, in synthetic complete (SC) medium. We find that nucleosome spacing is shorter in cells
lacking the H4 N-tail (***Figure 6A***), though not quite as short as in *isw1Δ* cells (***Figure 1B***), and both
mutants have poor phasing. These observations are consistent with the requirement of Isw1 for the H4
N-tail domain. However, removal of the H4 N-tail did not increase dinucleosome prevalence (***Figures 6B***
and S3), unlike loss of Isw1 (***Figure 4B***), indicating that loss of Isw1 and loss of the H4 N-tail are not
equivalent.

Figure 6. Deletion of the first 21 residues of the H4 N-terminal tail domain results in shorter global nucleosome spacing but no dinucleosome accumulation. **(A)** Comparison of the H4ΔN21 mutant with its isogenic wild type strain (WT-H4) (top panel). WT-H4 replicates (bottom panel). All yeast genes were aligned on the midpoints of their +1 nucleosomes. The dyad distribution was normalised to the global average (set at 1). WT-H4 replicate A: black line with grey fill in both plots. The average spacing in bp is shown for each replicate in the bottom right corner. **(B)** Heat map analysis of nucleosome occupancy (coverage) for all genes aligned on the midpoint of their average +1 nucleosome position and sorted by Pol II density on the coding region (CDS), using ChIP-seq data for the Rpb3 subunit (***Ocampo et al., 2016***). Data for replicate A are shown (see ***Figure S4*** for replicate B). The dashed line separates the top 300 most active genes from the rest. The scale bar is the same for all plots, ranging from blue (low nucleosome occupancy) to red (high occupancy).

Discussion

ISW1b is a primary nucleosome spacing enzyme in yeast

We and others have shown previously that Isw1 and Chd1 are both needed for normal global chromatin organization in yeast (*Gkikopoulos et al., 2011; Ocampo et al., 2016*). We proposed that these two enzymes compete to set nucleosome spacing in wild type cells, with Isw1 being dominant, setting wild type spacing, and Chd1 directing shorter spacing (*Ocampo et al., 2016*). In the absence of Isw1, the spacing is short, which we attributed to Chd1 activity. An important complication, which we have addressed here, is that there are two complexes containing the Isw1 ATPase subunit (*Vary et al., 2003*). We have determined their respective roles in chromatin organization in vivo: ISW1b and Chd1 are primarily responsible for global nucleosome spacing in wild type cells, whereas a role for ISW1a in directing intermediate spacing only becomes apparent when ISW1b is absent (*Figure 7A*). We note that INO80 also affects global spacing (*Qiu et al., 2020; van Bakel et al., 2013; Yen et al., 2012*), but it is not yet clear how its activity meshes with that of ISW1b and Chd1.

Figure 7. Dominant role of ISW1b in chromatin organization. (**A**) Global average nucleosome spacing is primarily determined by ISW1b (wild type cells), since loss of ISW1a has little effect (*ioc3Δ* mutant). In the absence of ISW1b (*ioc2Δ* and *ioc4Δ* mutants), intermediate spacing is observed, attributed to ISW1a, since the spacing is short in the absence of both ISW1a and ISW1b (*isw1Δ* mutant or *ioc2Δ ioc3Δ ioc4Δ* triple mutant). Short spacing is attributed to Chd1, because the *chd1Δ isw1Δ* double mutant has highly disrupted chromatin. (**B**) Dinucleosomes disrupt phasing and reduce spacing. Model to show the effects of dinucleosomes on chromatin organization. Top panel: Average nucleosome positions on a gene in wild type cells with regular spacing. Middle: Close-packed dinucleosomes (grey ovals) are high in the absence of both ISW1b and Chd1. These dinucleosomes preferentially involve the +2 nucleosome i.e., they are mostly +1/+2 or +2/+3 dinucleosomes. Nucleosomes farther down the gene (+3 etc.) may be regularly spaced relative to the dinucleosome, but because a linker is missing in the dinucleosome, downstream nucleosomes are out of phase with other nucleosomal arrays, resulting in poor phasing due to interference patterns from the different arrays. Bottom: Dinucleosomes are absent from the mononucleosome dyad density plots, resulting in depressed, flattened peaks. These effects are strongest in cells lacking both ISW1b and Chd1. Dinucleosomes that cannot be resolved by ISW1b can account for shorter spacing in *chd1Δ* cells.

A difficulty for our competition model is that, in the absence of Chd1, ISW1b is expected to win
the competition, resulting in wild type spacing. Instead, we observe that spacing is somewhat shorter in
cells lacking Chd1 (***Ocampo et al., 2016***) (***Figure 2A***). One possible explanation is that the dominance of
ISW1b depends on Chd1 and so, in its absence, we see an increased contribution from ISW1a, such that
the spacing represents the average of ISW1a and ISW1b activities. If so, cells having only ISW1b (the
chd1Δ ioc3Δ double mutant) are expected to have wild type spacing. However, the result is intermediate
spacing, very similar to the *chd1Δ* single mutant (***Figure 2***). Alternatively, ISW1b might only be able to
create nucleosome arrays with wild type spacing if Chd1 has already made arrays with short spacing,
although this model is inconsistent with in vitro data showing that purified ISW1b can space
nucleosomes by itself (***Vary et al., 2003***).
A more satisfying explanation involves dinucleosomes (***Figure 7B***). Consider a single gene. If
there are regularly spaced nucleosomes downstream of a close-packed +1/+2 dinucleosome (i.e. no
linker), these nucleosomes will be out of phase by one linker length with nucleosomes in cells with no
dinucleosome on this gene. Other cells may have a +2/+3 dinucleosome instead, which will alter the
positions of regularly spaced downstream nucleosomes to give a different phasing. The result is
generally weaker phasing (peak flattening) and a shift to shorter average spacing. Moreover, since the
dinucleosomes are not counted in the mononucleosome phasing pattern, there will be missing
occupancy around the +2 nucleosome, resulting in more pattern disruption (primarily a depressed +2
nucleosome peak). These effects will increase as the fraction of genes having a dinucleosome increases:
the *chd1Δ isw1Δ* double mutant and the *chd1Δ ioc2Δ ioc4Δ* triple mutant both have high dinucleosome
levels and extremely poor phasing. The *chd1Δ* single mutant has fewer dinucleosomes, resulting in a
smaller shift in the peaks to shorter spacing. These considerations can account for the shift to shorter
spacing in the absence of Chd1 and for the major disruption of chromatin in the absence of both Chd1
and ISW1b. Thus, we propose that the spacing is shorter in cells lacking Chd1 because ISW1b cannot
resolve all of the dinucleosomes.
377
Dinucleosome levels correlate with Pol II levels
Previously, we showed that Isw1 and/or Chd1 is required to resolve dinucleosomes, or to prevent their
formation (***Ocampo et al., 2019***). Here we show that both enzymes are important for separating
dinucleosomes and that, as observed for spacing activity, ISW1b is much more important than ISW1a. In
vitro, both ISW1a and ISW1b can move nucleosomes reconstituted on a pair of 601 nucleosome
positioning sequences farther apart, even if separated by a linker of only 4 bp (***Krajewski, 2014***),
indicating that both ISW1 complexes probably have close-packed dinucleosome resolving activity.
Interestingly, SWI/SNF may also be able to resolve dinucleosomes (***Dechassa et al., 2010***).
We observe a correlation between dinucleosome and Pol II levels on genes, such that the least
active genes have fewer dinucleosomes. The top ~300 most active genes represent an exception in that
they are only modestly enriched in dinucleosomes, relative to the genes immediately below them in the
heat map, which we attribute to the nucleosome loss that occurs at the most active genes (***Cole et al.,***
2014; Schwabish and Struhl, 2007, 2004). This loss may include both mononucleosomes and
dinucleosomes. There is an interesting conundrum concerning these most active genes, since they have
both very short nucleosome spacing (roughly ~155 bp), which implies that more nucleosomes are
present on a gene of a given length, and reduced nucleosome occupancy, which indicates fewer

nucleosomes. A possible resolution is that these genes have fewer nucleosomes which form clusters
with short spacing separated by gaps (i.e., long linkers).
The correlation between dinucleosomes and transcriptional activity suggests that Pol II might
form dinucleosomes as it transcribes along a gene, perhaps as part of a general tendency to push
nucleosomes closer together during transcription, possibly to free up the space needed for the Pol II
elongation complex itself, since the average linker is only ~20 bp. Some support for this hypothesis
comes from the observation that nucleosome spacing is longer in non-transcribing yeast cells than in
wild type cells (this experiment was possible using the Pol II large subunit *rpb1-1* mutant at the non-
permissive temperature) (***Weiner et al., 2010***), suggesting that transcription shortens spacing, although
the effect could be indirect.
In summary, we propose that transcription may result in some nucleosome loss, and some
clustering of the remaining nucleosomes with short spacing, including some close-packed
dinucleosomes. ISW1b and Chd1 are needed to space the nucleosomes out again after transcription.
Once a gene is transcribed, its chromatin organization is not restored if ISW1b and Chd1 are absent. It
seems likely that dinucleosomes might interfere with transcription and other processes. If so, regulating
dinucleosome formation might represent a novel mechanism for control of gene expression.
410
The Set2 methyltransferase makes a subtle contribution to global chromatin organization
We observe that the ancillary subunits of ISW1b, Ioc2 and Ioc4, are both necessary for ISW1b-
dependent chromatin organization, although Ioc2 makes a greater contribution, since it affects phasing
as well as spacing. We suggest that Ioc2 and Ioc4 cooperate to influence the remodeling activity of the
Isw1 subunit. Both are linked to certain histone modifications. Ioc4 binds to an H3-K36me3 peptide with
higher affinity than the unmethylated peptide, suggesting that Set2-mediated H3-K36me3 might be an
important regulator of ISW1b (***Maltby et al., 2012; Smolle et al., 2012; Smolle and Workman, 2013***).
Our data provide some evidence to support this proposal, because loss of Set2 has a similar but weaker
effect on chromatin organization as loss of Ioc4, although they are not quite equivalent.
Ioc2 has a PHD domain (***Smolle et al., 2012***) which binds H3-K4me3, accounting for the H3-
K4me3/Set1-dependence of Isw1 binding to chromatin at a subset of genes (***Santos-Rosa et al., 2003***).
However, our data indicate that the nucleosome spacing activity of the ISW1b complex, which requires
Isw1 binding to chromatin, is independent of Set1 (and therefore of H3-K4me3) at the global level, since
cells lacking Set1 have normal chromatin organization. We also note that, since loss of Set1 did not
result in global chromatin defects, the binding of Chd1 to H3K4me3 through one of its chromodomains
(***Pray-Grant et al., 2005***) is not critical for its nucleosome spacing function.
We note that neither ISW1 complex requires H3-K4me3 or H3-K36me3 to space nucleosomes in
vitro, since they are active on nucleosomes containing unmodified recombinant histones. On the other
hand, in vivo, H3-K36me3 and H3-K4me3 may have local effects on ISW1b activity, rather than global
effects, which we have not detected. These are probably not occurring at the most active genes because
their chromatin organization in *set1Δ* and *set2Δ* cells is similar to wild type (***Figure S2C***). Alternatively,
H3-K36me3 and H3-K4me3 recognition may affect other functions of ISW1b, such as gene repression
(***Pinskaya et al., 2009***) or mRNP quality control (***Babour et al., 2016***).
434
435

The H4 N-terminal tail domain is required for normal nucleosome spacing

The H4 tail domain is generally required by ISWI-like enzymes for activity (*Clapier et al., 2017, 2001; Hamiche et al., 2001; Paul and Bartholomew, 2018*). The ISWI ATPase subunit has an inhibitory AutoN domain that resembles the H4 tail, which is displaced by the H4 tail when ISWI binds to a nucleosome (*Clapier et al., 2017; Paul and Bartholomew, 2018*). The isolated yeast Isw1 subunit is inactive without Ioc subunits unless the AutoN domain is inactivated by mutation (*Yan et al., 2019*), implying that the Ioc subunits may regulate Isw1 activity through the AutoN domain. We find that the H4 tail deletion mutant exhibits shorter spacing, consistent with inactivation of ISW1b, but there is no increase in dinucleosomes. The most likely explanation for the non-equivalence of the H4Δ21 and *isw1Δ* mutations is that the H4 tail domain is involved in multiple processes, including interactions with other remodelers and proteins. This possibility is supported by the fact that the H4Δ21 mutant has a strong growth defect, unlike the various *ioc* mutants. Since Chd1 also makes contacts with the basic patch in the H4 tail (*Ferreira et al., 2007; Sundaramoorthy et al., 2018*), its remodeling activity may also be compromised, although its chromatin is clearly not as disrupted as in the *chd1Δ isw1Δ* double mutant. The H4 N-tail is also required by ISW2 in vitro (*Dang et al., 2006*), although global chromatin organization is unaffected in *isw2Δ* cells (*Gkikopoulos et al., 2011; Ocampo et al., 2016*). The low prevalence of dinucleosomes in the H4Δ21 mutant suggests that the H4 tail might be required for dinucleosome formation, possibly because the increased negative charge on nucleosomes lacking the H4 tail might limit how closely nucleosomes can approach one another (*Blank and Becker, 1995; Clark and Kimura, 1990*). Deletion of the H4 tail also results in the loss of multiple post-translational modification sites, preventing interaction with many chromatin factors.

Materials and methods

Plasmid construction. The primers used in this study are listed in *Table S1*. pRS-HHT1-HHF1 (p368) was constructed by insertion of the 1930-bp SacI-PstI fragment containing the *HHT1-HHF1* locus from p367 (**Eriksson et al., 2005**) at the same sites in pRS317 (*CEN ARS LYS2*) (ATCC 77157). pRS-HHT1-HHF1Δ21 (p730) was constructed by replacing the 752-bp AfeI-SacI fragment in p368 with a 689-bp AfeI-SacI fragment with the H4 N-terminal tail deletion (H4 begins Met-Leu22…) made by ligating two PCR fragments together (obtained using p368 as template with primers 1770/1771 and 1773/1812). The sequence was confirmed.

Yeast strain construction. The yeast strains used in this study are listed in *Table S2*. YDC507 (H4 N-terminal tail deletion) was constructed by transforming ROY1281 (**Dhillon and Kamakaka, 2000**), which carries plasmid pCC67 (2 micron origin *URA3 HHT1-HHT1 HTA1-HTB1*) (**Clark-Adams et al., 1988**), with p730 (*CEN ARS LYS2 HHT1-HHF1Δ21*), selection on plates made with synthetic complete medium without lysine (SC-lys) and then counter-selection against *URA3* with 5-fluoro-orotic acid (5-FOA) to evict pCC67. A wild type control strain (YDC101) was constructed in the same way using p368 (*CEN ARS LYS2 HHT1-HHF1*) instead of p730. The diploid strain YPE600 was made by crossing YTT196 (**Tsukiyama et al., 1999**) with YDC111 (**Kim et al., 2006**). YPE606 (*ioc3Δ*) was constructed by transforming YPE600 with an *ioc3Δ::KanMX* fragment made using primers 1918/1919 and genomic DNA from YTT645, followed by G418 selection on YPD plates and sporulation. YPE608 (*ioc4Δ*) was constructed by transforming YPE600 with an *ioc4Δ::HPH1* fragment made using primers 1922/1923 and genomic DNA from YTT827, followed by hygromycin selection on YPD plates and sporulation. YPE636 (*ioc2Δ*) was constructed by transforming YPE600 with an *ioc2Δ::URA3* fragment made using primers 1961/1962 and extended using primers 1965/1966 and wild type genomic DNA as template, followed by selection on SC-ura plates and sporulation. YPE654 (*ioc2Δ ioc3Δ*) was constructed by transforming YPE636 with the *ioc3Δ::KanMX* DNA fragment (see above). YPE655 (*ioc2Δ ioc4Δ*) was constructed by transforming YPE636 with the *ioc4Δ::HPH1* fragment (see above). YPE657 (*ioc2Δ ioc3Δ ioc4Δ*) was constructed by transforming YPE654 with the *ioc4Δ::HPH1* fragment. YPE712 (*chd1Δ ioc3Δ*) was obtained by crossing YJO486 (**Ocampo et al., 2016**) with YPE607. YJO487 was obtained by sporulation of YJO505 (**Ocampo et al., 2016**). YPE715 (*chd1Δ ioc2Δ ioc4Δ*) was constructed by crossing YJO487 with YPE655. YPE602 (*set1Δ*) was constructed by transforming YPE600 with a *set1Δ::NAT1* fragment made using primers 1926/1927 and genomic DNA from YTT1986, followed by nourseothricin selection on YPD plates and sporulation. YPE604 (*set2Δ*) was constructed by transforming YDC111 with a *set2Δ::TRP1* fragment made using primers 1914/1915 extended with 1916/17 and plasmid pFA6a-TRP1 (**Longtine et al., 1998**), followed by selection on SC-trp plates.

MNase-seq: Nuclei were prepared as described (**Cole et al., 2011**). MNase digestion of nuclei and construction of paired-end libraries was as described (**Chereji et al., 2017**), except that mononucleosomal DNA was not gel-purified, in order to retain mononucleosomes and dinucleosomes in the correct proportions (**Ocampo et al., 2019**). Biological replicate experiments were performed for all strains. MNase-seq data were analysed using scripts described by (**Ocampo et al., 2019**). Sequence data

499   for YJO484 (*chd1Δ isw1Δ*) are available at the GEO database (GSE117514 (***Ocampo et al., 2019***)). The
500   sequence data from this study have been submitted to the NCBI Gene Expression Omnibus (GEO;
501   http://www.ncbi.nlm.nih.gov/geo/) under accession number GSE156224.
502

Acknowledgements

We thank Alan Hinnebusch for helpful comments on the manuscript and Răzvan Chereji for help with
the bioinformatics. We thank Alex Naim for constructing p368, Toshi Tsukiyama for yeast strains, and
Rohinton Kamakaka and Namrita Dhillon for pCC67 (pRO149) and ROY1281. This research was
supported by the Intramural Research Program of the NIH (NICHD). We thank the NHLBI Core Facility
(Yan Luo, Poching Liu, and Jun Zhu) for paired-end sequencing. This study utilized the high performance
computational capabilities of the Biowulf Linux cluster at the National Institutes of Health.

References

Babour, A., Shen, Q., Dos-Santos, J., Murray, S., Gay, A., Challal, D., Fasken, M., Palancade, B., Corbett,
A., Libri, D., Mellor, J., Dargemont, C., 2016. The Chromatin Remodeler ISW1 Is a Quality Control
Factor that Surveys Nuclear mRNP Biogenesis. Cell 167, 1201-1214.e15.
doi:10.1016/j.cell.2016.10.048
Barisic, D., Stadler, M.B., Iurlaro, M., Schübeler, D., 2019. Mammalian ISWI and SWI/SNF selectively
mediate binding of distinct transcription factors. Nature 569, 136–140. doi:10.1038/s41586-019-
1115-5
Blank, T.A., Becker, P.B., 1995. Electrostatic mechanism of nucleosome spacing. J. Mol. Biol. 252, 305–
313. doi:10.1006/jmbi.1995.0498
Chereji, R.V., Clark, D.J., 2018. Major determinants of nucleosome positioning. Biophys. J. 114, 2279–
2289. doi:10.1016/j.bpj.2018.03.015
Chereji, R.V., Ocampo, J., Clark, D.J., 2017. MNase-Sensitive Complexes in Yeast: Nucleosomes and Non-
histone Barriers. Mol. Cell 65, 565-577.e3. doi:10.1016/j.molcel.2016.12.009
Clapier, C.R., Iwasa, J., Cairns, B.R., Peterson, C.L., 2017. Mechanisms of action and regulation of ATP-
dependent chromatin-remodelling complexes. Nat. Rev. Mol. Cell Biol. 18, 407–422.
doi:10.1038/nrm.2017.26
Clapier, C.R., Längst, G., Corona, D.F., Becker, P.B., Nightingale, K.P., 2001. Critical role for the histone H4
N terminus in nucleosome remodeling by ISWI. Mol. Cell. Biol. 21, 875–883.
doi:10.1128/MCB.21.3.875-883.2001
Clapier, C.R., Nightingale, K.P., Becker, P.B., 2002. A critical epitope for substrate recognition by the
nucleosome remodeling ATPase ISWI. Nucleic Acids Res. 30, 649–655. doi:10.1093/nar/30.3.649
Clark, D.J., Kimura, T., 1990. Electrostatic mechanism of chromatin folding. J. Mol. Biol. 211, 883–896.
doi:10.1016/0022-2836(90)90081-V
Cole, H.A., Howard, B.H., Clark, D.J., 2011. Activation-induced disruption of nucleosome position clusters
on the coding regions of Gcn4-dependent genes extends into neighbouring genes. Nucleic Acids
Res. 39, 9521–9535. doi:10.1093/nar/gkr643
Cole, H.A., Ocampo, J., Iben, J.R., Chereji, R.V., Clark, D.J., 2014. Heavy transcription of yeast genes

correlates with differential loss of histone H2B relative to H4 and queued RNA polymerases. Nucleic
Acids Res. 42, 12512–12522. doi:10.1093/nar/gku1013

Cui, F., Cole, H.A., Clark, D.J., Zhurkin, V.B., 2012. Transcriptional activation of yeast genes disrupts
intragenic nucleosome phasing. Nucleic Acids Res. 40, 10753–10764. doi:10.1093/nar/gks870

Dang, W., Kagalwala, M.N., Bartholomew, B., 2006. Regulation of ISW2 by concerted action of histone
H4 tail and extranucleosomal DNA. Mol. Cell. Biol. 26, 7388–7396. doi:10.1128/MCB.01159-06

Dechassa, M.L., Sabri, A., Pondugula, S., Kassabov, S.R., Chatterjee, N., Kladde, M.P., Bartholomew, B.,
2010. SWI/SNF has intrinsic nucleosome disassembly activity that is dependent on adjacent
nucleosomes. Mol. Cell 38, 590–602. doi:10.1016/j.molcel.2010.02.040

Dhillon, N., Kamakaka, R.T., 2000. A histone variant, Htz1p, and a Sir1p-like protein, Esc2p, mediate
silencing at HMR. Mol. Cell 6, 769–780. doi:10.1016/s1097-2765(00)00076-9

Eriksson, P.R., Mendiratta, G., McLaughlin, N.B., Wolfsberg, T.G., Mariño-Ramírez, L., Pompa, T.A.,
Jainerin, M., Landsman, D., Shen, C.-H., Clark, D.J., 2005. Global regulation by the yeast Spt10
protein is mediated through chromatin structure and the histone upstream activating sequence
elements. Mol. Cell. Biol. 25, 9127–9137. doi:10.1128/MCB.25.20.9127-9137.2005

Ferreira, H., Flaus, A., Owen-Hughes, T., 2007. Histone modifications influence the action of Snf2 family
remodelling enzymes by different mechanisms. J. Mol. Biol. 374, 563–579.
doi:10.1016/j.jmb.2007.09.059

Gangaraju, V.K., Bartholomew, B., 2007. Dependency of ISW1a chromatin remodeling on
extranucleosomal DNA. Mol. Cell. Biol. 27, 3217–3225. doi:10.1128/MCB.01731-06

Gkikopoulos, T., Schofield, P., Singh, V., Pinskaya, M., Mellor, J., Smolle, M., Workman, J.L., Barton, G.J.,
Owen-Hughes, T., 2011. A role for Snf2-related nucleosome-spacing enzymes in genome-wide
nucleosome organization. Science 333, 1758–1760. doi:10.1126/science.1206097

Hamiche, A., Kang, J.G., Dennis, C., Xiao, H., Wu, C., 2001. Histone tails modulate nucleosome mobility
and regulate ATP-dependent nucleosome sliding by NURF. Proc Natl Acad Sci USA 98, 14316–
14321. doi:10.1073/pnas.251421398

Ito, T., Bulger, M., Pazin, M.J., Kobayashi, R., Kadonaga, J.T., 1997. ACF, an ISWI-containing and ATP-
utilizing chromatin assembly and remodeling factor. Cell 90, 145–155. doi:10.1016/s0092-
8674(00)80321-9

Kim, Y., McLaughlin, N., Lindstrom, K., Tsukiyama, T., Clark, D.J., 2006. Activation of Saccharomyces
cerevisiae HIS3 results in Gcn4p-dependent, SWI/SNF-dependent mobilization of nucleosomes over
the entire gene. Mol. Cell. Biol. 26, 8607–8622. doi:10.1128/MCB.00678-06

Krajewski, W.A., 2014. Yeast Isw1a and Isw1b exhibit similar nucleosome mobilization capacities for
mononucleosomes, but differently mobilize dinucleosome templates. Arch. Biochem. Biophys. 546,
72–80. doi:10.1016/j.abb.2014.02.003

Krietenstein, N., Wal, M., Watanabe, S., Park, B., Peterson, C.L., Pugh, B.F., Korber, P., 2016. Genomic
Nucleosome Organization Reconstituted with Pure Proteins. Cell 167, 709-721.e12.
doi:10.1016/j.cell.2016.09.045

Lafon, A., Petty, E., Pillus, L., 2012. Functional antagonism between Sas3 and Gcn5 acetyltransferases
and ISWI chromatin remodelers. PLoS Genet. 8, e1002994. doi:10.1371/journal.pgen.1002994

Lieleg, C., Ketterer, P., Nuebler, J., Ludwigsen, J., Gerland, U., Dietz, H., Mueller-Planitz, F., Korber, P.,
2015. Nucleosome spacing generated by ISWI and CHD1 remodelers is constant regardless of

nucleosome density. Mol. Cell. Biol. 35, 1588–1605. doi:10.1128/MCB.01070-14

Lindstrom, K.C., Vary, J.C., Parthun, M.R., Delrow, J., Tsukiyama, T., 2006. Isw1 functions in parallel with

the NuA4 and Swr1 complexes in stress-induced gene repression. Mol. Cell. Biol. 26, 6117–6129.

doi:10.1128/MCB.00642-06

Longtine, M.S., McKenzie, A., Demarini, D.J., Shah, N.G., Wach, A., Brachat, A., Philippsen, P., Pringle,

J.R., 1998. Additional modules for versatile and economical PCR-based gene deletion and

modification in Saccharomyces cerevisiae. Yeast 14, 953–961. doi:10.1002/(SICI)1097-

0061(199807)14:10<953::AID-YEA293>3.0.CO;2-U

Luger, K., Mäder, A.W., Richmond, R.K., Sargent, D.F., Richmond, T.J., 1997. Crystal structure of the

nucleosome core particle at 2.8 A resolution. Nature 389, 251–260. doi:10.1038/38444

Lusser, A., Urwin, D.L., Kadonaga, J.T., 2005. Distinct activities of CHD1 and ACF in ATP-dependent

chromatin assembly. Nat. Struct. Mol. Biol. 12, 160–166. doi:10.1038/nsmb884

Maltby, V.E., Martin, B.J.E., Schulze, J.M., Johnson, I., Hentrich, T., Sharma, A., Kobor, M.S., Howe, L.,

2012. Histone H3 lysine 36 methylation targets the Isw1b remodeling complex to chromatin. Mol.

Cell. Biol. 32, 3479–3485. doi:10.1128/MCB.00389-12

Mellor, J., Morillon, A., 2004. ISWI complexes in Saccharomyces cerevisiae. Biochim. Biophys. Acta 1677,

100–112. doi:10.1016/j.bbaexp.2003.10.014

Morillon, A., Karabetsou, N., O'Sullivan, J., Kent, N., Proudfoot, N., Mellor, J., 2003. Isw1 chromatin

remodeling ATPase coordinates transcription elongation and termination by RNA polymerase II.

Cell 115, 425–435. doi:10.1016/s0092-8674(03)00880-8

Ocampo, J., Chereji, R.V., Eriksson, P.R., Clark, D.J., 2016. The ISW1 and CHD1 ATP-dependent chromatin

remodelers compete to set nucleosome spacing in vivo. Nucleic Acids Res. 44, 4625–4635.

doi:10.1093/nar/gkw068

Ocampo, J., Chereji, R.V., Eriksson, P.R., Clark, D.J., 2019. Contrasting roles of the RSC and ISW1/CHD1

chromatin remodelers in RNA polymerase II elongation and termination. Genome Res. 29, 407–417.

doi:10.1101/gr.242032.118

Paul, S., Bartholomew, B., 2018. Regulation of ATP-dependent chromatin remodelers:

accelerators/brakes, anchors and sensors. Biochem. Soc. Trans. 46, 1423–1430.

doi:10.1042/BST20180043

Pinskaya, M., Nair, A., Clynes, D., Morillon, A., Mellor, J., 2009. Nucleosome remodeling and

transcriptional repression are distinct functions of Isw1 in Saccharomyces cerevisiae. Mol. Cell. Biol.

29, 2419–2430. doi:10.1128/MCB.01050-08

Pointner, J., Persson, J., Prasad, P., Norman-Axelsson, U., Strålfors, A., Khorosjutina, O., Krietenstein, N.,

Svensson, J.P., Ekwall, K., Korber, P., 2012. CHD1 remodelers regulate nucleosome spacing in vitro

and align nucleosomal arrays over gene coding regions in S. pombe. EMBO J. 31, 4388–4403.

doi:10.1038/emboj.2012.289

Prajapati, H.K., Ocampo, J., Clark, D.J., 2020. Interplay among ATP-Dependent Chromatin Remodelers

Determines Chromatin Organization in Yeast. Biology (Basel) 9. doi:10.3390/biology9080190

Pray-Grant, M.G., Daniel, J.A., Schieltz, D., Yates, J.R., Grant, P.A., 2005. Chd1 chromodomain links

histone H3 methylation with SAGA- and SLIK-dependent acetylation. Nature 433, 434–438.

doi:10.1038/nature03242

Qiu, H., Biernat, E., Govind, C.K., Rawal, Y., Chereji, R.V., Clark, D.J., Hinnebusch, A.G., 2020. Chromatin

remodeler Ino80C acts independently of H2A.Z to evict promoter nucleosomes and stimulate
transcription of highly expressed genes in yeast. Nucleic Acids Res. doi:10.1093/nar/gkaa571
Richmond, T.J., 2012. Nucleosome recognition and spacing by chromatin remodelling factor ISW1a.
Biochem. Soc. Trans. 40, 347–350. doi:10.1042/BST20110748
Santos-Rosa, H., Schneider, R., Bernstein, B.E., Karabetsou, N., Morillon, A., Weise, C., Schreiber, S.L.,
Mellor, J., Kouzarides, T., 2003. Methylation of histone H3 K4 mediates association of the Isw1p
ATPase with chromatin. Mol. Cell 12, 1325–1332. doi:10.1016/s1097-2765(03)00438-6
Schwabish, M.A., Struhl, K., 2004. Evidence for eviction and rapid deposition of histones upon
transcriptional elongation by RNA polymerase II. Mol. Cell. Biol. 24, 10111–10117.
doi:10.1128/MCB.24.23.10111-10117.2004
Schwabish, M.A., Struhl, K., 2007. The Swi/Snf complex is important for histone eviction during
transcriptional activation and RNA polymerase II elongation in vivo. Mol. Cell. Biol. 27, 6987–6995.
doi:10.1128/MCB.00717-07
Simic, R., Lindstrom, D.L., Tran, H.G., Roinick, K.L., Costa, P.J., Johnson, A.D., Hartzog, G.A., Arndt, K.M.,
2003. Chromatin remodeling protein Chd1 interacts with transcription elongation factors and
localizes to transcribed genes. EMBO J. 22, 1846–1856. doi:10.1093/emboj/cdg179
Smolle, M., Venkatesh, S., Gogol, M.M., Li, H., Zhang, Y., Florens, L., Washburn, M.P., Workman, J.L.,
2012. Chromatin remodelers Isw1 and Chd1 maintain chromatin structure during transcription by
preventing histone exchange. Nat. Struct. Mol. Biol. 19, 884–892. doi:10.1038/nsmb.2312
Smolle, M., Workman, J.L., 2013. Transcription-associated histone modifications and cryptic
transcription. Biochim. Biophys. Acta 1829, 84–97. doi:10.1016/j.bbagrm.2012.08.008
Stockdale, C., Flaus, A., Ferreira, H., Owen-Hughes, T., 2006. Analysis of nucleosome repositioning by
yeast ISWI and Chd1 chromatin remodeling complexes. J. Biol. Chem. 281, 16279–16288.
doi:10.1074/jbc.M600682200
Sundaramoorthy, R., Hughes, A.L., El-Mkami, H., Norman, D.G., Ferreira, H., Owen-Hughes, T., 2018.
Structure of the chromatin remodelling enzyme Chd1 bound to a ubiquitinylated nucleosome. elife
7. doi:10.7554/eLife.35720
Thoma, F., Koller, T., Klug, A., 1979. Involvement of histone H1 in the organization of the nucleosome
and of the salt-dependent superstructures of chromatin. J. Cell Biol. 83, 403–427.
doi:10.1083/jcb.83.2.403
Tirosh, I., Sigal, N., Barkai, N., 2010. Widespread remodeling of mid-coding sequence nucleosomes by
Isw1. Gen. Biol. 11:R49. doi:10.1186/gb-2010-11-5-r49
Torigoe, S.E., Patel, A., Khuong, M.T., Bowman, G.D., Kadonaga, J.T., 2013. ATP-dependent chromatin
assembly is functionally distinct from chromatin remodeling. elife 2, e00863.
doi:10.7554/eLife.00863
Tsukiyama, T., Palmer, J., Landel, C.C., Shiloach, J., Wu, C., 1999. Characterization of the imitation switch
subfamily of ATP-dependent chromatin-remodeling factors in Saccharomyces cerevisiae. Genes
Dev. 13, 686–697. doi:10.1101/gad.13.6.686
Udugama, M., Sabri, A., Bartholomew, B., 2011. The INO80 ATP-dependent chromatin remodeling
complex is a nucleosome spacing factor. Mol. Cell. Biol. 31, 662–673. doi:10.1128/MCB.01035-10
van Bakel, H., Tsui, K., Gebbia, M., Mnaimneh, S., Hughes, T.R., Nislow, C., 2013. A compendium of
nucleosome and transcript profiles reveals determinants of chromatin architecture and

transcription. PLoS Genet. 9, e1003479. doi:10.1371/journal.pgen.1003479

Vary, J.C., Fazzio, T.G., Tsukiyama, T., 2004. Assembly of yeast chromatin using ISWI complexes. Meth.
Enzymol. 375, 88–102.

Vary, J.C., Gangaraju, V.K., Qin, J., Landel, C.C., Kooperberg, C., Bartholomew, B., Tsukiyama, T., 2003.
Yeast Isw1p forms two separable complexes in vivo. Mol. Cell. Biol. 23, 80–91.
doi:10.1128/mcb.23.1.80-91.2003

Weiner, A., Hughes, A., Yassour, M., Rando, O.J., Friedman, N., 2010. High-resolution nucleosome
mapping reveals transcription-dependent promoter packaging. Genome Res. 20, 90–100.
doi:10.1101/gr.098509.109

Wiechens, N., Singh, V., Gkikopoulos, T., Schofield, P., Rocha, S., Owen-Hughes, T., 2016. The Chromatin
Remodelling Enzymes SNF2H and SNF2L Position Nucleosomes adjacent to CTCF and Other
Transcription Factors. PLoS Genet. 12, e1005940. doi:10.1371/journal.pgen.1005940

Yamada, K., Frouws, T.D., Angst, B., Fitzgerald, D.J., DeLuca, C., Schimmele, K., Sargent, D.F., Richmond,
T.J., 2011. Structure and mechanism of the chromatin remodelling factor ISW1a. Nature 472, 448–
453. doi:10.1038/nature09947

Yan, L., Wu, H., Li, X., Gao, N., Chen, Z., 2019. Structures of the ISWI-nucleosome complex reveal a
conserved mechanism of chromatin remodeling. Nat. Struct. Mol. Biol. 26, 258–266.
doi:10.1038/s41594-019-0199-9

Yen, K., Vinayachandran, V., Batta, K., Koerber, R.T., Pugh, B.F., 2012. Genome-wide nucleosome
specificity and directionality of chromatin remodelers. Cell 149, 1461–1473.
doi:10.1016/j.cell.2012.04.036

Zhao, X., Muller, E.G., Rothstein, R., 1998. A suppressor of two essential checkpoint genes identifies a
novel protein that negatively affects dNTP pools. Mol. Cell 2, 329–340. doi:10.1016/s1097-
2765(00)80277-4

Supplementary Figures

A   Ioc3-bound genes

B   Ioc4-bound genes

C   Isw1-bound genes

Position relative to the +1 nucleosome (kb)

692
693
Figure S1. The chromatin organizations of Ioc3-bound, Ioc4-bound and Isw1-bound genes are not
significantly different from the unbound genes. Target genes for ISW1a (Ioc3-bound), ISW1b (Ioc4-
bound) and both complexes (Isw1-bound) are defined by ChIP-seq data (*Yen et al., 2012*). Average
nucleosome dyad density plots for bound and unbound genes: (**A**) Ioc3-bound (blue line) and non-bound
(red line) genes in wild type (WT) and *ioc3Δ* cells. (**B**) Ioc4-bound (blue line) and non-bound (red line)
genes in WT and *ioc4Δ* cells. (**C**) Isw1-bound (blue line) and non-bound (red line) genes in WT and *isw1Δ*
cells. All 5770 yeast genes were aligned on the midpoints of their +1 nucleosomes. The dyad distribution
was normalized to the global average (set at 1). Two biological replicate experiments (A and B) are
shown. The average spacing in bp is shown for bound genes (blue text) and non-bound genes (red text)
in the bottom right corner (measured by regression analysis of the first 5 nucleosome peaks, beginning
with the +1 nucleosome).

705
706 **Figure S2:** *continued on next page.*
707

Figure S2. Chromatin organization of the top 300 most transcriptionally active genes (defined by Pol II

density using ChIP-seq data for the Rpb3 subunit in wild type cells; ***Ocampo et al., 2016***). Average

nucleosome dyad density plots for the 300 most active genes (blue line) compared with the remaining

5470 genes (red line). Genes were aligned on the midpoints of their +1 nucleosomes. The dyad

distribution was normalized to the global average (set at 1). Two biological replicate experiments (A and

B) are shown in separate panels. The average spacing in bp is shown for the top 300 active genes (blue

text) and the other genes (red text), as measured by regression analysis of the first 5 nucleosome peaks,

beginning with the +1 nucleosome. (**A**) Wild type (WT), *ioc2Δ*, *ioc3Δ ioc4Δ*, *ioc2Δ*, *ioc2Δ ioc4Δ*, *ioc2Δ*

ioc3Δ ioc4Δ and *isw1Δ*. (**B**) *chd1Δ*, *chd1Δ ioc3Δ*, *chd1Δ ioc2Δ ioc4Δ* and *chd1Δ isw1Δ*. (**C**) *set1Δ* and

set2Δ. (**D**) Wild type H4 and H4*Δ21*.

Figure S3. Mononucleosome and dinucleosome occupancy plots for all strains. Comparison of biological

replicate experiments. Average nucleosome occupancy (coverage) plots for all genes. All 5770 yeast

genes were aligned on the midpoint of their average +1 nucleosome position. The occupancy was

normalized to the global average (set at 1) for dinucleosomes (250 - 350 bp; left panels) or

mononucleosomes (120 - 180 bp; right panels). Note different y-axis scales are used in the dinucleosome

and mononucleosome plots. Replicate A (black line); replicate B (red line).

Coordinate relative to +1 nucleosome dyad at 0 (from -1000 to +1000 nt)

728
729
730 **Figure S4:** *continued on next page*.
731

Coordinate relative to +1 nucleosome dyad at 0 (from -1000 to +1000 nt)

733
734
Figure S4. Dinucleosome formation correlates with RNA polymerase II levels, except for the most highly
transcribed genes. Heat map analysis of nucleosome occupancy (coverage) for all genes aligned on the
midpoint of their average +1 nucleosome position and sorted by Pol II density on the coding region
(CDS), using ChIP-seq data for the Rpb3 subunit (*Ocampo et al., 2016*). Data for replicate B are shown
(see *Figures 4, 5 and 6* for replicate A). The dashed line separates the top 300 most active genes from
the rest. The scale bar is the same for all plots, ranging from blue (low nucleosome occupancy) to red
(high occupancy). (**A**) wild type, (**B**) *isw1Δ*, (**C**) *chd1Δ isw1Δ*, (**D**) *ioc2Δ*, (**E**) *ioc3Δ*, (**F**) *ioc4Δ*, (**G**) *ioc2Δ*
ioc4Δ, (**H**) *ioc2Δ ioc3Δ ioc4Δ*, (**I**) *chd1Δ*, (**J**) *chd1Δ ioc3Δ*, (**K**) *chd1Δ ioc2Δ ioc4Δ*, (**L**) *set1Δ*, (**M**) *set2Δ*, (**N**)
wild type-H4 and (**O**) H4Δ21.
744
745

Supplementary Tables

Table S1. Primers used in this study.

1770 (H4Δ21)   ACTGTTCCGAGCGCTTCT
1771 (H4Δ21)   [5'-phosphate]-CATATTTTACTATATTATATTTGTTGCTTG
1773 (H4Δ21)   CGAATTGGAGCTCGGTAC
1812 (H4Δ21)   [5'-phosphate]-CTAAGAGATAACATCCAAGGTATTAC
1914 (set2)   AAAACTGCATAGTCGTGCTGTCAAACCTTTCTCCTTTCCTGGTTGTTGTTTTACGTGATCCGGATCCCCGGGTTAATTAA
1915 (set2)   ACAAGACTTCCTTTGGGACAGAAAACGTGAAACAAGCCCCAAATATGCATGTCTGGTTAAGAATTCGAGCTCGTTTAAAC
1916 (set2)   TCGCAGTAATAAGGACAGTATCCG
1917 (set2)   GCAGTTAACATGTAAATTTCCCGAT
1918 (ioc3)   TTCACAAGATAACTACGACCCTCC
1919 (ioc3)   GATAACAGTATGCCGTTTTACCAC
1922 (ioc4)   TTCTGGTTTGTTTCTTTGCCTCA
1923 (ioc4)   CATATACGGTGTTAAGATGATGACA
1926 (set1)   AGGTTCATTAAATGCTTGGCTT
1927 (set1)   TTCTGGAGCGTATCTAATTGCTTG
1961 (ioc2)
GTGAATCAGCGATCGATGAGAACGAAGAGAACTAGGAGTACAAGAAATGTAGGAGCGAGCTTGTTCTTGATTTGTGCCCCGTA
1962 (ioc2)   ATGAAGAACCAGTTGATATATATTTTACTATATACTTTTGCTGTGTATTTCTATATGAGGTGGTTTCAGGGTCCATAAAGC
1965 (ioc2)   AGTGAAATACTTAACAATTGAGCTG
1966 (ioc2)   GCTCCTACATTTCTTGTACTCCT

Table S2. Yeast strains used in this study. All are W303 *RAD5* background (***Zhao et al., 1998***).

Strain	Genotype	Source
YDC111	*MATa ade2-1 can1-100 leu2-3,112 trp1-1 ura3-1*	Kim 2006
YTT186	*MATa ade2-1 can1-100 his3-11,15 leu2-3,112 trp1-1 ura3-1 isw1Δ::ADE2*	Tsukiyama 1999
YTT196	*MATα ade2-1 can1-100 his3-11,15 leu2-3,112 trp1-1 ura3-1 isw2Δ::LEU2*	Tsukiyama 1999
YTT645	*MATa ade2-1 can1-100 his3-11,15 leu2-3,112 trp1-1 ura3-1 ioc3Δ::KanMX*	Tsukiyama Lab (unpub.)
YTT827	*MATa ade2-1 can1-100 his3-11,15 leu2-3,112 trp1-1 ura3-1 ioc4Δ::HPH1*	Tsukiyama Lab (unpub.)
YTT1986	*MATα ade2-1 can1-100 his3-11,15 leu2-3,112 trp1-1 ura3-1 set1Δ::NAT1*	Tsukiyama Lab (unpub.)
YJO482	*MATa ade2-1 can1-100 his3-11,15 leu2-3,112 trp1-1 ura3-1 chd1Δ::HIS3MX6*	Ocampo 2016
YJO484	*MATa ade2-1 can1-100 his3-11,15 leu2-3,112 trp1-1 ura3-1 isw1Δ::ADE2 chd1Δ::HIS3MX6*	Ocampo 2016
YJO486	*MATa ade2-1 can1-100 his3-11,15 leu2-3,112 trp1-1 ura3-1 isw2Δ::LEU2 chd1Δ::HIS3MX6*	Ocampo 2016
YJO487	*MATα ade2-1 can1-100 his3-11,15 leu2-3,112 trp1-1 ura3-1 isw2Δ::LEU2 chd1Δ::HIS3MX6*	This study
YJO505	*MATa/α ade2-1/ade2-1 can1-100/can1-100 his3-11,15/his3-11,15 leu2-3,112/leu2-3,112 trp1-1/trp1-1 ura3-1/ura3-1 ISW1/isw1Δ::ADE2 ISW2/isw2Δ::LEU2 RSC8/GALp-RSC8::KanMX CHD1/chd1Δ::HIS3MX6*	Ocampo 2016
YPE600	*MATa/α ade2-1/ade2-1 can1-100/can1-100 his3-11,15/HIS3 leu2-3,112/leu2-3,112 trp1-1/ trp1-1 ura3-1/ura3-1 isw2Δ::LEU2*	This study

792	YPE606	*MATa ade2-1 can1-100 leu2-3,112 trp1-1 ura3-1 ioc3Δ::KanMX*	This study
793	YPE607	*MATα ade2-1 can1-100 leu2-3,112 trp1-1 ura3-1 ioc3Δ::KanMX*	This study
794	YPE608	*MATa ade2-1 can1-100 leu2-3,112 trp1-1 ura3-1 ioc4Δ::HPH1*	This study
795	YPE636	*MATa ade2-1 can1-100 leu2-3,112 trp1-1 ura3-1 ioc2Δ::URA3*	This study
796	YPE654	*MATa ade2-1 can1-100 leu2-3,112 trp1-1 ura3-1 ioc2Δ::URA3 ioc3Δ::KanMX*	This study
797	YPE655	*MATa ade2-1 can1-100 leu2-3,112 trp1-1 ura3-1 ioc2Δ::URA3 ioc4Δ::HPH1*	This study
798	YPE657	*MATa ade2-1 can1-100 leu2-3,112 trp1-1 ura3-1 ioc2Δ::URA3 ioc3Δ::KanMX*	This study
799		*ioc4Δ::HPH1*	
800	YPE712	*MATa ade2-1 can1-100 his3-11,15 leu2-3,112 trp1-1 ura3-1 chd1Δ::HIS3MX6*	This study
801		*ioc3Δ::KanMX6*	
802	YPE715	*MATa ade2-1 can1-100 his3-11,15 leu2-3,112 trp1-1 ura3-1 chd1Δ::HIS3MX6*	This study
803		*ioc2Δ::URA3 ioc4Δ::HPH1*	
804	YPE602	*MATa ade2-1 can1-100 leu2-3,112 trp1-1 ura3-1 set1Δ::NAT1*	This study
805	YPE604	*MATa ade2-1 can1-100 leu2-3,112 trp1-1 ura3-1 set2Δ::TRP1*	This study
806	ROY1281	*MATα lys2 trp1 his3 leu2 ura3 hhf1-hht1Δ::LEU2 hhf2-hht2Δ::HIS3 pCC67*	Dhillon 2000
807	YDC507	*MATα lys2 trp1 his3 leu2 ura3 hhf1-hht1Δ::LEU2 hhf2-hht2Δ::HIS3 p730*	This study
808	YDC101	*MATα lys2 trp1 his3 leu2 ura3 hhf1-hht1Δ::LEU2 hhf2-hht2Δ::HIS3 p368*	This study

Supplementary References

Dhillon, N., Kamakaka, R.T., 2000. A histone variant, Htz1p, and a Sir1p-like protein, Esc2p, mediate
silencing at HMR. Mol. Cell 6, 769–780. doi:10.1016/s1097-2765(00)00076-9

Kim, Y., McLaughlin, N., Lindstrom, K., Tsukiyama, T., Clark, D.J., 2006. Activation of Saccharomyces
cerevisiae HIS3 results in Gcn4p-dependent, SWI/SNF-dependent mobilization of nucleosomes over
the entire gene. Mol. Cell. Biol. 26, 8607–8622. doi:10.1128/MCB.00678-06

Ocampo, J., Chereji, R.V., Eriksson, P.R., Clark, D.J., 2016. The ISW1 and CHD1 ATP-dependent chromatin
remodelers compete to set nucleosome spacing in vivo. Nucleic Acids Res. 44, 4625–4635.
doi:10.1093/nar/gkw068

Tsukiyama, T., Palmer, J., Landel, C.C., Shiloach, J., Wu, C., 1999. Characterization of the imitation switch
subfamily of ATP-dependent chromatin-remodeling factors in Saccharomyces cerevisiae. Genes
Dev. 13, 686–697. doi:10.1101/gad.13.6.686

Yen, K., Vinayachandran, V., Batta, K., Koerber, R.T., Pugh, B.F., 2012. Genome-wide nucleosome
specificity and directionality of chromatin remodelers. Cell 149, 1461–1473.
doi:10.1016/j.cell.2012.04.036

Zhao, X., Muller, E.G., Rothstein, R., 1998. A suppressor of two essential checkpoint genes identifies a
novel protein that negatively affects dNTP pools. Mol. Cell 2, 329–340. doi:10.1016/s1097-
2765(00)80277-4