Sampling SARS-CoV-2 proteomes for predicted CD8 T-cell epitopes

as a tool for understanding immunogenic breadth and rational

vaccine design

Jonathan Hare[1*], David Morrison[2] and Morten Nielsen[3]

[1]IAVI, 125 Broad Street, New York, New York, USA

[2]Bitefirst, South Walsham, Norfolk, UK

[3]Department of Health Technology, Technical University of Denmark, Lyngby, Denmark

*Correspondence. Corresponding author jhare@iavi.org

Keywords: SARS-CoV-2, CD8 T-cells, peptides, T-cell vaccine, predicted T-cell epitopes

Abstract

Predictive models for vaccine design have become a powerful and necessary resource for the

expeditiousness design of vaccines to combat the ongoing SARS-CoV-2 global pandemic. Here

we use the power of these predicted models to assess the sequence diversity of circulating

SARS-CoV-2 proteomes in the context of an individual's CD8 T-cell immune repertoire to

identify potential. defined regions of immunogenicity. Using this approach of expedited and

rational CD8 T-cell vaccine design, it may be possible to develop a therapeutic vaccine candidate

with the potential for both global and local coverage.

Introduction

The COVID-19 pandemic is a worldwide health emergency. The first cases were believed to

have occurred earlier than December 2019, and as of the middle of September there have been

more than 31,000,000 cases of SARS-CoV-2 infection have been reported worldwide and in

excess of 965,000 deaths (Update, ecdc.2020). Vaccine development against SARS-CoV-2 is

accelerating, and currently there are more than 130 vaccines in development with some

progressing to clinical trials in man (Mullard, 2020). Of these vaccines the majority are for a

prophylactic indication with the spike glycoprotein being the preferred antigen target.

The full repertoire of immune responses to COVID-19 in patients is still being evaluated but

recent publications indicate a significant role for cell mediated immunity in clearing SARS-CoV-

2 and conferring some level of protective immunity (Grifoni, Weiskopf, Ramirez, Mateus, et al.,

2020). This mirrors observations made for other coronavirus infections, SARS (Li et al., 2008),

MERS (Zhao et al., 2017) and other viral infections including Ebola (Sakabe et al., 2018) and

Lassa virus (Sullivan et al., 2020). Concurrent with vaccine design and understanding the

immune response in patients, significant efforts have been invested in understanding the virus

phylogeny as it progresses around the globe to identify key sequence changes that may influence

vaccine design. The data from this analysis appears to show that the majority of circulating

mutations are neutral or deleterious (van Dorp et al., 2020), although a D614G mutation

identified in the spike glycoprotein may cause an increase in susceptibility to infection (Korber et

al., 2020)

Linking novel sequence data to the use of existing predictive *in silico* tools for epitope

identification offers an intriguing approach that can complement existing vaccine design

strategies. Previously we hypothesized that incorporating potential immune recognition

information into established models may increase the likelihood of success. We have shown that

within a population, although HLA sequences show high levels of polymorphism, there are

conserved, over-represented alleles that can be used as representative of larger allele diversity

(Buggert et al., 2012; Hare, Fiore-Gartland, et al., 2020)

Previously we have applied NetMHCpan (Nielsen & Andreatta, 2016) as a proxy to identify

putative CD8 T-cell epitopes contained within the HIV transmitted founder virus (TFV)

identified from the Protocol C clinical cohort of sub Saharan and East Africa. We have shown

that it is possible to stratify and rank protein and/or proteome sequences for their contributions of

potential T-cell epitopes (Ed McGowan et al., 2020) . Here we propose to use the same approach

to evaluate a subset of global circulating SARS-CoV-2 sequences and, using the predefined

analysis applied to modeling HIV diversity, identify key regions within the SARS-CoV-2

proteome that could be included within a therapeutic T-cell vaccine.

Methods

For genes from each SARS-CoV-2 virus proteome sequence all 8-11mer peptides were

generated. The binding affinity of each peptide to the HLA alleles described above was predicted

using NetMHCpan-4.1.

Binding predictions below the peptide conservation threshold were read into PostgreSQLR for

analysis. The evaluated set of predicted binders was determined as peptides that appear in any

virus proteomes, above the peptide conservation threshold of 2.1%. First the virus proteome with

the largest number of unique predicted binders was identified. Next, the proteome that, when

combined with the previously selected proteome gave the highest increase in coverage of all

predicted peptide binders was included. This iterative process was repeated until 100% of all

predicted T-cell epitopes were accounted for. For comparison, set-building was performed a

second time using randomly selected proteomes instead of choosing the proteome that resulted in

the greatest increase of peptide coverage. For more information go to dataspace.iavi.org

Results

Our approach uses NetMHCpan to predict the HLA/peptide binding affinity for the selected

virus strains and then identify a set of close binding peptides that are common across the strains.

This algorithm was used to analyze 287 SARS-CoV-2 proteome sequences collected between

and 08 January and 02 April 2020 and downloaded from NCBI SARS-CoV-2 virus database.

The model parameters assessed included a minimum peptide conservation threshold of 2.1% (as

previously defined ((Ed McGowan et al., 2020)) and a rank binding cut-off of 2.0. Figure 1

summarizes the output dataset which includes only those peptides that are present in 6 or more

virus proteomes (epitope frequency) and have low rank binding scores although they may

represent multiple HLA/peptide interactions (see Table 1 for input sample data and full model

parameters).

These frequency and binding thresholds identified 11267 unique SARS-CoV-2-specific

predicted CD8 T-cell epitopes and identified a range of predicted binding profiles for the

different peptide-HLA interactions with the majority of peptides associating to selected HLA-A

and HLA-B alleles (Figure 1A & B -HLA Ids 1-36) and few primary associations to HLA-C

alleles.

These predicted peptides can be used to assess diversity by assigning a coverage gain value to

each sequence. These values can then be used to rank each virus proteome for the coverage it

provides within the sample population and by extension identify the sequences that are necessary

to obtain the optimum level of epitope restricted sequence coverage (Figure 2).

This model may be used to target and prioritize individual proteomes from which vaccine targets

could be derived. In our population of 287 SARS-CoV-2 proteomes each proteome offers an

individual coverage gain of 72.2- 98.3%. Using this model 8 key proteomes are required to reach

100% epitope coverage within the sample population. It would require an 8-fold increase in

virus sequences to achieve 100% coverage if sequences had been selected at random (n=68

p<0.0001).

Interestingly, of the 11267 unique predicted CD8 T-cell epitopes, 409 are contained in all 287

proteomes. These peptides map to 4 genes within the proteome with 392 peptides contained in

ORF1ab polypeptide and spike glycoprotein (Figure 3 and Supplementary Table 1).

Discussion

Evaluating predicted CD8 T-cell epitopes within the SARS-CoV-2 proteome revealed a high

level of conservation between proteomes, with each proteome representing 72 - 98.3% of all

predicted T-cell epitopes; however, using our model it is possible to reach 100% epitope

coverage using 8 defined proteomes. This is in contrast to HIV where it would take 83 proteomes

to achieve 100% epitope coverage on a comparable sample size (Ed McGowan et al., 2020) . The

enhanced conservation of epitopes is in concordance with the observed sequence diversity to

date and indicates that no bias has been introduced in to the analysis through modeling with

reduced input data.

Moreover, the elevated conservation can be used as a guide to identify regions within the

proteome that should be included within a therapeutic T-cell vaccine. Recent data has indicated

that naturally occurring T-cell responses in convalescent COVID-19 patients preferentially target

the spike glycoprotein and ORF1a polypeptide (Dong et al., 2020; Grifoni, Weiskopf, Ramirez,

Smith, et al., 2020). However, limitations in both these studies including that ORF1ab peptides

were not examined in one study, the sample size in both was small, peptides were generated from

the reference strain and the HLA Class 1 distribution was limited, may mean that potential

regions of interest have been overlooked.

We identified 409 predicted CD8 peptides that have 100% conservation within our sample set

with >90% of the predicted epitopes contained within either the ORF1ab polypeptide or spike

glycoprotein. Furthermore, the predicted epitopes appear to cluster within a ~550 a.a region of

the spike glycoprotein (amino acid 319-865) and within 2 regions of the ORF1ab polypeptide

(amino acid positions 2750-3250 and 4500-5500). Future experimental testing of these epitopes

would confirm whether natural infection induces CD8 T-cell responses targeting these regions,

but from an *in silico* perspective they offer a potential target for developing a therapeutic T-cell

vaccine that warrants further investigation.

Parameter	Values
SARS-COV-2 Proteomes	Brazil (1), China (13), France (1), Greece (1), Iran (1), South Africa (1), Spain (10), Sweden (1), Taiwan (1), Turkey (1), USA (238), Unknown (16), Vietnam (2)
Binding Threshold	1%
HLA allele contributions	46 alleles (16 x HLA-A, 19 x HLA-B, 11 x HLA-C)
HLA haplotype weighting	None
Rank Binding	<2.0
Peptide Conservation threshold (%)	2.1%
Peptide length	8, 9, 10 & 11mers

120   Table 1 Model Parameters and proteome sources

Figure 1. Affinity plots for all predicted peptides with conservation of ≥2.1% (n=11267). A –
Predicted peptide affinity (Rank Binding) versus peptide frequency within transmitted founder
proteome. B-Predicted peptide frequency versus primary associated HLA. C – Predicted peptide
affinity (Rank Binding) versus primary associated HLA

Figure 2. Cumulative coverage distribution plots of full length transmitted founder gag
sequences using a 3-select coverage model and a 2% Binding Threshold, 3-Select best (red)) and
3-Select random (blue).

Protein	No. Predicted peptides	Gene length (aa)
ORF1ab	352	7097
Spike	40	1274
ORF3a	0	276
Envelope	1	76
Matrix	0	223
ORF6	0	63
ORF7a	0	123
ORF7b	0	44
ORF8	0	64
Nucleocapsid	16	422
ORF10	0	39

Figure 3. Mapping of predicted epitopes with 100% conservation showing location and number of predicted responses across the proteome and cumulative number of precited epitopes per 250aa. A – Predicted responses mapped across the ORF1ab polyprotein (with NSP genes identified). B - Predicted responses mapped across the spike glycoprotein.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or

financial relationships that could be construed as a potential conflict of interest.

Author Contributions

JH wrote the manuscript and provided conceptual input. DM and MN developed and

implemented the analysis scripts and contributed to developing the manuscript.

Acknowledgements

This work was funded in part by IAVI and made possible by the support of the United States

Agency for International Development (USAID) and other donors. The full list of IAVI donors is

available at http://www.iavi.org. The contents of this manuscript are the responsibility of IAVI

and do not necessarily reflect the views of USAID or the US Government. We would also like to

thank Thiru Thangarajah (Genscript Inc.) for insight in to potential peptide strategies. This

publication was kindly accepted for pre-print publication by BioRxiv (Hare, Morrison, &

Nielsen, 2020)

References

Buggert, M., Norström, M. M., Czarnecki, C., Tupin, E., Luo, M., Gyllensten, K., … Karlsson,

A. C. (2012). Characterization of HIV-specific CD4+ T cell responses against peptides

selected with broad population and pathogen coverage. *PLoS ONE*, *7*(7).

https://doi.org/10.1371/journal.pone.0039874

Dong, T., Peng, Y., Mentzer, A. J., Liu, G., Yao, X., Yin, Z., … Goulder, P. (2020). Broad and

strong memory CD4+ and CD8+ T cells induced by SARS-CoV-2 in UK convalescent

COVID-19 patients. *BioRxiv*, 2020.06.05.134551.

https://doi.org/10.1101/2020.06.05.134551

Grifoni, A., Weiskopf, D., Ramirez, S. I., Mateus, J., Dan, J. M., Rydyznski Moderbacher, C., …

Silva, de. (2020). Journal Pre-proof Targets of T cell responses to SARS-CoV-2

coronavirus in humans with COVID-19 disease and unexposed individuals. *Cell*.

https://doi.org/10.1016/j.cell.2020.05.015

Grifoni, A., Weiskopf, D., Ramirez, S. I., Smith, D. M., Crotty, S., & Sette, A. (2020). Targets of

165 T Cell Responses to SARS-CoV-2 Coronavirus in Humans with COVID-19 Disease and

166 Unexposed Individuals. https://doi.org/10.1016/j.cell.2020.05.015

167 Hare, J., Fiore-Gartland, A., Mcgowan, E., Rosenthal, R., Hunter, E., Gilmour, J., & Nielsen, M.

168 (2020). Selective HLA restriction permits the evaluation and interpretation of immunogenic

169 breadth at comparable levels to autologous HLA.

170 https://doi.org/10.20944/preprints202008.0467.v1

171 Hare, J., Morrison, D., & Nielsen, M. (2020). Sampling SARS-CoV-2 proteomes for predicted

172 CD8 T-cell epitopes as a tool for understanding immunogenic breadth and rationale 2

173 vaccine design. *BioRxiv*, 2020.08.15.250647. https://doi.org/10.1101/2020.08.15.250647

174 Korber, B., Fischer, W., Gnanakaran, S. G., Yoon, H., Theiler, J., Abfalterer, W., … Group, S.

175 C.-19 G. (2020). Spike mutation pipeline reveals the emergence of a more transmissible

176 form of SARS-CoV-2. *BioRxiv*, 2020.04.29.069054.

177 https://doi.org/10.1101/2020.04.29.069054

178 Li, C. K., Wu, H., Yan, H., Ma, S., Wang, L., Zhang, M., … Xu, X.-N. (2008). T Cell Responses

179 to Whole SARS Coronavirus in Humans. *The Journal of Immunology*, *181*(8), 5490–5500.

180 https://doi.org/10.4049/jimmunol.181.8.5490

181 McGowan, Ed, Rosenthal, R., Fiore-Gartland, A., Macharia, G., Balinda, S., Kapaata, A., …

182 Hare, J. (2020). Utilizing Computational Machine Learning Tools to Understand

183 Immunogenic Breadth in the Context of a CD8 T-Cell Mediated 2 HIV Response 3.

184 *BioRxiv*, 2020.08.15.250589. https://doi.org/10.1101/2020.08.15.250589

185 McGowan, Edward, Rosenthal, R., Fiore-Gartland, A., Dalel, J., Streatfield, C., Coutinho, H., …

186 Hare, J. (n.d.). Utilizing Computational Machine Learning Tools to Understand

Immunogenic Breadth in the Context of a CD8 T-Cell Mediated HIV Response. *Submitted*.

Mullard, A. (2020). *COVID-19 vaccine development pipeline gears up*.

https://doi.org/10.1016/S0140-6736(20)31252-6

Nielsen, M., & Andreatta, M. (2016). NetMHCpan-3.0; improved prediction of binding to MHC

class I molecules integrating information from multiple receptor and peptide length datasets.

Genome Medicine, *8*(1), 33. https://doi.org/10.1186/s13073-016-0288-x

Sakabe, S., Sullivan, B. M., Hartnett, J. N., Robles-Sikisaka, R., Gangavarapu, K., Cubitt, B., …

Oldstone, M. B. A. (2018). Analysis of CD8+ T cell response during the 2013–2016 Ebola

epidemic in West Africa. *Proceedings of the National Academy of Sciences of the United*

States of America, *115*(32), E7578–E7586. https://doi.org/10.1073/pnas.1806200115

Sullivan, B. M., Sakabe, S., Hartnett, J. N., Ngo, N., Goba, A., Momoh, M., … Oldstone, M. B.

A. (2020). High crossreactivity of human T cell responses between Lassa virus lineages.

PLOS Pathogens, *16*(3), e1008352. https://doi.org/10.1371/journal.ppat.1008352

Update, E. (n.d.). COVID-19 situation update worldwide, as of 2 August 2020. Retrieved August

3, 2020, from https://www.ecdc.europa.eu/en/geographical-distribution-2019-ncov-cases

van Dorp, L., Acman, M., Richard, D., Shaw, L. P., Ford, C. E., Ormond, L., … Balloux, F.

(2020). Emergence of genomic diversity and recurrent mutations in SARS-CoV-2.

Infection, Genetics and Evolution, *83*, 104351.

https://doi.org/10.1016/j.meegid.2020.104351

Zhao, J., Alshukairi, A. N., Baharoon, S. A., Ahmed, W. A., Bokhari, A. A., Nehdi, A. M., …

Zhao, J. (2017). Recovery from the Middle East respiratory syndrome is associated with

208   antibody and T cell responses. *Science Immunology*, *2*(14).

209   https://doi.org/10.1126/sciimmunol.aan5393

211