1 **Bixafen, a succinate dehydrogenase inhibitor fungicide, causes microcephaly and**

2 **motor neuron axon defects during development**

4 Rahma Hassan-Abdi, Alexandre Brenet, and Nadia Soussi-Yanicostas*

7 Université de Paris, NeuroDiderot, Inserm, F-75019 Paris, France

8 **(*)Correspondence:** nadia.soussi@inserm.fr; Tel.: +00-33140-03-1931 (France)

14 **Running title:** Neurodevelopmental defects in bixafen-exposed embryos

20 **Keywords**: Succinate dehydrogenase inhibitors, SDHI, Fungicide, Zebrafish,

21 Neurodevelopment, *In vivo* imaging

Abstract

Succinate dehydrogenase inhibitors (SDHIs) are the most widely used fungicides against plant parasitic fungi. They act by blocking the enzyme succinate dehydrogenase (SDH), a key component of mitochondrial respiration, which is highly conserved throughout evolution. Recently, it has been reported that some SDHIs used in many fungicides do not only inhibit the SDH activity of target fungi but can also block several non-target human cells in *in vitro* models. This study reveals a lack of SDHI species specificity and so points to a major health risk for exposed organisms, including humans. Despite the frequent detection of SDHIs in the environment and on harvested products and their increasing use in modern agriculture, their potential toxic effects *in vivo*, especially on neurodevelopment, are still under-evaluated. Here, we assessed the neurotoxicity of bixafen, one of the latest-generation SDHIs, which had never been tested during neurodevelopment. For this purpose, we used a well-known vertebrate model for toxicity testing, namely zebrafish transparent embryos, and live imaging using transgenic lines labelling the brain and spinal cord. Here we show that bixafen causes microcephaly and defects on motor neuron axon outgrowth and their branching during development. Our findings show that the central nervous system is highly sensitive to bixafen, thus demonstrating for the first time *in vivo* that an SDHI, bixafen, widely used in agriculture, is neurotoxic in vertebrates and causes neurodevelopmental defects. This work adds to our knowledge of the toxic effect of SDHIs on neurodevelopment and may help us take the appropriate precautions to ensure protection against the neurotoxic effects of these substances

Introduction

Bixafen, a methyl-pyrazole carboxamide, is a fungicide widely used on cereal and rapeseed crops. It was initially approved and launched in the market in 2011 by Bayer (Ravichandra, 2018).Its high efficiency and fast penetration resulted in a significant rise in usage in Europe and the US. Currently, nearly thirty different products are authorized and present on the French market alone that contain bixafen as sole active substance or mixed with other fungicides. Bixafen belongs to the succinate dehydrogenase inhibitor (SDHI) family, the most widely used fungicides in agriculture as a treatment against a broad range of fungal diseases. It is categorized as one of the latest-generation SDHIs derived from carboxin. Bixafen, like all the latest generation SDHIs, acts in a similar way to carboxin, through inhibition of mitochondrial respiration chain complex II, also known as succinate dehydrogenase (SDH). The complex II targeted by SDHI is not specific to the targeted disease fungi but is also closely similar to that present in all plant and animals species, including humans. SDHI inhibitory action on mitochondrial activity has thus been shown to be effective in various models and human cells (Bénit et al., 2019). Additionally, these latest generation SDHIs have been shown to also inhibit mitochondrial respiration complex III (Bénit et al., 2019), suggesting more critical effects of these new fungicides than those expected. SDH activity has been shown to be irreplaceable in mitochondrial metabolism, and defects in its activity impair the cellular metabolome and functions (Bénit et al., 2014). A broad range of human diseases from neurodegeneration to cancer are related to SDH complex defects. Several studies have shown that SDH complex mutations are associated with several diseases, such as Leigh syndrome, an early-onset progressive neurodegenerative disease characterized by developmental delay, ataxia, seizures, and defects in the brain and spinal cord (Birch-Machin et al., 2000; Bourgeron, 2015; Finsterer, 2008; Horváth et al., 2006; Parfait et al., 2000; Van Coster et al., 2003).

SDH mutations have also been associated with several tumors, familial paraganglioma

syndrome, infantile leukoencephalopathy, thyroid and renal cancers, and neuroblastoma

(Ghezzi et al., 2009; Martin et al., 2007; Perry et al., 2006; Ricketts et al., 2009; Stratakis

and Carney, 2009; Timmers et al., 2009). One study showed that the SDHI fungicide

isopyrazam increased uterine tumor formation in rats (Yoshida et al., 2015). The

Norwegian scientific committee for food safety (VKM) has assessed the health and

environmental risk of Aviator Xpro EC225, a fungicide containing bixafen, and concluded

that the effects of bixafen in animals should be considered relevant for humans (VKM

Report, 2014). Also, M44, a bixafen metabolite, which has the potential to contaminate

groundwater, causes abnormalities in the rabbit fetus, suggesting that similar effects in

humans cannot be excluded (VKM Report, 2015).

Despite the frequent detection of SDHIs in the environment and on harvested products

(Abad-Fuentes et al., 2015; Añasco et al., 2010; Tanabe and Kawata, 2009; Tsuda et al.,

2009; Vu et al., 2016) and their increasing use in modern agriculture, their potential toxic

effects, especially on neurodevelopment, are still under-evaluated. Hence the need to

assess the neurotoxicity of these SDHI fungicides *in vivo* on neurodevelopment. In this

study, we tested the potential toxic effect of one of the latest-generation SDHIs, bixafen, in

zebrafish embryos. We observed that embryos exposed to bixafen developed a series of

defects during growth, including microcephaly and major defects in spinal cord motor

neuron axons and their branching, showing the toxicity of this SDHI on central nervous

system neurodevelopment. Our study gives insight into the environmental risk of bixafen

on neurodevelopment.

Results

1. Toxicity of bixafen toward zebrafish embryos

To assess the potential toxicity of bixafen, a succinate dehydrogenase inhibitor belonging to the pyrazole group of fungicides, 6 hours post-fertilization (hpf) zebrafish embryos were treated with increasing doses of bixafen from 0.1 μM to 1 μM. As there were no differences in survival percentage or embryo morphology between the 0.1% DMSO group and the fish water control group (data not shown), we used 0.1% DMSO for the control group. Exposure of zebrafish embryos to bixafen resulted in significant developmental abnormalities and mortality compared with the DMSO control. Mortality showed a concentration-response effect, with an increasing mortality rate with bixafen exposure concentration. The median lethal concentration (LC50) of bixafen for zebrafish embryos, causing a 50% mortality rate at 4 days post-fertilization (dpf), was calculated at 0.5 μM. The surviving embryos treated with 0.5 μM of bixafen displayed a curved tail (Figure 1E, red arrowhead) causing body shortening (Figure 1F, 33.8 ± 1.5 mm); and edemas around the heart, eye and head (Figure 1E, yellow arrowheads) causing a swollen head (Figure 1G, 38.1 ± 0.8 mm), compared to the DMSO control embryos (Figure 1C) body (Figure 1F, 37.2 ± 0.6 mm) and head (Figure 1G, 5.8 ± 0.2 mm).

We then determined the lowest-observed-adverse-effect-level (LOAEL) concentration of bixafen (0.3 μM), i.e. the lowest exposure level at which we observed the first noticeable body deformity observed in most bixafen exposed embryos, which was a slight edema around the heart. Embryos treated at the LOAEL (0.3 μM) displayed only a modest-size pericardial edema (Figure 1D, yellow arrowhead) but no significant difference in body size (Figure 1F, 35.7 ± 0.7 mm *vs.* 37.2 ± 0.6 mm) or head size (Figure 1G, 5.5 ± 0.4 mm *vs.* 5.8 ± 0.2 mm) after body and head measurements (Figure 1B). In this study, we used two

transgenic lines Tg(huC:G/U:RFP) and Tg(Olig2:eGFP) to analyze the embryos exposed to

LOAEL and LC50 concentrations and compare the resulting defects.

2. Brain defects in embryos exposed to bixafen

Several studies have reported that a number of SDHIs cause neurotoxicity (Wang et al.,

2020; Yao et al., 2018). We investigated the effect of bixafen exposure on the development

of the central nervous system. To assess the effect of bixafen on brain development

reliably under a fluorescence microscope, we used *elavl3:Gal4/5xUAS:RFP* double

transgenic embryos (hereafter called Tg(huC:G/U:RFP)), in which red fluorescence

protein (RFP) is expressed in postmitotic neurons, thus clearly delineating central

nervous system boundaries. We treated 6 hpf embryos from Tg(huC:G/U:RFP) transgenic

line with the determined LOAEL (0.3 µM) and LC50 (0.5 µM) concentrations, and imaged

live brain morphology at 30 hpf and 60 hpf (Figure 1A, N). Live imaging of embryo brains

treated at 30 hpf and 60 hpf showed a smaller brain structure (Figure 1H-M). These

observations were confirmed by quantifications of brain volume after three-dimensional

reconstruction of embryo brains based on RFP positive neurons (Imaris software,

Bitplane) at 30 hpf (Figure 1O; DMSO: $3.50 \pm 0.18 \times 10^6$ µm^3 , Bixafen 0.3µM: 2.06 ± 0.17

$\times 10^6$ µm^3, Bixafen 0.5µM: $1.43 \pm 0.23 \times 10^6$ µm^3) and 60 hpf (Figure 1P; DMSO: $15.37 \pm$

0.44×10^6 µm^3 , Bixafen 0.3µM: $10.35 \pm 0.52 \times 10^6$ µm^3, Bixafen 0.5µM: $5.66 \pm 0.73 \times 10^6$

µm^3). We also observed an absence of photoreceptors in the retina of embryos exposed

to the LC50 (Figure 1J, M, asterisks). Brain volume measurements were significantly

reduced in embryos treated with bixafen at LOAEL (Figure 1 I, L). This defect was

worsened at LC50 (Figure 1 J, M) compared to controls (Figure 1 H, K), showing that the

toxicity of bixafen led to microcephaly.

3. Defective spinal motor neuron axon outgrowth in embryos exposed to

bixafen

We found that more than 80% of embryos treated with LC50 displayed behavior defects

in a touch-response test (data not shown) compared to DMSO controls. We therefore

investigated whether these motor defects were due to abnormal development of motor

neurons or impaired pathfinding of their axons. To determine whether the spinal motor

neuron axons were affected in bixafen-treated embryos, we used transgenic line

Tg(Olig2:eGFP), in which eGFP is expressed in motor neuron progenitor cells, labeling

motor axons. Live embryos treated or not with bixafen were imaged by confocal

microscopy at 30 hpf and 60 hpf. We specifically analyzed caudal primary motor neuron

(CaPMN) axons, which are the most accessible (Figure 2A, H). At 30 hpf, in control

embryos and LOAEL treated embryos, 100% of spinal motor neuron axons extended

ventrally to contact their muscle targets (Figures 2B, C). By contrast, in LC50 treated

embryos, at 30 hpf, the spinal neuron axons exited the spinal cord and showed a severe

delay in outgrowth (Figure 2D). Length measurements of three-dimensionally

reconstructed CaPMN axons confirmed that these axons were significantly shorter in

embryos treated with bixafen at LOAEL and that this effect was worsened in embryos

exposed to bixafen LC50 concentrations (Figure 2I). At 60 hpf, in control embryos, the

CaPMN axons followed a stereotyped pathway, extending ventrally in the space between

the notochord and the myotome (Figure 2E, E'). At the ventral edge of the musculature,

each axon turned dorsally and laterally to grow along the dorsal myoseptum. However, in

bixafen-treated embryos at 60 hpf, axons showed a defective phenotype as determined by

their branching pattern at the choice point and/or the horizontal myoseptum (Figures 2

F, G, F', G'). At LC50, CaPMN axons extended aberrant branches projecting in all directions

instead of extending ventrally within the myotome (Figure 2G'). Hence treatment with

bixafen disorganized motor axon growth at LOAEL concentration and this effect was

worsened at the LC50 concentration. Length measurements of three-dimensionally

reconstructed CaPMN axons confirmed this observation, with significantly shorter axons

in embryos treated with bixafen at 30 hpf (Figure 1I; DMSO: 172.2 ± 3.03 μm, Bixafen

0.3μM: 97.1 ± 3.56 μm, Bixafen 0.5μM: 46.2 ± 2.3 μm) and 60 hpf (Figure 1J; DMSO: 406.3

± 7.65 μm, Bixafen 0.3μM: 272.9 ± 9.07 μm, Bixafen 0.5μM: 171.3 ± 11.52 μm). These data

show that exposure to bixafen has a drastic effect on the outgrowth of the spinal motor

axons.

Discussion

The developing central nervous system is especially vulnerable and sensitive to exposure to toxic chemicals in the environment. Succinate dehydrogenase inhibitors (SDHIs) are widely used as fungicides to control a broad range of fungal diseases, but it was not known whether these SDHIs had a toxic effect on the developing central nervous system. In this study we report evidence that exposure to an SDHI, bixafen, induces a series of neurodevelopmental defects.

In this study we used the zebrafish, a valuable vertebrate system for environmental health research. It enables real-time *in vivo* studies that address potential health hazards to human embryos, and in particular studies investigating neurodevelopment in a vertebrate model (Lein et al., 2005). Regarding brain development, we found that bixafen exposure at the median lethal concentration (LC50) caused major malformation phenotypes, including eye and brain defects leading to microcephaly. We also observed a pericardial edema, indicating that bixafen exerts cardiac toxicity. We also found that embryos treated with the lowest-observed-adverse-effect-level (LOAEL) concentration of bixafen showed a very discreet morphological phenotype sufficient to induce significant defects in brain size, indicating that the sublethal dose of bixafen could influence normal development. Microcephaly has been found in embryos exposed to many fungicides, including Maxim® XL (Svartz et al., 2016), and glyphosate (Paganelli et al., 2010). Moreover, bixafen-exposed embryos displayed absence of photoreceptors in the retina. All these findings suggest that the brain is very vulnerable to many toxic substances even at low concentrations. A number of studies have shown that prenatal and early childhood exposure to some pesticides is associated with neurodevelopmental and cognitive disorders (Muñoz-Quezada et al., 2013). Perinatal exposure of rats to organophosphorus pesticides alters brain morphometry (Veronesi and Pope, 1990). Similarly, our recent

studies have shown that embryos exposed to organophosphorus pesticides showed neurodevelopmental disorders, including altered excitatory/inhibitory synapse balance (Brenet et al., 2019).

Regarding the spinal cord, our data show that embryos exposed to bixafen show deficits in touch-induced swimming behavior and display a typical low-activity swimming behavior. As it is known that the motor neuron is a major cell type regulating swimming behavior in zebrafish during early life (Brustein et al., 2003), we investigated the development of the axons of the motor neurons in embryos exposed to bixafen. In the zebrafish embryos, there are two different kinds of spinal motor neurons, called primary and secondary motor neurons (Myers, 1985; Myers et al., 1986). The caudal primary motor neurons (CaPMN) have a soma located in the middle of each spinal cord and axons extending along a ventral pathway to innervate ventral axial muscle (Westerfield et al., 1986). In this study, we focused on the CaPMN axons because of their easy observation and distinct axon projections. In control embryos exposed to 0.1% DMSO, motor neurons extend their axonal trajectory in a highly stereotyped manner during development. CaPMN axons exit the spinal cord and extend toward the ventral muscle. However, in embryos exposed to bixafen, we observed a reduction in axon outgrowth and later defects in branching of motor axons. Similarly, we found that the low concentration of bixafen (LOAEL) was sufficient to induce defects in axon outgrowth and branching of motor neurons during development, indicating that the development of motor neuron axons is highly sensitive to bixafen. This study reveals motor neuron axon defects in embryos exposed to bixafen. It also provides data on the toxicity of bixafen fungicides to a nontarget organism, zebrafish embryos. It has been shown that chlorpyrifos-oxon pesticide disrupts motor neuron axons (Howard et al., 2005).

Bixafen is a methyl-pyrazole, one of the SDHI fungicides used to manage plant diseases

and inhibit the respiration of pathogenic fungi by blocking succinate complex II in the

mitochondrial respiratory chain. SDH is an enzyme involved in both oxidative

phosphorylation and the tricarboxylic acid cycle, two processes that generate energy. This

enzyme has been shown to be irreplaceable in mitochondrial and cell metabolism and also

to be highly conserved among all fungi, plants and animal species (Rehfus et al., 2016;

Veloukas et al., 2013; Yamashita and Fraaije, 2018). Therefore, any adverse change or

inhibition of SDH activity can lead to many diseases, including those that have been linked

to mitochondrial dysfunction (Birch-Machin et al., 2000; Bourgeron, 2015; Finsterer,

2008; Horváth et al., 2006; Parfait et al., 2000; Van Coster et al., 2003).

A recent study has questioned the specificity of these SDHIs, including bixafen. It showed

that the latest-generation SDHIs sold on the market could inhibit the SDH of several

species ranging from earthworms to human cells, evidence that these SDHIs are

nonspecific (Bénit et al., 2019). Several epidemiological studies published during the last

two decades suggest harmful effects of pesticides on human health (Merhi et al., 2007;

Weichenthal et al., 2010). Pesticide poisoning is a serious health problem that

disproportionately affects infants and children (Rauh et al., 2012). Pesticides are known

to cause millions of acute poisoning cases per year (Bertolote et al., 2006). Human

exposure to pesticides can occur environmentally, through consumption in food and

water (van den Berg et al., 2012). A number of studies show that prenatal and early

childhood exposure to some pesticides is associated with neurodevelopmental and

cognitive disorders (Muñoz-Quezada et al., 2013; Ross et al., 2013). Very little is known

about the consequences of exposure to SDHIs on human health. Currently, we know only

the consequences of the changes in their target, the SDH complex. Mutations in the SDH

complex are associated with several human diseases, including neurodegenerative

diseases and cancer (Ardissone et al., 2015; Birch-Machin et al., 2000; Bourgeron, 2015;

Finsterer, 2008; Horváth et al., 2006; Parfait et al., 2000; Van Coster et al., 2003; Martin et

al., 2007; Perry et al., 2006; Ricketts et al., 2009; Stratakis and Carney, 2009; Timmers et

al., 2009).

The fungicide bixafen exists as a pure compound or as an ingredient in 27 approved and

marketed fungicides. Our results will be used as basic data to study the complex mixtures

that contain bixafen. It is not known how bixafen can affect the development of the central

nervous system .Like for all SDHIs, the toxic effect of bixafen observed in this study may

be linked to the basic disruption of mitochondrial activity and accumulation of reactive

oxygen species caused on Complex II inhibition and subsequent oxidative DNA damage.

These severe brain and spinal cord malformations may also be related to the different

metabolic abnormalities reported in SDHI exposed animals (Graillot et al., 2012; Qian et

al., 2018; Wu et al., 2018; Yang et al., 2018). A recent work (Bénit et al., 2019) has shown

that bixafen also exerts an effect on mitochondrial respiration complex III, suggesting an

adverse effect of bixafen on cellular metabolism.

In summary, our study provides new evidence of bixafen toxicity on neurodevelopment

in a vertebrate model, and and may help us take the appropriate precautions to ensure

protection against the neurotoxic effects of these substances.

Materials and Methods

1. Ethics statement

All the animal experiments described in the present study were conducted at the French National Institute of Health and Medical Research (INSERM) UMR 1141 in Paris in accordance with European Union guidelines for the handling of laboratory animals (http://ec.europa.eu/environment/chemicals/lab_animals/home_en.htm). They were approved by the Direction Départementale de la Protection des Populations de Paris and the French Animal Ethics Committee under reference No. 2012-15/676-0069.

2. Zebrafish lines and maintenance

Zebrafish were maintained at 26.5 °C in 14 h light and 10 h dark cycles. Embryos were collected by natural spawning, and to avoid pigmentation, 0.003% 1-phenyl-2-thiourea (PTU) was added at 1 dpf (day post-fertilization). The following transgenic lines were used: Tg(elavl3:Gal4)[zf349] (Akerboom et al., 2012), Tg(5xUAS:RFP)[nkuasrfp1a] (Asakawa et al., 2008), Tg(Olig2:eGFP) (Shin et al., 2003).

3. Chemicals

Dimethyl sulfoxide (DMSO) hybrid-max sterile (Sigma, D2650) was diluted in fish water to a final concentration of 0.1% (v/v). Bixafen, (N-(3′,4′-dichloro-5-fluorobiphenyl-2-yl)-3-(difluoromethyl)-1-methylpyrazole-4-carboxamide) (Sigma, 32581) was dissolved at a stock concentration of 100 μM in pure DMSO (100%) and conserved at −20 °C in aliquots until use. 100 μM stocks were diluted in a 0.1% DMSO solution in fish water to final concentrations of 0.1 μM, 0.2 μM, 0.3 μM, 0.4 μM, 0.5 μM, 0.6 μM, 0.7 μM, 0.8 μM, 0.9 μM and 1.0 μM to determine lethality and deduce chemical toxicity values.

4. Toxicity assay

Transgenic lines, Tg(huC:G/U:RFP) or Tg (Olig2:eGFP) embryos were collected after spawning and washed twice with clean fish water. Fertilized and normal 6 hpf embryos were selected, distributed randomly in 24-well microplates, 20 embryos per well, following OECD zebrafish embryo toxicological assay directives. Each well was filled with 2 mL of solution for each condition, with 12 conditions in all: fish water, DMSO 0.1%, Bixafen diluted in DMSO 0.1% at final concentrations of 0.1 μM, 0.2 μM, 0.3 μM, 0.4 μM, 0.5 μM, 0.6 μM, 0.7 μM, 0.8 μM, 0.9 μM and 1.0 μM. Each day, dead embryos were selected under OECD guidelines (coagulation, no heartbeat or no somite) and discarded.

5. Embryo imaging

30 hpf or 60 hpf embryos were anesthetized with 112 μg/mL 3-aminobenzoic acid ethyl ester (tricaine, Sigma), immobilized in 1% low melting-point agarose in the center of a 35 mm glass-bottomed dish (Corning), and covered with fish water containing 112 μg/mL tricaine. For embryo morphology analysis, bright field images were captured using a stereomicroscope (Zeiss). Live imaging of transgenic lines Tg(huC:G/U:RFP) and Tg(Olig2:eGFP) was done using a Leica SP8 confocal scanning laser microscope equipped with a Leica 20x/0.75 multi-immersion objective.

6. Image analysis

Body and head measurements from bright field embryo images were made using the ruler tool in ImageJ. Brain volume and spinal neuron CaPMN axon length three-dimensional reconstructions and quantifications were analyzed using Imaris Measurement Pro (Bitplane Inc.).

7. Statistics

All statistics were obtained on Prism5 (GraphPad) and assessed using a Kruskal-Wallis

test followed by a Dunn post-test. All data are represented as means ± SEM.

Author Contributions:

R.H.A. performed the experiments and the analysis, designed the figures, and wrote the

original draft. A.B. performed data analysis. N.S.Y. supervised the project and wrote the

manuscript.

Funding: This work was supported by Institut National de la Santé et la Recherche

Médicale (INSERM), the National Center for Scientific Research (CNRS), and the French

National Research Agency (ANR-16-CE18-0010). Funding sources had no involvement in

study design, collection, analysis or interpretation of data, or decision to publish.

.

Acknowledgments: We thank Christiane Romain (Inserm UMR 1141), Olivier Bar

(Inserm UMR 1141) for their technical assistance.

Conflicts of Interest: The authors declare that the research was conducted in the absence

of any commercial or financial relationships that could be construed as a potential conflict

of interest.

Figure 1: Acute exposure to bixafen causes microcephaly.

(**A**) Illustration of the experimental setup used to analyze body morphology and brain size on acute exposure to bixafen. (**B**) Illustration of the measurements used to assess body size and brain size of embryos. (**C-E**) Images of 4 dpf embryos treated with DMSO (**C**), 0.3 µM (**D**) or 0.5 µM bixafen (**E**). (**D**) Image showing a small pericardial edema at the LOAEL (0.3 µM, **D**, yellow arrowhead) or severe pericardial and periocular edema at the LC50 (0.5 µM, **E**, yellow arrowheads) and a curved tail (**E**, red arrowhead). (**F**, **G**) Measurements of body (**F**) and head (**G**) size of 4 dpf embryos treated with DMSO (N=4), 0.3 µM (N=4) or 0.5 µM (N=4) bixafen. (**H-M**) Dorsal view images of Tg(huC:G/U:RFP) embryos at 30 hpf (**H-J**) or 60 hpf (**K-M**), treated with DMSO (**H**, **K**), 0.3 µM bixafen (**I**, **L**) or 0.5 µM bixafen (**J**, **M**), showing a microcephaly in both stages on exposure to bixafen, with an increase in severity at 0.5 µM. (**N**) Illustration of a zebrafish embryo; the red line marks the depth at which confocal imaging was performed on (**H-M**). (**O**, **P**) Quantifications of the brain size of embryos at 30 hpf (**O**) or 60 hpf (**P**), treated with DMSO (N(30hpf)=5; N(60hpf)=5), 0.3 µM bixafen (N(30hpf)=13; N(60hpf)=12) or 0.5 µM bixafen (N(30hpf)=6; N(60hpf)=5), showing significantly smaller brains in embryos exposed to bixafen at LOAEL concentration and worsened with increase in concentration LC50, confirming the observations in (**H-M**). ***$p < 0.001$; **$p < 0.01$; *$p < 0.05$. Scale bar : (**C-E**) = 5 mm ; (**H-M**) = 100 µm.

Figure 2: Acute exposure to bixafen alters motor neuron axon development.

(**A**) Illustration of the experimental setup used to analyze caudal primary motor neuron

(CaPMN) axons length on acute exposure to bixafen. (**B-G**) Three-dimensional

reconstructions of the lateral view of a section of Tg(Olig2:eGFP) embryo tails at 30 hpf

(**B-D**) or 60 hpf (**E-G**) treated with DMSO (vehicle, **B**, **E**), 0.3 μM bixafen (**C**, **F**) or 0.5 μM

bixafen (**D**, **G**), showing developmental defects of the CaPMN axon on exposure to bixafen

at 0.3 μM concentration that is worsened with the concentration increase (0.5 μM). (**E'-**

G') Close-ups extracted from respectively **E**, **F** and **G**, showing that bixafen causes

developmental defects of CaPMN axons, that is worsened by increasing concentrations.

(**H**) Illustration of a zebrafish embryo; the red line marks the depth at which confocal

imaging was performed on (**B-G**). (**I**, **J**) Quantifications of the CaPMN axon length of

embryos at 30 hpf (**I**) or 60 hpf (**J**), treated with DMSO (N(30hpf)=8; N(60hpf)=8), 0.3 μM

bixafen (N(30hpf)=17; N(60hpf)=17) or 0.5 μM bixafen (N(30hpf)=16; N(60hpf)=13),

showing significantly shorter axons in embryos exposed to bixafen at LOAEL (0.3 μM) and

worsened with increase in concentration (LC50 0.5 μM), confirming the observations in

(**B-G**). ***$p < 0.001$; **$p < 0.01$; *$p < 0.05$. Scale bar : (**B-G**) = 100 μm.

References

Abad-Fuentes, A., Ceballos-Alcantarilla, E., Mercader, J. V., Agulló, C., Abad-Somovilla, A.,

and Esteve-Turrillas, F. A. (2015). Determination of succinate-dehydrogenase-

inhibitor fungicide residues in fruits and vegetables by liquid chromatography–

tandem mass spectrometry. *Anal. Bioanal. Chem.* 407. doi:10.1007/s00216-015-

8608-3.

Akerboom, J., Chen, T.-W., Wardill, T. J., Tian, L., Marvin, J. S., Mutlu, S., et al. (2012).

Optimization of a GCaMP Calcium Indicator for Neural Activity Imaging. *J. Neurosci.*

32, 13819–13840. doi:10.1523/JNEUROSCI.2601-12.2012.

Añasco, N. C., Koyama, J., and Uno, S. (2010). Pesticide residues in coastal waters affected

by rice paddy effluents temporarily stored in a wastewater reservoir in southern

393  Japan. *Arch. Environ. Contam. Toxicol.* 58, 352–360. doi:10.1007/s00244-009-9364-

394  1.

395 Ardissone, A., Invernizzi, F., Nasca, A., Moroni, I., Farina, L., and Ghezzi, D. (2015).

396  Mitochondrial leukoencephalopathy and complex II deficiency associated with a

397  recessive SDHB mutation with reduced penetrance. *Mol. Genet. Metab. Reports* 5,

398  51–54. doi:10.1016/j.ymgmr.2015.10.006.

399 Asakawa, K., Suster, M. L., Mizusawa, K., Nagayoshi, S., Kotani, T., Urasaki, A., et al.

400  (2008). Genetic dissection of neural circuits by Tol2 transposon-mediated Gal4

401  gene and enhancer trapping in zebrafish. *Proc. Natl. Acad. Sci. U. S. A.* 105, 1255–

402  1260. doi:10.1073/pnas.0704963105.

403 B, V., and C, P. (1990). The Neurotoxicity of Parathion-Induced Acetylcholinesterase

404  Inhibition in Neonatal Rats. *Neurotoxicology* 11.

405 Bénit, P., Kahn, A., Chretien, D., Bortoli, S., Huc, L., Schiff, M., et al. (2019). Evolutionarily

406  conserved susceptibility of the mitochondrial respiratory chain to SDHI pesticides

407  and its consequence on the impact of SDHIs on human cultured cells. *PLoS One* 14.

408  doi:10.1371/journal.pone.0224132.

409 Bénit, P., Letouzé, E., Rak, M., Aubry, L., Burnichon, N., Favier, J., et al. (2014).

410  Unsuspected task for an old team: Succinate, fumarate and other Krebs cycle acids

411  in metabolic remodeling. *Biochim. Biophys. Acta - Bioenerg.* 1837, 1330–1337.

412  doi:10.1016/j.bbabio.2014.03.013.

413 Bertolote, J. M., Fleischmann, A., Eddleston, M., and Gunnell, D. (2006). Deaths from

414  pesticide poisoning: A global response. *Br. J. Psychiatry* 189, 201–203.

415  doi:10.1192/bjp.bp.105.020834.

416 Birch-Machin, M. A., Taylor, R. W., Cochran, B., Ackrell, B. A. C., and Turnbull, D. M.

417  (2000). Late-onset optic atrophy, ataxia, and myopathy associated with a mutation

of a complex II gene. *Ann. Neurol.* 48, 330–335. doi:10.1002/1531-

8249(200009)48:3<330::AID-ANA7>3.0.CO;2-A.

Bourgeron, T. (2015). From the genetic architecture to synaptic plasticity in autism

spectrum disorder. *Nat. Rev. Neurosci.* 16, 551–563. doi:10.1038/nrn3992.

Brenet, A., Somkhit, J., Hassan-Abdi, R., Yanicostas, C., Igert, A., Saurat, D., et al. (2019).

Preclinical zebrafish model for organophosphorus intoxication: neuronal

hyperexcitation, behavioral abnormalities and subsequent brain damages. *bioRxiv*,

2019.12.15.876649. doi:10.1101/2019.12.15.876649.

Brustein, E., Saint-Amant, L., Buss, R. R., Chong, M., McDearmid, J. R., and Drapeau, P.

(2003). Steps during the development of the zebrafish locomotor network. in

Journal of Physiology Paris, 77–86. doi:10.1016/j.jphysparis.2003.10.009.

Finsterer, J. (2008). Cognitive decline as a manifestation of mitochondrial disorders

(mitochondrial dementia). *J. Neurol. Sci.* 272, 20–33. doi:10.1016/j.jns.2008.05.011.

Ghezzi, D., Goffrini, P., Uziel, G., Horvath, R., Klopstock, T., Lochmüller, H., et al. (2009).

SDHAF1, encoding a LYR complex-II specific assembly factor, is mutated in SDH-

defective infantile leukoencephalopathy. *Nat. Genet.* 41, 654–656.

doi:10.1038/ng.378.

Graillot, V., Tomasetig, F., Cravedi, J. P., and Audebert, M. (2012). Evidence of the in vitro

genotoxicity of methyl-pyrazole pesticides in human cells. *Mutat. Res. - Genet.*

Toxicol. Environ. Mutagen. 748, 8–16. doi:10.1016/j.mrgentox.2012.05.014.

Horváth, R., Abicht, A., Holinski-Feder, E., Laner, A., Gempel, K., Prokisch, H., et al. (2006).

Leigh syndrome caused by mutations in the flavoprotein (Fp) subunit of succinate

dehydrogenase (SDHA). *J. Neurol. Neurosurg. Psychiatry* 77, 74–76.

doi:10.1136/jnnp.2005.067041.

Howard, A. S., Bucelli, R., Jett, D. A., Bruun, D., Yang, D., and Lein, P. J. (2005). Chlorpyrifos

exerts opposing effects on axonal and dendritic growth in primary neuronal

cultures. *Toxicol. Appl. Pharmacol.* 207, 112–124. doi:10.1016/j.taap.2004.12.008.

Lein, P., Silbergeld, E., Locke, P., and Goldberg, A. M. (2005). In vitro and other

alternative approaches to developmental neurotoxicity testing (DNT). in

Environmental Toxicology and Pharmacology (Elsevier), 735–744.

doi:10.1016/j.etap.2004.12.035.

Martin, C., Dubouchaud, H., Mosoni, L., Chardigny, J. M., Oudot, A., Fontaine, E., et al.

(2007). Abnormalities of mitochondrial functioning can partly explain the

metabolic disorders encountered in sarcopenic gastrocnemius. *Aging Cell* 6, 165–

177. doi:10.1111/j.1474-9726.2007.00271.x.

Merhi, M., Raynal, H., Cahuzac, E., Vinson, F., Cravedi, J. P., and Gamet-Payrastre, L.

(2007). Occupational exposure to pesticides and risk of hematopoietic cancers:

Meta-analysis of case-control studies. *Cancer Causes Control* 18, 1209–1226.

doi:10.1007/s10552-007-9061-1.

Muñoz-Quezada, M. T., Lucero, B. A., Barr, D. B., Steenland, K., Levy, K., Ryan, P. B., et al.

(2013). Neurodevelopmental effects in children associated with exposure to

organophosphate pesticides: A systematic review. *Neurotoxicology* 39, 158–168.

doi:10.1016/j.neuro.2013.09.003.

Myers, P. Z. (1985). Spinal motoneurons of the larval zebrafish. *J. Comp. Neurol.* 236,

555–561. doi:10.1002/cne.902360411.

Myers, P. Z., Eisen, J. S., and Westerfield, M. (1986). Development and axonal outgrowth

of identified motoneurons in the zebrafish. *J. Neurosci.* 6, 2278–2289.

doi:10.1523/jneurosci.06-08-02278.1986.

Paganelli, A., Gnazzo, V., Acosta, H., López, S. L., and Carrasco, A. E. (2010). Glyphosate-

based herbicides produce teratogenic effects on vertebrates by impairing retinoic

acid signaling. *Chem. Res. Toxicol.* 23, 1586–1595. doi:10.1021/tx1001749.

Parfait, B., Chretien, D., Rötig, A., Marsac, C., Munnich, A., and Rustin, P. (2000).

Compound heterozygous mutations in the flavoprotein gene of the respiratory

chain complex II in a patient with Leigh syndrome. *Hum. Genet.* 106, 236–243.

doi:10.1007/s004390051033.

Perry, C. G., Young, W. F., McWhinney, S. R., Bei, T., Stergiopoulos, S., Knudson, R. A., et al.

(2006). Functioning paraganglioma and gastrointestinal stromal tumor of the

jejunum in three women: Syndrome or coincidence. *Am. J. Surg. Pathol.* 30, 42–49.

doi:10.1097/01.pas.0000178087.69394.9f.

Qian, L., Cui, F., Yang, Y., Liu, Y., Qi, S., and Wang, C. (2018). Mechanisms of

developmental toxicity in zebrafish embryos (Danio rerio) induced by boscalid. *Sci.*

Total Environ. 634, 478–487. doi:10.1016/j.scitotenv.2018.04.012.

Rauh, V. A., Perera, F. P., Horton, M. K., Whyatt, R. M., Bansal, R., Hao, X., et al. (2012).

Brain anomalies in children exposed prenatally to a common organophosphate

pesticide. *Proc. Natl. Acad. Sci. U. S. A.* 109, 7871–7876.

doi:10.1073/pnas.1203396109.

Ravichandra, N. G. (2018). *Agrochemicals in Plant Disease Management*.

Rehfus, A., Miessner, S., Achenbach, J., Strobel, D., Bryson, R., and Stammler, G. (2016).

Emergence of succinate dehydrogenase inhibitor resistance of Pyrenophora teres in

Europe. *Pest Manag. Sci.* 72, 1977–1988. doi:10.1002/ps.4244.

Ricketts, C., Zeegers, M. P., Lubinski, J., and Maher, E. R. (2009). Analysis of germline

variants in CDH1, IGFBP3, MMP1, MMP3, STK15 and VEGF in familial and sporadic

renal cell carcinoma. *PLoS One* 4. doi:10.1371/journal.pone.0006037.

Ross, S. M. K., McManus, I. C., Harrison, V., and Mason, O. (2013). Neurobehavioral

problems following low-level exposure to organophosphate pesticides: A

systematic and meta-analytic review. *Crit. Rev. Toxicol.* 43, 21–44.

doi:10.3109/10408444.2012.738645.

Shin, J., Park, H. C., Topczewska, J. M., Madwsley, D. J., and Appel, B. (2003). Neural cell

fate analysis in zebrafish using olig2 BAC transgenics. *Methods Cell Sci.* 25, 7–14.

doi:10.1023/B:MICS.0000006847.09037.3a.

Stratakis, C. A., and Carney, J. A. (2009). The triad of paragangliomas, gastric stromal

tumours and pulmonary chondromas (Carney triad), and the dyad of

paragangliomas and gastric stromal sarcomas (Carney-Stratakis syndrome):

Molecular genetics and clinical implications. in *Journal of Internal Medicine* (J Intern

Med), 43–52. doi:10.1111/j.1365-2796.2009.02110.x.

Svartz, G., Meijide, F., and Pérez Coll, C. (2016). Effects of a fungicide formulation on

embryo-larval development, metamorphosis, and gonadogenesis of the South

American toad Rhinella arenarum. *Environ. Toxicol. Pharmacol.* 45, 1–7.

doi:10.1016/j.etap.2016.05.008.

Tanabe, A., and Kawata, K. (2009). Daily variation of pesticides in surface water of a

small river flowing through paddy field Area. *Bull. Environ. Contam. Toxicol.* 82,

705–710. doi:10.1007/s00128-009-9695-7.

Timmers, H. J. L. M., Gimenez-Roqueplo, A. P., Mannelli, M., and Pacak, K. (2009). Clinical

aspects of SDHx-related pheochromocytoma and paraganglioma. *Endocr. Relat.*

Cancer 16, 391–400. doi:10.1677/ERC-08-0284.

Tsuda, T., Nakamura, T., Inoue, A., and Tanaka, K. (2009). Pesticides in water, fish and

shellfish from littoral area of lake biwa. *Bull. Environ. Contam. Toxicol.* 82, 716–721.

doi:10.1007/s00128-009-9681-0.

Van Coster, R., Seneca, S., Smet, J., Van Hecke, R., Gerlo, E., Devreese, B., et al. (2003).

Homozygous Gly555Glu mutation in the nuclear-encoded 70 kDa flavoprotein gene

causes instability of the respiratory chain complex II. *Am. J. Med. Genet.* 120A, 13–18. doi:10.1002/ajmg.a.10202.

van den Berg, F., Tiktak, A., Heuvelink, G. B. M., Burgers, S. L. G. E., Brus, D. J., de Vries, F., et al. (2012). Propagation of Uncertainties in Soil and Pesticide Properties to Pesticide Leaching. *J. Environ. Qual.* 41, 253–261. doi:10.2134/jeq2011.0167.

Veloukas, T., Markoglou, A. N., and Karaoglanidis, G. S. (2013). Differential effect of SdhB gene mutations on the sensitivity to SDHI fungicides in botrytis cinerea. *Plant Dis.* 97, 118–122. doi:10.1094/PDIS-03-12-0322-RE.

VKM Report (2014). Health risk assessment of the fungicide Aviator Xpro EC 225 - Vitenskapskomiteen for mat og miljø. 1–19. Available at: https://vkm.no/english/riskassessments/allpublications/healthriskassessmentofthefungicideaviatorxproec225.4.62c3f31d15e03ed297212ed.html

VKM Report (2015). Risk assessment of the metabolite M44 of bixafen, one of the active substances in Aviator Xpro EC 225 - Vitenskapskomiteen for mat og miljø. 1–20. Available at: https://vkm.no/english/riskassessments/allpublications/riskassessmentofthemetabolitem44ofBixafenoneoftheactivesubstancesinaviatorxproec225.4.62c3f31d15e03ed2972a8d.html

Vu, H. T., Keough, M. J., Long, S. M., and Pettigrove, V. J. (2016). Effects of the boscalid fungicide Filan® on the marine amphipod Allorchestes compressa at environmentally relevant concentrations. *Environ. Toxicol. Chem.* 35, 1130–1137. doi:10.1002/etc.3247.

Wang, H., Meng, Z., Liu, F., Zhou, L., Su, M., Meng, Y., et al. (2020). Characterization of boscalid-induced oxidative stress and neurodevelopmental toxicity in zebrafish embryos. *Chemosphere* 238. doi:10.1016/j.chemosphere.2019.124753.

Weichenthal, S., Moase, C., and Chan, P. (2010). A review of pesticide exposure and

cancer incidence in the agricultural health study cohort. *Environ. Health Perspect.*

118, 1117–1125. doi:10.1289/ehp.0901731.

Westerfield, M., McMurray, J. V., and Eisen, J. S. (1986). Identified motoneurons and their

innervation of axial muscles in the zebrafish. *J. Neurosci.* 6, 2267–2277.

doi:10.1523/jneurosci.06-08-02267.1986.

Wu, S., Lei, L., Liu, M., Song, Y., Lu, S., Li, D., et al. (2018). Single and mixture toxicity of

strobilurin and SDHI fungicides to Xenopus tropicalis embryos.

doi:10.1016/j.ecoenv.2018.01.045.

Yamashita, M., and Fraaije, B. (2018). Non-target site SDHI resistance is present as

standing genetic variation in field populations of Zymoseptoria tritici. *Pest Manag.*

Sci. 74, 672–681. doi:10.1002/ps.4761.

Yang, Y., Dong, F., Liu, X., Xu, J., Wu, X., Qi, S., et al. (2018). Thifluzamide affects lipid

metabolism in zebrafish (Danio reio). *Sci. Total Environ.* 633, 1227–1236.

doi:10.1016/j.scitotenv.2018.03.302.

Yao, H., Yu, J., Zhou, Y., Xiang, Q., and Xu, C. (2018). The embryonic developmental effect

of sedaxane on zebrafish (Danio rerio). doi:10.1016/j.chemosphere.2018.01.046.

Yoshida, M., Inoue, K., and Takahashi, M. (2015). Predictive modes of action of pesticides

in uterine adenocarcinoma development in rats. *J. Toxicol. Pathol.* 28, 207–216.

doi:10.1293/tox.2015-0026.