1 **Evolution of multicellularity and unicellularity in yeast *S. cerevisiae* to study reversibility**
2 **of evolutionary trajectories.**
3
4

5 Phaniendra Alugoju[#], Anjali Mahilkar[#], and Supreet Saini*
6 Department of Chemical Engineering, Indian Institute of Technology Bombay, Mumbai 400 076
7
8
9 # Authors contributed equally to this study
10
11 *Corresponding author. Email: saini@che.iitb.ac.in, Phone: 091 22 2576 7216
12

Abstract

Adaptive trajectories of populations have been focus of number of studies. However, adaptive trajectories have not been studied in the context of reverse evolution. By reverse evolution, we mean a scenario where selection is reversed. In this work, we use evolution (and reversal from) of multicellularity in *S. cerevisiae* as a model to answer this question. When selected for fast-settling variants, multicellularity evolves rapidly in the organism. On reversing selection, unicellularity evolves from the multicellular clusters. However, the dynamic trajectories of the two processes are different. In this context, evolution is not reversed dynamically at a phenotypic level. The phenotypic reversal is not driven by reversal of the original mutations during the forward evolution. Overall, our results show that the dynamics of molecular and phenotypic trajectories of evolution are distinct, and reversal of selection leads to unique trajectories of phenotypic reversal.

Keywords

Multicellular yeast clusters, unicellularity, multicellularity, reverse evolution, evolutionary dynamics.

Introduction

Reversibility, in an evolutionary context, is often seen as return to ancestral state (1). Historically, investigation of reversibility of evolutionary processes has been an important theme in study of evolution (2-5). In microbial systems, this question has been addressed in a number of experimental studies, and has led to different answers (6, 7). In these experiments, a population is shifted from one environment to another, and adapted for a certain number of generations. The evolved population is then shifted back to the original environment, and its adaptation studied. In this context, reversibility is studied as the ability/dynamics of a population to re-adapt in an environment, when shifted from another (7). Towards this end, one of the most well studied systems has been evolution of sensitive bacteria in a population of resistant bacteria, when a stressor (antibiotic/phage) is removed from the environment (8-10). In the context of microbial systems, sequencing makes it possible to trace the reversibility of evolution at genetic or molecular level. In bacteria, when antibiotic resistance was removed, the original genotype was not restored (11, 12). On the other hand, evolution experiments with virus have demonstrated that phenotypic reversals can map to precise nucleotide reversals (6, 13). Enzymes, too, have been shown to recover activity, albeit via different mutational paths (14). The broad lesson from these experiments is that the longer the time spent in the novel environment, the more difficult the likelihood of reverse evolution becomes, when the population is re-introduced in the original environment.

However, reversibility in another context, has been largely unexplored. What happens to the evolutionary fate of a population, when the selection pressure is reversed? In this context, to study reversibility of evolution, we need systems where an exact and opposite selection pressure is applied on a system. Such a choice of a system is not trivial. For instance, is absence of an antibiotic the reverse of evolution in presence of (how much?) antibiotic? Or, is evolution in presence of glucose the reverse of evolution in glycerol? To answer this question, we study evolution (and reverse) of multicellularity in yeast, using settling (or lack of) as a selection force.

The evolutionary transition from single cells to multicellular organisms was one of the major events in the history of life on Earth (15). Eukaryotic multicellularity has evolved independently at least 25 times in distinct lineages (16-18). However, understanding the evolution of complex multicellular organisms from unicellular organisms is challenging, because the first steps in this process occurred in the deep past (16, 19). Mathematical models have demonstrated key steps involved in the transition from unicellular form to multicellular clusters (20-23). Experimentally, yeast *Saccharomyces cerevisiae* has been used for studying the origin of multicellularity in eukaryotes (24-32). These studies have demonstrated that under centrifugal selection, the transition from unicellularity to multicellularity is relatively quick. The mechanism of cluster formation in yeast is through post-division adhesion, but not through aggregation (24). This de novo origin

of snowflake yeast clusters is mainly due to mutations in transcription factor ACE2 gene (33, 34). This
method of growth ensures high relatedness among component cells of a multicellular cluster, therefore
reducing internal conflicts. Clusters reproduce by fragmentation especially when tension among cells in the
cluster exceeds the tensile strength of cellular adhesion (25). Apoptosis, a major determinant in the
development and somatic maintenance in metazoans, is also involved the reproduction of yeast
multiclusters (34).

Though largely unicellular, there are some yeast strains which exit as multicellular clumps in their ecological
niches. The multicellular existence is thought have been driven because of the benefit it provides under
conditions of stress (35). These clusters, when compared with the unicellular counterpart, were found to be
more resistant to oxidative and solvent stresses. This increased stress resistance however came at a cost
of lower growth rate, in the absence of stress.

Depending on the condition, yeast can show bidirectional transitions between unicellularity and
multicellularity (30, 36-38). A recent study showed that unicellular yeast evolve from wild type clump-forming
ancestors by propagating samples from suspension after larger clumps have settled (35). Thus, yeast
clusters serve as an excellent model for studying the dynamics of reverse evolutionary events.

In this work, we first evolved yeast multicellular clusters from unicellular yeast *S. cerevisiae* using settling
as a selection force. The multicellular cluster was verified to be because of mutation in ACE2 gene, and not
simply due to cell-cell adhesion. We show that the multicellular clusters perform better than the ancestor in
presence of various stresses. The multicellular cluster is then reverse evolved. Would reversal of selection
lead the population to re-trace its path on the phenotypic space to the ancestral phenotype? Or, would
availability of a large number of genetic solutions, in response to a precisely opposite environmental
selection, dictate that the reverse path is largely unpredictable? Our results from the reverse evolution
experiment show that, at the phenotypic level, the kinetics of evolution of unicellularity does not simply
retrace the path that was taken to multicellularity. The genetic solutions that help accomplish this change
are also distinct, and not because of mutations in ACE2. Study of bidirectional transitions between multi-
and unicellularity thus offer a model system to study reversibility in an evolutionary process.

Materials and Methods

Yeast strain and growth medium composition.

Yeast *Saccharomyces cerevisiae* haploid strain SC644 (*MATa MEL1ade1 ile trp1-HIII ura3-52*) was used

(39). Cells were grown in test tubes (25 x 150 mm) containing liquid YPD (1% yeast extract, 2% peptone

and 2% glucose) medium at 30°C. Unless stated otherwise, all cultures were grown in an orbital shaker at

a speed of 250 rpm for 24 h. Yeast extract and Peptone were procured from Himedia, Mumbai, India.

Glucose was procured from Sigma Aldrich, USA. All the chemicals used in this study were of highest

analytical grade.

Selection for multicellularity.

Multicellular yeast clusters were generated as described by Ratcliff et. al. 2012 (24) (Figure 1A). A single

isogenic yeast colony was inoculated into 10 ml of YPD medium and incubated at 30°C with 250 rpm for

24 h. The resulting culture was then used as a pre-inoculum for the initiation of the selection for

multicellularity. An aliquot (1 ml) of overnight grown yeast culture was transferred to a 1.5 ml microcentrifuge

tube and then centrifuged at a speed of 100g for 15 sec. The bottom 100 µL was transferred to a tube

containing fresh YPD media and allowed to grow (30°C, 250 rpm) for 24 h. This process was continued for

60 transfers. Six independent lines under centrifugal selection pressure and the control were studied in this

work. A control line was evolved in YPD, from which, 100 µL cultures was transferred into fresh YPD after

growth for 24 h (30°C, 24 h). In this control, the transfer was immediate without allowing the cells to settle

down. Glycerol freezer stocks, in 30% glycerol, were made for every tenth transfer and stored at -80°C.

Selection for unicellularity.

The frozen glycerol stocks of multicellular clusters obtained after 60 transfers were streaked onto YPD

plates and incubated at 30°C for 48 h (Figure 1B). Cells from a single colony were spread on glass slide

and a cover slip. Images were taken using a microscope (Labomed Binocular) under 40X magnification. A

single multicellular cluster was then inoculated into fresh YPD medium and incubated at 30°C for 24 h with

shaking. These cultures were used as ancestors for the reverse selection of unicellular phenotype from the

multicellular cluster phenotype. As described above, an aliquot (1 ml) of overnight grown yeast culture was

transferred to centrifuge tubes and centrifuged at a speed of 100xg for 15 sec. In this experiment, the top

100 µL was transferred into tubes containing fresh YDP medium every 24 h. Glycerol stocks were made

for every tenth transfer in 30% glycerol and stored at -80°C.

Characterization of colony phenotype.

Each of the overnight grown (24 h) yeast population streaked onto YPD plates. These plates were then incubated at 30°C until single colonies appear, at which point, images of the plates were taken (28).

Growth experiments.

Saturated yeast cultures, from both forward and reverse evolution experiments, were diluted in fresh media to get an initial optical density at 600 nm wavelength (OD600) to 0.1. The diluted culture was allowed to grow at 30°C and growth kinetics monitored in a Tecan Infinite M200 Pro microplate reader. The microplates were overlaid with a Breathe Easy membrane (Sigma).

To test growth of yeast clusters in ethanol stress, overnight cultures were diluted in fresh media to an OD600 of 0.1. Ethanol (to a final concentration of 0, 2.5%, 5%, and 10%) was added to the cultures and the cell growth was checked by measuring the OD600 using a 96-well plate reader. OD600 values were taken every one hour over a 30 h period.

Measurement of yeast cluster size.

Yeast populations from both forward and reverse evolution experiments were grown in 10 ml of fresh YPD media shaken at 250 rpm for 24 h. These overnight cultures were transferred to fresh YPD medium and grown until exponential phase (OD600 ~ 2). About 100 μl of exponentially growing cells were washed thrice with PBS and analyzed for FSC-A of each cells/cluster using flow cytometer. A minimum of 30,000 clusters per population were analyzed and the average of the FSC-A at the indicated tome of transfer time was calculated (40).

Settling assay.

Yeast cultures from forward evolution experiments were inoculated from single colonies into 10 ml YPD liquid cultures and grown overnight at 30°C. Each culture tube was vigorously vortexed and placed on a lab bench at room temperature to allow for settling. Images of each population during the settling were taken at different time intervals (0, 15, 30, and 60 min) (41).

Stress resistance assay.

Each of the yeast populations was grown at 30°C with 250 rpm shaking for 20 h. Cells were serially diluted to OD600 of 3, 1, 0.3 and 0.1. A 4 μl volume of these cultures was spotted onto YPD plates. In order to test the effect of cold and heat stress, a set of potted plates were also incubated at 20°C and 37°C, respectively.

170   A single plate was incubated at 30°C as a control. Images of the plates were taken at different days of
171   incubation.

Cloning of ACE2 gene in pT1-3B plasmid.

ACE2 promoter region and the coding sequence was amplified from the genomic DNA of the ancestor using
the primers (5'GGG GTC GAC TTA GTT AAC TCT ATC TAT TG3') and (5'GGG AAG CTT TTT TTG GCC
CTT AAG ACT AC3'). The PCR product and the plasmid pT1-3B (42) was digested with the restriction
enzymes Sal1 and HindIII (New England Biolab (NEB)). T4 DNA ligase was used to construct the resulting
plasmid pACE2, in which ACE2 expression was under its native promoter.

In addition to the ACE2 sequence from the ancestor, two other templates were used to construct plasmids
pACE2T40 and pACE2T50. For the first (pACE2T40), the ACE2 promoter and coding region was amplified
from the genomic DNA of cells in line 1 after 40 transfers. For the second (pACE2T50), the ACE2 promoter
and the coding region was amplified from the genomic DNA of cells in line 1 after 50 transfers.

Complementation of ACE2 was performed by cloning the promoter and the coding region from each of the
six lines after 60 transfers on the plasmid pT1-3B and transforming the resulting plasmid in an ACE2Δ
mutant. The ACE2 plasmids from each of the six lines are called pACE2L1, pACE2L2, pACE2L3, pACE2L4,
pACE2L5, and pACE2L6. The promoter and the coding region from each of the six lines was amplified
using the primers given above. The region was amplified from DNA isolated from a single colony on YPD
plates from each of the six lines.

ACE2 knockout.

The strain ΔACE2 was constructed by replacement of the ACE2 coding region with a hygromycin B
resistance gene. For this, the gene HPH was amplified using the template pUG75, with primers 5'ATG GAT
AAC GTT GTA GAT CCG TGG TAT ATA AAT CCC TCA GGC TTC GCG AAG CAG GTC GAC AAC CCT
TAA T3' and 5'TCA GAG AGC ATC AGT TTC GTT TGA AAG GGT GCG GTT CGA GTT TTG CTC GTA
GTG GAT CTG ATA TCA CCT A3'. The PCR product was transformed in the respective strain and single
colonies selected for on plates containing hygromycin B at a concentration 200 µg/ml. The replacement of
ACE2 with HPH was confirmed using primers 5'CTC AAG CAA CAG TTA AAG TGC3' and 5'TGT TAC TAT
TAT TTA TTA TG 3'.

Sequencing of ACE2 gene.

Genomic DNA was isolated from ancestor and six replicate populations from both forward and reverse
evolution as described previously (43). ACE2 gene was amplified using forward (5'ATG GAT AAC GTT

GTA GAT CCG TGG TAT3') and reverse (5'TCA GAG AGC ATC AGT TTC GTT TGA AAG3') primers and

sequenced. The coding sequence of the gene AMN1 was amplified using the primers (5'ATT TAT CAT TTT

CCT TTT CT3') and (5' TTA CTG GTG GTC TGT TGT GTA 3'). All sequencing was done by Eurofins

Scientific.

Results and discussion

Evolution of multicellularity on serial propagation under centrifugal selection

Unicellular to multicellular transition of yeast was studied using settling as selection (24). Figure 2A shows the representative images of yeast multicellular cluster populations during centrifugal selection. By 60 transfers, we note complex multicellular clusters in replicate populations (although cluster sizes vary across different lines). The dynamics of the change in phenomenology between the different independent lines is largely identical with each other in four of the six lines; but not the other two. Such parallelism in evolutionary experiments has been reported in the past (44-46). It has been suggested that even under controlled environmental growth conditions, the evolution of multicellular clusters generates spatial structure with increasing localized interactions and creates new spatial gradients (19). The evolution of multicellular clusters is driven by both chance and environmental changes (24, 27).

When streaked on YPD agar plates, the evolved lines exhibit an altered colony morphology. These plates were incubated at 30°C until we see single colonies. As shown in Figure 2B, yeast ancestor and multicellular cluster colonies exhibited a distinct surface morphology. Yeast multicellular colonies showed rugose (rough) colony morphology, whereas unicellular ancestor showed a smooth morphology. This is a result of the cluster structure dictating the colony surface structure in the evolved lines.

Dynamics of changes in yeast cluster size with serial propagation under centrifugal selection

Centrifugal selection favors the evolution of larger cluster size (24) and divergence in cluster size evolves rapidly during the transition from unicellularity to multicellularity (27). Therefore, in this study the average cluster size of yeast replicate populations was quantified as mean value of forward scatter area following every ten transfers using a flow cytometer. The size of the yeast clusters increased with concomitant increase in transfer time in most of the replicate populations (MC1-4 and MC6, but not MC5) (Figure 2C). While multicellularity evolved in all six replicate populations, it is also evident that multicellular clusters in the different lines did not evolve to the same size, and dynamics of evolution of the mean cluster size was not identical in multiple lines. Microbial evolution experiments have demonstrated that diversity may depend largely on environmental conditions (47) and ecological interactions between different genotypes (48).

Expectedly, the multicellular yeast clusters settles faster with increasing cluster size. It is evident from the Figure 3 that, four of the multicellular replicate populations (L1-L4) started settling faster after 40 transfers. However, replicate populations L5 showed moderate settling even after 60 transfers. The first snowflake yeast clusters formed are small during early transfer times, therefore they settle slowly. However, settling rate increased rapidly with the evolution of large cluster size during later stages of transfer times.

Multicellular clusters exhibit a tradeoff between growth rate and resistance to stress.

Multicellularity is associated with multiple benefits such as reduced predation and division of labor between cells. However, these benefits come with costs. Multicellular clusters showed lower growth rates relative to their unicellular counterparts because of limitations of space and resource acquisition (27). The geometric constraints such as low surface area to volume ratio and of spatial organization in multicellular clusters limit the extent of interactions with the nutrient resources in the environment, and subsequently affects the metabolism, growth rate and viability (49). It has been suggested that there may be a trade-off between individual cluster size and growth rate (27). Therefore, we investigate whether cluster phenotype has any effect on the growth kinetics. As shown in Figure 4A and 4B, multicellular cluster populations grow slower when compared to their unicellular counterparts. As per the size-complexity rule, bigger entities have a greater division of labor than smaller ones (50).

A transition to cluster form confers selective advantage over unicellularity in ecological niches (17, 51). Examples of this phenomenon include protection from phagocytosis (52), predators (53), environmental stressors (ethanol and antibiotics) (54) and uptake of harmful compounds (55). Clustering may be beneficial for more efficient exchange of external resources between component cells of a cluster by reducing interactions with non-cooperative individuals (or cheaters) (i.e. increased efficiency of cooperative feeding) (26). It has also been suggested that cluster levels selection is beneficial for the use of growth-promoting secretions as well as exclusion of cheaters (19). This division of labor can allow multicellular organisms to perform tasks more efficiently, and can facilitate the evolution of complex traits (1). We study the heat and solvent stress resistance of yeast clusters and compare it with the unicellular ancestor's performance in these environments.

Heat stress

Previous studies have reported that wild type yeast clusters strains were resistant to free/thaw treatment (31). We use spot assay to investigate the sensitivity of yeast cluster phenotype to cold (20°C) and heat stress (37°C). Multicellular yeast cluster lines (MC 1-6) were found to be resistant to heat stress (37°C) when compared to unicellular ancestor (Figure 4C). Even after five days of incubation, there is no significant growth of ancestor at indicated cell densities viz at OD600 of 0.3 and 0.1. Cold stress has no effect on the survival of both unicellular and multicellular cell lines.

Solvent stress.

Toxic effect of ethanol on yeast is mainly caused by damaging the mitochondrial DNA and inactivation of enzymes such as hexokinase and dehydrogenase (56). Yeast multicellular clusters showed longer lag phase when compared to unicellular ancestor (Figure 4D). None of the cultures grew on higher ethanol concentrations (5% and 10%) (Figure 5).

Mutations in ACE2 transcription factor leads to evolution of the multicellular phenotype.

Transition to multicellularity is typically associated with increases in the numbers of genes involved in three important processes such as cell differentiation (57), cell-cell communication (58, 59), and adhesion (17, 60, 61). The genetic basis for the evolution of multicellular cluster phenotype in yeast is due to disruption of ACE2 gene function. It has been reported that early (7-day) snowflake yeast clusters showed down-regulation of seven genes (CTS1, DSE4, DSE2, SUN4, DSE1, SCW11 and AMN1) which are important for daughter cell separation. Many of these genes act directly to degrade the bud neck septum. Previous studies have shown that the mutation or knockout of ACE2 gene prevents daughter–mother cell separation results in the formation of the snowflake phenotype (33, 62, 63). These experiments demonstrate that a single genetic mutation not only can create a new level of biological organization, but can also facilitate the evolution of higher-level complexity (34). Ratcliff et al reported non-synonymus mutations in that five of the ten replicate populations studied. In our experiments, all six lines acquired mutations in the ACE2 gene.

All six lines in our experiment, after 60 transfers, had mutations in the coding region of the gene (Figure 6). Line 1 had a SNP at position 400, which changed Lysine residue to a STOP codon and a non-synonymous change at position 862 (C to A; Pro to Thr). Line 2 had an insertion of nucleotide 'G' at position 1001 in the coding region of ACE2. Line 3 had a SNP (T to A) at position 1065, which changed a Tyrosine residue to a STOP codon. Line 4 had a T to G SNP at position 1038 in the coding region. This SNP changed a Tyrosine to a STOP codon. Thus, no functional copies of ACE2 were being produced in the cell in these four lines. Line 5 had a SNP at position 1134 in the coding region of ACE2 (C to G). This lead to a non-synonymous change in the coding region of the protein (Phe to Leu). Line 6 had SNPs at positions 554 (G to C; Ser to Thr) and 778 (C to G; Pro to Ala). To verify the effect of the mutant allele of ACE2 in dictating the multicellular cluster, the ACE2 promoter and the coding region from each of the six lines were transferred to an ACE2Δ mutant. Transferring the ACE2 allele lead to the multicellular phenotype in each of the six complementation experiment (Figure 7). We do not discount the possibility of other molecular determinants dictating the evolution of the multicellular cluster. However, our results show, mutations in ACE2 is a key molecular determinant in dictating the evolution of the multicellular phenotype. Sequencing of the ACE2 region in line 1 after 40 and 50 transfers revealed that the non-synonymous mutation (P to T) was acquired first. This mutation was first observed after 40 transfers. Thereafter, the A400T mutation was observed after 50 transfers. Thus, the increase in cluster size was concomitant with the accumulation of mutations at the ACE2 locus. The coding region of AMN1 was also sequenced in all six lines after 60 transfers. However, no mutation was identified in this region of the chromosome.

Reversal of centrifugal selection leads to evolution of unicellularity from multicellular clusters.

Compared to unicellular ancestor, multicellular clusters grow slower in culture media. The growth limitation

is due to increased competition and mass transfer limitations regarding access to resources (20). Further,

their complexity makes them more vulnerable to disruption (1). Given this, in what contexts can we study

evolution of unicellularity, starting with a multicellular cluster?

In the present study, we aimed at investigating the reverse evolution from multicellularity to unicellularity

using experimental evolution, under the same centrifugal selection pressure. Growing the multicellular cells

the same way, and after the same processing, the cells from the topmost layer of the liquid were transferred

to the next tube. In this way, the selection pressure in the experiment was exactly the opposite of the one

applied in the uni- to multicellular evolution. This reverse evolution experiment leads to several physiological

changes. Firstly, the average cell size in the clusters decreased rapidly and unicellular lines were evolved.

As a result, the colony morphology was restored from rugose to smooth colony morphology during reverse

evolution from multicellularity to unicellularity (Figure 8A). It has been suggested that both genotypic and

phenotypic factors can affect the colony morphology. Some cellular properties, such as cell adhesion, and

budding pattern, have also been associated with morphological changes in *S. cerevisiae* (64).

The trajectory of the cluster size in the reverse evolution is shown in Figure 8B. As shown in the Figure, in

most of the lines, the transition from multicellular to unicellular phenotype (transfers 60 to 120) has a

trajectory distinct from the original transition from unicellular to multicellular phenotype (transfer 0 to 60).

This data suggests that the evolutionary trajectory does not retrace its path, at a phenotypic level. Second,

the growth kinetics of the reverse evolved lines was restored closer to that of the ancestral strain (Figure

9). This is likely due to the transport limitations being eliminated in the reverse evolved unicellular lines.

To understand the molecular basis which trigger the reverse evolution, after 60 transfers, we sequenced

the ACE2 region of the reverse evolved lines. However, none of the four lines had acquired a mutation in

the ACE2 region, when compared to the corresponding sequence of the multicellular cluster that each line

had started from. Thus, the evolutionary transition from the multicellular to the unicellular phenotype is

brought about by mutations elsewhere on the chromosome.

Additionally, six independent lines from the multicellular clusters from line 1 were reverse evolved, selecting

for unicellularity. The premise behind this experiment was to test for repeatability of evolution and identify

paths where the phenotypic trajectory is re-traced. Our results demonstrate that significant differences exist

between the evolutionary paths traced by the six independent lines in the reverse evolution (Figure 10).

Moreover, none of the six paths is statistically identical with the path traced when the line was evolved for

multicellularity. Sequencing of the ACE2 coding region revealed that none of the six independent lines in

the reverse evolution had acquired a mutation. Thus, the molecular determinants of the reverse evolution

experiment lie elsewhere on the chromosome.

Conclusion

In this work, we report forward and reverse evolution between unicellularity and multicellularity in the yeast *S. cerevisiae.* In this context, selection of settling under gravity or centrifugation provides an environment where study of evolution, when selection is reversed can be studied. As has been reported previously, multicellularity readily evolves under the appropriate selection pressure. Of our six lines, four of them exhibit dynamics of cluster size increase quite close to each other. To that effect, the evolution of multicellularity is a repeatable and predictable process. Somewhat more surprisingly, all six lines acquire mutation at a single locus in the genome. This single set of mutations in the ACE2 gene is responsible for exhibition of the multicellularity trait (Figure 6). The mutations at this locus suggest that the selection coefficient associated with an ACE2 mutation is very high.

On reversal of the selection pressure, all six lines exhibit reversal of the multicellular phenotype. While the phenotype is completely reversed in these lines, the trajectory of the change in the cluster size does not mimic the forward evolution experiment. Moreover, sequencing the ACE2 region in the reverse evolved lines tells us that the unicellular phenotype did not evolve because of additional/reverse mutations in ACE2. Thus, the molecular basis of the reverse evolution experiment is at another locus/loci on the genome.

Our results also show that although the selection pressure of gravity leads to evolution of multicellularity, the multicellular phenotype confers advantages in the face of several environmental stresses too. This advantage has been observed in a number of studies (26, 65-67). Thus, in the context of evolution of multicellularity during the evolution, any of these stresses could provide an avenue for evolution of multicellularity, which is known to have evolved a number of times in the history of evolution of life on Earth (15). Reverse evolution from multicellularity to unicellularity has been studied in two different contexts (35, 68). In the first, natural isolates which formed multicellular clusters were reverse evolved for unicellularity. The authors report that mutations at the AMN1 (69) locus were responsible for evolution of unicellularity. The dynamics of this transition was, however, not reported. In a second study, reversal to unicellularity was evolved via propagation on solid media, in a structured environment. The selection forces driving this transition are, hence, not the opposite of those which drove evolution of multicellularity.

From our reverse evolution experiment, we note that the dynamic trajectory of reverse evolution is different from that of the forward one. In the context of reverse evolution, what is the genetics at the organismal level, which constrains this evolutionary process? (70, 71). If we assume that the strength of selection is reversed exactly in the reverse experiment, then what could lead to the difference in the trajectories? Several factors could be responsible for this. First, with the evolution of multicellularity, the size of the population in the culture media changes. Population size is an important determinant of the rate of

adaptation (72). Second, mutation rates across the genome are known to be variable (73). Hence, since in

this case, the molecular determinants of the forward and reverse evolution are different, these might be

subject to different mutation rates. Third, the number of adaptive mutations facilitating forward and reverse

evolution might be different, and as a result, the time of transition from one state to the other may be

different. This is especially so since the genetic backgrounds in which forward and reverse evolution of

multicellularity is being studied are different from each other. In this context, what role, if any, does epistasis

play in this process is also not known and can be a determinant in dictating the evolutionary trajectory of

the cluster size (74). Finally, how cluster size relates with the accumulation of mutations is not yet known.

The relationship between these two variables in the forward and reverse evolution experiment need not be

the same. Determining the answers to these questions related to the forward and backward adaptive

processes forms the basis of the future work in this direction.

Conflict of interest

Authors declare that there is no conflict of interest.

Acknowledgements.

PA was funded by the IITB Institute Post-Doctoral Fellowship. AM is supported by the Council of Scientific

and Industrial Research (CSIR), Government of India, as a Senior Research Fellow (09/087(0873)/2017-

EMR-I).

Figures.

Figure 1. (Top) The experimental evolution of multicellularity under centrifugal selection pressure.

(Bottom) Experimental plan for reverse evolution of unicellularity from multicellularity under centrifugal

selection pressure.

(A)

(B)

(C)

Figure 2. (A) The representative images of yeast multicellular cluster populations during centrifugal
selection. An aliquot of overnight grown yeast cultures (both ancestor and multicellular cluster populations
(MC1-MC6) from different transfer times (T10-T60) were placed on a microscopic slide covered with a cover
slip. Images were taken using a microscope under 40X magnification. **(B)** Smooth (left) and rugose (rough)
(right) morphologies of yeast unicellular and multicellular colonies, respectively. Overnight grown yeast
cultures (both unicellular ancestor and multicellular cluster populations) were streaked on to YPD agar
plates. Plates were allowed to incubate at 30°C until single colonies appear and images of the plates were
then taken. **(C)** Dynamics of changes in yeast cluster size after 60 rounds of centrifugal selection.
Exponentially growing yeast multicellular cluster populations (Line 1-6) were washed with PBS and
analyzed for FSC-A of each cells/cluster using flow cytometer. More than 30,000 clusters per population
were analyzed and the average of the FSC-A at the indicated time of transfer was calculated**.** The
experiment was performed three times and the average and the standard deviation is reported.

Figure 3. Yeast multicellular clusters settle faster with increase in transfer time. Six independent yeast lines (L1-L6) at different time points during forward evolution were inoculated from colonies into 10 mL YPD liquid cultures and grown overnight at 30°C. Each culture tube was vigorously vortexed and placed on a lab bench at room temperature to allow for settling. Images of each population during the settling were taken at different time intervals (0, 15, 30, and 60 min).

467 **(A)**   **(B)**

468
469 **(C)**   **(D)**

470
471
472

Figure 4. (A) Yeast multicellular cluster populations (empty square) show slow growth compared to
unicellular ancestor (closed circles). Overnight yeast clusters and ancestor lines were diluted to an initial
OD_{600} of 0.1 and the optical density was measured for every 1 h. The data represents average of six
independent repeats. **(B)** Relative growth rate of yeast clusters with respect to ancestor. The experiment
was performed in triplicate and the mean and the standard deviation are represented. **(C)** Yeast multicellular
cluster phenotype confers heat stress resistance. Overnight yeast clusters and ancestor lines were diluted
to OD_{600} of 3, 1, 0.3 and 0.1 and spotted on to YPD plates. Plates were incubated at indicated temperatures
and images were taken at different days. MC=multicellular cluster population. **(D)** Effect of ethanol on the
growth kinetics. Yeast multicellular cluster populations exhibit a shorter lag phase indicating resistance to
ethanol stress compared to unicellular ancestor phenotype. Overnight yeast clusters and ancestor lines

were diluted to an initial OD_{600} of 0.1 and exposed to ethanol (2.5 percent). Then the optical density was

measured for every 1 h. Data is shown from the average of the six independent lines. Note that the shorter

lag of the cluster's growth, when compared to the ancestor, is not present when the cultures are grown in

YPD (Figure 2A).

(A)

(B)

491 **Figure 5.** Growth curves of the ancestral and evolved lines in (A) 5% ethanol, and (B) 10% ethanol.

497 **Figure 6.** Mutations in ACE2 coding region in the six evolved lines after 60 transfers. The topmost figure

498 is for Line 1, and the bottommost for Line 6.

501 **Figure 7.** ACE2 mutant alleles from the evolved lines, complementing in an ACE2Δ deletion haploid strain.

503   (A)

504
505
506   (B)
507

508
509
Figure 8. (A) Reversion of rough colony morphology to smooth colony morphology during reverse
evolution from multicellularity to unicellularity (one of the six lines is shown)**.** Overnight grown yeast
cultures (both multicellular cluster population and population evolved from it during reverse evolution) were
streaked on to YPD agar plates. Plates were allowed to incubate at 30°C until single colonies appear and
images of the plates were taken. **(B)** Dynamics of trajectories during evolution of multicellularity (green) and
during evolution of unicellularity (red). Top row (left to right) is data for lines 1-3. Bottom row (left to right) is
data for lines 4-6.
517

520   **Figure 9.** Growth kinetics of ancestor, forward-, and reverse-evolved line. Data for one of the six lines is

521   shown. The data is average for six independent experiments. The standard deviation for the data is less

522   than 3% of all data points.

528 **Figure 10.** Trajectory of evolution of unicellularity starting from the multicellular cluster population in Line
529 1. Six independent repeats of reversal of multicellularity evolution exhibit distinct kinetics of reduction in
530 cluster size. Five of the six trajectories are distinct from reversal of the original unicellular to multicellular
531 cluster evolution (P<0.1, unpaired t-test).

538 **Figure 3.** Mutations in ACE2 coding region in the six evolved lines after 60 transfers. The topmost figure

539 is for Line 1, and the bottommost for Line 6.

(A)

543
544

545   **(B)**

548   **(C)**

549
550
551
Figure 4. (A) Reversion of rough colony morphology to smooth colony morphology during reverse evolution
from multicellularity to unicellularity. Overnight grown yeast cultures (both multicellular cluster population
and population evolved from it during reverse evolution) were streaked on to YPD agar plates. Plates were
allowed to incubate at 30°C until single colonies appear and images of the plates were taken. Images from
the multicellular line 1 (left) and reversed evolved Line1 (right) are shown. **(B)** Dynamic trajectory of
evolution of multicellularity (0-60 transfers) and evolution of unicellularity (60-120 transfers). Green shows

transfers 0-60 when lines are evolved for multicellularity. Red shows transfers 60-120, when lines are

evolved for unicellularity. The top row shows data for Lines 1-3 (left to right) and the bottom row for Lines

4-6 (left to right). The trajectories of evolution of multicellularity and that of its reversal are statistically

distinct from each other for lines 1, 2, 3, and 6 ($P<0.1$, unpaired t-test). **(C)** Cluster size for 6 independent

repeats of reverse evolution of multicellular clusters from Line 1. Five out of the six trajectories are

statistically different from the trajectory of evolution of multicellularity ($P<0.1$, unpaired t-test).

References

1. Anonymous, A Novel Laboratory Activity for Teaching about the Evolution of Multicellularity. *The American Biology Teacher* **76**, 81 (2014).

2. C. Darwin, *On the origin of species by means of natural selection, or preservation of favoured races in the struggle for life* (London : John Murray, 1859, 1859).

3. L. Dollo, The Laws of Evolution. *BULLETIN DE LA SOCIÉTÉ BELGE DE GÉOLOGIE* **VII**, 164-166 (1893).

4. S. D. Durrant, Simpson, George Gaylord. The Major Features of Evolution. Columbia University Press, New York City, New York. xx plus 434 pp., 52 figs., 28 tables. 1953. Price $7.50. *Journal of Mammalogy* **35**, 600-601 (1954).

5. R. C. Lewontin, Is Nature Probable or Capricious? *BioScience* **16**, 25-27 (1966).

6. W. D. Crill, H. A. Wichman, J. J. Bull, Evolutionary reversals during viral adaptation to alternating hosts. *Genetics* **154**, 27-37 (2000).

7. H. Teotónio, M. R. Rose, Variation in the reversibility of evolution. *Nature* **408**, 463-466 (2000).

8. R. E. Lenski, EXPERIMENTAL STUDIES OF PLEIOTROPY AND EPISTASIS IN ESCHERICHIA COLI. I. VARIATION IN COMPETITIVE FITNESS AMONG MUTANTS RESISTANT TO VIRUS T4. (1988).

9. R. E. Lenski, EXPERIMENTAL STUDIES OF PLEIOTROPY AND EPISTASIS IN ESCHERICHIA COLI. II. COMPENSATION FOR MALADAPTIVE EFFECTS ASSOCIATED WITH RESISTANCE TO VIRUS T4. (1988).

10. A. B. Duncan, S. Fellous, O. Kaltz, REVERSE EVOLUTION: SELECTION AGAINST COSTLY RESISTANCE IN DISEASE-FREE MICROCOSM POPULATIONS OF PARAMECIUM CAUDATUM. **65**, 3462-3474 (2011).

11. P. O. Larsson, Cost-effectiveness of misoprostol prevention. *Lakartidningen* **89**, 3657 (1992).

12. S. J. Schrag, V. Perrot, B. R. Levin, Adaptation to the fitness costs of antibiotic resistance in Escherichia coli. *Proc Biol Sci* **264**, 1287-1291 (1997).

13. J. J. Bull *et al.*, Exceptional convergent evolution in a virus. *Genetics* **147**, 1497-1507 (1997).

14. M. Kaltenbach, C. J. Jackson, E. C. Campbell, F. Hollfelder, N. Tokuriki, Reverse evolution leads to genotypic incompatibility despite functional and active site convergence. *Elife* **4** (2015).

15. S. A. West, R. M. Fisher, A. Gardner, E. T. Kiers, Major evolutionary transitions in individuality. *Proc Natl Acad Sci U S A* **112**, 10112-10119 (2015).

16. R. K. Grosberg, R. R. Strathmann, The Evolution of Multicellularity: A Minor Major Transition? *Annual Review of Ecology, Evolution, and Systematics* **38**, 621-654 (2007).

17. A. Rokas, The origins of multicellularity and the early history of the genetic toolkit for animal development. *Annual review of genetics* **42**, 235-251 (2008).

18.   L. Wolpert, E. Szathmáry, Multicellularity: Evolution and the egg. *Nature* **420**, 745-745 (2002).

19.   J. H. Koschwanez, K. R. Foster, A. W. Murray, Sucrose utilization in budding yeast as a model for
the origin of undifferentiated multicellularity. *PLoS biology* **9**, e1001122-e1001122 (2011).

20.   T. Pfeiffer, S. Bonhoeffer, An evolutionary scenario for the transition to undifferentiated
multicellularity. *Proceedings of the National Academy of Sciences* **100**, 1095-1098 (2003).

21.   I. Ispolatov, M. Ackermann, M. Doebeli, Division of labour and the evolution of multicellularity.
Proceedings. Biological sciences **279**, 1768-1776 (2012).

22.   M. Willensdorfer, ON THE EVOLUTION OF DIFFERENTIATED MULTICELLULARITY. *Evolution*
63, 306-323 (2009).

23.   A. Amado, C. Batista, P. R. A. Campos, A mechanistic model for the evolution of multicellularity.
bioRxiv 10.1101/115832, 115832 (2017).

24.   W. C. Ratcliff, R. F. Denison, M. Borrello, M. Travisano, Experimental evolution of multicellularity.
Proc Natl Acad Sci U S A **109**, 1595-1600 (2012).

25.   W. C. Ratcliff, J. T. Pentz, M. Travisano, TEMPO AND MODE OF MULTICELLULAR ADAPTATION
IN EXPERIMENTALLY EVOLVED SACCHAROMYCES CEREVISIAE. *Evolution* **67**, 1573-1581
(2013).

26.   J. H. Koschwanez, K. R. Foster, A. W. Murray, Sucrose utilization in budding yeast as a model for
the origin of undifferentiated multicellularity. *PLoS Biol* **9**, e1001122 (2011).

27.   M. Rebolleda-Gomez, W. Ratcliff, M. Travisano, *Adaptation and Divergence during Experimental*
Evolution of Multicellular Saccharomyces cerevisiae (2012), 10.7551/978-0-262-31050-5-ch014,
pp. 99-104.

28.   W. W. Driscoll, M. Travisano, Synergistic cooperation promotes multicellular performance and
unicellular free-rider persistence. *Nat Commun* **8**, 15707 (2017).

29.   S. Jacobeen *et al.*, Geometry, packing, and evolutionary paths to increased multicellular size. *Phys*
Rev E **97**, 050401 (2018).

30.   M. Rebolleda-Gomez, M. Travisano, The Cost of Being Big: Local Competition, Importance of
Dispersal, and Experimental Evolution of Reversal to Unicellularity. *Am Nat* **192**, 731-744 (2018).

31.   J. J. Kuzdzal-Fick, L. Chen, G. Balázsi, Disadvantages and benefits of evolved unicellularity versus
multicellularity in budding yeast. *Ecology and evolution* **9**, 8509-8523 (2019).

32.   M. D. Herron *et al.*, De novo origins of multicellularity in response to predation. *Scientific Reports*
9, 2328 (2019).

33.   B. Oud *et al.*, Genome duplication and mutations in ACE2 cause multicellular, fast-sedimenting
phenotypes in evolved Saccharomyces cerevisiae. *Proc Natl Acad Sci U S A* **110**, E4223-4231
(2013).

34.   W. C. Ratcliff, J. D. Fankhauser, D. W. Rogers, D. Greig, M. Travisano, Origins of multicellular
evolvability in snowflake yeast. *Nat Commun* **6**, 6102 (2015).

35.   J. J. Kuzdzal-Fick, L. Chen, G. Balazsi, Disadvantages and benefits of evolved unicellularity versus
639   multicellularity in budding yeast. *Ecol Evol* **9**, 8509-8523 (2019).
640   36.   A. S. Trigos, R. B. Pearson, A. T. Papenfuss, D. L. Goode, Altered interactions between unicellular
641   and multicellular genes drive hallmarks of transformation in a diverse range of solid tumors. *Proc*
642   *Natl Acad Sci U S A* **114**, 6406-6411 (2017).
643   37.   V. Rossetti, B. E. Schirrmeister, M. V. Bernasconi, H. C. Bagheri, The evolutionary path to terminal
644   differentiation and division of labor in cyanobacteria. *J Theor Biol* **262**, 23-34 (2010).
645   38.   B. E. Schirrmeister, A. Antonelli, H. C. Bagheri, The origin of multicellularity in cyanobacteria. *BMC*
646   *Evol Biol* **11**, 45 (2011).
647   39.   K. D. Breunig, P. Kuger, Functional homology between the yeast regulatory proteins GAL4 and
648   LAC9: LAC9-mediated transcriptional activation in Kluyveromyces lactis involves protein binding to
649   a regulatory sequence homologous to the GAL4 protein-binding site. *Mol Cell Biol* **7**, 4400-4406
650   (1987).
651   40.   J. Tan *et al.*, Copper oxide nanoparticles promote the evolution of multicellularity in yeast.
652   *Nanotoxicology* 10.1080/17435390.2018.1553253, 1-9 (2019).
653   41.   E. A. Hope, M. J. Dunham, Ploidy-regulated variation in biofilm-related phenotypes in natural
654   isolates of Saccharomyces cerevisiae. *G3 (Bethesda)* **4**, 1773-1786 (2014).
655   42.   T. E. Torchia, J. E. Hopper, Genetic and molecular analysis of the GAL3 gene in the expression of
656   the galactose/melibiose regulon of Saccharomyces cerevisiae. *Genetics* **113**, 229-246 (1986).
657   43.   S. Harju, H. Fedosyuk, K. R. Peterson, Rapid isolation of yeast genomic DNA: Bust n' Grab. *BMC*
658   *Biotechnology* **4**, 8 (2004).
659   44.   F. Bertels, C. Leemann, K. J. Metzner, R. R. Regoes, Parallel Evolution of HIV-1 in a Long-Term
660   Experiment. *Molecular Biology and Evolution* **36**, 2400-2414 (2019).
661   45.   R. Woods, D. Schneider, C. L. Winkworth, M. A. Riley, R. E. Lenski, Tests of parallel molecular
662   evolution in a long-term experiment with Escherichia coli. *Proceedings of the National*
663   *Academy of Sciences* **103**, 9107-9112 (2006).
664   46.   S. F. Bailey, F. Blanquart, T. Bataillon, R. Kassen, What drives parallel evolution?: How population
665   size and mutational variation contribute to repeated evolution. *Bioessays* **39**, 1-9 (2017).
666   47.   P. B. Rainey, M. Travisano, Adaptive radiation in a heterogeneous environment. *Nature* **394**, 69-
667   72 (1998).
668   48.   B. Kerr, M. A. Riley, M. W. Feldman, B. J. M. Bohannan, Local dispersal promotes biodiversity in a
669   real-life game of rock-paper-scissors. *Nature* **418**, 171-174 (2002).
670   49.   C. A. Solari, S. Ganguly, J. O. Kessler, R. E. Michod, R. E. Goldstein, Multicellularity and the
671   functional interdependence of motility and molecular transport. *Proceedings of the National*
672   *Academy of Sciences of the United States of America* **103**, 1353-1358 (2006).
673   50.   J. T. Bonner, Perspective: the size-complexity rule. *Evolution; international journal of organic*
674   *evolution* **58**, 1883-1890 (2004).

51.   J. T. Bonner, The origins of multicellularity. *Integrative Biology: Issues, News, and Reviews* **1**, 27-
36 (1998).

52.   M. E. Boraas, D. B. Seale, J. E. Boxhorn, Phagotrophy by a flagellate selects for colonial prey: A
possible origin of multicellularity. *Evolutionary Ecology* **12**, 153-164 (1998).

53.   R. H. Kessin, G. G. Gundersen, V. Zaydfudim, M. Grimson, How cellular slime molds evade
nematodes. *Proceedings of the National Academy of Sciences* **93**, 4857-4861 (1996).

54.   S. Smukalla *et al.*, FLO1 Is a Variable Green Beard Gene that Drives Biofilm-like Cooperation in
Budding Yeast. *Cell* **135**, 726-737 (2008).

55.   L. Váchová *et al.*, Architecture of developing multicellular yeast colony: spatio-temporal expression
of Ato1p ammonium exporter. *Environmental Microbiology* **11**, 1866-1877 (2009).

56.   H. W. Augustin, G. Kopperschläger, H. Steffen, E. Hofmann, Hexokinase as limiting factor of
anaerobic glucose consumption of Saccharomyces carlsbergensis NCYC74. *Biochimica et*
biophysica acta **110**, 437-439 (1965).

57.   D. E. Ferrier, P. W. Holland, Ancient origin of the Hox gene cluster. *Nature reviews. Genetics* **2**,
33-38 (2001).

58.   A. Kusserow *et al.*, Unexpected complexity of the Wnt gene family in a sea anemone. *Nature* **433**,
156-160 (2005).

59.   N. King, S. B. Carroll, A receptor tyrosine kinase from choanoflagellates: Molecular insights into
early animal evolution. *Proceedings of the National Academy of Sciences* **98**, 15032-15037 (2001).

60.   M. Abedin, N. King, The premetazoan ancestry of cadherins. *Science (New York, N.Y.)* **319**, 946-
948 (2008).

61.   S. A. Nichols, W. Dirks, J. S. Pearse, N. King, Early evolution of animal cell signaling and adhesion
genes. *Proceedings of the National Academy of Sciences of the United States of America* **103**,
12451-12456 (2006).

62.   L. King, G. Butler, Ace2p, a regulator of CTS1 (chitinase) expression, affects pseudohyphal
production in Saccharomyces cerevisiae. *Curr Genet* **34**, 183-191 (1998).

63.   H. J. McBride, Y. Yu, D. J. Stillman, Distinct regions of the Swi5 and Ace2 transcription factors are
required for specific gene activation. *J Biol Chem* **274**, 21029-21036 (1999).

64.   E. Gil de Prado *et al.*, Quantitative analysis of morphological changes in yeast colonies growing on
solid medium: the eccentricity and Fourier indices. *Yeast* **31**, 431-440 (2014).

65.   J. Smith, D. C. Queller, J. E. Strassmann, Fruiting bodies of the social amoeba Dictyostelium
discoideum increase spore transport by Drosophila. *BMC Evol Biol* **14**, 105 (2014).

66.   S. Smukalla *et al.*, FLO1 is a variable green beard gene that drives biofilm-like cooperation in
budding yeast. *Cell* **135**, 726-737 (2008).

67.   G. Zara, S. Zara, C. Pinna, S. Marceddu, M. Budroni, FLO11 gene length and transcriptional level
affect biofilm-forming ability of wild flor strains of Saccharomyces cerevisiae. *Microbiology*
(Reading) **155**, 3838-3846 (2009).

68.   M. Rebolleda-Gomez, M. Travisano, Adaptation, chance, and history in experimental evolution
713   reversals to unicellularity. *Evolution* **73**, 73-83 (2019).
714   69.   O. Fang *et al.*, Amn1 governs post-mitotic cell separation in Saccharomyces cerevisiae. *PLoS*
715   *Genet* **14**, e1007691 (2018).
716   70.   M. Travisano, J. A. Mongold, A. F. Bennett, R. E. Lenski, Experimental tests of the roles of
717   adaptation, chance, and history in evolution. *Science* **267**, 87-90 (1995).
718   71.   J. B. Losos, T. R. Jackman, A. Larson, K. Queiroz, L. Rodriguez-Schettino, Contingency and
719   determinism in replicated adaptive radiations of island lizards. *Science* **279**, 2115-2118 (1998).
720   72.   R. A. Neher, B. I. Shraiman, D. S. Fisher, Rate of adaptation in large sexual populations. *Genetics*
721   **184**, 467-481 (2010).
722   73.   S. Chattopadhyay *et al.*, High frequency of hotspot mutations in core genes of Escherichia coli due
723   to short-term positive selection. *Proc Natl Acad Sci U S A* **106**, 12412-12417 (2009).
724   74.   P. C. Phillips, Epistasis--the essential role of gene interactions in the structure and evolution of
725   genetic systems. *Nat Rev Genet* **9**, 855-867 (2008).