"Fishing for contact: Modeling perivascular glioma invasion in the zebrafish brain."

Umans, R.A.[1], ten Kate, M.[2], Pollock, C.[2], & Sontheimer, H.[1,2]

Center for Glial Biology in Health, Disease, and Cancer, The Fralin Biomedical Research Institute at VTC, Roanoke, VA, 24016, USA [1]
School of Neuroscience, Sandy Hall, 210 Drillfield Dr.,Virginia Tech, Blacksburg, VA, 24061, USA[2]
Corresponding author email: sontheim@vt.edu

Abstract

Glioblastoma multiforme (GBM) is a highly invasive, central nervous system (CNS) cancer for which there is no a cure. Invading tumor cells evade treatment, limiting the efficacy of current standard of care treatments. Understanding the underlying invasive behaviors that support tumor growth may allow for generation of novel GBM therapies. Zebrafish (*Danio rerio*) are attractive for genetics, live imaging, and in recent years, have emerged as a model system suitable for cancer biology research. While other groups have studied CNS tumors using zebrafish, few have concentrated on the invasive behaviors supporting the development of these diseases. Previous studies demonstrated that one of the main mechanisms of GBM invasion is perivascular invasion, i.e. single tumor cell migration along blood vessels. Here, we characterize phenotypes, methodology, and potential therapeutic avenues for utilizing zebrafish to model perivascular GBM invasion. Using patient derived xenolines or an adherent cell line, we demonstrate tumor expansion within the zebrafish brain. Within 24 hours post-intracranial injection, D54-MG-tdTomato glioma cells produce finger-like projections along the zebrafish brain vasculature. As few as 25-50 GBM cells were sufficient to promote single cell vessel co-option. Of note, these tumor-vessel interactions are CNS specific, and do not occur on pre-existing blood vessels when injected into the animal's peripheral tissue. Tumor-vessel interactions increase over time and can be pharmacologically disrupted through inhibition of Wnt signaling. Therefore, zebrafish serve as a favorable model system to study perivascular glioma invasion, one of the deadly characteristics that make GBM so difficult to treat.

Introduction

Brain tumors are one of the most devastating forms of cancer, as they grow within an organ of limited regenerative capacity and can be highly invasive. Additionally, drug delivery to brain tumors is challenging due to the blood-brain barrier (BBB), a physiological structure that limits the open flow of foreign compounds into the brain tissue. Glioblastoma multiforme (GBM) is the most common, primary, malignant brain tumor and is newly diagnosed in around 13,000 patients every year in the United States (Ostrom et al., 2017). The standard of care is a three-pronged approach of surgical resection, radiation, and chemotherapy (Stupp et al., 2005). Even after over a decade of implementation, this multifaceted treatment regimen extends survival to only 14-19

months (Feng et al., 2017). These aggressive therapies mainly target the main tumor mass that is most easily identified by MRI for surgical removal or therapeutics that target actively dividing cells. Even in the era of personalized medicine, every GBM case presents with a different set of mutations, making it difficult to standardize a therapeutic regimen that effectively treats all patients. Research efforts have shown that the genetic heterogeneity of GBM tumors results in varied patient response to specific therapies, and individual mutations may not be present in the majority of primary GBM patients (Wang X, 2013, Hegi et al., 2005, Mellinghoff et al., 2005, McNeill et al., 2018). Therefore, targeting how these brain tumor cells behave in the context of their mutations or environment may elucidate avenues for more promising therapies that have a wider reach within the GBM patient population (Umans and Sontheimer, 2018).

One of the main routes of GBM infiltration is cellular invasion within the tight spaces of the brain parenchyma. While the main neoplasm is targeted for surgical resection, migratory cells can remain undetected by MRI and form satellite tumors away from the original mass. The avenues in which these nomadic cells travel are via white matter tracts and blood vessels, with the perivascular space possessing little physical resistance (Cuddapah et al., 2014, Farin et al., 2006). Previous work using a mouse model intracranially injected with human GBM cells demonstrated human tumor cell tropism along the mouse brain vasculature and, after a 2-4 weeks of implantation, GBM cells caused micro-breaches at the BBB (Watkins et al., 2014). Furthermore, another GBM rodent model demonstrated the Wnt signaling cascade is necessary for glioma-vessel interactions (Griveau et al., 2018). However, *in vivo* mouse models require the injection of thousands of cells, use of immune-compromised animals, and performing advanced surgical and imaging techniques. In recent years, groups have established zebrafish (*Danio rerio*) specifically modeling the growth and invasion of brain tumors (Vittori et al., 2015, Welker et al., 2016). However some of these studies inject brain tumor cells into the yolk of the animal and not the brain, inject hundreds of cells which results in a massive cell clump, or have not carefully visualized the relationship these tumor cells have with the brain vasculature (Welker et al., 2016, Zeng et al., 2017, Yang et al., 2013). While these zebrafish models provide exciting avenues to study human disease in a simpler vertebrate system, most have looked at the molecular features of tumors and not the behaviors that make gliomas so hard to treat. The emergence of an *in vivo* model that could more easily visualize and manipulate perivascular interactions important for tumor invasion would greatly benefit the field of GBM biology.

In the present study, we hypothesize that zebrafish could be utilized to specifically model perivascular glioma invasion. To test and visualize this hypothesis, we intracranially injected red fluorescently labeled human glioma cells into the translucent zebrafish vascular reporter line *Tg(fli1a:eGFP)^{y1};casper*, whereby all blood vessels are labeled with enhanced green fluorescent protein (eGFP). We injected both an adult glioma cell line (D54-MG) and pediatric patient derived xenoline (D2159MG) into *Tg(fli1a:eGFP)^{y1};casper* zebrafish and witnessed both of these glioma lines attach to the zebrafish vasculature. Gliomas retain salient disease features in the zebrafish brain, such as attachment to secondary structures (blood vessels) and tissue necrosis. Injected animals were imaged daily and confocal microscopy revealed glioma cells

survive and expand within the zebrafish brain. Perivascular glioma invasion was dependent on the number of tumor cells injected. When more than 50 cells were present, tumors survived and grew but few cells contacted the vasculature. Tumor cell vessel co-option increased over time along the vessel and at vessel branch points. Glioma-vessel interactions were also dependent on the CNS microenvironment, as glioma cells injected into the animal's peripheral tissue failed to interact with pre-existing vasculature. Furthermore, we highlight the small molecule XAV939, an inhibitor of the Wnt signaling pathway, pharmacologically disrupted glioma cell-vascular interactions in zebrafish brain. These proof-of-principles studies demonstrate valuable features of this established zebrafish model of perivascular glioma invasion, which could lead to the discovery of newly needed GBM therapeutics.

Methods

Cell Culture: D54-MG human glioma adherent cell lines (WHO IV, glioblastoma multiforme; gifted by Dr. D. Bigner, Duke University, Durham, NC) were genetically modified to express tdTomato or eGFP in a previous study (Watkins et al., 2014). D54-MG cells were maintained in Dulbecco's modified Eagle's Media (DMEM) (Fisher #11320082) supplemented with 7% fetal bovine serum (Lot 105313, Aleken), and kept in a cell culture incubator with 10% CO_2 at 37°C

D2159MG and GBM22 patient-derived xenograft (PDX) lines (gifted by Darell D. Bigner, MD, PhD and Stephen T. Keir, DrPH, MPH, Duke University, Tisch Brain Tumor Center and Dr. Yancey Gillespie, University of Alabama at Birmingham, Birmingham, AL, USA, respectively) were from primary brain tissue and maintained by serial passage in the flank of athymic nude mice, as previously described (Watkins et al., 2014). PDX cells were grown in a DMEM complete media supplemented with B-27 (Fisher #12587010), Amphotericin (1:100; Fisher #BP264550), Sodium pyruvate (1:100;ThermoFisher #11360-070), EGF and FGF growth factors (10ng/ml each, final ;ThermoFisher #PHG0266 and #PHG0315), and Gentamicin (1:1,000;Fisher #BW17 518Z). PDX cells were maintained in a cell culture incubator with 10% CO_2 at 37°C. The D2159MG pediatric glioma cells were genetically modified with a lentivirus (MGH Vector Core) to constitutively express mCherry. To visualize unlabeled GBM22 cells, cells were labeled overnight with 5µM of the lipophilic DiL dye (Invitrogen #V22885).

For cell counting, D54-MG cells were dissociated using Trypsin-EDTA (ThermoFisher/LifeTech #25300054) and PDX cells were dissociated using Accutase (Sigma #A6964). Cells were counted on a hemocytometer and re-suspended at 25,000cells per µL in sterile PBS prior to microinjection.

Acute Mouse Brain Slice Invasion and Immunohistochemistry (IHC): Experiments were performed in accordance with the University of Alabama, Birmingham Institutional Animal Care and Use Committee (IACUC). *In situ* tumor invasion was performed as previously described (Watkins and Sontheimer, 2011). For IHC, 300µM brain sections were acutely sliced from 10 week old *NG2:dsRed* mice and seeded with 80,000 D54-MG-eGFP tumor cells on multi-well filter chambers (Fisher #08-771). Tumor cells

invaded brain slices for 4 hours on filter chambers in a 10% CO_2 cell culture incubator. Brain slices were fixed in 4% PFA and used for subsequent IHC. Slices were rinsed in PBST for 15 minutes, permeated with proteinase K (20mg/ml stock, 1:800) for 15 minutes at room temperature, and then blocked in 10% goat serum for 1 hour at room temperature. Sections were stained with a mouse-anti-Laminin primary antibody (Sigma L8271,1:500) overnight at 4°C. The next morning, slices were washed in PBST for 1.5 hours, stained with donkey-anti-mouse- Alexa 555 secondary antibody (Invitrogen A31570,1:500) for 3 hours at room temperature, washed in PBST for 1.5 hours, and mounted with Fluoromount (VWR 100502-406) on glass slides. Z-stacks were acquired with an Olympus Fluoview FV1000 confocal microscope and a 60X(NA1.42) oil objective.

Zebrafish: Adult zebrafish were maintained according to Virginia Tech IACUC guidelines on a 14/10 hour light/dark cycle in a Tecniplast system at 28.5°C, pH 7.0, and 1000 conductivity. The *Tg(fli1a:eGFP)[y1]* line was crossed onto the *casper* background (White et al., 2008) to generate genetically translucent vascular reporter lines. These animals were specific-pathogen free and came from the Sinnhuber Aquatic Research Laboratory at Oregon State University. The *Tg(glut1b:mCherry)* line was obtained from Dr. Michael Taylor's laboratory at the University of Wisconsin-Madison. For all experiments, embryos were collected from multiple pair-wise crosses. Embryos were maintained in embryo water (1.5g Instant ocean salt per 5L RO water) at 28.5°C. For studies with the *Tg(glut1b:mCherry)* line, 24hour post-fertilization embryos were treated with 200µM N-Phenylthiourea (Sigma # P7629) to prevent melanocyte formation. Embryos for tumor cell injections were gradually acclimated to develop at 32°C, as to accommodate a more desirable temperature for glioma cell growth.

Tumor Microinjections: Healthy animals (i.e. no developmental abnormalities) were screened before tumor implantation. 3 days post fertilization (dpf) *Tg(fli1a:eGFP)[y1];casper* or *Tg(glut1b:mCherry)* larvae were anesthetized with 0.04% MS-222 (Sigma #A5040) prior to microinjection. During anesthesia, thin wall, glass capillary needles with filament (World Precision Instruments #TW150F-4) were pulled with program 4 "Pro-Nuclear Injection" (Heat: 460, Pull:90, Vel: 70, Delay: 70, Pressure: 200, Ramp: 485) on a horizontal pipette puller (Sutter, Model P-1000). This program creates needles with a 0.7µM tip and a taper of 6-7mm. 1µL of Phenol Red (Sigma #P0290) was added to 9µL of the 25,000 glioma cell/µL solution mixture. Capillaries were positioned in a round, glass pipette holder and back loaded with 1uL of human glioma cell mixture. Each loaded needle was calibrated using a pneumatic pump and micrometer (Carolina Biological Supply #591430) as to calculate the bolus and corresponding cell number per injection. Animals were positioned dorsal side up for intracranial injection or on their sides for trunk injection in a homemade agarose injection mold and then microinjected with the desired amount of cells. After injections, animals recovered and were tracked in individual wells of a 24-well plate.

Live Confocal Microscopy: Animals were removed from multi-well plates with a glass Pasteur pipette and anesthetized with 0.04% MS-222 in individual 35mm petri dishes containing a 14mm glass coverslip (MatTek Corporation #P35G-1.5-14-C). A 1.2% low

melting point agarose (ThermoFisher #16520050) solution was made in embryo water for embedding. MS-222 was removed from the animal and a bolus of warm agarose the size of the coverslip was added to the animal/cover glass. A dissection probe was used to align animals either dorsal side down for brain imaging or sagitally for trunk imaging. After the agarose solidified, the petri dish was filled with 0.04% MS-222 to keep the animal immobilized and the dish was wrapped with parafilm along the edges. Live confocal microscopy was performed to take z-stacks at optimal section thickness of animals with D54-MG and D2159MG cells with an Olympus Fluoview FV1000 confocal microscope and a 40X(NA0.75) objective. The Nyquist optimal section thickness value for z-stacks was calculated through an algorithm in the Olympus Fluoview software for all images. For whole zebrafish brain imaging after GBM22 injections, a Nikon A1R confocal microscope with a 10X(NA 0.45) objective and 1.5 optical zoom was utilized. Because these microscopes are upright, the glass coverslip bottom dishes containing the animals were flipped upside down so that the glass side made contact with the objective. Supplemental Movie 1 was generated from a Maximum Projection Image in NIS-Elements software.

Small Molecule Treatment: After initial tumor implantation at 3dpf and subsequent imaging, tumors grew overnight before animals were unbiasedly sorted into treatment groups at 4dpf. The Wnt signaling inhibitor, XAV939, (Cayman Chemical #13596) was solubilized in dimethyl sulfoxide (DMSO) to generate 10mM stock solutions and used at a final concentration of 30µM in zebrafish egg water.

Glioma-Vessel Interaction Analysis: To assess the percentage of human glioma cells interacting with the vasculature, OIB z-stack files were analyzed in the Fluoview software. In brief, the total number of visible human glioma cells were initially counted. The orthogonal view option was used to detect whether the signal from a cell was associated with the signal from a blood vessel or blood vessel branch point as seen in both XZ and YZ planes. The total number of interacting cells were then divided by the total number of injected cells.

Tg(glut1b:mCherry) **Volumetric Analysis**: Fiji- ImageJ and NIS-Elements software were utilized to measure the volume and signal intensity of *glut1* in tumor-associated and non-tumor associated vessels. In brief, confocal z-stacks were opened as split channels in FIJI and tiff files were saved to import into NIS-Elements. After importing, channels were merged together, the document was calibrated, and files were saved as a ND2 file. Sections around vessels of interest were cropped out of the whole file and a line was drawn along a vessel where it was ensheathed by a tumor cell body. The rotating rectangle feature was utilized and this selected area underwent a binary threshold and 1X clean and smooth options. The same procedure and size line was drawn along a similarly placed vessel without a tumor cell in the contralateral brain as a paired control vessel to measure the volume and intensity in a similar manner. Measurements were generated in the NIS Elements "3D Object Measurements" table.

Statistical Analysis: For analysis of perivascular glioma invasion at developmental time points 3-7dpf, a One-way ANOVA with multiple comparisons test was performed

using GraphPad Prism software to determine *p*-values. For small molecule treatment experiments, a two-tailed**,** unpaired student's *t*-test was performed using GraphPad Prism software to determine *p*-values. For volumetric vessel and *glut1* fluorescence analyses, a two-tailed, paired student's *t*-test was performed using GraphPad Prism software to determine *p*-values.

<u>Results</u>

Human tumors retain salient features in a cell number-dependent manner in the zebrafish brain.

Previous studies demonstrated zebrafish as a suitable model to elucidate CNS cancer biology. However, most of these studies inject hundreds of cells into the animal, including the controversial U87-MG cell line, and did not examine perivascular glioma invasion (Yang et al., 2013, Zeng et al., 2017, Vargas-Patron et al., 2019, Allen et al., 2016). As a means to ask whether glioma-vessel interactions may occur in the zebrafish brain, we opted to inject fewer cells than previously published. Animals from the optically translucent *Tg(fli1a:eGFP)^{y1};casper* vascular reporter line were injected with 25-50 human D54-MG-tdTomato cells at 3 dpf, a time point when the BBB has started forming and sufficient CNS angiogenesis has occurred (Umans et al., 2017). Specific pathogen-free (SPF) animals were used for these brain tumor studies to avoid confounding factors such as *Pseudoloma neuropihila*, a pathogen that resides in the zebrafish CNS and is common in zebrafish colonies. Similar to the behavior of the D54-MG tumor line in acute mouse brain slices (**Figure 1A**), D54-MG glioma cells attached to the zebrafish brain vasculature within 24 hours post-intracranial injection (**Figure 1B,C**). We also performed microinjection into non-SPF animals and witnessed glioma-vascular interactions (**Supplemental Figure 1**), suggesting that while maintaining a SPF colony is the best practice for studying diseases of the CNS, it does not affect the growth of human brain cancer cells regarding the zebrafish brain vasculature. This data suggests conserved signaling mechanisms exist to promote the interaction between glioma cells and the developing zebrafish vessel network.

To determine whether glioma-vascular interactions would still be present after injecting a greater volume of cells, we intracranially injected a cell bolus >50 glioma cells in 3dpf *Tg(fli1a:eGFP)^{y1};casper* animals. We observed that the cells became more easily trapped within the ventricular space, had less of a "choice" to migrate within a 4 day timespan, and while present, had infrequent projections towards the surrounding vasculature compared to animals injected with 25-50 cells. (**Figure 1D-G**). These >50 cell compacted tumors were reminiscent of previously published zebrafish glioma studies that, due to cell number and size of the zebrafish brain versus human cells, could not examine the specific behaviors associated with perivascular glioma invasion (Zeng et al., 2017). However, when 25-50 cells were injected into the midbrain-hindbrain boundary at 3dpf, we witnessed more frequent extension of finger-like processes as witnessed in GBM mouse models and human brain (**Figure1D**) (Watkins et al., 2014). Other salient phenotypes such as brain necrosis were also dependent on cell number, as clumps with >50 cells would generate necrosis in the developing

zebrafish brain days post-injection (**Figure 1G**). While survival studies were not in the scope of this study, some animals would succumb to the disease 4 days post-injection (data not shown). Previous studies report a median survival time of 10 days post injection in zebrafish with glioma xenografts of the preferred cell number (~25-50 cells) used in our study (Welker et al., 2016). Taken together, these experiments reveal salient features of GBM and the process of perivascular invasion occur in the developing brain of our zebrafish glioma model.

Glioma cells actively invade the brain along the zebrafish vasculature.

To analyze the dynamics of perivascular glioma invasion in the developing zebrafish brain, we microinjected 25-50 D54-MG-tdTomato tumor cells into the midbrain-hindbrain boundary of 3dpf *Tg(fli1a:eGFP)^{y1};casper* larvae and monitored tumor behavior until 7dpf (**Figure 2A**). Because daily live imaging is easy to perform with the zebrafish model system, we wanted to assess tumor burden after the onset of injection and subsequent days. After tumor microinjection, animals recovered for at least one hour, were anesthetized, and then embedded for live confocal microscopy to assess initial tumor volume. As shown by representative images in **Figure 2**, the few D54-MG-tdTomato glioma cells implanted at 3dpf (**Figure 2B**) not only survived in the zebrafish brain but they also expanded and migrated along the vascular network by 7dpf (**Figure 2C**). These glioma-vessel interactions increased over time along the main vessel wall (**Figure 2D**) and at branch points (**Figure 2E**) as previously described in a GBM mouse model (Watkins et al., 2014). Furthermore, these interactions occurred with both pediatric or adult glioma cells as well as in adherent and PDX cell lines (**Figure 3**). With a variety of PDX lines at our disposal, we labeled the GBM22 PDX line with a lipophilic dye to visualize tumor cells post-intracranial injection. Of note, lipophilic dye did not permeate the cell and its finer processes after injection, and it was determined that tumor cells that are transgenically labelled to express a fluorophore are more suitable for these imaging studies (**Supplemental Figure 2**). These results demonstrate that even though glioma injections occurred in the developing brain, conserved signaling pathways support perivascular glioma invasion over time in our zebrafish model.

Tumor microenvironment affects perivascular glioma invasion in vivo.

Cancer arises when normal cells within an organ become misprogrammed. Therefore, it is logical to inject tumor cells in a tissue in which those cells naturally thrive. Prior studies have injected non-CNS cancer cells into the periphery of the zebrafish to assess tumor associations with blood vessels (Zhao et al., 2011, Stoletov et al., 2007). To validate the importance of the microenvironment in our perivascular GBM invasion model, we injected 25-50 D54-MG-tdTomato cells in the trunk near the dorsal fin of 3dpf *Tg(fli1a:eGFP)^{y1};casper* animals. Surprisingly, these glioma cells not only survived, but they dramatically moved rostrally towards the brain away from the dorsal fin area in 7dpf animals (**Figure 4A-C**). These cells extended long projections during migration (**Supplemental Movie 1**), and unlike when in the brain, these cells did not attach to non-lymphatic, pre-existing blood vessels. This data suggests there are unique trophic factors within the brain environment that attracts glioma cells to the zebrafish vascular

network. Interestingly, we observed instances of tumor-induced angiogenesis, a biology witnessed in other zebrafish cancer models (**Figure 4E,F**) (Gamble et al., 2018, Zhao et al., 2011). These experiments highlight the importance of the microenvironment when modeling and quantifying glioma-vessel interactions.

Gross vessel morphology and glut1 expression is not altered after acute tumor invasion.

After determining that tumor cells failed to attach to pre-existing, non-CNS vasculature, we wanted to further investigate other parameters post-invasion. Due to the reports of vascular tone alterations and BBB disruption in a GBM mouse model and patient tissue, we assessed the zebrafish brain vascular morphology after 4 days of glioma invasion (Liebner et al., 2000, Watkins et al., 2014). We first asked whether vessel volume was changed in our zebrafish model of perivascular glioma invasion. To answer this question, 25-50 D54-MG-eGFP cells were injected into the mid-brain hind-brain boundary of 3dpf *Tg(glut1b:mCherry)* animals. After 4 days of invasion, we performed live confocal imaging of injected animals. We then utilized volumetric analyses of vessels without tumor burden and vessels on the contralateral brain region contacted by D54-MG-eGFP tumor cells (**Figure 5A,B**). While tumor cells readily ensheathed vessels in the zebrafish brain, we did not find any difference in vessel volume 4 days post-invasion (**Figure 5C**). As the *Tg(glut1b:mCherry)* line expresses mCherry under the control of a reliable BBB marker, *glut1*, we next analyzed the mCherry fluorescence signal of tumor and non-tumor associated vessels in 7dpf animals (Umans et al., 2017). Data from fluorescence intensity analysis revealed that *glut1* expression did not differ between vessels co-opted by a tumor cell soma and vessels in a similar location on the contralateral side of the brain (**Figure 5D**). These live imaging results suggest that co-opted vessels are not visibly compromised and the BBB specific marker *glut1* was not altered after this acute time of invasion.

Glioma-vessel interactions can be pharmacologically inhibited in vivo.

In order to demonstrate whether glioma-vessel interactions could be pharmacologically disrupted *in vivo*, we sought to perform proof-of-principle experiments targeting pathways important for glioma-vessel interactions in GBM mouse models, such as Wnt signaling (Griveau et al., 2018). Therefore, we hypothesized that conserved pathways would maintain perivascular glioma invasion in zebrafish brain and by adding a small molecule inhibitor of Wnt signaling (XAV939), we predicted we would visualize fewer cellular interactions. To test this hypothesis, we intracranially injected 3dpf *Tg(fli1a:eGFP)^{y1};casper* animals with 25-50 D54-MG-tdTomato cells at the midbrain-hindbrain boundary and 4dpf animals were blindly sorted into DMSO vehicle control or XAV939 treatment groups. After 48 hours of bath application, confocal images revealed significantly fewer glioma-vessel interactions (21.57% ± 4.908) in XAV939 treated animals compared to control treated animals (41.5% ± 7.106) (**Figure 6C**). These studies suggest that conserved signaling pathways in mammals attract glioma cells to the zebrafish brain vasculature and this *in vivo* system is amenable to chemical perturbation.

Discussion

GBM is a fatal primary brain tumor studied in an extensive array of models. More recently, research groups have implemented xenograft studies to assess glioma invasion in the developing zebrafish. Zebrafish are genetically tractable, share a similar body plan and genetic conservation to mammals, and are optically transparent. The zebrafish does not develop a functional adaptive immune system until 4 weeks of age and therefore circumvents the need for requiring immunosuppressive chemicals or special animals for injection at early time points (Trede et al., 2004). The adult mouse brain contains roughly 700 times more neuronal cells than the larval zebrafish brain and thus fewer tumor cells can be injected to visualize similar phenotypes (Friedrich et al., 2013, Herculano-Houzel et al., 2006). These features have presented zebrafish to cancer scientists as an attractive model for dissecting tumor biology and finding new therapeutics (Zeng et al., 2017, Hason and Bartunek, 2019).

As the zebrafish model has emerged as a suitable system for dissecting cancer biology, these studies focus mainly on the process of tumor angiogenesis rather than the invasive behavior a tumor exhibits during migration (Stoletov et al., 2007). Furthermore, tumor cells are commonly loaded with lipophilic dyes, which compared to genetic expression of fluorescent proteins, are not as effective for thoroughly labelling cells and their finer processes for imaging studies (Zeng et al., 2017, Gamble et al., 2018). As it is now widely recognized in the field of GBM biology, gliomas are highly motile tumors known for rapid proliferation and migration. Establishing and characterizing a model of perivascular glioma invasion would open therapeutic possibilities towards a pathological tumor behavior commonly seen in a majority of GBM patients.

Our injection studies revealed the active process of perivascular glioma invasion along the zebrafish CNS vasculature *in vivo*. This biology is exciting for two reasons. First, we demonstrated that important glioma biology could be visualized in zebrafish with only 25-50 cells in 4 days. These experiments utilized significantly fewer cells in a fraction of the time compared to traditional mouse models (Watkins et al., 2014, Griveau et al., 2018). Secondly, a conserved biology must support the attraction of human cells to the developing zebrafish brain vascular network. One of the most critical elements in establishing a model of perivascular glioma invasion is cell number. Other research groups inject widely varied numbers of tumor cells, ranging, from fifty to hundreds of tumor cells. We found that fewer cells injected allowed the tumors to migrate. While glioma cells can shrink up to 30% of their volume to invade the mammalian brain, injecting fewer cells likely provides the glioma cells with more room to not only grow but also invade in a smaller space (Watkins and Sontheimer, 2011). By injecting only 25-50 cells, we observed glioma cells associating with the zebrafish brain vasculature within 24 hours post-implantation. This denotes a salient feature of GBM biology; tumor cell attachment to secondary structures during invasion. We also witnessed brain necrosis in animals injected with >50 cells, as well as animals succoumbing to their tumor burden (data not shown). Most interestingly, perivascular glioma invasion increased over time, from 3-7dpf. After only 4 days of growth, we witnessed a similar percentage of tumor-vessel interactions as previously demonstrated in a GBM mouse model that also utilized

implantation of D54-MG cells (Watkins et al., 2014). Furthermore, we observed tumor-vessel interactions with adherent cell lines and patient-derived xenolines. These experiments highlight the model's accessibility due to what resources may be available to a research group, as PDX lines are time and resource intensive to maintain. Of note, our study is one of the few to utilize pediatric xenoline cells and demonstrate that tumor cell-vessel interactions arise in the context of pediatric CNS cancer *in vivo*.

The extracellular milieu plays a major role during tissue development and for cellular interactions. When assessing the possibility of perivascular glioma invasion in the periphery of the zebrafish, we did not observe glioma-vessel interactions. Little tumor-induced angiogenesis occurred after intracranial injections, possibly due to the presence of vascular endothelial growth factor (VEGF) already in the developing zebrafish brain during active CNS angiogenesis (Liang et al., 2001). Moreover, while it is common for other cancers to metastasize into the brain, it is possible that specific tropism to the BBB and the surrounding microenvironment prevents CNS tumors from hematagenously spreading into the periphery. Glioma cells are known to release VEGF during tumor growth (Tate and Aghi, 2009). Because the intersegmental vessels (ISVs) are already patterned by the time we injected cells, it is possible new vessels formed due to tumor cell secreted VEGF. Researchers have witnessed angiogenesis towards non-CNS tumor cells implanted in the zebrafish, a common process that provides a tumor mass with nutrients necessary to maintain uncontrolled cell growth (Stoletov et al., 2007). Furthermore, in the context of CNS tumors, the BBB vasculature is specialized under a different set of signals than the vasculature within the periphery (Chow and Gu, 2015, Daneman et al., 2009). Interestingly, recent studies injecting non-CNS tumor cells into circulation of the developing zebrafish, demonstrated these cancers attached to ISVs unlike our brain tumor cells (Asokan et al., 2020). Therefore, the process of angiogenesis (the formation of new blood vessels) and perivascular invasion (attraction of cells to pre-existing vessels for cell migration) underscores different biology in GBM. While these studies support the examination of brain tumors within their tissue of origin, the zebrafish could provide an interesting model to dissect brain tumor preferences for the BBB versus peripheral vasculature attachment. Additionally, such experiments would be easily achievable with the zebrafish model system.

As tumor cells secret a myriad of destructive molecules during invasion, a theoretical side effect of perivascular glioma invasion is disruption of the vasculature's integrity (Hoelzinger et al., 2007). GBM models and morphological analysis of patient tissue have demonstrated a change in vessel response and BBB breach via the loss of tight junction proteins (Watkins et al., 2014, Liebner et al., 2000). While the BBB was not a direct focus of this study, we looked at a few potential effects of perivascular glioma invasion on the zebrafish CNS vasculature. Zebrafish possess the chemical and physical barriers necessary to make a functional BBB early in development, also making it easy to survey in our model of perivascular glioma invasion (Umans et al., 2017, van Leeuwen et al., 2018). We found that vessel volume and BBB integrity via expression of *glut1* was not altered in our zebrafish model of perivascular glioma invasion. Our *glut1* data aligns with a study of xenograft models and patient tissue

demonstrating normal expression of GLUT1 at vessels co-opted by tumor cells (Pacioni et al., 2019). While previous studies showed a decrease in tight junction protein expression, it would be an interesting focus to use this zebrafish model to see if changes in BBB proteins occur before or after BBB dysfunction during perivascular tumor invasion.

One of the ultimate goals of establishing a new zebrafish model of perivascular glioma invasion was to demonstrate the feasibility of using small molecules that could target mechanisms necessary for glioma-vessel interactions. The zebrafish is an ideal organism for *in vivo* cancer drug discovery, as animals can be successfully xenografted with tumors in a short time span and are amenable to high-throughput chemical assays (Letrado et al., 2018). In fact, zebrafish "avatars" are becoming successful platforms for drugs to make it into clinical trials (Cully, 2019). In the field of GBM biology, most studies assess overall tumor cell growth and few have investigated means for the chemical disruption of glioma-vessel communication. Previous research has begun to dissect pathways involved in perivascular glioma invasion mouse models, specifically through Wnt and Bradykinin signaling (Griveau et al., 2018, Montana and Sontheimer, 2011). Zebrafish express homologous proteins for both the Wnt and Bradykinin pathways but the function of these homologs in zebrafish cancer models remain unexplored (Wang et al., 2009, Bromée et al., 2005). We demonstrated that tumor-vessels interactions in the zebrafish brain can be disrupted by the Wnt signaling antagonist, XAV939, supporting the potential of BBB-specific cues that mediate perivascular glioma invasion (Daneman et al., 2009). These proof-of-principle small molecule experiments demonstrate the practicality of using this zebrafish model to identify compounds of interest that can target processes important to glioma invasion. Determining the underlying behavior that supports tumor growth would cultivate novel therapies, as migratory cell populations evade the current GBM standard of care. Therefore, this established zebrafish model of perivascular glioma invasion generates an additional platform for dissecting tumor cell biology and progressing cancer drug discovery.

Acknowledgements: The authors would like to acknowledge and thank Miss Joelle Martin for constructive criticism of the manuscript, Mr. Trè Mills for assistance with the volumetric analysis protocol, Dr. Emily Thompson for transducing the D2159MG xenoline for mCherry expression, the Pan laboratory for use of their stereoscope and intellectual discussion, and Dr. Michael Taylor for use of the *Tg(glut1b:mCherry)* line.

Figure legends

Figure 1. Glioma cells retain salient features in the zebrafish brain. (A) A maximum intensity projection of acute slice invasion with mouse brain seeded with 80,000 D54-MG-eGFP tumor cells. After 4 hours of invasion, slices were fixed and stained for the vessel protein Laminin (blue) to visualize the close association of glioma cells (green) during perivascular glioma invasion in this acute *ex vivo* model. Scale bar= 25μm. (B) The transgenic zebrafish vascular reporter line *Tg(fli1a:eGFP)[y1];casper* (green) was intracranially injected with 25-50 D54-MG-tdTomato glioma cells (red) at the midbrain-

hindbrain boundary at 3 days post-fertilization (dpf). 24h post-injection, a maximum intensity projection of an injected animal reveals tumor-vessel associations as seen in the acute slice model. Scale bar= 50µm. (C) An area (dotted white box in B) of a representative Z-plane from the confocal stack showing the corresponding orthogonal planes between the vessel (green) and glioma cell (red) signals. Scale bar 10µm. (D,E) Representative maximum intensity projection images of tumor cells (D, red) and the 6dpf zebrafish brain (E, brightfield) when 25-50 cells are injected. (F,G) Representative images of tumor cells (F, red) and the 6dpf zebrafish brain (G, brightfield) when >50 tumor cells are implanted. Note brain necrosis (G, red dotted area) develops when a large bolus of tumor cells populate the zebrafish brain parenchyma.

Figure 2. Glioma cells actively associate with the developing zebrafish brain over time. (A) A cartoon schematic of intracranial (IC) injection and subsequent *in vivo* imaging of our zebrafish perivascular glioma model. *Tg(fli1a:eGFP)^{y1};casper* animals were injected with 25-50 glioma cells in the midbrain-hindbrain boundary at 3dpf. After recovering, animals were imaged every day until 7dpf to montior tumor burden over time. (B,C) Representative maximum intensity projection confocal images of tumor cells implanted at 3dpf (B) and subsequent migration by 7dpf (C). Scale bar= 50µm.(D,E) Quantification of tumor-vessel interactions in the developing brain. (D) Quantification of D54-MG-tdTomato cells attached to blood vessels over time (**p<0.001,*p=<0.05). (E) Quantification of D54-MG-tdTomato cells at vessel branch points over time (****p=<0.0001,***p=<0.001,**p=<0.01). *n*= 18-21 animals per time point. A one-way ANOVA with Tukey's Multiple comparisons test was performed. Error bars represent mean with standard error of the mean.

Figure 3. Glioma-vascular interactions occur in the zebrafish brain with various GBM lines. An adult D54-MG-tdTomato cell line (A, red) and D2159MG pediatric xenoline (B, red) were intracranially implanted in 3dpf *Tg(fli1a:eGFP)^{y1}* larvae. Live confocal imaging of animals at 72 hours post-injection, reveals GBM growth and development in the larval zebrafish brain. Scale bar= 50µm.

Figure 4. Glioma cells do not migrate along pre-existing peripheral vessels but do evoke non-CNS angiogenesis. (A) Representative maximum intensity projection image of a 3dpf *Tg(fli1a:eGFP)^{y1};casper* animals (green) injected with 25-50 D54-MGtdTomato cells (red) into the trunk of the muscle (white asterisk) close to the dorsal fin (DF) to survey whether GBM would interact with non-CNS vasculature. (B) The same animal was imaged at 7dpf to survey tumor cell survival and movement from the point of injection (white asterisk). (C) Rostral from the point of injection in the same animal from (B), tumor cells moved down the trunk towards the brain and appear near the swim bladder (SB). *n*= 9 animals. (D) A representative image of trunk intersegmental vessel patterning in a 7dpf un-injected *Tg(fli1a:eGFP)^{y1};casper* sibling. (E) We noted instances where tumor cells (red) reaching outward also ensheathed new blood vessels (E,F arrows) that were created after tumor implantation, not seen in un-injected sibling controls. Scale bar=50µm.

Figure 5. Vessel volume and *glut1* expression is not altered after acute GBM invasion. (A,B) Representative 3D rendered volume views of a blood vessel from the contralateral brain (A) in a region similar to a vessel co-opted by a D54-MG-eGFP tumor cell in a 7dpf zebrafish brain (B). (C,D) The volume and signal intensity of *glut1* in tumor-associated and non-tumor associated vessels were measured in 3D rendered volume views. (C) Quantification of vessel volume. A two-tailed, paired student's t-test was performed. (D) Quantification of *glut1:mCherry* fluorescence. A two-tailed, paired student's t-test was performed. Error bars represent mean with standard error of the mean. *n*=8 animals total, 1-3 vessels per animal per group. Scale bar= 10µm.

Figure 6. Pharmacological inhibition of Wnt signaling diminishes tumor-vessel interactions. (A,B) Representative maximum intensity projections of tumor burden (red) in vehicle control (A) or 30µM XAV939 treated (B) animals. Animals were injected with D54-MG-tdTomato cells at 3dpf, unbiasedly sorted, and treated between 4-6dpf. (C) Quantification of tumor-vessel interactions after 48h of vehicle control (DMSO) or 30µM XAV939 treatment. A two-tailed, unpaired t-test was performed (p=<0.05). *n*= 6 animals per group. Error bars represent mean with standard error of the mean. Scale bar= 50µm.

Supplemental Figure 1. Tumor-vessel interactions are maintained in non-specific pathogen free zebrafish lines. (A) A maximum intensity project of a 7dpf *Tg(glut1b:mCherry)* larvae (red) implanted with D54-MG-eGFP tumor cells (green). (B) An area (dotted white box in A) of a representative Z-plane from the confocal stack showing the corresponding orthogonal planes, between the vessel (red) and glioma cell (green) signals. *n*=18 animals. Scale bar= 10µm.

Supplemental Figure 2. Labeling glioma cells with lipophilic dyes is not ideal for visualizing perivascular glioma invasion. (A) A maximum intensity projection of the whole brain of a 4dpf *Tg(fli1a:eGFP)y1;casper* (green) larvae with GBM22 cells (red) loaded with a lipophilic dye 24 hours post-injection. (B) A maximum intensity projection image of the red channel displaying the punctate staining of the human glioma cells. Scale bar= 100µm.

Supplemental Movie 1. Human glioma cells expand within in the periphery of the developing zebrafish larvae. A 3D rendered volume view movie of a 10dpf *Tg(fli1a:eGFP)^{y1};casper* zebrafish, 1 week post tumor implantation. Note that the tumor cells (red) do not interact with any pre-existing vasculature (green) and have migrated towards the rostral end of the animal, from the dorsal fin area past the swim bladder.

References

ALLEN, M., BJERKE, M., EDLUND, H., NELANDER, S. & WESTERMARK, B. 2016. Origin of the U87MG glioma cell line: Good news and bad news. *Science Translational Medicine,* 8**,** 354re3.

ASOKAN, N., DAETWYLER, S., BERNAS, S. N., SCHMIED, C., VOGLER, S., LAMBERT, K., WOBUS, M., WERMKE, M., KEMPERMANN, G., HUISKEN, J., BRAND, M. & BORNHÄUSER, M. 2020. Long-term in vivo imaging reveals tumor-specific dissemination and captures host tumor interaction in zebrafish xenografts. *Scientific Reports,* 10**,** 13254.

BROMÉE, T., KUKKONEN, J. P., ANDERSSON, P., CONLON, J. M. & LARHAMMAR, D. 2005. Pharmacological characterization of ligand-receptor interactions at the zebrafish bradykinin receptor. *British journal of pharmacology,* 144**,** 11-16.

CHOW, B. W. & GU, C. 2015. The molecular constituents of the blood-brain barrier. *Trends in neurosciences,* 38**,** 598-608.

CUDDAPAH, V. A., ROBEL, S., WATKINS, S. & SONTHEIMER, H. 2014. A neurocentric perspective on glioma invasion. *Nature reviews. Neuroscience,* 15**,** 455-465.

CULLY, M. 2019. Zebrafish earn their drug discovery stripes. *Nat Rev Drug Discov,* 18**,** 811-813.

DANEMAN, R., AGALLIU, D., ZHOU, L., KUHNERT, F., KUO, C. J. & BARRES, B. A. 2009. Wnt/β-catenin signaling is required for CNS, but not non-CNS, angiogenesis. *Proceedings of the National Academy of Sciences,* 106**,** 641-646.

FARIN, A., SUZUKI, S. O., WEIKER, M., GOLDMAN, J. E., BRUCE, J. N. & CANOLL, P. 2006. Transplanted glioma cells migrate and proliferate on host brain vasculature: A dynamic analysis. *Glia,* 53**,** 799-808.

FENG, E., SUI, C., WANG, T. & SUN, G. 2017. Temozolomide with or without Radiotherapy in Patients with Newly Diagnosed Glioblastoma Multiforme: A Meta-Analysis. *Eur Neurol,* 77**,** 201-210.

FRIEDRICH, R. W., GENOUD, C. & WANNER, A. A. 2013. Analyzing the structure and function of neuronal circuits in zebrafish. *Frontiers in neural circuits,* 7**,** 71-71.

GAMBLE, J. T., REED-HARRIS, Y., BARTON, C. L., LA DU, J., TANGUAY, R. & GREENWOOD, J. A. 2018. Quantification of glioblastoma progression in zebrafish xenografts: Adhesion to laminin alpha 5 promotes glioblastoma microtumor formation and inhibits cell invasion. *Biochem Biophys Res Commun,* 506**,** 833-839.

GRIVEAU, A., SEANO, G., SHELTON, S. J., KUPP, R., JAHANGIRI, A., OBERNIER, K., KRISHNAN, S., LINDBERG, O. R., YUEN, T. J., TIEN, A.-C., SABO, J. K., WANG, N., CHEN, I., KLOEPPER, J., LARROUQUERE, L., GHOSH, M., TIROSH, I., HUILLARD, E., ALVAREZ-BUYLLA, A., OLDHAM, M. C., PERSSON, A. I., WEISS, W. A., BATCHELOR, T. T., STEMMER-RACHAMIMOV, A., SUVÀ, M. L., PHILLIPS, J. J., AGHI, M. K., MEHTA, S., JAIN, R. K. & ROWITCH, D. H. 2018. A Glial Signature and Wnt7 Signaling Regulate Glioma-Vascular Interactions and Tumor Microenvironment. *Cancer cell,* 33**,** 874-889.e7.

HASON, M. & BARTUNEK, P. 2019. Zebrafish Models of Cancer-New Insights on Modeling Human Cancer in a Non-Mammalian Vertebrate. *Genes (Basel),* 10.

HEGI, M. E., DISERENS, A.-C., GORLIA, T., HAMOU, M.-F., DE TRIBOLET, N., WELLER, M., KROS, J. M., HAINFELLNER, J. A., MASON, W., MARIANI, L., BROMBERG, J. E. C., HAU, P., MIRIMANOFF, R. O., CAIRNCROSS, J. G., JANZER, R. C. & STUPP, R. 2005. MGMT Gene Silencing and Benefit from Temozolomide in Glioblastoma. *New England Journal of Medicine,* 352**,** 997-1003.

HERCULANO-HOUZEL, S., MOTA, B. & LENT, R. 2006. Cellular scaling rules for rodent brains. *Proceedings of the National Academy of Sciences of the United States of America,* 103**,** 12138-12143.

HOELZINGER, D. B., DEMUTH, T. & BERENS, M. E. 2007. Autocrine Factors That Sustain Glioma Invasion and Paracrine Biology in the Brain Microenvironment. *JNCI: Journal of the National Cancer Institute,* 99**,** 1583-1593.

LETRADO, P., DE MIGUEL, I., LAMBERTO, I., DÍEZ-MARTÍNEZ, R. & OYARZABAL, J. 2018. Zebrafish: Speeding Up the Cancer Drug Discovery Process. *Cancer Research,* 78**,** 6048.

LIANG, D., CHANG, J. R., CHIN, A. J., SMITH, A., KELLY, C., WEINBERG, E. S. & GE, R. 2001. The role of vascular endothelial growth factor (VEGF) in vasculogenesis, angiogenesis, and hematopoiesis in zebrafish development. *Mechanisms of Development,* 108**,** 29-43.

LIEBNER, S., FISCHMANN, A., RASCHER, G., DUFFNER, F., GROTE, E. H., KALBACHER, H. & WOLBURG, H. 2000. Claudin-1 and claudin-5 expression and tight junction morphology are altered in blood vessels of human glioblastoma multiforme. *Acta Neuropathologica,* 100**,** 323-331.

MCNEILL, R. S., STROOBANT, E. E., SMITHBERGER, E., CANOUTAS, D. A., BUTLER, M. K., SHELTON, A. K., PATEL, S. D., LIMAS, J. C., SKINNER, K. R., BASH, R. E., SCHMID, R. S. & MILLER, C. R. 2018. PIK3CA missense mutations promote glioblastoma pathogenesis, but do not enhance targeted PI3K inhibition. *PLOS ONE,* 13**,** e0200014.

MELLINGHOFF, I. K., WANG, M. Y., VIVANCO, I., HAAS-KOGAN, D. A., ZHU, S., DIA, E. Q., LU, K. V., YOSHIMOTO, K., HUANG, J. H. Y., CHUTE, D. J., RIGGS, B. L., HORVATH, S., LIAU, L. M., CAVENEE, W. K., RAO, P. N., BEROUKHIM, R., PECK, T. C., LEE, J. C., SELLERS, W. R., STOKOE, D., PRADOS, M., CLOUGHESY, T. F., SAWYERS, C. L. & MISCHEL, P. S. 2005. Molecular Determinants of the Response of Glioblastomas to EGFR Kinase Inhibitors. *New England Journal of Medicine,* 353**,** 2012-2024.

MONTANA, V. & SONTHEIMER, H. 2011. Bradykinin Promotes the Chemotactic Invasion of Primary Brain Tumors. *The Journal of Neuroscience,* 31**,** 4858.

OSTROM, Q. T., GITTLEMAN, H., LIAO, P., VECCHIONE-KOVAL, T., WOLINSKY, Y., KRUCHKO, C. & BARNHOLTZ-SLOAN, J. S. 2017. CBTRUS Statistical Report: Primary brain and other central nervous system tumors diagnosed in the United States in 2010-2014. *Neuro-oncology,* 19**,** v1-v88.

PACIONI, S., D'ALESSANDRIS, Q. G., BUCCARELLI, M., BOE, A., MARTINI, M., LAROCCA, L. M., BOLASCO, G., RICCI-VITIANI, L., FALCHETTI, M. L. &

PALLINI, R. 2019. Brain Invasion along Perivascular Spaces by Glioma Cells: Relationship with Blood-Brain Barrier. *Cancers,* 12**,** 18.

STOLETOV, K., MONTEL, V., LESTER, R. D., GONIAS, S. L. & KLEMKE, R. 2007. High-resolution imaging of the dynamic tumor cell vascular interface in transparent zebrafish. *Proceedings of the National Academy of Sciences of the United States of America,* 104**,** 17406-17411.

STUPP, R., MASON, W. P., VAN DEN BENT, M. J., WELLER, M., FISHER, B., TAPHOORN, M. J., BELANGER, K., BRANDES, A. A., MAROSI, C., BOGDAHN, U., CURSCHMANN, J., JANZER, R. C., LUDWIN, S. K., GORLIA, T., ALLGEIER, A., LACOMBE, D., CAIRNCROSS, J. G., EISENHAUER, E., MIRIMANOFF, R. O., EUROPEAN ORGANISATION FOR, R., TREATMENT OF CANCER BRAIN, T., RADIOTHERAPY, G. & NATIONAL CANCER INSTITUTE OF CANADA CLINICAL TRIALS, G. 2005. Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. *N Engl J Med,* 352**,** 987-96.

TATE, M. C. & AGHI, M. K. 2009. Biology of angiogenesis and invasion in glioma. *Neurotherapeutics : the journal of the American Society for Experimental NeuroTherapeutics,* 6**,** 447-457.

TREDE, N. S., LANGENAU, D. M., TRAVER, D., LOOK, A. T. & ZON, L. I. 2004. The Use of Zebrafish to Understand Immunity. *Immunity,* 20**,** 367-379.

UMANS, R. A., HENSON, H. E., MU, F., PARUPALLI, C., JU, B., PETERS, J. L., LANHAM, K. A., PLAVICKI, J. S. & TAYLOR, M. R. 2017. CNS angiogenesis and barriergenesis occur simultaneously. *Dev Biol,* 425**,** 101-108.

UMANS, R. A. & SONTHEIMER, H. 2018. Combating malignant astrocytes: Strategies mitigating tumor invasion. *Neuroscience research,* 126**,** 22-30.

VAN LEEUWEN, L. M., EVANS, R. J., JIM, K. K., VERBOOM, T., FANG, X., BOJARCZUK, A., MALICKI, J., JOHNSTON, S. A. & VAN DER SAR, A. M. 2018. A transgenic zebrafish model for the in vivo study of the blood and choroid plexus brain barriers using claudin 5. *Biology Open,* 7**,** bio030494.

VARGAS-PATRON, L. A., AGUDELO-DUEÑAS, N., MADRID-WOLFF, J., VENEGAS, J. A., GONZÁLEZ, J. M., FORERO-SHELTON, M. & AKLE, V. 2019. Xenotransplantation of Human glioblastoma in Zebrafish larvae: in vivo imaging and proliferation assessment. *Biology Open,* 8**,** bio043257.

VITTORI, M., MOTALN, H. & TURNŠEK, T. L. 2015. The Study of Glioma by Xenotransplantation in Zebrafish Early Life Stages. *Journal of Histochemistry & Cytochemistry,* 63**,** 749-761.

WANG X, C. J., YOU C, MAO Q 2013. Mutant TP53 enhances the resistance of glioblastoma cells to temozolomide by up-regulating O(6)-methylguanine DNA-methyltransferase. *Neurol Sci,* 34**,** 1421-1428.

WANG, X., LEE, J. E. & DORSKY, R. I. 2009. Identification of Wnt-responsive cells in the zebrafish hypothalamus. *Zebrafish,* 6**,** 49-58.

WATKINS, S., ROBEL, S., KIMBROUGH, I. F., ROBERT, S. M., ELLIS-DAVIES, G. & SONTHEIMER, H. 2014. Disruption of astrocyte–vascular coupling and the blood–brain barrier by invading glioma cells. *Nature Communications,* 5**,** 4196.

WATKINS, S. & SONTHEIMER, H. 2011. Hydrodynamic Cellular Volume Changes Enable Glioma Cell Invasion. *The Journal of Neuroscience,* 31**,** 17250.

WELKER, A. M., JAROS, B. D., PUDUVALLI, V. K., IMITOLA, J., KAUR, B. & BEATTIE, C. E. 2016. Standardized orthotopic xenografts in zebrafish reveal glioma cell-line-specific characteristics and tumor cell heterogeneity. *Disease Models & Mechanisms,* 9**,** 199.

WHITE, R. M., SESSA, A., BURKE, C., BOWMAN, T., LEBLANC, J., CEOL, C., BOURQUE, C., DOVEY, M., GOESSLING, W., BURNS, C. E. & ZON, L. I. 2008. Transparent adult zebrafish as a tool for in vivo transplantation analysis. *Cell stem cell,* 2**,** 183-189.

YANG, X.-J., CUI, W., GU, A., XU, C., YU, S.-C., LI, T.-T., CUI, Y.-H., ZHANG, X. & BIAN, X.-W. 2013. A Novel Zebrafish Xenotransplantation Model for Study of Glioma Stem Cell Invasion. *PLOS ONE,* 8**,** e61801.

ZENG, A., YE, T., CAO, D., HUANG, X., YANG, Y., CHEN, X., XIE, Y., YAO, S. & ZHAO, C. 2017. Identify a Blood-Brain Barrier Penetrating Drug-TNB using Zebrafish Orthotopic Glioblastoma Xenograft Model. *Scientific Reports,* 7**,** 14372.

ZHAO, C., WANG, X., ZHAO, Y., LI, Z., LIN, S., WEI, Y. & YANG, H. 2011. A novel xenograft model in zebrafish for high-resolution investigating dynamics of neovascularization in tumors. *PloS one,* 6**,** e21768-e21768.

48h DMSO

48h XAV939