BlueFeather, the singleton that wasn't: Shared gene content analysis supports expansion of *Arthrobacter* phage cluster FE

Stephanie Demo¶, Andrew Kapinos¶, Aaron Bernardino, Kristina Guardino, Blake Hobbs, Kimberly Hoh, Edward Lee, Iphen Vuong, Krisanavane Reddi, Amanda C. Freise, Jordan Moberg Parker*

¶*Denotes equal contributions*

Department of Microbiology, Immunology, and Molecular Genetics, University of California, Los Angeles (UCLA), Los Angeles, CA, USA

Keywords: bacteriophage, *Arthrobacter*, clustering, virology, microbiology, comparative genomics

*Corresponding Author:
Jordan Moberg Parker, jmobergparker @ ucla.edu

Brief Running Title
Arthrobacter phage BlueFeather

Abstract

Bacteriophages (phages) exhibit high genetic diversity, and the mosaic nature of the shared genetic pool makes quantifying phage relatedness a shifting target. Early parameters for clustering of related *Mycobacteria* and *Arthrobacter* phage genomes relied on nucleotide identity thresholds but, more recently, clustering of *Gordonia* and *Microbacterium* phages has been performed according to shared gene content. Singleton phages lack the nucleotide identity and/or shared gene content required for clustering newly sequenced genomes with known phages. Whole genome metrics of novel *Arthrobacter* phage BlueFeather, originally designated a putative singleton, showed low nucleotide identity but high amino acid and gene content similarity with *Arthrobacter* phages originally assigned to Clusters FE and FI. Gene content similarity revealed that BlueFeather shared genes with these phages in excess of the parameter for clustering *Gordonia* and *Microbacterium* phages. Single gene analyses revealed evidence of horizontal gene transfer between BlueFeather and phages in unique clusters that infect a variety of bacterial hosts. Our findings highlight the advantage of using shared gene content to study seemingly genetically isolated phages and have resulted in the reclustering of BlueFeather, a putative singleton, as well as former Cluster FI phages, into a newly expanded Cluster FE.

Introduction

Bacteriophages are ubiquitous biological entities with an estimated 10^{31} phage particles on Earth. Assuming an average length of 200 nm, they would extend 200 million light years if stacked head-to-tail (Wiles 2014). Phages are found in all ecosystems in which bacteria exist and function as drivers of bacterial evolution (Keen

2015). They exhibit horizontal gene transfer (HGT) with each other and with bacteria,

resulting in the diverse and mosaic nature of phage genomes (Pedulla et al. 2003).

Despite their incredible prevalence in the environment, phages remain largely

understudied (Miller-Ensminger et al. 2018).

Previous research on mycobacteriophages concluded that phages may exhibit a

continuum of diversity, wherein genes are constantly being shuffled amongst the phage

population, resulting in shared genes and sequences between different clusters (Pope et

al. 2015). The immense and ever-expanding diversity of phage genomes has historically

been categorized in terms of nucleotide sequence conservation, with a minimum 50%

nucleotide identity and 50% span length to at least one phage in a cluster to warrant

membership (Hatfull et al. 2010; Klyczek et al. 2017). A mass scale study on *Gordonia*

phages also identified a spectrum of genetic diversity, as clusters did not have clear

boundaries (Pope et al. 2017). Numerous phages lacked the requirement of 50%

nucleotide identity but shared many genes, suggesting a relatedness not captured by

nucleotide comparisons alone. This relatedness was confirmed with a gene content

network phylogeny and subsequently, the cluster assignment parameter for *Gordonia*

phages (Pope et al. 2017), and later for *Microbacterium* phages (Jacobs-Sera et al. 2020),

was adjusted to 35% shared gene content with at least one phage in a cluster.

Mycobacteriophages, as well as *Gordonia* and *Microbacterium* phages, exhibited this

spectrum; however, the extent of diversity varies depending on the current known

phage population, which in turn affects how clustering is carried out. *Arthrobacter*

phages were previously found to exchange genes more slowly than *Gordonia* phages,

and the 50% nucleotide clustering parameter was considered sufficient at the time (Pope

et al. 2017). Further studies on *Arthrobacter* phages found these phages to be

genetically isolated with highly variable gene content for phages that can infect a range

of host species. With this great diversity, nucleotide identity was used to separate

Arthrobacter phages into 10 distinct clusters and 2 singletons (Klyczek et al. 2017), and

this parameter has been considered sufficient to categorize the limited number of

Arthrobacter phages until recently.

Singleton phages can serve as the seeds to start new clusters or be extremely

distinct, as they lack the nucleotide identity and/or shared genes required for clustering

with known phages. In this study, the genome of novel *Arthrobacter* phage BlueFeather

was examined for nucleotide and amino acid identity with other known phages.

BlueFeather lacked sufficient nucleotide conservation for clustering according to

nucleotide-based parameters, and was thus designated a putative singleton. Phage

BlueFeather did, however, have notable amino acid conservation and shared gene

content with other *Arthrobacter* phages previously assigned to Clusters FE and FI,

suggesting it may not be as isolated as its putative singleton status implied. The

outcomes of this research on phage BlueFeather provided evidence for the reclustering

of phage BlueFeather, as well as phages formerly assigned to Cluster FI, into a newly

expanded Cluster FE.

Materials and methods

Sample collection and direct isolation

Soil was collected from Los Angeles, CA in a residential area located at 34.05638889° N, 118.445010000° W. Direct isolation of phages was performed by shaking a soil sample and 2X PYCa broth (Yeast Extract 1 g/L, Peptone 15 g/L, 4.5mM $CaCl_2$, Dextrose 0.1%) in conical tubes at 250 RPM at 25°C for 1.5 hours. After incubation, the solution was filtered through a 0.22 µm syringe and spotted onto *Arthrobacter globiformis* B-2979 (*A. globiformis*). Plaque purifications were performed and a high titer lysate was filter-sterilized to be used in subsequent characterization experiments.

Transmission electron microscopy

Transmission electron microscopy (TEM) was performed on BlueFeather lysate. The sample was placed onto a carbon-coated electron microscope grid and stained with 1% uranyl acetate. Phage particles were visualized using the CM120 Instrument (Philips, Amsterdam, Netherlands) and micrographs were captured. Phage head and tail lengths were measured using ImageJ (Schneider, Rasband, and Eliceiri 2012).

Genome sequencing and assembly

Viral DNA was isolated with the Wizard® DNA Clean-Up System (cat # A7280, Promega, WI, USA). Sequencing libraries were constructed with the NEBNext® Ultra™ II DNA Library Prep kit (New England Biolabs, MA, USA), and sequenced by Illumina-MiSeq at the Pittsburgh Bacteriophage Institute to an approximate shotgun coverage of 3538x. Genome assembly and finishing were performed as previously described (Russell 2018).

Gene annotation

Genomes were annotated as described previously (Pope and Jacobs-Sera 2018) using DNA Master (http://cobamide2.bio.pitt.edu/) and PECAAN (https://pecaan.kbrinsgd.org/) for auto-annotation. GLIMMER (Delcher et al. 1999) and GeneMark (Besemer and Borodovsky 2005) were used to predict protein-coding regions along with their start and stop sites. Manual annotation was performed using Phamerator (Cresawn et al. 2011), Starterator (*SEA-PHAGES/Starterator* [2016] 2020), and host-trained and self-trained GeneMark coding potential maps to support or refute auto-annotation predictions (Besemer and Borodovsky 2005). Gene functions were determined using PhagesDB BLAST (https://phagesdb.org/blastp/), NCBI BLAST (Altschul et al. 1990), HHpred (Soding, Biegert, and Lupas 2005) and CDD (Marchler-Bauer et al. 2014). The presence of transmembrane proteins was determined using TMHMM (Krogh et al. 2001) and TOPCONS (Tsirigos et al. 2015). The annotated complete genome was deposited to GenBank under the accession number MT024867.

Gene content comparisons

Phage genomes used in this study are available from phagesdb.org (Russell and Hatfull 2017). Gepard was used to perform sequence analysis to identify regions of homology between nucleotide sequences or amino acid sequences of different phages (Krumsiek, Arnold, and Rattei 2007). Concatenated whole genome nucleotide and whole amino acid sequences were used to create dot plots with word sizes of 15 and 5, respectively.

SplitsTree was used to generate a network phylogeny in order to reveal the genetic distance between *Arthrobacter* phages (Huson 1998). BlueFeather and up to 10

representative phages from each *Arthrobacter* cluster were selected from the

Actino_Draft database (version 366) for comparison.

The gene content calculator on PhagesDB (https://phagesdb.org/genecontent/)

was used to calculate Gene Content Similarity (GCS), the percentage of shared genes

in phams (groups of genes with related sequences), between BlueFeather, Cluster FE,

and former Cluster FI phages (Cresawn et al. 2011). Gene Content Dissimilarity (GCD)

and maximum GCD gap (MaxGCDGap) were calculated using scripts described

previously (Pope et al. 2017). Heatmaps and scatter plots were created using Prism 8.0.0

(GraphPad Software, San Diego, California, USA) and were used for quantitative

analysis and visualization of GCS and GCD values.

PhagesDB Pham View was used to gather information about phages with genes

in the same phams as BlueFeather's (Russell and Hatfull 2017). PECAAN was used to

obtain the nucleotide sequences for each BlueFeather gene

(https://discover.kbrinsgd.org). The BiologicsCorp online GC content calculator was

used for each gene in the genome (https://www.biologicscorp.com/tools/GCContent/).

Results

BlueFeather is a siphovirus with a short genome

Phage BlueFeather was isolated from a soil sample via direct isolation on *A.*

globiformis B-2979 at 25°C and had a bullseye plaque morphology 5 mm in diameter

(Fig 1a). Transmission electron microscopy (TEM) at 67,000X magnification showed an

average phage capsid diameter and tail length of 48 ± 8 nm and 156 ± 53 nm,

respectively (Fig 1b). The long, flexible, non-contractile tail suggested BlueFeather's

classification as a *Siphoviridae* (Yuan and Gao 2017).

BlueFeather's genome had a length of 16,302 bp, 64.30% GC content, and

genome ends with 15 base 3' sticky overhangs (CCACGGTTCCCGTCC). Phages that

infect *Arthrobacter* hosts have genome lengths that range from 15,319 bp (Toulouse) to

70,265 bp (PrincessTrina) (Klyczek et al. 2017). The average *Arthobacter* phage genome

length (as of May 2020) was 46,968 bp with a standard deviation of 20,619 bp and a

median length of 53,859 bp, suggesting that most *Arthrobacter* phages have genomes

notably larger than that of BlueFeather. BlueFeather's genome contained 25 manually

annotated genes; 18 were of known function, 6 were orphams–meaning they have not

been identified in any other known phage–and 1 was a reverse gene (Fig 2). The left

arm of the genome had highly conserved genes amongst siphoviral *Arthrobacter*

phages, such as those encoding terminase, portal protein, head-to-tail adapter, and tail

proteins (Klyczek et al. 2017). Tail tube and sheath genes were absent, confirming the

classification of BlueFeather as a siphovirus. Genes characteristic of the lytic life cycle,

such as lysin A and holin, were identified; however, there were no genes that would

indicate BlueFeather's ability to undergo a lysogenic life cycle, suggesting that

BlueFeather is not a temperate phage (Shi et al. 2012).

Dot plot comparisons revealed synonymous substitutions in

BlueFeather's genome

Phage BlueFeather was originally classified as a singleton on PhagesDB due to

low nucleotide identity with other known phages. Nucleotide and amino acid dot plots

were created to qualitatively compare BlueFeather to the most similar *Arthrobacter*

phages, including those in Cluster FE (Corgi, Idaho, Noely) and former Cluster FI

(Whytu, Yavru), as identified by BLASTn. Due to the limited number of sequenced

Arthrobacter phages, many of the clusters have few members. Of the 28 *Arthrobacter*

clusters on PhagesDB (as of May 2020), 17 clusters have between 2-4 phages

(including the former Cluster FI). As expected, phages originally assigned to the same

cluster had alignments indicating large regions of nucleotide similarity (Grose and

Casjens 2014), while comparison of BlueFeather's genome to phages originally assigned

to Clusters FE and FI revealed no homologous sequences (Fig 3a). Unexpectedly, dot

plot analysis of concatenated amino acid sequences with a word size of 5 revealed

numerous regions of amino acid sequence similarity between these phages (Fig 3b).

This reflects, at present, perhaps one of the clearest examples in which a group of

phages lack nucleotide identity while sharing considerable amino acid identity.

Gene similarity demonstrates a close relationship between

BlueFeather and phages of Cluster FE and former Cluster FI

Gene Content Similarity (GCS) is a key metric in quantifying phage genetic

relationships and is calculated by averaging the number of shared genes between two

phages (Mavrich and Hatfull 2017). GCS was calculated for BlueFeather, Cluster FE

phages, and phages originally assigned to Cluster FI. BlueFeather shared over 35% of

genes with most Cluster FE phages, and over 55% of genes with former Cluster FI

phages. Over 35% of genes were shared in each pairwise comparison performed, with

the exception of phage Idaho (34.18% shared gene content with BlueFeather) (Fig 4a).

Given that BlueFeather was originally determined to be a singleton, it was surprising to

find GCS greater than the recently adopted threshold of 35% for clustering other phage

populations (Jacobs-Sera et al. 2020; Pope et al. 2017). Gene Content Dissimilarity (GCD)

is the opposite of GCS and was used to calculate the maximum GCD gap

(MaxGCDGap), a metric that represents the degree of isolation between a phage and a

selected phage population (Pope et al. 2017). GCD was calculated for BlueFeather and

all *Arthrobacter* phages. There was a MaxGCDGap of 44.45% between BlueFeather

and Whytu, indicating a relatively high degree of separation between BlueFeather and

the rest of the *Arthrobacter* phage population (Fig 4b). *Arthrobacter* phages which

exhibited pairwise GCD values with BlueFeather of less than 1 were found in Clusters

AN, AU, AM, AZ, AV, AL, FE, AO, FH, FF, and former Cluster FI, indicating shared

gene content. GCD was then calculated for BlueFeather and all known phages in the

PhagesDB Actino_draft database (Fig 4c). Similar to the *Arthrobacter* GCD plot, phages

assigned to Cluster FE and former Cluster FI were the least dissimilar to BlueFeather. It

is notable that in this comparison, there were 63 additional phages ranging from 0.959

to 0.975 GCD, meaning BlueFeather shares a low number of genes with many non-

Arthrobacter phages. Non-*Arthrobacter* phages which exhibited pairwise GCD values

with BlueFeather of less than 1 were found in *Microbacterium* phage Cluster EE,

Mycobacterium phage Clusters N, I, P, and the singleton IdentityCrisis, as well as

Gordonia phage Clusters DT, CW and the singleton GMA4.

To compare the relationships between the *Arthrobacter* phage population as

whole and the phages assigned to Cluster FE, former Cluster FI, and BlueFeather, a

SplitsTree network phylogeny of the phams from each *Arthobacter* phage cluster was

generated to examine the genetic distance between the phages. As expected,

BlueFeather was shown to be more genetically similar to phages originally assigned to

Clusters FE and FI than to any other *Arthrobacter* phage clusters (Fig 5). BlueFeather

demonstrated a closer pham similarity to former Cluster FI phages Whytu and Yavru

than to Cluster FE phages Idaho, Noely and Corgi; however, these phages altogether

formed a distinct branch from the rest of the phages sampled and together comprise the

newly expanded Cluster FE.

BlueFeather genome exhibits evidence of horizontal gene

transfer

Given that BlueFeather shares genes with phages infecting distinct hosts, we

investigated its genome for potential evidence of horizontal gene transfer (HGT). A

whole genome heatmap was created using common metrics for evidence of HGT for

each gene in the genome. As of March 2020, 4 genes in BlueFeather were considered

to have the most convincing evidence for HGT based on GC content and prevalence in

phages that infect unique bacterial hosts: genes 2, 15, 19, and 24 (Fig 6).

Typically, viral genes have about the same (Bohlin and Pettersson 2019) or slightly

lower GC content (de Melo et al. 2019) compared to their bacterial hosts, suggesting that

genes with higher GC content may have been horizontally transferred. BlueFeather had

an overall average GC content of 64.30% and *Arthrobacter globiformis* mrc11 was

found to have an overall GC content of 65.9% (Sahoo et al. 2019). BlueFeather gene-

specific average GC contents ranged from 59.30% to 70.30%, and genes with

maximum average GC contents were considered for HGT. This included genes 15 and

24 with GC contents of 70.3% and 70.1%, respectively.

It is increasingly understood that phages infecting different hosts may share

considerable gene content through processes such as HGT (Pope et al. 2017). For each

gene in the BlueFeather genome, we calculated the number of unique isolation hosts for

phages possessing a pham found in BlueFeather. Gene 2 belongs to a pham with

member genes found in phages that infect *Gordonia malaquae* BEN700 and

Arthrobacter sp. ATCC 21022. Gene 15 belongs to a pham with member genes found in

phages that infect *A. globiformis* B-2979, *A. sp.* ATCC 21022, *Mycobacterium*

smegmatis mc²155, *G. malaquae* BEN700, and *Gordonia rubripertincta* NRRL B-16540.

Gene 19 was the only reverse gene in the BlueFeather genome, and this gene was only

found in BlueFeather and in phages infecting *Microbacterium foliorum* NRRL B-24224

SEA and *Microbacterium paraoxydans* NWU1.

Discussion

Our research was focused on the genomic and evolutionary relationships

between the novel *Arthrobacter* phage BlueFeather and other known phages,

particularly those originally assigned to Clusters FE and FI. Previous studies have

shown that new clusters can be formed when novel phages are found to be similar to

former singletons, as demonstrated by the formation of Cluster AS from *Arthrobacter*

singleton Galaxy (Klyczek et al. 2017). BlueFeather, originally designated as a putative

singleton phage, exhibits over 35% GCS with nearly all phages originally in Cluster FE

and over 55% GCS with those formerly assigned to Cluster FI. The conservation of

amino acids, rather than nucleotides, suggests a history of purifying selection via many

synonymous mutations in which deleterious mutations were filtered out (Ngandu et al.

2008). Moreover, this presents some of the most clear evidence to date of highly

conserved amino acid sequences despite the absence of significant nucleotide

conservation amongst related phages.

There is a high degree of synteny between these phages as well. While

BlueFeather's original designation as a singleton would imply low genomic relatedness

to other phages (Pope et al. 2015), gene similarity between BlueFeather and phages

from Cluster FE and former Cluster FI–in excess of 35%–indicates conservation of gene

functions and genome architecture despite extensive divergence of nucleotide identity.

While *Arthrobacter* phages have been clustered according to nucleotide identity in the

past (Klyczek et al. 2017), this study on BlueFeather, Cluster FE, and the former Cluster

FI highlights the importance of continually reevaluating clustering parameters,

particularly when different parameters may result in different cluster assignments.

Moreover, BlueFeather has the smallest genome of all *Arthrobacter* singletons, and it is

possible that clustering parameters may also need to take genome size into account.

Given that GCS reflects the number of genes shared as a proportion of the total number

of genes for each phage, the same number of shared genes would yield higher GCS in

comparisons between smaller genomes.

Gene content dissimilarity demonstrated that BlueFeather has a MaxGCDGap of

44.45% with phage Whytu, which was originally assigned to Cluster FI. BlueFeather

was found to be least dissimilar with Cluster FE and former Cluster FI phages; this was

supported by a network phylogeny of representative *Arthrobacter* phages that indicated

great diversity between clusters, but revealed that phage BlueFeather forms a distinct

branch with Cluster FE and former Cluster FI phages. Additionally, many phages were

found to share between 0-10% GCS with BlueFeather. While this is too low to warrant a

significant phylogenetic relationship, it reinforced the observed continuum of diversity in

phage populations. Previous research found *Arthrobacter* phage clusters to be very

discrete (Klyczek et al. 2017). Even so, this low yet seemingly widespread display of

shared genes, as well as BlueFeather's unexpected relationships with Cluster FE and

former Cluster FI phages, provides new insight into the genetic landscape of

Arthrobacter phages. Few phages were previously assigned to Cluster FE and the

former Cluster FI, representing only 5 of the 306 sequenced and manually annotated

Arthrobacter phages (as of May 2020). On the other hand, there are 1,906 sequenced

Mycobacterium phages (as of May 2020), which has allowed for a more thorough

investigation of the mycobacteriophage continuum of diversity. As more *Arthrobacter*

phages are sequenced, we expect to observe similar trends in these host-dependent

genetic landscapes.

Unlike singleton phages that are replete with orphams (Klyczek et al. 2017; Pope et

al. 2015), the BlueFeather genome, originally designated as a putative singleton, is

composed predominantly of genes with known functions that have been assigned to

phams. BlueFeather has less than half as many genes as current *Arthrobacter*

singletons and contains highly conserved genes required for viral mechanisms. These

vital functional genes have been more thoroughly studied and as a result, are more

likely to be found in phams with predicted functions (Cresawn et al. 2011). Additionally,

given that pham assignments are performed on the basis of amino acid identity, it is

unsurprising that many of the phams containing these vital functional genes are shared

amongst BlueFeather, Cluster FE, and former Cluster FI phages, despite the lack of

significant nucleotide identity in gene encoding sequences.

Markers of horizontal gene transfer (HGT) included unexpectedly high GC

content, as well as multiple bacterial hosts on which phages sharing genes with

BlueFeather were isolated (Pope et al. 2015). BlueFeather shared phams with a

multitude of non-*Arthrobacter* phages from various clusters, which allowed us to identify

multiple regions as having evidence for HGT. These potential HGT events serve to

magnify phage diversity and promote the phenomenon of genetic mosaicism.

BlueFeather serves as yet another example of the highly intricate mosaic relationships

which exist among phages, a feature of the genetic landscape which makes phage

taxonomy an increasingly difficult task.

In sum, this research has led to the reclustering of BlueFeather and phages

formerly assigned to Cluster FI into a newly expanded Cluster FE. Recent observations

in which there appear to be limited nucleotide conservation but high shared gene

content, as observed in this newly expanded cluster, support the notion that clustering

methods should be continually reevaluated and optimized as more phages are

sequenced (Pope et al. 2017). This study thus provides valuable insight into the

continuum of diversity amongst *Arthrobacter* phages, while also supporting a 35%

shared gene content clustering parameter as was previously adopted for *Gordonia* and

Microbacterium phages (Pope et al. 2017; Jacobs-Sera et al. 2020). Further investigation

into novel phages is essential to understand the complex phage landscape. As more

Arthrobacter phages are discovered, it is likely that we will discover many more phages

like BlueFeather which belong to clusters whose close relationships become apparent

only through the lens of shared gene content.

Acknowledgments

We thank Rebecca A. Garlena and Daniel A. Russell at the Pittsburgh

Bacteriophage Institute for genome sequence and assembly; and Travis Mavrich,

Welkin Pope, Debbie Jacobs-Sera, and Graham Hatfull with the HHMI Science

Education Alliance-Phage Hunters Advancing Genomics and Evolutionary Science

(SEA-PHAGES) program for programmatic support. The authors acknowledge the use

of instruments at the Electron Imaging Center for NanoMachines supported by NIH

(1S10RR23057 to ZHZ) and CNSI at UCLA.

Author contributions

354   A.B., B.H., E.L., I.V., K.G., K.H., and S.D. performed experiments and drafted the

355   paper; S.D., A.K., J.M.P. and A.F. revised the paper; and J.M.P. and A.F. supervised

356   the research. This research was funded in part by the Microbiology, Immunology and

357   Molecular Genetics Department, and the Dean of Life Sciences Division at UCLA.

Author disclosure statement

360   The authors declare that there is no conflict of interest regarding the publication

361   of this article.

Figures

Fig 1. BlueFeather is a Siphovirus. (A) A picked plaque was suspended in phage buffer, then tested in a plaque assay for purification. Plaque morphology was consistent with bullseye plaques around 5 mm in diameter. **(B)** TEM image of BlueFeather at 67,000X magnification. The capsid was estimated to be 48 ± 8 nm and the tail 156 ± 53 nm.

372
373 **Fig 2. BlueFeather genome shares little nucleotide similarity but many phams**
374 **with Cluster FE and former Cluster FI.** The BlueFeather genome is linear with a
375 relatively small length of 16 kbp. Of the 25 identified ORFs, 18 were of known function,
376 6 were orphams and 1 was a reverse gene. BlueFeather had little BLASTn homology to
377 its most similar phages, as indicated by the limited orange and yellow shading.

379
380
381
Fig 3. Dot plots suggest shared amino acids but not nucleotides. Whole genomes
and proteomes for each phage were concatenated and dot plots were created using
Gepard. Original cluster information is denoted along the top of each figure, with phage
BlueFeather indicated by *. **(A)** A whole genome dot plot with word size of 15 indicates
strong intracluster nucleotide similarities with both FE and former FI phages. No

intercluster nucleotide similarities were observed, indicating BlueFeather does not share
significant nucleotide sequences with any of these phages. **(B)** A whole proteome dot
plot with a word size of 5 indicated the same intracluster amino acid similarities seen in
the genome dot plot, but there were also amino acid similarities observed between
BlueFeather, Cluster FE, and former Cluster FI phages. BlueFeather appeared to have
greater amino acid similarity with phages originally assigned to Cluster FI.
393

394
395
396
Fig 4. BlueFeather shares the most phams with phages originally assigned to
Cluster FE and former Cluster FI (A) Gene Content Similarity (GCS) between
BlueFeather, Cluster FE and the former Cluster FI was calculated with the PhagesDB
GCS calculator using the number of shared phams. There was high intracluster GCS,
and BlueFeather showed higher GCS values with former Cluster FI. **(B)** Gene Content
Dissimilarity (GCD) output values of all pairwise comparisons of BlueFeather and all
Arthrobacter phages (305), ordered by magnitude. Cluster FE and former Cluster FI
were found to be least dissimilar to BlueFeather, with a MaxGCDGap of 44.45%,
between BlueFeather and Whytu. **(C)** GCD output values of all pairwise comparisons of
BlueFeather and all phages in PhagesDB (3381). MaxGCDGap remained at 44.45%.
There are no non-*Arthrobacter* phages that are less dissimilar to BlueFeather than
Whytu. BlueFeather shares up to 10% of genes with at least 63 non-*Arthrobacter*
phages.

Fig 5. Expanded Cluster FE includes BlueFeather and former FI phages. A
SplitsTree was generated in order to group *Arthrobacter* phages based on pham
similarity. Ten representative phages from each cluster were selected to measure
evolutionary relatedness. While there is great diversity of *Arthrobacter* phages,
BlueFeather forms a relatively small branch with phages originally assigned to Cluster
FE and the former Cluster FI. These phages, boxed in blue, comprise the expanded FE
Cluster.

Fig 6. Evidence of horizontal gene transfer in the BlueFeather genome. The GC
content for each gene in BlueFeather's genome ranged from 59.30%-70.30%, with an
average of 64.30%. The number of unique isolation hosts that were represented in each
pham ranged from 1-4. Genes with unexpectedly high values were considered to be the
result of horizontal gene transfer. There were four genes with the most convincing
evidence, indicated by *.

428
429
References
431

Altschul, Stephen F., Warren Gish, Webb Miller, Eugene W. Myers, and David J. Lipman. 1990.
"Basic Local Alignment Search Tool." *Journal of Molecular Biology* 215 (3): 403–10.
https://doi.org/10.1016/S0022-2836(05)80360-2.
Besemer, John, and Mark Borodovsky. 2005. "GeneMark: Web Software for Gene Finding in
Prokaryotes, Eukaryotes and Viruses." *Nucleic Acids Research* 33 (Web Server issue):
W451-454. https://doi.org/10.1093/nar/gki487.
Bohlin, Jon, and John H.-O. Pettersson. 2019. "Evolution of Genomic Base Composition: From
Single Cell Microbes to Multicellular Animals." *Computational and Structural*
Biotechnology Journal 17: 362–70. https://doi.org/10.1016/j.csbj.2019.03.001.
Cresawn, Steven G., Matt Bogel, Nathan Day, Deborah Jacobs-Sera, Roger W. Hendrix, and
Graham F. Hatfull. 2011. "Phamerator: A Bioinformatic Tool for Comparative
Bacteriophage Genomics." *BMC Bioinformatics* 12: 395. https://doi.org/10.1186/1471-
2105-12-395.
Delcher, Arthur L., Douglas Harmon, Simon Kasif, Owen White, and Steven L. Salzberg. 1999.
"Improved Microbial Gene Identification with GLIMMER." *Nucleic Acids Research* 27
(23): 4636–41. https://doi.org/10.1093/nar/27.23.4636.
Grose, Julianne H., and Sherwood R. Casjens. 2014. "Understanding the Enormous Diversity of
Bacteriophages: The Tailed Phages That Infect the Bacterial Family
Enterobacteriaceae." *Virology* 468–470 (November): 421–43.
https://doi.org/10.1016/j.virol.2014.08.024.
Hatfull, Graham F., Deborah Jacobs-Sera, Jeffrey G. Lawrence, Welkin H. Pope, Daniel A.
Russell, Ching-Chung Ko, Rebecca J. Weber, et al. 2010. "Comparative Genomic
Analysis of 60 Mycobacteriophage Genomes: Genome Clustering, Gene Acquisition,
and Gene Size." *Journal of Molecular Biology* 397 (1): 119–43.
https://doi.org/10.1016/j.jmb.2010.01.011.
Huson, D. H. 1998. "SplitsTree: Analyzing and Visualizing Evolutionary Data." *Bioinformatics*
(Oxford, England) 14 (1): 68–73. https://doi.org/10.1093/bioinformatics/14.1.68.
Jacobs-Sera, Deborah, Lawrence A. Abad, Richard M. Alvey, Kirk R. Anders, Haley G. Aull,
Suparna S. Bhalla, Lawrence S. Blumer, et al. 2020. "Genomic Diversity of
Bacteriophages Infecting Microbacterium Spp." Edited by Axel Cloeckaert. *PLOS ONE*
15 (6): e0234636. https://doi.org/10.1371/journal.pone.0234636.
Keen, Eric C. 2015. "A Century of Phage Research: Bacteriophages and the Shaping of Modern
Biology: Cause to Reflect." *BioEssays* 37 (1): 6–9.
https://doi.org/10.1002/bies.201400152.
Klyczek, Karen K., J. Alfred Bonilla, Deborah Jacobs-Sera, Tamarah L. Adair, Patricia Afram,
Katherine G. Allen, Megan L. Archambault, et al. 2017. "Tales of Diversity: Genomic and
Morphological Characteristics of Forty-Six Arthrobacter Phages." Edited by Raymond
Schuch. *PLOS ONE* 12 (7): e0180517. https://doi.org/10.1371/journal.pone.0180517.
Krogh, A., B. Larsson, G. von Heijne, and E. L. Sonnhammer. 2001. "Predicting
Transmembrane Protein Topology with a Hidden Markov Model: Application to Complete
Genomes." *Journal of Molecular Biology* 305 (3): 567–80.
https://doi.org/10.1006/jmbi.2000.4315.
Krumsiek, J., R. Arnold, and T. Rattei. 2007. "Gepard: A Rapid and Sensitive Tool for Creating
Dotplots on Genome Scale." *Bioinformatics* 23 (8): 1026–28.
https://doi.org/10.1093/bioinformatics/btm039.

Marchler-Bauer, Aron, Myra K. Derbyshire, Noreen R. Gonzales, Shennan Lu, Farideh Chitsaz,
Lewis Y. Geer, Renata C. Geer, et al. 2014. "CDD: NCBI's Conserved Domain
Database." *Nucleic Acids Research*, gku1221.
Mavrich, Travis N., and Graham F. Hatfull. 2017. "Bacteriophage Evolution Differs by Host,
Lifestyle and Genome." *Nature Microbiology* 2 (9): 17112.
https://doi.org/10.1038/nmicrobiol.2017.112.
Melo, Anna Cristhina Carmine de, Amanda da Mata Gomes, Fernando L. Melo, Daniel M. P.
Ardisson-Araújo, Agueda Palmira Castagna de Vargas, Valessa Lunkes Ely, Elliot W.
Kitajima, Bergmann M. Ribeiro, and José Luiz Caldas Wolff. 2019. "Characterization of a
Bacteriophage with Broad Host Range against Strains of Pseudomonas Aeruginosa
Isolated from Domestic Animals." *BMC Microbiology* 19 (1): 134.
https://doi.org/10.1186/s12866-019-1481-z.
Miller-Ensminger, Taylor, Andrea Garretto, Jonathon Brenner, Krystal Thomas-White, Adriano
Zambom, Alan J. Wolfe, and Catherine Putonti. 2018. "Bacteriophages of the Urinary
Microbiome." *Journal of Bacteriology* 200 (7). https://doi.org/10.1128/JB.00738-17.
Ngandu, Nobubelo K, Konrad Scheffler, Penny Moore, Zenda Woodman, Darren Martin, and
Cathal Seoighe. 2008. "Extensive Purifying Selection Acting on Synonymous Sites in
HIV-1 Group M Sequences." *Virology Journal* 5 (1): 160. https://doi.org/10.1186/1743-
422X-5-160.
Pedulla, Marisa L., Michael E. Ford, Jennifer M. Houtz, Tharun Karthikeyan, Curtis Wadsworth,
John A. Lewis, Debbie Jacobs-Sera, et al. 2003. "Origins of Highly Mosaic
Mycobacteriophage Genomes." *Cell* 113 (2): 171–82. https://doi.org/10.1016/s0092-
8674(03)00233-2.
Pope, Welkin H, Charles A Bowman, Daniel A Russell, Deborah Jacobs-Sera, David J Asai,
Steven G Cresawn, William R Jacobs Jr, et al. 2015. "Whole Genome Comparison of a
Large Collection of Mycobacteriophages Reveals a Continuum of Phage Genetic
Diversity." Edited by Roberto Kolter. *ELife* 4 (April): e06416.
https://doi.org/10.7554/eLife.06416.
Pope, Welkin H., and Deborah Jacobs-Sera. 2018. "Annotation of Bacteriophage Genome
Sequences Using DNA Master: An Overview." In *Bacteriophages*, edited by Martha R.J.
Clokie, Andrew M. Kropinski, and Rob Lavigne, 1681:217–29. Methods in Molecular
Biology. New York, NY: Springer New York. https://doi.org/10.1007/978-1-4939-7343-
9_16.
Pope, Welkin H., Travis N. Mavrich, Rebecca A. Garlena, Carlos A. Guerrero-Bustamante,
Deborah Jacobs-Sera, Matthew T. Montgomery, Daniel A. Russell, Marcie H. Warner,
Science Education Alliance-Phage Hunters Advancing Genomics and Evolutionary
Science (SEA-PHAGES), and Graham F. Hatfull. 2017. "Bacteriophages of Gordonia
Spp. Display a Spectrum of Diversity and Genetic Relationships." *MBio* 8 (4).
https://doi.org/10.1128/mBio.01069-17.
Russell, Daniel A. 2018. "Sequencing, Assembling, and Finishing Complete Bacteriophage
Genomes." In *Bacteriophages: Methods and Protocols, Volume 3*, edited by Martha R.J.
Clokie, Andrew M. Kropinski, and Rob Lavigne, 109–25. Methods in Molecular Biology.
New York, NY: Springer New York. https://doi.org/10.1007/978-1-4939-7343-9_9.
Russell, Daniel A., and Graham F. Hatfull. 2017. "PhagesDB: The Actinobacteriophage
Database." *Bioinformatics (Oxford, England)* 33 (5): 784–86.
https://doi.org/10.1093/bioinformatics/btw711.
Sahoo, Dinabandhu, N. Jusna Devi, N. Ngashangva, P. Momota, Y. Rojeena, and S. Indira
Devi. 2019. "Draft Genome Sequence of Arthrobacter Globiformis Mrc11, an
Antimicrobial Agent Isolated from a Khangkhui Cave Deposit." Edited by Catherine
Putonti. *Microbiology Resource Announcements* 8 (11): MRA.01620-18, e01620-18.
https://doi.org/10.1128/MRA.01620-18.

Schneider, Caroline A, Wayne S Rasband, and Kevin W Eliceiri. 2012. "NIH Image to ImageJ:
25 Years of Image Analysis." *Nature Methods* 9 (7): 671–75.
https://doi.org/10.1038/nmeth.2089.
SEA-PHAGES/Starterator. (2016) 2020. Python. SEA-PHAGES Program.
https://github.com/SEA-PHAGES/starterator.
Shi, Yibo, Ning Li, Yaxian Yan, Hengan Wang, Yan Li, Chengping Lu, and Jianhe Sun. 2012.
"Combined Antibacterial Activity of Phage Lytic Proteins Holin and Lysin from
Streptococcus Suis Bacteriophage SMP." *Current Microbiology* 65 (1): 28–34.
https://doi.org/10.1007/s00284-012-0119-2.
Soding, J., A. Biegert, and A. N. Lupas. 2005. "The HHpred Interactive Server for Protein
Homology Detection and Structure Prediction." *Nucleic Acids Research* 33 (Web
Server): W244–48. https://doi.org/10.1093/nar/gki408.
Tsirigos, Konstantinos D., Christoph Peters, Nanjiang Shu, Lukas Käll, and Arne Elofsson.
2015. "The TOPCONS Web Server for Consensus Prediction of Membrane Protein
Topology and Signal Peptides." *Nucleic Acids Research* 43 (W1): W401-407.
https://doi.org/10.1093/nar/gkv485.
Wiles, Siouxsie. 2014. "Monday Micro – 200 Million Light Years of Viruses?!" Sciblogs. May 8,
2014. https://sciblogs.co.nz/infectious-thoughts/2014/08/05/monday-micro-200-million-
light-years-of-viruses/.
Yuan, Yihui, and Meiying Gao. 2017. "Jumbo Bacteriophages: An Overview." *Frontiers in*
Microbiology 8 (March). https://doi.org/10.3389/fmicb.2017.00403.
549
550

A

B

20 nm