Impacts of Crenarchaeota and Halobacterota on sulfate reduction in the subtropical mangrove ecosystem as revealed by SMDB analysis

Shuming Mo [1, †], Jinhui Li [1, †], Bin Li [2], Ran Yu [1], Shiqing Nie [1], Zufan Zhang [1], Jianping Liao [3], Qiong Jiang [1], Bing Yan [2, *], and Chengjian Jiang [1, 2 *]

[1] State Key Laboratory for Conservation and Utilization of Subtropical Agro-bioresources, Guangxi Research Center for Microbial and Enzyme Engineering Technology, College of Life Science and Technology, Guangxi University, Nanning 530004, China.

[2] Guangxi Key Lab of Mangrove Conservation and Utilization, Guangxi Mangrove Research Center, Guangxi Academy of Sciences, Beihai 536000, China.

[3] School of Computer and Information Engineering, Nanning Normal University, Nanning 530299, China.

† These authors contributed equally to this work.

*: Corresponding Author:

Tel: +86-771-3270736; Fax: +86-771-3237873

Email: jiangcj0520@vip.163.com (CJ); gxybing@tom.com (BY)

Abstract

Sulfate reduction is an important process in the sulfur cycle. However, the relationship between this process and the genotype of microorganisms involved in subtropical mangrove ecosystems is poorly understood. A professional and efficient gene integration database of sulfur metabolism has not been established yet. In this study, a sulfur metabolism gene integrative database (SMDB) had been constructed successfully. The database achieved high coverage, fast retrieval, and low false positives. Then the sulfate reduction by microorganisms in subtropical mangroves ecosystem had been evaluated quickly and accurately with SMDB database. Relative enrichment of sulfate adenylyltransferase indicated that environmental factors select for a partial dissimilatory sulfate reduction process. Furthermore, the sulfate reduction community compensates by producing certain sulfate-reduction genes in response to high-sulfur environments. Taxonomic assignment of dissimilatory sulfate-reduction genes revealed that Crenarchaeota and Halobacterota are completely responsible for this process. Sulfite reductase can help the community cope with the toxic sulfite produced by these Archaea phyla. Collectively, these findings demonstrated that Halobacterota and Crenarchaeota play essential roles in dissimilatory sulfate reduction.

Keywords: sulfate-reduction gene families, mangrove sediment, sulfate-reduction genotype, sulfur metabolism gene integrative database

1 Introduction

Dissimilatory sulfate reduction, which results in the conversion of sulfate (SO_4^{2-}) to sulfide (S^{2-}), is an important reaction in the sulfur cycle (Wenk et al., 2018). The study of dissimilatory sulfate reduction can reveal the occurrence of all dissimilatory sulfate-reduction genes in a community. However, sulfate reduction lacks a complete pathway, a condition that might be a common occurrence (Bukhtiyarova et al., 2019).

The high occurrence of partial sulfate reduction implies that environmental conditions can affect gene occurrence. Gene families, including ATP sulfurylase (*sat*), ATP sulfurylase (APS) reductase (*aprAB*), and dissimilatory sulfite reductase (*dsrABC*), are involved in the canonical dissimilatory sulfate-reduction pathway (Jochum et al., 2018, Anantharaman et al., 2018). Recently, the gene family of dissimilatory sulfite reductase beta subunit (*dsrB)* has been applied to study the diversity of sulfate-reducing bacteria (SRB) (Vavourakis et al., 2019). Dissimilatory sulfate reduction is primarily driven by SRB, and a complete absence of oxygen (O_2) is vital for SRB to gain energy (Wu et al., 2017).

To the best of our knowledge, a professional sulfur database has not been constructed yet. Different databases, such as Clusters of Orthologous Groups (COG) (Galperin et al., 2015), Kyoto Encyclopedia of Genes and Genomes (KEGG) (Kanehisa et al., 2016), Evolutionary Genealogy of Genes: Non-supervised Orthologous Groups (eggNOG) (Huerta-Cepas et al., 2016), SEED subsystems (Overbeek et al., 2005), and M5nr (Wilke et al., 2012), are widely used for functional assignment of predicted genes. However, the application of these databases in analyzing sulfur cycle via shotgun

metagenome sequencing data has several obstacles. First, few of these databases

contain a complete database of gene families involved in sulfur cycle. Second, the use

of these databases obtains a wrong relative abundance of gene families because of

similar functional domains of the genes. For example, certain gene families, such as

polysulfide reductase chain A (*psrA)* and thiosulfate reductase (*phsA)*, share high

sequence similarity (Stoffels et al., 2012). This condition often leads to the mismatch

interpretation of potential ecological processes. Finally, searching for sulfur cycle genes

in large databases is time consuming. Therefore, a specific database of genes related to

the sulfur cycle pathway should be established to address these problems. This novel

database can also be applied to shotgun metagenomics research.

Mangrove sediments are usually characterized as anoxic with high levels of sulfur and

salt and rich in nutrients (Ferreira et al., 2010). Sulfate reduction is one of the most

active processes occurring in mangrove ecosystem (Lin et al., 2019). Dissimilatory

sulfate reduction drives the formation of enormous quantities of reducing sulfide. The

study of sulfate reduction in mangroves is interesting and important, but this endeavor

has several limitations. First, although the diversity of sulfate-reducing bacteria has

been studied in mangrove ecosystems, an understanding of sulfate reduction in these

ecosystems remains insufficient (Wu et al., 2019). Second, researchers have discovered

culturable sulfate reduction of single strains via genomic analysis in other ecological

environments (Spring et al., 2019), but similar studies have been rarely conducted in

mangrove ecosystems. Finally, the relationship between sulfate reduction and the

genotype of microorganisms involved in this process in mangroves are also poorly

understood. Furthermore, the environmental conditions that select for dissimilatory

sulfate-reduction gene families for a more frequent reliance on sulfate reduction remain

unclear.

Therefore, in this study, we created a manually managed database that primarily gathers

most of the publicly available sulfur cycle gene families to address the limitations of

available public databases in sulfur cycle analysis. An integrated database, namely,

sulfur metabolism gene integrative database (SMDB), covering sulfur cycle gene

families was established. SMDB was applied to samples from the subtropical mangrove

ecosystem of Beibu Gulf in China to generate functional profiles. And then the sulfate

reduction in the mangrove ecosystem had been evaluated. This study presents unique

microbial features as a consequence of adapting to distinct environmental conditions to

dissimilatory sulfate-reduction process.

2 Materials and methods

2.1 Sulfur metabolism gene integrative database development

A manually created integrated database was constructed to profile sulfur cycle genes

from shotgun metagenomes, as the described with Tu et al. (2019), with slight

modifications. The database was expected to bear the following characteristics: all

sulfur gene families should be accurate and false positives should be minimized in

database searching. The framework is shown in Supplementary Fig. S1.

2.1.1 Sulfur metabolism gene integrative database sources

The universal protein (UniProt) database (http://www.uniprot.org/) was utilized to

retrieve core database sequences of sulfur cycle gene families by using keywords. Other

databases were used for retrieving nontarget homologous sequences. The other

databases included COG (ftp://ftp.ncbi.nih.gov/pub/COG/COG2014/data), eggNOG

(http://eggnogdb.embl.de/download/eggnog_4.5/), KEGG

(http://www.genome.jp/kegg/), and M5nr

(ftp://ftp.metagenomics.anl.gov/data/M5nr/current/M5nr.gz). In addition, the

homologous sequences from RefSeq non-redundant proteins (NR) database

(ftp://ftp.ncbi.nlm.nih.gov/blast/db/) were also added in the core constructed database.

These databases were selected because they are widely used in metagenome research.

2.1.2 Core database for constructing sulfur cycle genes

The KEGG and MetaCyc databases (Caspi et al., 2016) were referenced to manually

retrieve gene families involved in sulfur cycle. Only those genes that had been

experimentally confirmed to be involved in sulfur metabolism were collected through

an extensive search of the literature. This step was followed by searching the Uniprot

database with their keywords to identify the corresponding annotated sulfur metabolism

gene families. For gene families with vague definitions (e.g., *cysQ* and *MET17*), the

full protein name was added in the keywords to exclude sequences containing vague

annotations. These datasets were retained as the core database for sulfur cycle genes.

2.1.3 Full sulfur metabolism gene integrative database construction

To reduce false positives in database searching and increase the comprehensiveness of

the core database, COG, eggNOG, KEGG, M5nr, and NR databases were used to

retrieve nontarget homologous sequences to construct a full database. The full sulfur

metabolism gene integrative database included the core database of sulfur cycle gene

families and their nontarget homologous gene families. The sequence files of this

integrative database were clustered using the CD-HIT program (version 4.6) (Fu et al.,

2012) at 95% and 100% identity cutoffs. Representative sequences from this integrative

database were then selected to construct SMDB. SMDB database had been deposited

in https://github.com/taylor19891213/sulfur-metabolism-gene-database on January 8th,

2020. The SMDB website (http://smdb.gxu.edu.cn/) had been online since June 22th,

2020.

2.2 Case study

2.2.1 Sampling sites and sediment collection

The ShanKou mangrove sediment in Beihai City, China

(21°29′25.74″N, 109°45′49.43″E) was selected as the sampling location

(Supplementary Fig. S4). Rhizosphere samples (RS) from near-root were designated as

RS1, RS2, and RS3. Nonrhizosphere samples (NRS) from nonrhizosphere were

labelled as NRS1, NRS2, and NRS3. Nonmangrove samples (MS) were assigned as

NMS1, NMS2, and NMS3. Mangrove samples (MS) included groups of RS and NRS.

Several sediment properties, such as total organic carbon (TOC), total organic nitrogen

(TN) content, nutrients (NH_4^+, NO_3^-), and available sulfur (AS), were determined.

Redox potential (ORP) was measured in situ by using a portable ORP meter (BPH-220,

Bell, China). Soil suspension was obtained by centrifugation to determine pH by using

a pH meter (PHS-2C; Sanxin, China) and salinity by using a salinity meter (PAL-06S;

Atago, Japan). TOC, TN, TP, AS, NH_4^+, sulfide, Fe, and NO_3^- were determined

following Chinese standard methods (State Bureau of Technical Supervision, 1994).

2.2.2 DNA extraction and high-throughput sequencing

DNA was extracted from the sediments by using the FastDNA SPIN kit for soil (MP

Biomedicals, USA) according to the manufacturer's instructions. At least 6 μg DNA

from each sample was submitted to the Shanghai Majorbio Bio-Pharm Technology Co.,

Ltd. (Shanghai, China) for sequencing on an Illumina platform. The data output from

each DNA sample was over 10 Gb. The metagenomic sequencing data were deposited

to the NCBI SRA database under the BioProject PRJCA002311.

2.2.3 Shotgun metagenomic sequence processing and analysis

Initial quality assurance/quality control, including trimming sequencing adapters and

bar codes from sequence reads, was performed. Adapter sequences were removed using

SeqPrep v1.33 (https://github.com/jstjohn/SeqPrep). Moreover, sequences less than

100 bp, sequences with quality of less than 20, and reads containing an N base were

removed using Sickle v1.2 (Joshi NA, 2011). Finally, clean reads were created.

For functional annotation, the clean reads were aligned against the online SMDB

database website (http://smdb.gxu.edu.cn/). The blast software of the SMDB database

is DIAMOND v0.9.14 (Buchfink et al., 2015) with parameters set as an e-value cutoff

of 1×10^{-5} and sequence identity cutoff of 60%. The online website of the SMDB

database output was converted to m8 blast format. Then, the reads were assigned as

being from a particular gene if the aligned region was >25 amino acids with an identity

of >60%. Gene families of dissimilatory sulfate reduction were screened out. A

correlation heat map was used to visualize the composition of dissimilatory sulfate

reduction across all nine samples.

2.2.4 16S read extraction and characterization

Reads from the 16S rRNA encoding region of bacterial genomes were extracted using

SortMeRNA (Kopylova et al., 2012) by running the software in filtering mode

(https://github.com/biocore/sortmerna/). The reference database was compiled by Brian

Bushnell   from   Silva   database   (https://www.arb-

silva.de/no_cache/download/archive/release_138/Exports/). The reads extracted using

SortMeRNA were clustered with vsearch (Rognes et al., 2016) with an id = 0.97. The

clusters were then matched using vsearch to the Silva SSUref NR99 dataset to assign

their taxonomy.

3 Results

3.1 Characteristics of sulfur metabolism gene integrative database

A total of 175 gene (sub)families covering 11 sulfur metabolism pathways, including

assimilatory sulfate reduction, thiosulfate disproportionation, sulfide oxidation,

dissimilatory sulfate reduction, sulfite oxidation, sulfur oxidation, sulfur reduction,

tetrathionate oxidation, tetrathionate reduction, thiosulfate oxidation, and organic

degradation/synthesis, were recruited in SMDB (Supplementary Table S1). SMDB

obtained 115,321 and 395,737 representative sequences at 95% and 100% identity

cutoffs, respectively.

To demonstrate the necessity of building a manually created sulfur metabolism gene

database, we first compared the coverage and accuracy of sulfur metabolism gene

(sub)families in SMDB with those of existing public databases. Among the 175 gene (sub)families we recruited in SMDB, only 118, 157, 144, 169, and 172 could be found in these gene (sub)families in COG, eggNOG, KEGG, M5nr, and NR databases, respectively (Fig. 1). The accuracy of SMDB containing sulfur gene (sub)families exceeded that of public databases (Supplementary Fig. S2). We characterized sulfur metabolism gene (sub)families by searching nine shotgun metagenomic datasets obtained by sampling the mangrove environment against SMDB, COG, eggNOG, KEGG, M5nr, and NR databases. Subsequently, the number of sulfur metabolism gene families detected by searching against SMDB far exceeded that of the five other databases (Supplementary Fig. S3).

3.2 Abundance and diversity of sulfur (sub) gene families

A total of 165 different sulfur gene (sub)families were annotated. The sulfur gene (sub)families in each sample ranged from 165 (sample RS3) to 160 (sample RS3) (Supplementary Table S2). The abundance of pathways showed that the organic sulfur degradation/synthesis pathway in these samples was the highest (Supplementary Fig. S6). The pathways of sulfur cycle gene (sub) families showed that dissimilatory sulfate reduction and sulfide oxidation were significantly high in mangrove sediments ($p < 0.05$, Supplementary Fig. S8). However, the pathways of assimilatory sulfate reduction, tetrathionate reduction, tetrathionate oxidation, sulfide oxidation, and sulfite oxidation were significantly high in RS ($p < 0.05$, Supplementary Fig. S8). Hence, we focused on the analysis of dissimilatory sulfate reduction in mangrove ecosystems.

3.3 Bacterial community composition

To study the distribution of dissimilatory sulfate reduction in microbial communities, we used SortMeRNA (Kopylova et al., 2012) to extract 16S rRNA reads. The taxonomic assignments of the reads assigned to 16S rRNA genes indicated that members of Proteobacteria and Desulfobacterota were equally dominant, with compositions ranging from 31% to 39%, across all reads from each sample (Supplementary Fig. S10). Other phyla, such as Crenarchaeota and Chloroflexi, comprised approximately 5% of the total reads. Halobacterota accounted for approximately 1.2% of the total reads (Supplementary Fig. S10).

3.4 Genes for dissimilatory sulfate reduction

The prevalence of dissimilatory sulfate-reduction genes at the sites was initially analyzed by considering the complete dissimilatory sulfate-reduction pathway. Accordingly, reads coming from genes encoding sulfate adenylyl transferase (*sat*), adenylyl sulfate reductase subunit A (*aprA*), and dissimilatory sulfite reductase (*dsrA and dsrB*) were obtained. In the MS and NMS groups, adenylyl sulfate reductase-associated electron transfer protein (*qmoA*) and *dsr* subunit C (*dsrC*) were significantly high in mangrove samples ($p < 0.05$, Fig. 3A). However, sulfite reductase (NADPH), flavoprotein alpha-component (*cysJ*), phosphoadenosine phosphosulfate reductase (*cysH*), sulfite reductase (NADPH) hemoprotein beta-component (*cysI*), sulfite reductase (*sir*), and sulfite reduction-associated complex DsrMKJOP proteins *dsrP* and *dsrO* were significantly high in rhizosphere samples, whereas anaerobic sulfite

reductase (*asrB* and *asrC*) and *dsrB* were significantly low in rhizosphere samples ($p <$

0.05, Fig. 3B).

Reads obtained from *sat* and *aprA* were more frequent than reads from *qmoA*, and *cysI*

(Figs. 2C and 2D). The largest number of reads was assigned to *sat*, accounting for

10%–12% of the total dissimilatory sulfate-reduction reads in all samples. Reads

assigned to *qmoA* did not exceed 1% of the total dissimilatory sulfate-reduction reads

(Supplementary Fig. S7).

3.5 Taxonomic assignment of dissimilatory sulfate-reduction genes

Proteobacteria were the predominant sources of 16S rRNA reads. Therefore, taxonomic

assignments were done at the phylum level to obtain good process of dissimilatory

sulfate-reduction genes families (Table 1). The numbers of dissimilatory sulfate-

reduction genes of Crenarchaeota and Halobacterota exceeded those of other taxonomic

assignments, whereas the remaining genes were assigned to Firmicutes, Proteobacteria,

and Desulfobacterota. This finding was somewhat unexpected because the 16S rRNA

gene analysis showed that these samples were as prevalent as Proteobacteria (Figs. 2A

and 2B). Proteobacteria were the sources of reads assigned to *sat,*

dimethylsulfoniopropionate lyase *(cysQ)*, *dsrA*, and *dsrB* (Table 1). The high

abundance of *aprA* was obtained from Crenarchaeota and Halobacterota (Table 1). *Sat*

and *dsrA* were mainly obtained from Halobacterota. In addition, *asr* and *cys* gene

families were obtained from Crenarchaeota, Halobacterota, and Proteobacteria (Table

1).

3.6 Sediment properties

To determine the environmental factors that likely shape the structure and composition of dissimilatory sulfate-reduction genes in mangrove sediments, we generated a correlation heat map (Supplementary Fig. S9). Certain sediment properties, such as pH, TOC, AS, ORP, NH_4^+, NO_3^-, TN, TP, Fe, salinity, and sulfide content, were determined (Supplementary Fig. S5). The concentrations of AS, Fe, TOC, and TN were significantly higher in the mangrove samples than those in the nonmangrove samples. Moreover, the concentrations of TOC, TN, and TP were significantly higher in the rhizosphere samples than those in the nonrhizosphere samples (Supplementary Tables S5 and S6, $p < 0.05$).

Result showed significant correlations between sediment properties and dissimilatory sulfate-reduction gene (sub) families ($p < 0.01$, Supplementary Fig. S9). pH was negatively correlated with *qmoA*. Sulfide content was negatively correlated with adenylylsulfate reductase subunit B (*aprB)*. Results of Pearson correlation analysis of the pathways are summarized in Supplementary Table S4. Results showed no significant correlation between environmental factors and the dissimilatory sulfate-reduction pathway. AS was negatively correlated with the organic degradation/synthesis pathway (Supplementary Table S4).

4 Discussion

Sulfur cycle is one of the most important biogeochemical cycles in Earth ecosystems. In this study, we developed a manually created database for the fast and accurate profiling of sulfur metabolism genes from shotgun metagenome sequencing data. The

developed SMDB addresses the bottleneck encountered in the previous public

databases. First, SMDB has a precise definition for sulfur metabolism gene families.

Unlike other database generations (Huerta-Cepas et al., 2016), gene families in SMDB

were manually retrieved by using keywords combined with sequence similarity. Typical

examples include the *phsA* and *psrA* genes that encode particle thiosulfate reductase

and polysulfide reductase, respectively, but these genes share high sequence similarity

(Stoffels et al., 2012). Hence, distinguishing the activities of these two enzymes during

genome annotation is difficult. Second, SMDB presents a comprehensive coverage of

gene families in sulfur metabolism. We aimed to provide the latest knowledge and

research progress in sulfur metabolism studies, such as aerobic dimethyl sulfide (DMS)

degradation (Koch and Dahl, 2018). The gene families *TST/MPST/sseA* for K01011

display considerable sequence similarity in KEGG, but they are distinct in two sulfur

transferase subfamilies. Finally, SMDB considers the problem of false positives. Herein,

this issue was addressed by including homologous gene families from multiple public

databases. Most importantly, the small size of SMDB reduces the computational cost

required to output the functional profiling of files of sulfur metabolism gene families.

We applied SMDB in characterizing sulfur metabolism gene (sub)families in mangrove

sediment environment to demonstrate its potential applications in microbial ecology

studies. The metagenomics survey described herein provided strong evidence that in

neutral pH, anoxic, high-sulfur soil, the environment in the Beibu Gulf of China favored

the enrichment of dissimilatory sulfate-reduction genotypes. AS concentrations were

consistently higher in the mangrove zones than in the nonmangrove zones ($p < 0.05$,

Supplementary Table S5), suggesting that the concentrations of available sulfur

determine the abundance of dissimilatory sulfate-reduction reads. Spearman correlation

coefficients of abiotic factors and dissimilatory sulfate-reduction genes were calculated by SPSS (George and Mallery, 2013). We found that sulfide content limited the abundance of *aprB* (Supplementary Table S3), suggesting that sulfate reduction is inhibited by sulfide content when it exceeds a certain threshold. However, our current understanding of dissimilatory sulfate reduction suggests that AS contents are insufficient to explain gene trends, and available O_2 is also an important factor. Mangrove sediments are usually characterized as anoxic (Ferreira et al., 2010). Mangrove conditions favoring dissimilatory sulfate reduction were probably of sufficient duration to contribute to the patterns observed herein, but a more detailed study of sediment properties should be conducted for confirmation.

This pattern supported the hypothesis that environmental conditions act on individual genes in the pathway and favor the occurrence of partial dissimilatory sulfate reduction. In all samples, the reads from *sat* occurred at higher levels than that from dissimilatory sulfate-reduction genes (Figs. 2C and 2D). This finding was unexpected because previous analysis of the occurrence of inorganic sulfur cycle genes in sequence genomes found that *aprA* occurs at higher frequencies than the other gene-encoding proteins required for dissimilatory sulfate-reduction genes (Vavourakis et al., 2019). This condition is probably due to the simultaneous sulfate reduction and sulfate oxidation processes (Watanabe et al., 2013). The protein product of *sat* is responsible for the activation of sulfate to adenosine 5′-phosphosulfate in the first step of dissimilatory sulfate reduction. Owing to the unique ecological environments of subtropical mangrove ecosystems, the class Proteobacteria of Betaproteobacteria is the dominant SRB group in mangrove (Varon-Lopez et al., 2014). The high abundance of

sat and *cysQ* gene families were derived from Proteobacteria possibly because special

environmental conditions in mangrove sediments choose to convert sulfate to APS

(Table 1, Figs. 2A–2D), indicating that the first step of dissimilatory sulfate reduction

is mainly performed by Proteobacteria. In addition, Crenarchaeota and Halobacterota

had relatively complete gene families of dissimilatory sulfate reduction in this study

(Table 1). The reads of *aprA* in mangrove zones were basically obtained from

Crenarchaeota and Halobacterota, suggesting that the function of aprA in mangrove

ecosystems may be mainly performed by these Archaea phyla. Other studies have

shown that Crenarchaeota and Halobacterota were involved in sulfur metabolism

(López-López et al., 2015, Cho et al., 2018). These findings showed that unique

ecological environments selected these microbial communities for dissimilatory sulfate

reduction.

Notably, sulfite is cytotoxic above a certain threshold if not rapidly metabolized and

can wreak havoc at the cellular and whole-plant levels. By providing the gene families

of *asrC, asrA*, *asrB, cysI, cysJ*, and *cysH* that reduces sulfite levels to protect the

members of the local community (Bender et al., 2019). These activities are required for

the biosynthesis of L-cysteine from sulfate (Zeghouf and Chemistry, 2000). If sulfite

reductase is absent, most organisms in the community would be inhibited and grow

slowly. The sediment properties (e.g., OPR, Fe) results in the release of sulfides from

sediments (Shi et al., 2016). Other studies have shown that sulfides in mangrove

sediments are eventually converted to L-cysteine (Lin et al., 2019).

We depicted the process of sulfur metabolism in mangrove sediments. Sulfur

metabolism in mangrove sediments includes biological and abiotic processes (Fig. 4).

For abiotic processes, Fe(III) is buried and acts as an oxidant for sulfide in deeper

sediment layers where it partly binds the produced sulfide as iron sulfide (FeS) and

pyrite (FeS_2) (Wasmund et al., 2017). In this study, the concentration of Fe was

consistently higher in the mangrove zones than that in the nonmangrove zones ($p < 0.05$,

Supplementary Table S5). FeS can be oxidized into various oxidation states of sulfur.

For biological processes, the role of microorganisms includes both oxidation and

reduction of sulfur compounds. Dissimilatory sulfate reduction involves the reduction

of sulfate to sulfide where the electron acceptor is elemental sulfur. By contrast, the

reduced sulfide is assimilated in proteins and amino acids in assimilatory sulfate

reduction (Fuentes-Lara et al., 2019). Sulfide oxidation by O_2, NO_3^-, Fe^{3+}, or Mn^{4+} as

electron acceptors produces sulfur. In addition, sulfur undergoes coupled oxidation and

reduction of sulfur compounds (thiosulfate, sulfite, and sulfur) to sulfate and sulfide.

DMS　　was　　derived　　from　　the　　degradation　　of　　organic　　sulfur　　(e.g.,

dimethylsulfoniopropionate).

The mechanism of dissimilatory sulfate-reduction metabolic pattern in mangrove

ecosystem was constructed (Fig. 5). The figure shows the relative abundances of gene-

encoding enzymes involved in dissimilatory sulfate reduction. The reduced sulfur

compounds were likely oxidized to sulfite by reverse dissimilatory sulfite reductase

(encoded by *rdsrAB*) and thiosulfate sulfur transferase (Fig. 5A). EC 2.7.7.4 (sulfate

adenylyltransferase) had the highest abundance among the enzymes. EC 1.8.99.2

(adenylyl-sulfate reductase) and EC 1.8.99.5 (dissimilatory sulfite reductase) had

higher abundance in mangrove samples than in nonmangrove samples. Hence, the results indicated that the majority of sulfate was converted into APS in dissimilatory sulfate reduction pathway owing to the special environment of the mangrove ecosystem. The model of dissimilatory sulfite reductase also supports the hypothesis that mangrove samples prefer dissimilatory sulfate reduction. In rhizosphere samples, the abundance of EC 1.8.99.5 was low, whereas that of EC 1.8.1.2 (assimilatory sulfite reductase) was high (Fig. 5B). This finding indicated that rhizosphere microorganisms are conducive to the assimilation of sulfate to L-cysteine. Moreover, this result highlighted the importance of mangrove plants in sulfur cycle.

5 Conclusions

In sum, a high-coverage database, namely, SMDB, was successfully constructed. This integrative database contains 175 gene (sub)families and covers 11 sulfur metabolism processes. SMDB was able to quickly and accurately analyze sulfur metabolism. The subtropical mangrove ecosystem favored dissimilation sulfate reduction. Proteobacteria convert sulfate to APS owing to environmental conditions in mangrove sediments. However, Crenarchaeota and Halobacterota were found to be responsible for dissimilatory sulfate reduction. This study provides a theoretical basis for the mechanism of sulfur cycle metabolism in subtropical mangrove wetland ecosystems.

CRediT authorship contribution statement

Conceptualization, C.J. and B.Y.; Methodology, S.M., B.Y. and B.L.; Software, B.Y., S.N., and J.H.; Validation, Z.Z, S.N., R.Y., and J.H.; Formal Analysis, B.Y. and C.J.;

Investigation, B.Y.; Resources, C.J. and B.L.; Data Curation, J.L. and B.Y.; Writing—

Original Draft Preparation, S.M.; Writing—Review and Editing, C.J.; Supervision, Q.J.

and B.Y.; Project Administration, C.J.; Funding Acquisition, C.J. and B.Y.

Declaration of competing interest

The authors declare that they have no known personal relationships or competing

financial interests that could have influenced the work conducted in this study.

Acknowledgements

This research was supported by the Science and Technology Basic Resources

Investigation Program of China (Grant No. 2017FY100704), the Natural Science Fund

for Distinguished Young Scholars of Guangxi Zhuang Autonomous Region of China

(Grant No. 2019GXNSFFA245011), the Natural Science Foundation of Guangxi

Zhuang Autonomous Region of China (Grant No. 2018GXNSFAA050090,

2017GXNSFBA198101).

References

Anantharaman, K. et al. (2018) Expanded diversity of microbial groups that shape the

dissimilatory sulfur cycle. ISME J, 12, 1715-1728. DOI: 10.1038/s41396-018-0078-0

Bender, D. et al. (2019) Impaired mitochondrial maturation of sulfite oxidase in a

patient with severe sulfite oxidase deficiency. Hum Mol Genet, 28, 2885-2899. DOI:

10.1093/hmg/ddz109

Buchfink, B. et al. (2015) Fast and sensitive protein alignment using DIAMOND.

Nat Methods, 12, 59-60. DOI: 10.1038/nmeth.3176

Bukhtiyarova, P.A. et al. (2019) Isolation, characterization, and genome insights into

an anaerobic sulfidogenic Tissierella bacterium from Cu-bearing coins. Anaerobe, 56, 66-77. DOI: 10.1016/j.anaerobe.2019.02.012

Caspi, R., et al. (2016) The MetaCyc database of metabolic pathways and enzymes and the BioCyc collection of pathway/genome databases, Nucleic Acids Res, 44(D1), D471-D480. DOI: 10.1093/nar/gkv1164

State Bureau of Technical Supervision. (1996) Standardization Administration of the People's Republic of China. GB/T 4459.6-1996. (in Chinese).

Cho, E.S. et al. (2018) *Haloplanus rallus sp.* nov., a halophilic archaeon isolated from crude solar salt. Int J Syst Evol Microbiol, 68, 3226-3231. DOI: 10.1099/ijsem.0.002970

Ferreira, T.O. et al. (2010) Spatial patterns of soil attributes and components in a mangrove system in Southeast Brazil (Sao Paulo). J Soils Sediments, 10, 995-1006. DOI: 10.1007/s11368-010-0224-4

Fu, L. et al. (2012) CD-HIT: accelerated for clustering the next-generation sequencing data. Bioinformatics, 28, 3150-3152. DOI: 10.1093/bioinformatics/bts565

Fuentes-Lara, L.O. et al. (2019) From Elemental Sulfur to Hydrogen Sulfide in Agricultural Soils and Plants. Molecules, 24, 2282. DOI: 10.3390/molecules24122282

Galperin, M.Y. et al. (2015) Expanded microbial genome coverage and improved protein family annotation in the COG database. Nucleic Acids Res, 43, D261-269. DOI: 10.1093/nar/gku1223

George, D. and Mallery, P. (2013) IBM SPSS Statistics 21 Step by Step: A Simple Guide and Reference. Pearson, 2013, p 416.

Hausmann, B. et al. (2019) Draft Genome Sequence of *Desulfosporosinus sp.* Strain

Sb-LF, Isolated from an Acidic Peatland in Germany. Microbiol Resour Announc, 8,

e00428-19. DOI: 10.1128/MRA.00428-19

Huerta-Cepas, J. et al. (2016) eggNOG 4.5: a hierarchical orthology framework with

improved functional annotations for eukaryotic, prokaryotic and viral sequences.

Nucleic Acids Res, 44, D286-D293. DOI: 10.1093/nar/gkv1248

Jochum, L.M. et al. (2018) Single-Cell Genomics Reveals a Diverse Metabolic

Potential of Uncultivated Desulfatiglans-Related Deltaproteobacteria Widely

Distributed in Marine Sediment. Front Microbiol, 9, 2038. DOI:

10.3389/fmicb.2018.02038

Jorgensen, B.B. et al. (2019) The Biogeochemical Sulfur Cycle of Marine Sediments.

Front Microbiol, 10, 849. DOI: 10.3389/fmicb.2019.0084

Joshi NA, F.J. (2011) Sickle: A sliding-window, adaptive, quality-based trimming tool

for FastQ files [Software]. Available at https://github.com/najoshi/sickle.

Kanehisa, M. et al. (2016) KEGG as a reference resource for gene and protein

annotation. Nucleic Acids Res, 44, D457-D462. DOI: 10.1093/nar/gkv1070

Koch, T. and Dahl, C. (2018) A novel bacterial sulfur oxidation pathway provides a new

link between the cycles of organic and inorganic sulfur compounds. ISME J, 12, 2479-

2491. DOI: 10.1038/s41396-018-0209-7

Kopylova, E. et al. (2012) SortMeRNA: fast and accurate filtering of ribosomal RNAs

in metatranscriptomic data. Bioinformatics, 28, 3211-3217. DOI:

10.1093/bioinformatics/bts611

Lin, X.L. et al. (2019) Mangrove Sediment Microbiome: Adaptive Microbial

Assemblages and Their Routed Biogeochemical Processes in Yunxiao Mangrove

National Nature Reserve, China. Microb Ecol, 78, 57-69. DOI: 10.1007/s00248-018-

1261-6

López-López, O. et al. (2015) Metagenomics of an Alkaline Hot Spring in Galicia

(Spain): Microbial Diversity Analysis and Screening for Novel Lipolytic Enzymes.

Front Microbiol, 6, 1291. DOI: 10.3389/fmicb.2015.01291

Overbeek, R. et al. (2005) The subsystems approach to genome annotation and its use

in the project to annotate 1000 genomes. Nucleic Acids Research, 33, 5691-5702. DOI:

10.1093/nar/gki866

Rognes, T. et al. (2016) VSEARCH: a versatile open source tool for metagenomics.

PeerJ, 4, e2584. DOI: 10.7717/peerj.2584

Santos, A.A. et al. (2015) A protein trisulfide couples dissimilatory sulfate reduction to

energy conservation. Science, 350, 1541-1545. DOI: 10.1126/science.aad3558

Shi, J.C. et al. (2016) The Variation Characteristic of Sulfides and VOSc in a Source

Water Reservoir and Its Control Using a Water-Lifting Aerator. Int J Environ Res Public

Health, 13, 427. DOI: 10.3390/ijerph13040427

Spring, S. et al. (2019) Sulfate-Reducing Bacteria That Produce Exopolymers Thrive

in the Calcifying Zone of a Hypersaline Cyanobacterial Mat. Front Microbiol, 10, 862.

DOI: 10.3389/fmicb.2019.00862

Stoffels, L. et al. (2012) Thiosulfate Reduction in Salmonella enterica Is Driven by the

Proton Motive Force. J Bacteriol, 194, 475-485. DOI: 10.1128/JB.06014-11

Tu, Q.C. et al. (2019) NCycDB: a curated integrative database for fast and accurate

metagenomic profiling of nitrogen cycling genes. Bioinformatics, 35, 1040-1048. DOI:

10.1093/bioinformatics/bty741

Varon-Lopez, M. et al. (2014) Sulphur-oxidizing and sulphate-reducing communities

in Brazilian mangrove sediments. Environ Microbiol, 16, 845-855. DOI: 10.1111/1462-

2920.12237

Vavourakis, C.D. et al. (2019) Metagenomes and metatranscriptomes shed new light on

the microbial-mediated sulfur cycle in a Siberian soda lake. BMC Biol, 17:69. DOI:

10.1186/s12915-019-0688-7

Wasmund, K. et al. (2017) The life sulfuric: microbial ecology of sulfur cycling in

marine sediments. Environ Microbiol Rep, 9, 323-344. DOI: 10.1111/1758-2229.12538

Watanabe, T. et al. (2013) Diversity of sulfur-cycle prokaryotes in freshwater lake

sediments investigated using *aprA* as the functional marker gene. Syst Appl Microbiol,

36, 436-443. DOI: 10.1016/j.syapm.2013.04.009

Wenk, C.B. et al. (2018) Electron carriers in microbial sulfate reduction inferred from

experimental and environmental sulfur isotope fractionations. ISME J, 12, 495-507.

DOI: 10.1038/ismej.2017.185

Wilke, A. et al. (2012) The M5nr: a novel non-redundant database containing protein

sequences and annotations from multiple sources and associated tools. BMC

Bioinformatics 13, 141. DOI: 10.1186/1471-2105-13-141

Wu, J. et al. (2017) Oxygen Reduction Reaction Affected by Sulfate-Reducing Bacteria:

Different Roles of Bacterial Cells and Metabolites. Indian J Microbiol, 57, 344-350.

DOI: 10.1007/s12088-017-0667-z

Wu, S. et al. (2019) Depth-related change of sulfate-reducing bacteria community in

mangrove sediments: The influence of heavy metal contamination. Mar Pollut Bull,

140, 443-450. DOI: 10.1016/j.marpolbul.2019.01.042.

Chemistry, M. et al. (2000) A Simplifed Functional Version of the *Escherichia coli*

Sulfite Reductase. J Biol Chem, 275, 37651-37656. DOI: 10.1074/jbc.M005619200

Table 1. Taxonomic assignments of dissimilatory sulfate-reduction genes.

Phylum	RS1	RS2	RS3	NRS1	NRS2	NRS3	NMS1	NMS2	NMS3
Crenarchaeota	*aprA;asrA ; dsrK; dsrN*	*aprB; cysJ; cysN; dsrB; nrnA*	*dsrN; nrnA*	*aprA;dsrB; dsrK*	*cysJ; nrnA*	*cysN; cysQ*	*asrC; cysH; cysN; nrnA*	*asrB; asrC; cysD; cysH; cysQ; dsrK; nrnA; sat*	*asrB; cysI; nrnA*
Halobacterota	*cysJ*	*nrnA; asrA; asrC; cysJ; dsrA; dsrM; sat*	*aprA;aprB; cysD;dsrA; dsrB; sat*	*aprA;cysQ; dsrA;dsrB; sat*	*aprA; asrA; cysD; cysH*	*dsrA;dsrM; nrnA*	*asrC; cysD; cysN;cysQ; dsrK; dsrM; dsrO; nrnA; sat*	*asrA; asrB; asrC; cysI; cysN; cysQ; dsrA; dsrB; dsrM; dsrP; nrnA; sat*	*dsrP*
Amorphea	NA	*cysN*	cysD	NA	*cysN; cysNC*	NA	NA	*aprA; dsrA*	NA
Firmicutes	NA	*cysN; dsrB*	NA	NA	*cysN*	NA	NA	*cysH; cysNC*	NA
Chloroflexi	NA	NA	*dsrN*	NA	NA	NA	*dsrA*	NA	NA
Iainarchaeota	NA	NA	NA	*asrA*	NA	NA	NA	NA	NA
Proteobacteria	NA	NA	NA	*cysQ*	*sat*	NA	*asrC*	*dsrA; dsrB; dsrM*	NA
Verrucomicrobiota	NA	NA	NA	NA	*nrnA*	NA	NA	NA	NA
Thermoplasmatota	NA	NA	NA	NA	NA	NA	*sat*	*dsrN*	NA
Actinobacteriota	NA	NA	NA	NA	NA	NA	NA	*asrC; cysD; cysQ; dsrB; nrnA*	NA
Bacteroidota	NA	NA	NA	NA	NA	NA	NA	*aprA; asrA; sir*	NA
Deferrisomatota	NA	NA	NA	NA	NA	NA	NA	*aprA*	NA
Desulfobacterota	NA	NA	NA	NA	NA	NA	NA	*cysQ; sat; sir*	NA
Euryarchaeota	NA	NA	NA	NA	NA	NA	NA	*aprA; dsrK*	NA
Planctomycetota	NA	NA	NA	NA	NA	NA	NA	*aprB*	NA
Spirochaetota	NA	NA	NA	NA	NA	NA	NA	*asrB*	*dsrM*
Zixibacteria	NA	NA	NA	NA	NA	NA	NA	*dsrJ*	NA
Gemmatimonadota	NA	NA	NA	NA	NA	NA	NA	NA	*asrB*

Figure Legends

Figure 1. Sulfur cycle gene (sub)families were detected using SMDB, NR, M5nr, KEGG, eggNOG, COG databases.

Figure 2. Comparison of taxonomic assignments of 16S reads extracted from the metagenomes according to A) mangrove and nonmangrove or B) rhizosphere and nonrhizosphere groups. Comparison of read numbers of genes associated with dissimilatory sulfate reduction grouped into C) mangroves and nonmangrove and D) rhizosphere and nonrhizosphere. Rates are expressed as the relative frequency of reads for each gene multiplied by the average of the total read counts for all samples. Error bars represent 95% confidence interval, and statistically significant groupings are indicated by different letters above the bars ($p < 0.05$).

Figure 3. Significant differences in sulfur (sub) gene families between mangrove and nonmangrove sediments (A) and rhizosphere and nonrhizosphere sediments (B).

Figure 4. Conceptual depiction of sulfur metabolism in mangrove ecosystem, including reactions of inorganic and organic sulfur compounds and transformations of sulfur compounds of intermediate oxidation states (sulfur cycle intermediates, SCI). Inorganic sulfur compounds include oxidation and reduction of sulfur compounds. Black lines depict biological sulfur transformations by microorganisms. Green lines depict abiotic reaction-mediated sulfur transformations to pyrite (FeS_2). Sulfur compounds are depicted within yellow eclipses.

Figure 5. Model for the pathway of dissimilatory sulfate reduction. A) Relative abundances of the gene-encoding enzymes involved in dissimilatory sulfate reduction.

The EC numbers of enzymes are boxed. The relative abundances of enzymes in the two

samples are shown in the nearby color bar, in which the two segments from left to right

represent MS and NMS, respectively. B) Relative abundances of the gene-encoding

enzymes involved in dissimilatory sulfate reduction. The EC numbers of enzymes are

boxed. The relative abundances of genes in the two samples are shown in the nearby

color bar, in which the two segments from left to right represent RS and NRS,

respectively.

Figure 1

Figure 3

Figure 4

Figure 5