Genomic selection for any dairy breeding program via optimized investment in phenotyping and genotyping

Jana Obšteter[1*], Janez Jenko[2], Gregor Gorjanc[3,4]

[1]Department of Animal Science, Agricultural Institute of Slovenia, Ljubljana, Slovenia

[2]Geno Breeding and A.I. Association, Hamar, Norway

[3]The Roslin Institute and Royal (Dick) School of Veterinary Studies, University of Edinburgh,
Edinburgh, United Kingdom

[4]Biotechnical Faculty, University of Ljubljana, Ljubljana, Slovenia

* Correspondence:
Jana Obšteter, Department of Animal Science, Agricultural Institute of Slovenia, Ljubljana, Slovenia
jana.obsteter@kis.si
10

11   **Keywords: genomic selection, dairy breeding program, small populations, optimized**
12   **investment, return on investment**

13
14   Manuscript includes 6982 words, 1 table and 3 figures.

Abstract

This paper evaluates the potential of maximizing genetic gain in dairy cattle breeding by optimizing
investment into phenotyping and genotyping. Conventional breeding focuses on phenotyping
selection candidates or their close relatives to maximize selection accuracy for breeders and quality
assurance for producers. Genomic selection decoupled phenotyping and selection and through this
increased genetic gain per year compared to the conventional selection. Although genomic selection
is established in well-resourced breeding programs, small populations and developing countries still
struggle with the implementation. The main issues include the lack of training animals and lack of
financial resources. To address this, we simulated a case-study of a small dairy population with a
number of scenarios with equal resources yet varied use of resources for phenotyping and
genotyping. The conventional progeny testing scenario had 11 phenotype records per lactation. In
genomic scenarios, we reduced phenotyping to between 10 and 1 phenotype records per lactation and
invested the saved resources into genotyping. We tested these scenarios at different relative prices of
phenotyping to genotyping and with or without an initial training population for genomic selection.
Reallocating a part of phenotyping resources for repeated milk records to genotyping increased
genetic gain compared to the conventional scenario regardless of the amount and relative cost of
phenotyping, and the availability of an initial training population. Genetic gain increased by
increasing genotyping, despite reduced phenotyping. High-genotyping scenarios even saved
resources. Genomic scenarios expectedly increased accuracy for young non-phenotyped male and
female candidates, but also cows. This study shows that breeding programs should optimize
investment into phenotyping and genotyping to maximise return on investment. Our results suggest
that any dairy breeding program using conventional progeny testing with repeated milk records can
implement genomic selection without increasing the level of investment.

1 Introduction

This paper evaluates the potential of maximizing genetic gain in dairy cattle breeding by optimizing investment into phenotyping and genotyping. Breeding programs strive to maximize genetic gain, which is a function of selection intensity, accuracy of selection, genetic variation, and generation interval. The conventional dairy breeding program uses an expensive and time-consuming progeny test. Genomic selection (Meuwissen et al., 2001; Schaeffer, 2006) achieves the same genetic progress faster and cheaper through a substantially reduced generation interval, increased accuracy of selection for young animals, and increased selection intensity of males (Schaeffer, 2006; Obšteter et al., 2019). Despite lower accuracy of sire selection compared to the conventional selection, genomic selection doubles the rate of genetic gain per year in dairy cattle (Wiggans et al., 2017).

All breeding programs operate with a limited resources allocated to breeding activities with the aim to maximize return on investment. Genomic selection is now a de-facto standard in leading breeding programs but is still challenging to implement in small national breeding programs or in developing countries. While leading breeding programs can service some small national breeding, developing countries require tailored breeding goals that will respond to the rapidly rising demand for a sustainable dairy production in local environment (Ducrocq et al., 2018; Marshall et al., 2019; Mrode et al., 2019). We hypothesize that these breeding programs need to evaluate priorities and could optimize the allocation of resources for phenotyping and genotyping to maximize return on investment. We base this hypothesis on the following simple examples (Table S1).

The accuracy of conventional (pedigree-based) estimates of breeding values increases with increasing heritability and increasing number of phenotype records per animal or its closest relatives (e.g. Mrode, 2005). For example, for a female-expressed trait with 0.25 heritability, the accuracy as a function of the number of repeated records per lactation (n) is 0.89 (n=10), 0.81 (n=5), 0.70 (n=2) and 0.62 (n=1). The corresponding accuracies for 100 sires tested on the total of 10,000 daughters are respectively 0.98 (n=10), 0.97 (n=5), 0.96 (n=2) and 0.93 (n=1).

The accuracy of genome-based estimates of breeding values similarly increases with increasing heritability and increasing number of phenotype records per genotyped animal. It also increases with increasing training population, decreasing genetic distance between training and prediction individuals, and decreasing number of effective genome segments (Daetwyler et al., 2008; Clark et al., 2011; Goddard et al., 2011). Following the previous example, assume 10,000 effective genome segments, 0.25 heritability, and a training population of 10,000 cows. The accuracy of genomic prediction for non-phenotyped animals as a function of the number of repeated records per lactation in the training population (n) is 0.76 (n=10), 0.71 (n=5), 0.63 (n=2), or 0.56 (n=1). These examples show diminishing returns with repeated phenotyping and a scope for optimizing return on investment in genomic breeding programs.

The accuracy of genomic prediction as a function of the number of genotyped and phenotyped cows (N) and the number of repeated records per lactation (n) is then 0.84 (N=20,000, n=5), 0.90 (N=50,000, n=2), or 0.93 (N=100,000, n=1). While these genomic prediction accuracies are lower

than with progeny testing, shorter generation interval enables larger genetic gain per unit of time
(Schaeffer, 2006).

Previous studies also explored the value of adding females to the training population (Van Grevenhof
et al., 2012; Gonzalez-Recio et al., 2014). They concluded that accuracy has diminishing returns with
increasing the number of genotyped and phenotyped animals in the training population, hence
additional females are most valuable when training population is small. However, real breeding
programs involve overlapping generations, individuals with a mix of phenotype, pedigree, and
genotype information, various selection intensities, and other dynamic components. Thus, evaluating
the optimal allocation of resources into phenotyping and genotyping is beyond these simple
examples.

The above examples suggest that repeated phenotyping could serve as an internal financial reserve to
enable dairy breeding programs to implement genomic selection and maximize return on investment.
In dairy breeding the most repeatedly and extensively recorded phenotypes are milk production traits.
There are different milk recording methods that differ in the recording frequency (International
Committee for Animal Recording, 2017). The recording interval ranges from daily recording to
recording every nine weeks, which translates to between 310 and 5 records per lactation. The
different recording methods have different costs, which vary considerably between recording
systems, countries, and even their regions. For example, some organizations require payment of a
participation fee plus the cost per sample, while others include the fee in the sample cost, or cover the
costs in other ways. There is also a huge variance in the way dairy breeding programs are funded.
Some are funded by farmers and breeding companies while in some countries data recording or
breeding are subsidized by state or charities. There are also situations where farmers do not record
phenotypes and all genetic progress is generated in "research" nucleus herds.

The aim of this study was to evaluate the potential of maximizing genetic gain by optimizing
investment into phenotyping and genotyping in dairy breeding programs. Since milk production traits
are example of repeated phenotypes with diminishing returns, we aimed to optimize investment into
milk recording and genotyping. To this end we have compared a dairy breeding program with
conventional progeny testing and several genomic breeding programs under equal financial
resources. To implement genomic selection, we reduced the number of milk records per lactation and
invested the saved resources into genotyping. We compared the breeding programs in a case-study of
a small cattle population where implementing genomic selection is challenging. The results show that
reallocating a part of phenotyping resources to genotyping increases genetic gain regardless of the
cost and amount of genotyping, and the availability of an initial training population.

2   Methods

The study aimed to evaluate the effect of different investment into phenotyping and genotyping with
a simulation of a case-study of a small dairy breeding program. The simulation mimicked a real dairy
cattle population of ~30,000 animals analyzed in Obšteter et al. (2019). Here we evaluated 36
genomic scenarios against the conventional scenario, all with equal available resources, but varying
extent of phenotyping and genotyping. The conventional scenario implemented progeny testing and

collected 11 phenotype records per lactation, while genomic scenarios reduced phenotyping and
invested the saved resources into genotyping. The genomic scenarios differed in i) the number of
phenotype records per lactation; ii) the relative cost of phenotyping and genotyping; and iii) the
availability of an initial training population. All scenarios were compared on genetic gain and
accuracy of selection.

120 2.1 Simulation of the base population, phenotype and historical
121   breeding

The simulation mimicked a small dairy cattle breeding program of ~30,000 animals with ~10,500
cows. The introduction of effective genomic selection in such populations is challenging due to costs
of assembling a training population and limited number of training animals. We use this population
as a case-study to optimize investment into phenotyping and genotyping. The dairy breeding program
aimed to improve production traits, which we simulated as a single polygenic trait. We used a
coalescent process to simulate genome comprised of 10 cattle-like chromosomes, each with 10^8 base
pairs, 1,000 randomly chosen causal loci, and 2,000 randomly chosen marker loci. We sampled the
effects of causal loci from a normal distribution and use them to calculate animal's breeding value
(a_i) for dairy performance (y_{ijkl}), which was affected also by a permanent environment (p_i), herd (h_j),
herd-year (hy_{jk}), herd-test-day (htd_{jkl}), and residual environment (e_{ijkl}) effects:
$y_{ijkl} = a_i + p_i + h_j + hy_{jk} + htd_{jkl} + e_{ijkl}$.

We sampled permanent environment effects from a normal distribution with zero mean and variance
equal to the base population additive genetic variance (σ^2_a). We sampled herd, herd-year, and herd-
test-day effects each from a normal distribution with zero mean and variance of 1/3 σ^2_a. Finally, we
sampled residual environment effects from a normal distribution with zero mean and variance of σ^2_a.
This sampling scheme gave a trait with 0.25 heritability and 0.50 repeatability. With the simulated
genome and phenotype architecture we have initiated a dairy cattle breeding program and ran it for
20 years of conventional selection with progeny-testing based on 11 cow phenotype records per
lactation. The detailed parameters of the simulation are described in Obšteter et al. (2019). In
summary, in the breeding program we selected 3,849 out of 4,320 new-born females. We selected
139 bull dams out of cows in the second, third, and fourth lactation. We generated 45 male calves
from matings of bull dams and progeny tested sires (elite matings). Out of these we chose 8 for
progeny testing of which 4 were eventually selected as sires for widespread insemination of cows.
We made all selection decisions based on pedigree-based estimates of breeding values. The 20 years
represented historical breeding and provided a starting point for evaluating future breeding scenarios,
which we ran for additional 20 years.

148 2.2 Scenarios

We evaluated 36 genomic scenarios with varying extent of phenotyping and genotyping against the
conventional scenario, all with equal resources. The conventional scenario continued the breeding
scheme from the historical breeding. It used progeny testing and 11 phenotype recordings per
lactation (named C11), corresponding to the standard ICAR recording interval of 4 weeks
(International Committee for Animal Recording, 2017). We assumed that this scenario represented

the total resources for generating data. We then created genomic scenarios by distributing resources
between phenotyping and genotyping - we reduced phenotyping and invested the saved resources
into genotyping. In genomic scenarios we selected females as in the conventional scenario and males
on genomic estimates of breeding values. The number of genomically tested male candidates varied
according to the genotyping resources in a specific scenario. We selected 5 males with the highest
genomic estimates of breeding value as sires for widespread insemination of cows. We evaluated the
genomic scenarios with a varying number of phenotype records per lactation, relative cost of
phenotyping to genotyping, and the availability of an initial training population.

In genomic scenarios we reduced the number of phenotype records per lactation to between 10 and 1.
Three of them followed ICAR standards of 9, 8, and 5 records per lactation, corresponding to
recording intervals of 5, 6, and 9 weeks. Additionally, we tested the non-standard 10, 2, and 1 records
per lactation. We named the scenarios as "GX" with X being the number of records per lactation.
Genomic scenarios next varied the relative cost of phenotyping ($P) to genotyping ($G). We
compared the cost of 11 phenotype records per lactation to the cost of one genotype. By increasing
the number of records per lactation the cost per record decreased by 6%. The cost of genotyping
remained constant. Based on a survey of several breeding programs, milk recording organizations,
and genotyping providers we have considered three cost ratios of $P:$G: 2:1, 1:1, and 1:2. The
reduction in phenotyping and the relative cost of phenotyping to genotyping dictated the number of
genotyped animals (Table 1). For example, assume that the cost per milk record is €1.78 with 11
records per lactation (in total €19.55). Further assume that the cost per milk record is €1.89 with 10
records per lactation (in total €18.91). This second scenario saves €0.64 that can be used for
genotyping. Applying this to our simulated populations with 10,852 active cows we could genotype
356 animals every year when $P:$G = 1:1.

We invested the saved resources into genotyping females and males in ratio 7:1 based on our
previous work (Obšteter et al., 2019). We genotyped first parity cows. This maximized the accuracy
of genomic prediction, since it reduced genetic distance between training and prediction population,
prevented the loss of investment with culled heifers, and minimized the time to obtain a phenotype
linked to a genotype. To maximize the genetic gain, we genotyped male calves from elite matings
and other high parent average matings. In scenarios where the resources for genotyping females were
larger than the cost of genotyping all first parity cows, we did not reallocate the excess of resources
to genotyping males for consistency (we saved resources in those cases).

Lastly, we created scenarios with and without an initial training population for genomic prediction.
When we assumed an initial training population was available, we genotyped all active cows
(10,852) and progeny tested sires (100) before the first genomic selection of males. When an initial
training population was not available, we yearly genotyped a designated number of first parity cows
until the training population reached 2,000 cows. Once we reached this goal, we started to genotype
both females and males as specified in Table 1. At that point we started genomic selection of males.

2.3 Estimation of breeding values

We selected animals based on their breeding values estimated from a pedigree or single-step genomic
repeatability model with breeding value, permanent environment, and herd-year as random effects.
We did not fit the herd-test-day effect as data structure of this small population did not enable its
accurate estimation. We estimated breeding values once a year with blupf90 (Misztal et al., 2018)
with default settings. In the estimation we included all available phenotype and pedigree records for
all active, phenotyped, or genotyped animals, and additional three generations of their ancestors. We
used at most 25,000 genotype records due to a limit in the academic software version. When we
accumulated more than 25,000 genotyped animals, we removed genotypes of the oldest animals in
favor of the latest genotyped cows and male selection candidates.

201   2.4 Analysis of scenarios

All scenarios had equal resources. We compared the scenarios based on their final genetic gain,
which indicated return on investment, accuracy of selection, and selection intensity. We measured the
genetic gain as an average true breeding value by year of birth and standardized it to have zero mean
and unit standard genetic deviation in the first year of comparison. We measured the accuracy of
breeding values as the correlation between true and estimated breeding values. We measured the
accuracy separately for four groups of animals: i) male candidates (genotyped and non-phenotyped);
ii) sires (currently used in artificial insemination); iii) female candidates (non-genotyped and
non-phenotyped); and iv) cows (all active phenotyped cows and bull dams). We computed the
intensities of two-stage selection (i) by integrating standard bivariate distribution

$$f(x,y,\rho) = \frac{1}{2\pi\sqrt{1-\rho}} e^{\frac{-1}{2(1-\rho)}(x^2+y^2-2\rho\cdot x\cdot y)} \text{ as } i = \frac{\int\limits_{T_x}^{x}\int\limits_{T_y}^{y} f(x,y,\rho)\cdot x\cdot dy\cdot dx}{p_2}, \text{ where } x \text{ is the parent average,}$$

y is the genomic breeding value, and ρ is the correlation between the variables (Young, 1964; Jopson
et al., 2004). We computed the correlation ρ by dividing the scenario-specific accuracy of parent
average by the accuracy of genomic prediction. T_x is the standardized cut-off for the proportion of all
new-born male calves selected for genomic testing (p_1) based on parent average in the first selection
stage. T_y is the standardized cut-off for the proportion of all newborn males selected as sires (p_2)
based on genomic breeding values in the second selection stage. We repeated simulation of the base
population and each scenario 10 times and summarized them with mean and standard deviation
across the replicates. We used Tukey's multiple comparison test to test the significance of the
difference between means.

221   3   Results

Genomic scenarios increased the genetic gain compared to the conventional scenario regardless of
the number of phenotype records per lactation, relative cost of phenotyping to genotyping, and the
availability of an initial training population. Genomic scenarios with an initial training population
increased the genetic gain of the conventional scenario by up to 143%, despite reduced phenotyping.
Genetic gain increased with increasing investment into genotyping. Genomic scenarios increased
accuracy for non-phenotyped male and female candidates, and cows. Scenarios without an initial
training population showed the same trends for genetic gain and accuracy. Although these scenarios

had a slightly smaller genetic gain due to delayed implementation of genomic selection, they still
increased the genetic gain of the conventional scenario by up to 134%. We present these results in
more details in the following sub-sections separately for settings with and without an initial training
population available.

3.1 Genetic gain with an initial training population

Genomic scenarios with an initial training population increased the genetic gain of the conventional
scenario using the same resources. The genetic gain increased with increasing investment in
genotyping, despite reduced phenotyping (Figure 1 and Table S2). We show the corresponding
intensities of sire selection in Table S3. In the $P:$G = 1:1 setting, the genomic scenarios increased
the genetic gain of the conventional scenario between 79% and 143%. By reducing the number of
phenotype records from 11 (C11) to 10 per lactation (G10), we saved resources for genotyping 355
animals per year (45 male candidates). This small change increased male selection intensity from
0.10 to 0.26 and coupled with a shorter generation interval increased the genetic gain by 79% (from
3.01 to 5.41). By reducing the number of phenotype records to nine or eight per lactation (G9 or G8),
we respectively saved resources to genotype 800 or 1,345 animals per year (100 or 165 male
candidates). This respectively increased male selection intensity to 0.39 or 0.49, and genetic gain by
109% or 120% (from 3.01 to 6.30 or 6.62). We achieved the highest genetic gain, between 135% and
143% of the conventional scenario (between 7.07 and 7.33), when we collected between five and one
phenotype records per lactation. In these three scenarios we saved resources for genotyping between
3,230 and 3,850 (all) cows and between 465 and 1,125 male candidates per year, and achieved male
selection intensity between 0.77 and 0.92.

Changing the relative cost of phenotyping to genotyping did not change the trend in genetic gain. In
the G10 scenario of the $P:$G = 1:2 setting we yearly genotyped 182 animals (22 male candidates)
and increased the genetic gain by 80% (from 3.01 to 5.43). In the G10 scenario of the $P:$G = 2:1
setting we yearly genotyped 710 animals (90 males candidates) and increased the genetic gain by
116% (from 3.01 to 6.50). When we maximized the investment into genotyping (G1), we genotyped
between 565 and 2,245 male candidates and all females. This achieved a comparable genetic gain,
between 136% and 143% of the conventional scenario, regardless of the relative cost of phenotyping
to genotyping and male selection intensities.

The high-genotyping scenarios achieved the observed genetic gain without using all the resources
(marked bold in Table S2). In these scenarios the resources designated to genotyping females
exceeded the cost of genotyping all females. These savings could cover additional 11 phenotype
records per lactation for between 169 and 5,950 animals, or between 85 and 11,900 additional
genotypes.

In Figure 1 we also show the growth of the training population for genomic prediction. The training
population started with ~10,000 individuals and grew until reaching 25,000 individuals. The increase
was not linear through all generations, since the procedure for choosing the training animals changed
when the training population exceed 25,000 (only latest females and male candidates included)

3.2 Accuracy with an initial training population

Compared to the conventional scenario, genomic scenarios increased accuracy for young non-phenotyped and genotyped male, non-phenotyped and non-genotyped female candidates, and cows, but decreased accuracy for sires (Figure 2 and Table S4). In the $P:$G=1:1 setting, the accuracy for young genomically tested male candidates ranged between 0.90 and 0.91 regardless of the amount of phenotyping and genotyping. This accuracy was between 0.53 and 0.54 higher compared to the pre-selection for progeny testing and between 0.03 and 0.04 lower compared to the sire selection in the conventional scenario. In contrast, the accuracy for already selected sires decreased with decreasing investment into phenotyping and was between 0.11 and 0.23 lower than in conventional scenario. We observed the lowest accuracy for sires (0.63) when we invested the most into genotyping (G1) and the highest (0.75) when we invested the most into phenotyping (G10).

The accuracy for non-genotyped female candidates and cows increased with increasing genotyping, despite reduced phenotyping. We observed the highest accuracy for female candidates (0.55 - 0.57) and cows (0.77 - 0.79) when we recorded between five and one phenotype record per lactation and invested the rest into genotyping. Compared to the conventional scenario, the genomic scenarios increased the accuracy between 0.03 and 0.11 for female candidates, and between 0.11 and 0.29 for cows.

Changing the relative cost of phenotyping to genotyping affected primarily the accuracy for female candidates and cows. In the majority of scenarios the accuracy increased with decreasing the relative cost of genotyping, which enabled more genotyping. We observed the largest difference of 0.06 for female candidates and 0.12 for cows when we changed the relative cost of phenotyping from half to twice the cost of genotyping. Changing the relative costs, however, did not change the accuracy trends.

3.3 Scenarios without an initial training population

3.3.1 Genetic gain

Genomic scenarios without an initial training population also increased the genetic gain of the conventional scenario using the same resources. The trends were in line with what we observed with an initial training population, that is, increasing genotyping increased genetic gain despite reduced phenotyping (Figure 3). However, all corresponding scenarios achieved between 2% and 28% smaller genetic gain than when an initial training population was available (Table S2).

In the $P:$G setting, genomic scenarios increased the genetic gain of the conventional scenario between 51% and 131%. Compared to when we had an initial training population, the corresponding scenarios achieved between 2% and 16% lower genetic gain. This difference was the largest when we invested the least into genotyping (G10). In this scenario we needed six years to build a training population of 2,000 cows and implement genomic selection, since we only genotyped 355 cows per year. We observed the smallest difference in the scenario that collected two phenotype records per lactations (G2) and implemented genomic selection in the first year.

Changing the relative cost of phenotyping to genotyping did not change the overall trend. In the $P:
$G=1:2 setting, the genomic scenarios increased genetic gain of the conventional scenario between
31% and 126%. That was between 4% and 28% less than corresponding scenarios with an initial
training population. In the $P:$G=2:1 setting, the genomic scenarios increased the genetic gain of the
conventional scenario between 86% and 134%, which was between 3% and 14% less than
corresponding scenarios with an initial training population.

3.3.2 Accuracy

Similar to the scenarios with an initial training population, genomic scenarios without it increased the
accuracy for non-phenotyped male and female candidates, and cows (Figure 3 and Table S4). In the
$P:$G=1:1 setting the accuracy for male candidates ranged between 0.84 and 0.91. In contrast to
scenarios with an initial training population, the accuracy increased with increasing investment into
genotyping. The accuracy for sires ranged between 0.64 and 0.74. Contrary to when we had an initial
training population, we observed no clear trend of either increasing or decreasing accuracy with
decreasing investment into phenotyping. For female candidates the accuracy ranged between 0.47
and 0.56, and for cows between 0.56 and 0.76. For both the accuracies followed the trends of when
we had an initial training population - increasing genotyping increased the accuracy.

Changing the relative cost of phenotyping to genotyping affected the accuracy for non-genotyped
female candidates, cows and male candidates. Decreasing the relative cost of genotyping to
phenotyping increased the accuracy in the majority of the scenarios, particularly the low-genotyping
ones.

4   Discussion

Our results show that any dairy breeding program using conventional progeny testing with repeated
milk records can implement genomic selection without extra costs. While breeding programs have
established funding for phenotyping, not all of them have established funding for genotyping. We
show that by reallocating a part of phenotyping resources into genotyping, breeding programs can
implement genomic selection and substantially increase genetic gain regardless of the amount and
cost of genotyping, and availability of an initial training population. However, increasing investment
in genotyping has diminishing returns, which suggests that breeding programs should optimize the
investment into phenotyping and genotyping to maximize return on investment for selection and
management. The results raise four discussion points: 1) how optimizing the investment in
phenotyping and genotyping affects genetic gain; 2) how optimizing the investment in phenotyping
and genotyping affects accuracy; 3) implications for dairy breeding programs; and 4) limitations of
the study. We first discuss the results under equal cost of phenotyping and genotyping, and an initial
training population available. We then discuss changes at different costs and no initial training
population.

4.1 Genetic gain with an initial training population

4.1.1 Genomic vs. conventional selection

Implementing genomic selection by optimizing the investment in phenotyping and genotyping
increased genetic gain compared to the conventional selection, mainly due to reduced generation
interval in sire selection paths. This improvement is in agreement with previous theoretical studies
(Schaeffer, 2006; Pryce et al., 2010; Obšteter et al., 2019). Empirical studies confirm this; in the US
Holstein population the generation interval for the sires of sires and sires of dams paths recently
decreased between 25% and 50% compared to the conventional selection (García-Ruiz et al., 2016).
Van Grevenhof et al. (2012) also showed that when genomic selection halves the generation interval,
a training population with ~2,000 individuals with own performance or ~3,500 individuals with ten
progeny gives comparable response as conventional selection for a trait with intermediate heritability.

Another major advantage of the genomic scenarios was increased intensity of sire selection. A costly
and lengthy progeny-testing limits the number of tested male candidates in conventional selection.
Genomic selection significantly reduces the cost of testing (Schaeffer, 2006) and thus allows for
testing more male candidates. In the US Holstein population, genomic selection improved the
selection differential for all traits, particularly for traits with low heritability, such as health and
fertility (García-Ruiz et al., 2016).

4.1.2 Increasing the investment into genotyping

Genetic gain increased with increased investment into genotyping. This was mainly due to higher
intensity of sire selection, since more resources for genotyping allowed us to test more male
candidates while selecting the same number. A larger investment into genotyping also increased the
update and total size of the training population, which increased the accuracy of female selection (we
discuss this in the next sub-section).

The genetic gain had diminishing relationship with investment into genotyping. This has important
implications for dairy breeding programs, since they use phenotypes also for management, and we
discuss this separately. The results showed that investing resources of more than six phenotype
records into genotyping did not significantly improve the genetic gain. There are three reasons for
this. First, increasing female training population has diminishing relationship with genetic gain (Van
Grevenhof et al., 2012; Gonzalez-Recio et al., 2014). Since scenarios with an initial training
population started with ~10,000 genotyped and phenotyped cows, enlarging the training population
had a marginal effect. Consequently, the accuracy of sire selection in genomic scenario was high
regardless of the amount of genotyping. Second, increasing investment into genotyping did not
proportionally increase the size of the training population due to the limit of 25,000 animals of the
training population. And third, the intensity of sire selection had diminishing return with increasing
number of genotyped male candidates. This agrees with Reiner-Benaim et al. (Reiner-Benaim et al.,
2017) that showed an increased genetic gain with increasing the number of tested male candidates,
but with a diminishing return. While they achieved the maximum profit with four selected sires out
of 1,721 tested candidates, they achieved 99% or 90% of the maximum profit with respectively 740
or 119 tested candidates. The same three reasons enabled comparable maximum genetic gain
regardless of the relative price of phenotyping to genotyping. In general, selecting less than 2% of the

tested males and updating the training population with at least 35% of first-parity cows resulted in the
maximum genetic gain.

While genetic gain increases with the number of cows in training population, it does not increase
with the number of repeated records. The scenarios with the largest genetic gain therefore had a
training population with many cows and few repeated records (Figure S1). However, since we used
the single-step genomic prediction, the phenotypes of the non-genotyped animals contributed to the
estimation as well. Effectively, all scenarios thus operated with the same number of phenotyped
animals.

We should emphasize, that some of the high-genotyping scenarios achieved the observed genetic
gain at a lower total cost, since they could not use all the saved resources for genotyping females in
the studied population. The saved resources could be invested back into phenotyping females for
milk production or novel traits, genotyping more male candidates, or other breeding actions.

4.2 Accuracy with an initial training population

Despite reduced phenotyping, genomic scenarios increased the accuracy for young non-phenotyped
calves and cows. When the accuracy of parent average is already high, genomic prediction increases
primarily the accuracy of the Mendelian sampling term. But when the accuracy of parent average is
low, such as for the offspring of parents with little or no own or progeny information, genomic
information increases accuracy both for the parent average and the Mendelian sampling term
(Daetwyler et al., 2007; Wolc et al., 2011).

4.2.1 Accuracy for males

For male candidates, genomic prediction more than doubled the accuracy compared to the parent
average used for pre-selection of male calves for progeny testing. This is in agreement with two-fold
accuracy increase in dairy (Schaeffer, 2006) and layers (Wolc et al., 2011). Within the genomic
scenarios, the accuracy for male candidates was high regardless of the amount of genotyping and
phenotyping for two reasons. First, the accuracy of their parent average was high, since we tested
offspring of elite matings. Second, starting with an initial 10,000 training population gave an
adequate accuracy that was additionally boosted by using all available information jointly through the
single-step genomic prediction. Using single-step genomic prediction also removed the bias due to
pre-selection (Jibrila et al., 2020).

In contrast, reducing phenotyping decreased the accuracy of already selected sires. We believe this is
due to two reasons. First, since sires are the very best animals, their breeding values are close
together in the tail of the distribution. Due to small differences between the sires, each additional
phenotypic record helps to differentiate them and thus increases the accuracy. Second, as we invested
more into genotyping, the training population grew quicker and reached the limit of 25,000. At this
point we removed sires' in favor of cows' genotypes, hence prediction for sires depended only on
daughters' data and no longer on their own genotype. However, since this is the accuracy after the
selection has already been made, it is not of great interest for breeding.

4.2.2 Accuracy for females

Genomic scenarios increased the accuracy for cows compared to the conventional scenario. Besides
increasing the accuracy of Mendelian sampling term, using genomic information increases genetic
connectedness between individuals from different management units (Yu et al., 2017; Powell et al.,
2019). Increased connectedness in turn increases the accuracy of prediction regardless of the
heritability and the number of causal loci or markers (Yu et al., 2018). This improvement is
important because we selected bull dams for elite mating from cows.

The accuracy for cows increased with increasing investment into genotyping, despite reduced
phenotyping due to three reasons. First, more cows had both genomic and phenotypic information,
which increased the accuracy of their estimated breeding values. Second, more genotyped cows
increased genetic connectedness (Yu et al., 2018). And third, investing more into genotyping
translated into larger training population and its yearly update. As shown by previous studies (Van
Grevenhof et al., 2012; Gonzalez-Recio et al., 2014), the accuracy of genomic prediction increases
with increasing the size of a female training population. They showed that the accuracy of 0.70 is
achieved with ~20,000 animals as in our study. Same studies shown that, as with genetic gain,
accuracy had a diminishing return with the size of the training population. We observed plateau in
accuracy when we invested more than six phenotype records (out of eleven) into genotyping.

Accuracy for female candidates followed the trend for cows, but at lower values. Female candidates
were not genotyped nor phenotyped, hence their accuracy mainly reflected the accuracy of their
parent average. Increasing genotyping increased the accuracy for cows and in turn increased the
accuracy of female candidate`s parent average. The benefit of this increase was not large, since the
intensity of female selection was low. However, there is potential for this benefit to be larger with
sexed semen and embryo transfer.

4.3 Scenario without an initial training population

4.3.1 Genetic gain

We also considered that some populations do not have access to an initial training population and
have to initialize one themselves. These genomic scenarios still increased genetic gain compared to
the conventional scenario, but achieved lower genetic gain than corresponding scenarios with an
initial training population available. Increasing the investment into genotyping compensated for
starting without a training population in two ways. First, it shortened the time to obtain the targeted
2,000 genotypes required to implement genomic selection down to one year in high-genotyping
scenarios. Second, it shortened the time to build a training population in which an additional record
had negligible effect on accuracy (Gonzalez-Recio et al., 2014).

4.3.2 Accuracy

Accuracy in scenarios without an initial training population closely followed the trends of the
corresponding scenarios with an initial training population available. We observed minor differences
in the low genotyping scenarios that had reduced accuracy for male candidates and sires. We attribute
this to a smaller training population. Buch et al. (2012) showed that for new traits and large-scale

recording, we can achieve 75% of the maximum genomic accuracy within first two to three years of
recording. In our study we shortened this period even more by including the historical data through
the single-step genomic prediction.

4.4 Implications

The results suggest that any dairy breeding program using conventional progeny testing with repeated
milk records can implement genomic selection without extra costs by optimizing the investment of
resources into breeding actions. Here we propose funding the genotyping with a part of resources for
milk recording, since we can manipulate the number of repeated records. Breeding programs could
reduce phenotyping for a different trait that they record repeatedly and is perhaps less crucial for
management. They could also reallocate the funds from another breeding action.

Additionally, we could optimize which individuals to genotype and phenotype, as well as the
computational costs . Selective phenotyping was shown to increase the accuracy of genomic
prediction up to 20% with small sample sizes in plant breeding (Heslot & Feoktistov, 2017; Akdemir
& Isidro-Sánchez, 2019). Similarly, selective genotyping of cows from the distribution tails has been
shown to increase the accuracy of genomic prediction by 15% (Jenko et al., 2017). We expect this
would further increase the return on investment, but increase the complexity of optimization.
Regarding computing costs, the problem of a large number of genotypes can be alternatively solved
by using methods with reduced computational costs. Examples of such methods are the algorithm for
proven and young (Misztal et al., 2014) and singular value decomposition of the genotype matrix
(Ødegård et al., 2018). Also, as shown in our study, we can achieve large genetic gain with a
relatively small training population of recent genotypes.

The economic efficiency of breeding programs strongly depends on which stakeholders fund which
breeding action. Different programs have different investment schemes, often complex. The
scenarios presented in this paper are of little value for programs where funding for phenotyping and
genotyping is disconnected. Similarly, optimizing the investment into phenotyping is not of interest
for breeding programs with abundant use of automated milking systems where the cost of
phenotyping does not depend on the number of records. However, in populations with small herds
the use of automated milking systems is limited. Further on, genomic selection could benefit some
settings more than others. For example, genomic information is especially important for generating
genetic connectedness in systems with small herd sizes, geographically dispersed farms, and limited
use of artificial insemination, often found in low- to mid-income countries (Powell et al., 2019). The
same benefits are expected for small ruminant programs that do not actively exchange sires between
herds (Kasap et al., 2018).

High cost of genotyping diminishes the benefit of the proposed solutions. The relative cost of
phenotyping to genotyping at which genomic selection is not beneficial to conventional selection
depends on a number of factors: i) the number of females in the recorded population, since it dictates
the savings from reducing the phenotyping; ii) the number of phenotype records the breeders are
willing to sacrifice; iii) the availability of an initial training population; and iv) the ratio of genotyped
males and females. In our case, if genotyping was ten-times more expensive than phenotyping with

11 records, we would save resources to genotype between 36 (ten phenotypic records) and 900 (one
phenotypic record) animals. While such numbers of genotyped animals could allow for efficient
genomic selection if we had an initial training population and genotyped only males, it would
probably not be viable if we had to build the training population ourselves.

We did not account for the benefits of genotyping besides genomic selection. Genomic information
has additional value for i) parentage verification, parentage discovery, or correction of parentage
errors; ii) management of monogenic diseases and traits, which can prevent economic losses caused
by spreading lethal alleles or create economic gain by adding value to the products; iii) better
monitoring and control of inbreeding (Sonesson et al., 2012) and optimization of matings (Obšteter et
al., 2019); iv) determination of animals' breed composition, which serves to identify the most
appropriate breed or cross-breed in a production system, or improve the structure of the training
population (Marshall et al., 2019). Additional uses of genotypes increase the return on investment
beyond what we measured in this study.

4.5 Limitations of the study

4.5.1 Reducing the number of phenotype records

Balancing phenotyping and genotyping can lead to conflicts between managing production (short-
term goal) and achieving genetic gain (long-term goal). Producers use phenotype records to manage
animals' health, reproduction and feed composition, which affect the success of dairy cattle
production in the short-term. Besides managing production, milk recording is also important from an
environmental perspective (Verbič et al., 2019), but so is genetic improvement. In general, about half
of phenotypic improvement is due to management and half due to selection (Dekkers & Hospital,
2002).

The ownership of the data that drives dairy improvement is a matter of discussion in many breeding
programs. When data collection is subsidized, the data are usually free for management and genetic
improvement. Finding optimal frequency of recording for management and genetic improvement in
such systems is crucial for optimal return on investment. When data collection is paid by producers,
they can be available to breeding organizations for free or at a cost. Depending on the cost breeding
organizations can save some resources by purchasing smaller number of repeated phenotypes per
individual and genotype more selection candidates instead.

While genotype data is currently largely used for selection, the same data could also support
management (predicting feed requirements, disease liability, etc.). Therefore, evaluating the value of
phenotype and genotype data is complex and beyond the scope of this study. One possible way
forward would be to compare variance between herd-test day effects and genetic variance to contrast
the value of managing production and genetic gain in addition to comparing phenotypic and genetic
trends (Dekkers & Hospital, 2002).

In practice, test day records are also used to compute the 305-day milk yield (International
Committee for Animal Recording, 2020). The longest sampling interval tested in our study and still
approved by ICAR was nine weeks, which yielded five records per lactation. Previous studies

showed that the correlation of predicting 305-day milk using five records per lactation with using
either weekly or eleven records per lactation was between 0.98 and 0.99 (Pool and Meuwissen, 1999;
Berry et al. 2005). However, some studies showed that prediction using less than eleven records can
yield substantial bias (Gantner et al., 2008)

4.5.2 Single additive trait

We simulated milk yield as a single polygenic trait with additive genetic as well as herd, permanent
environment, and residual environmental effects. We did not simulate nor account for non-additive
genetic effects that also affect dairy performance (Fuerst & Sölkner, 1994; Ertl et al., 2014; Jiang et
al., 2017). We note that we simulated permanent environment effects, which capture non-additive
genetic effects and other individual specific environmental effects. We also simulated milk yield in
different lactations as a single trait with constant heritability, whereas genetic correlation between
different lactations and through the lactation is not unity (Meyer, 1984; Swalve & Vleck, 1987; Dong
& van Vleck, 1989). If genetic correlation is less than one, the repeatability of the phenotype
decreases and the value of a repeated record and its contribution to accuracy diminishes.

We simulated a trait with heritability of 0.25, since the majority of production traits recorded
repeatedly show intermediate heritability. Previous studies also provide insights in how changing the
heritability of the phenotype would affect the results. On one hand, at a lower heritability we would
need more females in the training population until the contribution of additional female was
negligible (Gonzalez-Recio et al., 2014). On the other hand, genomic selection is less affected by the
heritability than conventional selection and hence more beneficial for traits with low heritability
(Lillehammer et al., 2011; García-Ruiz et al., 2016).

4.5.3 Genomic selection of females

We did not use genomic selection for females, nor did we use reproductive technologies such as
sexing semen or embryo transfer. This would further decrease the generation interval, increase
selection intensity on female side, and in turn increase genetic gain of genomic scenarios even more
(Pryce et al., 2010; García-Ruiz et al., 2016). Such an implementation of genomic selection requires
only a minor modification of the design used in this study - genotyping heifers instead of first-parity
cows. However, reproductive technologies require a larger modification and investment. Some of the
scenarios saved resource and could invest into these technologies.

5   Author Contributions

561 JO designed the study, ran the simulation, analyzed the data, interpreted the results, and wrote the
562 manuscript. JJ participated in study design, results interpretation, and manuscript revision. GG has
563 initiated and supervised the work and contributed to all its stages.

6   Funding

JO acknowledges support from Agricultural Institute of Slovenia and core financing of Slovenian
Research Agency (grant P4-0133). GG acknowledges support from the BBSRC to The Roslin
Institute (BBS/E/D/30002275) and The University of Edinburgh's Data-Driven Innovation
Chancellor's fellowship.

7   Acknowledgments

The authors acknowledge I. Pocrnić (The Roslin Institute, University of Edinburgh) for his help in
interpreting the results and comments on the manuscript.

8   References

Akdemir, D., and J. Isidro-Sánchez. 2019. Design of training populations for selective phenotyping
in genomic prediction. Sci Rep. 9:1446. https://doi.org/10.1038/s41598-018-38081-6

Berry, D. P., V. E. Olori, A. R. Cromie, R. F. Veerkamp, M. Rath, and P. Dillon. 2005. Accuracy of
predicting milk yield from alternative milk recording schemes. Animal Science, 801:53–60.
https://doi.org/10.1079/ASC34880053

Buch, L. H., M. Kargo, P. Berg, J. Lassen, and A. C. Sørensen. 2012. The value of cows in reference
populations for genomic selection of new functional traits. Animal. 6:880–886.
https://doi.org/10.1017/S1751731111002205

Clark, S. A., J. M. Hickey, and J. H. J. van der Werf. 2011. Different models of genetic variation and
their effect on genomic evaluation. Genet. Sel. Evol. 43:18. https://doi.org/10.1186/1297-9686-43-18

Daetwyler, H. D., B. Villanueva, P. Bijma, and J. A. Woolliams. 2007. Inbreeding in genome-wide
selection. Journal of Animal Breeding and Genetics, 124:369–376.

Daetwyler, H. D., B. Villanueva, and J. A. Woolliams. 2008. Accuracy of Predicting the Genetic
Risk of Disease Using a Genome-Wide Approach. PLoS ONE, 3:e3395.
https://doi.org/10.1371/journal.pone.0003395

Dekkers, J. C. M. and F. Hospital. 2002. The use of molecular genetics in the improvement of
agricultural populations. Nat Rev Genet. 3:22–32. https://doi.org/10.1038/nrg701

Dong, M. C. and L. D. van Vleck. 1989. Correlations Among First and Second Lactation Milk Yield
and Calving Interval. J. Dairy Sci. 72:1933–1936. https://doi.org/10.3168/jds.S0022-0302(89)79313-
9

Ducrocq, V., D. Laloe, M. Swaminathan, X. Rognon, M. Tixier-Boichard, and T. Zerjal. 2018.
Genomics for Ruminants in Developing Countries: From Principles to Practice. Front Genet. 9:251.
https://doi.org/10.3389/fgene.2018.00251

Ertl, J., A. Legarra, Z. G. Vitezica, L. Varona, C. Edel, R. Emmerling, and K. U. Götz. 2014.
Genomic analysis of dominance effects on milk production and conformation traits in Fleckvieh
cattle. Genet Sel Evol. 46:40. https://doi.org/10.1186/1297-9686-46-40

Fuerst, C. and J. Sölkner. 1994. Additive and Nonadditive Genetic Variances for Milk Yield,
Fertility, and Lifetime Performance Traits of Dairy Cattle. J Dairy Sci. 77:1114–1125. https://doi.org/
10.3168/jds.S0022-0302(94)77047-8

Gantner, V., S. Jovanovac, N. Raguž, M. Klopčič and D. Solić. 2008. Prediction of lactation milk
yield using various milk recording methods. Biotechnol Anim Husb. 24: 9–18.

García-Ruiz, A., J. B. Cole, P. M. VanRaden, G. R. Wiggans, F. J. Ruiz-López, and C. P. V. Tassell.
2016. Changes in genetic selection differentials and generation intervals in US Holstein dairy cattle
as a result of genomic selection. PNAS. 113:E3995–E4004. https://doi.org/10.1073/pnas.1519061113

Goddard, M. 2009. Genomic selection: Prediction of accuracy and maximisation of long term
response. Genetica. 136:245–257. https://doi.org/10.1007/s10709-008-9308-0

Goddard, M. E., B. J. Hayes, and T. H. E. Meuwissen. 2011. Using the genomic relationship matrix
to predict the accuracy of genomic selection. J Anim Breed Genet. 128:409–421.
https://doi.org/10.1111/j.1439-0388.2011.00964.x

Gonzalez-Recio, O., M. P. Coffey, and J. E. Pryce. 2014. On the value of the phenotypes in the
genomic era. J Dairy Sci. 97:7905–7915. https://doi.org/10.3168/jds.2014-8125

Habier, D., J. Tetens, F. R. Seefried, P. Lichtner, and G. Thaller. 2010. The impact of genetic
relationship information on genomic breeding values in German Holstein cattle. Genet Sel Evol.
42:5. https://doi.org/10.1186/1297-9686-42-5

Heslot, N. and V. Feoktistov. 2017. Optimization of selective phenotyping and population design for
genomic prediction. BioRxiv. 172064. https://doi.org/10.1101/172064

ICAR DNA Working Group. 2017. ICAR Guidelines for Parentage Verification and Parentage
Discovery Based on SNP Genotypes. The International Committee for Animal Recording.

International Committee for Animal Recording. 2017. Section 02—Cattle Milk Recording.
Overview. In ICAR Guidelines. ICAR.

International Committee for Animal Recording. 2020. Section 02—Cattle Milk Recording. Procedure
2—Computing Lactation Yield. In ICAR Guidelines. ICAR.

Jenko, J., G. R. Wiggans, T. A. Cooper, S. A. E. Eaglen, W. G. de L. Luff, M. Bichard, R. Pong-
Wong, and J. A. Woolliams. 2017. Cow genotyping strategies for genomic selection in a small dairy
cattle population. J Dairy Sci. 100:439–452. https://doi.org/10.3168/jds.2016-11479

Jiang, J., B. Shen, J. R. O'Connell, P. M. VanRaden, J. B. Cole, and L. Ma. 2017. Dissection of
additive, dominance, and imprinting effects for production and reproduction traits in Holstein cattle.
BMC Genomics. 18:425. https://doi.org/10.1186/s12864-017-3821-4

Jibrila, I., J. ten Napel, J. Vandenplas, R. F. Veerkamp, and M. P. L. Calus. 2020. Investigating the
impact of preselection on subsequent single-step genomic BLUP evaluation of preselected animals.
Genet Sel Evol. 52:42. https://doi.org/10.1186/s12711-020-00562-6

Jopson, N. B., P. R. Amer, and J. M. McEwan. 2004: Comparison of two-stage selection breeding
programmes for terminal sire sheep. Pages 212-216 in Proceedings of the New Zealand Society of
Animal Production 64.

Kasap, A., B. Mioc, J. M. Hickey, and G. Gorjanc. 2018. Genetic connectedness in the U.S. sheep
industry. Page 590 in Book of Abstracts of the 69th Annual Meeting of the European Federation of
Animal Science. Wageningen Academic Publishers, Wageningen, Netherlands.
https://doi.org/10.3920/978-90-8686-871-1

Lidauer, M., E. A. Mäntysaari, and I. Strandén. 2003. Comparison of test-day models for genetic
evaluation of production traits in dairy cattle. Livest Prod Sci. 79: 73–86.
https://doi.org/10.1016/S0301-6226(02)00142-2

Lillehammer, M., T. H. E.Meuwissen, and A. K. Sonesson. 2011. A comparison of dairy cattle
breeding designs that use genomic selection. J Dairy Sci. 94:493–500.
https://doi.org/10.3168/jds.2010-3518

Marshall, K., J. P. Gibson, O. Mwai, J. M. Mwacharo, A. Haile, T. Getachew, R. Mrode, and S. J.
Kemp. 2019. Livestock Genomics for Developing Countries – African Examples in Practice. Front
Genet, 10:297. https://doi.org/10.3389/fgene.2019.00297

Meuwissen, T. H. E., B. J. Hayes, and M. E. Goddard. 2001. Prediction of total genetic value using
genome-wide dense marker maps. Genetics. 157:1819–1829.

Meyer, K. 1984. Estimates of genetic parameters for milk and fat yield for the first three lactations in
British Friesian cows. Animal Science. 38:313–322. https://doi.org/10.1017/S0003356100041519

Misztal, I., A. Legarra, and I. Aguilar. 2014. Using recursion to compute the inverse of the genomic
relationship matrix. J Dairy Sci. 97:3943–3952. https://doi.org/10.3168/jds.2013-7752

Misztal, I., S. Tsuruta, D. A. L. Lourenco, Y. Masuda, I. Aguilar, A. Legarra, and Z. Vitezica. 2018.
Manual for BLUPF90 family programs. University of
Georgia. http://nce.ads.uga.edu/wiki/doku.php?id=documentation

Mrode, R. A. 2005. Linear Models for the Prediction of Animal Breeding Values (Second edition).
CAB International, Wallingford, UK.

Mrode, R., J. M. K. Ojango, A. M. Okeyo, and J. M. Mwacharo. 2019. Genomic Selection and Use
of Molecular Tools in Breeding Programs for Indigenous and Crossbred Cattle in Developing
Countries: Current Status and Future Prospects. Front Genet. 9:964.
https://doi.org/10.3389/fgene.2018.00694

Obšteter, J., J. Jenko, J. M. Hickey, and G. Gorjanc. 2019. Efficient use of genomic information for
sustainable genetic improvement in small cattle populations. J Dairy Sci. 102:9971–9982.
https://doi.org/10.3168/jds.2019-16853

Ødegård, J., U. Indahl, I. Strandén, and T. H. E. Meuwissen. 2018. Large-scale genomic prediction
using singular value decomposition of the genotype matrix. Genet Sel Evol. 50:6.
https://doi.org/10.1186/s12711-018-0373-2

Pool, M. H. and T. H. E. Meuwissen. 1999. Prediction of Daily Milk Yields from a Limited Number
of Test Days Using Test Day Models. J Dairy Sci. 82:1555–1564. https://doi.org/10.3168/jds.S0022-
0302(99)75383-X

Powell, O., R. Mrode, C. R. Gaynor, M. Johnsson, G. Gorjanc, and J. M. Hickey. 2019. Genomic
data enables genetic evaluation using data recorded on LMIC smallholder dairy farms. BioRxiv.
827956. https://doi.org/10.1101/827956

Pryce, J. E., M. E. Goddard, H. W. Raadsma, and B. J. Hayes. 2010. Deterministic models of
breeding scheme designs that incorporate genomic selection. J Dairy Sci. 93:5455–5466.
https://doi.org/10.3168/jds.2010-3256

Reiner-Benaim, A., E. Ezra, and J. I. Weller. 2017. Optimization of a genomic breeding program for
a moderately sized dairy cattle population. J Dairy Sci. 100:2892–2904.
https://doi.org/10.3168/jds.2016-11748

Schaeffer, L. R. 2006. Strategy for applying genome-wide selection in dairy cattle. J Anim Breed
Genet. 123:218–223. https://doi.org/10.1111/j.1439-0388.2006.00595.x

Sonesson, A. K., J. A. Woolliams, and T. H. E. Meuwissen. 2012. Genomic selection requires
genomic control of inbreeding. Genet Sel Evol. 44:27. https://doi.org/10.1186/1297-9686-44-27

Swalve, H. and L. D. V. Vleck. 1987. Estimation of Genetic (Co) Variances for Milk Yield in First
Three Lactations Using an Animal Model and Restricted Maximum Likelihood. J Dairy Sci. 70:842–
849. https://doi.org/10.3168/jds.S0022-0302(87)80082-6

Van Grevenhof, E. M., J. A. M. Van Arendonk, and P. Bijma. 2012. Response to genomic selection:
The Bulmer effect and the potential of genomic selection when the number of phenotypic records is
limiting. Gene Sel Evoul. 44:26. https://doi.org/10.1186/1297-9686-44-26

Verbič, J., J. Jenko, J. Jeretina, and D. Babnik. 2019. Milk urea concentration as a tool to reduce the
nitrogen footprint of milk production in conditions of small scale farming. In Towards Precision
Livestock Husbandry and Its Potential to Mitigate Ammonia and GHG Emissions: Abstracts' Leaflet.
The 4th liveAGE meeting, Galilee, Israel.

Wiggans, G. R., J. B. Cole, S. M. Hubbard, and T. S. Sonstegard. 2017. Genomic Selection in Dairy
Cattle: The USDA Experience. Annu Rev Anim Biosci. 5:309-327. https://doi.org/10.1146/annurev-
animal-021815-111422

Wolc, A., J. Arango, P. Settar, J. E. Fulton, N. P. O'Sullivan, R. Preisinger, D. Habier, R. Fernando,
D. J. Garrick, and J. C. Dekkers. 2011. Persistence of accuracy of genomic estimated breeding values
over generations in layer chickens. Genet Sel Evol. 43:23. https://doi.org/10.1186/1297-9686-43-23

Young, S. 1964. Multi-stage selection for genetic gain. Heredity. 19**:**131–145.
https://doi.org/10.1038/hdy.1964.11

Yu, H., M. L. Spangler, R. M. Lewis, and G. Morota. 2017. Genomic Relatedness Strengthens
Genetic Connectedness Across Management Units. G3 (Bethesda). 7:3543–3556.
https://doi.org/10.1534/g3.117.300151

Yu, H., M. L. Spangler, R. M. Lewis, and G. Morota. 2018. Do stronger measures of genomic
connectedness enhance prediction accuracies across management units? J Anim Sci. 96: 4490–4500.
https://doi.org/10.1093/jas/sky316

Table 1: Number of genotyped animals per year by scenario and relative cost of phenotyping to
genotyping ($P:$G). Scenarios are named "G" for genomic, followed by the number of phenotype
records per lactation. For the $P:$G we compared the cost of 11 phenotype records per lactation to
the cost of one genotype. The number of phenotype records and the $P:$G dictated the number of
genotyped animals. We genotyped females (F) and males (M) in 7:1 ratio. We genotyped the females
to update and increase the training population and males for selection.

Relative cost	Scenario					
	G10	G9	G8	G5	G2	G1
$P:$G = 1:2	160 F	350 F	590 F	1,610 F	3,230 F	3,850 F
	22 M	50 M	85 M	235 M	465 M	565 M
$P:$G = 1:1	310 F	700 F	1,180 F	3,230 F	3,850 F	3,850 F
	45 M	100 M	165 M	465 M	925 M	1,125 M
$P:$G = 2:1	620 F	1,400 F	2,360 F	3,850 F	3,850 F	3,850 F
	90 M	200 M	335 M	925 M	1,845 M	2,245 M

718
719 **Figure 1. Genetic gain and training population size by scenario and relative cost of phenotyping**
720 **to genotyping ($P:$G) with an initial training population (TP).** The figure presents the means
721 (lines) and 95% confidence intervals (polygons) across 10 replicates for the conventional (C) and
722 genomic (G) scenarios, with numbers indicating the number of phenotype records per lactation. For
723 the $P:$G we compared the cost of 11 phenotype records per lactation to the cost of one genotype.

Figure 2. Accuracy by scenario with an initial training population and equal cost of phenotyping and genotyping. The figure presents the means (lines) and 95% confidence intervals (error bars) across 10 replicates for the conventional (C) and genomic (G) scenarios with numbers indicating the number of phenotype records per lactation. The cost of phenotyping was defined as the cost of 11 phenotype records per lactation, and the cost of genotyping as the cost of one genotype. Conventional selection implemented two-stage selection for males, hence we present the accuracy of pre-selection for progeny testing (empty point) and the accuracy of sire selection (solid point).

Figure 3. Genetic gain, training population size, and accuracy by scenario without an initial training population (TP) and equal cost of phenotyping and genotyping. The figure presents the means (lines or points) and 95% confidence intervals (polygons or error-bars) across 10 replicates for the conventional (C) and genomic (G) scenarios with numbers indicating the number of phenotype records per lactation. The cost of phenotyping was defined as the cost of 11 phenotype records per lactation, and the cost of genotyping as the cost of one genotype.The red line marks the condition of required 2,000 training animals to start genomic selection. Conventional selection implemented two-stage selection for males, hence we present the accuracy of the pre-selection for progeny testing (empty point) and the accuracy of sire selection (solid point).

Supplementary Material

1. Supplementary Figures and Tables

a. Supplementary Tables

Table S1 Accuracy of conventional and genomic selection with varying number of phenotypes and phenotyped animals. NoRec = Number of phenotypic records per lactation, NoDaughters = number or daughters per sire, r_{sire} = accuracy for sires, r_{cows} = accuracy for cows, $r_{non-pheno}$ = accuracy for non-phenotyped animals, NoPhenoCows = number of phenotyped cows, NoPhenoTotal = total number of phenotypes (number of phenotypes per lactation times the number of phenotyped cows).

NoPheno	NoDaughters	r_{sires}	r_{cows}	$r_{non-pheno}$	NoPhenoCows	NoPhenoTotal
Conventional selection, 100 sires						
Variable resources for phenotyping						
1	100	0.93	0.62	0.56	10,000	10,000
2	100	0.96	0.70	0.59	10,000	20,000
5	100	0.97	0.81	0.64	10,000	50,000
10	100	0.98	0.89	0.66	10,000	100,000
Fixed resources for phenotyping						
1	1000	0.99	0.63	0.59	100,000	100,000
2	500	0.99	0.71	0.61	50,000	100,000
5	200	0.99	0.82	0.64	20,000	100,000
10	100	0.98	0.89	0.66	10,000	100,000
Genomic selection						
Variable resources for phenotyping						
1	-	-	0.62	0.56	10,000	10,000
2	-	-	0.70	0.63	10,000	20,000
5	-	-	0.81	0.71	10,000	50,000
10	-	-	0.89	0.76	10,000	100,000
Fixed resources for phenotyping						
1	-	-	0.63	0.93	100,000	100,000
2	-	-	0.71	0.90	50,000	100,000
5	-	-	0.82	0.84	20,000	100,000
10	-	-	0.89	0.76	10,000	100,000

Table S2 Genetic gain by scenario, relative cost of phenotyping to genotyping ($P:$G), and
availability of an initial training population. The table presents the means and standard
deviations (subscript) across 10 replicates for the conventional (C) and genomic (G) scenarios, with
numbers indicating the number of phenotype records per lactation. For the $P:$G we compared the
cost of 11 phenotype records per lactation to the cost of one genotype. The scenarios in bold did not
spend all the available resources. Lower-case letters denote statistically significant differences
between scenarios within the same $P:$G and upper-case letters between different $P:$G within the
same scenario.

		Relative cost of phenotyping (P) to genotyping (G)		
	Scenario	$P:$G = 1:2	$P:$G = 1:1	$P:$G = 2:1
	C11	$3.01_{0.22}^{a,A}$	$3.01_{0.22}^{a,A}$	$3.01_{0.22}^{a,A}$
With an initial training population	G10	$5.43_{0.20}^{b,A}$	$5.41_{0.29}^{b,A}$	$6.50_{0.20}^{b,B}$
	G9	$5.58_{0.26}^{b,A}$	$6.30_{0.17}^{c,B}$	$7.02_{0.24}^{c,C}$
	G8	$6.35_{0.25}^{c,A}$	$6.62_{0.25}^{d,B}$	$7.02_{0.17}^{c,C}$
	G5	$6.78_{0.21}^{d,A}$	$7.07_{0.20}^{e,B}$	$\mathbf{7.26_{0.19}^{c,B}}$
	G2	$7.13_{0.29}^{e,A}$	$\mathbf{7.33_{0.26}^{e,A}}$	$\mathbf{7.28_{0.17}^{c,A}}$
	G1	$\mathbf{7.11_{0.16}^{e,A}}$	$7.27_{0.28}^{e,A}$	$7.24_{0.22}^{c,A}$
Without an initial training population	G10	$3.93_{0.22}^{b,A}$	$4.54_{0.14}^{b,B}$	$5.61_{0.25}^{b,C}$
	G9	$4.64_{0.18}^{c,A}$	$5.75_{0.28}^{c,B}$	$6.52_{0.17}^{c,C}$
	G8	$5.61_{0.28}^{d,A}$	$6.24_{0.19}^{d\,B}$	$6.70_{0.25}^{cd,C}$
	G5	$6.43_{0.21}^{e,A}$	$6.90_{0.22}^{e,B}$	$\mathbf{7.05_{0.27}^{de,B}}$
	G2	$6.81_{0.28}^{f,A}$	$\mathbf{6.96_{0.17}^{e,A}}$	$\mathbf{7.00_{0.30}^{de,A}}$
	G1	$\mathbf{6.78_{0.29}^{f,A}}$	$6.92_{0.26}^{e,A}$	$7.01_{0.23}^{e,A}$

Table S3 Intensity of sire selection by scenario and relative cost of phenotyping to genotyping
($P:$G). PA = parent average, gEBV = genomic breeding value. We are presenting the bivariate
intensities for a two-stage selection. The pre-selection step selects the animals with the highest
parent average out of all available new born males to send into testing (progeny of genomical). The
selection step selects the sires with the highest breeding values out of all tested males to use in
artificial insemination. The scenarios are named C/G for conventional/genomic with numbers
indicating the number of phenotype records per lactation. For the $P:$G we compared the cost of 11
phenotype records per lactation to the cost of one genotype.

		Relative cost of phenotyping (P) to genotyping (G)		
	Scenario	$P:$G = 1:2	$P:$G = 1:1	$P:$G = 2:1
	C11	0.10	0.10	0.10
With an initial training population	G10	0.17	0.27	0.37
	G9	0.28	0.39	0.55
	G8	0.38	0.49	0.68
	G5	0.59	0.77	0.91
	G2	0.77	0.91	0.98
	G1	0.83	0.92	0.99
Without an initial training population	G10	0.17	0.25	0.36
	G9	0.28	0.43	0.56
	G8	0.37	0.55	0.71
	G5	0.58	0.77	0.91
	G2	0.76	0.89	0.98
	G1	0.83	0.94	0.99

Table S4 Selection accuracy by scenario, relative cost of phenotyping to genotyping ($P:$G),
and the availability of an initial training population. The table presents the means and standard
deviations (subscript) across 10 replicates for the conventional (C) and genomic (G) scenarios, with
numbers indicating the number of phenotype records per lactation. For the $P:$G we compared the
cost of 11 phenotype records per lactation to the cost of one genotype. Conventional selection
implemented two-stage selection for males, hence we present the accuracy of pre-selection for
progeny testing (S1) and the accuracy sire selection (S2). In genomic scenarios the male candidates
were genotyped and non-phenotyped. We also present the accuracy for sires currently used in
artificial insemination (sires), for non-genotyped and non-phenotyped females (female candidates),
and for all active phenotyped cows and bull dams (cows). Lower-case letters denote statistically
significant differences between scenarios within the same $P:$G and upper-case letters between
different $P:$G within the same scenario. Stars denote statistically significant difference between
corresponding scenarios with and without an initial training population.

Scenario	With an initial training population			Without an initial training population		
	$P:$G=1:2	$P:$G=1:1	$P:$G=2:1	$P:$G=1:2	$P:$G=1:1	$P:$G=2:1
Male candidates						
C11, S1	$0.37_{0.04}^{a,A}$	$0.37_{0.04}^{a,A}$	$0.37_{0.04}^{a,A}$	$0.37_{0.04}^{a,A}$	$0.37_{0.04}^{a,A}$	$0.37_{0.04}^{a,A}$
C11, S2	$0.94_{0.01}^{b,A}$	$0.94_{0.01}^{b,A}$	$0.94_{0.01}^{b,A}$	$0.94_{0.01}^{b,A}$	$0.94_{0.01}^{b,A}$	$0.94_{0.01}^{b,A}$
G10	$0.89_{0.03}^{c,A}$	$0.90_{0.02}^{bc,AB}$	$0.91_{0.01}^{bc,B}$	$0.81_{0.03}^{b,A}$ *	$0.84_{0.01}^{b,B}$ *	$0.87_{0.01}^{b,C}$ *
G9	$0.90_{0.03}^{bc,A}$	$0.91_{0.02}^{bc,A}$	$0.91_{0.01}^{bc,A}$	$0.85_{0.02}^{c,A}$ *	$0.87_{0.01}^{bc,B}$ *	$0.90_{0.01}^{bc,C}$ *
G8	$0.91_{0.01}^{bc,A}$	$0.91_{0.01}^{bc,A}$	$0.91_{0.01}^{bc,A}$	$0.86_{0.01}^{cd,A}$ *	$0.89_{0.01}^{c,B}$ *	$0.90_{0.01}^{bc,B}$
G5	$0.91_{0.01}^{bc,A}$	$0.91_{0.00}^{bc,A}$	$0.91_{0.01}^{bc,A}$	$0.90_{0.01}^{d,A}$	$0.91_{0.01}^{c,A}$	$0.91_{0.01}^{c,A}$
G2	$0.91_{0.01}^{bc,A}$	$0.91_{0.00}^{bc,A}$	$0.90_{0.01}^{bc,A}$	$0.90_{0.01}^{d,A}$	$0.90_{0.01}^{c,A}$	$0.90_{0.01}^{bc,A}$
G1	$0.89_{0.01}^{c,A}$	$0.90_{0.01}^{c,A}$	$0.89_{0.01}^{c,A}$	$0.89_{0.01}^{cd,A}$	$0.89_{0.01}^{c,A}$	$0.89_{0.01}^{bc,A}$
Sires						
C11	$0.86_{0.05}^{a,A}$	$0.86_{0.05}^{a,A}$	$0.86_{0.05}^{a,A}$	$0.86_{0.05}^{a,A}$	$0.86_{0.05}^{a,A}$	$0.86_{0.05}^{a,A}$
G10	$0.75_{0.04}^{b,A}$	$0.75_{0.03}^{b,A}$	$0.73_{0.05}^{b,A}$	$0.67_{0.08}^{bc,A}$ *	$0.68_{0.05}^{cde,A}$ *	$0.67_{0.06}^{b,A}$ *
G9	$0.76_{0.04}^{b,A}$	$0.72_{0.06}^{bc,AB}$	$0.69_{0.05}^{c,A}$	$0.70_{0.05}^{b,A}$ *	$0.72_{0.05}^{bc,A}$	$0.71_{0.05}^{b,A}$
G8	$0.76_{0.03}^{b,A}$	$0.69_{0.05}^{cd,B}$	$0.68_{0.06}^{c,B}$	$0.71_{0.05}^{b,A}$ *	$0.74_{0.05}^{b,A}$ *	$0.70_{0.07}^{b,A}$
G5	$0.68_{0.07}^{c,A}$	$0.67_{0.08}^{de,A}$	$0.69_{0.04}^{c,A}$	$0.68_{0.05}^{bc,A}$	$0.69_{0.05}^{cd,A}$	$0.69_{0.03}^{b,A}$
G2	$0.67_{0.05}^{c,A}$	$0.67_{0.05}^{de,A}$	$0.67_{0.04}^{c,A}$	$0.65_{0.06}^{c,A}$	$0.64_{0.07}^{e,A}$	$0.69_{0.05}^{b,A}$
G1	$0.66_{0.06}^{c,A}$	$0.63_{0.05}^{e,A}$	$0.67_{0.04}^{c,A}$	$0.67_{0.04}^{bc,A}$	$0.67_{0.03}^{de,A}$	$0.69_{0.05}^{b,A}$
Female candidates						
C11	$0.45_{0.02}^{a,A}$	$0.45_{0.02}^{a,A}$	$0.45_{0.02}^{a,A}$	$0.45_{0.02}^{a,A}$	$0.45_{0.02}^{a,A}$	$0.45_{0.02}^{a,A}$
G10	$0.48_{0.01}^{ab,A}$	$0.48_{0.01}^{ab,A}$	$0.51_{0.01}^{b,B}$	$0.46_{0.02}^{ab,A}$ *	$0.47_{0.02}^{ab,AB}$	$0.49_{0.01}^{b,B}$ *
G9	$0.49_{0.02}^{b,A}$	$0.50_{0.01}^{b,B}$	$0.52_{0.01}^{b,C}$	$0.47_{0.02}^{ab,A}$ *	$0.49_{0.02}^{bc,B}$	$0.52_{0.01}^{bc,C}$
G8	$0.51_{0.01}^{b,A}$	$0.51_{0.01}^{b,A}$	$0.54_{0.01}^{bc,B}$	$0.49_{0.02}^{bc,A}$ *	$0.52_{0.01}^{cd,B}$	$0.53_{0.01}^{cd,C}$
G5	$0.51_{0.01}^{bc,A}$	$0.55_{0.01}^{c,B}$	$0.57_{0.01}^{c,C}$	$0.52_{0.01}^{cd,A}$	$0.55_{0.01}^{de,B}$	$0.57_{0.01}^{d,C}$
G2	$0.55_{0.01}^{cd,A}$	$0.57_{0.01}^{c,B}$	$0.57_{0.01}^{c,B}$	$0.55_{0.01}^{d,A}$	$0.56_{0.02}^{e,AB}$	$0.57_{0.01}^{d,B}$
G1	$0.56_{0.01}^{d,A}$	$0.56_{0.01}^{c,A}$	$0.56_{0.01}^{c,A}$	$0.55_{0.01}^{d,A}$	$0.56_{0.01}^{e,A}$	$0.56_{0.01}^{d,A}$
Cows						
C11	$0.48_{0.03}^{a,A}$	$0.48_{0.03}^{a,A}$	$0.48_{0.03}^{a,A}$	$0.48_{0.03}^{a,A}$	$0.48_{0.03}^{a,A}$	$0.48_{0.03}^{a,A}$
G10	$0.56_{0.02}^{b,A}$	$0.59_{0.02}^{b,B}$	$0.63_{0.01}^{b,C}$	$0.53_{0.01}^{b,A}$ *	$0.56_{0.01}^{b,B}$ *	$0.61_{0.01}^{b,C}$ *
G9	$0.59_{0.03}^{bc,A}$	$0.63_{0.02}^{c,B}$	$0.70_{0.01}^{c,C}$	$0.57_{0.02}^{bc,A}$ *	$0.62_{0.02}^{c,B}$	$0.68_{0.02}^{c,C}$ *
G8	$0.62_{0.02}^{c,A}$	$0.67_{0.02}^{c,B}$	$0.74_{0.02}^{d,C}$	$0.60_{0.02}^{c,A}$ *	$0.66_{0.01}^{d,B}$	$0.73_{0.02}^{d,C}$
G5	$0.70_{0.02}^{d,A}$	$0.77_{0.01}^{d,B}$	$0.79_{0.02}^{e,C}$	$0.69_{0.02}^{d,A}$	$0.76_{0.01}^{e,B}$	$0.78_{0.02}^{e,B}$
G2	$0.76_{0.02}^{e,A}$	$0.79_{0.02}^{d,B}$	$0.78_{0.01}^{e,AB}$	$0.76_{0.01}^{e,A}$	$0.77_{0.02}^{e,A}$ *	$0.77_{0.01}^{de,A}$
G1	$0.77_{0.02}^{e,A}$	$0.77_{0.02}^{d,A}$	$0.77_{0.01}^{de,A}$	$0.76_{0.01}^{e,A}$	$0.76_{0.02}^{e,A}$	$0.76_{0.02}^{de,A}$

b. Supplementary Figures

Figure S1 The number of animals and repeated phenotypes in the training population by
scenario and relative cost of phenotyping to genotyping ($P:$G) with an initial training
population (TP). The scenarios are named C/G for conventional/genomic with numbers indicating
the number of phenotype records per lactation. For the $P:$G we compared the cost of 11
phenotype records per lactation to the cost of one genotype. In our simulation, scenarios traded
repeated phenotype records for genotypes. Hence, the scenarios with the largest training population
collected the least repeated records. These were also the scenarios that achieved the highest genetic
gain.