1 **Gene co-expression network analysis in human spinal cord highlights mechanisms**
2 **underlying amyotrophic lateral sclerosis susceptibility**
3

4 Jerry C. Wang[1], Gokul Ramaswami[1], Daniel H. Geschwind[1,2,3,4*]
5
6 Affiliations
7 [1] Program in Neurogenetics, Department of Neurology, David Geffen School of Medicine,
8 University of California, Los Angeles, Los Angeles, CA, United States of America
9
10 [2] Center for Autism Research and Treatment, Semel Institute, David Geffen School of Medicine,
11 University of California, Los Angeles, Los Angeles, CA, United States of America
12
13 [3] Department of Human Genetics, David Geffen School of Medicine, University of California, Los
14 Angeles, Los Angeles, CA, United States of America
15
16 [4] Institute for Precision Health, David Geffen School of Medicine, University of California, Los
17 Angeles, Los Angeles, CA, United States of America
18
*. Correspondence should be addressed to D.H.G: dhg@mednet.ucla.edu

Abstract

Amyotrophic lateral sclerosis (ALS) is a neurodegenerative disease defined by motor neuron

(MN) loss. Multiple genetic risk factors have been identified, implicating RNA and protein

metabolism and intracellular transport, among other biological mechanisms. To achieve a

systems-level understanding of the mechanisms governing ALS pathophysiology, we built gene

co-expression networks using RNA-sequencing data from control human spinal cord samples,

identifying 13 gene co-expression modules, each of which represents a distinct biological

process or cell type. Analysis of four RNA-seq datasets from a range of ALS disease-associated

contexts reveal dysregulation in numerous modules related to ribosomal function, wound

response, and leukocyte activation, implicating astrocytes, oligodendrocytes, endothelia, and

microglia in ALS pathophysiology. To identify potentially causal processes, we partitioned

heritability across the genome, finding that ALS common genetic risk is enriched within two

specific modules, SC.M4, representing genes related to RNA processing and gene regulation,

and SC.M2, representing genes related to intracellular transport and autophagy and enriched in

oligodendrocyte markers. Top hub genes of this module include ALS-implicated risk genes such

as KPNA3, TMED2, and NCOA4, the latter of which regulates ferritin autophagy, implicating this

process in ALS pathophysiology. These unbiased, genome-wide analyses confirm the utility of a

systems approach to understanding the causes and drivers of ALS.

Introduction

Amyotrophic lateral sclerosis (ALS) is an adult-onset neurodegenerative disease defined by motor neuron (MN) loss and results in paralysis and eventually death 3-5 years following initial disease presentation[1]. In 5-10% of cases, patients exhibit a Mendelian pattern of inheritance and the disease segregates within multigenerational affected families (familial ALS)[2]. However, in 90% of cases, the underlying cause of ALS is unknown and disease occurrence is classified as sporadic (sporadic ALS)[2]. Most genetic risk in ALS is polygenic, indicating a complex genetic architecture[3]. Given this complex genetic architecture with contributions of rare and common alleles, we took a systems level approach to understand the gene expression signatures associated with ALS pathophysiology.

Weighted gene co-expression network analysis (WGCNA)[4] has been a powerful method for organizing and interpreting wide-ranging transcriptional alterations in neurodegenerative diseases. In Alzheimer's disease (AD), transcriptomic and proteomic studies have revealed a pattern of dysregulation implicating immune- and microglia-specific processes[5,6]. More broadly across tauopathies including AD, frontotemporal dementia (FTD), progressive supranuclear palsy (PSP), molecular network analyses have identified gene modules and microRNAs mediating neurodegeneration, highlighting its utility in prioritizing molecular targets for therapeutic interventions[7–10]. In ALS, previous studies have studied the transcriptional landscape in sporadic and C9orf72 (familial) ALS post-mortem brain tissues, finding a pattern of dysregulation in gene expression, alternative splicing, and alternative polyadenylation that disrupted RNA processing, neuronal functioning, and cellular trafficking[11]. However, the generalizability of these findings is limited due to relatively small sample sizes (n=26). Numerous studies have previously examined the transcriptional landscape of the spinal cord, which hosts axons from the upper motor neurons (UMN) and somae of the lower motor neuron

(LMN) and is the location of LMN degeneration in ALS[12–17]. However, a rigorous interrogation of

gene co-expression networks in the spinal cord has not yet been fully explored.

To investigate gene expression signatures associated with ALS in the spinal cord, we generated

gene co-expression networks using RNA-sequencing of 62 neurotypical human cervical spinal

cord samples from the GTEx consortium[18]. We characterized these co-expression modules

through gene ontology (GO) enrichment analysis, which reveals that the modules represent a

diverse range of biological processes. To identify modules enriched with ALS genetic risk

factors, we queried enrichment of rare, large effect risk mutations previously described in the

literature, as well as common genetic variation[19,20]. The second approach is genome-wide and

therefore unbiased, which focuses on the enrichment of small effect common risk variants

identified from a Genome Wide Association Study (GWAS) of ALS[21]. By partitioning SNP based

heritability and performing enrichment testing for genes with rare mutations, we find significant

overlap of genes harboring rare and common variation in SC.M4, which represents genes

involved in RNA processing and epigenetic regulation. We also find enrichment of ALS common

genetic risk variants in SC.M2, which represents genes involved in intracellular transport,

protein modification, and cellular catabolic processes. We extend these results by

demonstrating the dysregulation of these modules in data from multiple published human ALS

patient and animal model tissues[12,13,22,23]. We identify modules related to ribosomal function,

wound response, and leukocyte activation that are dysregulated across this diverse set of ALS-

related datasets. The set of genes underlying these modules provides a rich set of targets for

exploring new mechanisms for therapeutic development.

Results

Co-expression network analysis as a strategic framework for elucidating ALS susceptibility

We reasoned that because ALS preferentially affects spinal cord and upper motor neurons, co-

expression networks from spinal cords of control subjects would provide an unbiased context for

understanding potential convergence of ALS risk genes. We used the Genotype-Tissue

Expression (GTEx) Project[18] control cervical spinal cord samples to generate a co-expression

network as a starting point for a genome wide analysis to identify specific pathways or cell types

affected by ALS genetic risk. First, we identified gene co-expression modules and assigned

each module a biological, as well as cell-type identity[24,25]. Next, we leveraged multiple datasets

identifying ALS genetic risk factors[19–21,26–28] to comprehensively assess which pathways and

cell-types are susceptible to ALS genetic risk. Finally, we compared these predictions with

human ALS and animal model datasets[12,13,22,23] to develop a mechanistic understanding of ALS

etiology and progression (**Fig. 1**).

Generation of human spinal cord gene co-expression network

We utilized RNA-sequencing data from the GTEx consortium to generate a co-expression

network of human spinal cord (**Methods**). Co-expression networks are highly sensitive to

biological and technical confounders[25], which we controlled for by calculating the correlation of

biological and technical covariates with the top expression principal components (PCs)

(**Methods, Supplementary Fig. S1a-c**). We identified 11 technical and biological covariates:

seqPC1-5, RNA integrity number (RIN), Hardy's death classification scale, age, sampling

center, ethnicity, and ischemic time as significant drivers of expression (**Supplementary Fig.**

S1a). We regressed out the effect of these covariates using a linear model (**Methods**) and

verified that the regressed expression dataset was no longer correlated with technical and

biological covariates (**Supplementary Fig. S1b**).

We applied WGCNA to construct a co-expression network (**Fig. 2a, Supplementary Fig. S2;**

Supplementary Data 1; Methods) identifying 13 modules. To verify that the modules were not

driven by confounding covariates, we confirmed that the module eigengenes were not

correlated with the major technical and biological covariates (**Supplementary Fig. S1d**). We

also found that randomly sampled genes from within each module were significantly more inter-

correlated when compared to a background set, verifying the co-expression of genes within

each module (**Supplementary Fig. S1e**). To biologically annotate our co-expression modules,

we performed gene ontology (GO) term enrichment and identified the top hubs (**Fig. 2c-g, 3c-d,**

Supplementary Fig. S3). GO analysis revealed that co-expression modules represented

coherent biological processes that we categorized into five distinct groupings (**Fig. 2b**),

encompassing morphogenesis and development, epigenetics and gene regulation, neuronal

signaling, immune activation, and intracellular transport, metabolism, and sensing functions.

Most modules were significantly enriched for markers of specific cell types, suggesting that they

represent gene co-expression within major cell classes, including SC.M1 (astrocytes), SC.M2

(oligodendrocytes), SC.M3 (endothelial cells), SC.M9 and SC.M11 (neurons), SC.M10

(ependymal), and SC.M12 and SC.M13 (microglia) (**Supplementary Fig. S4**). These groupings,

both with regards to cell type and molecular pathways, represent diverse biological functions

that play key roles in human spinal cord biology.

Enrichment of ALS genetic risk in intracellular transport/autophagy module SC.M2 and RNA

processing/gene regulation module SC.M4

To address how these modules fit within the context of known ALS susceptibility factors, we

assessed whether a literature-curated set of well-known, large effect size, familial ALS risk

genes[19,20] were enriched within our co-expression modules (**Fig. 3a**). We found a nominally

significant enrichment with SC.M4 (OR = 2.52, P = 0.0124). However, the FDR adjusted p-value

was not significant (FDR= 0.174), likely due to the small number of familial ALS risk genes

identified to date. SC.M4 was highly enriched with GO terms related to RNA processing and

epigenetic regulation (**Fig. 3c**), consistent with the known role of several ALS risk genes in RNA

metabolism and function. To prioritize interacting partners to ALS risk genes found in SC.M4,

we leveraged a direct protein-protein interaction (PPI) network using the top 300 hub genes of

SC.M4, finding a significant network that includes ALS risk genes such as TARDBP and

EWSR1 (**Fig. 3e**, **Methods**, p = 0.001).

The small number (~50) of large effect size risk genes identified to date in ALS makes the

detection of strong enrichments difficult. To complement the rare risk gene enrichment

analyses, we performed an unbiased genome-wide analysis, partitioning the common SNP-

based heritability across the co-expression modules using stratified LD score-regression[29]

based on summary statistics from the most recent GWAS of ALS (ALS n= 20,806; CTL n =

59,804)[21] (**Fig. 3b**). We found significant enrichment of ALS common risk variants in the SC.M2

(FDR = 0.047) and SC.M4 modules (FDR = 0.041). SC.M2 was enriched with GO terms related

to intracellular transport, protein modification, and cellular catabolic processes (**Fig. 3d**).

Additionally, SC.M2 was significantly enriched with oligodendrocyte markers (**Supplementary**

Fig. S4, Oligodendrocytes_mature: OR = 5.3, FDR < 10e-5; Oligodendrocytes_myelinating: OR

= 2.9, FDR < 0.05), while SC.M4 showed no specific cell type enrichment. Generation of a direct

protein-protein interaction (PPI) network using the top 300 hub genes of SC.M2 reveals a

significant network that highlights SC.M2 hub genes that interact with known ALS risk genes

such as TMED2 and KPNA3[27] (**Fig. 3f**, **Methods**, p = 0.018).

As a comparison, we tested for the enrichment of common risk variants from Inflammatory

Bowel Disorder (IBD)[30] (N=86,640) across modules (**Fig. 3b**). We found a significant enrichment

of IBD common risk variants (FDR = 0.047) in module, SC.12, which is a microglial/immune

module, consistent with the known disease biology of IBD[31]. As a negative control, we

performed enrichments of common risk variants from Autism Spectrum Disorder (ASD)[32,33]

(N=15,954) and found no enrichment in any module. Further, when we compared with

Alzheimer's Disease (AD)[34] (N=455,258), Frontotemporal Dementia (FTD)[35] (N=12,928), and

Progressive Supranuclear Palsy (PSP)[36] (N=12,308) (**Fig. 3b**), we also found no significant

enrichments for these diseases' common risk variants, demonstrating the specificity of the ALS

GWAS enrichment for modules SC.M2 and SC.M4. Overall, these genetic enrichment analyses

implicate dysfunction in RNA processing and intracellular transport, including in

oligodendrocytes, as putative drivers of ALS pathophysiology in the human spinal cord.

Enrichment of ALS genetic modifiers in ribosomal-associated module SC.M6

As an orthogonal approach to assess which genes and pathways are mediators of ALS

pathophysiology, we leveraged genetic screen datasets from three studies of modifiers of

C9orf72 and FUS toxicity in yeast[26–28]. We assessed our human spinal cord modules for

enrichment with modifiers of C9orf72 and FUS toxicity identified from these screens. We found

significant enrichment of C9orf72 and FUS toxicity suppressors in SC.M6 (**Fig. 4a**), a module

representing genes involved in ribosomal function (**Fig. 2e**), but without any cell type specific

enrichment (**Supplementary Fig. S4**). This finding is consistent with previous observations that

modifiers of Glycine-Arginine$_{100}$ (GR$_{100}$) DPR and FUS toxicity are enriched with gene ontology

terms related to ribosomal biogenesis[26,28].

We reasoned that SC.M6 could be leveraged to prioritize highly interconnected hub genes

within the module that may warrant additional scrutiny as points of convergence for C9orf72 and

FUS pathology. To address this, we first identified the interacting partners for the top 100

SC.M6 hub genes by generating an indirect protein-protein interaction (PPI) network (**Methods**).

We identified 30 modifiers of C9orf72 and FUS toxicity that were present in this indirect PPI

network, either as hubs or as interacting partners (**Fig. 4b**)[26–28]. These modifiers include

ribosomal proteins (RPL and RPS-related genes), intracellular transporters of ribosomal

components (TNPO1), ribosomal assembly proteins (NOB1), and RNA-binding proteins

(HABP4) (**Fig. 4b**), indicating specific aspects of ribosomal dysfunction shared between C9orf72

and FUS toxicity. Indeed, previous studies[26] have suggested that the overlap in modifiers of

C9orf72 and FUS toxicity could be due to the sequence similarity between the

arginine/glycine/glycine (RGG) domain in FUS and the GR_{100} sequence expressed by

C9orf72[37,38].

Dysregulation of immune-related modules in postmortem ALS spinal cord

Until this point, we had assessed co-expression modules for their relevance to ALS based on

enrichment of genetic risk factors. To assess whether the co-expression modules identified

using the GTEx dataset were directly dysregulated in ALS patients, we processed bulk RNA-

sequencing data from a previous study of postmortem control and ALS cervical spinal cord[13].

We corrected the data for potential confounders such as age, ethnicity, sex, and sequencing

variability and assessed module preservation (**Supplementary Fig. S5a-b**, **Methods**). We

identified 10 of the 13 modules generated from GTEx as preserved in the postmortem control

and ALS cervical spinal cord ($Z_{summary} > 2$) with four (SC.M2, SC.M4, SC.M5, and SC.M8)

strongly preserved ($Z_{summary} > 10$). The remaining three modules (SC.M9, SC.M11, and SC.M10)

were not preserved ($Z_{summary} < 2$) (**Supplementary Fig. S5c**).

We adopted two approaches to assess which modules were dysregulated in ALS in the human

spinal cord. The first assessed enrichments of differentially expressed genes (DEGs) between

postmortem control and ALS cervical spinal cord samples[13] (**Methods**). We found a significant

enrichment of DEGs upregulated in cervical spinal cord from ALS patients in SC.M12 (FDR =

5.92e-39) and of downregulated DEGs in SC.M3 (FDR = 0.00152) (**Fig. 5a**). SC.12 is highly

enriched with microglial markers and GO terms related to immune response (**Fig. 2f,**

Supplementary Fig. S4a), whereas SC.M3 is highly enriched with endothelial markers and GO

terms related to inflammation and morphogenesis (**Fig. 2c, Supplementary S4a**).

For our second approach, we assessed if the spinal cord modules were themselves differentially

expressed between ALS and control samples by looking at the relationship between the

eigengenes (first principle component; **Methods**) of each module and disease status

(**Supplementary Fig. S5d**). We found that the SC.M1 eigengene was significantly upregulated

in ALS samples. SC.M1 was enriched with astrocyte markers and GO terms related to wound

response (**Supplementary Fig. S3a, S4a**). Overall, our analyses identify a marked upregulation

of neuro-immune cells in the human post-mortem spinal cord, including astrocytes and

microglia, similar to previous reports[13,22]. However, we note that genes differentially expressed

in postmortem ALS spinal cord are not enriched in causal genetic risk, suggesting that these

changes are not causal drivers.

Dysregulation of immune and neuronal related modules in dissected human motor neurons

primed for degeneration

A major disadvantage of utilizing postmortem human tissues is that the molecular pathology is

indicative of late-stage disease. In particular, immune hyperactivation and infiltration is

consistently seen in neurodegenerative disorders including ALS. However, it is unclear if these

processes are disease causing or merely a reaction to neurodegenerative cell death[39]. To

identify early-stage differences in human spinal cord potentially representative of causal factors,

we leveraged a previously published dataset of laser capture micro-dissected motor neurons

deemed to be primed for degeneration[22]. In this study, the authors isolated motor neurons from

ALS patients' postmortem spinal cord, but in regions that did not exhibit any signs of

degeneration at the time of death. We processed and corrected the data for potential

confounders such as age, gender, postmortem interval, RNA integrity, and sequencing

variability[22] (**Supplementary Fig. S6a-b**). We identified 9 of the 13 modules generated from

GTEx as preserved in the micro-dissected motor neuron dataset ($Z_{summary} > 2$) with two (SC.M3

and SC.M6) strongly preserved ($Z_{summary}$ > 10). The remaining four modules (SC.M2, SC.M4,

SC.M8, and SC.M10) were not preserved ($Z_{summary}$ < 2) (**Supplementary Fig. S6c**).

We assessed the enrichment of DEGs between ALS and control micro-dissected motor neurons

in our modules. We found enrichment of up-regulated DEGs within SC.M1, SC.M3, SC.M6, SC.

M12, and SC.M13, modules related to astrocyte, endothelial, and microglial functions

(**Supplementary Fig. S4a**). We found enrichment of down-regulated DEGs within SC.M7 and

SC.M9 (**Fig. 5b**), two modules related to synaptic signaling (**Fig. 2d, Supplementary Fig. S3b**),

possibly marking degeneration at early stages of ALS. We also assessed the relationship

between the module eigengene and disease status, finding significant up-regulation of SC.M1,

SC.M3, SC.M5, SC.M6, SC.12, and SC.M13 in the ALS samples (**Supplementary Fig. S6d**).

Despite performing laser-capture micro-dissection of motor neurons, our results indicate that the

predominant signal from this dataset highlights the infiltration of neuro-immune cells, including

astrocytes and microglia in the progression of ALS pathophysiology.

Dysregulation of a ribosome-related module in iPSC-derived motor neurons from ALS8 patients

To gain further insights into processes dysregulated in motor neurons and avoid the pro-

immune phenotypes observed in previous postmortem studies[13,22], we leveraged a RNA-seq

dataset generated from iPSC-derived motor neurons from ALS8 patients[23]. In this study, iPSC-

derived motor neurons were differentiated from five patients carrying pathogenic variants in the

VAPB gene and three unaffected control individuals. We processed and corrected this dataset

for potential confounders such as age, sex, sampled tissue, and sequencing variability

(**Supplementary Fig. S7a-b**). We identified 10 of the 13 modules generated from GTEx as

preserved in the iPSC-derived motor neuron dataset ($Z_{summary}$ > 2) with two (SC.M7 and SC.M9)

strongly preserved ($Z_{summary}$ > 10). The remaining three modules (SC.M5, SC.M12, and SC.M13)

were not preserved ($Z_{summary}$ < 2) (**Supplementary Fig. S7c**).

We assessed the enrichment of DEGs between iPSC-derived MNs from severe ALS versus

control individuals in our modules. We found modest enrichment of up-regulated DEGs within

SC.M1, SC.M7, SC.M9, and SC.M11 (OR = 1.77, 1.76, 2.09, 3.84 respectively), modules

related to astrocyte and neuronal functions (**Fig. 5c, Supplementary Fig. S4**). We found a

strong and pronounced enrichment of down-regulated DEGs within SC.M6 (OR = 27.28), a

ribosomal-related module (**Fig. 2e, 5c**). We also assessed the relationship of module eigengene

to disease status, revealing up-regulation of SC.M2 and SC.M7 and down-regulation of SC.M1

(**Supplementary Fig. S7d**). By analyzing gene expression in iPSC-derived motor neurons from

ALS patients, we identified striking downregulation of ribosomal function, represented by

SC.M6, as a prominent dysregulated feature specific to MNs.

Dysregulation of immune and neuronal-related modules in SOD1-G93A (ALS) mice

Finally, as an orthogonal approach to identify temporal differences in ALS, we leveraged a

spatial transcriptomics dataset from spinal cord of SOD1-G93A mice over four time points

during disease progression[12,40,41]. To assess which modules were dysregulated over the

progression of disease, we performed an enrichment analysis between the human spinal cord

modules and mouse DEGs stratified by time point (**Methods**). Consistent with our findings in

postmortem tissues (**Fig. 5a-b, Supplementary Fig. S6e**), immune response modules (SC.M12

and SC.M13) were highly enriched with up-regulated DEGs as early as postnatal day 70, which

is the time of clinical disease onset in this model (**Fig. 5c, 2f, Supplementary Fig. S3f**). We

also noted a strong enrichment of down-regulated DEGs in neuronal signaling modules (SC.M7,

SC.M9, and SC.M11) (**Supplementary Fig. S3b, Fig. 2d, Supplementary Fig. S3e**) by

postnatal day 120 (p120 - end-stage ALS), which is consistent with motor neuron degeneration.

Interestingly, we found enrichment of down-regulated DEGs in SC.M2 at p100 (symptomatic

ALS), providing evidence that SC.M2 is dysregulated at symptomatic stages of ALS in this

model system (**Fig. 5c, Supplementary Fig. S9**). We also noted an enrichment of up-regulated

DEGs in SC.M6 at p120 (end-stage ALS), suggesting dysregulation of SC.M6 in the end-stages

of ALS in this model system (**Fig. 5c, Supplementary Fig. S8**). Overall, enrichment analyses in

SOD1-G93A (ALS) mice highlight immune and neuronal-related modules as key disrupted

pathways in the spinal cord and implicates the dysregulation of SC.M2, a module enriched with

ALS genetic risk, and SC.M6, a module enriched with ALS genetic modifiers, in later stages of

disease progression (**Fig. 5e**).

Discussion

To help elucidate the biological mechanisms underlying ALS pathophysiology in an unbiased

genome-wide manner, we generated co-expression networks using RNA-sequencing of control

human cervical spinal cord samples from the GTEx Consortium. We identified 13 total co-

expression modules each representing distinct aspects of biology in the control spinal cord

including inflammation, neurotransmission, RNA processing, and cellular metabolism, as well as

major cell classes. We find that ALS genetic risk factors are enriched in SC.M4, a co-expression

module representing genes involved in RNA processing and epigenetics and SC.M2, which is

enriched in oligodendrocyte markers and genes involved in intracellular transport, protein

modification, and cellular catabolic processes.

It is notable that genes within both SC.M2 and SC.M4 have previously been shown to cause

rare Mendelian forms of ALS or have been strongly linked with risk. For example, SC.M4, which

is enriched in RNA processing, harbors RNA binding proteins mutated in ALS including TDP-43

and FUS/TLS, which are abnormally aggregated and mis-localized in ALS affected neurons,

leading to the loss of normal RNA binding function with incitement of neurotoxicity[42]. One of the

top hubs of the SC.M4 module is PTPN23, a non-receptor-type tyrosine phosphatase that is a

key regulator of the survival motor neuron (SMN) complex, which helps assemble small

ribonucleoproteins particles (snRNPs) and mediates pre-mRNA processing in motor neurons[43].

Furthermore, two other top hubs in SC.M2, KPNA3 and TMED2, are nucleocytoplasmic

transporters that act as modifiers of dipeptide repeat (DPR) toxicity in cases of C9orf72

mutations[27]. NCOA4, another top hub in SC.M2, is a key mediator of ferritinophagy and has

been implicated in neurodegeneration[44,45]. Taken together, the presence of ALS-linked risk

genes as hubs in SC.M2 and SC.M4 further underscores the importance of aberrant RNA

processing[42] and intracellular transport[46] as processes likely driving ALS pathophysiology.

We also find enrichment of genetic modifiers of ALS in SC.M6, indicating the importance of

ribosomal proteins and elongation factors (**Fig. 4B**) in mitigating ALS pathophysiology. These

factors have been suggested to act by reducing translation of toxic dipeptide repeat (DPRs)

expressed off of the GGGGCC hexanucleotide repeat expansion found in C9orf72[26]. We found

SC.M6 to be up-regulated with disease status in the SOD1-G93A mouse and dissected human

motor neurons primed for neurodegeneration (**Fig. 5b, d**), datasets with predominant glial

infiltration. Interestingly, we found a striking down-regulation of SC.M6 in iPSC-derived motor

neurons from ALS8 patients (**Fig. 5c**), a dataset without contaminating glial signal. This

suggests that role of this co-expression module is complex and cell-type dependent in ALS.

Across three of four ALS tissue and model datasets analyzed, we noted the presence of a

strong pro-immune phenotype (**Fig. 5**). While immune dysregulation is a key molecular feature

of the spinal cord by mid- to late-stage ALS, it remains unclear whether this disease signature is

primary or secondary[47,48]. Our analyses focused on ALS genetic risk factors, which serve as

causal anchors to identify, in an unbiased manner, specific modules that implicate processes

involved at the early stages of ALS pathophysiology. Furthermore, immune dysregulation is a

common feature in post-mortem tissue from other neurodegenerative and neuropsychiatric

disorders[8,49,50]; however, the enrichment of genetic risk within upregulated immune genes is

complex. For neuropsychiatric disorders such as autism and schizophrenia, the strongest

genetic enrichments are within neuronal genes[49]. In contrast, the strongest genetic risk

enrichments for Alzheimer's disease are within microglia[51]. For ALS, a recent assessment of

genetic risk within different cell types implicates oligodendrocytes[51] which is consistent with our

finding of risk enrichment within the oligodendrocyte-enriched module, SC.M2 (**Fig. 3b**).

A major question still remains concerning the specificity of MN degeneration in ALS. Our results

support the hypothesis that ALS is a non-cell-autonomous disease[52], as evidenced by SC.M1,

SC.M2, SC.M3, SC.12, and SC.M13 being enriched with astrocyte, oligodendrocyte, endothelia,

and microglia markers (**Supplementary Fig. S4**). Oligodendrocytes have critical roles through

the metabolic support of neurons and neurotransmission in the central nervous system[53]. A

previous study has shown that oligodendrocytes harboring familial and sporadic, but not

C9orf72 variants of ALS, induce motor neuron death via a SOD1-dependent mechanism, but

can be rescued via lactate supplementation[54]. These previous observations corroborate our

findings that highlight the potential deleterious effect of dysregulation in the intracellular

transport and autophagy system on oligodendrocyte - MN dynamics in ALS. Furthermore,

neuro-immune responses mediated by astrocytes and microglia also play a critical role in ALS

pathogenesis[55,56] and disruptions of blood spinal cord barrier and damage to the endothelial

cells have been observed in electron micrographs from ALS[57]. Overall, our analyses support the

notion that ALS is multifactorial disorder, implicating a number of cell-types and mitigating

mechanisms, acting at different stages of disease. Invariably, understanding of cell-type specific

mechanisms governing MN degeneration will involve future interrogation of the transcriptome

and epigenome from ALS spinal cord at single-cell resolutions. Additionally, emerging RNA-

sequencing datasets, including from the TargetALS foundation (http://www.targetals.org),

contain larger sample sizes of ALS and control spinal cord that will help elucidate the cellular

dynamics underlying ALS pathophysiology.

Methods

Initial processing of RNA-sequencing datasets

The latest version of GTEx data (version 7) was obtained from the dbGaP accession number

phs000424.v7.p2 on 10/10/2017 and pre-processed as previously described[18]. Genes in the

GTEx dataset containing fewer than 10 samples with a TPM > 1 were excluded from

downstream analysis on the basis of unreliable quantification for very lowly expressed genes.

We removed 4 outlier samples in the GTEx dataset that had sample-sample connectivity Z-

scores of > 2 as previously described[58]. Our starting GTEx dataset consisted of 62 samples and

14577 genes.

RNA-sequencing data[13] from postmortem cervical spinal cord (Brohawn et al.) was downloaded

using the SRP accession number SRP064478 on 5/24/2019. RNA-sequencing data[22] from laser

capture microdissected (LCM) MNs primed for degeneration (Krach et al.) was downloaded

using the SRP accession number SRP067645 on 6/27/2019. RNA-sequencing data[23] from

iPSC-derived motor neurons from ALS8 patients (Oliveira et al.) was downloaded using the

SRP accession number SRP223674 on 10/18/2020. Reads were aligned to the reference

genome GRCh37 using STAR (version 2.5.2b) and quantified using RSEM (version 1.3.0) with

Gencode v19 annotations. Genes in the Brohawn et al., Krach et al., and Oliveira et al. datasets

with zero variance in expression were excluded from further analysis. All expression data was

quantified in TPM (transcript per million reads) and TPM values +1 were $\log_2$ transformed to

stabilize their variance. The starting Brohawn et al., Krach et al., and Oliveira et al. datasets

consisted of 33050, 37138, and 14171 genes respectively. Krach et al. metadata tables were

transcribed from the manuscript and missing values for PMI (postmortem interval) were imputed

as the mean of all available PMI values.

Identification and regression of biological and technical confounders from gene expression

datasets

Gene expression values were subjected to dimensionality reduction using principal component

analysis (PCA) with data centering and scaling. The top four components for the GTEx dataset,

which collectively captures 70.6% of the total variance, were assigned to be expression principal

components (PCs) 1-4. The top five components for the Brohawn et al., Krach et al., and

Oliveira et al. dataset, which collectively capture 55.9%, 42.3%, and 56.3% of the total variance

respectively, were assigned to be expression principal components (PCs) 1-5. Sequencing Q/C

metrics were calculated using PicardTools (version 2.5.0) and dimensionally reduced using PCA

with centering and scaling. Sequencing Q/C metrics were aggregated manually for Brohawn et

al. and Krach et al. datasets and aggregated using MultiQC[59] for the Oliveira et al. dataset

(**Supplementary Data 2-4**). The top five components of sequencing metrics for the GTEx,

Brohawn et al., Krach et al., and Oliveira et al. datasets, which collectively capture 75.1%, 91%,

90.4%, and 89.0% of the total variance respectively, were selected and assigned to be

sequencing principal components (seqPCs) 1-5.

Numerical covariates of technical or biological importance were correlated with the expression

PCs using Spearman's correlation. Binary covariates were numerically encoded and multi-level

categorical covariates of technical or biological importance were binarized using dummy

variables. The model fit (R^2) was used to assess correlations. We identified technical or

biological covariates with a meaningful influence on gene expression if they were correlated with

an R > 0.3 to any of the top expression PCs.

For the GTEx expression dataset, a linear model was fit for each gene as follows: GTEX.Expr ~

seqPC1 + seqPC2 + seqPC3 + seqPC4 + seqPC5 + RIN + DTHHRDY + AGE + SMCENTER +

TRISCHD and the effect of all covariates were regressed out. RIN indicates RNA integrity

number, DTHHRDY represents the Hardy Scale, SMCENTER represents the center for

sampling, and TRISCHD represents ischemic time. Ethnicity was not included in the linear

model because it was collinear with SMCENTER (**Supplementary Fig. S1c**).

For the Brohawn et al. dataset, a linear model was fit for each gene as follows: Expr ~ seqPC1

+ seqPC2 + seqPC3 + seqPC4 +seqPC5 + Ethnicity + Sex + Age + Disease and the effect of all

covariates except disease were regressed out.

For the Krach et al. dataset, a linear model was fit for each gene as follows: Expr ~ Diagnosis +

Age + Gender + RIN + PMI + seqPC1 + seqPC2 + seqPC3 + seqPC4 + seqPC5 and the effect

of all covariates except diagnosis were regressed out.

For the Oliveira et al. dataset, a linear model was fit for each gene as follows: Expr ~ Disease +

Age + Sex + Tissue + seqPC1 + seqPC2 + seqPC3 + seqPC4 + seqPC5 and the effect of all

covariates except disease were regressed out.

Weighted Gene Co-expression Network Analysis (WGCNA)

We used WGCNA[4,58] to generate a gene co-expression network using the GTEx dataset. We

chose a soft power of 10 using the function "pickSoftThreshold" in the WGCNA R package to

achieve optimal scale free topology. The Topological Overlap Matrix (TOM) was generated

using the function "TOMsimilarity" in the WGCNA R package. A hierarchical gene clustering

dendrogram was constructed using the TOM-based dissimilarity and module selection was

tested using multiple permutations of the tree cutting parameters: minimum module size, cut

height, and deep split. Minimum module size (mms) was assessed at 50, 100, and 150. Module

merging (dcor) was assessed at 0.1, 0.2, and 0.25. Deep split (DS) was assessed at 2 and 4.

By visual inspection, we utilized the following parameters: minimum module size = 100, module

merging = 0.1, and deep split = 2 to generate the network. We assigned unique module

identifiers (SC.M1-13) for each module color except the grey module which represents genes

that were not co-expressed.

Pairwise Spearman's correlations were calculated using 1000 randomly sampled genes within

each module and a matched background set. If a module contained less than 1000 genes, all

genes in the module were sampled.

Enrichment of gene sets in co-expression modules

We analyzed a collection of gene sets generated by previous genetic studies of ALS. For large

effect size ALS risk genes, we obtained 50 ALS literature curated ALS risk genes from two

studies[19,20]. For the determination of cell-type specificity, we obtained the top 50 differentially

expressed genes per cell-type by fold change from single-nuclei RNA sequencing of postnatal

day 2 mouse spinal cord[24]. For the human ALS spinal cord differentially expressed gene (DEG)

enrichments, we obtained 265 DEGs (172 up- and 93 down-regulated) at FDR < 0.10 using the

union of DESeq2 and EdgeR analyses[13]. For the dissected motor neurons primed for

neurodegeneration, we obtained 1600 DEGs (1290 up- and 310 down-regulated) at FDR < 0.05

from the report[22]. For the iPSC-derived motor neurons from ALS patients, we chose to examine

the severe ALS vs. control comparison and filtered gene biotype for protein-coding genes,

obtaining 653 DEGs (364 up- and 289 down-regulated) at FDR < 0.01 from the report[23]. For

DEGs from SOD1-G93A mice, we obtained genes with a Bayes Factor > 3 (corresponding to

FDR = 0.1) identified through a spatial transcriptomics study[12].

Logistic regression was performed to calculate an odds ratio and p-value for enrichment of each

co-expression module with a test gene set. The p-values were FDR-corrected for the number of

modules analyzed.

Enrichment of common ALS genetic risk variants in co-expression modules

Stratified LD score regression[29] was performed to evaluate the enrichment of common risk

variants from GWAS studies of ALS[21] (N=80,610), Inflammatory Bowel Disorder (IBD)[30]

(N=86,640), Autism Spectrum Disorder (ASD)[32,33] (N=15,954), Alzheimer's Disease (AD)[34]

(N=455,258), Frontotemporal Dementia (FTD)[35] (N=12,928), and Progressive Supranuclear

Palsy (PSP)[36] (N=12,308) within each co-expression module. We used the full baseline model of

53 functional categories per calculation of partitioned heritability (1000 Genomes Phase 3). We

defined genomic regions of each co-expression module as their constituent gene bodies plus a

flanking region of +/- 10 KB.

Gene Ontology Enrichments

Gene Ontology enrichment analysis was performed using the gProfiler package[60] with

hierarchical filtering set to "moderate", domain size set to "annotated", src filter set to "GO:BP"

and "GO:MF", electronic annotations (IEAs) included, maximum p-value set to 0.05, correction

method set to "FDR", and a maximum set size of 1000. To reflect the weight of each gene within

a co-expression module, genes were ordered by their module membership (kME) during the

enrichment calculations.

Network visualization of module hub genes

500 For each module, an undirected and weighted network amongst the top 30 module hub genes

501 (by kME) was visualized using the igraph package in R[61].

Protein-protein interaction (PPI) networks

PPI networks were produced using the Disease Association Protein-Protein Evaluator

(DAPPLE)[62]. Two hundred and thirty eight of the top 300 SC.M4 hub genes, 238 of the top 300

SC.M2 hub genes, 84 of top 100 SC.M6 hub genes as ranked by Module Membership (kME)

were found in the DAPPLE database and seeded into the shown PPI network (**Fig. 3e, 3f, 4b**).

All networks were produced from 1,000 permutations (within-degree node-label permutation).

SC.M6, SC.M4, and SC.M2 contain the following network properties respectively: direct edges

count (p = 0.001, p =0.001, p = 0.001), seed direct degrees mean (p = 0.001, p = 0.001, p =

0.018), seed indirect degrees mean (p = 0.001, p = 0.006, p = 0.006), and the CI degrees mean

(p = 0.001, p = 0.132, p = 0.332).

Module preservation analyses

Module preservation for co-expression modules in the Brohawn et al. and Krach et al. datasets

were performed as previously described using the modulePreservation function in the WGCNA

R package[58]. We performed 200 permutation tests using a randomSeed of 1.

Correlations of module eigengenes to biological and technical covariates

To evaluate the correlations of module eigengenes to biological and technical covariates in the

Brohawn et al. and Krach et al. datasets, a Student p-value was calculated using the function

"corPvalueStudent" in the WGCNA R package. P-values were FDR corrected across the

modules and plotted using the function "labeledHeatmap" in the WGCNA R package. To

account for multiple biological replicates per patient in the Oliveira et al. dataset, a mixed effect

linear model was fitted for each module eigengene and each of the following covariates:

Disease, Age, Sex, Tissue, seqPC1, seqPC2, seqPC3, seqPC4, and seqPC5. The covariate

"Isolate", which uniquely identifies each individual, was used as the random effect.

Data Availability

Module assignments are provided as Supplementary Data 1 and the full range of values

underlying the barplots and heatmaps in Figures 3a-b, 4a, 5a-c, and Supplementary Figures S4,

S5d, S6d, S7d, S8, S9 are provided as a Source Data File.

Code Availability

Underlying R code to run WGCNA is available at [https://github.com/dhglab/ALS-Gene-Network-

Manuscript].

References

1. Ravits, J. *et al.* Deciphering amyotrophic lateral sclerosis: What phenotype, neuropathology and genetics are telling us about pathogenesis. *Amyotroph. Lateral Scler. Front. Degener.* **14**, 5–18 (2013).
2. Taylor, J. P., Brown, R. H. & Cleveland, D. W. Decoding ALS: from genes to mechanism. *Nature* **539**, 197–206 (2016).
3. Hardiman, O. *et al.* Amyotrophic lateral sclerosis. *Nat. Rev. Dis. Primer* **3**, 17071 (2017).
4. Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network analysis. *BMC Bioinformatics* **9**, 559 (2008).
5. Zhang, B. *et al.* Integrated Systems Approach Identifies Genetic Nodes and Networks in Late-Onset Alzheimer's Disease. *Cell* **153**, 707–720 (2013).
6. Mostafavi, S. *et al.* A molecular network of the aging human brain provides insights into the pathology and cognitive decline of Alzheimer's disease. *Nat. Neurosci.* **21**, 811–819 (2018).
7. Rexach, J., Swarup, V., Chang, T. & Geschwind, D. *Dementia risk genes engage gene networks poised to tune the immune response towards chronic inflammatory states*. http://biorxiv.org/lookup/doi/10.1101/597542 (2019) doi:10.1101/597542.
8. Swarup, V. *et al.* Identification of evolutionarily conserved gene networks mediating neurodegenerative dementia. *Nat. Med.* **25**, 152–164 (2019).
9. Johnson, E. C. B. *et al.* Large-scale proteomic analysis of Alzheimer's disease brain and cerebrospinal fluid reveals early changes in energy metabolism associated with microglia and astrocyte activation. *Nat. Med.* **26**, 769–780 (2020).
10. Swarup, V. *et al.* Identification of Conserved Proteomic Networks in Neurodegenerative Dementia. *Cell Rep.* **31**, 107807 (2020).
11. Prudencio, M. *et al.* Distinct brain transcriptome profiles in C9orf72-associated and sporadic ALS. *Nat. Neurosci.* **18**, 1175–1182 (2015).
12. Maniatis, S. *et al.* Spatiotemporal dynamics of molecular pathology in amyotrophic lateral sclerosis. *Science* **364**, 89–93 (2019).
13. Brohawn, D. G., O'Brien, L. C. & Bennett, J. P. RNAseq Analyses Identify Tumor Necrosis Factor-Mediated Inflammation as a Major Abnormality in ALS Spinal Cord. *PLOS ONE* **11**, e0160520 (2016).
14. D'Erchia, A. M. *et al.* Massive transcriptome sequencing of human spinal cord tissues provides new insights into motor neuron degeneration in ALS. *Sci. Rep.* **7**, 10046 (2017).
15. Schiffer, D., Cordera, S., Cavalla, P. & Migheli, A. Reactive astrogliosis of the spinal cord in amyotrophic lateral sclerosis. *J. Neurol. Sci.* **139**, 27–33 (1996).
16. Phatnani, H. P. *et al.* Intricate interplay between astrocytes and motor neurons in ALS. *Proc. Natl. Acad. Sci.* **110**, E756–E765 (2013).
17. Barham, C. *et al.* RNA-Seq Analysis of Spinal Cord Tissues from hPFN1G118V Transgenic Mouse Model of ALS at Pre-symptomatic and End-Stages of Disease. *Sci. Rep.* **8**, 13737 (2018).
18. GTEx Consortium. Genetic effects on gene expression across human tissues. *Nature* **550**, 204–213 (2017).
19. Eykens, C. & Robberecht, W. The genetic basis of amyotrophic lateral sclerosis: recent breakthroughs. *Advances in Genomics and Genetics* https://www.dovepress.com/the-genetic-basis-of-amyotrophic-lateral-sclerosis-recent-breakthrough-peer-reviewed-fulltext-article-AGG (2015) doi:10.2147/AGG.S57397.
20. Chia, R., Chiò, A. & Traynor, B. J. Novel genes associated with amyotrophic lateral sclerosis: diagnostic and clinical implications. *Lancet Neurol.* **17**, 94–102 (2018).
21. Nicolas, A. *et al.* Genome-wide Analyses Identify KIF5A as a Novel ALS Gene. *Neuron* **97**, 1268-1283.e6 (2018).

22. Krach, F. *et al.* Transcriptome–pathology correlation identifies interplay between TDP-43
and the expression of its kinase CK1E in sporadic ALS. *Acta Neuropathol. (Berl.)* **136**,
405–423 (2018).
23. Oliveira, D. *et al.* Different gene expression profiles in iPSC-derived motor neurons from
ALS8 patients with variable clinical courses suggest mitigating pathways for
neurodegeneration. *Hum. Mol. Genet.* **29**, 1465–1475 (2020).
24. Rosenberg, A. B. *et al.* Single-cell profiling of the developing mouse brain and spinal
cord with split-pool barcoding. *Science* **360**, 176–182 (2018).
25. Parikshak, N. N., Gandal, M. J. & Geschwind, D. H. Systems biology and gene networks
in neurodevelopmental and neurodegenerative disorders. *Nat. Rev. Genet.* **16**, 441–458
(2015).
26. Chai, N. & Gitler, A. D. Yeast screen for modifiers of C9orf72 poly(glycine-arginine)
dipeptide repeat toxicity. *FEMS Yeast Res.* **18**, (2018).
27. Jovičić, A. *et al.* Modifiers of C9orf72 dipeptide repeat toxicity connect nucleocytoplasmic
transport defects to FTD/ALS. *Nat. Neurosci.* **18**, 1226–1229 (2015).
28. Sun, Z. *et al.* Molecular Determinants and Genetic Modifiers of Aggregation and Toxicity
for the ALS Disease Protein FUS/TLS. *PLoS Biol.* **9**, e1000614 (2011).
29. Finucane, H. K. *et al.* Partitioning heritability by functional annotation using genome-wide
association summary statistics. *Nat. Genet.* **47**, 1228–1235 (2015).
30. Liu, J. Z. *et al.* Association analyses identify 38 susceptibility loci for inflammatory bowel
disease and highlight shared genetic risk across populations. *Nat. Genet.* **47**, 979–986
(2015).
31. Xavier, R. J. & Podolsky, D. K. Unravelling the pathogenesis of inflammatory bowel
disease. *Nature* **448**, 427–434 (2007).
32. Grove, J. *et al. Common risk variants identified in autism spectrum disorder*.
http://biorxiv.org/lookup/doi/10.1101/224774 (2017) doi:10.1101/224774.
33. The Autism Spectrum Disorders Working Group of The Psychiatric Genomics
Consortium. Meta-analysis of GWAS of over 16,000 individuals with autism spectrum
disorder highlights a novel locus at 10q24.32 and a significant overlap with
schizophrenia. *Mol. Autism* **8**, 21 (2017).
34. Jansen, I. E. *et al.* Genome-wide meta-analysis identifies new loci and functional
pathways influencing Alzheimer's disease risk. *Nat. Genet.* **51**, 404–413 (2019).
35. Ferrari, R. *et al.* Frontotemporal dementia and its subtypes: a genome-wide association
study. *Lancet Neurol.* **13**, 686–699 (2014).
36. Chen, J. A. *et al.* Joint genome-wide association study of progressive supranuclear palsy
identifies novel susceptibility loci and genetic correlation to neurodegenerative diseases.
Mol. Neurodegener. **13**, 41 (2018).
37. Boeynaems, S. *et al.* Phase Separation of C9orf72 Dipeptide Repeats Perturbs Stress
Granule Dynamics. *Mol. Cell* **65**, 1044-1055.e5 (2017).
38. Ozdilek, B. A. *et al.* Intrinsically disordered RGG/RG domains mediate degenerate
specificity in RNA binding. *Nucleic Acids Res.* **45**, 7984–7996 (2017).
39. Cooper-Knock, J. *et al.* A data-driven approach links microglia to pathology and
prognosis in amyotrophic lateral sclerosis. *Acta Neuropathol. Commun.* **5**, 23 (2017).
40. Ripps, M. E., Huntley, G. W., Hof, P. R., Morrison, J. H. & Gordon, J. W. Transgenic
mice expressing an altered murine superoxide dismutase gene provide an animal model
of amyotrophic lateral sclerosis. *Proc. Natl. Acad. Sci.* **92**, 689–693 (1995).
41. Gurney, M. *et al.* Motor neuron degeneration in mice that express a human Cu,Zn
superoxide dismutase mutation. *Science* **264**, 1772 (1994).
42. Polymenidou, M. *et al.* Misregulated RNA processing in amyotrophic lateral sclerosis.
Brain Res. **1462**, 3–15 (2012).

43. Husedzinovic, A. *et al.* The catalytically inactive tyrosine phosphatase HD-PTP/PTPN23
is a novel regulator of SMN complex localization. *Mol. Biol. Cell* **26**, 161–171 (2015).
44. Mancias, J. D., Wang, X., Gygi, S. P., Harper, J. W. & Kimmelman, A. C. Quantitative
proteomics identifies NCOA4 as the cargo receptor mediating ferritinophagy. *Nature*
509, 105–109 (2014).
45. Quiles del Rey, M. & Mancias, J. D. NCOA4-Mediated Ferritinophagy: A Potential Link to
Neurodegeneration. *Front. Neurosci.* **13**, 238 (2019).
46. Kim, H. J. & Taylor, J. P. Lost in Transportation: Nucleocytoplasmic Transport Defects in
ALS and Other Neurodegenerative Diseases. *Neuron* **96**, 285–297 (2017).
47. McCombe, P. A. & Henderson, R. D. The Role of Immune and Inflammatory
Mechanisms in ALS. *Curr. Mol. Med.* **11**, 9 (2011).
48. McCombe, P. A. The Peripheral Immune System and Amyotrophic Lateral Sclerosis.
Front. Neurol. **11**, 12 (2020).
49. Gandal, M. J. *et al.* Shared molecular neuropathology across major psychiatric disorders
parallels polygenic overlap. *Science* 6 (2018).
50. Morabito, S., Miyoshi, E., Michael, N. & Swarup, V. Integrative genomics approach
identifies conserved transcriptomic networks in Alzheimer's disease. *Hum. Mol. Genet.*
29, 2899–2919 (2020).
51. Nott, A. *et al.* Brain cell type–specific enhancer–promoter interactome maps and disease
risk association. *Science* **366**, 1134–1139 (2019).
52. Ilieva, H., Polymenidou, M. & Cleveland, D. W. Non–cell autonomous toxicity in
neurodegenerative disorders: ALS and beyond. *J. Cell Biol.* **187**, 761–772 (2009).
53. Simons, M. & Nave, K.-A. Oligodendrocytes: Myelination and Axonal Support. *Cold
Spring Harb. Perspect. Biol.* **8**, a020479 (2016).
54. Ferraiuolo, L. *et al.* Oligodendrocytes contribute to motor neuron death in ALS via
SOD1-dependent mechanism. *Proc. Natl. Acad. Sci.* **113**, E6496–E6505 (2016).
55. Yamanaka, K. & Komine, O. The multi-dimensional roles of astrocytes in ALS. *Neurosci.
Res.* **126**, 31–38 (2018).
56. Geloso, M. C. & D'Ambrosi, N. The Dual Role of Microglia in ALS: Mechanisms and
Therapeutic Approaches. *Front. Aging Neurosci.* **9**, 10 (2017).
57. Garbuzova-Davis, S. *et al.* Impaired blood–brain/spinal cord barrier in ALS patients.
Brain Res. **1469**, 114–128 (2012).
58. Horvath, S. *Weighted Network Analysis: Applications in Genomics and Systems Biology*.
(Springer-Verlag, 2011).
59. Ewels, P., Magnusson, M., Lundin, S. & Käller, M. MultiQC: summarize analysis results
for multiple tools and samples in a single report. *Bioinformatics* **32**, 3047–3048 (2016).
60. Reimand, J. *et al.* g:Profiler—a web server for functional interpretation of gene lists
(2016 update). *Nucleic Acids Res.* **44**, W83–W89 (2016).
61. Csárdi, G. & Nepusz, T. The igraph software package for complex network research. in
(2006).
62. Rossin, E. J. *et al.* Proteins Encoded in Genomic Regions Associated with Immune-
Mediated Disease Physically Interact and Suggest Underlying Biology. *PLoS Genet.* **7**,
e1001273 (2011).
63. Carbon, S. *et al.* AmiGO: online access to ontology and annotation data. *Bioinformatics*
25, 288–289 (2009).
683
684

Acknowledgements

We thank Christopher Hartl, William Pembroke, Timothy S. Chang, and other members of the

Geschwind Laboratory for their technical assistance and manuscript feedback. We also thank

Aaron D. Gitler and members of his laboratory for their stimulating discussions and assistance

with the genetic modifier screen data. Funding for this work was provided by the Boyer

Endowment Award, Helga K. & Walter Oppenheimer Endowment Award, and the MacDowell

Endowment Award through the UCLA Undergraduate Research Center - Sciences (to J.C.W),

NIMH 1F32MH114620 (to G.R), and NIMH 5R01MH109912 (to D.H.G).

Author Contributions

J.C.W, G.R, and D.H.G planned the analyses and wrote the manuscript. Analyses were

performed by J.C.W with supervision from G.R. and D.H.G.

Competing Financial Interests

The authors declare no competing financial interests.

1.) Generate spinal cord network and annotate co-expression modules

Initial Processing of GTEx Data

Cell-Type Identity via Markers from Single Cell Datasets [Rosenberg et al. 2018] ← Generation of Co-Expression Modules from Control Spinal Cord (WGCNA) → Gene Ontology

2.) Test for genetic drivers by risk allele, risk gene, and modifier screen enrichments

ALS Common Risk Variant Enrichment → SC.M2 & SC.M4

ALS Rare Variant Risk Gene Enrichment → SC.M4

Enrichment of Modifiers of C9orf72 and FUS Toxicity → SC.M6

3.) Identify modules with dysregulation in human ALS and model system data

Bulk-Tissue sALS [Brohawn et al. 2016] → SC.M3,12

LCM of Presymptomatic MN [Krach et al. 2018] → SC.M1, 3, 6, 7, 9, 12, 13

iPSC-derived MN from ALS8 Patients [Oliveira et al. 2020] → SC.M1, 6, 7, 9, 11

Spatial Transcriptomics in SOD1-G93A (ALS) Mouse [Maniatis et al. 2019] → SC.M1, 2, 3, 6, 7, 9, 11, 12, 13

Figure 1. Overview of spinal cord co-expression network analysis.

Flow diagram of analyses in this study. A co-expression network was generated using the GTEx

cervical spinal cord expression data and Weighted Gene Co-Expression Network Analysis

(WGCNA). Modules were annotated by their associated biological processes, hub genes, and

cell-type identities. Modules were then assessed for their enrichment of ALS genetic risk factors

and analyzed for differential expression in RNA-seq data from human ALS tissues and model

system studies[12,13,22,23].

Figure 2. Construction of co-expression network in control cervical spinal cord.

(A) Dendrogram of the topological overlap matrix for gene co-expression dissimilarity. Co-

expression module assignments are shown in the "Merged Colors" track (See Supplementary

Fig. S2 for optimization of tree-cutting parameters). Genes that failed to be clustered into a co-

expression module were grouped within the grey module.

(B) Thirteen co-expression modules were categorized into five broad molecular themes based

on Gene Ontology (GO) term enrichment and assigned a unique module identifier (SC.M1-13).

(C, D, E, F, G) Top 30 hub genes and 300 connections for SC.M3, SC.M9, SC.M6, SC.M12,

and SC.M5 respectively (left panels). Top 5 enriched GO terms for each of the respective

modules (right panels). Red lines mark an FDR corrected p-value threshold of 0.05.

A. ALS Rare Variant Risk Gene Enrichment

p=0.0124, FDR=0.174

Overlapping ALS Risk Genes in SC.M4

FUS	C21orf2
SPG11	EWSR1
SETX	SS18L1
TARDBP	ZNF512B
ATXN2	

B. ALS Common Variant Enrichment

-log₁₀(FDR): $-\log_{10}(FDR)$

Columns: ALS (N=80,610), IBD (N=86,640), ASD (N=15,954), AD (N=455,258), FTD (N=12,928), PSP (N=12,308)

SC.M2: 0.047 (ALS)
SC.M4: 0.041 (ALS)
SC.M12: 0.047 (ALS)

C. SC.M4

Bar chart categories (x-axis: $-\log_{10}(FDR)$):
- histone modification
- RNA processing
- peptidyl–lysine modification
- mRNA processing
- regulation of gene expression, epigenetic

D. SC.M2

Bar chart categories (x-axis: $-\log_{10}(FDR)$):
- cellular protein catabolic process
- protein modification by small protein conjugation or removal
- Golgi vesicle transport
- GDP binding
- macroautophagy

E. Key

Blue	histone modification
Orange	RNA processing
Red	peptidyl-lysine modification
Green	mRNA processing
Yellow	regulation of gene expression, epigenetic

p = 0.001

F. Key

Blue	cellular protein catabolic process
Orange	protein modification by small protein conjugation or removal
Red	golgi vesicle transport
Green	GDP binding
Yellow	macroautophagy

p = 0.018

Figure 3. Enrichment of ALS genetic risk factors in intracellular transport/autophagy module

SC.M2 and RNA processing/gene regulation module SC.M4.

(A) Enrichment of literature curated large-effect ALS risk genes[19,20] within co-expression

modules (left panel). Red line marks a baseline odds ratio of 1. SC.M4 is marked with an

asterisk as having a nominal enrichment p-value < 0.05. Literature-curated large-effect ALS risk

genes within the SC.M4 module (right panel).

(B) Partitioned heritability enrichments of common risk variants for ALS[21], IBD[30], ASD[32,33], AD[34],

FTD[35], and PSP[36] within co-expression modules. Text values of enrichments with FDR

corrected p-value < 0.05 are shown.

(C, D) Top 30 hub genes and 300 connections for SC.M4 and SC.M2 respectively (left panels).

Highlighted in red are genes that have been previously implicated in ALS pathophysiology[27,43–

45]. Top 5 enriched GO terms for each of the respective modules (right panels). Red lines mark

an FDR corrected p-value threshold of 0.05.

(E, F) Direct protein-protein interaction (PPI) network of the top 300 hub genes from modules

SC.M4 and SC.M2 respectively. Gene vertices are color coded according to their Gene

Ontology terms as shown[63]. P-value of the PPI network is derived from 1,000 permutations.

Highlighted in pink are genes that have been previously implicated in ALS pathophysiology[27,43–

45].

A. Genetic Modifier Enrichments

B.

p = 0.001

Key

Blue	SC.M6 Hub
Orange	C9orf72 Suppressor
Red	C9orf72 Enhancer
Green	FUS Enhancer
Yellow	FUS Suppressor

Figure 4. Modifiers of C9orf72 and FUS toxicity converge on top hubs of ribosomal-associated

module SC.M6.

(A) Enrichment of genetic modifiers of C9orf72 and FUS toxicity[26–28] within co-expression

modules. Odds ratios of enrichments with FDR corrected p-value < 0.05 are shown. FDR

corrected p-values themselves are shown within parentheses.

(B) Direct protein-protein interaction (PPI) network of the top 100 hub genes from module

SC.M6. Gene vertices are color coded as shown. P-value of the PPI network is derived from

1,000 permutations.

A. Bulk-Tissue sALS DEG Enrichment
B. LCM Presymptomatic MNs DEG Enrichment
C. ALS8 iPSC-derived MNs DEG Enrichment

D. Enrichment of DEGs from SOD1-G93A (ALS) Mouse

E.

Module	Number of Genes	Functional Annotations	Cell Type Enrichments	Genetic Enrichments	Tissue Dysregulation	Direction of Change	Stage of Change
SC.M1	522	Response to wounding, tube development, morphogenesis	Astrocyte	None	Laser microdissected presymptomatic human ALS MNs	Up-regulated	Presymptomatic
					iPSC-derived motor neurons from ALS8 patients	Up-regulated	N/A
					SOD1-G93A (ALS) mouse	Up-regulated	Onset - end stage
SC.M2	1610	Autophagy, intracellular transport, oligodendrocyte identity	Oligodendrocyte	Common ALS risk variants	SOD1-G93A (ALS) mouse	Down-regulated	Symptomatic
SC.M3	530	Cardiovascular development, morphogenesis, angiogenesis	Endothelial	None	Bulk Tissue sALS	Down-regulated	End stage
					Laser microdissected presymptomatic human ALS MNs	Up-regulated	Presymptomatic
					SOD1-G93A (ALS) mouse	Up-regulated	Symptomatic - end stage
SC.M4	1199	RNA processing, gene regulation	None	Rare and common ALS risk variants	None	None	None
SC.M6	168	Ribosomal-related	None	Genetic modifiers of C9orf72 and FUS cytotoxicity	Laser microdissected presymptomatic human ALS MNs	Up-regulated	Presymptomatic
					iPSC-derived motor neurons from ALS8 patients	Down-regulated	N/A
					SOD1-G93A (ALS) mouse	Up-regulated	End stage
SC.M12	157	Leukocyte activation, immune response	Microglia	None	Bulk Tissue sALS	Up-regulated	End stage
					Laser microdissected presymptomatic human ALS MNs	Up-regulated	Presymptomatic
					SOD1-G93A (ALS) mouse	Up-regulated	Onset - end stage

Figure 5. Validation of co-expression module dynamics in human ALS and model system

datasets.

(A) Enrichments of 112 up-regulated and 70 down-regulated DEGs from bulk-tissue RNA-

sequencing of postmortem ALS cervical spinal cord[13] within co-expression modules. Red line

marks a baseline odds ratio of 1. * (FDR corrected p-value < 0.05). ** (FDR corrected p-value <

0.01). *** (FDR corrected p-value < 0.001).

(B) Same as Fig. 5a except for using 1290 up-regulated and 310 down-regulated DEGs from

RNA-sequencing of laser capture microdissections of motor neurons primed for degeneration [22].

(C) Same as Fig. 5a except for using 364 up-regulated and 289 down-regulated DEGs from

RNA-sequencing of iPSC-derived motor neurons from ALS8 patients[23].

(D) Enrichment of DEGs within each disease stage of the SOD1-G93A mouse[12] within co-

expression modules. p30, p70, p100, and p120 represent postnatal age in days at time of

spatial transcriptomic sequencing. Odds ratios of enrichments with FDR corrected p-value <

0.05 are shown. * (FDR corrected p-value < 0.05). ** (FDR corrected p-value < 0.01). *** (FDR

corrected p-value < 0.001).

(E) Summary of the key findings from this study regarding modules relevant to ALS

pathophysiology.