The conserved ER-transmembrane protein TMEM39A coordinates with vesicular COPII to promote collagen secretion and prevent ER stress

Running title: The TMEM39A protein family is essential for collagen secretion

Zhe Zhang[1], Shuo Luo[1], Guilherme Oliveira Barbosa[1], Meirong Bai[1], Thomas B. Kornberg[1,2] and Dengke K. Ma[1, 3, *]

[1]Cardiovascular Research Institute, University of California San Francisco, San Francisco, CA 94158

[2]Department of Biochemistry and Biophysics, University of California San Francisco, San Francisco, CA 94158

[3]Department of Physiology, University of California San Francisco, San Francisco, CA 94158

*Correspondence: Dengke.Ma@ucsf.edu

1
2

Abstract

Dysregulation of collagen production and secretion contributes to aging and tissue

fibrosis of major organs. How premature collagen proteins in the endoplasmic

reticulum (ER) route in specialized vesicles for secretion remains to be fully elucidated.

Here, we report that TMEM39, an ER-localized transmembrane protein, regulates

production and secretory cargo trafficking of procollagen. We identify the *C. elegans*

orthologue TMEM-39 from an unbiased RNAi screen and show that deficiency of

tmem-39 leads to striking defects in cuticle collagen production and constitutively high

ER stress response. RNAi knockdown of the *tmem-39* orthologue in *Drosophila* causes

similar defects in collagen secretion in fat body cells. The cytosolic domain of the

human orthologue TMEM39A binds to Sec23A, a vesicle coat protein that drives

collagen secretion and vesicular trafficking. We propose that roles of TMEM39 in

collagen secretion and preventing ER stress are likely evolutionarily conserved.

Keywords

18 TMEM-39, Sec23A, Collagen, Fibrosis, COPII vesicles

Introduction

Collagen is the major molecular component of connective tissues, and the most abundant protein in animals (Shoulders and Raines, 2009). Collagen dysregulation causes many human disorders, including autoimmune diseases, brittle bone diseases (too little collagen), tissue fibrosis (too much collagen) and ageing (Ewald et al., 2015; Humphreys, 2018; McKleroy et al., 2013; Murtha et al., 2017; Timpane et al., 2016; Wynn and Ramalingam, 2012). The multi-step biosynthesis of mature collagen by the cell is a complex process and involves procollagen gene transcription and protein translation, posttranslational modification, assembly into trimers inside the endoplasmic reticulum (ER), vesicular secretion from ER, extracellular peptide cleavage and cross-linking into collagen fibers (Mienaltowski and Birk, 2014; Shoulders and Raines, 2009).

Specific mechanisms responsible for secretion of procollagen still remain poorly understood. In general, specialized intracellular vesicles defined by the coat protein complex II (COPII) transport most secreted proteins, including procollagen, from the ER to the Golgi apparatus (Barlowe and Helenius, 2016; Jensen and Schekman, 2011). Sec23, Sec24, Sec13 and Sec31 comprise key COPII coat proteins, while the transport protein particle (TRAPP) complex acts a key tethering factor for COPII vesicle in route to the Golgi (Barlowe, 2020; Barrowman et al., 2010; D'Arcangelo et al., 2013). Typical COPII vesicles are 60 to 80 nm in diameter, which is not sufficient for transporting procollagen trimers with up to 300 to 400 nm in length (Malhotra and Erlmann, 2015). In mammals, large-size COPII-coated vesicles may transport procollagen from the ER

to the Golgi apparatus. TANGO1, a transmembrane protein at the ER exit site, is a key

linker to mediate formation of specialized collagen-transporting vesicle and recruitment

of procollagen (Malhotra and Erlmann, 2015; Malhotra et al., 2015; Santos et al., 2015).

The N-terminal SH3-like domain of TANGO1 binds to the collagen chaperone HSP47

in the ER lumen, recruiting procollagens to the ER exit site (Ito and Nagata, 2019). Its

C-terminal proline-rich domain (PRD) servers as a COPII receptor by interacting with

the inner shell of Sec23/Sec24 (Wilson et al., 2011). The coil-coil domain of TANGO1

forms a stable complex with cTAE5 and SEC12, which is particularly enriched around

large COPII carriers for procollagen (Yuan et al., 2018). Through its membrane helices,

TANGO1 organizes ER exit sites by creating a lipid diffusion barrier and an export

conduit for collagen (Raote et al., 2020). While requirement of TANGO1 for secretion

may depend on specific collagen types, it remains unclear whether TANGO1's

functions are broadly conserved in all animals (Matsui et al., 2020; Zhang et al., 2020).

Caenorhabditis elegans produces over 180 collagen members that constitute the

cuticle and basement membranes, encodes conserved homologs of COPII/TRAPP

proteins, yet lacks apparent TANGO1 homologs (Chioran et al., 2017; Kramer, 2005;

Page and Johnstone, 2007; Teuscher et al., 2019). This indicates that evolutionarily

conserved and TANGO1-independent mechanisms may exist in *C. elegans* to regulate

procollagen secretion. From a genome-wide RNAi screen for genes affecting ER stress

response, we previously identified *tmem-131* that defines a broadly conserved family

of proteins important for procollagen assembly and secretion (Zhang et al., 2020).

Mutations in specific collagen genes, conserved COPII/TRAPP-encoding homologues,

and impairment of collagen biosynthetic pathway components are known to result in a

range of phenotypes including ER stress response, abnormal cuticle-associated

morphology (Blister and Dumpy), and early death or growth arrest (Page and

Johnstone, 2007). *tmem-131* mutants exhibit such phenotypes typical for genes

required for collagen secretion (Zhang et al., 2020), while many other genes of

unknown functions from our initial screen remain uncharacterized.

Here, we characterize another *C. elegans* gene *tmem-39* that encodes an ER

transmembrane protein and is essential for cuticle collagen production. The deficiency

of TMEM-39 protein in *C. elegans* impairs cuticle integrity and secretion of COL-19, an

adult-specific cuticle collagen protein (Thein et al., 2003). We show that the *Drosophila*

ortholog of *tmem-39, CG13016,* is also essential for collagen secretion. From yeast-

two-hybrid (Y2H) screen, we find that the cytoplasmic loop domain of human

TMEM39A binds to Sec23A, the inner shell component of the COPII coating complex.

We demonstrate that SEC-23 and other COPII proteins are also essential for collagen

secretion in *C. elegans*. Our findings suggest that TMEM39 coordinates with

TMEM131 and COPII transport vesicles in the ER, and its roles in collagen secretion

and preventing ER stress are likely evolutionarily conserved in multicellular animals.

Results

Genome-wide RNAi screen identifies *tmem-39* regulating ER stress response in

C. elegans

We identified *D1007.5*, the sole *tmem-39* homologue in *C. elegans*, from a genome-

wide RNAi screen for genes affecting the abundance of transgenic reporter *asp-*

*17*p::GFP, which is up-regulated by temperature stress and down-regulated by ER

stress (Zhang et al., 2020). RNAi against *tmem-39* fully suppressed the *asp-17*p::GFP

reporter expression (Fig. 1A). Ensembl gene tree analysis shows that TMEM39 family

proteins are evolutionarily conserved from *C. elegans* to humans (Fig. 1B). Multiple

human genetic studies have identified the human homologue *TMEM39A* as a

susceptibility locus for various autoimmune diseases (Tran et al., 2017). TMEM39A is

also overexpressed in glioblastoma cells and considered as a brain tumor-specific

marker (Park et al., 2017). A recent study reported that TMEM39A is an ER-localized

transmembrane protein that regulates autophagy (Miao et al., 2020). How TMEM-39

regulates ER stress response remains unknown.

In this work, we first confirmed that RNAi against *tmem-39* in *C. elegans* caused a fully

penetrant and strong up-regulation of *hsp-4*p::GFP in hypodermis (Fig. 1C). *hsp-*

*4*p::GFP is a well-established reporter for unfolded protein response (UPR) caused by

ER stress in *C. elegans* (Harding et al., 2003). Loss-of-function of IRE-1, an ER stress-

sensing protein, abolished *hsp-4*p::GFP induction in *tmem-39* RNAi treated animals

(Fig. 1D). To verify the *tmem-39* RNAi phenotype, we used CRISPR/Cas9 to generate

a *C. elegans* null allele *dma258* carrying a 2750 bp deletion of the entire coding

sequences (Fig. 1E, Table S1). *dma258* mutants exhibited abnormally elevated

transcriptional level of *hsp-4*p::GFP (Fig. 1F). Besides constitutively activated *hsp-*

*4*p::GFP transcription, we also noticed that TMEM-39 deficient animals by RNAi or

dma258 were small and dumpy. Together, these results suggest that TMEM-39 as an

ER-localized transmembrane protein is required for maintaining normal homeostasis

of ER-resident proteins.

Loss of *tmem-39* impairs cuticle collagen secretion in *C. elegans*

To identify potential protein clients in the ER regulated by TMEM-39, we examined 24

various translational reporters of ER-resident secreted and transmembrane proteins

(Fig. S1, Table S2) and found that *tmem-39* RNAi knock-down strongly reduced COL-

19::GFP reporter abundance (Fig. 2A). COL-19 is a *C. elegans* exoskeleton collagen

that is secreted by the underlying hypoderm and required for integral structure of the

cuticle (Page and Johnstone, 2007). The C-terminal GFP-tagged COL-19 reporter

enables highly robust and tractable visualization of the cuticle morphology and to

identify defects in the collagen production machinery (Thein et al., 2003).

Using confocal microscopy to characterize the structure of hypodermal cuticle, we

found that in the control RNAi animals, COL-19::GFP is enriched in the hypoderm,

constituting regular annular furrows and lateral alae of the cuticle (Fig. 2B). In the

tmem-39 RNAi animals, COL-19::GFP is apparently aggregated in the intracellular

region of hypoderm, and largely absent in the cuticle (Fig. 2C). We further analyzed

the abundance and composition of COL-19::GFP proteins by Western blot. Besides

deficiency in overall COL-19::GFP abundance, *tmem-39* RNAi markedly increased the

soluble "premature" monomeric procollagens, while decreased the insoluble fractions

of cross-linked multimer and "mature" monomer of COL-19::GFP (Fig. 2D).

To examine possible involvement of *tmem-39* in collagen gene transcription, we used

RNAi to knock-down *tmem-39* in animals with the *col-19*p::GFP transcriptional reporter

in which *GFP* expression is driven by the promoter of *col-19*. In contrast to the fully

penetrant decrease of overall COL-19::GFP protein abundance, the transcriptional

activity of the *col-19* promoter was not affected by *tmem-39* (Fig. 2C and 2E). We also

evaluated the mRNA level of *col-19* by quantitative reverse transcription polymerase

chain reaction (qRT-PCR) and found that the *dma258* mutant displayed a mild increase

of *col-19* mRNA level, likely caused by a regulatory feedback loop in transcriptional

regulation of collagen genes (Fig. 2F). The *dma258* mutant fully recapitulated the

tmem-39 RNAi phenotype in defective COL-19::GFP secretion (Fig. 2G).

There are two main collagen-enriched tissues in *C. elegans*, the cuticle (exoskeleton)

and basement membranes (Kramer, 2005). *tmem-39* RNAi had no effect on the

production of mCherry-tagged EMB-9 (Ihara et al., 2011), a Collagen IV α1 on

basement membranes (Fig. S1T and Table S2). We found that loss of *tmem-39*

specifically affected collagens in cuticle, as exemplified by COL-19:GFP, LON-3::GFP

and COL-101::GFP (Fig. S2). Electron microscopy (EM) analysis revealed striking

reduction of cuticle thickness in *dma258* mutants than in wild type (Fig. 2H). We also

noticed that TMEM-39 deficient animals were small in size and dumpy, more sensitive

to cuticle-disrupting osmotic stresses and developed more slowly. Taken together,

these results demonstrate essential roles of TMEM-39 in collagen secretion, proper

cuticle formation and preventing ER stress that is induced by premature collagen

accumulation in *C. elegans*.

Evolutionarily conserved roles of TMEM39 family proteins for collagen secretion

TMEM39 family proteins are evolutionarily conserved among multicellular animals, and

invertebrate model organisms have one ortholog each, named *D1007.5* and *CG13016,*

in *C. elegans* and *Drosophila*, respectively (Fig. 1B). We determined whether functions

of TMEM39 family proteins in collagen secretion are evolutionarily conserved in

Drosophila. We visualized collagen secretion in fat body cells of the *Lsp2> Col4a1:RFP*

transgenic fly (Liu et al., 2017; Pastor-Pareja and Xu, 2011), and generated transgenic

RNAi to knock-down *Drosophila CG13016*, the sole *TMEM39* ortholog (Fig. 3A). The

physiological function of *Drosophila* fat body cells is to secrete collagen to the insect

blood, hemolymph. Confocal microscopy analysis of COL4A1::RFP revealed that the

Collagen type IV alpha 1::RFP proteins were strikingly accumulated in fat body cells of

CG13016 knock-down flies but not in control (Fig. 3B). Collagen accumulation caused

by *CG13016* RNAi indicates the role of TMEM39 family proteins in collagen secretion

is evolutionarily conserved also in *Drosophila*.

Since the sequence and function of TMEM39 appear to be highly conserved, we next

characterized the localization and protein interactors of human TMEM39A. The

vertebrate TMEM39 family consists of two paralogues, TMEM39A and TMEM39B

(Park et al., 2017). The *TMEM39B* gene is only conserved in vertebrates, and is likely

produced by the duplication of an ancestral form of *TMEM39A* (Kumar et al., 2017).

Consistent with a recent study (Miao et al., 2020), our confocal imaging of Hela cells

transiently transfected with reporters of GFP::TMEM39A and mCherry-tagged ER

markers indicated that TMEM39A localized to the ER (Fig. 4A). Interestingly,

TMEM39B appears to interact with the SARS-CoV-2 ORF9C protein (Gordon et al.,

2020), which localizes to ER-derived vesicles in HeLa cells (Baruah et al., 2020).

Human TMEM39A cytoplasmic loop domain interacts with Sec23A

Predicted by the TOPCONS program, TMEM-39 contains putatively eight transmembrane segments and two large cytoplasmic loops (Fig. 4B). We further used the Y2H screen to search for human proteins that could interact with the conserved first loop domain (198-298 a.a.) and the second loop domain (337-420 a.a.) of TMEM39A (Fig. 4C). Among the prey cDNA clones identified from the Y2H, Sec23A was confirmed to interact with the second loop domain of TMEM39A (Fig. 4D).

The cDNA clone from the Y2H library encodes the C-terminal 583-765 a.a. of Sec23A, encompassing the Gelsolin repeat and C-terminal actin depolymerization factor-homology domain (Fig. 4C). Sec23A is a core component of the COPII vesicle coating complex, which forms SEC23-SEC24 heterodimers in the inner shell of the COPII coat to select specific cargo molecules (Bi et al., 2002; Khoriaty et al., 2018). Mutations in human Sec23A cause an autosomal recessive disease, named Cranio-lenticulo-sutural dysplasia (CLSD) (Khoriaty et al., 2018). The disease manifests with skeletal abnormalities, dysmorphic facial features and calvarial hypomineralization, features thought to result from defects in collagen secretion (Boyadjiev et al., 2011). Consistent with recent studies using the CoIP assay to demonstrate association between TMEM39A and Sec23A (Miao et al., 2020), we found that TMEM39A interacted with Sec23A but not Sec24D in Y2H assays (Fig. 4D). These results indicate that the TMEM39A cytoplasmic loop domain interacts with Sec23A, which forms an inner-shell heterodimer with Sec24 to select procollagen proteins destined for secretion.

We next examined the loss-of-function phenotype of *sec-23*. RNAi knock-down of *sec-*

23, the *C. elegans* homolog of *Sec23A*, strongly reduced COL-19::GFP secretion in

the cuticle and increased its aggregation in the intracellular region of hypoderm (Fig.

4E). RNAi of *sec-23* also led to strong *hsp-4*p::GFP induction indicating constitutively

activated ER stress response (Fig. 4F). RNAi against genes encoding other

components of COPII also recapitulated the COL-19::GFP defects and *hsp-4*p::GFP

induction phenotype (Fig. 4E, 4F, S3-4 and Table 1). These results show that the loop

region of TMEM39A specifically binds to the COPII inner-shell component Sec23A and

its *C. elegans* homolog *sec-23* is also essential for collagen production *in vivo*.

Importance of the conserved YR motif in the TMEM39A cytoplasmic loop domain

By amino acid sequence alignment, we identified two Tyrosine-Arginine (YR) residues

in the second cytoplasmic loop domain of TMEM39A that are highly evolutionarily

conserved among all examined species from invertebrates to vertebrates (Fig. 5A). We

hypothesize that the conserved YR motif is important for interaction between

TMEM39A and Sec23A. To confirm the prediction, we substituted the YR motif of

TMEM39A into Alanine-Alanine (AA). Using Y2H assays, we found that such

substitution in TMEM39A strongly attenuated its interaction with Sec23A (Fig. 5B).

To determine whether the YR motif of TMEM-39 is physiologically important *in vivo*,

we generated a *C. elegans* knock-in allele *dma312* carrying a precise replacement of

YR residues into AA (Fig. 5C). Ce-YR replacement mutants recapitulated the *tmem-*

39(dma258) or *tmem-39* RNAi phenotype and exhibited abnormally strong *hsp-*

*4*p::GFP induction but did not affect the mRNA abundance of *col-19* (Fig. 5D). As

mutations, including synonymous mutations, can alter the secondary structure of an

mRNA and affect its stability (Duan et al., 2003), we measured the overall mRNA

abundance of *tmem-39* by qRT-PCR. We found that the YR-AA mutant slightly

increased *tmem-39* mRNA levels, similarly as the null allele likely caused by a

transcriptional feedback regulation (Fig. 5D). Together, our findings support the

essential role of the YR motif in the cytoplasmic loop domain of TMEM39A for its

interaction with Sec23, and homologues of both proteins are required for collagen

secretion and preventing ER stress in *C. elegans* (Fig. 5E).

Discussion

Our study identifies an ER transmembrane protein TMEM-39 in *C. elegans* with

essential roles in collagen secretion. Such roles are likely evolutionarily conserved in

animals. We propose that the TMEM39 cytoplasmic loop domain binds to Sec23A as

the core component of COPII coating complex to facilitate procollagen transport from

ER to Golgi (Fig. 5E). Phenotypic similarities of losses of TMEM-39 and TMEM-131,

another ER transmembrane protein we recently identified (Zhang et al., 2020), suggest

that both proteins cooperate in collagen secretion by assembling premature collagen

and recruiting COPII/TRAPPIII complexes for sequential ER-to-Golgi cargo transport.

By yeast-two-hybrid assays, we found that the TMEM39A cytoplasmic loop domain can

interact with the Sec23A. RNAi knock-down of *sec-23* and most other COPII genes

recapitulated the *tmem-39* loss-of-function phenotypes in constitutively high ER stress

response, defective collagen secretion and sensitivity to osmolality stress in *C. elegans*

(Table 1). However, we also noticed that RNAi knock-down of many COPII related

genes, such as *sec-23, sec-24.1, npp-20, sar-1, sec-12, rab-5 and trpp-8* caused more

severe phenotypes than *tmem-39* RNAi, leading to lethality or developmental arrest

that prevent collagen phenotype analysis (Table 1). However, treatment with these

RNAi starting from L4-stage for animals transferred from normal conditions to RNAi led

to robust COL-19::GFP phenotype (Fig. 4E and Fig. S3). Shorter duration of RNAi

treatment may explain milder collagen defective phenotype for *sec-31*, *npp-20* and

sec-12 (Fig. S3E, H and I). Compared with most COPII-related genes, *tmem-39* null

mutants exhibit similar collagen secretion defects but are nonetheless viable,

supporting the notion that TMEM-39 acts with COPII in collagen secretion but may

have more specialized roles in facilitating secretion of specific client proteins including

collagen COL-19 and the PtdIns(4)P Phosphatase SAC1 (Miao et al., 2020).

Besides Sec23A, additional interactors, such as Dctn6, was also identified from Y2H

screens with the TMEM39A cytoplasmic domain as bait. We verified the interaction

between full-length Dctn6 (1-190 a.a.) and TMEM39A (337-420 a.a.) (Fig. 4D). DCTN6

was identified as a subunit of the dynactin protein complex (Eckley et al., 1999).

Interestingly, dynactin is an essential cofactor of cytoplasmic dynein motor to transport

a variety of cargos and organelles along the microtubule-based cytoskeleton (Reck-

Peterson et al., 2018; Urnavicius et al., 2015). In mammalian cells, ER-to-Golgi

transport proceeds by cargo assembly into COPII-coated ER export sites (ERES)

followed by vesicular/tubular transport along microtubule tracks toward the Golgi in a

dynein/dynactin-dependent manner (Watson et al., 2005). Sec23p directly interacts

with the dynactin complex (Watson et al., 2005), indicating that TMEM39A may also

participate in a Sec23/DCTN6 complex to facilitate COPII coat assembly and

subsequent dynein/dynactin-dependent transport. Test of this model and determination

of the underlying mechanism in relation to TMEM131's role in collagen secretion await

further investigations.

Mammalian genomes encode two TMEM39 family proteins, TMEM39A and TMEM39B.

TMEM39A is a susceptibility locus associated with various autoimmune diseases and

highly up-regulated in brain tumors (Park et al., 2017; Tran et al., 2017). TMEM39B

was recently found to interact with the SARS-CoV-2 ORF9C protein, which localizes

to ER-derived vesicles (Baruah et al., 2020; Gordon et al., 2020). It remains unclear

whether *TMEM39A* and *TMEM39B* exhibit functional redundancy in collagen secretion

or pathological processes in human diseases. With single *tmem39* orthologue for each,

model organisms *C. elegans* and *Drosophila* may continue to provide insights into

functions and mechanisms of action of this protein family. Future elucidation of novel

and evolutionarily conserved roles of mammalian TMEM39 proteins in physiological

and pathological processes will likely lead to therapeutic targets and strategies for

alleviating diseases associated with this protein family in humans.

Materials and Methods

Worm Strains

The Bristol N2 strain was used as the wild type strain, and genotypes of other strains used are as follow: *zcIs4* [*hsp-4*p::GFP] V, *ire-1(zc14)* II; *zcIs4* V, *dmaIs10* [*asp-17*p::GFP; *unc-54*p::mCherry] X and *kaIs12* [*col-19*::GFP], *tmem-39(dma258)* and *tmem-39(dma312)* I. Extrachromosomal array transgenic strains *dmaEx169* [*rpl-28*p::T19D2.1::mCherry; *unc-122*p::GFP], *dmaEx153* [*rpl-28*p::Y73E7A.8::mCherry; *unc-122*p::GFP], and *dmaEx152* [*rpl-28*p::F23H12.5::mCherry; *unc-122*p::GFP] were generated as extrachromosomal arrays as described (Mello et al., 1991).

The precise *tmem-39(dma258)* knock-out strain was generated by CRISPR/Cas9 methods (Zhao et al., 2014; Zhao et al., 2016), and *tmem-39(dma312)* YR→AA was generated by the amplified sgRNA method (Ward, 2015). Translational fluorescent reporters used by *tmem-39* RNAi knock-down to identify a phenotype include: *bcIs39* [*lim-7*p::*ced-1*::GFP+*lin-15*(+)], *caIs618* [*eff-1*p::*eff-1*::GFP], *dnSi4* [*gna-1*p::GFP + Cbr-*unc-119*(+)], *juEx1111* [*spon-1*::vGFP], *lrp-1(ku156)eqIs1* [*lrp-1*p::*lrp-1*::GFP] I; *rrf-3(pk1426)* II, *muIs49* [*egl-20*::GFP+*unc-22*(+)], *nIs590* [*fat-7*p::*fat-7*::GFP], *nuIs26* [*cat-1*::GFP], *osIs60* [*unc-54*p::*mig-23*::GFP; *unc-119*(+)], *osIs66* [*myo-3*p::eGFP::*wrk-1*], *sqIs11* [*lgg-1*p::mCherry::GFP::*lgg-1*+*rol-6*(+)], *osIs77* [*unc-54*p::RFP::SP12;*unc-119*(+)], *pwIs503* [*vha-6*p::*mans*::GFP+Cbr-*unc-119*(+)], *qyIs44* [*emb-9*p::*emb-9*::mCherry], *rhIs23* [GFP::*him-4*], *veIs13 [col-19::GFP + rol-6(+)] V; let-7(mn112) unc-3(e151) X; mgEx725 [lin-4::let-7 + ttx-3::RFP], vkEx1243* [*nhx-2*p::ubiquitin-V::mCherry+*myo-2*p::GFP], *vkEx1256* [*nhx-2*p::*cpl-1*::YFP], *vkEx1260* [*nhx-2*p::*cpl-*

297   *1*::YFP], *vkEx1879* [*nhx-2*p::*cpl-1*(W32A Y35A)::YFP] and *xnIs96* [*hmr-1*p::*hmr-*

298   *1*::GFP].

Worm maintenance

We used standard methods to grow *C. elegans* strains on *E. coli* at 20 °C, and animals

were maintained in standard nematode growth medium (NGM) plates (Brenner, 1974).

Worms were synchronized by bleaching the gravid adults, and RNAi feeding induced

knock-down was performed as previously described (Kamath and Ahringer, 2003). For

RNAi colonies that show lethality or larvae arrest phenotype, around 20-30 P0 L4

animals were transferred from normal NGM plates to RNAi plates, and grew for 2-3

days to observe the P0 phenotype.

Imaging

Digital automated epifluorescence microscopes (EVOS, Life Technologies) and SPE

confocal microscope (Leica) were used to obtain fluorescence images. Animals at the

same stage were randomly picked from the plate, and transferred to an 4% agar pad

with 10 mM sodium azide and 1 mM levamisole in M9 solution (31742-250MG, Sigma-

Aldrich) on a slide for imaging. Identical setting and conditions were used to compare

experimental groups with control.

Western blot analysis of proteins

Animals at the same stage from the control and experiment groups were picked (N>30)

into 20 μL Laemmli Sample Buffer and lysed directly for Western blot analysis. Protein

samples were run with 15% SDS-PAGE (Bio-Rad, 4561084), and then transferred to

the nitrocellulose membrane (Bio-Rad, 1620167). The membranes were detected by

antibodies against GFP (A02020, Abbkine), Tubulin (Sigma, T5168) and H3 (Abcam,

ab1791).

Quantitative RT-PCR

Worm total RNA was extracted by following the protocols of Quick-RNA MiniPrep kit

(Zymo Research, R1055). And cDNA was reverse transcribed by the reverse

transcriptase mix kit (BioTools, B24408). Using SYBR Green Supermix (Thermo Fisher

Scientific, FERK1081), the Real-time PCR was performed by on Roche LightCycler96

(Roche, 05815916001) system. Ct values of specific genes were normalized to the *C.*

elegans housekeeping gene *act-1* levels. Results are presented as fold changes to

respective references. Statistical significance was determined with t-test, using

GraphPad Prism 7. Primer sequences are listed in Supplementary Table 1B.

***Drosophila* experiments**

Fly strains included: UAS-Cg25C:RFP.2.1/CyO; Lsp2-Gal4/TM6B, and UAS-

CG13016_dsRNA (Vienna Drosophila Resource Center ID# 42509/GD). Lsp2-Gal4

specifically expresses in the fat body. Flies expressing Collagen:RFP in fat body were

crossed to either wild type or UAS-CG13016_dsRNA flies. Wandering stage third instar

larvae were picked out. Fat body was dissected and fixed in PFA 4%, stained with

DAPI, and mounted for imaging by confocal microscopy.

Yeast two hybrid assay

The cDNA coding sequences of the first and second cytoplasmic loop domain of

human TMEM39A were cloned into the pGBKT7 vector and screened against a

normalized universal human cDNA library (Clontech, 630481), following instructions of

the Matchmaker® Gold Yeast Two-Hybrid System (Clontech, 630489). Verification of

positive colonies was achieved by co-transforming wild-type or YR-mutant TMEM39A

loop domain (in pGBKT7 Vector) with genes of interest (in pGADT7 Vector) following

the instruction of YeastMaker™ Yeast Transformation System 2 (Clontech, 630439) as

well as plasmids from re-cloned cDNA.

Fluorescent imaging of Hela cells

Hela cells were seeded in 24-well plates with cover glass, each with three replicates

(Fisher Scientific, 22293232). Cells were transiently transfected with GFP-tagged

human *TMEM39A* full-length cDNA in the FUGW plasmid backbone, and the

endoplasmic reticulum localization marker mCherry-ER-3 (Addgene: 55041) for 2 days.

After 1xPBS washing for once, cells were treated by 4% formaldehyde solution for 10

mins. With 1xPBS washing for three times, cells were treated with 0.2% Triton X-100

in 1xPBS solution for 15 mins. Following 1xPBS washing for three times, the cover

slide with cell samples was sealed on the microscope slide with Fluoroshield Mounting

Medium with DAPI (Thermo Fisher Scientific, NC0200574) for imaging by confocal

microscopy.

Acknowledgements

We thank the *Caenorhabditis* Genetics Center for *C. elegans* strains. The work was supported by NIH grants R01GM117461, Pew Scholar Award, and Packard Fellowship in Science and Engineering (D.K.M). We thank Drs. Hong Zhang, Kamran Atabai and Matthew Shoulders for discussion and Dr. Jose Pastor-Pareja for the transgenic *Drosophila* line: w*; UAS-Cg25C.RFP.2.1. The authors declare that they have no competing interests and all participated experimental design, execution or data interpretation and analysis. Z.Z. and D.K.M wrote the paper.

Abbreviations

a.a., amino acid; CLSD, Cranio-lenticulo-sutural dysplasia; COPII, Coat protein complex II; C-terminus, carboxylic acid-terminus; EM, Electron microscopy; ER, endoplasmic reticulum; ERES, ER export sites; GWASs, Genome-wide association studies; NGM, nematode growth media; nt, nucleotide; N-terminus, amino-terminus; PRD, proline-rich domain; qRT-PCR, quantitative reverse transcription polymerase chain reaction TRAPP, transport protein particle; Y2H, yeast-two-hybrid;

Compliance with ethics guidelines

Zhe Zhang, Shuo Luo, Guilherme Oliveira Barbosa, Meirong Bai, Thomas B. Kornberg and Dengke K. Ma declare that they have no conflict of interests.

All institutional and national guidelines for the care and use of laboratory animals were followed.

References

Barlowe, C. (2020). Twenty-five years after coat protein complex II. Mol Biol Cell *31*, 3-6.

Barlowe, C., and Helenius, A. (2016). Cargo Capture and Bulk Flow in the Early Secretory Pathway. Annu Rev Cell Dev Biol *32*, 197-222.

Barrowman, J., Bhandari, D., Reinisch, K., and Ferro-Novick, S. (2010). TRAPP complexes in membrane traffic: convergence through a common Rab. Nat Rev Mol Cell Biol *11*, 759-763.

Baruah, C., Devi, P., and Sharma, D.K. (2020). Sequence analysis and structure prediction of SARS-CoV-2 accessory proteins 9b and ORF14: evolutionary analysis indicates close relatedness to bat coronavirus.

Bi, X., Corpina, R.A., and Goldberg, J. (2002). Structure of the Sec23/24-Sar1 pre-budding complex of the COPII vesicle coat. Nature *419*, 271-277.

Boyadjiev, S.A., Kim, S.D., Hata, A., Haldeman-Englert, C., Zackai, E.H., Naydenov, C., Hamamoto, S., Schekman, R.W., and Kim, J. (2011). Cranio-lenticulo-sutural dysplasia associated with defects in collagen secretion. Clin Genet *80*, 169-176.

Brenner, S. (1974). The genetics of Caenorhabditis elegans. Genetics *77*, 71-94.

Chioran, A., Duncan, S., Catalano, A., Brown, T.J., and Ringuette, M.J. (2017). Collagen IV trafficking: The inside-out and beyond story. Dev Biol *431*, 124-133.

D'Arcangelo, J.G., Stahmer, K.R., and Miller, E.A. (2013). Vesicle-mediated export from the ER: COPII coat function and regulation. Biochim Biophys Acta *1833*, 2464-2472.

Duan, J., Wainwright, M.S., Comeron, J.M., Saitou, N., Sanders, A.R., Gelernter, J., and Gejman, P.V. (2003). Synonymous mutations in the human dopamine receptor D2 (DRD2) affect mRNA stability and synthesis of the receptor. Hum Mol Genet *12*, 205-216.

Eckley, D.M., Gill, S.R., Melkonian, K.A., Bingham, J.B., Goodson, H.V., Heuser, J.E., and Schroer, T.A. (1999). Analysis of dynactin subcomplexes reveals a novel actin-related protein associated with the arp1 minifilament pointed end. J Cell Biol *147*, 307-320.

Ewald, C.Y., Landis, J.N., Porter Abate, J., Murphy, C.T., and Blackwell, T.K. (2015). Dauer-independent insulin/IGF-1-signalling implicates collagen remodelling in longevity. Nature *519*, 97-101.

Gordon, D.E., Jang, G.M., Bouhaddou, M., Xu, J., Obernier, K., White, K.M., O'Meara,
M.J., Rezelj, V.V., Guo, J.Z., Swaney, D.L., *et al.* (2020). A SARS-CoV-2 protein
interaction map reveals targets for drug repurposing. Nature *583*, 459-468.

Harding, H.P., Zhang, Y., Zeng, H., Novoa, I., Lu, P.D., Calfon, M., Sadri, N., Yun, C.,
Popko, B., Paules, R., *et al.* (2003). An integrated stress response regulates amino
acid metabolism and resistance to oxidative stress. Mol Cell *11*, 619-633.

Humphreys, B.D. (2018). Mechanisms of Renal Fibrosis. Annu Rev Physiol *80*, 309-
326.

Ihara, S., Hagedorn, E.J., Morrissey, M.A., Chi, Q., Motegi, F., Kramer, J.M., and
Sherwood, D.R. (2011). Basement membrane sliding and targeted adhesion remodels
tissue boundaries during uterine-vulval attachment in Caenorhabditis elegans. Nat Cell
Biol *13*, 641-651.

Ito, S., and Nagata, K. (2019). Roles of the endoplasmic reticulum-resident, collagen-
specific molecular chaperone Hsp47 in vertebrate cells and human disease. J Biol
Chem *294*, 2133-2141.

Jensen, D., and Schekman, R. (2011). COPII-mediated vesicle formation at a glance.
Journal of cell science *124*, 1-4.

Kamath, R.S., and Ahringer, J. (2003). Genome-wide RNAi screening in
Caenorhabditis elegans. Methods *30*, 313-321.

Khoriaty, R., Hesketh, G.G., Bernard, A., Weyand, A.C., Mellacheruvu, D., Zhu, G.,
Hoenerhoff, M.J., McGee, B., Everett, L., Adams, E.J., *et al.* (2018). Functions of the
COPII gene paralogs SEC23A and SEC23B are interchangeable in vivo. Proc Natl
Acad Sci U S A *115*, E7748-E7757.

Kramer, J.M. (2005). Basement membranes. WormBook, 1-15.

Kumar, S., Stecher, G., Suleski, M., and Hedges, S.B. (2017). TimeTree: A Resource
for Timelines, Timetrees, and Divergence Times. Mol Biol Evol *34*, 1812-1819.

Liu, M., Feng, Z., Ke, H., Liu, Y., Sun, T., Dai, J., Cui, W., and Pastor-Pareja, J.C. (2017).
Tango1 spatially organizes ER exit sites to control ER export. J Cell Biol *216*, 1035-
1049.

Malhotra, V., and Erlmann, P. (2015). The pathway of collagen secretion. Annu Rev
Cell Dev Biol *31*, 109-124.

Malhotra, V., Erlmann, P., and Nogueira, C. (2015). Procollagen export from the
endoplasmic reticulum. Biochem Soc Trans *43*, 104-107.

Matsui, Y., Hirata, Y., Wada, I., and Hosokawa, N. (2020). Visualization of Procollagen IV Reveals ER-to-Golgi Transport by ERGIC-independent Carriers. Cell Struct Funct *45*, 107-119.

McKleroy, W., Lee, T.H., and Atabai, K. (2013). Always cleave up your mess: targeting collagen degradation to treat tissue fibrosis. Am J Physiol Lung Cell Mol Physiol *304*, L709-721.

Mello, C.C., Kramer, J.M., Stinchcomb, D., and Ambros, V. (1991). Efficient gene transfer in C.elegans: extrachromosomal maintenance and integration of transforming sequences. EMBO J *10*, 3959-3970.

Miao, G., Zhang, Y., Chen, D., and Zhang, H. (2020). The ER-Localized Transmembrane Protein TMEM39A/SUSR2 Regulates Autophagy by Controlling the Trafficking of the PtdIns(4)P Phosphatase SAC1. Mol Cell *77*, 618-632 e615.

Mienaltowski, M.J., and Birk, D.E. (2014). Structure, physiology, and biochemistry of collagens. Adv Exp Med Biol *802*, 5-29.

Murtha, L.A., Schuliga, M.J., Mabotuwana, N.S., Hardy, S.A., Waters, D.W., Burgess, J.K., Knight, D.A., and Boyle, A.J. (2017). The Processes and Mechanisms of Cardiac and Pulmonary Fibrosis. Front Physiol *8*, 777.

Page, A.P., and Johnstone, I.L. (2007). The cuticle. WormBook, 1-15.

Park, J., Lee, H., Tran, Q., Mun, K., Kim, D., Hong, Y., Kwon, S.H., Brazil, D., Park, J., and Kim, S.H. (2017). Recognition of Transmembrane Protein 39A as a Tumor-Specific Marker in Brain Tumor. Toxicol Res *33*, 63-69.

Pastor-Pareja, J.C., and Xu, T. (2011). Shaping cells and organs in Drosophila by opposing roles of fat body-secreted Collagen IV and perlecan. Dev Cell *21*, 245-256.

Raote, I., Ernst, A.M., Campelo, F., Rothman, J.E., Pincet, F., and Malhotra, V. (2020). TANGO1 membrane helices create a lipid diffusion barrier at curved membranes. Elife *9*.

Reck-Peterson, S.L., Redwine, W.B., Vale, R.D., and Carter, A.P. (2018). The cytoplasmic dynein transport machinery and its many cargoes. Nat Rev Mol Cell Biol *19*, 382-398.

Santos, A.J., Raote, I., Scarpa, M., Brouwers, N., and Malhotra, V. (2015). TANGO1 recruits ERGIC membranes to the endoplasmic reticulum for procollagen export. Elife *4*.

Shoulders, M.D., and Raines, R.T. (2009). Collagen structure and stability. Annu Rev Biochem *78*, 929-958.

Teuscher, A.C., Jongsma, E., Davis, M.N., Statzer, C., Gebauer, J.M., Naba, A., and
Ewald, C.Y. (2019). The in-silico characterization of the Caenorhabditis elegans
matrisome and proposal of a novel collagen classification. Matrix Biology Plus *1*,
100001.

Thein, M.C., McCormack, G., Winter, A.D., Johnstone, I.L., Shoemaker, C.B., and
Page, A.P. (2003). Caenorhabditis elegans exoskeleton collagen COL-19: an adult-
specific marker for collagen modification and assembly, and the analysis of organismal
morphology. Dev Dyn *226*, 523-539.

Timpane, S., Brandling-Bennett, H., and Kristjansson, A.K. (2016). Autoimmune
collagen vascular diseases: Kids are not just little people. Clin Dermatol *34*, 678-689.

Tran, Q., Park, J., Lee, H., Hong, Y., Hong, S., Park, S., Park, J., and Kim, S.H. (2017).
TMEM39A and Human Diseases: A Brief Review. Toxicol Res *33*, 205-209.

Urnavicius, L., Zhang, K., Diamant, A.G., Motz, C., Schlager, M.A., Yu, M., Patel, N.A.,
Robinson, C.V., and Carter, A.P. (2015). The structure of the dynactin complex and its
interaction with dynein. Science *347*, 1441-1446.

Ward, J.D. (2015). Rapid and precise engineering of the Caenorhabditis elegans
genome with lethal mutation co-conversion and inactivation of NHEJ repair. Genetics
199, 363-377.

Watson, P., Forster, R., Palmer, K.J., Pepperkok, R., and Stephens, D.J. (2005).
Coupling of ER exit to microtubules through direct interaction of COPII with dynactin.
Nature cell biology *7*, 48-55.

Wilson, D.G., Phamluong, K., Li, L., Sun, M., Cao, T.C., Liu, P.S., Modrusan, Z.,
Sandoval, W.N., Rangell, L., Carano, R.A.*, et al.* (2011). Global defects in collagen
secretion in a Mia3/TANGO1 knockout mouse. J Cell Biol *193*, 935-951.

Wynn, T.A., and Ramalingam, T.R. (2012). Mechanisms of fibrosis: therapeutic
translation for fibrotic disease. Nat Med *18*, 1028-1040.

Yuan, L., Kenny, S.J., Hemmati, J., Xu, K., and Schekman, R. (2018). TANGO1 and
SEC12 are copackaged with procollagen I to facilitate the generation of large COPII
carriers. Proc Natl Acad Sci U S A *115*, E12255-E12264.

Zhang, Z., Bai, M., Barbosa, G.O., Chen, A., Wei, Y., Luo, S., Wang, X., Wang, B.,
Tsukui, T., Li, H.*, et al.* (2020). Broadly conserved roles of TMEM131 family proteins in
intracellular collagen assembly and secretory cargo trafficking. Sci Adv *6*, eaay7667.

Zhao, P., Zhang, Z., Ke, H., Yue, Y., and Xue, D. (2014). Oligonucleotide-based
targeted gene editing in C. elegans via the CRISPR/Cas9 system. Cell Res *24*, 247-
250.

Zhao, P., Zhang, Z., Lv, X., Zhao, X., Suehiro, Y., Jiang, Y., Wang, X., Mitani, S., Gong,

H., and Xue, D. (2016). One-step homozygosity in precise gene editing by an improved

CRISPR/Cas9 system. Cell Res *26*, 633-636.

Figure legends

Fig. 1. TMEM-39 regulates ER stress response in *C. elegans*. (A), Exemplar

fluorescence images for *asp-17*p::GFP with control and *tmem-39* RNAi. Scale bars: 20

μm. (B), Ensembl gene tree analysis for TMEM39 protein family and amino acid

sequence alignment among metazoan species (adapted from https://m.ensembl.org/).

(C-D), Exemplar fluorescence and bright-field images for the UPR reporter *hsp-*

*4*p::GFP with control and *tmem-39* RNAi in wild type (C) and *ire-1*(D) mutants. Scale

bars: 20 μm. (E), Schematic of *tmem-39* gene structure with the *dma258* deletion

generated by CRISPR-Cas9. Bold and underlined are sgRNA target sequence from

the *tmem-3*9 regions. The location of the deleted *tmem-39* is marked with "^",

generating a precise 2750 bp deletion. (F) qRT-PCR measurements of *hsp-4*p::GFP

mRNA levels in wild-type and *dma258* mutants. ***P < 0.001 (n ≥ 3 biological

replicates).

Fig. 2. TMEM-39 is essential for collagen secretion and cuticle formation in *C. elegans*. (A), Exemplar epifluorescence image of *col-19*::GFP with control and *tmem-39* RNAi. Three to four animals were shown to indicate representative reporter expression with around 50 animals observed. (B-C) Exemplar confocal fluorescence images of COL-19::GFP with indicated phenotypic penetrance of control RNAi (B) and *tmem-39* RNAi (C) in wild-type animals. Scale bars: 20 μm. (D), Exemplar Western blot analysis of COL-19::GFP proteins from total lysates of wild type animals with control and *tmem-39* RNAi. IB, immunoblotting. The arrow indicates soluble premature monomers; triangles indicate insoluble mature monomers and cross-linked COL-19::GFP. (E), Exemplar fluorescence images of *col-19* transcriptional reporter (*col-19* promoter-driven GFP) with indicated phenotypic penetrance of control RNAi and *tmem-39* RNAi in wild-type animals, indicating no significant difference in GFP expression. Scale bars: 20 μm. (F), qRT-PCR quantification of endogenous col-19 mRNA levels in wild-type and *dma258* mutants. ***$P < 0.001$ (n ≥ 3 biological replicates).

Fig. 3. Evolutionarily conserved roles of TMEM39 family proteins for collagen

secretion in *Drosophila*. (A), Schematic of generating fat body cell specific CG13016

knock-down strains in *Drosophila.* Lsp2-Gal4 specifically expresses in the fat body.

Wandering stage third instar larvae were picked out for imaging analysis. The

Drosophila images are created by BioRender.com. (B), Exemplar confocal images of

transgenic *Drosophila* fat body cells showing collagen COL4A1 secretion is normal

with control RNAi (left), and collagen accumulation with *tmem39/CG13016* RNAi. scale

bar, 100 μm.

Fig. 4. Human TMEM39A cytoplasmic loop domain interacts with Sec23A and

COPII components regulate collagen secretion in *C. elegans*. (A), Exemplar

confocal fluorescence images of HeLa cells co-transfected with GFP-tagged

TMEM39A and mCherry-tagged ER marker (ER3). Scale bars, 5 μm. (B) Schematic of

human TMEM39A transmembrane domain predicted by the TOPCONS program, with

cytosolic localization in red (two long cytoplasmic loop domains labeled with rectangles,

loop1 in green and loop2 in blue) and ER localization in blue. (C), Schematic of Y2H

screens identifying the human Sec23A C-terminal domain as a binder of the second

cytoplasmic loop domain of TMEM39A. (D), Y2H assays of yeast colony growth after

prey and bait vectors retransformation to verify the interaction of human Sec23A C-

termini (a.a. 583-765), Sec24D full length (a.a. 1-1032) and Dctn6 full length (a.a. 1-

190) with TMEM39A loop1 (a.a. 198-298) and loop2 (a.a. 337-420). (E-F) Exemplar

confocal fluorescence images of COL-19::GFP (E) and *hsp-4*p::GFP with indicated

phenotypic penetrance of wild-type with control RNAi and COPII components *sec-23*

and *trpp-3* RNAi. Scale bars: 20 μm.

Fig. 5. The conserved YR motif in cytoplasmic loop region is essential for the

function of TMEM39A protein. (A), Multiple sequence alignment of TMEM39A from

major representative animal species (by COBALT program), with conserved YR

residues indicated in the second cytoplasmic loop domains. (B), Y2H assays of yeast

colony growth after prey and bait vectors retransformation to verify the interaction

between human Sec23A C-termini (a.a. 583-765) with human wild type and YR mutant

TMEM39A cytoplasmic loop domain (a.a. 337-420). (C), Schematic of precise

replacement for YR motif coding sequence in *tmem-39* gene generated by CRISPR-

Cas9. (D), qRT-PCR measurements of endogenous *col-19*, *tmem-39* and *hsp-4*p::GFP

mRNA levels in wild-type and *tmem-39* YR mutants. ***$P < 0.001$ (n ≥ 3 biological

replicates). (E), Schematic model showing TMEM39A regulation of collagen secretion.

The second cytoplasmic loop domain of TMEM39A interacts with the core COPII

coating component Sec23A. TMEM131 binds to COL1A2 to facilitate assembly of

procollagen trimers and TRAPP III activation of Rab GTPase, in coordination with

TMEM39A to promote the ER-to-Golgi transport of procollagen cargo in COPII. Uso1

interacts with the COPII vesicle to promote targeting to the Golgi apparatus.

Table 1. RNAi phenotypic analysis of COPII related genes in *hsp-4*p::GFP (ER stress) and COL-19::GFP (collagen) .

Gene	Description	ER	collagen	Phenotype
tmem-39	Recruit Sec23A	+++	----	Smaller, disrupting
tmem-131	Recruit TRAPPC8, procollagen	+++	----	Smaller, disrupting
sec-23	COPII component	+++	----	L1-L2 arrest
sec-24.1	COPII component	+++	----	L1-L2 arrest
sec-24.2	COPII component	N.D.	N.D.	N.D.
npp-20	Sec13, COP II component	+++	N.D.	L4 arrest, disrupting
sec-31	COPII component	+	N.D.	N.D.
sar-1	GTPase, interact with sec-12	+++	--	L4 arrest, disrupting
sec-12	GTP-binding Sar1 protein	+++	N.D.	L1-L2 arrest
rab-1	Rab GTPase	+++	----	L1-L2 arrest
trpp-3	TRAPPIII component	++	----	N.D.
trpp-6	TRAPPIII component	N.D.	----	N.D.
trpp-8	TRAPPIII component	N.D.	----	sterile
uso-1	USO1 vesicle transport factor	+++	N.D.	N.D.

+ indicates *hsp-4*p::GFP fluorescent reporter level induction, and – indicates COL-19::GFP level reduction under different RNAi conditions; qualitative changes were followed up by quantitative fluorescence (n≥ 50 biological replicates) and western blot analysis for verification. N.D., no difference.

Fig. 1

609　**Fig. 2**

615 **Fig. 3**

A

P1   *w*; UAS-Cg25C.RFP.2.1; Lsp2-Gal4* X   *y v; UAS-CG13016_dsRNA*
$\dfrac{}{Cyo,wp}$ $\dfrac{}{TM6B}$

Fat body specific enhancer | Gal4

UAS | TMEM39 dsRNA

+

UAS binding sites | Col4a1:RFP

B

Control   *tmem39* RNAi

Fat body cells

100 µm

Drosophila Lsp2>Collagen::RFP

616 **Fig. 4**

618 **Fig. 5**

622   **Supplementary Materials**

623   **Figures. S1 to S4**

624   **Fig. S1**

Refer to Extended Data Fig. S1 (continued)

Fig. S1. *tmem-39* **RNAi knock-down for screen of phenotypic defects of different**

translational fluorescent reporters. (A-V), Exemplar fluorescence images showing

translational reporters for (A) *wrk-1*, (B) *gna-1*, (C) *hmr-1*, (D) *mans*, (E) *eff-1*, (F-G)

cpl-1, (H) *mig-23*, (I) *fat-7*, (J) *ced-1*, (K) *egl-20*, (L) *ubiquitin-V*, (M) *Y73E7A.8*, (N)

spon-1, (O) *him-4*, (P) *SP12*, (Q) *lgg-1*, (R) *F23H12.5*, (S) *lrp-1*, (T) *emb-9*, (U) *T19D2.1*

and (V) *cat-1* in wild-type animals by control and *tmem-39* RNAi at 20 °C (n = 3-4 for

each reporters). Scale bars: 20 µm.

Fig. S2

650 **Fig. S2.** *tmem-39* **RNAi knock-down other cuticle collagen translational**

651 **fluorescent reporters.** (A-B), Exemplar fluorescence images showing translational

652 reporters for (A) *col-101 and* (B*) lon-3.* In wild-type animals at 20 °C (n = 3-4 for each

653 reporters). Scale bars: 20 μm.

654
655

Fig. S3

A Ctrl

(100%, n=50)
col-19::GFP

B *sar-1*

(60%, n=50)

C *sec-24.1*

(70%, n=50)

D *sec-24.2*

(100%, n=50)

E *sec-31*

(100%, n=50)
col-19::GFP

F *trpp-6*

(80%, n=50)

G *trpp-8*

(90%, n=50)

H *npp-20*

(100%, n=50)

I *sec-12*

(100%, n=50)

J *rab-1*

(100%, n=50)

K *tmem-131*

(100%, n=50)

Fig. S3. RNAi knock-down other COP II component genes for screen genes

involved in collagen production and degradation. (A-J), Exemplar fluorescence

images of *col-19* translational reporter for (A) control, (B) *sar-1*, (C) *sec-24.1*, (D) *sec-*

24.2, (E) *sec-31*, (F) *trpp-6*, (G) *trpp-8*, (H) *npp-20*, (I) *sec-12,* (J) *rab-1* and (K) *tmem-*

131 RNAi in wild-type animals at 20 °C. Scale bars: 20 μm.

Fig. S4

Fig. S4. RNAi knock-down COPII component genes for screen genes involved in ER stress response. (A-J), Exemplar fluorescence images of *hsp-4*p::GFP transcriptional reporter for (A) control, (B) *sec-24.1*, (C) *sar-1*, (D) *npp-20*, (E) *tmem-131*, (F) *sec-24.2*, (G) *sec-31*, (H) *pdi-2*, (I) *trpp-8* and (J) *uso-1*RNAi in wild-type animals at 20 °C. Scale bars: 20 μm.

Table S1A. Primers and oligos used in genomic editing.

Primer	Sequence (5'-3')
sgRNA-EcoRI-F	ttgtaaaacgacggccagtgaattc**CTCCAAGAACTCGTACAAA AATG**
sgRNA-HindIII-R	ctatgaccatgattacgccaagctt**CACAGCCGACTATGTTTG**
TMEM-39	**GCCGCCTCGAAGACGAGTGC**GTTTAAGAGCTATGC
sgRNA1 F:	TGGAAACAGC
TMEM-39	**GCACTCGTCTTCGAGGCGGC**AAACATTTAGATTTGC
sgRNA1 R:	AATTCAATTA
TMEM-39	**GATCTACGAACCTTCTCAAG**GTTTAAGAGCTATGCT
sgRNA2 F:	GGAAACAGC
TMEM-39	**CTTGAGAAGGTTCGTAGATC**AAACATTTAGATTTGCA
sgRNA2 R:	ATTCAATTA
TMEM-39 YR	CCTCCTATTGCGAGATGTCTTG**GCTCGATAGCACCT**
sgRNA F:	**GCCGTC**GTTTAAGAGCTATGCTGG

sgRNA cloning (left column label spanning the above rows)

Primer	Sequence (5'-3')
TMEM-39 deletion oligo:	aagat**ATGCCGCCTCGAAGACGAG**^**AAGAGGATCTTC TCCTTATGCATCAACTTCAACAAGAACGATAA**
TMEM-39 YR oligo:	ACGCCATGGAGCGAGTTTTGTCTCTATAATGACGGA GAGACTGTTCAAATGCCGGACGGCAGGTGC**GCAGC A**GCCAAATCTTCGAATTCGATAAGAACTGTGGCCGC ACATCCGGAAAGTTCTCGGCATAACACTTTCTTTgta

Donor oligos (left column label spanning the above rows)

674 Primers were used for PCR with the Addgene plasmid #46169 as template, except YR

675 with the Addgene Plasmid #61253 as template. Bold and underlined are sgRNA target

676 sequence from the *tmem-39* regions. The location of the deleted *tmem-39* is marked

677 with "^", generating a precise 2750 bp deletion. And the nucleotides change for YR

678 codons are labeled in red (**TATCG**A→**GCAGC**A).

Table S1B. Primers in Genotyping and RT-PCR.

	Primer	Sequence (5'-3')
Geno-typing	TMEM-39 Del screen F:	tacagaaccgagaaggtcac
	TMEM-39 Del screen R:	tcacaattgggtagtaccac
	TMEM-39 Del screen R2:	GTGTGAACTGAATATCCGGC
	TMEM-39 YR screen F1:	ATCTTCTACACCGAGCAGCT
	TMEM-39 YR screen R1:	ATGCCGAGAACTTTCCGGAT
	COL-19::GFP screen F：	TTCCAGGACAAAAGGGAGAG
	COL-19::GFP screen R：	TCTCGAGAAGCATTGAACAC
RT-PCR	act-1 RT-F:	CATCCCAGTTGGTGACGATA
	act-1 RT-R:	TCGGTATGGGACAGAAGGAC
	gfp RT-F:	TGTTCCATGGCCAACACTTG
	gfp RT-R:	ACGTGTCTTGTAGTTCCCGT
	gfp RT-R:	ACGTGTCTTGTAGTTCCCGT
	col-19 RT-F:	TACTTGTGTGCGTTCTTGCC
	col-19 RT-R:	TTGGGTTGATGTGCTTGCTC
	tmem-39 RT-F:	GCTTCAATCCCAAGAGCGAG
	tmem-39 RT-R:	GACTTCGGAAGCCACCAAAG

Table S2. Reporters examined in phenotypic screen for *tmem-39* RNAi

Genotype	Reporter	Control	*tmem-39*
hsp-4p::GFP	*hsp-4* transcriptional	*	****
col-19p::col-19::GFP	*col-19* translational	****	*
cat-1p::cat-1::GFP	*cat-1* translational	**	**
eff-1p::eff-1::GFP	*eff-1* translational	*	**
emb-9p::emb-9::mCherry	*emb-9* translational	***	***
lrp-1p::lrp-1::GFP	*lrp-1* translational	**	**
fat-7p::fat-7::GFP	*fat-7* translational	***	***
gna-1p::GFP::gna-1	*gna-1* translational	**	**
him-4p::GFP::him-4	*him-4* translational	**	***
hmr-1p::hmr-1::GFP	*hmr-1* translational	***	***
lgg-1p::mCherry::GFP::lgg-1	*lgg-1* translational	**	**
lim-7p::ced-1::GFP	*ced-1* translational	**	**
myo-3p:: EGFP::wrk-1	*wrk-1* translational	***	****
nhx-2p::cpl-1::YFP	*cpl-1* translational	**	***
nhx-2p::cpl-1(W32A Y35A)::YFP	*cpl-1(W32A Y35A)* translational	***	***
nhx-2p::ubiquitin-V::mCherry	ubiquitin-V proteins	***	***
rpl-28p::F23H12.5::mCherry	F23H12.5 translational	****	****
rpl-28p::T19D2.1::mCherry	T19D2.1 translational	****	****
rpl-28p::Y73E7A.8::mCherry	Y73E7A.8 translational	****	****
spon-1p::spon-1::vGFP	*spon-1* translational	**	**
unc-22p::egl-20::GFP	*egl-20* translational	*	***
unc-54p::RFP::SP12	ER membrane RFP marker	**	***
unc-54p::mig-23::GFP	*mig-23* translational	***	***
vha-6p::mans::GFP	intestinal GFP marker for the Golgi	***	***

683 * indicates fluorescent reporter levels under control and *tmem-39* RNAi conditions;

684 qualitative changes were followed up by quantitative fluorescence (n≥ 20 biological

685 replicates) and Western blot for verification.