The anti-sigma factor MucA is required for viability in *Pseudomonas aeruginosa*

Melissa C. Schofield[1], Daniela Rodriguez[1], Amanda A. Kidman[1], Erin K. Cassin[1], Lia A.

Michaels[1], Elizabeth A. Campbell[2], Peter A. Jorth[3] and Boo Shan Tseng[1]*

[1] School of Life Sciences, University of Nevada Las Vegas, Las Vegas, NV, USA

[2] Rockefeller University, New York, NY, USA

[3] Departments of Pathology and Laboratory Medicine, Medicine, and Biomedical Sciences,

Cedars-Sinai Medical Center, Los Angeles, CA, USA

Running title: MucA is required in *P. aeruginosa*

* Corresponding author

Email: boo.tseng@unlv.edu

Keywords: AlgU, cystic fibrosis, sigma factor competition, envelope stress

ABSTRACT

During decades-long infections in the cystic fibrosis (CF) airway, *Pseudomonas aeruginosa*

undergoes selection. One bacterial genetic adaptation often observed in CF isolates is *mucA*

mutations. MucA inhibits the sigma factor AlgU. Mutations in *mucA* lead to AlgU misregulation,

resulting in a mucoid phenotype that is associated with poor CF disease outcomes. Due to its

ability to be mutated, *mucA* is assumed to be dispensable for bacterial viability. Here we show

that, paradoxically, a portion of *mucA* is essential in *P. aeruginosa*. We demonstrate that *mucA*

is no longer required in a strain lacking *algU*, that *mucA* alleles encoding for proteins that do not

bind to AlgU are insufficient for viability, and that *mucA* is no longer essential in mutant strains

containing AlgU variants with reduced sigma factor activity. Furthermore, we found that

overexpression of *algU* prevents cell growth in the absence of MucA, and that this phenotype

can be rescued by overproduction of RpoD, the housekeeping sigma factor. Together, these

results suggest that in the absence of MucA, the inability to regulate AlgU activity results in the

loss of bacterial viability. Finally, we speculate that essentiality of anti-sigma factors that

regulate envelope function may be a widespread phenomenon in bacteria.

INTRODUCTION

The major cause of death in people with cystic fibrosis (CF), a human autosomal recessive

genetic disease, is respiratory failure due to chronic lung infection. *Pseudomonas aeruginosa* is

a prevalent CF respiratory pathogen (Cystic Fibrosis Foundation Patient Registry, 2019). The

CF lung environment selects for mucoid *P. aeruginosa* mutants, which overproduce the

exopolysaccharide alginate and are associated with poor disease prognosis (Douglas *et al.*,

2009, Emerson *et al.*, 2002, Farrell *et al.*, 2009, Henry *et al.*, 1992, Li *et al.*, 2005, Nixon *et al.*,

2001, Parad *et al.*, 1999, Pedersen *et al.*, 1992, Strateva *et al.*, 2010). Conversion to the mucoid

phenotype in clinical *P. aeruginosa* isolates, which is thought to be advantageous for chronic

infection, is often caused by *mucA* mutations (Martin *et al.*, 1993).

MucA is an anti-sigma factor to the alternative sigma factor AlgU (also known as AlgT, σ^E, or

σ^{22}), which responds to envelope stress (Damron & Goldberg, 2012, Govan & Deretic, 1996).

MucA is a transmembrane protein that sequesters AlgU away from RNA polymerase (RNAP)

via its N-terminus (Li *et al.*, 2019, Schurr *et al.*, 1996, Xie *et al.*, 1996). The C-terminus of MucA

is in the periplasm, where it is protected from proteolysis by MucB (Mathee *et al.*, 1997, Schurr

et al., 1996). The envelope stress response is controlled via a regulated intramembrane

proteolysis cascade: MucB dissociates from MucA upon stress detection, allowing the proteases

AlgW and MucP to cleave MucA from the inner membrane (Qiu *et al.*, 2007, Wood & Ohman,

2009). In the cytoplasm, MucA is further degraded by ClpXP, releasing AlgU to interact with

RNAP and activate the AlgU regulon (Qiu *et al.*, 2008). AlgU regulates at least 350 genes,

including those responsible for the production of itself, MucA, and the alginate biosynthetic

enzymes (Firoved & Deretic, 2003, Schulz *et al.*, 2015, Wood & Ohman, 2009). This system is

homologous to the well-studied envelope stress response in *Escherichia coli*: MucA shares 28%

identity (72% similarity) to the anti-sigma factor RseA. The cognate sigma factor of the *E. coli*

RseA, called RpoE, is homologous to and functionally interchangeable with AlgU (Yu *et al.*,

1995), with 66% identity (93% similarity).

For several reasons, *mucA* is assumed to be dispensable for *P. aeruginosa* viability. First, after

being released from the cell membrane, MucA is presumed to be fully degraded in the

cytoplasm by ClpXP (Qiu *et al.*, 2008). Second, *mucA* mutations often arise in CF clinical

isolates (Boucher *et al.*, 1997, Candido Cacador *et al.*, 2018, Ciofu *et al.*, 2008, Martin *et al.*,

1993, Pulcrano *et al.*, 2012), and many laboratory *mucA* mutants exist in the literature

(Gallagher *et al.*, 2011, Liberati *et al.*, 2006, Skurnik *et al.*, 2013, Turner *et al.*, 2015). Third,

there are three published strains in which the entirety of *mucA* is removed from the genome

(Intile *et al.*, 2014, Jones *et al.*, 2010, Pritchett *et al.*, 2015). Paradoxically, three different whole-

genome studies in two different strains of *P. aeruginosa* identified *mucA* as an essential gene

(Lee *et al.*, 2015, Liberati *et al.*, 2006, Skurnik *et al.*, 2013). Additionally, it has been anecdotally

noted that *mucA* could not be deleted from the *P. aeruginosa* PAO1 genome (Panmanee *et al.*,

2019). To investigate this paradox, we systematically attempted to delete *mucA* from *P.*

aeruginosa using allelic exchange. Our results show that a portion of *mucA* was required for

bacterial viability in multiple *P. aeruginosa* strains, that *mucA* was no longer essential in a strain

lacking *algU*, and that *mucA* alleles that encode for proteins that do not interact with AlgU were

insufficient to rescue viability or led to a growth defect. We found that *algU* mutations encoding

for a less active sigma factor could relieve *mucA* essentiality, and that AlgU overproduction in

the absence of MucA was toxic. Interestingly, our works suggests that *mucA* essentiality can be

suppressed by increasing the levels of the housekeeping sigma factor RpoD. Together, our

results strongly suggest that the unregulated activity of AlgU itself, in the absence of MucA,

leads to bacterial cell death.

RESULTS

***mucA* is essential for viability in a diverse set of *P. aeruginosa* isolates**

To determine if *mucA* is essential, we attempted to delete the gene from various *P. aeruginosa*

strains, using a modified allelic exchange protocol to turn it into a robust assay (Fig S1). An

allele of *mucA* missing >95% of the coding region was introduced into *P. aeruginosa*. Using

PCR-confirmed isolates containing both the endogenous and deletion alleles in the genome, we

counter-selected for loss of one allele and then determined which *mucA* allele isolates resolved

to via PCR. For non-essential genes, we should observe isolates that resolved to the

endogenous or deletion allele. However, for essential genes, only cells that resolved to the

endogenous allele should be isolated, as cells that resolved to the deletion allele should not

survive in the absence of the gene. For statistical power, we screened ≥125 isolates for strains

resolving only to the endogenous *mucA*. We performed this assay on a diverse set of wild-type

P. aeruginosa strains, including four laboratory, five clinical and two environmental isolates

(Table 1). These strains are not only isolated from diverse locations, but they also vary in their

colony morphology and their exopolysaccharide production profile (Colvin *et al.*, 2012). For all

strains tested, all observed isolates resolved to the endogenous *mucA* allele ($p < 0.0001$,

Fisher's exact test), strongly suggesting that *mucA* is essential in wild-type *P. aeruginosa*. Since

all tested strains required *mucA* for viability, we continued our experiments using PAO1 as a

representative strain.

Depletion of MucA leads to cell death

To evaluate the effect of nutrient conditions on *mucA* essentiality, we tested the effect of

depleting MucA on *P. aeruginosa* viability in different media. We engineered a strain lacking the

native *mucA* and containing a chromosomally-integrated rhamnose-inducible copy (Δ*mucA*

*att*Tn7::P*rhaBAD*-*mucA*). We determined viability of this strain grown without rhamnose over time.

Since cells would cease producing MucA in the absence of rhamnose, the MucA present in the

Figure S1. Allelic exchange assay.
(A) Schematic of assay. To delete the endogenous *mucA* (short red box with *mucA*) from the genome, the deletion vector pBT396, which contains an allele of *mucA* that is missing >95% of the coding region (tall red box with delta), was introduced into *P. aeruginosa* via conjugation. Regions of homology (tall blue boxes) to the genome (short blue boxes) approximately 400 base pairs in length flank the deletion allele and allow for recombination into the genome (X) to create a merodiploid. This first recombination event (1st) can be selected for using antibiotics due to integration of the vector backbone, which contains a gentamicin resistance marker. At least six merodiploids were confirmed via PCR to ensure the genome contained both the deletion and endogenous *mucA* allele. Merodiploids can then undergo a second recombination event (2nd) that will lead to the loss of one of the two alleles, resolving to either the endogenous or deletion allele, which results in a cell that is either wild-type (top) or a deletion mutant (bottom), respectively. For each of the six confirmed merodiploids, we counter-selected for the second recombination event via the loss of *sacB*, a vector backbone marker. Using PCR, eight isolates per merodiploid were tested to determine which allele each isolate resolved to. If a gene is non-essential, we expect to observe both wild type or deletion mutants. However, if a gene is essential, we expect to isolate only those cells that resolved to wild type, as the cells cannot survive with the deletion allele. If we are unable to delete *mucA* from the first replicate of this experiment, we perform two additional biological replicates, for a minimum total of 125 colonies screened. If all isolates resolve to wild-type, we deem *mucA* to be essential in that strain background ($p < 0.0001$, Fisher's exact test). **(B)** Representative image of merodiploid confirmation. The PCR products corresponding to the endogenous allele and the deletion allele are indicated with an arrow labelled "WT" and "Δ," respectively. Positive (WT ctrl; PAO1) and negative (Δ*mucA* ctrl; BTPa355) controls are included. PCR products from six representative merodiploids are shown. **(C)** Representative image of PCR products from isolates after the second recombination. Image is labeled as in (B). Top, the PCR products of 8 isolates from a strain in which *mucA* was deemed essential. Bottom, the PCR products of 8 isolates from a strain in which *mucA* was not essential.

Table 1. *mucA* is essential in a variety of wild-type *P. aeruginosa* strains.

Strain background	Description[†]	Number of isolates resolved to	
		WT	Δ*mucA*
PAO1	Laboratory strain, Class II	168	0*
PA14	Laboratory strain, Class I	147	0*
PA103	Laboratory strain, unclassified	361	0*
PAK	Laboratory strain, unclassified	191	0*
CF127	Mucoid CF isolate, unclassified	136	0*
CF18	Non-mucoid CF isolate, unclassified	129	0*
CF27	Rugose CF Isolate, Class IV	152	0*
X13273	Blood isolate, Class II	142	0*
X24509	UTI isolate, Class II	221	0*
MSH10	Water isolate, Class III	146	0*
E2	Tomato plant isolate, Class II	141	0*

[†] Class identification is based on the exopolysaccharide expression profile as described in (Colvin *et al.*, 2012): Class I, Pel-dominant matrix; Class II, Psl-dominant matrix; Class III, exopolysaccharide redundant matrix users; and Class IV, matrix overproducers. CF, cystic fibrosis; UTI, urinary tract infection.

* $p < 0.0001$, Fischer's exact test

cells at the start of the experiment would be depleted as the cells divided. In all four media we

tested, if rhamnose was not removed, the cells increased in density by ~2-logs over time. In

comparison, when rhamnose was removed, the cells lost viability over time in all four media (Fig

1), showing that *mucA* is essential in *P. aeruginosa* in various nutritional environments.

Alginate biosynthesis is not solely responsible for *mucA* essentiality

Clinical isolates with *mucA* mutations are mucoid due to the overproduction of alginate (Martin

et al., 1993). To determine if alginate overproduction is responsible for *mucA* essentiality, we

attempted to delete *mucA* from a strain lacking *algD*, a key alginate biosynthesis gene (Deretic

et al., 1987). We were unable to delete *mucA* in this background (Table S1).

Expression of the alginate biosynthesis genes is controlled by three AlgU-regulated transcription

factors that are active in mucoid cells: AlgB, AlgR, and AmrZ (Martin *et al.*, 1994, Wozniak &

Ohman, 1994, Wozniak *et al.*, 2003). Since these transcription factors regulate many genes in

addition to those involved in alginate biosynthesis (Huang *et al.*, 2019, Jones *et al.*, 2014, Kong

et al., 2015, Leech *et al.*, 2008), we tested if overexpression of these three regulons underlies

mucA essentiality. We could not delete *mucA* from strains lacking these transcription factors

(Table S1). We conclude that eliminating alginate biosynthesis and the expression of ~50%

AlgU regulon do not alleviate *mucA* essentiality.

The first 50 amino acids of MucA are necessary and sufficient for cell viability

While we were unable to delete *mucA* using an allele lacking >95% of the coding region, *mucA*

is often mutated in clinical isolates and many *mucA* transposon mutants exist (Fig S2),

suggesting that only a portion of *mucA* is essential. Based on the co-crystal structures of MucA

with AlgU and MucB (Li *et al.*, 2019), the first 78 residues of MucA interact with AlgU, and the

last 48 residues of MucA interact with MucB. We attempted to delete the endogenous *mucA*

Figure 1. Depletion of MucA results in cell death.
Viable colony counts of PAO1 Δ*mucA* attTn7::P*rhaBAD*-*mucA* over time in **(A)** LB, **(B)** PIB, **(C)** SCFM and **(D)** VBMM with (+ rha; gray squares) or without (– rha; red circles) 0.05% rhamnose. Viable colony counts after incubation in the indicated condition and time was determined by plating the cells on LB with 0.05% rhamnose to allow cells to recover and grow. Hash, broken y-axis; error bars, SEM (N=3). Asterisk, statistically different from that at the same time point grown in the presence of rhamnose (p < 0.01, N=3, mixed model ANOVA with post-hoc Bonferroni test).

Table S1. Eliminating alginate biosynthesis does not alleviate *mucA* essentiality.

PAO1	Number of isolates resolved to	
	WT	Δ*mucA*
WT	168	0*
Δ*algD*	144	0*
Δ*algB*	158	0*
Δ*algR*	153	0*
Δ*amrZ*	131	0*
Δ*algB* Δ*algR* Δ*amrZ*	133	0*

* $p < 0.0001$, Fischer's exact test

Figure S2. Published mutations in *mucA*.
Top, schematic of MucA with the regions indicated encoding for the AlgU binding domain (green, AlgU BD), the transmembrane domain (blue, TM), and the MucB binding domain (purple, MucB BD). Hash marks on top indicate the residue number in MucA. Bottom, sites of *mucA* mutations that lead to the production of a truncated protein in published isolates. Each line represents one or more individual isolates described in the references on the right. The mutations for the upper five references (Boucher *et al.*, 1997, Candido Cacador *et al.*, 2018, Ciofu *et al.*, 2008, Martin *et al.*, 1993, Pulcrano *et al.*, 2012) are in cystic fibrosis clinical isolates of *P. aeruginosa*, while the mutations in the last reference (Turner *et al.*, 2015) are random transposon insertions in the laboratory PAO1 and PA14 strains.

from a series of strains containing an ectopic chromosomally-integrated *mucA*, driven by its

native promoter and encoding for truncation products (Fig 2A). We were unable to probe

whether equivalent amounts of protein were produced across our strains, since our truncated

versions of MucA are cleaved out of the membrane and degraded in the cytosol, as previously

shown (Qiu *et al.*, 2008). However, we expect that the ectopic alleles are expressed at similar

levels prior to endogenous *mucA* deletion, as they are under the same promoter and at the

same genomic site. These strains with two *mucA* alleles were non-mucoid, showing that the

endogenous *mucA* allele, encoding full-length protein, is dominant over the ectopic alleles that

encode for truncated MucA. We used a Δ*algD* background to make strains easier to manipulate,

as deleting the endogenous *mucA* would result in alginate overproduction in some strains.

We then attempted to delete the endogenous *mucA* from these strains. Since the flanking

regions of the two alleles differ, our deletion allele specifically targets the native allele. We were

able to easily recover isolates that resolved to the *mucA* deletion allele from strains containing

ectopic alleles encoding for the full length MucA (aa 1-194), as well as MucA aa 1-143, aa 1-

110, aa 1-75, and aa 1-62 (Fig 2B). Interestingly, while the endogenous *mucA* could be deleted

from strains containing an ectopic *mucA* that encoded for aa 1-50 (Fig 2B), isolates resolving to

the deletion allele grew up more slowly than those resolving to the native allele. This suggests

that while this allele is sufficient for viability, it is not well tolerated. All tested isolates of strains

carrying *mucA* alleles encoding shorter products (aa 1-40 and aa 1-24) resolved to the native

allele. Furthermore, we were unable to delete the native *mucA* from a strain with an ectopic

mucA that encodes for aa 51-194 (Fig 2B). These results suggest that the first 50 amino acids

of MucA are necessary and sufficient for *P. aeruginosa* cell viability. Consistent with these

results, reported *mucA* mutations (Boucher *et al.*, 1997, Candido Cacador *et al.*, 2018, Ciofu *et*

al., 2008, Martin *et al.*, 1993, Pulcrano *et al.*, 2012, Turner *et al.*, 2015) almost entirely fall

outside the region of *mucA* encoding the first 50 residues (Fig S2).

Figure 2. The first 50 amino acids of MucA are necessary and sufficient for viability.
(A) Schematic of MucA encoded by the ectopic *mucA* alleles in PAO1 Δ*algD* strains tested for *mucA* essentiality in (B). MucA aa 1-143 lacks the MucB binding domain and is the product of the common *mucA22* allele. MucA 1-110 lacks the entire periplasmic domain. MucA aa 1-75 contains most of the AlgU binding domain, while shorter truncations contain only parts of the AlgU binding domain (MucA aa 1-62, 1-50, 1-40, and 1-24). Green, AlgU binding domain (AlgU BD); blue, the transmembrane domain (TM); and purple, the MucB binding domain (MucB BD).
(B) Frequency of observed isolates resolving to the endogenous *mucA* allele (WT, black) or the deletion allele (Δ*mucA*, white) in the allelic exchange assay. Super-imposed on the bars are the number of isolates that were tested in each category. Asterisk, p < 0.0001; Fisher's exact test. Caret, slower growing isolates.

The interaction of MucA with AlgU is required for cell survival

Since the only described function for MucA is to inhibit AlgU, we hypothesized that *mucA*

essentiality is rooted in its regulation of AlgU. To test this, we attempted to delete *mucA* from a

strain lacking *algU* (Δ*algU*). We were able to do so, with 16 of the tested isolates resolving to the

deletion allele out of the 48 colonies tested (see Fig 4A), showing that *mucA* essentiality is *algU*-

dependent.

Because our data suggest that the AlgU-binding domain of MucA is required for viability (Fig

2B), we hypothesized that the physical interaction between MucA and AlgU is necessary for cell

survival. In the co-crystal structure (Li *et al.*, 2019), four residues in MucA make more than one

hydrogen bond with AlgU: D15, E22, R42, and E72 (Fig 3A). We engineered alleles that encode

for MucA D15A, E22A, R42A, or E72A to maximally affect the hydrogen bonding while limiting

the effects on overall protein structure. We used MucA aa 1-75 as a base because the co-

crystal structure includes the first 78 residues of MucA. As described above, an allele encoding

MucA aa 1-75 was sufficient for viability (Fig 2B). We were unable to delete the native *mucA*

from strains carrying alleles encoding the D15A and R42A substitutions (Fig 3B). While we were

able to delete the native *mucA* from the strain containing MucA E22A, isolates that resolved to

the deletion allele grew up slower than those that resolved to the wild-type allele, suggesting

that the cells do not tolerate this allele well. As expected, due to being outside the required

region of MucA (Fig 2B), we were able to delete the native *mucA* from a strain containing MucA

E72A with 9 of 48 tested isolates resolving to the deletion allele.

To determine if the D15A, E22A, and R42A substitutions affect the ability of MucA to interact

with AlgU, we used a yeast-two hybrid assay. Using beta-galactosidase activity as a proxy for

their interaction, wild-type MucA aa 1-75 strongly interacted with AlgU (Fig 3C). This interaction

Figure 3. The physical interaction of MucA and AlgU is required for survival.
(A) Four residues of MucA make greater than one predicted hydrogen bond with AlgU. The MucA-AlgU co-crystal structure (PDB 6IN7; Li et al., 2019) with the cytosolic domain of MucA (aa 1-78; red) and Regions 2 and 4 of AlgU (green, $AlgU_2$; yellow, $AlgU_4$) is shown. The residues of MucA that are predicted to make more than one hydrogen bond with AlgU (grey) are labeled in the inset. Black dotted lines, predicted hydrogen bonds; red atoms, oxygen; blue atoms, nitrogen. **(B)** Frequency of observed isolates resolving to the endogenous *mucA* allele (WT, black) or the deletion allele (Δ*mucA*, white) in the allelic exchange assay, using PAO1 Δ*algD att*Tn7::P*algU*-*mucA*, where *mucA* encodes for the indicated substitution. Super-imposed on the bars are the number of isolates that were tested in each category. Asterisk, $p < 0.0001$; Fisher's exact test. Caret, slower growing isolates. **(C)** Substitution of MucA residues at its interface with AlgU abolish their binding via yeast two-hybrid. The first 75 residues of MucA were fused to the Gal4 DNA-binding domain (DBD-MucA$^{1-75}$) and AlgU was fused to the Gal4 activation domain (AD-AlgU). Interaction of MucA and AlgU led to *lacZ* expression. Beta-galactosidase activity (in Miller units) was used as a proxy for the protein interaction strength. WT, wild-type protein sequence; –, no fusion protein included; hash, broken y-axis; error bars, SEM (N=3); letters, statistical groupings ($p < 0.01$; biological triplicate with technical quadruplicates; ANOVA with post-hoc Tukey HSD).

was dependent on the presence of both MucA and AlgU, and the effect was not directional (Fig

S3). In comparison, the D15A, E22A, and R42A MucA mutants failed to interact with AlgU (Fig

3C). We conclude that the interaction of MucA and AlgU are required for viability.

Mutations that reduce AlgU activity alleviate the requirement for MucA

If *mucA* essentiality is due to its inhibition of the AlgU regulon, strains containing a mutant AlgU

with lower affinity for DNA should alleviate *mucA* essentiality because the regulon expression is

reduced in such mutants. We engineered Δ*algU* strains with an ectopic chromosomally-

integrated *algU* allele encoding such DNA-binding mutants (K57A and N81A), based on

homology to *E. coli* RpoE mutants that have decreased *in vitro* transcriptional activity

(Campagne *et al.*, 2014). As described above, we were able to delete *mucA* from a Δ*algU* strain

(Fig 4A). This phenotype could be rescued via the ectopic addition of a wild-type *algU* allele, as

we were no longer able to delete *mucA* from such a strain. In contrast, *mucA* could be deleted

from strains carrying alleles encoding AlgU K57A or N81A, showing that these alleles failed to

rescue the Δ*algU* phenotype (Fig 4A). To determine the effect of these substitutions on AlgU

activity, we induced envelope stress using D-cycloserine (Wood *et al.*, 2006) and measured

AlgU activity using a plasmid-borne *gfp* reporter driven under the AlgU-regulated *algD* promoter

(Damron *et al.*, 2009). Similar to what is seen for RpoE (Campagne *et al.*, 2014), these mutant

strains had reduced AlgU activity upon induction of envelope stress (Fig 5A,B).

Strains with the entirety of *mucA* deleted from *P. aeruginosa* PAO1, PAK, and PA103 exist

(Intile *et al.*, 2014, Jones *et al.*, 2010, Pritchett *et al.*, 2015). Since we were unable to delete

mucA from these strain backgrounds (Table 1), we sequenced the genomes of these Δ*mucA*

strains to determine if they contain suppressor mutations that allowed for their survival in the

absence of *mucA*. In PAO1 Δ*mucA* (Pritchett *et al.*, 2015), while the native *mucA* was deleted,

the strain contained a second full-length copy of *algU* and a *mucA* allele that encoded for aa 1-

DBD	Krev1	Krev1	MucA$^{1-75}$	MucA$^{1-75}$	–	AlgU	–	AlgU
AD	Ral^{WT}	Ral^{Mut}	AlgU	–	AlgU	MucA$^{1-75}$	MucA$^{1-75}$	–
Avg	2.832	0.017	17.142	0.008	0.010	6.833	0.017	0.009
SEM	0.402	0.010	0.221	0.001	0.004	0.355	0.006	0.003

Figure S3. MucA aa 1-75 and AlgU interact via yeast two-hybrid assay.
The indicated proteins were fused to the Gal4 DNA binding domain (DBD; "bait") or the Gal4 activation domain (AD; "prey"). Interaction of the bait and prey proteins drive the expression of *lacZ*. Beta-galactosidase activity (in Miller units) was used as a proxy for the protein interaction strength. The average (Avg) and standard error of the mean (SEM) of each bar are indicated (N=3). As positive and negative controls, Krev1 is known to interact with wild-type RalGDS (Ral^{WT}), but not the mutant RalGDS (Ral^{Mut}) via yeast two-hybrid. AlgU, construct encoding full-length wild-type AlgU; MucA$^{1-75}$, construct encoding only the first 75 residues of wild-type MucA; –, no fusion protein included; error bars, SEM (N=3); letters, statistical groups with the different letters representing statistically different groups (p < 0.01; biological triplicate with technical quadruplicates; ANOVA with post-hoc Tukey HSD).

Figure 4. Mutations in AlgU can suppress *mucA* essentiality.
(A) Frequency of observed isolates resolving to the endogenous *mucA* allele (WT, black) or the deletion allele (Δ*mucA*, white) in the allelic exchange assay, using PAO1 Δ*algU att*Tn7::P*algU*-*algU*, where *algU* encodes for the indicated substitution. Super-imposed on the bars are the number of isolates that were tested in each category. Asterisk, p < 0.0001; Fisher's exact test.
(B) Schematic of mutations seen in revertants that could grow in the absence of *mucA*. Revertants were selected by growing PAO1 Δ*mucA att*Tn7::P*rhaBAD*-*mucA* on media lacking rhamnose. Blue rectangles, multi-base pair deletions; left arrow, deletion extends into the promoter; blue triangles, single base pair deletions; black triangles, nonsense mutations; white triangles, duplications resulting in 3 or 4 amino acid insertions; red triangles, missense mutations. See Table S2 for a full description of the *algU* mutations.

Figure 5. AlgU mutants that suppress *mucA* essentiality have decreased activity.
(A) Fold-induction of GFP in PAO1 wild-type (WT) and Δ*algU* carrying a plasmid with either a promoter-less *gfp* (vector) or a *gfp* driven by the *algD* promoter (*algD* reporter), which is positively regulated by AlgU, after a 2h D-cycloserine treatment to activate envelope stress. GFP fluorescence was normalized to cell density and was then divided by the signal from untreated cells to determine the fold-induction. Hash, broken y-axis; error bars, SEM (N=3);

letters, statistical groupings (p < 0.01; biological triplicate with technical quadruplicates; ANOVA with post-hoc Tukey HSD). **(B)** Fold-induction of GFP in PAO1 Δ*algU att*Tn7::P*algU*-*algU*, where *algU* encodes for the indicated substitution. The experiment and statistics are as described in (A). WT, the wild-type AlgU sequence. **(C)** Fold-induction of GFP in the revertants isolated from PAO1 Δ*mucA att*Tn7::P*rhaBAD*-*mucA* (parental) that can grow in the absence of MucA. The experiment and statistics are as described in (A), except all cells were grown in the presence of 0.05% rhamnose to allow for comparison to the parental strain, which grows only in the presence of rhamnose. The isolate identification numbers are shown and are grouped based on their *algU* mutation. **(D)** Viable colony counts of PAO1 Δ*mucA* attTn7::P*rhaBAD*-*mucA* carrying the *algD* reporter over time in LB with (+ rha; gray squares) or without (– rha; red circles) 0.05% rhamnose. Hash, broken y-axis; error bars, SEM (N=3). Asterisk, statistically different from that at the same time point grown in the presence of rhamnose (p < 0.01, N=3, mixed model ANOVA with post-hoc Bonferroni test). **(E)** Normalized GFP fluorescence of the samples in (D). GFP fluorescence was normalized to cell density. The statistical analysis is as described in (D).

155 elsewhere in the genome. As expected (Fig 2B), we were able to delete the native *mucA*

allele from a PAO1 strain containing an ectopic allele that encoded for MucA aa 1-155 (with 5 of

the 28 tested isolates resolving to the deletion allele), confirming that the ectopic *mucA* allele in

the published PAO1 Δ*mucA* strain is sufficient for viability in the absence of the endogenous

mucA. The PAK and PA103 Δ*mucA* strains (Intile *et al.*, 2014, Jones *et al.*, 2010) contain

missense mutations in *algU,* which result in A58T and E46G, respectively. We replicated these

mutations in PAO1 by inserting *algU* alleles encoding these substitutions in a Δ*algU* strain. We

were able to delete *mucA* from these strains (Fig 4A). These results confirm that the *algU* alleles

in the published PAK and PA103 Δ*mucA* strains suppress *mucA* essentiality. Using our reporter

assay, we found that strains carrying these AlgU substitutions had reduced sigma factor activity

(Fig 5B). We note that our reporter assay is not very sensitive to low levels of AlgU activity. Both

PAK and PA103 Δ*mucA* strains are mucoid (Intile *et al.*, 2014, Jones *et al.*, 2010), suggesting

that AlgU A58T and E46G are not completely inactive. Nevertheless, our results show that

these AlgU mutants have significantly less transcriptional activity than the wild-type protein (Fig

5A,B).

To identify additional suppressors of *mucA* essentiality, we used the Δ*mucA att*Tn7::P*rhaBAD*-

mucA strain. This strain was not viable in the absence of rhamnose, but natural revertants arose

at a frequency of less than 1 in 10^9 colony forming units. We sequenced the *algU* gene in 25

revertants, since AlgU mutants can suppress *mucA* essentiality (Fig 4A). All 25 isolates

contained mutations in *algU* that are predicted to be hypomorphic (Fig 4B, Table S2). There

were 10 revertants with deletions or nonsense mutations of *algU*, encoding either no product or

a truncated product completely lacking Region 4 of the sigma factor. There were 6 revertants

that contained multi-base pair duplications in *algU* that would lead to the insertion of 3 or 4

amino acids in Region 2 helix 3 of the sigma factor. There were 9 revertants containing

missense mutations, 8 of which were unique, encoding the following substitutions: D18G, A21V,

Y29C, A47T, D49G, Y59C, N81D, and R174G. Using a model of σ^E in complex with the RNAP

core and the promoter element, we expect these insertions and substitutions to affect sigma

factor folding, RNAP core interactions, or promoter interactions (Fig S4). Using our *algD*

reporter assay in these natural revertants, we saw that all 25 revertants had much lower AlgU

activity than the parental strain upon induction of envelope stress (Fig 5C), as predicted.

Together, these data strongly suggest that *algU* mutations that encode for a protein with

reduced transcriptional activity allow *P. aeruginosa* to survive in the absence of *mucA*.

If *mucA* essentiality is due to AlgU inhibition, AlgU activity should increase as MucA is depleted

from cells, concomitant with the decrease in viability. To test this, the viability and fluorescence

(as a proxy for AlgU activity) of a Δ*mucA att*Tn7::P*rhaBAD*-*mucA* strain containing our *algD*

reporter plasmid was measured over time in the absence of rhamnose. This strain had a similar

viability to that seen in Figure 2 (Fig 5D), and as MucA was depleted, the AlgU activity

increased (Fig 5E). These results show the cell death observed upon MucA depletion is

correlated with a dramatic increase in AlgU activity.

Overexpression of *algU* in the absence of *mucA* leads to a growth defect

Our results suggest that *mucA* essentiality is due to unregulated AlgU activity. We reasoned that

overexpression of *algU* should be toxic. However, *algU* can be overexpressed in wild-type *P.*

aeruginosa that contain *mucA* (Qiu *et al.*, 2008, Schulz *et al.*, 2015). We therefore examined the

effect of overproducing AlgU in wild type, Δ*algU*, and Δ*algU* Δ*mucA* strains, using a

chromosomally integrated arabinose-inducible *algU*. Of note, the expression of *algU* is

completely dependent on the inducer in the Δ*algU* and Δ*algU* Δ*mucA* strains, which lack positive

feedback of AlgU on its own expression. Our results show that in the absence of inducer when

AlgU is not overexpressed, all three strains had similar growth rates (Fig 6A). In the presence of

Table S2. Description of changes found in *algU* of PAO1 Δ*mucA att*Tn7::P*rhaBAD*-*mucA* revertants that can grow in the absence of *mucA* expression.

	Isolate	Genomic Change to *algU*[†]	Effect on AlgU[‡]
Multi-base pair deletions			
	#1-10	831242-831394 del	No product made
	#2-9	831298-831491 del	No product made
	#2-7	831467-831473 del	I56fs X97
	#2-11	831469-831502 del	K57fs X88
	#2-6	831595-831715 del	V99fs X107
Single base pair deletions			
	#2-14	831467 del	I56fs X99
	#1-1	831510 del	Y72fs X99
	#2-10	831538 del	I80fs X99
Nonsense mutations			
	#2-1	831433 C>T	Q45X
	#2-8	831521 G>A	W74X
Multi-base pair insertions			
	#1-9	831427-831435 ins GACGCCCAG	D43-A44 ins DAQ
	#1-8	831430-831438 ins GCCCAGGAA	A44-Q45 ins AQE
	#1-11	831436-831444 ins GAAGCCCAG	E46-A47 ins EAQ
	#2-4	831436-831444 ins GAAGCCCAG	E46-A47 ins EAQ
	#1-5	831445-831456 ins GACGTAGCCCAG	D49-V50 ins DVAQ
	#1-6	831451-831459 ins GCCCAGGAA	A51-Q52 ins AQE
Missense mutations			
	#1-3	831353 A>G	D18G
	#2-12	831362 C>T	A21V
	#2-3	831386 A>G	Y29C
	#2-2	831439 G>A	A47T
	#1-2	831446 A>G	D49G
	#2-13	831446 A>G	D49G
	#1-7	831476 A>G	Y59C
	#2-5	831541 A>G	N81D
	#1-4	831820 C>G	R174G

[†] genomic location based on the Pseudomonas Genome Database (Winsor *et al.*, 2016) for PAO1; bp, base pair; del, deletion; ins, insertion.

[‡] fs, frameshift; ins, insertion; X, stop codon.

Figure S4. Location of affected AlgU residues in revertants that can grow in the absence of MucA.

Model of the RNAP holoenzyme containing σ^E bound to the promoter. Gray, RNAP core; cyan, β flap; pink, β' coiled coil; green, σ^E Region 2; yellow, σ^E Region 4; blue/light blue, promoter element. Insets, location of residues that are substituted in the revertants with missense *algU* mutations. The side chain of the affected residue is in white. Substitution of A21 and A47 (V in model) likely affect protein packing and folding of AlgU. Substitution of D18, Y29, and Y59 would reduce predicted intra- and inter-molecular hydrogen bonding and likely affect AlgU folding. Substitution of D49, N81, and R174 would likely affect the interaction of AlgU with the RNAP core or the promoter. Dashed yellow lines, hydrogen bonds; red atoms, oxygen; blue atoms, nitrogen; orange, phosphorus.

1% arabinose (i.e. high expression of *algU*), all three strains had a growth defect with the strain

lacking *mucA* failing to grow at all.

To determine if the Δ*algU* Δ*mucA att*Tn7::P*araBAD*-*algU* strain can grow under lower levels of *algU*

induction, we tested a range of inducer concentrations (Fig 6B, Table S3). We found that

although there was a growth defect, Δ*algU* Δ*mucA att*Tn7::P*araBAD*-*algU* was able to grow in the

presence of 0.1% arabinose. The drop in growth rate in comparison to the no arabinose

condition was statistically larger for the Δ*algU* Δ*mucA att*Tn7::P*araBAD*-*algU* than for the other two

strains (N=3, $p < 0.05$, ANOVA with post hoc Tukey HSD), since this strain lacks the ability to

produce any MucA to reduce AlgU activity. These results strongly suggest that in the absence of

mucA, while a certain low level of AlgU activity is tolerated, high AlgU activity is fatal to the cell.

The above experiments were performed using only one medium. To determine if *algU*

overexpression in the absence of *mucA* leads to a growth defect under other nutrient conditions,

we determined the growth rate of the Δ*algU* Δ*mucA att*Tn7::P*araBAD*-*algU* strain in other media.

The Δ*algU* Δ*mucA att*Tn7::P*araBAD*-*algU* strain failed to grow in the presence of 1% arabinose for

all four media we tested (Fig S5), showing that AlgU overproduction is toxic under various

nutrient conditions.

Expression of *rpoD* can rescue the growth defect of *algU* overexpression

AlgU competes for RNAP binding with RpoD, the essential primary sigma factor (Yin *et al.*,

2013). We, therefore, tested if the growth defect of high *algU* expression in the Δ*algU* Δ*mucA*

background could be ameliorated by overexpressing *rpoD* under the same arabinose-inducible

promoter. We tested growth in the presence of 2% arabinose, since these new strains contain

two arabinose-inducible promoters, in contrast to the strains described in Figure 6. We

confirmed that RpoD was overproduced in these new strains at similar levels upon arabinose

Figure 6. Overexpression of *algU* leads to a growth defect.
(A) Growth curves of PAO1 strains containing an arabinose-inducible copy of *algU* in wild-type (purple squares), Δ*algU* (blue diamonds), and Δ*algU* Δ*mucA* (red circles) backgrounds in LB. Open symbols represent conditions in the absence of arabinose; closed symbols, with 1% arabinose. Error bars, SD (n=18). **(B)** Growth rate of strains as described in (A), with increasing induction of *algU*. Strains were grown in LB with 0%, 0.1%, 0.25%, and 1% arabinose ([Ara]). Error bars, SEM (N=3). Asterisk, statistically different from the same strain grown without arabinose ($p < 0.01$, N = 3, two-way ANOVA with post-hoc Bonferroni). See Table S3 for full statistical comparisons.

Table S3. Statistical differences in growth rate among groups under conditions of *algU* induction.

Medium: LB	Strain	*att*Tn7::P*araBAD*-*algU*				Δ*algU* *att*Tn7::P*araBAD*-*algU*				Δ*algU* Δ*mucA* *att*Tn7::P*araBAD*-*algU*			
Strain	Ara	0	0.1	0.25	1	0	0.1	0.25	1	0	0.1	0.25	1
*att*Tn7::P*araBAD*-*algU*	0	▓	*	*	*	▓	▓	▓	▓		▓	▓	▓
	0.1	▓	▓		*	▓		▓	▓	▓		▓	▓
	0.25	▓	▓	▓		▓	▓		▓	▓	▓	*	▓
	1	▓	▓	▓	▓	▓	▓	▓	*	▓	▓	▓	*
Δ*algU* *att*Tn7::P*araBAD*-*algU*	0	▓	▓	▓	▓	▓		*	*		▓	▓	▓
	0.1	▓	▓	▓	▓	▓	▓	*	*	▓	*	▓	▓
	0.25	▓	▓	▓	▓	▓	▓	▓	*	▓	▓	*	▓
	1	▓	▓	▓	▓	▓	▓	▓	▓	▓	▓	▓	
Δ*algU* Δ*mucA* *att*Tn7::P*araBAD*-*algU*	0	▓	▓	▓	▓	▓	▓	▓	▓	▓	*	*	*
	0.1	▓	▓	▓	▓	▓	▓	▓	▓	▓	▓	*	*
	0.25	▓	▓	▓	▓	▓	▓	▓	▓	▓	▓	▓	
	1	▓	▓	▓	▓	▓	▓	▓	▓	▓	▓	▓	

Medium: PIB	Strain	*att*Tn7::P*araBAD*-*algU*		Δ*algU* *att*Tn7::P*araBAD*-*algU*		Δ*algU* Δ*mucA* *att*Tn7::P*araBAD*-*algU*	
Strain	Ara	0	1	0	1	0	1
*att*Tn7::P*araBAD*-*algU*	0	▓	*		▓		▓
	1	▓	▓			▓	
Δ*algU* *att*Tn7::P*araBAD*-*algU*	0	▓	▓	▓	*		▓
	1	▓	▓	▓	▓	▓	
Δ*algU* Δ*mucA* *att*Tn7::P*araBAD*-*algU*	0	▓	▓	▓	▓	▓	*
	1	▓	▓	▓	▓	▓	▓

Medium: SCFM	Strain	*att*Tn7::P*araBAD*-*algU*		Δ*algU* *att*Tn7::P*araBAD*-*algU*		Δ*algU* Δ*mucA* *att*Tn7::P*araBAD*-*algU*	
Strain	Ara	0	1	0	1	0	1
*att*Tn7::P*araBAD*-*algU*	0	▓	*		▓		▓
	1	▓	▓			▓	
Δ*algU* *att*Tn7::P*araBAD*-*algU*	0	▓	▓	▓	*		▓
	1	▓	▓	▓	▓	▓	
Δ*algU* Δ*mucA* *att*Tn7::P*araBAD*-*algU*	0	▓	▓	▓	▓	▓	*
	1	▓	▓	▓	▓	▓	▓

Medium: VBMM	Strain	*att*Tn7::P*araBAD*-*algU*		Δ*algU* *att*Tn7::P*araBAD*-*algU*		Δ*algU* Δ*mucA* *att*Tn7::P*araBAD*-*algU*	
Strain	Ara	0	1	0	1	0	1
*att*Tn7::P*araBAD*-*algU*	0	▓	*		▓	*	▓
	1	▓	▓			▓	*
Δ*algU* *att*Tn7::P*araBAD*-*algU*	0	▓	▓	▓	*		▓
	1	▓	▓	▓	▓	▓	*
Δ*algU* Δ*mucA* *att*Tn7::P*araBAD*-*algU*	0	▓	▓	▓	▓	▓	*
	1	▓	▓	▓	▓	▓	▓

Significance was determined via a two-way ANOVA with post-hoc Bonferroni. Ara, concentration of arabinose (%). Asterisk, $p < 0.05$. Open box, no statistical difference.

Figure S5. Overexpression of *algU* in the absence of *mucA* causes a growth defect in various media.
Growth rate of indicated strains grown in (A) LB, (B) PIB, (C) SCFM, and (D) VBMM with (+) or without (–) 1% arabinose (relative to PAO1 *att*Tn7::P$_{araBAD}$-*algU* grown in the absence of arabinose). Error bars, SEM (N=3). Asterisk, statistically different from the same strain grown in the absence of arabinose (p < 0.01, N = 3, two-way ANOVA with post-hoc Bonferroni). See Table S3 for full statistical comparisons. LB contains tryptone and yeast extract, both of which provide protein hydrolates, as the main carbon source. The main carbon source of PIB is also protein hydrolates, but from pancreatic gelatin digest. While PIB also contains glycerol, which can serve as a carbon source, it is not a preferred carbon source for *Pseudomonas aeruginosa*. SCFM is a defined medium that contains amino acids as the primary carbon source, and mimics the nutrients found in the CF lung environment. VBMM is a minimal medium that contains citrate as the sole carbon source.

induction (Fig S6). As expected (Fig 6), Δ*algU* Δ*mucA* *att*Tn7::P*araBAD*-*algU* failed to grow in the

presence of 2% arabinose. The Δ*algU* Δ*mucA* *att*Tn7::P*araBAD*-*algU* *att*CTX::P*araBAD*-*rpoD* strain,

however, was able to grow with arabinose. Furthermore, the growth rate of this strain with

arabinose was indistinguishable from that of Δ*algU* Δ*mucA* *att*CTX::P*araBAD*-*rpoD* (Table S4).

There are two major explanations for how sigma factor competition may lead to toxicity during

algU overexpression. First, the AlgU overproduction may reduce housekeeping gene expression

due to the limited availability of RNAP to interact with RpoD. Second, *algU* overexpression

results in high AlgU activity, which may lead to toxic expression of the AlgU regulon. We,

therefore, constructed a strain containing an inducible *algU* encoding a K57A substitution in the

Δ*algU* Δ*mucA* background. AlgU K57A exhibited reduced sigma factor activity (Fig 5A,B) and

based on its location in the crystal structure (Campagne *et al.*, 2014), we do not expect this

substitution to affect RNAP affinity. In the presence of arabinose, the Δ*algU* Δ*mucA*

*att*Tn7::P*araBAD*-*algU* K57A strain exhibited a growth rate greater than that of the strain

overexpressing wild-type AlgU, but significantly less than that of the parental Δ*algU* Δ*mucA*

strain (Fig 7, Table S4). RpoD overproduction in this background returned growth to a

statistically similar level as that of a strain overexpressing RpoD alone (Table S4). Assuming

that the K57A substitution does not affect AlgU affinity for RNAP, these results suggest that

AlgU overexpression may lead to toxic expression of the AlgU regulon.

DISCUSSION

Contrary to its assumed dispensability, our work shows that *mucA* is essential in a variety of *P.*

aeruginosa wild-type strains and under various nutrient conditions (Table 1, Fig 1). Our data

strongly suggest that unchecked AlgU activity in the absence of MucA leads to cell death (Fig

8). Under non-stress conditions, MucA inhibits AlgU. Under envelope stress or in strains

containing *mucA* mutations, the anti-sigma factor is cleaved. We propose that while this cleaved

Figure S6. RpoD production in strains with an ectopic inducible *rpoD* allele.
Expression of *rpoD* was inferred via semi-quantitative Western blot in the indicated strains (top). RNAP was used as a loading control (bottom). Strains were grown in LB with (+) or without (–) 2% arabinose. The relative amount of RpoD is indicated, where 1.00x represents RpoD levels in the parental strain without arabinose (± SD; N=3). While the strains produce significantly more RpoD in the presence of arabinose relative to the same strain in the absence of arabinose (p < 0.01), there is no statistical difference in the amount of RpoD produced from the induced strains containing an ectopic *rpoD* allele (two-way ANOVA with post-hoc Bonferroni).

Figure 7. Growth defect caused by *algU* overexpression can be rescued via *rpoD* overexpression.

Growth rate of indicated strains grown in LB with (+) or without (−) 2% arabinose. Error bars, SEM (N=3). Asterisk, statistically different from the same strain grown without arabinose ($p < 0.01$, N = 3, two-way ANOVA with post-hoc Bonferroni). See Table S4 for full statistical comparisons.

Table S4. Statistical differences in growth rate among groups overexpressing *algU* and/or *rpoD*.

Strain	Ara	ΔalgUΔmucA P_BAD-rpoD		ΔalgUΔmucA P_BAD-algU^WT		ΔalgUΔmucA P_BAD-algU^WT P_BAD-rpoD		ΔalgUΔmucA P_BAD-algU^K57A		ΔalgUΔmucA P_BAD-algU^K57A P_BAD-rpoD	
		0	2	0	2	0	2	0	2	0	2
ΔalgUΔmucA	0		▨		▨		▨		▨		▨
ΔalgUΔmucA	2	▨		▨	*	▨		▨	*	▨	
ΔalgUΔmucA P_BAD-rpoD	0	▨			▨		▨		▨		▨
ΔalgUΔmucA P_BAD-rpoD	2	▨	▨	▨	*	▨		▨	*	▨	
ΔalgUΔmucA P_BAD-algU^WT	0	▨	▨	▨	*		▨		▨		▨
ΔalgUΔmucA P_BAD-algU^WT	2	▨	▨	▨	▨	▨	*	▨	*	▨	*
ΔalgUΔmucA P_BAD-algU^WT P_BAD-rpoD	0	▨	▨	▨	▨	▨	*		▨		▨
ΔalgUΔmucA P_BAD-algU^WT P_BAD-rpoD	2	▨	▨	▨	▨	▨	▨	▨		▨	
ΔalgUΔmucA P_BAD-algU^K57A	0	▨	▨	▨	▨	▨	▨	▨	*		▨
ΔalgUΔmucA P_BAD-algU^K57A	2	▨	▨	▨	▨	▨	▨	▨	▨	▨	
ΔalgUΔmucA P_BAD-algU^K57A P_BAD-rpoD	0	▨	▨	▨	▨	▨	▨	▨	▨	▨	*
ΔalgUΔmucA P_BAD-algU^K57A P_BAD-rpoD	2	▨	▨	▨	▨	▨	▨	▨	▨	▨	▨

Significance was determined via a two-way ANOVA with post-hoc Bonferroni. Ara, concentration of arabinose (%). Asterisk, $p < 0.05$. Open box, no statistical difference.

Figure 8. Unchecked AlgU activity in the absence of MucA leads to bacterial cell death. Under conditions of no envelope stress (left), the full-length MucA (red) binds strongly to AlgU (green star), leaving very little free AlgU to interact with the RNAP (grey oval). Therefore, the AlgU regulon (green arrow) is in the "off" state (red X). Under conditions of envelope stress or in strains containing *mucA* mutations that lead to a truncated product (middle), a cleaved cytosolic form of MucA is produced. This form can still interact with and inhibit AlgU. However, the strength of the interaction is weaker than with the full-length MucA, allowing for a pool of free AlgU that can then bind to and recruit RNAP to the promoters of its regulon. Under such conditions, the AlgU regulon, which includes *algU* and *mucA*, is activated and in the "on" state (green up arrows). However, because of the feedback on MucA, negative regulation of AlgU activity is still present. In the absence of *mucA* (right), there is no MucA to inhibit AlgU. All of the AlgU is free to interact with the core RNAP. This leads to high expression of the AlgU regulon and overproduction of AlgU itself. Under such strong positive feedback and in the absence of the negative regulator, the unchecked AlgU activity leads to cell death.

cytosolic form of MucA does not inhibit AlgU to the same extent as the full-length protein, it is

still able to inhibit AlgU to some degree. Although AlgU is active under such conditions, because

mucA is positively regulated by AlgU, this negative feedback keeps AlgU activity under control.

In comparison, this ability to control the positive feedback of AlgU on its own expression and

activity is lost in the absence of MucA, which we propose is the cause of cell death. Supporting

this model are several lines of evidence. First, *mucA* essentiality is rooted in its interaction with

algU (Fig 2-3), strongly suggesting that the ability of MucA to inhibit AlgU is required for viability.

Second, AlgU mutants with decreased transcriptional activity can suppress *mucA* essentiality

(Fig 4-5), supporting the idea that decreasing the positive feedback of AlgU on its own

expression allows for survival in the absence of MucA. Lastly, *algU* overexpression led to a

growth defect, which was lethal at high levels in the absence of *mucA* (Fig 6), suggesting that

high AlgU activity leads to cell death when MucA is not present. Overall, our work strongly

suggests that *mucA* essentiality is caused by unchecked AlgU activity, in agreement with

previous studies suggesting that overproduction of AlgU is toxic (Cross *et al.*, 2020,

Hershberger *et al.*, 1995, Schurr *et al.*, 1994).

AlgU overproduction may be toxic because of reduced housekeeping gene expression (due to

sigma factor competition with RpoD) and/or the increased expression of the AlgU regulon.

Increasing RpoD levels alleviate the toxicity in both cases, by increasing the RNAP-RpoD

complexes and reducing the RNAP-AlgU complexes. Our data does not definitively distinguish

between these two mechanisms, which may not be mutually exclusive. Supporting a role for

decreased housekeeping functions in AlgU toxicity, *mucA* was still essential in a strain lacking

the three major AlgU-regulated transcription factors AlgB, AlgR, and AmrZ (Table S1), which

regulate ~50% of the AlgU regulon (Huang *et al.*, 2019, Jones *et al.*, 2014, Leech *et al.*, 2008,

Schulz *et al.*, 2015). However, we cannot exclude that a combination of genes in the other half

of the AlgU regulon may be responsible for *mucA* essentiality. Supporting the hypothesis that

increased AlgU regulon expression causes toxicity, mutations in *algU*, which is AlgU-regulated,

suppress *mucA* essentiality (Fig 4). While this autoregulation of AlgU is important for *mucA*

essentiality under physiological conditions, it does not lead to toxicity per se, since

overexpressing *algU* in a strain lacking the positive feedback led to a severe growth defect (Fig

6). Under such conditions, overproduction of AlgU K57A, which has reduced sigma factor

activity (Fig 5B), caused a less severe growth defect than wild-type AlgU overproduction (Fig 7).

Assuming that RNAP affinity is not affected by this AlgU substitution, this suggests that high

AlgU activity and the ensuing regulon expression may be the mechanism underlying toxicity.

Work in *Bacillus subtilis* has shown that the toxicity of unregulated SigM, an alternative sigma

factor, is due specifically to membrane protein overproduction, which is alleviated by a gain-of-

function mutation to the membrane insertase YidC1 (Zhao *et al.*, 2019b). Since only ~80 of the

~200 AlgU-regulated predicted membrane proteins are regulated by AlgB, AlgR, and AmrZ

(Huang *et al.*, 2019, Jones *et al.*, 2014, Leech *et al.*, 2008, Schulz *et al.*, 2015), it is possible that

a gain-of-function YidC mutant may suppress *mucA* essentiality in *P. aeruginosa*. Such an

experiment would help distinguish between the potential mechanisms underlying AlgU toxicity.

Anti-sigma factor essentiality is not unique to *P. aeruginosa*. In the defined transposon mutant

library for *Vibrio cholerae*, an interruption in the *mucA* homolog *rseA* does not exist, suggesting

that this anti-sigma factor may be essential (Cameron *et al.*, 2008). Similarly, the *Pseudomonas*

syringae mucA homolog is deemed essential based on transposon insertion analysis (Helmann

et al., 2019). Furthermore, in *Mycobacterium tuberculosis*, the gene encoding the anti-sigma

factor RslA is identified as essential (Griffin *et al.*, 2011). Similar to our results (Fig 4), *rslA* can

be deleted from *M. tuberculosis* lacking the gene encoding its cognate sigma factor SigL, which

regulates genes involved in cell envelope processes (Dainese *et al.*, 2006). Finally, there are

two anti-sigma factors, YhdL and YxlC, in *B. subtilis* that are necessary for bacterial viability

(Horsburgh & Moir, 1999, Koo *et al.*, 2017, Mendez *et al.*, 2012). The *yhdL* essentiality is

notable as this anti-sigma factor negatively regulates SigM, an envelope stress response sigma

factor (Horsburgh & Moir, 1999). Its essentiality could be suppressed by overproduction of SigA,

the *B. subtilis* housekeeping sigma factor (Zhao *et al.*, 2019a) and by a hyperactive form of the

membrane insertase YidC1 (Zhao *et al.*, 2019b). Interestingly, the YidC1 mutant ameliorated the

toxicity of unregulated SigM by reducing the secretion stress associated with high membrane

protein production (Zhao *et al.*, 2019b). Taken together with our data, while not universal (as

there are no essential anti-sigma factors in *E. coli* (Baba *et al.*, 2006)), we speculate that

essentiality of anti-sigma factors that regulate cell envelope function may be a widespread

phenomenon in bacteria and that their essentiality is due to the increased production of

membrane proteins when their sigma factors are not held in check.

In clinical CF isolates of *P. aeruginosa*, *mucA* mutations that lead to C-terminal truncations are

common (Fig S2) and are expected to result in proteins that retain partial AlgU-inhibitory

function. Interestingly, there is one *P. aeruginosa* CF isolate that is reported to have a *mucA*

mutation that would result in a truncation within the first 50 aa of the protein (Boucher *et al.*,

1997). Our sequencing data of the published Δ*mucA* strains (Intile *et al.*, 2014, Jones *et al.*,

2010, Pritchett *et al.*, 2015) suggests that this isolate may contain a suppressor mutation that

allows it to survive in the absence of a functional MucA. Nonetheless, in CF clinical isolates,

mucA mutations that lead to C-terminal truncations are the norm, while null mutations in *mucA*

with a presumed hypomorphic *algU* allele is rare (Fig S2). Furthermore, our results (Fig 4) agree

with the literature showing that mucoid isolates with *mucA* mutations revert to a non-mucoid

state via changes to *algU* (Ciofu *et al.*, 2008, DeVries & Ohman, 1994, Sautter *et al.*, 2012,

Schurr *et al.*, 1994). While these *algU* secondary site mutations are found in non-mucoid CF

isolates with *mucA* mutations, such isolates are detected at a lower frequency (Ciofu *et al.*,

2008). As suggested by Ciofu and associates, this may be due to the importance of AlgU for the

survival of *P. aeruginosa* in the CF lung environment, suggesting that reducing AlgU activity

may increase *P. aeruginosa* eradication from the CF airway. We propose that the MucA-AlgU

interaction may serve as a good therapeutic target. Our results show that destabilizing the

MucA-AlgU interaction results in bacterial cell death or mutations in *algU* that result in reduced

sigma factor activity and likely reduced mucoid conversion, of which either outcome could be

beneficial in the treatment of *P. aeruginosa* CF lung infections.

METHODS AND MATERIALS

Bacterial strains and growth conditions. The bacterial strains, plasmids, and oligonucleotides

used for this study are in Tables S5-S7. The construction of strains is described in

Supplementary Information. Bacteria were grown at 37°C in LB with shaking or on semi-solid LB

media, unless otherwise noted.

Allelic exchange assay. The protocol, depicted in Figure S1, was modified from (Hmelo *et al.*,

2015). Briefly, a vector containing the *mucA* deletion allele flanked by approximately equal

length homology regions is introduced via conjugation. Merodiploids were selected on semi-

solid VBMM (Vogel & Bonner, 1956) with 60 mg/L gentamicin. Using OMS118 and OMS119,

PCR was performed on at least six isolates to confirm that the presence of both the wild-type

and deletion alleles of *mucA*. Confirmed merodiploids were then individually streaked on NSLB

(10 g/L tryptone, 5 g/L yeast extract) with 10% sucrose semi-solid media for counterselection.

PCR was performed on eight colonies per merodiploid, using OBT601 and OBT602 to

determine the resolution to either the endogenous or deletion allele.

MucA depletion assay. Cells were grown in LB with 0.05% rhamnose at 37°C with shaking to

an OD_{600} of 0.3. After washing, the culture was then divided in two, half resuspended with

rhamnose and half without in its respective media base (LB, PIB, SCFM, or VBMM). Cells were

incubated at 37°C with shaking. At the indicated time points, two aliquots were removed from

each culture, serially diluted and plated in triplicate onto LB agar plates with 0.05% rhamnose

for recovery. Colonies were counted and $\log_{10}$-transformed CFU/mL of the culture was

calculated.

Yeast two-hybrid assay. The ProQuest Two-Hybrid System (Invitrogen) was used, per

manufacturer's instructions. Briefly, yeast containing the Gal4 activation domain-based prey and

the Gal4 DNA-binding domain-based bait vectors were grown overnight in SD-Leu-Trp broth

(Clontech). The OD_{600} value was recorded. ONPG (VWR) was added to lysed cells, and the

mixture was incubated at 37°C until a light yellow color was achieved. The incubation time was

recorded. The OD_{420} of the supernatants was determined using a Synergy Hybrid HTX

Microplate Reader (Bio-Tek Instruments). Beta-galactosidase activity was determined using

Miller units, based on the following equation: 1000 x OD_{420}/(time x culture volume x OD_{600}).

AlgU activity assay. Strains of interest were transformed with a plasmid-borne *algD* reporter

(pBT435) via electroporation (Choi *et al.*, 2006). Strains were grown overnight in LSLB with 50

mg/L gentamicin and 0.05% rhamnose (where indicated) at 37°C with shaking. The overnight

culture was diluted 1:100 and grown to an OD_{600} of 0.3. Cells were then treated with fresh 400

mg/L D-cycloserine for 2 h at 37°C with shaking, as previously described (Wood *et al.*, 2006).

Cells were then pelleted and resuspended in PBS. GFP fluorescence and OD_{600} was

determined using a Synergy Hybrid HTX Microplate Reader (Bio-Tek Instruments). To

normalize the data, the GFP fluorescence was divided by the OD_{600} for each data point. To

determine the fold induction, the ratios for the treated samples were divided by the average of

that for the same strain not treated with D-cycloserine. The resulting ratios of the triplicate

samples were averaged.

Natural revertant assay. *P. aeruginosa* Δ*mucA att*Tn7::P$_{rhaBAD}$-*mucA* was grown in LB with

0.05% rhamnose at 37°C with shaking to an OD$_{600}$ of 1. Cells were washed and plated on semi-

solid LB plates without rhamnose. Plates were incubated for 24-48 h at 37°C. For isolates that

grew on LB without rhamnose, *algU* was Sanger sequenced.

ACKNOWLEGDEMENTS

The authors would like to thank Cai Tao and Maria Alexandra Ledesma for technical assistance;

and Joe J. Harrison for critical discussions of this work. MCS, AAK, DR, and BT are supported

by the Cystic Fibrosis Foundation (TSENG19I0). MCS and EKC are supported by University of

Nevada Las Vegas doctoral graduate research assistantships. EAC and PAJ are supported by

the National Institutes of Health (R01GM114450 and K22AI127473, respectively). PAJ is also

supported by the Cystic Fibrosis Foundation (JORTH17F5). The authors have no conflicts of

interest to disclose.

AUTHOR CONTRIBUTIONS

Conceptualization: MCS, BST; Acquisition, analysis, or interpretation of data: MCS, DR, AAK,

EKC, LAM, EAC, PAJ, BST; Writing – original draft: MCS, BST; and Writing – review and

editing: MCS, DR, LAM, EAC, PAJ, BST.

REFERENCES

Baba, T., Ara, T., Hasegawa, M., Takai, Y., Okumura, Y., Baba, M., Datsenko, K.A., Tomita, M.,

Wanner, B.L., and Mori, H. (2006) Construction of Escherichia coli K-12 in-frame, single-

gene knockout mutants: the Keio collection. *Mol Syst Biol* **2**: 2006 0008.

Boucher, J.C., Yu, H., Mudd, M.H., and Deretic, V. (1997) Mucoid Pseudomonas aeruginosa in

cystic fibrosis: characterization of muc mutations in clinical isolates and analysis of

clearance in a mouse model of respiratory infection. *Infect Immun* **65**: 3838-3846.

Cameron, D.E., Urbach, J.M., and Mekalanos, J.J. (2008) A defined transposon mutant library

and its use in identifying motility genes in Vibrio cholerae. *Proc Natl Acad Sci U S A* **105**:

8736-8741.

Campagne, S., Marsh, M.E., Capitani, G., Vorholt, J.A., and Allain, F.H. (2014) Structural basis

for -10 promoter element melting by environmentally induced sigma factors. *Nat Struct*

Mol Biol **21**: 269-276.

Candido Cacador, N., Paulino da Costa Capizzani, C., Gomes Monteiro Marin Torres, L.A.,

Galetti, R., Ciofu, O., da Costa Darini, A.L., and Hoiby, N. (2018) Adaptation of

Pseudomonas aeruginosa to the chronic phenotype by mutations in the algTmucABD

operon in isolates from Brazilian cystic fibrosis patients. *PLoS One* **13**: e0208013.

Choi, K.H., Kumar, A., and Schweizer, H.P. (2006) A 10-min method for preparation of highly

electrocompetent Pseudomonas aeruginosa cells: application for DNA fragment transfer

between chromosomes and plasmid transformation. *J Microbiol Methods* **64**: 391-397.

Ciofu, O., Lee, B., Johannesson, M., Hermansen, N.O., Meyer, P., and Hoiby, N. (2008)

Investigation of the algT operon sequence in mucoid and non-mucoid Pseudomonas

aeruginosa isolates from 115 Scandinavian patients with cystic fibrosis and in 88 in vitro

non-mucoid revertants. *Microbiology* **154**: 103-113.

Colvin, K.M., Irie, Y., Tart, C.S., Urbano, R., Whitney, J.C., Ryder, C., Howell, P.L., Wozniak,

D.J., and Parsek, M.R. (2012) The Pel and Psl polysaccharides provide Pseudomonas

aeruginosa structural redundancy within the biofilm matrix. *Environ Microbiol* **14**: 1913-

1928.

Cross, A.R., Raghuram, V., Wang, Z., Dey, D., and Goldberg, J.B. (2020) Overproduction of the

AlgT sigma factor is lethal to mucoid Pseudomonas aeruginosa. *Journal of Bacteriology*.

Cystic Fibrosis Foundation Patient Registry, (2019) 2018 Annual Data Report. In. Bethesda,

MD: Cystic Fibrosis Foundation, pp.

Dainese, E., Rodrigue, S., Delogu, G., Provvedi, R., Laflamme, L., Brzezinski, R., Fadda, G.,

Smith, I., Gaudreau, L., Palu, G., and Manganelli, R. (2006) Posttranslational regulation

of Mycobacterium tuberculosis extracytoplasmic-function sigma factor sigma L and roles

in virulence and in global regulation of gene expression. *Infect Immun* **74**: 2457-2461.

Damron, F.H., and Goldberg, J.B. (2012) Proteolytic regulation of alginate overproduction in

Pseudomonas aeruginosa. *Mol Microbiol* **84**: 595-607.

Damron, F.H., Qiu, D., and Yu, H.D. (2009) The Pseudomonas aeruginosa sensor kinase KinB

negatively controls alginate production through AlgW-dependent MucA proteolysis. *J*

Bacteriol **191**: 2285-2295.

Deretic, V., Gill, J.F., and Chakrabarty, A.M. (1987) Gene algD coding for GDPmannose

dehydrogenase is transcriptionally activated in mucoid Pseudomonas aeruginosa. *J*

Bacteriol **169**: 351-358.

DeVries, C.A., and Ohman, D.E. (1994) Mucoid-to-nonmucoid conversion in alginate-producing

Pseudomonas aeruginosa often results from spontaneous mutations in algT, encoding a

putative alternate sigma factor, and shows evidence for autoregulation. *J Bacteriol* **176**:

6677-6687.

Douglas, T.A., Brennan, S., Gard, S., Berry, L., Gangell, C., Stick, S.M., Clements, B.S., and

Sly, P.D. (2009) Acquisition and eradication of P. aeruginosa in young children with

cystic fibrosis. *Eur Respir J* **33**: 305-311.

Emerson, J., Rosenfeld, M., McNamara, S., Ramsey, B., and Gibson, R.L. (2002)

Pseudomonas aeruginosa and other predictors of mortality and morbidity in young

children with cystic fibrosis. *Pediatr Pulmonol* **34**: 91-100.

Farrell, P.M., Collins, J., Broderick, L.S., Rock, M.J., Li, Z., Kosorok, M.R., Laxova, A., Gershan,

W.M., and Brody, A.S. (2009) Association between MucoidPseudomonasInfection and

Bronchiectasis in Children with Cystic Fibrosis. *Radiology* **252**: 534-543.

Firoved, A.M., and Deretic, V. (2003) Microarray analysis of global gene expression in mucoid

Pseudomonas aeruginosa. *J Bacteriol* **185**: 1071-1081.

Gallagher, L.A., Shendure, J., and Manoil, C. (2011) Genome-scale identification of resistance

functions in Pseudomonas aeruginosa using Tn-seq. *MBio* **2**: e00315-00310.

Govan, J.R., and Deretic, V. (1996) Microbial pathogenesis in cystic fibrosis: mucoid

Pseudomonas aeruginosa and Burkholderia cepacia. *Microbiol Rev* **60**: 539-574.

Griffin, J.E., Gawronski, J.D., Dejesus, M.A., Ioerger, T.R., Akerley, B.J., and Sassetti, C.M.

(2011) High-resolution phenotypic profiling defines genes essential for mycobacterial

growth and cholesterol catabolism. *PLoS Pathog* **7**: e1002251.

Helmann, T.C., Deutschbauer, A.M., and Lindow, S.E. (2019) Genome-wide identification of

Pseudomonas syringae genes required for fitness during colonization of the leaf surface

and apoplast. *Proc Natl Acad Sci U S A* **116**: 18900-18910.

Henry, R.L., Mellis, C.M., and Petrovic, L. (1992) Mucoid Pseudomonas aeruginosa is a marker

of poor survival in cystic fibrosis. *Pediatr Pulmonol* **12**: 158-161.

Hershberger, C.D., Ye, R.W., Parsek, M.R., Xie, Z.D., and Chakrabarty, A.M. (1995) The algT

(algU) gene of Pseudomonas aeruginosa, a key regulator involved in alginate

biosynthesis, encodes an alternative sigma factor (sigma E). *Proc Natl Acad Sci U S A*

92: 7941-7945.

Hmelo, L.R., Borlee, B.R., Almblad, H., Love, M.E., Randall, T.E., Tseng, B.S., Lin, C., Irie, Y.,

Storek, K.M., Yang, J.J., Siehnel, R.J., Howell, P.L., Singh, P.K., Tolker-Nielsen, T.,

Parsek, M.R., Schweizer, H.P., and Harrison, J.J. (2015) Precision-engineering the

Pseudomonas aeruginosa genome with two-step allelic exchange. *Nat Protoc* **10**: 1820-

1841.

Horsburgh, M.J., and Moir, A. (1999) sigmaM, an ECF RNA polymerase sigma factor of Bacillus

subtilis 168, is essential for growth and survival in high concentrations of salt. *Molecular*

Microbiology **32**: 41-50.

Huang, H., Shao, X., Xie, Y., Wang, T., Zhang, Y., Wang, X., and Deng, X. (2019) An integrated

genomic regulatory network of virulence-related transcriptional factors in Pseudomonas

aeruginosa. *Nature Communications* **10**.

Intile, P.J., Diaz, M.R., Urbanowski, M.L., Wolfgang, M.C., and Yahr, T.L. (2014) The AlgZR

two-component system recalibrates the RsmAYZ posttranscriptional regulatory system

to inhibit expression of the Pseudomonas aeruginosa type III secretion system. *J*

Bacteriol **196**: 357-366.

Jones, A.K., Fulcher, N.B., Balzer, G.J., Urbanowski, M.L., Pritchett, C.L., Schurr, M.J., Yahr,

T.L., and Wolfgang, M.C. (2010) Activation of the Pseudomonas aeruginosa AlgU

regulon through mucA mutation inhibits cyclic AMP/Vfr signaling. *J Bacteriol* **192**: 5709-

5717.

Jones, C.J., Newsom, D., Kelly, B., Irie, Y., Jennings, L.K., Xu, B., Limoli, D.H., Harrison, J.J.,

Parsek, M.R., White, P., and Wozniak, D.J. (2014) ChIP-Seq and RNA-Seq reveal an

AmrZ-mediated mechanism for cyclic di-GMP synthesis and biofilm development by

Pseudomonas aeruginosa. *PLoS Pathog* **10**: e1003984.

Kong, W., Zhao, J., Kang, H., Zhu, M., Zhou, T., Deng, X., and Liang, H. (2015) ChIP-seq

reveals the global regulator AlgR mediating cyclic di-GMP synthesis in Pseudomonas

aeruginosa. *Nucleic Acids Res* **43**: 8268-8282.

Koo, B.M., Kritikos, G., Farelli, J.D., Todor, H., Tong, K., Kimsey, H., Wapinski, I., Galardini, M.,

Cabal, A., Peters, J.M., Hachmann, A.B., Rudner, D.Z., Allen, K.N., Typas, A., and

Gross, C.A. (2017) Construction and Analysis of Two Genome-Scale Deletion Libraries

for Bacillus subtilis. *Cell Syst* **4**: 291-305 e297.

Lee, S.A., Gallagher, L.A., Thongdee, M., Staudinger, B.J., Lippman, S., Singh, P.K., and

Manoil, C. (2015) General and condition-specific essential functions of Pseudomonas

aeruginosa. *Proc Natl Acad Sci U S A* **112**: 5189-5194.

Leech, A.J., Sprinkle, A., Wood, L., Wozniak, D.J., and Ohman, D.E. (2008) The NtrC Family

Regulator AlgB, Which Controls Alginate Biosynthesis in Mucoid Pseudomonas

aeruginosa, Binds Directly to the algD Promoter. *Journal of Bacteriology* **190**: 581-589.

Li, S., Lou, X., Xu, Y., Teng, X., Liu, R., Zhang, Q., Wu, W., Wang, Y., and Bartlam, M. (2019)

Structural basis for the recognition of MucA by MucB and AlgU in Pseudomonas

aeruginosa. *The FEBS Journal* **286**: 4982-4994.

Li, Z., Kosorok, M.R., Farrell, P.M., Laxova, A., West, S.E., Green, C.G., Collins, J., Rock, M.J.,

and Splaingard, M.L. (2005) Longitudinal development of mucoid Pseudomonas

aeruginosa infection and lung disease progression in children with cystic fibrosis. *JAMA*

293: 581-588.

Liberati, N.T., Urbach, J.M., Miyata, S., Lee, D.G., Drenkard, E., Wu, G., Villanueva, J., Wei, T.,

and Ausubel, F.M. (2006) An ordered, nonredundant library of Pseudomonas aeruginosa

strain PA14 transposon insertion mutants. *Proc Natl Acad Sci U S A* **103**: 2833-2838.

Martin, D.W., Schurr, M.J., Mudd, M.H., Govan, J.R., Holloway, B.W., and Deretic, V. (1993)

Mechanism of conversion to mucoidy in Pseudomonas aeruginosa infecting cystic

fibrosis patients. *Proc Natl Acad Sci U S A* **90**: 8377-8381.

Martin, D.W., Schurr, M.J., Yu, H., and Deretic, V. (1994) Analysis of promoters controlled by

the putative sigma factor AlgU regulating conversion to mucoidy in Pseudomonas

aeruginosa: relationship to sigma E and stress response. *J Bacteriol* **176**: 6688-6696.

Mathee, K., McPherson, C.J., and Ohman, D.E. (1997) Posttranslational control of the algT

(algU)-encoded sigma22 for expression of the alginate regulon in Pseudomonas

aeruginosa and localization of its antagonist proteins MucA and MucB (AlgN). *J Bacteriol*

179: 3711-3720.

Mendez, R., Gutierrez, A., Reyes, J., and Marquez-Magana, L. (2012) The extracytoplasmic

function sigma factor SigY is important for efficient maintenance of the Spbeta prophage

that encodes sublancin in Bacillus subtilis. *DNA Cell Biol* **31**: 946-955.

Nixon, G.M., Armstrong, D.S., Carzino, R., Carlin, J.B., Olinsky, A., Robertson, C.F., and

Grimwood, K. (2001) Clinical outcome after early Pseudomonas aeruginosa infection in

cystic fibrosis. *J Pediatr* **138**: 699-704.

Panmanee, W., Su, S., Schurr, M.J., Lau, G.W., Zhu, X., Ren, Z., McDaniel, C.T., Lu, L.J.,

Ohman, D.E., Muruve, D.A., Panos, R.J., Yu, H.D., Thompson, T.B., Tseng, B.S., and

Hassett, D.J. (2019) The anti-sigma factor MucA of Pseudomonas aeruginosa: Dramatic

differences of a mucA22 vs. a DeltamucA mutant in anaerobic acidified nitrite sensitivity

of planktonic and biofilm bacteria in vitro and during chronic murine lung infection. *PLoS*

One **14**: e0216401.

Parad, R.B., Gerard, C.J., Zurakowski, D., Nichols, D.P., and Pier, G.B. (1999) Pulmonary

outcome in cystic fibrosis is influenced primarily by mucoid Pseudomonas aeruginosa

infection and immune status and only modestly by genotype. *Infect Immun* **67**: 4744-

4750.

Pedersen, S.S., Hoiby, N., Espersen, F., and Koch, C. (1992) Role of alginate in infection with

mucoid Pseudomonas aeruginosa in cystic fibrosis. *Thorax* **47**: 6-13.

Pritchett, C.L., Little, A.S., Okkotsu, Y., Frisk, A., Cody, W.L., Covey, C.R., and Schurr, M.J.

(2015) Expression analysis of the Pseudomonas aeruginosa AlgZR two-component

regulatory system. *J Bacteriol* **197**: 736-748.

Pulcrano, G., Iula, D.V., Raia, V., Rossano, F., and Catania, M.R. (2012) Different mutations in

mucA gene of Pseudomonas aeruginosa mucoid strains in cystic fibrosis patients and

their effect on algU gene expression. *New Microbiol* **35**: 295-305.

Qiu, D., Eisinger, V.M., Head, N.E., Pier, G.B., and Yu, H.D. (2008) ClpXP proteases positively

regulate alginate overexpression and mucoid conversion in Pseudomonas aeruginosa.

Microbiology **154**: 2119-2130.

Qiu, D., Eisinger, V.M., Rowen, D.W., and Yu, H.D. (2007) Regulated proteolysis controls
mucoid conversion in Pseudomonas aeruginosa. *Proc Natl Acad Sci U S A* **104**: 8107-
8112.

Sautter, R., Ramos, D., Schneper, L., Ciofu, O., Wassermann, T., Koh, C.L., Heydorn, A.,
Hentzer, M., Hoiby, N., Kharazmi, A., Molin, S., Devries, C.A., Ohman, D.E., and
Mathee, K. (2012) A complex multilevel attack on Pseudomonas aeruginosa algT/U
expression and algT/U activity results in the loss of alginate production. *Gene* **498**: 242-
253.

Schulz, S., Eckweiler, D., Bielecka, A., Nicolai, T., Franke, R., Dotsch, A., Hornischer, K.,
Bruchmann, S., Duvel, J., and Haussler, S. (2015) Elucidation of sigma factor-
associated networks in Pseudomonas aeruginosa reveals a modular architecture with
limited and function-specific crosstalk. *PLoS Pathog* **11**: e1004744.

Schurr, M.J., Martin, D.W., Mudd, M.H., and Deretic, V. (1994) Gene cluster controlling
conversion to alginate-overproducing phenotype in Pseudomonas aeruginosa: functional
analysis in a heterologous host and role in the instability of mucoidy. *J Bacteriol* **176**:
3375-3382.

Schurr, M.J., Yu, H., Martinez-Salazar, J.M., Boucher, J.C., and Deretic, V. (1996) Control of
AlgU, a member of the sigma E-like family of stress sigma factors, by the negative
regulators MucA and MucB and Pseudomonas aeruginosa conversion to mucoidy in
cystic fibrosis. *J Bacteriol* **178**: 4997-5004.

Skurnik, D., Roux, D., Aschard, H., Cattoir, V., Yoder-Himes, D., Lory, S., and Pier, G.B. (2013)
A comprehensive analysis of in vitro and in vivo genetic fitness of Pseudomonas
aeruginosa using high-throughput sequencing of transposon libraries. *PLoS Pathog* **9**:
e1003582.

Strateva, T., Petrova, G., and Mitov, I. (2010) Antimicrobial activity of tobramycin against

respiratory cystic fibrosis Pseudomonas aeruginosa isolates from Bulgaria. *J Chemother*

22: 378-383.

Turner, K.H., Wessel, A.K., Palmer, G.C., Murray, J.L., and Whiteley, M. (2015) Essential

genome of Pseudomonas aeruginosa in cystic fibrosis sputum. *Proc Natl Acad Sci U S A*

112: 4110-4115.

Vogel, H.J., and Bonner, D.M. (1956) Acetylornithinase of Escherichia coli: partial purification

and some properties. *J Biol Chem* **218**: 97-106.

Wood, L.F., Leech, A.J., and Ohman, D.E. (2006) Cell wall-inhibitory antibiotics activate the

alginate biosynthesis operon in Pseudomonas aeruginosa: Roles of sigma (AlgT) and

the AlgW and Prc proteases. *Mol Microbiol* **62**: 412-426.

Wood, L.F., and Ohman, D.E. (2009) Use of cell wall stress to characterize sigma 22 (AlgT/U)

activation by regulated proteolysis and its regulon in Pseudomonas aeruginosa. *Mol*

Microbiol **72**: 183-201.

Wozniak, D.J., and Ohman, D.E. (1994) Transcriptional analysis of the Pseudomonas

aeruginosa genes algR, algB, and algD reveals a hierarchy of alginate gene expression

which is modulated by algT. *J Bacteriol* **176**: 6007-6014.

Wozniak, D.J., Sprinkle, A.B., and Baynham, P.J. (2003) Control of Pseudomonas aeruginosa

algZ expression by the alternative sigma factor AlgT. *J Bacteriol* **185**: 7297-7300.

Xie, Z.D., Hershberger, C.D., Shankar, S., Ye, R.W., and Chakrabarty, A.M. (1996) Sigma

factor-anti-sigma factor interaction in alginate synthesis: inhibition of AlgT by MucA. *J*

Bacteriol **178**: 4990-4996.

Yin, Y., Withers, T.R., Wang, X., and Yu, H.D. (2013) Evidence for sigma factor competition in

the regulation of alginate production by Pseudomonas aeruginosa. *PLoS One* **8**:

e72329.

Yu, H., Schurr, M.J., and Deretic, V. (1995) Functional equivalence of Escherichia coli sigma E

and Pseudomonas aeruginosa AlgU: E. coli rpoE restores mucoidy and reduces

sensitivity to reactive oxygen intermediates in algU mutants of P. aeruginosa. *J Bacteriol*

177: 3259-3268.

Zhao, H., Roistacher, D.M., and Helmann, J.D. (2019a) Deciphering the essentiality and

function of the anti-sigma(M) factors in Bacillus subtilis. *Mol Microbiol* **112**: 482-497.

Zhao, H., Sachla, A.J., and Helmann, J.D. (2019b) Mutations of the Bacillus subtilis YidC1

(SpoIIIJ) insertase alleviate stress associated with sigmaM-dependent membrane

protein overproduction. *PLoS Genet* **15**: e1008263.