Genetic basis for the essentiality of a bacterial anti-sigma factor involved in virulence

Melissa C. Schofield[1], Daniela Rodriguez[1], Amanda A. Kidman[1], Lia A. Michaels[1], Elizabeth A.

Campbell[2], Peter A. Jorth[3] and Boo Shan Tseng[1]*

[1] School of Life Sciences, University of Nevada Las Vegas, Las Vegas, NV, USA

[2] Rockefeller University, New York, NY, USA

[3] Departments of Pathology, Medicine, and Biomedical Sciences, Cedars-Sinai Medical Center,

Los Angeles, CA, USA

Running title: MucA is required in *P. aeruginosa*

* Corresponding author

Email: boo.tseng@unlv.edu

Keywords: MucA, AlgU, sigma factor competition, cystic fibrosis

ABSTRACT

During decades-long infections in the cystic fibrosis (CF) airway, *Pseudomonas aeruginosa* undergoes selection. One such bacterial genetic adaptation frequently observed in CF isolates is mutation of *mucA.* MucA inhibits the sigma factor AlgU. In clinical isolates, *mucA* mutations lead to AlgU misregulation, resulting in a mucoid phenotype that is associated with poor CF disease outcomes. Here we show that, paradoxically, a portion of the *mucA* gene is essential for *P. aeruginosa* viability. We demonstrate that *mucA* is no longer essential in a strain lacking *algU*, that *mucA* alleles that encode for proteins that do not bind to AlgU are insufficient for bacterial viability, and that *mucA* is no longer essential in mutant strains containing AlgU variants with reduced sigma factor activity. Finally, we found that overexpression of *algU* prevents cell growth in the absence of MucA, and that this phenotype can be rescued by overproduction of RpoD, the housekeeping sigma factor. Together, these results suggest that in the absence of MucA, the inability to regulate AlgU activity may inhibit essential housekeeping functions, resulting in the loss of bacterial viability. Finally, we speculate that essentiality of anti-sigma factors that regulate envelope function may be a widespread phenomenon in bacteria**.**

INTRODUCTION

The major cause of death in people with cystic fibrosis (CF), a human autosomal recessive

genetic disease, is respiratory failure due to chronic lung infection. *Pseudomonas aeruginosa* is

one of the most prevalent respiratory pathogens of CF patients (Cystic Fibrosis Foundation

Patient Registry, 2019). The CF lung environment selects for mucoid *P. aeruginosa* mutants,

which overproduce the exopolysaccharide alginate and are associated with poor disease

prognosis (Emerson *et al.*, 2002, Henry *et al.*, 1992, Li *et al.*, 2005, Parad *et al.*, 1999, Pedersen

et al., 1992). Conversion to the mucoid phenotype in clinical *P. aeruginosa* isolates, which is

thought to be advantageous for chronic infection, is frequently caused by *mucA* mutations

(Martin *et al.*, 1993).

MucA is an anti-sigma factor to the alternative sigma factor AlgU (also known as AlgT, σ^E, or

σ^{22}), which responds to envelope stress (Damron & Goldberg, 2012, Govan & Deretic, 1996).

MucA is a transmembrane protein that sequesters the sigma factor away from RNA polymerase

(RNAP) via its N-terminal interaction with AlgU (Li *et al.*, 2019, Schurr *et al.*, 1996, Xie *et al.*,

1996). The C-terminus of MucA is in the periplasm, where it is protected from proteolysis via an

interaction with MucB (Mathee *et al.*, 1997, Schurr *et al.*, 1996). The envelope stress response

is controlled via a regulated intramembrane proteolysis cascade: MucB dissociates from MucA

upon detection of stress, allowing the proteases AlgW and MucP to cleave MucA from the inner

membrane (Qiu *et al.*, 2007, Wood & Ohman, 2009). In the cytoplasm, MucA is further

degraded by ClpXP, releasing AlgU to interact with RNAP and activate the AlgU regulon (Qiu *et*

al., 2008). AlgU regulates at least 350 genes, including those responsible for the production of

itself, its cognate anti-sigma factor MucA, and the alginate biosynthetic enzymes (Firoved &

Deretic, 2003, Schulz *et al.*, 2015, Wood & Ohman, 2009). This system is homologous to the

well-studied envelope stress response in *Escherichia coli*: MucA shares 28% identity (72%

similarity) to the anti-sigma factor RseA, and the cognate sigma factor of the *E. coli* RseA, called

RpoE, is homologous to and functionally interchangeable with AlgU (Yu *et al.*, 1995), with 66%

identity (93% similarity).

For several reasons, *mucA* is assumed to be dispensable for *P. aeruginosa* viability. First, after

being released from the cell membrane, MucA is presumed to be fully degraded in the

cytoplasm by ClpXP (Qiu *et al.*, 2008). Second, *mucA* mutations commonly arise in CF clinical

isolates (Boucher *et al.*, 1997, Candido Cacador *et al.*, 2018, Ciofu *et al.*, 2008, Martin *et al.*,

1993, Pulcrano *et al.*, 2012), and many laboratory *mucA* mutants exist in the literature

(Gallagher *et al.*, 2011, Liberati *et al.*, 2006, Skurnik *et al.*, 2013, Turner *et al.*, 2015). Third,

there are three published strains in which the entirety of *mucA* is removed from the genome

(Intile *et al.*, 2014, Jones *et al.*, 2010, Pritchett *et al.*, 2015). Paradoxically, three different whole-

genome studies of essential genes in two different strains of *P. aeruginosa* identified *mucA* as

an essential gene, based on the low observed frequency of transposon insertions in this gene

(Lee *et al.*, 2015, Liberati *et al.*, 2006, Skurnik *et al.*, 2013). To investigate this paradox, we

systematically attempted to delete *mucA* from *P. aeruginosa* using allelic exchange. Our results

show that a portion of *mucA* is required for bacterial viability in multiple *P. aeruginosa* strains.

Furthermore, our work shows that *mucA* was no longer essential in a strain lacking *algU*, and

that *mucA* alleles that encode for proteins that do not interact with AlgU were insufficient to

rescue viability or led to a growth defect. We found that *algU* mutations, which encode for a less

active sigma factor, can relieve *mucA* essentiality, and that overproduction of AlgU in the

absence of MucA is lethal. Interestingly, our works suggests that *mucA* essentiality can be

suppressed by increasing the levels of the housekeeping sigma factor RpoD. Together, our

results strongly suggest that the unregulated activity of AlgU itself, in the absence of MucA,

leads to bacterial cell death, which is in part due to sigma factor competition with RpoD.

RESULTS

***mucA* is essential for viability in a diverse set of *P. aeruginosa* isolates**

To determine if *mucA* is essential, we attempted to delete the gene from various *P. aeruginosa*

strains using a modified allelic exchange protocol to turn it into a robust assay (Fig S1). We

introduced a deletion allele of *mucA*, which is missing >95% of the coding region, into *P.*

aeruginosa and selected for merodiploids. Via PCR, we confirmed that isolates contained both

the endogenous and deletion alleles of *mucA* in the genome. Using six confirmed merodiploids,

we then counter-selected for loss of one of the two alleles. Isolates were then tested via PCR to

determine which *mucA* allele the isolate resolved to. If a gene is non-essential, isolates that

resolved to either the wild-type or deletion allele should be observed. However, if a gene is

essential, then only the cells that resolved to the wild-type allele should be isolated, as cells that

resolved to the deletion allele cannot survive due to the absence of the gene. To give statistical

power to our assay, we performed three biological replicates for strains from which we were

unable to delete *mucA*, for a minimum total of 125 isolates screened. We performed this assay

on a diverse set of wild-type *P. aeruginosa* strains, including four common laboratory isolates as

well as seven clinical and environmental isolates (Table 1). These strains are not only isolated

from diverse locations, but they also vary in their colony morphology and their

exopolysaccharide production profile (Colvin *et al.*, 2012). For all strains tested, all observed

isolates resolved to the wild-type *mucA* allele ($p < 0.0001$, Fisher's exact test), strongly

suggesting that *mucA* is essential in a variety of wild-type *P. aeruginosa* strains. Since all tested

strains required *mucA* for viability, we continued our experiments using the laboratory isolate

PAO1 as a representative strain.

Alginate biosynthesis is not solely responsible for *mucA* essentiality

Clinical isolates containing *mucA* mutations are mucoid due to their overproduction of alginate

(Martin *et al.*, 1993). To determine if alginate overproduction is responsible for *mucA*

essentiality, we attempted to delete *mucA* from a strain lacking *algD*, a key alginate biosynthesis

gene (Deretic *et al.*, 1987). Using our allelic exchange assay, we were unable to delete *mucA* in

this background (Table S1).

Expression of the alginate biosynthesis genes is controlled by three AlgU-regulated transcription

factors that are active in mucoid cells: AlgB, AlgR, and AmrZ (Martin *et al.*, 1994, Wozniak &

Ohman, 1994, Wozniak *et al.*, 2003). Since these transcription factors regulate a suite of genes

in addition to those involved in alginate biosynthesis (Huang *et al.*, 2019, Jones *et al.*, 2014,

Kong *et al.*, 2015, Leech *et al.*, 2008), we tested if overexpression of these three regulons

underlies *mucA* essentiality by attempting to delete *mucA* from strains lacking these

transcription factors, either individually or all together. We found that we could not delete *mucA*

in any of these backgrounds (Table S1). Altogether, our data support the conclusion that

eliminating alginate biosynthesis and the expression of large parts of the AlgU regulon do not

alleviate *mucA* essentiality.

The first 50 amino acids of MucA are necessary and sufficient for cell viability

While we were unable to delete *mucA* using an allele lacking >95% of the coding region, *mucA*

is commonly mutated in clinical isolates and many *mucA* transposon mutants exist (Fig S2).

These data suggest that only a portion of *mucA* is essential. To determine the essential portion

of *mucA*, we first constructed a series of strains containing an ectopic chromosomally-integrated

mucA, driven by its native promoter. We then attempted to delete the endogenous wild-type

mucA from these strains. Although we designed these ectopic alleles to encode for a protein

with a His-tag, we were unable to probe whether equivalent amounts of protein were produced

across our strains, since truncated versions of MucA are cleaved in the cytosol, as shown in

previous work (Qiu *et al.*, 2008). However, we expect that the ectopic alleles produce similar

protein levels, as they are all under the same promoter and located in the same genomic site.

Since we expected that the resultant Δ*mucA* strain would be mucoid in some cases, we used a

Δ*algD* background to eliminate alginate production, which made the strains easier to

manipulate.

Mutant MucA proteins of varying lengths were ectopically expressed to determine the necessary

portion of MucA. Full-length MucA is 194 amino acids (aa). Based on the co-crystal structures of

MucA with AlgU and MucB (Li *et al.*, 2019), the first 78 residues of MucA interact with AlgU, and

the last 48 residues of MucA interact with MucB. We engineered *mucA* alleles that encode for

various truncated proteins (Fig 1A). We then attempted to delete the endogenous *mucA* from

these strains. Since the flanking regions of these two alleles differ, our deletion allele specifically

targets the native allele. We were able to easily recover isolates that resolved to the *mucA*

deletion allele from strains containing ectopic alleles that encode for the full length MucA (aa 1-

194), as well as MucA aa 1-143, aa 1-110, aa 1-75, and aa 1-62 (Fig 1B). These results show

that the requirement for *mucA* can be complemented *in trans*, and suggest that the gene

product of *mucA* is required for viability, irrespective of its genomic location.

Interestingly, while we were able to delete the endogenous *mucA* from strains containing an

ectopic *mucA* that encoded for aa 1-50 (Fig 1B), isolates that resolved to the deletion allele

grew up more slowly than those that resolved to the wild-type allele. This suggests that while

this allele is sufficient for viability, it is not well tolerated by the cells. All tested isolates of strains

carrying *mucA* alleles encoding shorter products (aa 1-40 and aa 1-24) resolved to the wild-type

allele. These results show that while an allele encoding the first 50 residues of MucA is sufficient

for cell viability, alleles encoding for shorter lengths are not.

To determine if the first 50 residues of MucA are necessary for viability, we constructed a strain

with an ectopic *mucA* that encodes for aa 51-194. We were unable to delete the native *mucA*

from this strain (Fig 1B). Together, these results suggest that the first 50 amino acids of MucA

are necessary and sufficient for cell viability in *P. aeruginosa*. Consistent with these results,

while many clinical and engineered *mucA* mutations have been reported (Boucher *et al.*, 1997,

Candido Cacador *et al.*, 2018, Ciofu *et al.*, 2008, Martin *et al.*, 1993, Pulcrano *et al.*, 2012,

Turner *et al.*, 2015), no mutations fall within the region of *mucA* that encodes for the first 50

residues (Fig S2). Furthermore, this region of MucA corresponds to the AlgU binding domain,

suggesting that the physical interaction between the two proteins is important for cell viability.

Depletion of MucA leads to cell death

To evaluate the effect of the nutrient conditions on *mucA* essentiality, we tested the effect of

depleting MucA on *P. aeruginosa* cell viability in different media. To deplete MucA from cells, we

engineered a strain lacking the native *mucA* and containing a rhamnose-inducible copy (Δ*mucA*

*att*Tn7::P$_{rhaBAD}$-*mucA*). We removed rhamnose from growing Δ*mucA* *att*Tn7::P$_{rhaBAD}$-*mucA* cells

and determined the viability of cells over time based on their ability to recover on medium

containing rhamnose. Since cells would cease producing MucA in the absence of rhamnose, the

MucA present in the cells at the start of the experiment would deplete as the cells divided. In all

four media we tested, if rhamnose was not removed at the beginning of the experiment, the

cells increased in density by ~2-logs over the duration of the experiment. In comparison, when

rhamnose was removed, the Δ*mucA* *att*Tn7::P$_{rhaBAD}$-*mucA* cells lost viability over time in all four

media (Fig 2). Together, these results suggest that *mucA* is essential in *P. aeruginosa*

independent of the nutrient environment.

The interaction of MucA with AlgU is required for cell survival

Since the only described function for MucA is to inhibit AlgU, we hypothesized that *mucA*

essentiality is rooted in its regulation of AlgU. In such a case, *mucA* essentiality should require

algU. Therefore, we attempted to delete *mucA* from a strain lacking *algU* (Δ*algU*). We found that

we were able to delete *mucA* in the absence of *algU*, with 16 of the tested isolates resolving to

the deletion allele out of the 48 colonies tested (see Fig 4A), showing that *mucA* essentiality is

algU-dependent.

Because our data suggest that the AlgU-binding domain of MucA is required for viability (Fig

1B), we hypothesized that the physical interaction between MucA and AlgU is necessary for cell

survival. In the co-crystal structure of MucA and AlgU (Li *et al.*, 2019), four residues in MucA

make more than one hydrogen bond with AlgU: D15, E22, R42, and E72 (Fig 3A). Although E72

likely contributes to the binding energy between these two proteins, it is unlikely that E72 is

critical to the MucA-AlgU interaction, since the first 50 amino acids of MucA are sufficient for

viability (Fig 1B). Therefore, we engineered alleles that encode for MucA D15A, E22A, or R42A

to maximally affect the hydrogen bonding while limiting the effects on overall protein structure.

We used MucA aa 1-75 as a base because the co-crystal structure includes the first 78 residues

of MucA. We tested the effect of these substitutions on the MucA-AlgU interaction via a yeast-

two hybrid assay. Using beta-galactosidase activity as a proxy for their interaction, wild-type

MucA aa 1-75 strongly interacted with AlgU (Fig 3B). This interaction was dependent on the

presence of both MucA and AlgU, and the effect was not directional (Fig S3). In comparison, the

D15A, E22A, and R42A MucA mutants failed to interact with AlgU (Fig 3B). To determine the

effects of these mutations on viability, we engineered strains carrying an ectopic *mucA* encoding

these substitutions and then attempted to delete the native *mucA*. As described above, an allele

encoding MucA aa 1-75 was sufficient for viability (Fig 1B). However, we were unable to delete

the endogenous *mucA* from strains carrying alleles encoding the D15A and R42A substitutions,

suggesting that the mutant protein is not sufficient for viability (Fig 3C). For the strain carrying a

mucA allele encoding the E22A substitution, we were able to delete the endogenous *mucA*.

However, isolates that resolved to the deletion allele in this strain grew up slower than those

that resolved to the wild-type allele. Similar to the *mucA* allele encoding only the first 50 amino

acids, these results suggest that while MucA E22A is sufficient for viability, the cells do not

tolerate it well.

Mutations that reduce AlgU activity alleviate the requirement for MucA

Since MucA inhibits AlgU from binding to RNAP, we hypothesized that *mucA* essentiality is due

to its inhibition of the AlgU regulon, which includes *algU* itself. To start to test this hypothesis,

we reasoned that strains containing a mutant AlgU with lower affinity for DNA would alleviate

mucA essentiality, since expression of the AlgU regulon would inherently be lower in such

mutants. Therefore, we engineered Δ*algU* strains with an ectopic chromosomally-integrated

algU allele encoding such DNA-binding mutants. Based on homology to *E. coli* RpoE mutants

that have decreased *in vitro* transcriptional activity (Campagne *et al.*, 2014), we engineered

algU alleles that encode for K57A and N81A. As described above, we were able to delete *mucA*

from a Δ*algU* strain (Fig 4A). This phenotype could be rescued via the ectopic addition of a wild-

type *algU* allele, as we were no longer able to delete *mucA* from such a strain. In contrast,

mucA could be readily deleted from strains carrying alleles encoding AlgU K57A or N81A,

showing that such mutant alleles failed to rescue the Δ*algU* phenotype (Fig 4A). To determine

the effect of these substitutions on AlgU activity, we induced envelope stress using D-

cycloserine (Wood *et al.*, 2006) and measured AlgU activity using a plasmid-borne *gfp* reporter

driven under the AlgU-regulated *algD* promoter (Damron *et al.*, 2009). Similar to what is seen for

RpoE (Campagne *et al.*, 2014), these mutant strains had reduced AlgU activity upon induction

of envelope stress, relative to the strain containing a wild-type *algU* (Fig S4A,B). These results

suggest that AlgU mutants with reduced sigma factor activity can suppress *mucA* essentiality.

As mentioned above, others have reported strains in which the entirety of *mucA* is deleted from

P. aeruginosa PAO1, PAK, and PA103 (Intile *et al.*, 2014, Jones *et al.*, 2010, Pritchett *et al.*,

2015). Since we were unable to delete *mucA* from these three strain backgrounds (Table 1), we

hypothesized that these Δ*mucA* strains may contain suppressor mutations that allowed for their

survival in the absence of *mucA*. To identify these suppressors, we sequenced the genomes of

these Δ*mucA* isolates. In the PAO1 Δ*mucA* strain (Pritchett *et al.*, 2015), we found that while the

native *mucA* was deleted, the strain contained a second full-length copy of *algU* and a *mucA*

allele that encoded for MucA aa 1-155 elsewhere in the genome. As expected from our data

(Fig 1B), we were able to delete the native *mucA* allele from a PAO1 strain containing an

ectopic *mucA* allele that encoded for MucA aa 1-155 (with 5 of the 28 tested isolates resolving

to the deletion allele), confirming that the ectopic *mucA* allele in the published PAO1 Δ*mucA*

strain is sufficient for viability in the absence of the endogenous *mucA*. In comparison, the PAK

and PA103 Δ*mucA* strains (Intile *et al.*, 2014, Jones *et al.*, 2010) contain missense mutations in

algU, which results in AlgU A58T and E46G, respectively. We replicated these mutations in

PAO1 by inserting *algU* alleles encoding these substitutions in a Δ*algU* strain. We were able to

delete *mucA* from these strains (Fig 4A). These results confirm that the *algU* alleles in the

published PAK and PA103 Δ*mucA* strains suppress *mucA* essentiality. To examine the activity

of these mutant proteins, we used our reporter assay. Similar to what we saw with the AlgU

DNA-binding mutants, we found that strains carrying these AlgU substitutions had reduced

sigma factor activity relative to that of strains carrying wild-type AlgU (Fig S4B). We note that

our reporter assay does not appear to be very sensitive to low levels of AlgU activity. Both the

published PAK and PA103 Δ*mucA* strains are mucoid (Intile *et al.*, 2014, Jones *et al.*, 2010),

suggesting that AlgU A58T and E46G are not completely inactive. While we cannot distinguish

this low level of activity from the null case, our results show that nevertheless these AlgU

mutants have significantly less transcriptional activity than the wild-type protein (Fig S4A,B).

To identify additional suppressors of *mucA* essentiality, we used the Δ*mucA att*Tn7::P$_{rhaBAD}$-

mucA strain. This strain was not viable when grown in the absence of rhamnose, but natural

revertants arose at a frequency of less than 1 in 10^9 colony forming units. We sequenced the

algU gene in 25 of these revertants, since our work shows that AlgU mutants can suppress

mucA essentiality (Fig 4A). All 25 isolates contained mutations in *algU* that are expected to

encode for sigma factors with reduced activity (Fig 4B, Table S2). There were 10 revertants with

deletions or nonsense mutations of *algU*, encoding either no product or a truncated product

completely lacking Region 4 of the sigma factor. There were 6 revertants that contained multi-

base pair duplications in *algU* that would lead to the insertion of 3 or 4 amino acids in Region 2

helix 3 of the sigma factor. There were 9 revertants containing missense mutations, 8 of which

were unique, encoding the following substitutions: D18G, A21V, Y29C, A47T, D49G, Y59C,

N81D, and R174G. Using a model of σ^E in complex with the RNAP core and the promoter

element, we expect these insertions and substitutions to affect sigma factor folding, RNAP core

interactions, or promoter element interactions (Fig S5). To determine the effect of these 25

mutations on AlgU activity, we then tested AlgU function using our *algD* reporter assay in these

natural revertants. As expected, we saw that all 25 revertants had much lower AlgU activity than

the parental strain upon induction of envelope stress with D-cycloserine (Fig S4C). Together,

these data strongly suggest that *algU* mutations that encode for a protein with reduced

transcriptional activity allow *P. aeruginosa* to survive in the absence of *mucA*.

Overexpression of *algU* in the absence of *mucA* is lethal

Collectively, our results suggest that *mucA* essentiality is due to unregulated AlgU activity. We

therefore reasoned that overexpression of *algU* should be lethal. Supporting this hypothesis, the

toxic effects of *algU* overexpression have been previously noted (Cross *et al.*, 2020,

Hershberger *et al.*, 1995, Schurr *et al.*, 1994). However, studies have shown that *algU* can be

overexpressed in wild-type *P. aeruginosa* (Qiu *et al.*, 2008, Schulz *et al.*, 2015). It is important to

note that these wild-type strains contain *mucA*, which is positively regulated by AlgU. We

therefore examined the effect of overproducing AlgU in WT, Δ*algU*, and Δ*algU* Δ*mucA* strains,

using a chromosomally integrated arabinose-inducible *algU*. Of note, the Δ*algU* and Δ*algU*

Δ*mucA* strains lack the positive feedback of AlgU on its own expression, and the expression of

algU is completely dependent on the inducer. Our results show that in the absence of inducer

when AlgU is not overexpressed, all three strains had similar growth rates (Fig 5A). In the

presence of 1% arabinose (i.e. high expression of *algU*), all three strains had a growth defect.

However, in comparison to the other two strains, the strain lacking *mucA* failed to grow at all in

the presence of this high inducer concentration.

To determine if the Δ*algU* Δ*mucA att*Tn7::P$_{araBAD}$-*algU* strain can grow under lower levels of *algU*

induction, we tested a range of inducer concentrations (Fig 5B, Table S3). We found that

although there was a growth defect, Δ*algU* Δ*mucA att*Tn7::P$_{araBAD}$-*algU* was able to grow in the

presence of 0.1% arabinose, suggesting that this strain was able to tolerate low levels of *algU*

induction. Furthermore, as expected, the drop in growth rate in comparison to the no arabinose

condition was statistically larger for the Δ*algU* Δ*mucA att*Tn7::P$_{araBAD}$-*algU* than for the other two

strains (N=3, $p < 0.05$, ANOVA with post hoc Tukey HSD), since this strain lacks the ability to

produce any MucA to reduce AlgU activity. Together, these results strongly suggest that in the

absence of *mucA*, while a certain level of AlgU activity is tolerated, high AlgU activity is fatal to

the cell.

The above experiments were performed using only one medium (LB). To determine if

overexpression of *algU* in the absence of *mucA* is lethal under other nutrient conditions, we

determined the growth rate of the Δ*algU* Δ*mucA att*Tn7::P$_{araBAD}$-*algU* strain in other media.

Similar to the results in LB (Fig 5), Δ*algU* Δ*mucA att*Tn7::P$_{araBAD}$-*algU* strain failed to grow in the

presence of 1% arabinose for the three other media we tested (Fig S6), suggesting that AlgU

overproduction is toxic independent of the nutrient conditions.

Expression of *rpoD* can rescue lethality of *algU* overexpression.

It has been suggested that AlgU competes for RNAP binding with RpoD, the essential primary

sigma factor (Yin *et al.*, 2013). If high concentrations of AlgU outcompetes RpoD for RNAP, this

could reduce the expression of essential housekeeping genes and lead to toxicity in AlgU-

overexpressing cells. If this is true, we hypothesized that overexpressing *rpoD* with *algU* could

ameliorate the growth defect of high *algU* expression in the Δ*algU* Δ*mucA att*Tn7::P*araBAD*-*algU*

strain. We inserted into this strain a copy of *rpoD* under the same arabinose-inducible promoter.

We chose to test growth in the presence of 2% arabinose, since this new strain contains two

arabinose-inducible promoters, in contrast to the strains described above in Fig 5. Growth with

2% arabinose caused a growth defect for all strains tested, even in a strain lacking an ectopic

arabinose-inducible gene (Fig 6). As expected from our results (Fig 5), Δ*algU* Δ*mucA*

*att*Tn7::P*araBAD*-*algU* failed to grow in the presence of 2% arabinose. This lethality defect,

however, was rescued in the Δ*algU* Δ*mucA att*Tn7::P*araBAD*-*algU att*CTX::P*araBAD*-*rpoD* strain.

Furthermore, the growth rate of this strain in the presence of arabinose was indistinguishable

from that of Δ*algU* Δmuc*A* and Δ*algU* Δ*mucA att*CTX::P*araBAD*-*rpoD* (Table S4). Since increasing

RpoD levels rescued the lethality associated with high AlgU levels, these results suggest that in

the absence of MucA, the ensuing sigma factor competition of AlgU with RpoD for RNAP

contributes to the viability defect, and that MucA may be required to rein in AlgU in order to

reduce the interference with the essential function of RpoD.

DISCUSSION

Although it has been assumed that *mucA* can be deleted without affecting bacterial viability, our

work shows that *mucA* is essential in a variety of *P. aeruginosa* wild-type strains and under

various nutrient conditions. Our results strongly suggest that the interaction of MucA with its

cognate sigma factor AlgU is required for viability, and that AlgU mutants with reduced sigma

factor activity can suppress *mucA* essentiality. Furthermore, overproduction of AlgU is

detrimental to cell growth, which can be rescued by increasing RpoD production. Together, our

data strongly suggest that unchecked AlgU activity in the absence of MucA leads to cell death.

Our data point to the negative regulation of AlgU being necessary for cell survival in *P.*

aeruginosa (Fig 7). Under non-stress conditions, AlgU is bound to and inhibited by a full-length

MucA. Under envelope stress or in strains containing *mucA* mutations, the anti-sigma factor is

cleaved. We propose that while this cleaved cytosolic form of MucA does not inhibit AlgU to the

same extent as the full-length protein, it is still able to interact with and inhibit AlgU to some

degree. Although AlgU is active under such conditions, because *mucA* is positively regulated by

AlgU, this negative feedback allows the cell to keep AlgU activity under control. In comparison,

in the absence of MucA, this ability to control AlgU is lost. We propose that the strong positive

feedback of AlgU on its own expression is what leads to cell death and *mucA* essentiality.

Supporting this model are several lines of evidence. First, *mucA* essentiality is rooted in its

interaction with *algU* (Figs 1 and 3), strongly suggesting that the ability of MucA to inhibit AlgU is

required for viability. Second, deletion of AlgU-regulated genes, including those encoding the

well-characterized transcription factors AlgB (Wozniak & Ohman, 1994), AlgR (Martin *et al.*,

1994, Wozniak & Ohman, 1994), and AmrZ (Wozniak *et al.*, 2003), do not allow for *mucA*

deletion (Table S1), supporting that *algU* itself is the lethal component of the AlgU regulon.

Third, AlgU mutants can suppress *mucA* essentiality, and all 25 natural revertants that we

screened contained mutations in *algU* (Fig 4). The strains carrying these mutations had

decreased AlgU transcriptional activity (Fig S4), supporting the idea that decreasing the positive

feedback of AlgU on its own expression allows the cell to survive in the absence of MucA.

Lastly, overexpression of *algU* leads to a growth defect, which is lethal at high levels in the

absence of *mucA* (Fig 5) and can be rescued by overexpression of *rpoD* (Fig 6), suggesting that

sigma factor competition between AlgU and RpoD may be partially responsible for lethality in

the absence of MucA.

While our data supports our model that the AlgU-regulated gene critical for *mucA* essentiality is

algU itself, we cannot exclude the possibility that the regulation of another single gene or a

combination of genes within the AlgU regulon is responsible for the *mucA* requirement. This

possibility, however, seems less likely for two reasons. First, all the revertants we tested that

could grow in the absence of *mucA* expression contained *algU* mutations (Table S2). If another

single AlgU-regulated gene were responsible for *mucA* essentiality, we would have expected to

find a revertant with such a mutation. Second, we were unable to delete *mucA* from a strain

lacking three AlgU-regulated transcription factors (Table S1). Genes regulated by AlgB, AlgR, or

AmrZ can account for approximately fifty percent of the AlgU regulon (Huang *et al.*, 2019, Jones

et al., 2014, Leech *et al.*, 2008, Schulz *et al.*, 2015), suggesting that it is unlikely that deleting a

combination of AlgU-regulated genes can alleviate the necessity for *mucA*. Nonetheless, we are

actively searching for non-*algU* suppressors of *mucA* essentiality.

Our data show that the anti-sigma factor MucA is required for viability in various *P. aeruginosa*

strains (Table 1). However, the requirement of an anti-sigma factor for bacterial viability is not

unique to *P. aeruginosa*. While there do not appear to be any essential anti-sigma factors in *E.*

coli, as all of them are individually deleted in the Keio collection (Baba *et al.*, 2006), there are

many occurrences in other organisms. There are two anti-sigma factors, YhdL and YxlC, in

Bacillus subtilis that are necessary for bacterial viability (Horsburgh & Moir, 1999, Koo *et al.*,

2017, Mendez *et al.*, 2012). The essentiality of *yhdL* is notable as this anti-sigma factor

negatively regulates SigM, a sigma factor involved in the envelope stress response (Horsburgh

& Moir, 1999). Similarly, in *Mycobacterium tuberculosis*, the gene encoding the anti-sigma factor

RslA is identified as essential due to an underrepresentation of interruptions in a genome-wide

transposon insertional analysis (Griffin *et al.*, 2011). However, similar to our results (Fig 4), *rslA*

can be deleted from *M. tuberculosis* lacking the gene encoding its cognate sigma factor SigL,

which regulates genes involved in cell envelope processes (Dainese *et al.*, 2006). Furthermore,

in the defined transposon mutant library for *Vibrio cholerae*, an interruption in the *mucA*

homolog *rseA* does not exist, suggesting that this anti-sigma factor may be essential in *V.*

cholerae, similar to in *P. aeruginosa* (Cameron *et al.*, 2008). Finally, the *Pseudomonas syringae*

mucA homolog is deemed essential based on a reduced number of transposon insertions

(Helmann *et al.*, 2019). Therefore, although the *E. coli mucA* homolog *rseA* is not required for

cell viability (De Las Penas *et al.*, 1997, Missiakas *et al.*, 1997), these data, together with our

results, suggest that essentiality of anti-sigma factors that regulate cell envelope function may

be a widespread phenomenon in bacteria.

Our work strongly suggests that lethality in the absence of *mucA* is caused by unchecked AlgU

activity (Figs 5 and 6), in agreement with previous studies suggesting that overproduction of

AlgU is toxic (Cross *et al.*, 2020, Hershberger *et al.*, 1995, Schurr *et al.*, 1994). Our data

suggests that sigma factor competition may play a role in this lethality (Fig 6). Competition

between AlgU and the primary sigma factor RpoD has been previously suggested (Yin *et al.*,

2013). Overexpression of RpoD suppresses alginate biosynthesis in a mucoid CF isolate.

Additionally, the activity of AlgU increases in a mutant with decreased RpoD activity (Yin *et al.*,

2013). Expanding to other organisms, in *B. subtilis*, the lethality defect associated with deleting

yhdL, which encodes for an anti-sigma factor, can be rescued via overproduction of the primary

sigma factor SigA, similar to our results (Fig 6). Zhao and colleagues suggest that the lethality of

this strain is due to the YhdL-cognate sigma factor, SigM, outcompeting SigA for RNAP (Zhao *et*

al., 2019). Based on these findings, and our data showing that RpoD can rescue the lethality of

high AlgU levels (Fig 6), we speculate that sigma factor competition may be involved in other

instances where anti-sigma factors that regulate cell envelope functions have been found to be

essential.

The interaction of MucA and AlgU is of interest, as *mucA* mutations are common in clinical CF

isolates of *P. aeruginosa* (Boucher *et al.*, 1997, Candido Cacador *et al.*, 2018, Ciofu *et al.*, 2008,

Martin *et al.*, 1993, Pulcrano *et al.*, 2012). Our results (Fig 4) agree with the literature showing

that mucoid isolates with *mucA* mutations revert to a non-mucoid state via changes to *algU*

when the cells are grown under standard laboratory conditions (Ciofu *et al.*, 2008, DeVries &

Ohman, 1994, Sautter *et al.*, 2012, Schurr *et al.*, 1994). While these *algU* secondary site

mutations are found in non-mucoid CF isolates with *mucA* mutations, such isolates are detected

at a lower frequency (Ciofu *et al.*, 2008). As suggested by Ciofu and associates, this may be

due to the importance of AlgU for the survival of *P. aeruginosa* in the CF lung environment,

suggesting that reducing AlgU activity may increase *P. aeruginosa* eradication from the CF

airway. Based on our data and the published literature, we propose that the MucA-AlgU

interaction may serve as a good therapeutic target. Our results show that destabilizing the

MucA-AlgU interaction results in bacterial cell death or mutations in *algU* that likely result in

reduced sigma factor activity and reduced mucoid conversion, of which either outcome could be

beneficial in the treatment of *P. aeruginosa* CF lung infections.

METHODS AND MATERIALS

Bacterial strains and growth conditions. The bacterial strains, plasmids, and oligonucleotides

used for this study are in Tables S5-S7. The construction of strains is described in Supporting

Information. Bacteria were grown at 37°C in LB with shaking or on semi-solid LB media, unless

otherwise noted.

Allelic exchange assay. The protocol, depicted in Fig S1, was modified from (Hmelo *et al.*,

2015). Briefly, after introduction of the *mucA* deletion vector, merodiploids were selected on

semi-solid VBMM (Vogel & Bonner, 1956) with 60 mg/L gentamicin. Using OMS118 and

OMS119, PCR was performed on at least six isolates to confirm that the presence of both the

wild type and deletion alleles of *mucA*. Confirmed merodiploids were then individually streaked

on NSLB (10 g/L tryptone, 5 g/L yeast extract) with 10% sucrose semi-solid media for

counterselection. PCR was performed on eight colonies per merodiploid, using OBT601 and

OBT602 to determine the resolution to either the WT or deletion allele.

MucA depletion assay. Cells were grown in LB with 0.05% rhamnose at 37°C with shaking to

an OD_{600} of 0.3. After washing, the culture was then divided in two, half resuspended with

rhamnose and half without in its respective media base (LB, PIB, SCFM, or VBMM). Cells were

incubated at 37°C with shaking. At the indicated time points, two aliquots were removed from

each culture, serially diluted and plated in triplicate onto LB agar plates with 0.05% rhamnose

for recovery. Colonies were counted and log_{10}-transformed CFU/mL of the culture was

calculated.

Yeast two-hybrid assay. The ProQuest Two-Hybrid System (Invitrogen) was used, per

manufacturer's instructions. Briefly, yeast containing the Gal4 activation domain-based prey and

the Gal4 DNA binding domain-based bait vectors were grown overnight in SD-Leu-Trp broth

(Clontech). The OD_{600} value was recorded. ONPG (VWR) was added to lysed cells, and the

mixture was incubated at 37°C until a light yellow color was achieved. The incubation time was

recorded. The OD_{420} of the supernatants was determined using a Synergy Hybrid HTX

Microplate Reader (Bio-Tek Instruments). Beta galactosidase activity was determined using

Miller units, based on the following equation: 1000 x OD_{420}/(time x culture volume x OD_{600}).

Natural revertant assay. *P. aeruginosa* Δ*mucA att*Tn7::P*rhaBAD*-*mucA* was grown in LB with

0.05% rhamnose at 37°C with shaking to an OD_{600} of 1. Cells were washed and plated on semi-

solid LB plates without rhamnose. Plates were incubated for 24-48 h at 37°C. For isolates that

grew on LB without rhamnose, *algU* was Sanger sequenced.

ACKNOWLEGDEMENTS

The authors would like to thank Cai Tao and Maria Alexandra Ledesma for technical assistance;

and Joe J. Harrison for critical discussions of this work. MCS, AAK, DR, and BT are supported

by the Cystic Fibrosis Foundation (TSENG19I0). MCS is also supported by a University of

Nevada Las Vegas doctoral graduate research assistantship. EAC and PAJ are supported by

the National Institutes of Health (R01GM114450 and K22AI127473, respectively). PAJ is also

supported by the Cystic Fibrosis Foundation (JORTH17F5).

AUTHOR CONTRIBUTIONS

Conceptualization: MCS, BST; Acquisition, analysis, or interpretation of data: MCS, DR, AAK,

LAM, EAC, PAJ, BST; Writing – original draft: MCS, BST; and Writing – review and editing:

MCS, DR, LAM, EAC, PAJ, BST.

REFERENCES

Baba, T., Ara, T., Hasegawa, M., Takai, Y., Okumura, Y., Baba, M., Datsenko, K.A., Tomita, M.,

Wanner, B.L., and Mori, H. (2006) Construction of Escherichia coli K-12 in-frame, single-

gene knockout mutants: the Keio collection. *Mol Syst Biol* **2**: 2006 0008.

Boucher, J.C., Yu, H., Mudd, M.H., and Deretic, V. (1997) Mucoid Pseudomonas aeruginosa in

cystic fibrosis: characterization of muc mutations in clinical isolates and analysis of

clearance in a mouse model of respiratory infection. *Infect Immun* **65**: 3838-3846.

Cameron, D.E., Urbach, J.M., and Mekalanos, J.J. (2008) A defined transposon mutant library

and its use in identifying motility genes in Vibrio cholerae. *Proc Natl Acad Sci U S A* **105**:

8736-8741.

Campagne, S., Marsh, M.E., Capitani, G., Vorholt, J.A., and Allain, F.H. (2014) Structural basis

for -10 promoter element melting by environmentally induced sigma factors. *Nat Struct*

Mol Biol **21**: 269-276.

Candido Cacador, N., Paulino da Costa Capizzani, C., Gomes Monteiro Marin Torres, L.A.,

Galetti, R., Ciofu, O., da Costa Darini, A.L., and Hoiby, N. (2018) Adaptation of

Pseudomonas aeruginosa to the chronic phenotype by mutations in the algTmucABD

operon in isolates from Brazilian cystic fibrosis patients. *PLoS One* **13**: e0208013.

Ciofu, O., Lee, B., Johannesson, M., Hermansen, N.O., Meyer, P., and Hoiby, N. (2008)

Investigation of the algT operon sequence in mucoid and non-mucoid Pseudomonas

aeruginosa isolates from 115 Scandinavian patients with cystic fibrosis and in 88 in vitro

non-mucoid revertants. *Microbiology* **154**: 103-113.

Colvin, K.M., Irie, Y., Tart, C.S., Urbano, R., Whitney, J.C., Ryder, C., Howell, P.L., Wozniak,

D.J., and Parsek, M.R. (2012) The Pel and Psl polysaccharides provide Pseudomonas

aeruginosa structural redundancy within the biofilm matrix. *Environ Microbiol* **14**: 1913-

1928.

Cross, A.R., Raghuram, V., Wang, Z., Dey, D., and Goldberg, J.B. (2020) Overproduction of the

AlgT sigma factor is lethal to mucoid Pseudomonas aeruginosa. *Journal of Bacteriology*.

Cystic Fibrosis Foundation Patient Registry, (2019) 2018 Annual Data Report. In*.* Bethesda,

MD: Cystic Fibrosis Foundation, pp.

Dainese, E., Rodrigue, S., Delogu, G., Provvedi, R., Laflamme, L., Brzezinski, R., Fadda, G.,

Smith, I., Gaudreau, L., Palu, G., and Manganelli, R. (2006) Posttranslational regulation

of Mycobacterium tuberculosis extracytoplasmic-function sigma factor sigma L and roles

in virulence and in global regulation of gene expression. *Infect Immun* **74**: 2457-2461.

Damron, F.H., and Goldberg, J.B. (2012) Proteolytic regulation of alginate overproduction in

Pseudomonas aeruginosa. *Mol Microbiol* **84**: 595-607.

Damron, F.H., Qiu, D., and Yu, H.D. (2009) The Pseudomonas aeruginosa sensor kinase KinB

negatively controls alginate production through AlgW-dependent MucA proteolysis. *J*

Bacteriol **191**: 2285-2295.

De Las Penas, A., Connolly, L., and Gross, C.A. (1997) The sigmaE-mediated response to

extracytoplasmic stress in Escherichia coli is transduced by RseA and RseB, two

negative regulators of sigmaE. *Mol Microbiol* **24**: 373-385.

Deretic, V., Gill, J.F., and Chakrabarty, A.M. (1987) Gene algD coding for GDPmannose

dehydrogenase is transcriptionally activated in mucoid Pseudomonas aeruginosa. *J*

Bacteriol **169**: 351-358.

DeVries, C.A., and Ohman, D.E. (1994) Mucoid-to-nonmucoid conversion in alginate-producing

Pseudomonas aeruginosa often results from spontaneous mutations in algT, encoding a

putative alternate sigma factor, and shows evidence for autoregulation. *J Bacteriol* **176**:

6677-6687.

Emerson, J., Rosenfeld, M., McNamara, S., Ramsey, B., and Gibson, R.L. (2002)

Pseudomonas aeruginosa and other predictors of mortality and morbidity in young

children with cystic fibrosis. *Pediatr Pulmonol* **34**: 91-100.

Firoved, A.M., and Deretic, V. (2003) Microarray analysis of global gene expression in mucoid

Pseudomonas aeruginosa. *J Bacteriol* **185**: 1071-1081.

Gallagher, L.A., Shendure, J., and Manoil, C. (2011) Genome-scale identification of resistance

functions in Pseudomonas aeruginosa using Tn-seq. *MBio* **2**: e00315-00310.

Govan, J.R., and Deretic, V. (1996) Microbial pathogenesis in cystic fibrosis: mucoid

Pseudomonas aeruginosa and Burkholderia cepacia. *Microbiol Rev* **60**: 539-574.

Griffin, J.E., Gawronski, J.D., Dejesus, M.A., Ioerger, T.R., Akerley, B.J., and Sassetti, C.M.

(2011) High-resolution phenotypic profiling defines genes essential for mycobacterial

growth and cholesterol catabolism. *PLoS Pathog* **7**: e1002251.

Helmann, T.C., Deutschbauer, A.M., and Lindow, S.E. (2019) Genome-wide identification of

Pseudomonas syringae genes required for fitness during colonization of the leaf surface

and apoplast. *Proc Natl Acad Sci U S A* **116**: 18900-18910.

Henry, R.L., Mellis, C.M., and Petrovic, L. (1992) Mucoid Pseudomonas aeruginosa is a marker

of poor survival in cystic fibrosis. *Pediatr Pulmonol* **12**: 158-161.

Hershberger, C.D., Ye, R.W., Parsek, M.R., Xie, Z.D., and Chakrabarty, A.M. (1995) The algT

(algU) gene of Pseudomonas aeruginosa, a key regulator involved in alginate

biosynthesis, encodes an alternative sigma factor (sigma E). *Proc Natl Acad Sci U S A*

92: 7941-7945.

Hmelo, L.R., Borlee, B.R., Almblad, H., Love, M.E., Randall, T.E., Tseng, B.S., Lin, C., Irie, Y.,

Storek, K.M., Yang, J.J., Siehnel, R.J., Howell, P.L., Singh, P.K., Tolker-Nielsen, T.,

Parsek, M.R., Schweizer, H.P., and Harrison, J.J. (2015) Precision-engineering the

Pseudomonas aeruginosa genome with two-step allelic exchange. *Nat Protoc* **10**: 1820-

1841.

Horsburgh, M.J., and Moir, A. (1999) sigmaM, an ECF RNA polymerase sigma factor of Bacillus

subtilis 168, is essential for growth and survival in high concentrations of salt. *Molecular*

Microbiology **32**: 41-50.

Huang, H., Shao, X., Xie, Y., Wang, T., Zhang, Y., Wang, X., and Deng, X. (2019) An integrated

genomic regulatory network of virulence-related transcriptional factors in Pseudomonas

aeruginosa. *Nature Communications* **10**.

Intile, P.J., Diaz, M.R., Urbanowski, M.L., Wolfgang, M.C., and Yahr, T.L. (2014) The AlgZR

two-component system recalibrates the RsmAYZ posttranscriptional regulatory system

to inhibit expression of the Pseudomonas aeruginosa type III secretion system. *J*

Bacteriol **196**: 357-366.

Jones, A.K., Fulcher, N.B., Balzer, G.J., Urbanowski, M.L., Pritchett, C.L., Schurr, M.J., Yahr,

T.L., and Wolfgang, M.C. (2010) Activation of the Pseudomonas aeruginosa AlgU

regulon through mucA mutation inhibits cyclic AMP/Vfr signaling. *J Bacteriol* **192**: 5709-

5717.

Jones, C.J., Newsom, D., Kelly, B., Irie, Y., Jennings, L.K., Xu, B., Limoli, D.H., Harrison, J.J.,

Parsek, M.R., White, P., and Wozniak, D.J. (2014) ChIP-Seq and RNA-Seq reveal an

AmrZ-mediated mechanism for cyclic di-GMP synthesis and biofilm development by

Pseudomonas aeruginosa. *PLoS Pathog* **10**: e1003984.

Kong, W., Zhao, J., Kang, H., Zhu, M., Zhou, T., Deng, X., and Liang, H. (2015) ChIP-seq

reveals the global regulator AlgR mediating cyclic di-GMP synthesis in Pseudomonas

aeruginosa. *Nucleic Acids Res* **43**: 8268-8282.

Koo, B.M., Kritikos, G., Farelli, J.D., Todor, H., Tong, K., Kimsey, H., Wapinski, I., Galardini, M.,

Cabal, A., Peters, J.M., Hachmann, A.B., Rudner, D.Z., Allen, K.N., Typas, A., and

Gross, C.A. (2017) Construction and Analysis of Two Genome-Scale Deletion Libraries

for Bacillus subtilis. *Cell Syst* **4**: 291-305 e297.

Lee, S.A., Gallagher, L.A., Thongdee, M., Staudinger, B.J., Lippman, S., Singh, P.K., and

Manoil, C. (2015) General and condition-specific essential functions of Pseudomonas

aeruginosa. *Proc Natl Acad Sci U S A* **112**: 5189-5194.

Leech, A.J., Sprinkle, A., Wood, L., Wozniak, D.J., and Ohman, D.E. (2008) The NtrC Family

Regulator AlgB, Which Controls Alginate Biosynthesis in Mucoid Pseudomonas

aeruginosa, Binds Directly to the algD Promoter. *Journal of Bacteriology* **190**: 581-589.

Li, S., Lou, X., Xu, Y., Teng, X., Liu, R., Zhang, Q., Wu, W., Wang, Y., and Bartlam, M. (2019)

Structural basis for the recognition of MucA by MucB and AlgU in Pseudomonas

aeruginosa. *The FEBS Journal* **286**: 4982-4994.

Li, Z., Kosorok, M.R., Farrell, P.M., Laxova, A., West, S.E., Green, C.G., Collins, J., Rock, M.J.,

and Splaingard, M.L. (2005) Longitudinal development of mucoid Pseudomonas

aeruginosa infection and lung disease progression in children with cystic fibrosis. *JAMA*

293: 581-588.

Liberati, N.T., Urbach, J.M., Miyata, S., Lee, D.G., Drenkard, E., Wu, G., Villanueva, J., Wei, T.,

and Ausubel, F.M. (2006) An ordered, nonredundant library of Pseudomonas aeruginosa

strain PA14 transposon insertion mutants. *Proc Natl Acad Sci U S A* **103**: 2833-2838.

Martin, D.W., Schurr, M.J., Mudd, M.H., Govan, J.R., Holloway, B.W., and Deretic, V. (1993)

Mechanism of conversion to mucoidy in Pseudomonas aeruginosa infecting cystic

fibrosis patients. *Proc Natl Acad Sci U S A* **90**: 8377-8381.

Martin, D.W., Schurr, M.J., Yu, H., and Deretic, V. (1994) Analysis of promoters controlled by

the putative sigma factor AlgU regulating conversion to mucoidy in Pseudomonas

aeruginosa: relationship to sigma E and stress response. *J Bacteriol* **176**: 6688-6696.

Mathee, K., McPherson, C.J., and Ohman, D.E. (1997) Posttranslational control of the algT

(algU)-encoded sigma22 for expression of the alginate regulon in Pseudomonas

aeruginosa and localization of its antagonist proteins MucA and MucB (AlgN). *J Bacteriol*

179: 3711-3720.

Mendez, R., Gutierrez, A., Reyes, J., and Marquez-Magana, L. (2012) The extracytoplasmic

function sigma factor SigY is important for efficient maintenance of the Spbeta prophage

that encodes sublancin in Bacillus subtilis. *DNA Cell Biol* **31**: 946-955.

Missiakas, D., Mayer, M.P., Lemaire, M., Georgopoulos, C., and Raina, S. (1997) Modulation of

the Escherichia coli sigmaE (RpoE) heat-shock transcription-factor activity by the RseA,

RseB and RseC proteins. *Mol Microbiol* **24**: 355-371.

Parad, R.B., Gerard, C.J., Zurakowski, D., Nichols, D.P., and Pier, G.B. (1999) Pulmonary

outcome in cystic fibrosis is influenced primarily by mucoid Pseudomonas aeruginosa

infection and immune status and only modestly by genotype. *Infect Immun* **67**: 4744-

4750.

Pedersen, S.S., Hoiby, N., Espersen, F., and Koch, C. (1992) Role of alginate in infection with

mucoid Pseudomonas aeruginosa in cystic fibrosis. *Thorax* **47**: 6-13.

Pritchett, C.L., Little, A.S., Okkotsu, Y., Frisk, A., Cody, W.L., Covey, C.R., and Schurr, M.J.
(2015) Expression analysis of the Pseudomonas aeruginosa AlgZR two-component
regulatory system. *J Bacteriol* **197**: 736-748.

Pulcrano, G., Iula, D.V., Raia, V., Rossano, F., and Catania, M.R. (2012) Different mutations in
mucA gene of Pseudomonas aeruginosa mucoid strains in cystic fibrosis patients and
their effect on algU gene expression. *New Microbiol* **35**: 295-305.

Qiu, D., Eisinger, V.M., Head, N.E., Pier, G.B., and Yu, H.D. (2008) ClpXP proteases positively
regulate alginate overexpression and mucoid conversion in Pseudomonas aeruginosa.
Microbiology **154**: 2119-2130.

Qiu, D., Eisinger, V.M., Rowen, D.W., and Yu, H.D. (2007) Regulated proteolysis controls
mucoid conversion in Pseudomonas aeruginosa. *Proc Natl Acad Sci U S A* **104**: 8107-
8112.

Sautter, R., Ramos, D., Schneper, L., Ciofu, O., Wassermann, T., Koh, C.L., Heydorn, A.,
Hentzer, M., Hoiby, N., Kharazmi, A., Molin, S., Devries, C.A., Ohman, D.E., and
Mathee, K. (2012) A complex multilevel attack on Pseudomonas aeruginosa algT/U
expression and algT/U activity results in the loss of alginate production. *Gene* **498**: 242-
253.

Schulz, S., Eckweiler, D., Bielecka, A., Nicolai, T., Franke, R., Dotsch, A., Hornischer, K.,
Bruchmann, S., Duvel, J., and Haussler, S. (2015) Elucidation of sigma factor-
associated networks in Pseudomonas aeruginosa reveals a modular architecture with
limited and function-specific crosstalk. *PLoS Pathog* **11**: e1004744.

Schurr, M.J., Martin, D.W., Mudd, M.H., and Deretic, V. (1994) Gene cluster controlling
conversion to alginate-overproducing phenotype in Pseudomonas aeruginosa: functional
analysis in a heterologous host and role in the instability of mucoidy. *J Bacteriol* **176**:
3375-3382.

Schurr, M.J., Yu, H., Martinez-Salazar, J.M., Boucher, J.C., and Deretic, V. (1996) Control of

AlgU, a member of the sigma E-like family of stress sigma factors, by the negative

regulators MucA and MucB and Pseudomonas aeruginosa conversion to mucoidy in

cystic fibrosis. *J Bacteriol* **178**: 4997-5004.

Skurnik, D., Roux, D., Aschard, H., Cattoir, V., Yoder-Himes, D., Lory, S., and Pier, G.B. (2013)

A comprehensive analysis of in vitro and in vivo genetic fitness of Pseudomonas

aeruginosa using high-throughput sequencing of transposon libraries. *PLoS Pathog* **9**:

e1003582.

Turner, K.H., Wessel, A.K., Palmer, G.C., Murray, J.L., and Whiteley, M. (2015) Essential

genome of Pseudomonas aeruginosa in cystic fibrosis sputum. *Proc Natl Acad Sci U S A*

112: 4110-4115.

Vogel, H.J., and Bonner, D.M. (1956) Acetylornithinase of Escherichia coli: partial purification

and some properties. *J Biol Chem* **218**: 97-106.

Wood, L.F., Leech, A.J., and Ohman, D.E. (2006) Cell wall-inhibitory antibiotics activate the

alginate biosynthesis operon in Pseudomonas aeruginosa: Roles of sigma (AlgT) and

the AlgW and Prc proteases. *Mol Microbiol* **62**: 412-426.

Wood, L.F., and Ohman, D.E. (2009) Use of cell wall stress to characterize sigma 22 (AlgT/U)

activation by regulated proteolysis and its regulon in Pseudomonas aeruginosa. *Mol*

Microbiol **72**: 183-201.

Wozniak, D.J., and Ohman, D.E. (1994) Transcriptional analysis of the Pseudomonas

aeruginosa genes algR, algB, and algD reveals a hierarchy of alginate gene expression

which is modulated by algT. *J Bacteriol* **176**: 6007-6014.

Wozniak, D.J., Sprinkle, A.B., and Baynham, P.J. (2003) Control of Pseudomonas aeruginosa

algZ expression by the alternative sigma factor AlgT. *J Bacteriol* **185**: 7297-7300.

Xie, Z.D., Hershberger, C.D., Shankar, S., Ye, R.W., and Chakrabarty, A.M. (1996) Sigma

673   factor-anti-sigma factor interaction in alginate synthesis: inhibition of AlgT by MucA. *J*

674   *Bacteriol* **178**: 4990-4996.

675   Yin, Y., Withers, T.R., Wang, X., and Yu, H.D. (2013) Evidence for sigma factor competition in

676   the regulation of alginate production by Pseudomonas aeruginosa. *PLoS One* **8**:

677   e72329.

678   Yu, H., Schurr, M.J., and Deretic, V. (1995) Functional equivalence of Escherichia coli sigma E

679   and Pseudomonas aeruginosa AlgU: E. coli rpoE restores mucoidy and reduces

680   sensitivity to reactive oxygen intermediates in algU mutants of P. aeruginosa. *J Bacteriol*

681   **177**: 3259-3268.

682   Zhao, H., Roistacher, D.M., and Helmann, J.D. (2019) Deciphering the essentiality and function

683   of the anti-sigma(M) factors in Bacillus subtilis. *Mol Microbiol* **112**: 482-497.

685 **TABLES**

686 **Table 1. *mucA* is essential in a variety of wild-type *P. aeruginosa* strains.**

Strain background	Description[†]	Number of isolates resolved to	
		WT	ΔmucA
PAO1	Laboratory strain, Class II	168	0*
PA14	Laboratory strain, Class I	147	0*
PA103	Laboratory strain, unclassified	361	0*
PAK	Laboratory strain, unclassified	191	0*
CF127	Mucoid CF isolate, unclassified	136	0*
CF18	Non-mucoid CF isolate, unclassified	129	0*
CF27	Rugose CF Isolate, Class IV	152	0*
X13273	Blood isolate, Class II	142	0*
X24509	UTI isolate, Class II	221	0*
MSH10	Water isolate, Class III	146	0*
E2	Tomato plant isolate, Class II	141	0*

687 [†] Class identification is based on the exopolysaccharide expression profile as described in

688 (Colvin *et al.*, 2012): Class I, Pel-dominant matrix; Class II, Psl-dominant matrix; Class III,

689 exopolysaccharide redundant matrix users; and Class IV, matrix overproducers. CF, cystic

690 fibrosis; UTI, urinary tract infection.

691 * $p < 0.0001$, Fischer's exact test

 FIGURE LEGENDS

Figure 1. The first 50 amino acids of MucA are necessary and sufficient for viability.

(A) Schematic of MucA encoded by the ectopic *mucA* alleles in the PAO1 Δ*algD* strains tested

for *mucA* essentiality in (B). MucA aa 1-143 lacks the MucB binding domain and is the product

of the common *mucA22* allele. MucA 1-110 lacks the entire periplasmic domain. MucA aa 1-75

contains most of the AlgU binding domain, while shorter truncations contain only parts of the

AlgU binding domain (MucA aa 1-62, 1-50, 1-40, and 1-24). Green, AlgU binding domain (AlgU

DB); blue, the transmembrane domain (TM); and purple, the MucB binding domain (MucB BD).

(B) Frequency of observed isolates resolving to the endogenous wild-type *mucA* allele (WT,

black) or the deletion allele (Δ*mucA*, white) in the allelic exchange assay. Super-imposed on the

bars are the number of isolates that were tested in each category. Asterisk, $p < 0.0001$; Fisher's

exact test. Caret, slower growing isolates.

Figure 2. Depletion of MucA results in cell death.

Viable colony counts of PAO1 Δ*mucA* attTn7:: *rhaSR*-P*rhaBAD*-*mucA*-10His over time in **(A)** LB,

(B) PIB, **(C)** SCFM and **(D)** VBMM with (+ rha; red circles) or without (– rha; gray squares)

0.05% rhamnose. Viable colony counts after incubation in the indicated condition and time was

determined by plating the cells on LB with 0.05% rhamnose to allow cells to recover and grow.

Hash, broken y-axis; error bars, SEM (N=3). Asterisk, statistically different from that at the same

time point grown in the presence of rhamnose ($p < 0.01$, N=3, mixed model ANOVA with post-

hoc Bonferroni test).

Figure 3. The physical interaction of MucA and AlgU is required for survival.

(A) Four residues of MucA make greater than one predicted hydrogen bond with AlgU. The

MucA-AlgU co-crystal structure (PDB 6IN7; Li et al., 2019) with the cytosolic domain of MucA

(aa 1-78; red) and Regions 2 and 4 of AlgU (green, AlgU$_2$; yellow, AlgU$_4$) is shown. The

residues of MucA (grey) that are predicted to make more than one hydrogen bond with AlgU are

labeled in the inset. Black dotted lines, predicted hydrogen bonds; red atoms, oxygen; blue

atoms, nitrogen. **(B)** Substitution of MucA residues at its interface with AlgU abolish their binding

via yeast two-hybrid. The first 75 residues of MucA were fused to the Gal4 DNA binding domain

(DBD-MucA$^{1-75}$) and AlgU was fused to the Gal4 activation domain (AD-AlgU). Interaction of

MucA and AlgU led to *lacZ* expression. Beta-galactosidase activity (in Miller units) was used as

a proxy for the protein interaction strength. WT, wild-type protein sequence; –, no fusion protein

included; hash, broken y-axis; error bars, SEM (N=3); letters, statistical groups with the different

letters representing statistically different groups ($p < 0.01$; biological triplicate with technical

quadruplicates; ANOVA with post-hoc Tukey HSD). **(C)** Frequency of observed isolates

resolving to the endogenous wild-type *mucA* allele (WT, black) or the deletion allele (Δ*mucA*,

white) in the allelic exchange assay, using PAO1 Δ*algD att*Tn7::P$_{algU}$-*mucA*, where *mucA*

encodes for the indicated substitution. Super-imposed on the bars are the number of isolates

that were tested in each category. Asterisk, $p < 0.0001$; Fisher's exact test. Caret, slower

growing isolates.

Figure 4. AlgU mutants with reduced sigma factor activity suppress *mucA* essentiality.

(A) Frequency of observed isolates resolving to the endogenous wild-type *mucA* allele (WT,

black) or the deletion allele (Δ*mucA*, white) in the allelic exchange assay, using PAO1 Δ*algU*

*att*Tn7::P$_{algU}$-*algU*, where *algU* encodes for the indicated substitution. Super-imposed on the

bars are the number of isolates that were tested in each category. Asterisk, $p < 0.0001$; Fisher's

exact test. **(B)** Schematic of mutations seen in revertants that could grow in the absence of

mucA. Revertants were selected by growing PAO1 Δ*mucA att*Tn7::P$_{rhaBAD}$-*mucA* on media

lacking rhamnose. Blue rectangles, multi-base pair deletions; left arrow, deletion extends into

the promoter; blue triangles, single base pair deletions; black triangles, nonsense mutations;

white triangles, duplications resulting in 3 or 4 amino acid insertions; red triangles, missense

mutations. See Table S2 for a full description of the *algU* mutations.

Figure 5. Overexpression of *algU* is lethal in the absence of *mucA*.

(A) Growth curves of PAO1 strains containing an arabinose-inducible copy of *algU* in wild-type

(purple squares), Δ*algU* (blue diamonds), and Δ*algU* Δ*mucA* (red circles) backgrounds in LB.

Readings are normalized to wild-type PAO1 (lacking an inducible *algU* gene). Open symbols

represent conditions in the absence of arabinose; closed symbols, with 1% arabinose. Error

bars, SD (n=18). Asterisk, statistically different from the same strain grown in the absence of

arabinose (p < 0.01, N = 3, ANOVA with post-hoc Tukey HSD). **(B)** Growth rate of strains as

described in (A), with increasing induction of *algU* (relative to PAO1 *att*Tn7::P$_{araBAD}$-*algU* grown

in the absence of arabinose). Strains were grown in LB with 0%, 0.1%, 0.25%, and 1%

arabinose ([Ara]). Error bars, SEM (N=3). Asterisk, statistically different from the same strain

grown in the absence of arabinose (p < 0.05, N = 3, ANOVA with post-hoc Tukey HSD). See

Table S3 for full statistical comparisons.

Figure 6. Lethality caused by *algU* overexpression can be rescued via *rpoD* expression.

Growth rate of indicated strains grown in LB with (+) or without (−) 2% arabinose (relative to

PAO1 Δ*algU* Δ*mucA* grown in the absence of arabinose). Error bars, SEM (N=3). Asterisk,

statistically different from the same strain grown in the absence of arabinose (p < 0.05, N = 3,

ANOVA with post-hoc Tukey HSD). See Table S4 for full statistical comparisons.

Figure 7. Unchecked AlgU activity in the absence of MucA leads to bacterial cell death.

Under conditions of no envelope stress (left), the full-length MucA (red) binds strongly to AlgU

(green star), leaving very little free AlgU to interact with the RNAP (grey oval). Therefore, the

AlgU regulon (green arrow) is in the "off" state (red X). Under conditions of envelope stress or in

strains containing *mucA* mutations that lead to a truncated product (middle), a cleaved cytosolic

form of MucA is produced. This form can still interact with and inhibit AlgU. However, the

strength of the interaction is weaker than with the full-length MucA, allowing for a pool of free

AlgU that can then bind to and recruit RNAP to the promoters of its regulon. Under such

conditions, the AlgU regulon, which includes *algU* and *mucA*, is activated and in the "on" state

(green up arrows). However, because of the feedback on MucA, negative regulation of AlgU

activity is still present in the cell. In the absence of *mucA* (right), there is no MucA to bind to and

regulate AlgU. All of the AlgU is free to interact with the core RNAP. This leads to high

expression of the AlgU regulon and overproduction of AlgU itself. Under such strong positive

feedback and in the absence of the negative regulator, the unchecked activity of AlgU leads to

cell death.

Figure 1. The first 50 amino acids of MucA are necessary and sufficient for viability.
(A) Schematic of MucA encoded by the ectopic *mucA* alleles in the PAO1 Δ*algD* strains tested for *mucA* essentiality in (B). MucA aa 1-143 lacks the MucB binding domain and is the product of the common *mucA22* allele. MucA 1-110 lacks the entire periplasmic domain. MucA aa 1-75 contains most of the AlgU binding domain, while shorter truncations contain only parts of the AlgU binding domain (MucA aa 1-62, 1-50, 1-40, and 1-24). Green, AlgU binding domain (AlgU DB); blue, the transmembrane domain (TM); and purple, the MucB binding domain (MucB BD). **(B)** Frequency of observed isolates resolving to the endogenous wild-type *mucA* allele (WT, black) or the deletion allele (Δ*mucA*, white) in the allelic exchange assay. Super-imposed on the bars are the number of isolates that were tested in each category. Asterisk, p < 0.0001; Fisher's exact test. Caret, slower growing isolates.

Figure 2. Depletion of MucA results in cell death.
Viable colony counts of PAO1 Δ*mucA* attTn7:: *rhaSR*-P*rhaBAD*-*mucA*-10His over time in **(A)** LB, **(B)** PIB, **(C)** SCFM and **(D)** VBMM with (+ rha; red circles) or without (– rha; gray squares) 0.05% rhamnose. Viable colony counts after incubation in the indicated condition and time was determined by plating the cells on LB with 0.05% rhamnose to allow cells to recover and grow. Hash, broken y-axis; error bars, SEM (N=3). Asterisk, statistically different from that at the same time point grown in the presence of rhamnose ($p < 0.01$, N=3, mixed model ANOVA with post-hoc Bonferroni test).

Figure 3. The physical interaction of MucA and AlgU is required for survival.
(A) Four residues of MucA make greater than one predicted hydrogen bond with AlgU. The MucA-AlgU co-crystal structure (PDB 6IN7; Li et al., 2019) with the cytosolic domain of MucA (aa 1-78; red) and Regions 2 and 4 of AlgU (green, $AlgU_2$; yellow, $AlgU_4$) is shown. The residues of MucA (grey) that are predicted to make more than one hydrogen bond with AlgU are labeled in the inset. Black dotted lines, predicted hydrogen bonds; red atoms, oxygen; blue atoms, nitrogen. **(B)** Substitution of MucA residues at its interface with AlgU abolish their binding via yeast two-hybrid. The first 75 residues of MucA were fused to the Gal4 DNA binding domain (DBD-MucA$^{1-75}$) and AlgU was fused to the Gal4 activation domain (AD-AlgU). Interaction of MucA and AlgU led to *lacZ* expression. Beta-galactosidase activity (in Miller units) was used as a proxy for the protein interaction strength. WT, wild-type protein sequence; –, no fusion protein included; hash, broken y-axis; error bars, SEM (N=3); letters, statistical groups with the different letters representing statistically different groups ($p < 0.01$; biological triplicate with technical quadruplicates; ANOVA with post-hoc Tukey HSD). **(C)** Frequency of observed isolates resolving to the endogenous wild-type *mucA* allele (WT, black) or the deletion allele ($\Delta mucA$, white) in the allelic exchange assay, using PAO1 $\Delta algD$ *att*Tn7::P$_{algU}$-*mucA*, where *mucA* encodes for the indicated substitution. Super-imposed on the bars are the number of isolates that were tested in each category. Asterisk, $p < 0.0001$; Fisher's exact test. Caret, slower growing isolates.

Figure 4. AlgU mutants with reduced sigma factor activity suppress *mucA* essentiality.
(A) Frequency of observed isolates resolving to the endogenous wild-type *mucA* allele (WT, black) or the deletion allele (Δ*mucA*, white) in the allelic exchange assay, using PAO1 Δ*algU* *att*Tn7::P*algU*-*algU*, where *algU* encodes for the indicated substitution. Super-imposed on the bars are the number of isolates that were tested in each category. Asterisk, p < 0.0001; Fisher's exact test. **(B)** Schematic of mutations seen in revertants that could grow in the absence of *mucA*. Revertants were selected by growing PAO1 Δ*mucA* *att*Tn7::P*rhaBAD*-*mucA* on media lacking rhamnose. Blue rectangles, multi-base pair deletions; left arrow, deletion extends into the promoter; blue triangles, single base pair deletions; black triangles, nonsense mutations; white triangles, duplications resulting in 3 or 4 amino acid insertions; red triangles, missense mutations. See Table S2 for a full description of the *algU* mutations.

Figure 5. Overexpression of *algU* is lethal in the absence of *mucA*.
(A) Growth curves of PAO1 strains containing an arabinose-inducible copy of *algU* in wild-type (purple squares), Δ*algU* (blue diamonds), and Δ*algU* Δ*mucA* (red circles) backgrounds in LB. Readings are normalized to wild-type PAO1 (lacking an inducible *algU* gene). Open symbols represent conditions in the absence of arabinose; closed symbols, with 1% arabinose. Error bars, SD (n=18). Asterisk, statistically different from the same strain grown in the absence of arabinose ($p < 0.01$, N = 3, ANOVA with post-hoc Tukey HSD). **(B)** Growth rate of strains as described in (A), with increasing induction of *algU* (relative to PAO1 *att*Tn7::P$_{araBAD}$-*algU* grown in the absence of arabinose). Strains were grown in LB with 0%, 0.1%, 0.25%, and 1% arabinose ([Ara]). Error bars, SEM (N=3). Asterisk, statistically different from the same strain grown in the absence of arabinose ($p < 0.05$, N = 3, ANOVA with post-hoc Tukey HSD). See Table S3 for full statistical comparisons.

Figure 6. Lethality caused by *algU* overexpression can be rescued via *rpoD* expression. Growth rate of indicated strains grown in LB with (+) or without (−) 2% arabinose (relative to PAO1 Δ*algU* Δ*mucA* grown in the absence of arabinose). Error bars, SEM (N=3). Asterisk, statistically different from the same strain grown in the absence of arabinose ($p < 0.05$, N = 3, ANOVA with post-hoc Tukey HSD). See Table S4 for full statistical comparisons.

Figure 7. Unchecked AlgU activity in the absence of MucA leads to bacterial cell death. Under conditions of no envelope stress (left), the full-length MucA (red) binds strongly to AlgU (green star), leaving very little free AlgU to interact with the RNAP (grey oval). Therefore, the AlgU regulon (green arrow) is in the "off" state (red X). Under conditions of envelope stress or in strains containing *mucA* mutations that lead to a truncated product (middle), a cleaved cytosolic form of MucA is produced. This form can still interact with and inhibit AlgU. However, the strength of the interaction is weaker than with the full-length MucA, allowing for a pool of free AlgU that can then bind to and recruit RNAP to the promoters of its regulon. Under such conditions, the AlgU regulon, which includes *algU* and *mucA*, is activated and in the "on" state (green up arrows). However, because of the feedback on MucA, negative regulation of AlgU activity is still present in the cell. In the absence of *mucA* (right), there is no MucA to bind to and regulate AlgU. All of the AlgU is free to interact with the core RNAP. This leads to high expression of the AlgU regulon and overproduction of AlgU itself. Under such strong positive feedback and in the absence of the negative regulator, the unchecked activity of AlgU leads to cell death.