The Role of Colony Temperature in the Entrainment of Circadian Rhythms of Honey Bee Foragers

Authors:

Manuel A. Giannoni-Guzmán[1], Emmanuel Rivera[2], Janpierre Aleman-Rios[3], Alexander M. Melendez Moreno[3], Melina Perez Ramos[2], Darimar Loubriel[1], Darell Moore[4], Tugrul Giray[3] and Jose L. Agosto-Rivera[3]

Affiliations:

1. Department of Biological Sciences, Vanderbilt University, Nashville, TN, USA

2. Department of Biology, Brandeis University, Waltham, MA, USA

3. Department of Biology, University of Puerto Rico Rio Piedras Campus, San Juan, PR

4. Department of Biological Sciences, East Tennessee State University, Johnson City, TN, USA

Corresponding author: Manuel A. Giannoni-Guzmán

e-mail: manuel.giannoni10@gmail.com

Abstract

Honey bees utilize their circadian rhythms to accurately predict the time of day. This ability allows foragers to remember the specific timing of food availability and its location for several days. Previous studies have provided strong evidence toward light/dark cycles being the primary Zeitgeber for honey bees. Recent work in our laboratory described large individual variation in the endogenous period length of honey bee foragers from the same colony and differences in the endogenous rhythms under different constant temperatures. In this study, we further this work by examining temperature inside the honey bee colony. By placing temperature and light data loggers at different locations inside the colony we uncovered that temperature oscillates with a 24-hour period at the periphery of the colony. We then simulated this temperature oscillation in the laboratory and found that using the temperature cycle as a Zeitgeber, foragers present large individual differences in the phase of locomotor rhythms with respect to temperature. Moreover, foragers successfully entrain to these simulated temperature cycles and advancing the cycle by six hours, resulted in changes in the phase of locomotor activity for the most foragers in the assay. The results shown in this study highlight the importance of temperature as a potential Zeitgeber in the field. Future studies will examine the possible functional and evolutionary role of the observed phase differences of circadian rhythms.

Introduction:

Circadian rhythms are biological clocks that synchronize endogenous processes to daily

changes in environmental cues, such as light and temperature (Panda, Hogenesch, & Kay, 2002).

While circadian rhythms were originally measured as daily behavioral patterns, such as locomotor

activity, at the molecular level, circadian rhythms are driven by a set of proteins that generate

negative feedback loops that regulate the transcription, translation and post translational

modifications of a large number of genes (Blau, 2001; Cyran et al., 2003; Dunlap, 1999; Gardner,

Hubbard, Hotta, Dodd, & Webb, 2006). These feedback loops can be measured by oscillations in

the expression levels of the genes that make up the clock (Ko & Takahashi, 2006; Takahashi,

1999).Furthermore, changes in light/dark cycles, temperature, food and other factors that serve

as Zeitgebers can alter the behavioral and molecular cycles (Hut & Beersma, 2011; Moore &

Rankin, 1993; von Frisch, Aschoff, Frisch, & Aschoff, 1987).

In honey bees the circadian clock is associated with various complex behavioral

processes that drive the survival and fitness of the colony. Drones and queens (the reproductive

castes in the colony) mate at a specific time during the day (Galindo-Cardona et al., 2012).

Foragers, who go out in search of different resources during daylight, learn the time and location

of a food source and anticipate its availability on following days (Moore, 2001; Moore & Doherty,

2009; von Frisch, 1967). Also, foragers remember the sun's location throughout the day and are

able to navigate on cloudy days in the absence of direct sunlight (Goodwin & Lewis, 1987; von

Frisch, 1967). Although these and other processes rely on precise timing of the circadian clock,

few studies have examined the potential environmental cues that entrain circadian rhythms in

nature.

The first studies of honey bee circadian rhythms looked at foraging rhythms at the colony

level (H. Buttel-Reepen, 1900; Forel, 1910; Wahl, 1932, 1933). Studies examining the potential

Zeitgebers that entrain circadian rhythms in honey bee colonies and individuals came to the

conclusion that light/dark cycles are the primary time giver (W Beier, 1968; Wilhelm Beier &

Lindauer, 1970; Detrain, Deneubourg, & Pasteels, 1999; Renner, 1960). At the group level,

studies where groups of individuals or whole colonies trained to visit a specific food source were

translocated to a different time zone (ex. New York to California) presented foraging activity at

the time relative to their original light/dark cycles and would entrain over time to the new light/dark

regiment (Renner, 1959). At the individual level studies examining the entrainment to light/dark

cycles of foragers show that light successfully entrains the locomotor rhythms of foragers and that

the lights-off transition determines the phase of activity in foragers (Moore & Rankin, 1985, 1993;

Spangler, 1972). However, the fact that honey bee colonies and individuals entrain to light/dark

cycles does not exclude the possibility to entrainment by other environmental cues.

An important aspect of hive maintenance is that honey bees socially regulate colony

temperature keeping it optimally at 35°C (Heinrich, 1980). At the behavioral level this is achieved

by fanning (ventilation) and water evaporation inside the colony (Kronenberg & Heller, 1982;

Seeley, 1974). Researchers examining the mechanisms underlying this thermoregulation found

that individual variation at the genetic level is associated with differences in worker's fanning

response threshold to temperature (Jones, Myerscough, Graham, & Oldroyd, 2004). This

variation results in different patrilines fanning at different temperatures and, therefore, promoting

the stability of brood nest temperature (Jones et al., 2004). Although one of the main challenges

to colony thermoregulation is heat associated with the sun, which changes throughout the day,

whether individual variation in circadian rhythms play a role in thermoregulation remains unknown.

Thermoregulation of honey bee colonies is essential for colony performance and survival

(Heinrich , 1981; Heinrich, 1993). Experiments examining the effects of low temperatures (28-

30°C) on brood development revealed that at these temperatures can result in shriveled wings or

other malformations, while brood kept at high temperatures (38°C-40°C) present high mortality

rates ((Heinrich, 1993; Himmer, 1927). Experiments following up on this work, show that pupal

development at 32°C, which is only 3°C less than the optimal core temperature, results in

significant deficits on waggle dance behavior and learning and memory assays, compared to bees

raised at 36°C (Tautz, Maier, Groh, Rossler, & Brockmann, 2003). Nonetheless, little is known

about the effects of colony temperature on the physiological regulation of the circadian rhythms

of honey bees.

Researchers explored the entrainment of circadian rhythms by square-wave temperature

cycles and found that cycles with amplitudes greater than 9°C could successfully entrain circadian

rhythms of foragers in the laboratory (Moore & Rankin, 1993). However, whether temperature

cycles strong enough to entrain circadian rhythms exist inside the colony has not been thoroughly

studied. Changes in environmental temperature in the laboratory have been shown to have strong

effects on the endogenous period length of foragers (Fuchikawa & Shimizu, 2007; Giannoni-

Guzmán et al., 2014). Furthermore, recent work has revealed that there is a large degree of

individual differences in the endogenous period length of the circadian clock of foragers

(Giannoni-Guzmán et al., 2014). This large variation in the endogenous period length could be

adaptive at the colony level and could result in individual differences in the phase of the circadian

clock to time givers.

The main goal of this study was to ascertain if temperature changed on a daily basis inside

honey bee colonies and if these changes (temperature cycles) were capable of entraining the

circadian rhythms of honey bee foragers. In order to do this, we measured light and temperature

at various locations inside of the colony to examine whether temperature and light present daily

oscillations inside honey bee colonies. We then explored the effects of temperature cycles

observed in the colony on the circadian rhythms of honey bees in the laboratory. To determine if

these cycles were capable of entraining the circadian rhythms we advanced the temperature cycle

after 6 days of locomotor activity measurements and observed its effects on the phase of the

locomotor rhythms.

Material and Methods

Colony Light and Temperature Measurements:

Light and temperature measurements were carried out using HOBO® pendant data

loggers (UA-002-64). Four loggers were placed inside the colony at the center, entrance,

periphery and top, while one was placed outside the hive as shown (Figure 1A). The colony in

which we collected these measurements was a two story colony in good health with a naturally

mated queen laying eggs. With the exception of 30 minutes during July 11, 2013, when data was

uploaded from the pendants and the batteries were replaced, temperature and light

measurements were continuously collected in 30-minute intervals from June 13, 2012 until

September 2, 2014. Temperature and light data presented was averaged as 48 hours for

comparison with double plotted actograms (Figure 1).

Forager Collection:

All of the bees utilized in our experiments came from colonies in good health that had naturally

mated queens and were laying eggs at the time of the experiments. For each experiment, foragers

were collected returning to the colony by blocking the entrance with 8-mesh wire screen and

capturing them in tubes as previously described (Giray, Galindo-Cardona, & Oskay, 2007).

Collected bees were provided with food and water during transportation to UPR Rio Piedras

campus (30-40min car ride). Once in the lab bees were anesthetized and placed in locomotor

activity monitors as previously described (Giannoni-Guzmán et al., 2014).

Experiment 1: Simulating temperature cycles of the colony in the laboratory

Locomotor activity recordings were performed inside an environmental chamber (Percival,

I-30BLL), where temperature was programed to oscillate with an amplitude of 9.2°C, as observed

in the periphery of the colony (Figure 1C). To insure that the temperature was oscillating in the

same manner (or as close as possible) as the observed oscillation in nature, the incubator was

set up and calibrated 2 weeks prior to the experiment. Age was controlled by paint marking

individuals 1-day old workers and returning them to the colony on October 7[th] 2014. Locomotor

activity recordings began at 21 days of age and took place from October 28[th] until November 7[th]

2014. Phase analysis of locomotor activity was performed on days 6-11 of locomotor recording,

to allow 5-6 days needed for for bees to acclimate to square wave temperature cycles (Moore &

Rankin, 1993).

Experiment 2: 6-hour phase advance of simulated temperature cycles

Locomotor activity assays were conducted using the the same environmental conditions

as experiment 1 for the first 6 days of the experiment. On the 7[th] day the environmental

temperature cycles were advanced by 6 hours and were kept this way until the end of the

experiment (Figure 3A).

Data analysis:

Locomotor activity data was processed using MatLab® toolboxes developed in Jeffrey

Hall's laboratory (Levine et al., 2002). Period length and rhythm strength were calculated using

autocorrelation analysis. The locomotor activity was then compared to the environmental cycles

to which bees were exposed using cross correlation analysis. Changes in the phase of locomotor

rhythms after temperature phase advance were analyzed via Repeated Measures MANOVA.

Results:

Previous work has shown that temperature is a successful Zeitgeber for the circadian

clock of honey bees in the laboratory (Moore & Rankin, 1993). However, whether temperature

cycles capable of entraining the circadian rhythm of honey bees are present inside the colony

was unknown. To examine if temperature cycles are present in honey bee colonies, we placed 5

data loggers at different locations of the colony (center, entrance, top, periphery and outside) and

recorded light and temperature measurements in 30-minute intervals (Figure 1A). Our results

revealed that temperature oscillates inside honey bee colonies in a daily manner, with amplitudes

greater than 7 degrees (Figure 1). Specifically, we observed that at the center of the colony

temperature was maintained within 35±0.7 ˚C, while daily temperature cycles at the entrance

(amplitude=7.3˚C), periphery (amplitude=9.2˚C) and top (amplitude= 16.3˚C) of the colony were

detected. The control data logger that was placed outside the colony detected strong temperature

and light cycles with greater amplitude than those detected in the colony (Figure 1E). In addition,

the presence of light inside the colony was only detected by the logger at the entrance and was

200 times less at its highest peak than that detected by the logger outside the hive (Figure 1E,

F).

156
Figure 1. Temperature oscillates with a 24h period in the periphery of the colony. A) Top and
side view of a two story honey bee colony presenting the positions of the 5 sensors used to measure
environmental temperature and light, inside and outside the hive. Average environmental data logged
at the **B)** center, **C)** periphery, **D)** top, **E)** entrance and **F)** outside the colony. Temperature was plotted
on the left y-axis (black line), while light is plotted on the right Y-axis (blue line). As sensors move
further away from the center of the colony the amplitude of temperature oscillations increases from
0.7˚C at the center of the colony up to 16.3˚C in the top of the colony.

Given the previously reported effects of environmental temperature on the endogenous

rhythm of honey bees (Fuchikawa & Shimizu, 2007; Giannoni-Guzmán et al., 2014), we examined

the effect of the observed temperature cycles on the locomotor rhythms of foragers. In order to

examine the possible effects of temperature cycles on the circadian rhythm of foragers, we

simulated the temperature cycle of the periphery of the colony in the laboratory and measured

the locomotor rhythms of foragers. This particular temperature oscillation was chosen due to its

amplitude (9.2°C), since square-wave cycles with amplitudes less than 9°C were unable to entrain

foragers in a previous study (Moore & Rankin, 1993). We hypothesized that exposing foragers to

simulated temperature cycles would result in their locomotor activity rhythms syncing with the

phase of the temperature cycle. We were surprised to uncover that honey bee foragers exposed

to the simulated cycles presented individual differences in the phase of their locomotor rhythms

with respect to the temperature cycles (ψ) (Figure 2). While there were differences in the phase

of the clock with respect to temperature, most individuals in our sample were phase advanced

(Figure 2A), while some presented a delayed phase (Figure 2B). Phase quantification for each

individual was performed via cross-correlation of the locomotor activity and the temperature cycle

(Figure 2, panel (iii)). Frequency distribution of the phase of individuals revealed that more than

60 percent of foragers are phase advanced to the temperature cycle (Figure 2C). In addition,

period length of foragers was on average 24 hours with little variation (0.06h SEM) (Figure 2D).

This small variation of period length is significantly smaller to that of foragers under constant

conditions (23.8±0.19 h), and suggests that these temperature cycles may be entraining the

circadian rhythm of foragers.

A) Advanced forager

B) Delayed forager

C) Phase Distribiution

D) Period length Distribiution

Figure 2. Simulating peripheral temperature oscillations is associated with strong rhythmicity and large individual differences in circadian phase. Representative examples of foragers that presented **A)** phase advance or **B)** phase delay with respect to the simulated temperature cycles in controlled laboratory conditions. **i)** Double plotted actogram of the simulated temperature cycle (red histograms) at the top and locomotor activity patterns below (blue histograms). **ii)** Autocorrelation analysis was utilized to determine period length and overall rhythm strength of locomotor rhythms (Levine et al., 2002). **iii)** Phase of the locomotor rhythms with respect to the temperature cycles (ψ) was quantified using cross-correlation analysis, which compares the distribution of locomotor activity with the temperature cycle and yields the phase as the lag (h). **C)** Frequency distribution of the phase of locomotor activity with respect to temperature cycle. **D)** Frequency distribution of period length of foragers exposed to simulated temperature cycles.

199   To test if the observed temperature cycles were capable of entraining the locomotor

200 rhythms of foragers, we performed a 6-hour phase advance to the temperature cycle on the 7[th]

201 day of locomotor activity measurements (Figure 3A). We hypothesized that if the temperature

202 cycles observed in the colony were capable of entraining circadian rhythms, then shifting the

203 temperature cycle would result in changes in the phase of locomotor activity of bees exposed to

204 this change. Consistent with our hypothesis we observed that some individuals shifted their

205 locomotor rhythms as temperature was shifted (Figure 3B), while some foragers seemed to be

206 unaffected by the temperature shift (Figure 3C). Repeated measures analysis of the phase before

207 and after the temperature advance resulted in average locomotor activity phase advance of 2

208 hours (Repeated Measures MANOVA, $F=1.62$, $df=1$, $p<<0.01$, $n=144$) (Figure 3D). Frequency

209 distribution of the differences in the phase before and after the 6hr advance, revealed differences

210 in the response of locomotor rhythms to temperature (Figure 3E). Taken together, these results

211 suggest that temperature cycles observed in the colony are capable of entraining locomotor

212 rhythms of honey bee foragers, but individual differences in the response to temperature cycle

213 changes suggest differences in the sensitivity of the clock to changes in temperature.

214
Figure 3. Simulated peripheral temperature serves as a Zeitgeber to the circadian rhythm of
foragers. A) Double plotted simulated temperature cycles. From day 7 onward temperature cycle
inside the incubator were advanced by 6 hours. This resulted in some individuals displaying changes
in their locomotor activity patterns **B)**, while some individuals presented no response to the change in
the environmental temperature cycle **C)**. **D)** Repeated measures MANOVA utilized to analyze the
effects of the 6hr temperature advance on the phase of locomotor rhythms, revealed significant
differences between the phase before the advance and the phase after said advance (F=1.62, df=1,
p<<0.01, n=144). **E)** Frequency distribution of the change in phase locomotor activity rhythms after 6-
hour temperature phase advance.

Discussion:

Here we show that at the periphery of the colony, where foragers spend time, temperature

significantly oscillates when a 24hrs period (Figure 1). Simulating this temperature signal in the

laboratory is sufficient to entrain the circadian clock of foragers, suggesting that temperature could

be an important Zeitgeber in the colony (Figure 2, 3). Interestingly, we found that there are large

individual differences in the phase of locomotor rhythms with respect to the temperature cycles.

These differences may provide the basis for a shift strategy suggested by a previous study (Kraus,

Gerecke, & Moritz, 2011). Taken together, temperature is an important environmental cue inside

the colony and could possibly entrain the circadian rhythms of foragers in nature.

Before this study, the circadian clock of honey bees was thought to be entrained mainly

by light/dark cycles (Moore & Rankin, 1985; Renner, 1959, 1960). This idea of light as the main

Zeitgeber of honey bee foragers results from the combination of field observations of foraging,

translocation experiments where a colony was moved from the east to the west coast of the United

States, and locomotor activity measurements performed in the laboratory (Renner, 1959).

However, these experiments do not take into account temperature changes that occur within the

colony. Our discovery that temperature cycles strong enough to entrain circadian rhythms of

foragers are present inside the colony (Figure 1) opens a novel avenue of research where

temperature could potentially influence the rhythms of bees in a natural setting.

Consistent with this theory Moore and Rankin proposed that light and temperature may

have a synergistic effect on the circadian clock system in honey bees (Moore and Rankin 1993).

At this point it is unknown which of the environmental Zeitgebers (temperature or light) plays a

more significant role in the entrainment of circadian rhythms in the hive. Experiments where both

light and temperature cycles show that some individuals respond strongly to light, while others

present locomotor activity during times when light is on and temperature high (Moore & Rankin,

1993). Since foragers have been shown to stay inside the colony after visiting their food source

(Körner, 1940; Moore, Siegfried, Wilson, & Rankin, 1989; Seeley, 1995; H. B. von Buttel-Reepen,

1903; von Frisch, 1940), it is possible that while inside the colony foragers rely on temperature

cycles to accurately tell time and use light input as a Zeitgeber foraging. This hypothesis stems

from the lack of light inside most of the colony. In order to test this hypothesis, further studies

exploring the functional role of temperature cycles in the colony as well as the mechanisms of

circadian entrainment in honey bees are needed.

Consistent with the results from the Moore and Rankin (1993) study, we observed that

honey bee foragers entrain to temperature cycles. To our knowledge, this is the first time that

gradual temperature cycles, simulating those observed in the field, have been utilized in the

laboratory. In addition, by simulating the peripheral temperature cycles we found that foragers

present a large degree of individual variation in the phase of locomotor rhythms with respect to

temperature (Figure 2). Although further studies are needed to understand the functional and

evolutionary role of this variation, we can speculate that phase differences of the individual honey

bee foragers may result in differences in the temporal allocation of tasks. This hypothesis would

be in line with the results of a recent study that suggests the existence of shift work in honey bees

and its genetic component (Kraus et al., 2011). Alternatively, the observed differences in the

phase of circadian rhythms could be the result of previous entrainment to an outside stimulus,

such as nectar and pollen. This can be evidenced by the ability of foragers to successfully be

trained for several days to a food source (von Frisch et al., 1987).

When performing a 6-hour advance on the temperature cycle we observed that some

individual's locomotor activity was altered with an average change in phase of -2 hours, while in

other cases it remained unchanged (Figure 3). This variation in the change of locomotor activity

patterns may reflect differences in the perception of temperature by the circadian system of

foragers. Although the mechanisms for light and temperature input to the clock of honey bees

remain to be elucidated, a study in *Drosophila melanogaster* larvae shows that light and

temperature signaling occurs via alternate mechanisms (Picot, Klarsfeld, Chélot, Malpel, &

Rouyer, 2009). Further studies examining the pathways of light and temperature to the circadian

rhythms in honey bees may provide insight into the observed differences in the response to

temperature shifts and possible roles of temperature entrainment in nature.

In conclusion, temperature oscillations as observed in the colony serve as a Zeitgeber for

the honey bee circadian rhythms. Future studies will explore the importance of temperature as a

time giver in the wild. Individual differences in the phase of circadian rhythms are loosely

reminiscent to those of circadian chronotypes in human populations and with further studies,

honey bees could be a potential model for these differences in human populations. Research into

the functional relevance of these phase differences in the colony may provide a better

understanding of the evolution of circadian rhythms and honey bees as a eusocial species. Taken

together, temperature plays several important roles in the honey bee colony that range a

developmental to the entrainment of circadian rhythms. Future experiments should not ignore

temperature conditions since as shown in this and previous studies, environmental temperature

can have major consequences on honey bee behavior.

Acknowledgements:

We would like to thank Dr. Arian Avalos and Emmanuel Rivera, for help with the

experiments. Thanks Dr. Luis de Jesus for their comments and suggestions. We would also like

to recognize the director, Manuel Diaz and the personnel of the Gurabo Experimental Agriculture

Station of the University of Puerto Rico at Mayaguez for use of facilities at ''Casa Amarilla''. This

work was sponsored by the National Science Foundation (NSF) awards 1026560, 1633184,

1707355 and the National Institute of Health (NIH) 2R25GM061151-13, P20GM103475 .

References:

Beier, W. (1968). Beeinflussung der inneren Uhr der Bienen durch Phasenverschiebung des
Licht-Dunkel-Zeitgebers. *Z Bienenforsch*, *9*, 356–378. JOUR.

Beier, W., & Lindauer, M. (1970). Der Sonnenstand als Zeitgeber für die Biene. *Apidologie*, *1*(1),
5–28. JOUR.

Blau, J. (2001). The Drosophila circadian clock: what we know and what we don't know.
Seminars in Cell & Developmental Biology, *12*(4), 287–293.
http://doi.org/10.1006/scdb.2001.0256

Buttel-Reepen, H. (1900). *Sind die bienen reflexmaschinen?: Experiementelle beiträge zur*
biologie der honigbiene. BOOK, Arthur Georgi.

Cyran, S., Buchsbaum, A., Reddy, K., Lin, M.-C., Glossop, N. R. J., Hardin, P. E., … Blau, J.
(2003). vrille , Pdp1 , and dClock Form a Second Feedback Loop in the Drosophila
Circadian Clock. *Cell*, *112*, 329–341. Retrieved from
http://www.sciencedirect.com/science/article/pii/S0092867403000746

Detrain, C., Deneubourg, J. L., & Pasteels, J. M. J. M. (1999). *Information Processing in Social*
Insects. Springer Science & Business Media. Retrieved from
https://books.google.com/books?id=N9iw1nLPaacC&pgis=1

Dunlap, J. C. (1999). Molecular Bases for Circadian Rhythms . *Cell*, *96*, 271–290. Journal
Article.

Forel, A. (1910). Das Sinnesleben der Insekten: eine Sammlung von experimentellen und
kritischen Studien über Insektenpsychologie. JOUR.

Fuchikawa, T., & Shimizu, I. (2007). Effects of temperature on circadian rhythm in the Japanese
honeybee, Apis cerana japonica. *Journal of Insect Physiology*, *53*(11), 1179–1187. Journal
Article. http://doi.org/10.1016/j.jinsphys.2007.06.013

Galindo-Cardona, A., Carolina Monmany, A., Moreno-Jackson, R., Rivera-Rivera, C., Huertas-
Dones, C., Caicedo-Quiroga, L., & Giray, T. (2012). Landscape analysis of drone
congregation areas of the honey bee, Apis mellifera. *Journal of Insect Science*, *12*. Journal
Article.

Gardner, M. J., Hubbard, K. E., Hotta, C. T., Dodd, A. N., & Webb, A. A. (2006). How plants tell
the time. *Biochem J*, *397*(1), 15–24. Journal Article. http://doi.org/10.1042/BJ20060484

Giannoni-Guzmán, M. A., Avalos, A., Marrero Perez, J., Otero Loperena, E. J., Kayım, M.,
Medina, J. A., … Agosto-Rivera, J. L. (2014). Measuring individual locomotor rhythms in
honey bees, paper wasps and other similar-sized insects. *The Journal of Experimental*
Biology, *217*(Pt 8), 1307–15. http://doi.org/10.1242/jeb.096180

Giray, T., Galindo-Cardona, A., & Oskay, D. (2007). Octopamine influences honey bee foraging
preference. *Journal of Insect Physiology*, *53*(7), 691–698. Journal Article.
http://doi.org/10.1016/j.jinsphys.2007.03.016

Goodwin, R. M., & Lewis, D. (1987). Honeybees use a biological clock to incorporate sun
positions in their waggle dances after foraging under heavy overcast skies. *New Zeland*
Entomologist, *10*, 138–140. Journal Article.

Heinrich, B. (1980). Mechanisms of body-temperature regulation in honeybees, Apis mellifera:
II. Regulation of thoracic temperature at high air temperatures. *Journal of Experimental*
Biology, *85*(1), 73–87. Retrieved from http://jeb.biologists.org/cgi/content/abstract/85/1/73

Heinrich, B. (1993). *The Hot-Blooded Insects*. Berlin, Heidelberg: Springer Berlin Heidelberg.
http://doi.org/10.1007/978-3-662-10340-1

HEINRICH, B. (1981). The Mechanisms and Energetics of Honeybee Swarm Temperature
Regulation. *J. Exp. Biol.*, *91*(1), 25–55. Retrieved from

http://jeb.biologists.org/content/91/1/25.short

Himmer, A. (1927). Ein Beitrag zur Kenntnis des Wärmehaushalts im Nestbau sozialer
Hautflügler. *Zeitschrift für vergleichende Physiologie*, *5*(2), 375–389.
http://doi.org/10.1007/BF00338020

Hut, R. A., & Beersma, D. G. (2011). Evolution of time-keeping mechanisms: early emergence
and adaptation to photoperiod. *Philosophical Transactions of the Royal Society of London.*
Series B, Biological Sciences, *366*(1574), 2141–2154. Journal Article.
http://doi.org/10.1098/rstb.2010.0409

Jones, J. C., Myerscough, M. R., Graham, S., & Oldroyd, B. P. (2004). Honey bee nest
thermoregulation: diversity promotes stability. *Science*, *305*(5682), 402–404. Journal
Article. http://doi.org/10.1126/science.1096340

Ko, C. H., & Takahashi, J. S. (2006). Molecular components of the mammalian circadian clock.
Hum Mol Genet, *15 Spec No*, R271-7. Journal Article. http://doi.org/10.1093/hmg/ddl207

Körner, I. (1940). Zeitgedächtnis und Alarmierung bei den Bienen. *Zeitschrift Für Vergleichende*
Physiologie, *27*(4), 445–459. JOUR.

Kraus, F. B., Gerecke, E., & Moritz, R. F. a. (2011). Shift work has a genetic basis in honeybee
pollen foragers (Apis mellifera L.). *Behavior Genetics*, *41*(2), 323–328. Journal Article.
http://doi.org/10.1007/s10519-010-9382-9

Kronenberg, F., & Heller, H. C. (1982). Colonial thermoregulation in honey bees (Apis mellifera).
Journal of Comparative Physiology ? B, *148*(1), 65–76. http://doi.org/10.1007/BF00688889

Moore, D. (2001). Honey bee circadian clocks: behavioral control from individual workers to
whole-colony rhythms. *Journal of Insect Physiology*, *47*, 843–857. Journal Article.

Moore, D., & Doherty, P. (2009). Acquisition of a time-memory in forager honey bees. *Journal of*
Comparative Physiology. A, Neuroethology, Sensory, Neural, and Behavioral Physiology,
195(8), 741–751. Journal Article. http://doi.org/10.1007/s00359-009-0450-7

Moore, D., & Rankin, M. A. (1985). Circadian locomotor rhythms in individual honey bees.
Physiological Entomology, *10*, 191–197. Journal Article.

Moore, D., & Rankin, M. A. (1993). Light and Temperature entrainment of a locomotor rhythm in
honeybees. *Physiological Entomology*, *18*, 271–278. Journal Article.

Moore, D., Siegfried, D., Wilson, R., & Rankin, M. A. (1989). The Influence of Time of Day on
the Foraging Behavior of the Honeybee, Apis mellifera. *Journal of Biological Rhythms*, *4*(3),
305–325. Journal Article.

Panda, S., Hogenesch, J. B., & Kay, S. a. (2002). Circadian rhythms from flies to human.
Nature, *417*(6886), 329–35. http://doi.org/10.1038/417329a

Picot, M., Klarsfeld, A., Chélot, E., Malpel, S., & Rouyer, F. (2009). A role for blind DN2 clock
neurons in temperature entrainment of the Drosophila larval brain. *The Journal of*
Neuroscience : The Official Journal of the Society for Neuroscience, *29*(26), 8312–20.
http://doi.org/10.1523/JNEUROSCI.0279-08.2009

Renner, M. (1959). Über ein weiteres Versetzungsexperiment zur Analyse des Zeitsinnes und
der Sonnenorientierung der Honigbiene. *Zeitschrift Für Vergleichende Physiologie*, *42*(5),
449–483. JOUR.

Renner, M. (1960). The contribution of the honey bee to the study of time-sense and
astronomical orientation. In *Cold Spring Harbor symposia on quantitative biology* (Vol. 25,

pp. 361–367). CONF, Cold Spring Harbor Laboratory Press.

Seeley, T. D. (1974). Atmospheric carbon dioxide regulation in honey-bee (Apis mellifera)
colonies. *Journal of Insect Physiology*, *20*(11), 2301–2305. http://doi.org/10.1016/0022-
1910(74)90052-3

Seeley, T. D. (1995). The Wisdom of the Hive. *Harvard University Press, ….* Retrieved from
http://scholar.google.com/scholar?hl=en&btnG=Search&q=intitle:The+Wisdom+of+the+Hiv
e#2

Spangler, H. G. (1972). Daily Activity Rhythms of Individual Worker and Drone Honey Bees.
Annals of the Entomological Society of America, *65*(5), 1073–1076. Journal Article.

Takahashi, J. (1999). Genetic Analysis of Daily Activity in Humans and Mice, 387567. Retrieved
from
http://oai.dtic.mil/oai/oai?verb=getRecord&metadataPrefix=html&identifier=ADA387567

Tautz, J., Maier, S., Groh, C., Rossler, W., & Brockmann, A. (2003). Behavioral performance in
adult honey bees is influenced by the temperature experienced during their pupal
development. *Proceedings of the National Academy of Sciences of the United States of
America*, *100*(12), 7343–7347. Journal Article. http://doi.org/10.1073/pnas.1232346100

von Buttel-Reepen, H. B. (1903). *Die stammesgeschichtliche entstehung des bienenstaates
sowie beiträge zur lebensweise der solitären u. sozialen bienen (hummeln, meliponinen,
etc.).: Vortrag gehalten auf dem Zoologen-kongress in Giessen (1902). Stark erweitert, mit
anmerkungen und zusätz*. BOOK, G. Thieme.

von Frisch, K. (1940). Die Tänze und das Zeitgedächtnis der Bienen im Widerspruch.
Naturwissenschaften, *28*(5), 65–69. JOUR.

von Frisch, K. (1967). *The Dance Language and Orientation of Bees*. Book, Cambridge, MA.:
Harvard University Press.

von Frisch, K., Aschoff, J., Frisch, B., & Aschoff, J. (1987). Circadian rhythms in honeybees:
entrainment by feeding cycles. *Physiological Entomology*, *12*(1), 41–49. Journal Article.
http://doi.org/10.1111/j.1365-3032.1987.tb00722.x

Wahl, O. (1932). Neue Untersuchungen über das Zeitgedächtnis der Bienen. *Zeitschrift für
vergleichende Physiologie*, *16*(3), 529–589. http://doi.org/10.1007/BF00338333

Wahl, O. (1933). Beitrag zur Frage der biologischen Bedeutung des Zeitgedächtnisses der
Bienen. *Zeitschrift für vergleichende Physiologie, 18*(4), 709–717.
http://doi.org/10.1007/BF00338365