A toolkit for studying *Varroa* genomics and transcriptomics: Preservation, extraction, and sequencing library preparation

Nonno Hasegawa[1], Maeva Techer[1], Alexander S. Mikheyev[1,2]

1 Okinawa Institute of Science and Technology, 1919-1 Tancha Onna-son, 904-0495 Okinawa, Japan
2 Australian National University, Canberra, ACT, 2600, Australia

Corresponding Authors
Nonno Hasegawa nonno.hasegawa3@oist.jp
Alexander S. Mikheyev alexander.mikheyev@anu.edu.au

Abstract

Background:

The honey bee parasite, *Varroa destructor*, is the leading cause of global honey bee population declines. In addition to being an obligate ectoparasitic mite, *Varroa* carries several viruses that infect honey bees and act as the proximal causes of colony collapses. Yet, until recently, the study of *Varroa* has been limited by the availability of genomic tools. Lab- and field-based methods exploiting such methods are still nascent. This study developed a set of methods for preserving *Varroa* DNA and RNA from the field to the lab and for processing them into sequencing libraries. We performed preservation testing experiments in which *Varroa* mites were immersed in TRIzol, RNAlater, and absolute ethanol, as well as preservation periods up to 21 days post-treatment to assess DNA and RNA integrity.

Results:

For both DNA and RNA, mites preserved in TRIzol and RNAlater at room temperature degraded within 10 days post-treatment; mites preserved in ethanol at room temperature and 4°C remained intact through 21 days. *Varroa* mite DNA and RNA libraries were created and sequenced for ethanol preserved samples, 15 and 21 days post-treatment. All DNA sequences mapped to *V. destructor* at above 95% on average,

while RNA sequences mapped to *V. destructor*, but also sometimes to high levels of the deformed wing

virus, and other taxa.

Conclusion:

Ethanolic preservation of field-collected mites is inexpensive, simple and allows them to be shipped and

processed successfully in the lab for a wide variety of sequencing applications. It appears to preserve

RNA from both *Varroa* and at least some of the viruses it vectors.

Keywords (max 10):

Genomics, transcriptomics, *Varroa destructor*, preservation method, sequencing

Background

Honey bees are one of the most economically significant agricultural resources, contributing to crop

pollination as well as providing products such as honey, propolis, and beeswax, which also contribute to

the economy [1]. The global decline of honey bee populations harms agricultural output, as some studies

suggest honey bees contribute greatly to the agricultural economy. For instance, the agricultural

department of Canada reported that in 2016, the honey bee pollination contributed 2.5 billion CAD to

Canadian crops [2]. While honey bee declines are multifactorial, they have been accelerated by the

pandemic spread of the ectoparasitic mite, *Varroa destructor*, which jumped hosts from the closely

related eastern honey bee (*Apis cerana*) [1, 3]. Until recently, few genomic resources for *Varroa* existed,

and how it evolves post-host switch remains poorly understood [4, 5]. Furthermore, ˆ*Varroa* mites vector

several viruses that impair the honey bee immune system, cause underdevelopment in honey bees, as well

as cognitive impairment [6]. Viruses transmitted by *Varroa* are believed to be the primary driver of the

decline in honey bee populations around the world, exacerbating parasitism by *Varroa* mites themselves

[1]. Many *Varroa*-associated viruses have been identified, most notably deformed wing virus (DWV) [1, 7–9], acute bee paralysis virus [8, 9], Kashmir bee virus [8, 9], and the black queen cell virus [8, 9]. However, much remains unknown about the biology of both *Varroa* and viruses it carries, as standard molecular methods to study them using genomic and transcriptomic tools have not been established [9].

The majority of honey bee viruses are single-stranded, positive-sense RNA viruses. These commonly remain dormant, leaving the bees asymptomatic; thus, field diagnosis of honey bee viruses remains challenging [6, 9]. To understand viral loads of colonies, beekeepers must send samples to facilities that are equipped with proper instrumentation for diagnosis [9]. The optimal sampling method for live organisms for laboratory analysis is snap freezing and transporting with dry ice, which is frequently not possible for field workers and beekeepers [10]. Previous work exploring alternative storage conditions for honey bee RNA yielded degraded RNA in 70% ethanol, and whole honey bee RNA started degrading within a week in RNAlater [10]. Campbell *et al.* [11] explored preservation of *Varroa* mites in RNAlater and its effect on RNA integrity. That study suggested that only when the mites are pierced does RNA remain intact, and did not explore other preservation solutions. Thus, there is limited published research regarding storage conditions of *Varroa* mites and their effect on RNA and DNA integrity generally. Also, existing protocols for DNA and RNA extraction require pooling several mites to obtain enough material [11, 12]. We propose a new method to extract both DNA and RNA from a single mite, sufficient in both quality and quantity for downstream analysis, such as next-generation sequencing. This method allows analysis of individuals, which can be used to map viruses present in each individual rather than pooled samples. Standardizing a method to analyze viruses present in each *Varroa* mite will allow biogeographical mapping to help visualize and track virus trends on a global scale. Here, we explore *Varroa* mite preservation conditions in TRIzol, RNAlater, and absolute ethanol for storage periods of up to 21 days, and their effects on DNA and RNA quality by mapping to the *V. destructor* reference genome [5] and transcriptome. We propose a more affordable and feasible method for fieldworkers who have limited immediate access to laboratory facilities or equipment.

Results

DNA Quality and Library preparation

Extracted *Varroa* mite DNA exhibited no trend for A260/280 values among the four treatments. A260/280 values among all samples ranged from 1.76 to 3.35, with a mean of 2.76, indicating high quality. We inspected samples treated with TRIzol and RNAlater using DNA electrophoresis, and found that they were heavily degraded 10 days post-treatment. Mean total DNA yield was 49.7 ± 58 (s.d.) ng. DNA generally remains intact over days or weeks of storage at room temperature; thus, our target for downstream analysis was the control group and ethanol groups from 21 days post-treatment to further analyze the quality by library prep, sequencing, and mapping to *V. destructor*. The DNA library was successfully produced using the Nextera XT library preparation kit (catalog #FC-131-1096), uniquely indexed for sequencing

RNA Quality and Library preparation

Varroa mite total RNA was extracted with a mean yield of $2{,}157 \pm 2{,}570$ (s.d.) ng. RNA electrophoresis showed that RNA in samples preserved in ethanol remained intact even at room temperature, while RNAlater and TRIzol preserved samples had degraded heavily in both 15- and 21-day samples (Fig. 1). Control group samples, which were snap-frozen and stored at -80°C, were also well-preserved. As RNAlater fails to penetrate the mite unless the exoskeleton is punctured, degradation of RNA was expected [11]. *Varroa* mites float to the surface in TRIzol, as they are less dense than the solution; thus, RNA degradation was also expected in these samples. Total RNA libraries were prepared using the NEB low-input RNA library preparation kit (catalog #E6420), uniquely indexed for sequencing

DNA and RNA Sequence Mapping

Varroa mite DNA libraries acquired on average, ~231,000 ± 49,000 (s.d.) sequences while RNA libraries

averaged approximately 447,000 ± 248,000 (s.d.) sequences per library. Given the short-read length used

in sequencing, inspection with FastQC revealed that there were essentially no adapters and that sequence

quality was consistently high, so raw reads were not processed further. DNA libraries of mites preserved

in ethanol for 15 and 21 days at room temperature and 4°C aligned to the *V. destructor* reference genome

with a median of 95.49 ± 1.16 (s.d.) % (Fig. 2). Of the mapped reads, approximately 5.5 ± 0.74 (s.d.) %

were PCR duplicates, an artifact of library preparation. Since ribosomal RNA depletion was not

performed before library preparation, we also conducted alignments to 18S (accession FJ911866.1) and

28S (FJ911801.1) rRNA to ensure the ribosomal RNA was not abundant, possibly swamping the RNA

sample. On average there was less than 0.001% rRNA, allowing us to proceed with general mapping to

Varroa mite gene models. RNA sequences from *Varroa* mites stored in ethanol at both room temperature

and 4°C mapped inconsistently to *V. destructor*. Mapping to the *V. destructor* reference library varied

from as little as 2% to as high as 76%. We used Kraken2 [13] to classify reads to other organisms, which

mapped to various species (Fig. 2). Taxon classification with Kraken2 suggested higher mapping to DWV

for individuals that didn't map well to the *V. destructor* reference library.

Discussion

Transporting samples from the field to the lab for RNA processing often involves considerable stress and

logistical complexity. To facilitate this process, we assessed effects of different preservation conditions

on *Varroa* mite total DNA and RNA integrity. Surprisingly, *Varroa* mite total DNA and RNA libraries

mapped well to *V. destructor* preserved well for 21 days post-treatment (possibly longer) when stored in

absolute ethanol at room temperature and 4°C. This was a significant finding since for honey bees [10]

and spiders [14], ethanol does not preserve RNA well. While butterfly [15], honey bee [10], and *Varroa*

mite [11] RNA preserved well in RNAlater when specimens were ground, sliced, or pierced, intact

specimens generally degraded within a few days post-treatment. Similar to these findings, we found that

intact *Varroa* mite DNA and RNA stored in TRIzol and RNAlater started to degrade within 10 days, and

possibly earlier.

As previously reported [16], DNA preserves well over longer periods compared to RNA, as long as the

solution penetrates the specimen. *Varroa* mites floated to the surface in TRIzol; thus, mite DNA was not

well preserved. Similarly, mites immersed in RNAlater were not well preserved because the solution does

not penetrate the cuticle. Through RNA electrophoresis, we found that *Varroa* mite total RNA was

completely degraded 2 weeks post-treatment (possibly earlier) when stored in TRIzol or RNAlater (Fig.

1). Surprisingly, RNA samples 21 days post-treatment immersed in ethanol at room temperature remained

largely intact, allowing appropriate NGS library preparation and analysis. Both ethanol RNA samples at

room temperature and 4°C mapped either to *V. destructor* or DWV, with a minimum amount of rRNA.

Chen *et al*. [10] stated that whole honey bee RNA did not preserve well in RNAlater without slicing or

crushing 1-week post-treatment, which is consistent with our results. Several other studies also suggest

that spiders [14] and butterflies [15] preserve well in RNAlater when specimens are crushed. However,

RNA integrity varies when stored in ethanol between organisms. Spiders only preserved well when

crushed, and not when immersed in ethanol as whole organisms [14].

Mites must be punctured [11] if these solutions are to successfully preserve sample RNA, as for honey

bees [10]. However, it is not practical for beekeepers to puncture each specimen, particularly in the field,

and TRIzol and RNAlater are only available through vendors in laboratory reagents, and are much more

expensive than ethanol. For these reasons, we recommend the use of ethanol, which is readily available

from chemical suppliers. *Varroa* mites provided high-quality DNA and RNA for NGS analysis for weeks

when stored in ethanol at room temperature or 4°C, which should be sufficient for collection and

shipping. Though we terminated the experiment at three weeks, it seems likely that both DNA and RNA

are stable for much longer, given the minimal degradation observed at this time.

Although RNA libraries mapped well to *V. destructor* and DWV, there was still a portion of RNA

sequence that was not classified. These unclassified sequences may contain new, unidentified RNA

viruses that the *Varroa* mites are vectoring. These unclassified sequences also may belong to organisms

that have not been sequenced or may represent artifacts of library preparation. Though it is beyond the

scope of our current study, and not entirely feasible with 50-bp reads used in these experiments, exploring

the unclassified region of RNA sequences will be beneficial in understanding the complex host-parasite

relationship of honey bees and *Varroa* mites, opening new avenues to honey bee health.

There are several limitations we must take into account. Most importantly, our sample size is limited to

n=6 per treatment per time point. However, the number of samples we processed for library preparation

and sequencing resulted in high-quality mapping to the reference genome, and none of the samples

deviated strongly from the rest, suggesting we have a good representation of the treatments. The quality

of ethanol must also be considered. In laboratory experiments, we require high-grade alcohol for quality

control of samples. So, while we used high-grade absolute ethanol in this study, this quality of ethanol is

not readily available worldwide. We suspect that lower quality ethanol, particularly methanol

contamination, may harm preservation.

Although both DNA and RNA integrity were retained at room temperature when immersed in ethanol, it

is more realistic and practical to store and ship samples within 2 weeks (15 days post-treatment). We

advise beekeepers and field workers to store mite samples in the refrigerator when possible, and at room

temperature when this is not possible. Samples may also be shipped without refrigeration, reducing

shipping costs. Additionally, as we observed 21 days of stable storage, shipping delays should be

considered if RNA integrity must be maintained for longer periods, for high-quality NGS and other

downstream analyses. In short, we found that a possible alternative to snap freezing *Varroa* mites and

storing at -80°C is immersion in absolute ethanol for up to 21 days. By immersing in absolute ethanol,

Varroa mite DNA and RNA are well preserved at both room temperature and at 4°C, which allow more
flexible sampling and storage conditions. We believe that methods presented in this study will lead to
insights in *Varroa* genomics and population biology, as well as facilitating studies of the viruses it
vectors.

Conclusion

We found that *Varroa* mite DNA and RNA were adequately preserved in absolute ethanol for up to 21
days, and produced high quality DNA and RNA libraries when sequenced, mapping to the *Varroa*
genome and to other taxa. On the contrary, when *Varroa* mites were preserved in TRIzol and RNAlater,
the mite DNA and RNA degraded within the first 10 days, possibly earlier, likely as a result of poor
penetration through the exoskeleton Ethanolic preservation of *Varroa* mites is inexpensive and uses a
readily available reagent, thus allowing specimens to be shipped and processed for a wide variety of
sequencing applications. In addition, ethanol also preserves viruses that *Varroa* vectors, most notably
DWV. We propose that ethanolic preservation can replace cryopreservation, providing a more tractable
method for preserving DNA and RNA quality.

Material and Methods

Mite Collection

In March 2020, *Varroa* mites were collected from managed hives in Onna village, Okinawa by removing
honey bees from 2 frames onto a tray with icing sugar, and shaking the tray to remove mites from honey
bees (Fig. 3). From the site of collection to the laboratory, mites were in icing sugar. Once at the
laboratory, we discarded dead mites. Mites were divided into the following treatment preservation
conditions: 1) snap-frozen and stored at -80°C 2), immersed in absolute ethanol at room temperature, 3)
immersed in absolute ethanol at 4°C, 4) immersed in TRIzol at room temperature, and 5) immersed in
RNAlater at room temperature. When immersed in a solution, 500 µL was used and each treatment

consisted of 6 mites placed in separate tubes. Each treatment group was subjected to DNA and RNA extractions at intervals of 5, 10, 15, and 21 days for subsequent DNA and RNA quantity and quality evaluations. Snap-freezing and storage at -80°C was chosen for the control group as it is the widely used method for specimen preservation [6, 14, 15]

Mite Preparation for Extractions

Mites were removed from the solution in which they were stored to unused 1.5-mL tubes (Eppendorf catalog #0020125215) and chilled in liquid nitrogen. A clean, autoclaved pestle (Sigma Aldrich catalog #Z359947) was also chilled in liquid nitrogen to grind each mite in its tube. By visual inspection, ground mite powder that remained on the pestle was used for DNA extraction, ensuring that about half of the mite homogenates remain in the tubes for RNA extraction (Fig. 3).

DNA Extraction

Mite DNA was extracted by using the QIAamp DNA Micro kit according to the manufacturer's protocol. DNA quantity was determined using the Qubit fluorometer, and quality was evaluated by the A280 and A260 ratio using Nanodrop 2000c. Eluted DNA was stored at -20°C until further applications.

DNA Extraction

Mite DNA was extracted using the QIAamp DNA Micro kit, according to the manufacturer's protocol. DNA quantity was determined using a Qubit fluorometer, and quality was evaluated by the A280 / A260 ratio using a Nanodrop (Thermo Scientific). Eluted DNA was stored at -20°C until further applications.

RNA Extraction

Mite RNA was extracted using TRIzol according to the manufacturer's protocol. The protocol was modified to use 500 µL of TRIzol instead of 1 mL, and half the volume of all other reagents. Total RNA quality and quantity were evaluated using the Nanodrop.

Library Preparation

DNA Library Preparation

DNA libraries were prepared using a Nextera XT DNA Library Prep kit, according to the manufacturer's protocol, with optimization for mite DNA, using reagents 20% of their specified volume. DNA was visualized by running electrophoresis on 1% agarose gels, for 20 min at 135 V

RNA Library Preparation

RNA libraries were prepared using an NEBNext Single Cell/ Low Input RNA Library kit, according to the manufacturer's protocol. Purified cDNA was indexed with i5 and i7 primers, then purified and size selected for a range of 400 to 2000 bp using 11% and 11.5% PEG and DynaBeads [17].

Both DNA and RNA libraries, as well as total RNA, were analyzed with a Bioanalyzer 2100 or a 4200 Tapestation. For Bioanalyzer, high-sensitivity DNA kits and RNA pico kits were used, while for the Tapestation, a high-sensitivity D5000 kit was used.

Sequencing and Analysis

A MiSeq (Illumina) was used to perform both DNA and RNA sequencing, at 50 cycles and read with single read-only. We were interested in validating the protocol and estimating mapping percentages, so longer read lengths or higher coverage were not necessary. Raw sequence data from MiSeq were first analyzed with FastQC [18] for quick quality control to see if adapter removal was necessary and to ensure

that sequenced data was of sufficient quality. DNA sequence data were then analyzed using Bowtie2 [19],

Samtools [20], and Picard tools [21] to map to the reference genome (Vdes_3.0) and to identify

duplicates. RNA sequence data were analyzed using Bowtie2 and Samtools, and then taxonomically

classified using Kraken [13].

Data availability

Raw sequence data are available under DDBJ/NCBI BioProject PRJDB10252.

Abbreviations

252 DWV - deformed wing virus

Declarations

Ethics approval and consent to participate. Not applicable.

Consent for publication. All authors have consented to publish this work.

Competing interests. None.

Funding. ASM was supported by a Future Fellowship from the Australian Research Council
(FT160100178) and a Kakenhi Grant-in-Aid for Scientific Research from the JSPS (18H02216). This
work as additionally funded by the Okinawa Institute of Science and Technology Graduate University

Author contributions. All authors designed the study. NH and MT performed the laboratory work. NH
performed the analysis and wrote the first manuscript draft, to which all authors contributed.

Acknowledgements. We are grateful to Miyuki Suenaga for help in the lab, the sequencing centre at
OIST, and to the Honey & Coral project of the Onna Village for help in maintaining the bee colonies. We
are grateful to Alexandra Sébastien for conducting pilot experiments that helped guide this project.

REFERENCES

1. McMenamin AJ, Genersch E. Honey bee colony losses and associated viruses. Current Opinion in Insect Science. 2015;8:121–
9.

2. Mukezangango J, Page S. Statistical overview of the Canadian honey and bee industry and the economic contribution of honey
bee pollination, 2016. Horticulture and Cross Sectoral Division, Agriculture and Agri-Food Canada. 2017.

3. Vanengelsdorp D, Meixner MD. A historical review of managed honey bee populations in Europe and the United States and
the factors that may affect them. J Invertebr Pathol. 2010;103 Suppl 1:S80–95.

4. Eliash N, Mikheyev A. Varroa mite evolution: a neglected aspect of worldwide bee collapses? Curr Opin Insect Sci.
2020;39:21–6.

5. Techer MA, Rane RV, Grau ML, Roberts JMK, Sullivan ST, Liachko I, et al. Divergent evolutionary trajectories following
speciation in two ectoparasitic honey bee mites. Communications Biology. 2019;2:357.

6. Navajas M, Migeon A, Alaux C, Martin-Magniette M, Robinson G, Evans J, et al. Differential gene expression of the honey
bee Apis mellifera associated with Varroa destructor infection. BMC Genomics. 2008;9:301.

7. de Miranda JR, Genersch E. Deformed wing virus. J Invertebr Pathol. 2010;103 Suppl 1:S48–61.

8. Remnant EJ, Shi M, Buchmann G, Blacquière T, Holmes EC, Beekman M, et al. A Diverse Range of Novel RNA Viruses in
Geographically Distinct Honey Bee Populations. J Virol. 2017;91. doi:10.1128/JVI.00158-17.

9. Dietemann V, Nazzi F, Martin SJ, Anderson DL, Locke B, Delaplane KS, et al. Standard methods for varroa research. J Apic
Res. 2013;52:1–54.

10. Chen Y, Evans J, Hamilton M, Feldlaufer M. The influence of RNA integrity on the detection of honey bee viruses:
molecular assessment of different sample storage methods. J Apic Res. 2007;46:81–7.

11. Campbell EM, McIntosh CH, Bowman AS. A Toolbox for Quantitative Gene Expression in Varroa destructor: RNA
Degradation in Field Samples and Systematic Analysis of Reference Gene Stability. PLoS One. 2016;11:e0155640.

12. Gusachenko ON, Woodford L, Balbirnie-Cumming K, Campbell EM, Christie CR, Bowman AS, et al. Green Bees: Reverse
Genetic Analysis of Deformed Wing Virus Transmission, Replication, and Tropism. Viruses. 2020;12. doi:10.3390/v12050532.

13. Wood DE, Salzberg SL. Kraken: ultrafast metagenomic sequence classification using exact alignments. Genome Biol.
2014;15:R46.

14. Kono N, Nakamura H, Ito Y, Tomita M, Arakawa K. Evaluation of the impact of RNA preservation methods of spiders for de
novo transcriptome assembly. Mol Ecol Resour. 2016;16:662–72.

15. Gayral P, Weinert L, Chiari Y, Tsagkogeorga G, Ballenghien M, Galtier N. Next-generation sequencing of transcriptomes: a
guide to RNA isolation in nonmodel animals. Mol Ecol Resour. 2011;11:650–61.

16. Post RJ, Flook PK, Millest AL. Methods for the preservation of insects for DNA studies. Biochem Syst Ecol. 1993;21:85–92.

17. Tin MM-Y, Economo EP, Mikheyev AS. Sequencing degraded DNA from non-destructively sampled museum specimens for
RAD-tagging and low-coverage shotgun phylogenetics. PLoS One. 2014;9:e96793.

18. Bioinformatics B. FastQC: A quality control tool for high throughput sequence data. Babraham Bioinformatics. 2010.
http://www.bioinformatics.babraham.ac.uk/projects/fastqc/. Accessed 5 Aug 2020.

19. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods. 2012;9:357–9.

20. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence Alignment/Map format and SAMtools.
Bioinformatics. 2009;25:2078–9.

21. Broad Institute. Picard Toolkit. Broad Institute, GitHub Repository. 2020. http://broadinstitute.github.io/picard/. Accessed 5
Aug 2020.

326   **Figures**

327
Figure 1: Representative bioanalyzer result of *Varroa* mite total RNA, extracted 21 days post-treatment.

Control samples that were snap-frozen and stored at -80°C show minimal noise and a clean 18S peak,

while ethanol samples at room temperature and 4°C also showed a similar 18S peak; however, with more

degradation products. Mites stored in RNAlater and TRIzol had degraded and did not show a peak at 18S,

indicating that RNA preservation was not successful. This suggests that weeks of storage in ethanol, even

at room temperature, have little effect on RNA integrity.

Figure 2: RNA sequence alignment of *Varroa* mite total RNA to *V. destructor*, DWV, and other viruses.

Some samples had fewer *V. destructor* reads, compared to DWV. Other viruses remained at a lower

mapping percentage, and some reads did not map to any reference genome. Reads that remained

unclassified may be organisms that have not yet been sequenced, but could also be library preparation

artifacts, new RNA viruses for which *Varroa* mites serve as vectors, or microbes.

Figure 3: Generalized experimental workflow. 1) *Varroa* mites were collected from an apiary located at

the Onna village office in Okinawa, Japan, March, 2020. Mites were shaken off the honey bees by

sprinkling powdered sugar on the frame. Bees were collected into a tray and shaken to remove mites. The

tray containing powdered sugar and *Varroa* mites was transported back to the laboratory, within 30

minutes, discarding dead mites on arrival. 2) Individual mites were placed in separate 1.5-mL

microcentrifuge tubes and 3) were immersed in 500 µL of a preservative solution or snap-frozen and

stored at -80 °C until processing. 4) Incubation periods were 5, 10, 15, and 21 days in respective

preservation methods. 5) Preservation solution was discarded after the incubation period, and 6) samples

were immersed in liquid nitrogen for a minute, then 7) crushed with a sterile pestle that was also

immersed in liquid nitrogen. Each sample was split in two tubes, 8a) mite dust on the pestle was washed

with ATL buffer into a new tube and used for DNA extraction with a QIAamp DNA extraction kit and

8b) TRIzol was added to the tube containing mite dust and subjected to RNA extraction using TRIzol.