Spike protein mutational landscape in India: Could Muller's ratchet be a future game-

changer for COVID-19?

Rachana Banerjee[1], Kausik Basak[1], Anamika Ghosh[2], Vyshakh Rajachandran[3], Kamakshi

Sureka[1], Debabani Ganguly[1*], Sujay Chattopadhyay[1*]

[1]Centre for Health Science and Technology, JIS Institute of Advanced Studies and Research

Kolkata, JIS University, Kolkata, 700091, West Bengal, India

[2]Department of Chemistry, Indian Institute of Engineering Science and Technology, Shibpur,

Howrah, 711103, West Bengal, India

[3]School of Biotechnology, Amrita Vishwa Vidyapeetham, Kollam, 690525, Kerala, India

[*]Corresponding authors.

Address for correspondence:

Email: sujayc@jisiasr.org, debabani@jisiasr.org

Abstract

The dire need of effective preventive measures and treatment approaches against SARS-

CoV-2 virus, causing COVID-19 pandemic, calls for an in-depth understanding of its

evolutionary dynamics with attention to specific geographic locations, since lockdown and

social distancing to prevent the virus spread could lead to distinct localized dynamics of virus

evolution within and between countries owing to different environmental and host-specific

selection pressures. To decipher any correlation between SARS-CoV-2 evolution and its

epidemiology in India, we studied the mutational diversity of spike glycoprotein, the key

player for the attachment, fusion and entry of virus to the host cell. For this, we analyzed the

sequences of 630 Indian isolates as available in GISAID database till June 07, 2020, and

detected the spike protein variants to emerge from two major ancestors – Wuhan-Hu-1/2019

and its D614G variant. Average stability of the docked spike protein – host receptor (S-R)

complexes for these variants correlated strongly (R^2=0.96) with the fatality rates across

Indian states. However, while more than half of the variants were found unique to India, 67%

of all variants showed lower stability of S-R complex than the respective ancestral variants,

indicating a possible fitness loss in recently emerged variants, despite a continuous increase

in mutation rate. These results conform to the sharply declining fatality rate countrywide (>7-

fold during April 11 – June 28, 2020). Altogether, while we propose the potential of S-R

complex stability to track disease severity, we urge an immediate need to explore if SARS-

CoV-2 is approaching mutational meltdown in India.

Author summary

Epidemiological features are intricately linked to evolutionary diversity of rapidly evolving

pathogens, and SARS-CoV-2 is no exception. Our work suggests the potential of average

stability of complexes formed by the circulating spike mutational variants and the human host

receptor to track the severity of SARS-CoV-2 infection in a given region. In India, the

stability of these complexes for recent variants tend to decrease relative to their ancestral

ones, following countrywide declining fatality rate, in contrast to an increasing mutation rate.

We hypothesize such a scenario as nascent footprints of Muller's ratchet, proposing large-

scale population genomics study for its validation, since this understanding could lead to

therapeutic approaches for facilitating mutational meltdown of SARS-CoV-2, as experienced

earlier for influenza A virus.

Introduction

The emergence and rapid global spread of novel severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), the causative agent of the coronavirus disease 2019 (COVID-19), has led to an unprecedented worldwide public health crisis, crossing the species barrier and disseminating rapidly through the global population [1]. Like any other rapidly evolving emerging pathogens, effective preventive measures and treatment approaches call for an in-depth understanding of the evolutionary dynamics of SARS-CoV-2 and its association with the epidemiological data [2, 3], especially with distinct attention to specific geographic regions. Nature and strength of selection pressures acting on the pathogen can vary across different ethnicities, countries, or even among specific administrative regions (such as states or provinces) within a country like India, China, USA, largely because of the implementation of social distancing via lockdown as a preventive measure against the geographic spread of the pathogen between and within countries.

The transmembrane spike glycoprotein protrudes from the viral surface and is responsible for viral attachment, fusion and entrance into the host cells, thereby establishing the infection [4, 5]. The two most notable genetic features of this protein are (a) the binding of its surface unit (S1) to the human cellular receptor angiotensin converting enzyme 2 (hACE2), and (b) the fusion of viral membrane with a host cell membrane via its transmembrane unit (S2). The S1 subunit contains the receptor-binding domains (RBD) supporting the stabilization of the membrane-anchored state of the S2 subunit along with its fusion machinery [6-11]. Some previous works have acknowledged that extensive and irreversible conformational changes stimulate the cleavage of spike protein, thereby activating it for membrane fusion [11-17]. Though it is not yet confirmed if the differences owing to conformational changes are aiding to the expansion of SAR-CoV2 varieties increasing or decreasing its infectivity or transmissibility, but the researchers confirmed the

spike proteins to be the key pathogenic determinant that differentiates SAR-CoV2 from other

SARS-related coronaviruses [17]. It is predicted that mutations in the spike protein increases

or decreases protein glycosylation, thereby enhancing or reducing the viral uptake by host

cells [18-22]. Although the RBD in the spike protein is known as the most important player

to recognize the host receptor, it is highly likely that the region outside of this C-terminal

domain in S1 subunit as well as the domains of S2 subunit also influence the host-receptor

binding allosterically.

The present study analyses spike protein variants from Indian isolates to understand

the evolutionary patterns and driving forces behind the emerging SARS-CoV-2 infection and

its potential epidemiological footprints in India. We detected a strong correlation between the

average stability of complexes formed by the circulating spike protein variants with the host

receptor (S-R complex) and the disease severity of a given location, suggesting the S-R

complex stability as a potential marker to assess the severity of the disease. Importantly,

majority of the emerging variants showed decreased stability, indicating accumulation of

deleterious mutations in the spike protein. This conforms to the declining fatality rates of the

disease countrywide. Could the fixation of these deleterious mutations in the population lead

to mutational meltdown, following Muller's ratchet dynamics of evolution?

Methods

Analysis of sequence diversity and reconstruction of phylogeny

The average pairwise nucleotide diversity (π) and the rates of synonymous (dS) and

nonsynonymous (dN) mutations for the spike protein-coding genes were calculated using

MEGA version X [23]. TimeZone software [24] was used to reconstruct the maximum-

likelihood based phylogeny to map the protein variants and identify convergent amino acid

changes (i.e. repeated independent or phylogenetically unlinked mutations at the same amino

acid positions). The spike gene sequence from Wuhan-Hu-1/2019 genome was used as

reference to detect the orthologs in the sequenced Indian genomes based on a threshold value

of 95% for both nucleotide sequence diversity and gene length coverage.

Identification of spike protein variants unique to India

All 17529 spike protein sequences from worldwide isolates available till May 9, 2020

in the GISAID database (https://www.gisaid.org/) were downloaded. We implemented CD-

HIT Suite [25, 26] to cluster all the spike protein sequences considering 100% amino acid

sequence identity as ortholog clustering criteria, and detected a total of 3706 clusters. Of

these clusters, we considered 3577 clusters matching the complete length of spike protein

(1273 amino acids), using the sequence from Wuhan-Hu-1/2019 (GenBank accession number

MN908947) as reference. These 3577 spike protein sequences were aligned using ClustalW

program [27, 28]. The resulting alignment was compared with Indian mutational variants

mapped in the previous step to distinguish the variants unique to Indian isolates.

Analysis of state-wise diversity of spike protein variants

For each Indian state, we computed the number of spike protein variants and the

frequency of each of those variants. The state-wise calculation of variant diversity was

performed using Simpson's index [29].

Modeling of spike protein – hACE2 complex variants

Variants of the ancestral spike protein (Wuhan-Hu-1/2019) were built by homology

modeling using Swiss modeler [30] with the aid of available templates (residue range: 27-

1147) using Wuhan-Hu-1/2019 isolate as reference [17, 31]. X-ray crystal structure of the

human ACE2 was used from the complex of receptor binding domain (RBD) of spike protein

with hACE2 (PDB code: 6lzg) [32]. We docked the RBD (residue range: 331-524) of spike

protein (residue range: 27-1147) mutants to the binding site of hACE2 using HADDOCK

webserver [33, 34] by providing binding site information [31, 35]. HADDOCK score (often

mentioned as docking score in the text), with some arbitrary unit, signifies a measure

determined by weighted sum of intermolecular interactions, such as electrostatic and van der

Waals interactions between protein and ligand, desolvation energy, restraint violation energy

and the buried surface area upon binding. For each docked complex HADDOCK score was

estimated and VMD [36] was used to visualize the structures.

Results

Two major ancestors circulating in India lead to a burst of spike protein variants.

We identified a total of 630 isolates with complete gene sequences encoding the spike

protein based on the submissions of Indian SARS CoV-2 genome sequences to GISAID till

June 7, 2020 (S1 Table). The samples analyzed were isolated from 17 states and 2 union

territories of India, collectively called as 'states' hereafter. We found a countrywide average

pairwise nucleotide diversity (π) of 0.048±0.02%. The rates of synonymous (or silent) and

nonsynonymous (or amino acid replacement) mutations were found to be 0.097±0.05% and

0.033±0.02% respectively. Phylogenetic analysis showed the Wuhan-Hu-1/2019 variant of

the spike protein as the most ancestral one, as expected, while the D614G variant emerged

from the Wuhan-Hu-1/2019 variant appeared to be another stable variant circulating in the

Indian population. Since both these variants have established themselves in the worldwide

population as two major ancestors of SARS CoV-2 spike protein variants, we here onwards

will refer the Wuhan-Hu-1/2019 and D614G variants as ancestor 1 and ancestor 2

respectively.

Apart from giving rise to ancestor 2, the ancestor 1 led to a total of 20 variants (Fig 1

and S2 Table). Of these, the variant K77M evolved further to yield three more variants

isolated from three different states (Bihar, Tamil Nadu and Telengana), suggesting the

emergence of K77M as another stable variant. On the other hand, the ancestor 2 showed

about twice more diversity by giving rise to 47 variants (Fig 1 and S2 Table). In this ancestor

2 clade, several variants (L5F, T22I, L54F, G261S, T572I, E583D, Q677H, A706S, H1083Q)

indicated their stability in the population via mutating further, giving rise to additional

variants. Besides, although the ancestral variants were predominant in the population

circulating in India, we detected a total of 16 variants of spike proteins that were represented

by multiple isolates (Fig 1), from 2 to as many as 9 isolates, indicating the possible fixation

of some of these variants in the population irrespective of the stability of S-R complex. Of

these, an array of mutations in 8 variants (at positions 5, 54, 78, 558, 574, 583, 677, 1243)

showed their convergent nature, where those mutations at the same positions were

phylogenetically unlinked, i.e. repeated independently (S2 Table). Interestingly, 53% of the

total set of variants detected in Indian population was found unique, i.e. not found in 17529

worldwide genomic isolates analyzed from the GISAID database till May 09, 2020 (S2

Table). We believe that this considerable level of uniqueness could be an expected scenario

in almost all geographical regions where a newly emerging, rapidly evolving viral pathogen

tries to adapt to a new host, and many of these variants might be detrimental to the fitness of

the organism. However, specific positive selection pressures could also play a role in this

mutational pattern which needs to be studied separately in some greater depth.

Figure 1. Schematic representation of the diversity of spike protein variants circulating in India, using maximum likelihood-based phylogeny reconstruction. Each node represents a specific spike protein variant, while the node-size and the number inside depict the frequency of that variant. The red or green color of each arrow indicates the higher or lower stability index respectively of the S-R complex for each variant than the major ancestral variant it emerged from (either Ancestor 1 or Ancestor 2). The black arrows lead to the variants for which the docking scores could not be determined either because of the presence of at least one variation outside the available template region for docking [17, 31], or due to non-existing isolate in the lone hypothetical node with H1083Q mutation denoted by black color. This black node signifies a variant with no available isolate in our dataset, while it gives rise to two derived variants, H1083Q:R78M and H1083Q:E583D, for which representative isolates were available.

The average stability index of S-R complex correlates strongly with the fatality rates in a given location.

We next looked into the distribution of these spike protein variants across Indian states (S3 Table). As we computed the diversity based on both richness and evenness of spike variants, some of the states like Maharashtra, Odisha, West Bengal and Gujarat demonstrated significantly higher ($P<0.05$) diversity of circulating variants than most of the remaining states. However, while we had 196, 97, 75 and 73 sequenced isolates from Gujarat, Telengana, Delhi and Maharashtra respectively, the remaining states were represented by even lower sample size (ranging from 1 to 48 isolates. Of these, Delhi variants showed lowest diversity, significantly different from both Maharashtra ($P=0.0002$) and Gujarat ($P=0.0006$), though not from Telengana ($P=0.16$). On the other hand, Maharashtra variants presented much higher diversity than Telengana ($P=0.017$) or Gujarat ($P=0.21$).

It is expected that these variations might be in response to strong selection pressures

acting on the spike protein, especially its S1 subunit which, being a major immunogenic

target for the host, plays the pivotal role to evade the host immune response and to offer a

successful viral entry. Therefore, the mutational variations in spike proteins can essentially

affect the stability of the S-R complex. To assess this, we modeled each of the spike protein

variants, and then docked to the binding site of host receptor, hACE2, using HADDOCK

webserver [33, 34] of data-driven docking algorithm by providing binding site information as

the same was already established from the crystal structure [32, 35]. In the circulating

variants, we detected a significant excess (P=0.035) of mutations in the S1 subunit (with 41

mutations) compared to the S2 subunit (with 22 mutations). The mutation positions in the

secondary structures of the analyzed variants are detailed in the S1 Dataset in supporting

information.

The docking score (HADDOCK score) of each variant on hACE2 is hereafter

designated as the stability index of S-R complex. More negative is the docking (HADDOCK)

score, higher is the stability of the S-R complex [37]. Under the assumption that better

stability would lead to better invasion of the virus into the host, we hypothesize that such a

stability index of a given spike protein variant could be linked to the severity of viral

pathogenicity. To test this hypothesis, we measured the severity as the 'fatality rate'

calculated simply as the ratio of the number of deceased people to the number of recovered in

a given state (available from the Government of India website: www.mygov.in/corona-

data/covid19-statewise-status/).

While we aimed to estimate an average stability index of a given state based on the

stability indices of circulating variants in that location, we were handicapped with the

available sample size and the information of collection diversity. Considering this issue, we

restricted our study to the states having 50 or more sequenced samples for analysis.

Therefore, we could assess the association of average stability index with fatality rates for

three states and one union territory (Maharashtra, Gujarat, Telengana and Delhi) which

qualified our sample size threshold. Importantly, these four regions harbored 70% of all

samples analyzed across 19 states, while their variant diversity ranged from the highest to one

of the lowest (S3 Table).

We detected a strong exponential correlation (R^2=0.96) between the average stability

index of circulating spike variants of the region with the fatality rate in that region (Fig 2).

While Telengana and Delhi showed comparable average stability index values with ~7%

fatality rates, Maharashtra and Gujarat had exponentially higher stability index values (i.e.,

more negative docking or HADDOCK scores) with 8% and 9% fatality rates respectively. It

is highly plausible that the spike protein, as the primary controller of both the attachment to

the host cell surface and the initiation of infection by fusing the viral and the host cell

membranes, would be represented by variants with varying efficiency of the virus to enter

human cells, and to get transmitted among people [17]. However, our conclusions based on

the available initial data are premature due to low sample size per location and lack of direct

evidence for the correlation between spike protein variant's docking score and the pathogen's

contribution to host fatality, thereby warranting population-level robust association analysis

and experimental validations.

Figure 2. Heat map distribution across four Indian states with >50 sequenced isolates

based on average stability index. The average stability index for a particular state denotes

the averaged value of docking scores / HADDOCK scores of S-R complexes for all

circulating variants. The values of average stability index and fatality rate in Indian states are

plotted to fit an exponential function (R^2=0.96).

The emerging spike protein variants showing reduced stability of S-R complex are significantly abundant.

Of 630 isolates analyzed, Ancestors 1 and 2 were represented by 253 and 248 isolates, suggesting their steady circulation across India. However, the remaining ones, i.e. more than 20% isolates represented relatively recently emerged variants out of two ancestral variants. Interestingly, a quick look at Fig 1 showed that, for majority of the variants derived from the Ancestor 1 and Ancestor 2, the stability was reduced (having less negative stability index values) relative to their respective ancestors. We therefore plotted the trend of those emerging variants with reference to their ancestral backgrounds (Fig 3). Significant majority (χ^2 P=0.03) of the variants that emerged from the two ancestral variants showed reduced stability (having less negative docking scores) from their respective ancestors. This picture got even more prominent (χ^2 P=0.009) when we looked into exclusively the variants which were detected multiple times in the dataset, i.e. represented by more than one isolate (sometimes collected from different states), thereby suggesting possible fixation of those variants in the population (S2 Table). Interestingly, 13 of these variants with multiple occurrences accumulated mutations exclusively in S1 subunit, while only 3 variants showed all mutations in S2 subunit, which might be suggesting an increased selection pressure in the S1 subunit because of its key role in the viral entry and the presence of the receptor binding domain.

Figure 3. Stability index (i.e., docking score or HADDOCK score) plot of S-R complexes for spike protein variants emerging from (a) Ancestor 1 (Wuhan-Hu-1/2019 variant) and (b) Ancestor 2 (D614G variant). The blue dotted line is used as a reference to denote the stability index value for respective ancestral variants. More negative is the value, higher is the stability level. The red dotted rectangular block includes the variants that are represented by multiple isolates in our dataset (S2 Table).

303   At this point, we can propose that the relatively recent variants emerging from the

304 ancestors in India are losing their ability on an average to form a stable complex with the

305 human receptor as compared with their ancestors, which could possibly result in lower

306 countrywide fatality rate, if we combine our earlier observation of the direct correlation

307 between average stability index and fatality rate. Conforming to this proposal, we found that,

308 after an initial steady increase of fatality rate for the first four weeks, the fatality rate reached

309 a peak at 38.2% on April 11, 2020, followed by a continuous sharp decline at 5.3% on June

310 28, 2020 until when the data were available (Fig 4), fitting a power law function (R^2=0.94)

311 (data source: http://covidindiaupdates.in/). This was despite the fact that the rate of mutations

312 was found to be increasing during the three-month period from March till May 2020 during

313 which our analyzed samples were collected. Interestingly, the increase in nonsynonymous

314 (i.e. amino acid replacement) mutation rate (from 0.027% in March to 0.033% in May 2020)

315 was detected to be about 14% higher than the increase in synonymous mutation rate (from

316 0.091% in March to 0.097% in May 2020), suggesting a stronger selection pressure for amino

317 acid changes in the spike protein.

319 **Figure 4. Countrywide fatality rates (the number of deaths / the number of recovered**

320 **cases) over time in India.** The date of first available data (March 13, 2020), the date after

321 which the decline of fatality rate started (April 11, 2020), and the date until which the

322 analyzed samples were collected (May 27, 2020) are denoted by blue dotted lines. The

323 declining fatality rate curve is best fitted by a power law function (R^2=0.94).

Could Muller's ratchet be a player in shaping SARS-CoV-2 evolutionary dynamics in India?

In evolutionary genetics, Muller's ratchet signifies accumulation of deleterious mutation in a population leading to 'mutational meltdown', leading to a gradual extinction of that population [38]. In general, genetic mutations that provide adaptive advantages are fixed in the population by natural selection whereas deleterious ones are wiped off from the population. However, an accelerated mutation rate puts huge mutational pressure, and natural selection is unable to wash out these deleterious mutations, retaining the newly formed variants within the population and thereby leading to their fixation. This irretrievable evolutionary mechanism is coined as Muller's ratchet by evolutionary biologists [39]. When more and more deleterious mutations are accumulated and become permanent in the population, this results in 'mutational meltdown' or ultimate loss of the population [40, 41].

It is known that the apparent tendency to directly correlate the high mutation rate of virus with its infectivity and transmissibility is without merit [42]. On the other side, it has been suggested that Muller's ratchet, via mutational meltdown, could be a key player in leading the SARS-CoV-2 population to gradual extinction due to accumulation and fixation of deleterious mutations in future [40, 43]. On one hand, we observed that the stability of S-R complex is directly linked to the fatality rates, while the continuous emergence of variants from the ancestral ones were found to be less stable compared to their ancestors. As we combine these findings with the countrywide data of a sharp decline in the fatality rate over time, we propose the possibility of mutational meltdown in action for SARS-CoV-2 in India, indicating Muller's ratchet as a plausible game-changer for COVID-19 scenario here in near future.

We understand that, only in three months, we cannot expect a radical increase in the mutation rate to give rise to any significant accumulation of deleterious mutations that could

have offered a prominent picture of mutational meltdown. However, our results altogether point toward the trend, thereby suggesting the potential of future studies in this otherwise overlooked domain of microbial dynamics, which could in turn lead to a possibility of a successful therapeutic approach [40, 43].

Discussion

Our work has combined genetic and epidemiological data of SARS-CoV2 in India to decipher a direct correlation between the average stability of S-R complexes for the circulating spike protein variants and the fatality rate of a geographic region. The docking score of S-R complex, designated here as the stability index, is estimated by protein-protein docking based on intermolecular interactions, such as electrostatic and van der Waals interactions, desolvation and restraint violation energies along with buried surface area upon binding via detection of the correct binding pose. This score, in essence, quantifies the stability of the docked complex by optimizing global minimum energy conformation of the complex [44, 45]. On the other side, while it is probable that a better stability of S-R complex could lead to an increased viral load and maybe an increased infectivity, previous works showed that the spike protein – hACE2 complex is crucial for viral pathogenesis by causing acute lung damage [46, 47], which suggests a direct link between S-R complex stability and fatality rate. However, the robustness of the potential of S-R complex stability index for the spike protein variants as a tracker of fatality rate or disease severity needs to be studied in greater depths with more structured region-specific patient data (that are primarily available to selected government/non-government agencies) in connection with larger population-level sequence datasets for the given locations.

As expected for any fast-moving pathogen outbreaks [42], we find an increasing rate of mutations, while the extent of increase is much higher for the nonsynonymous (amino acid replacement) changes. Interestingly, the S-R complex of a significant majority of variants tend to lose stability relative to their two most stable ancestral variants. Alongside, the countrywide data show a continuous sharp decline in the fatality rate after an initial surge. Therefore, keeping in mind the observed correlation between the S-R complex stability and fatality rate across Indian states, here we pose an open question for future research: Does the phylodynamics of SARS-CoV-2 in India indicate any nascent action of Muller's ratchet where the otherwise deleterious mutations tend to get fixed in the population as natural selection remains unable to purge them due to excessive mutational pressures? If so, there lies an immense potential of using therapeutics that could facilitate such a process of mutational meltdown, as was demonstrated earlier for influenza A virus [48, 49]. Therefore, future large-scale population genomics analysis supported by epidemiological information is of high importance to explore this question for India as well as for other countries across the globe to develop efficient analytical methods, thereby guiding better surveillance programs, prevention and treatment management of COVID-19.

References

1. Coronaviridae Study Group of the International Committee on Taxonomy of V. The species Severe acute respiratory syndrome-related coronavirus: classifying 2019-nCoV and naming it SARS-CoV-2. Nat Microbiol. 2020;5(4):536-44. Epub 2020/03/04. doi: 10.1038/s41564-020-0695-z.

2. Holmes EC, Nee S, Rambaut A, Garnett GP, Harvey PH. Revealing the history of infectious disease epidemics through phylogenetic trees. Philos Trans R Soc Lond B Biol Sci. 1995;349(1327):33-40. Epub 1995/07/29. doi: 10.1098/rstb.1995.0088.

3.   Pybus OG, Rambaut A. Evolutionary analysis of the dynamics of viral infectious

disease. Nat Rev Genet. 2009;10(8):540-50. Epub 2009/07/01. doi: 10.1038/nrg2583.

4.   Andersen KG, Rambaut A, Lipkin WI, Holmes EC, Garry RF. The proximal origin of

SARS-CoV-2. Nat Med. 2020;26(4):450-2. Epub 2020/04/15. doi: 10.1038/s41591-

020-0820-9.

5.   Tortorici MA, Veesler D. Structural insights into coronavirus entry. Adv Virus Res.

2019;105:93-116. Epub 2019/09/17. doi: 10.1016/bs.aivir.2019.08.002.

6.   Gui M, Song W, Zhou H, Xu J, Chen S, Xiang Y, et al. Cryo-electron microscopy

structures of the SARS-CoV spike glycoprotein reveal a prerequisite conformational

state for receptor binding. Cell Res. 2017;27(1):119-29. Epub 2016/12/23. doi:

10.1038/cr.2016.152.

7.   Kirchdoerfer RN, Cottrell CA, Wang N, Pallesen J, Yassine HM, Turner HL, et al.

Pre-fusion structure of a human coronavirus spike protein. Nature.

2016;531(7592):118-21. Epub 2016/03/05. doi: 10.1038/nature17200.

8.   Pallesen J, Wang N, Corbett KS, Wrapp D, Kirchdoerfer RN, Turner HL, et al.

Immunogenicity and structures of a rationally designed prefusion MERS-CoV spike

antigen. Proc Natl Acad Sci U S A. 2017;114(35):E7348-E57. Epub 2017/08/16. doi:

10.1073/pnas.1707304114.

9.   Song W, Gui M, Wang X, Xiang Y. Cryo-EM structure of the SARS coronavirus

spike glycoprotein in complex with its host cell receptor ACE2. PLoS Pathog.

2018;14(8):e1007236. Epub 2018/08/14. doi: 10.1371/journal.ppat.1007236.

10.   Walls AC, Tortorici MA, Bosch BJ, Frenz B, Rottier PJM, DiMaio F, et al. Cryo-

electron microscopy structure of a coronavirus spike glycoprotein trimer. Nature.

2016;531(7592):114-7. Epub 2016/02/09. doi: 10.1038/nature16988.

11.   Walls AC, Tortorici MA, Snijder J, Xiong X, Bosch BJ, Rey FA, et al. Tectonic

conformational changes of a coronavirus spike glycoprotein promote membrane

fusion. Proc Natl Acad Sci U S A. 2017;114(42):11157-62. Epub 2017/10/27. doi:

10.1073/pnas.1708727114.

12.   Belouzard S, Chu VC, Whittaker GR. Activation of the SARS coronavirus spike

protein via sequential proteolytic cleavage at two distinct sites. Proc Natl Acad Sci U

S A. 2009;106(14):5871-6. Epub 2009/03/27. doi: 10.1073/pnas.0809524106.

13.   Heald-Sargent T, Gallagher T. Ready, set, fuse! The coronavirus spike protein and

acquisition of fusion competence. Viruses. 2012;4(4):557-80. Epub 2012/05/17. doi:

10.3390/v4040557.

14.   Millet JK, Whittaker GR. Host cell entry of Middle East respiratory syndrome

coronavirus after two-step, furin-mediated activation of the spike protein. Proc Natl

Acad Sci U S A. 2014;111(42):15214-9. Epub 2014/10/08. doi:

10.1073/pnas.1407087111.

15.   Millet JK, Whittaker GR. Host cell proteases: Critical determinants of coronavirus

tropism and pathogenesis. Virus Res. 2015;202:120-34. Epub 2014/12/03. doi:

10.1016/j.virusres.2014.11.021.

16.   Park JE, Li K, Barlan A, Fehr AR, Perlman S, McCray PB, Jr., et al. Proteolytic

processing of Middle East respiratory syndrome coronavirus spikes expands virus

tropism. Proc Natl Acad Sci U S A. 2016;113(43):12262-7. Epub 2016/10/30. doi:

10.1073/pnas.1608147113.

17.   Walls AC, Park YJ, Tortorici MA, Wall A, McGuire AT, Veesler D. Structure,

Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. Cell.

2020;181(2):281-92 e6. Epub 2020/03/11. doi: 10.1016/j.cell.2020.02.058.

18. Brufsky A, Lotze MT. DC/L-SIGNs of hope in the COVID-19 pandemic. J Med
Virol. 2020. Epub 2020/05/07. doi: 10.1002/jmv.25980.

19. Gordon DE, Jang GM, Bouhaddou M, Xu J, Obernier K, White KM, et al. A SARS-
CoV-2 protein interaction map reveals targets for drug repurposing. Nature.
2020;583(7816):459-68. Epub 2020/05/01. doi: 10.1038/s41586-020-2286-9.

20. Jia Y, Shen G, Zhang Y, Huang K-S, Ho H-Y, Hor W-S, et al. Analysis of the
mutation dynamics of SARS-CoV-2 reveals the spread history and emergence of
RBD mutant with lower ACE2 binding affinity. bioRxiv. 2020:2020.04.09.034942.
doi: 10.1101/2020.04.09.034942.

21. Korber B, Fischer W, Gnanakaran S, Yoon H, Theiler J, Abfalterer W, et al. Spike
mutation pipeline reveals the emergence of a more transmissible form of SARS-CoV-
2. bioRxiv. 2020:2020.04.29.069054. doi: 10.1101/2020.04.29.069054.

22. Yao H, Lu X, Chen Q, Xu K, Chen Y, Cheng L, et al. Patient-derived mutations
impact pathogenicity of SARS-CoV-2. medRxiv. 2020:2020.04.14.20060160. doi:
10.1101/2020.04.14.20060160.

23. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: Molecular Evolutionary
Genetics Analysis across Computing Platforms. Mol Biol Evol. 2018;35(6):1547-9.
Epub 2018/05/04. doi: 10.1093/molbev/msy096.

24. Chattopadhyay S, Paul S, Dykhuizen DE, Sokurenko EV. Tracking recent adaptive
evolution in microbial species using TimeZone. Nat Protoc. 2013;8(4):652-65. Epub
2013/03/09. doi: 10.1038/nprot.2013.031.

25. Huang Y, Niu B, Gao Y, Fu L, Li W. CD-HIT Suite: a web server for clustering and
comparing biological sequences. Bioinformatics. 2010;26(5):680-2. Epub 2010/01/08.
doi: 10.1093/bioinformatics/btq003.

26.   Li W, Godzik A. Cd-hit: a fast program for clustering and comparing large sets of

473   protein or nucleotide sequences. Bioinformatics. 2006;22(13):1658-9. Epub

474   2006/05/30. doi: 10.1093/bioinformatics/btl158.

475   27.   Higgins DG, Sharp PM. CLUSTAL: a package for performing multiple sequence

476   alignment on a microcomputer. Gene. 1988;73(1):237-44. Epub 1988/12/15. doi:

477   10.1016/0378-1119(88)90330-7.

478   28.   Rice P, Longden I, Bleasby A. EMBOSS: the European Molecular Biology Open

479   Software Suite. Trends Genet. 2000;16(6):276-7. Epub 2000/05/29. doi:

480   10.1016/s0168-9525(00)02024-2.

481   29.   Simpson EH. Measurement in Diversity. Nature. 1949;163:688. doi:

482   https://doi.org/10.1038/163688a0.

483   30.   Waterhouse A, Bertoni M, Bienert S, Studer G, Tauriello G, Gumienny R, et al.

484   SWISS-MODEL: homology modelling of protein structures and complexes. Nucleic

485   Acids Res. 2018;46(W1):W296-W303. Epub 2018/05/23. doi: 10.1093/nar/gky427.

486   31.   Wrapp D, Wang N, Corbett KS, Goldsmith JA, Hsieh CL, Abiona O, et al. Cryo-EM

487   structure of the 2019-nCoV spike in the prefusion conformation. Science.

488   2020;367(6483):1260-3. Epub 2020/02/23. doi: 10.1126/science.abb2507.

489   32.   Wang Q, Zhang Y, Wu L, Niu S, Song C, Zhang Z, et al. Structural and Functional

490   Basis of SARS-CoV-2 Entry by Using Human ACE2. Cell. 2020;181(4):894-904 e9.

491   Epub 2020/04/11. doi: 10.1016/j.cell.2020.03.045.

492   33.   van Dijk M, Wassenaar TA, Bonvin AM. A Flexible, Grid-Enabled Web Portal for

493   GROMACS Molecular Dynamics Simulations. J Chem Theory Comput.

494   2012;8(10):3463-72. Epub 2012/10/09. doi: 10.1021/ct300102d.

495   34.   van Zundert GCP, Rodrigues J, Trellet M, Schmitz C, Kastritis PL, Karaca E, et al.

496   The HADDOCK2.2 Web Server: User-Friendly Integrative Modeling of

Biomolecular Complexes. J Mol Biol. 2016;428(4):720-5. Epub 2015/09/28. doi:

10.1016/j.jmb.2015.09.014.

35. Lan J, Ge J, Yu J, Shan S, Zhou H, Fan S, et al. Structure of the SARS-CoV-2 spike

receptor-binding domain bound to the ACE2 receptor. Nature. 2020;581(7807):215-

20. Epub 2020/04/01. doi: 10.1038/s41586-020-2180-5.

36. Humphrey W, Dalke A, Schulten K. VMD: visual molecular dynamics. J Mol Graph.

1996;14(1):33-8, 27-8. Epub 1996/02/01. doi: 10.1016/0263-7855(96)00018-5.

37. Pantsar T, Poso A. Binding Affinity via Docking: Fact and Fiction. Molecules.

2018;23(8). Epub 2018/08/01. doi: 10.3390/molecules23081899.

38. Muller HJ. The Relation of Recombination to Mutational Advance. Mutat Res.

1964;106:2-9. Epub 1964/05/01. doi: 10.1016/0027-5107(64)90047-8.

39. Felsenstein J. The evolutionary advantage of recombination. Genetics.

1974;78(2):737-56. Epub 1974/10/01.

40. Jensen JD, Lynch M. Considering mutational meltdown as a potential SARS-CoV-2

treatment strategy. Heredity (Edinb). 2020;124(5):619-20. Epub 2020/04/07. doi:

10.1038/s41437-020-0314-z.

41. Lynch M, Burger R, Butcher D, Gabriel W. The mutational meltdown in asexual

populations. J Hered. 1993;84(5):339-44. Epub 1993/09/01. doi:

10.1093/oxfordjournals.jhered.a111354.

42. Grubaugh ND, Petrone ME, Holmes EC. We shouldn't worry when a virus mutates

during disease outbreaks. Nat Microiol. 2020;5(4):529-30. Epub 2020/02/20. doi:

10.1038/s41564-020-0690-4.

43. Brufsky A, Lotze MT. Ratcheting down the virulence of SARS-CoV-2 in the COVID-

19 pandemic. J Med Virol. 2020. Epub 2020/05/28. doi: 10.1002/jmv.26067.

44.   Halperin I, Ma B, Wolfson H, Nussinov R. Principles of docking: An overview of

522   search algorithms and a guide to scoring functions. Proteins. 2002;47(4):409-43. Epub

523   2002/05/10. doi: 10.1002/prot.10115.

524   45.   Moreira IS, Fernandes PA, Ramos MJ. Protein-protein docking dealing with the

525   unknown. J Comput Chem. 2010;31(2):317-42. Epub 2009/05/23. doi:

526   10.1002/jcc.21276.

527   46.   Kuba K, Imai Y, Rao S, Gao H, Guo F, Guan B, et al. A crucial role of angiotensin

528   converting enzyme 2 (ACE2) in SARS coronavirus-induced lung injury. Nat Med.

529   2005;11(8):875-9. Epub 2005/07/12. doi: 10.1038/nm1267.

530   47.   Lumbers ER, Delforce SJ, Pringle KG, Smith GR. The Lung, the Heart, the Novel

531   Coronavirus, and the Renin-Angiotensin System; The Need for Clinical Trials. Front

532   Med (Lausanne). 2020;7:248. Epub 2020/06/24. doi: 10.3389/fmed.2020.00248.

533   48.   Ormond L, Liu P, Matuszewski S, Renzette N, Bank C, Zeldovich K, et al. The

534   Combined Effect of Oseltamivir and Favipiravir on Influenza A Virus Evolution.

535   Genome Biol Evol. 2017;9(7):1913-24. Epub 2017/09/01. doi: 10.1093/gbe/evx138.

536   49.   Penisson S, Singh T, Sniegowski P, Gerrish P. Dynamics and Fate of Beneficial

537   Mutations Under Lineage Contamination by Linked Deleterious Mutations. Genetics.

538   2017;205(3):1305-18. Epub 2017/01/20. doi: 10.1534/genetics.116.194597.

Supporting information Captions

S1 Table: List of the sequenced isolates analyzed from GISAID database (www.gisaid.org) available

till June 7, 2020. These isolates included complete gene sequence for the spike protein. The mutations

of spike protein variant for each isolate was mentioned along with the docking score of S-R complex

for the variant. The docking scores of the variants with mutations between amino acid positions 1-26

and 1148-1273 were not determined because of the unavailability of the modeling template (1, 2) and

therefore mentioned as NA (not available).

S2 Table: Countrywide and state-wise frequency distribution matrix of circulating variants along with

the docking score of S-R complex for each variant. The docking scores of the variants with mutations

between amino acid positions 1-26 and 1148-1273 were not determined because of the unavailability

of the modeling template (1, 2) and therefore mentioned as NA (not available). The variant mutation

positions marked by asterisks denote convergent nature of these mutations, suggesting their

independent accumulation multiple times along phylogenetically unlinked allelic backgrounds. The

variants that were detected to be uniquely present in Indian population as compared to a total of 17529

worldwide genomic isolates available from the GISAID database till May 9, 2020.

S3 Table: State-wise distribution of the type and frequency of spike protein variants, along with the

calculated diversity based on Simpson's index (3).

Dataset S1: Positional information of the spike protein mutations in the docked S-R complex.

Fig. 1. Schematic representation of the diversity of spike protein variants circulating in India, using maximum likelihood-based phylogeny reconstruction. Each node represents a specific spike protein variant, while the node-size and the number inside depict the frequency of that variant. The red or green color of each arrow indicates the higher or lower stability index respectively of the S-R complex for each variant than the major ancestral variant it emerged from (either Ancestor 1 or Ancestor 2). The black arrows lead to the variants for which the docking scores could not be determined either because of the presence of at least one variation outside the available template region for docking(18, 42), or due to non-existing isolate in the lone hypothetical node with H1083Q mutation denoted by black color. This black node signifies a variant with no available isolate in our dataset, while it gives rise to two derived variants, H1083Q:R78M and H1083Q:E583D, for which representative isolates were available.

592 Fig. 2. Heat map distribution across four Indian states with >50 sequenced isolates based on average
593 stability index. The average stability index for a particular state denotes the averaged value of docking
594 scores / HADDOCK scores of S-R complexes for all circulating variants. The values of average stability
595 index and fatality rate in Indian states are plotted to fit an exponential function (R^2=0.96).
596

598
599
600

Fig. 3. Stability index (i.e., docking score or HADDOCK score) plot of S-R complexes for spike protein variants emerging from (a) Ancestor 1 (Wuhan-Hu-1/2019 variant) and (b) Ancestor 2 (D614G variant). The blue dotted line is used as a reference to denote the stability index value for respective ancestral variants. More negative is the value, higher is the stability level. The red dotted rectangular block includes the variants that are represented by multiple isolates in our dataset (*SI appendix*, Table S2).

614 **Fig. 4. Countrywide fatality rates (the number of deaths / the number of recovered cases) over time in**
615 **India.** The date of first available data (March 13, 2020), the date after which the decline of fatality rate
616 started (April 11, 2020), and the date until which the analyzed samples were collected (May 27, 2020) are
617 denoted by blue dotted lines. The declining fatality rate curve is best fitted by a power law function
618 (R^2=0.94).
619