Fitting side-chain NMR relaxation data using molecular simulations

Felix Kümmerer[1†], **Simone Orioli**[1,2†], **David Harding-Larsen**[1], **Falk Hoffmann**[3‡], **Yulian Gavrilov**[1], **Kaare Teilum**[1], **Kresten Lindorff-Larsen**[1*]

*For correspondence:
lindorff@bio.ku.dk (KLL)

[†]These authors contributed equally to this work

Present address: [‡]Institute of Biomaterial Science and Berlin-Brandenburg Center of Regenerative Therapies, Helmholtz-Zentrum Geesthacht, Kantstrasse 55, D-14513 Teltow, Germany

[1]Structural Biology and NMR Laboratory, Linderstrøm-Lang Centre for Protein Science, Department of Biology, University of Copenhagen. Ole Maaløes Vej 5, DK-2200 Copenhagen N, Denmark; [2]Structural Biophysics, Niels Bohr Institute, Faculty of Science, University of Copenhagen, Copenhagen, Denmark.; [3]Theoretical Chemistry, Ruhr University Bochum, D-44780 Bochum, Germany

Abstract   Proteins display a wealth of dynamical motions that can be probed using both experiments and simulations. We present an approach to integrate side chain NMR relaxation measurements with molecular dynamics simulations to study the structure and dynamics of these motions. The approach, which we term ABSURDer (Average Block Selection Using Relaxation Data with Entropy Restraints) can be used to find a set of trajectories that are in agreement with relaxation measurements. We apply the method to deuterium relaxation measurements in T4 lysozyme, and show how it can be used to integrate the accuracy of the NMR measurements with the molecular models of protein dynamics afforded by the simulations. We show how fitting of dynamic quantities leads to improved agreement with static properties, and highlight areas needed for further improvements of the approach.

Introduction

Proteins are dynamical entities and a detailed understanding of their function and biophysical properties requires an accurate description of both their structure and dynamics. Nuclear magnetic resonance (NMR) experiments, in particular, have the ability to probe and quantify dynamical properties on a wide range of time scales and at atomic resolution. Computationally, molecular dynamics (MD) simulations also make it possible to probe both the structure and dynamics of proteins. Indeed, NMR and MD simulations may fruitfully be combined in a number of ways (*Case, 2002*; *Lindorff-Larsen et al., 2005*), including using experiments for validating or improving the force fields used in simulations (*Norgaard et al., 2008*; *Li and Brüschweiler, 2010*, *2011*), or for using the simulations as a tool to interpret the experiments (*Nodet et al., 2009*; *Brookes and Head-Gordon, 2016*; *Chen et al., 2019*; *Vasile and Tiana, 2019*; *Bottaro et al., 2020*).

A particularly tight integration between NMR and MD simulations has involved NMR spin relaxation experiments, which probe motions on the ps-to-ns timescales. Indeed, already soon after the first reported MD simulation of a protein (*McCammon et al., 1977*), such simulations were compared to NMR order parameters (*Lipari et al., 1982*). Integration between spin relaxation and MD is aided by the fact that MD can probe the relevant time scales and that the physical processes leading to spin relaxation are relatively well understood, and can therefore be modelled computationally.

The capability to reach given timescales does not, however, automatically translate into a perfect match between experiments and the simulated observables (*van Gunsteren et al., 2018*; *Bot-*

taro and Lindorff-Larsen, 2018). Indeed, significant discrepancies between the two can emerge because of multiple factors, such as insufficient sampling (*Bernardi et al., 2015*), inaccurate description of the observable, i.e. the so-called forward model (*Cordeiro et al., 2017*), the imperfection of the underlying force fields (*Rauscher et al., 2015*; *Henriques et al., 2015*; *Robustelli et al., 2018*; *Piana et al., 2020*; *Nerenberg and Head-Gordon, 2018*) or a combination of these factors.

NMR spin relaxation experiments are generally interpreted using variants of the so-called model free formalism (*Halle and Wennerström, 1981*; *Lipari and Szabo, 1982*; *Clore et al., 1990*) that interprets the experimental data using generalized order parameters that describe the amplitudes of the motions and time scales associated with those motions. A common approach to compare MD and NMR relaxation experiments involves calculating order parameters from the simulations and comparing to those extracted from the experiments. The results of such comparisons have shown that while MD simulations generally give a relatively good agreement with order parameters for the backbone amide groups, the agreement for methyl bearing side chains is more variable (*Bremi et al., 1997*; *Skrynnikov et al., 2002*; *Best et al., 2005*; *Showalter et al., 2007*; *O'Brien et al., 2016*; *Bowman, 2016*).

A more detailed and richer integration, however, involves calculating the NMR relaxation rates directly from the simulations and comparing these to experiments. Recently, an approach was developed to calculate deuterium NMR relaxation rates in methyl bearing side chains from MD simulations (*Hoffmann et al., 2018b*, *2020*). Comparison to experiments revealed systematic deviations from simulations with several different force fields, and corrections to the methyl torsion potential were developed and shown to increase agreement with experiments (*Hoffmann et al., 2018a*, *2020*). Despite these improvements, however, the agreement remains imperfect and further improvements are desirable.

One possible way to reduce the systematic error coming from force field inaccuracies is to introduce a bias in the ensemble generated by MD simulations to improve agreement with the experimental data. Such a bias may either be introduced during the simulation by adding a correction to the force field that depends on the experimental measurement, or after the simulation has been completed through a procedure known as *reweighting*. Using biased simulations it is, for example, possible to construct conformational ensembles that are in agreement with backbone and side chain order parameters from NMR spin relaxation (*Best and Vendruscolo, 2004*; *Lindorff-Larsen et al., 2005*).

Most implementations of such methods for biasing against experiments work by updating some 'prior' information, typically encoded in the MD force field, with information from the experiments by changing the weights of each conformation so as to improve the agreement with experiments when calculated with these new weights. Often these problems are highly underdetermined in that there are many more parameters (weights) to be determined than experimental measurements. Thus, an important ingredient in many methods is a framework to avoid overfitting by balancing the information from the force field with that from the data, and many approaches apply Bayesian or Maximum Entropy formalisms for this purpose (reviewed in *Orioli et al.* (*2020*)).

Although different methods do so in different ways and with different assumptions, most reweighting methods share a key commonality: they can only deal with static (equilibrium) observables, i.e. quantities that can be expressed as ensemble averages over a set of configurations (*Bonomi et al., 2017*; *Orioli et al., 2020*). Evidently, this complicates the generation of conformational ensembles that match the spin relaxation data, which depend both on the amplitudes and time scales of the motions.

A few approaches have been described to fit NMR relaxation data directly. For example, the isotropic reorientational eigenmode dynamics (iRED) approach (*Prompers and Brüschweiler, 2002*) uses a covariance matrix to calculate spin relaxation, and may be applied to blocked simulations thus indirectly taking into account the time scales of the motions (*Gu et al., 2014*). Recently, a more direct approach has been described to reweight MD simulations against NMR spin relaxation data (*Salvi et al., 2016*, *2019*). The basic idea is to split one or more simulations into smaller

parts (blocks), each of which are long enough to contain information about both the amplitudes and time scales of the motions that lead to spin relaxation. By averaging over such blocks, one can then estimate NMR relaxation parameters that may be compared to experiments. Similar to the reweighting methods described above for individual configurations, one can instead use reweighting of the blocks and thus determine weights that improve agreement with the experimental data without having to decompose it into order parameters. This approach, called ABSURD (Average Block Selection Using Relaxation Data), has been applied to the analysis of backbone NMR relaxation of intrinsically disordered proteins that are difficult to analyse using conventional model free techniques.

We here describe a new approach to interpret NMR relaxation data, focusing our application on side chain dynamics. Our approach builds upon the ABSURD method, and includes two extensions. First, we use our recently described methods to calculate side chain NMR relaxation parameters from molecular dynamics simulations (*Hoffmann et al., 2018b*). Second, we extend ABSURD by including an entropy restraint term in the optimization that helps avoid overfitting. We thus term our method ABSURDer (Average Block Selection Using Relaxation Data with Entropy Restraints). We applied ABSURDer to study the fast timescale dynamics of T4 lysozyme (T4L), a protein which has been the subject of numerous experimental and computational studies. We validate and describe our method using synthetic data and then apply it to data from NMR experiments. We find that reweighting simulations with ABSURDer improves the agreement between methyl relaxation rates in synthetic data sets coming from both the same and a different force field. Furthermore, we find that these reweighted trajectories also show improvements in the agreement of equilibrium observables, such as rotamer distributions, coming from the two data sets. Using experimental NMR data for reweighting, we show similar improvements although to a lesser extent, and we discuss possible origins of this observation.

Results and Discussion

Overview of the ABSURDer approach

The workflow in ABSURDer consists of three separate steps (Fig. 1). Steps 1 and 2 correspond to a previously described method for calculating side chain relaxation, and step 3 corresponds to the ABSURD approach with the inclusion of an additional restraint to decrease the risk of overfitting, and to balance the information in the force field with that in the data.

First, we performed three independent $5\,\mu s$-long MD simulations to sample the structure and dynamics of T4L. We here use the recently optimized Amber ff99SB*-ILDN force field with the TIP4P/2005 water model and a modification of the methyl torsion potential, based on CCSD(T) coupled cluster quantum chemical calculations of isolated dipeptides, that improves agreement with NMR relaxation data (*Hoffmann et al., 2018a*). See the Methods section for further details about all simulations.

Second, we divided the total of $15\,\mu s$ simulations into 1500 $10\,ns$ blocks, and calculated NMR relaxation parameters for each of these independently (see Methods, and below for a more detailed discussion on the origin of this block size). Briefly described we: (i) calculate internal correlation functions $C_{int}(t)$ for methyl C-H bonds; (ii) use backbone motions to describe the global overall tumbling motion; (iii) combine internal and global rotational motions to calculate the spectral density function $J(\omega)$ and (iv) extract NMR relaxation parameters from this. We here focus on three NMR relaxation parameters $R(D_z)$, $R(D_y)$ and $R(3D_z^2 - 2)$, and calculate these rates for each methyl group and each of the 1500 blocks. As done previously (*Hoffmann et al., 2018b*) we exclude residue ALA146 from the analyses.

The third step in our approach consists of reweighting the MD simulations to improve agreement with the experimental data. Our approach combines the ABSURD method with concepts from Bayesian/Maximum Entropy ensemble refinement. The goal here is to assign to each of the 1500 $10\,ns$ simulations a different weight (w_i; $i = 1, \dots, 1500$) so as to improve agreement with the

Figure 1. Schematic representation of the ABSURDer workflow. Steps 1 involves sampling protein dynamics using one or more longer MD simulations and calculating bond vector orientations for the resulting trajectories. These are then divided into blocks and in Step 2 we calculate correlation functions, spectral densities and NMR relaxation rates for each block. In Step 3, we optimize the agreement between the average calculated rates and experimentally measured values by changing the weights of the different blocks. For details see main text, methods and code online.

experimental data, as quantified by the χ^2 between experimental and calculated NMR relaxation rates. Such an approach is, however, prone to overfitting and does not necessarily utilize the information about the conformational landscape encoded in the molecular force field. Thus, we and others have previously described methods that circumvent this problem by adding an entropy restraint to the χ^2 minimization (*Gull and Daniell, 1978*; *Cesari et al., 2018*; *Köfinger et al., 2019*; *Bottaro et al., 2020*; *Orioli et al., 2020*), by instead minimizing the functional $\mathcal{L}(\mathbf{w}) = \chi^2(\mathbf{w}) - \theta S_{rel}(\mathbf{w})$ (see Methods). In this equation S_{rel} represents the relative entropy that measures how different the weights are from the initial (uniform) weights, and thus how much the final set of simulations differs from the initial set. Typically, we represent this as $\phi_{eff} = \exp(S_{rel})$, which represents the effective fraction of the original 1500 blocks (sub-simulations) that contribute to the final ensemble. In these equations the parameter θ sets the balance between fitting the data (minimizing χ^2) and keeping as much as possible of the original simulation (maximizing ϕ_{eff}). We refer the reader to previous literature about the methods overall and how best to select this parameter (*Andrae et al., 2010*; *Hummer and Köfinger, 2015*; *Bottaro et al., 2018*; *Cesari et al., 2018*; *Köfinger et al., 2019*; *Crehuet et al., 2019*; *Chen et al., 2019*; *Bottaro et al., 2020*; *Orioli et al., 2020*).

To describe and understand how well ABSURDer works, we have applied it both to synthetic and experimental data for T4L. We used data of increasing complexity in an attempt to disentangle problems arising from the sampling, the force field, the forward model to calculate relaxation data and our approach to fit the data. In all cases, we employed the 1500 $10\,\mathrm{ns}$ blocks coming from the three $5\,\mu\mathrm{s}$ simulations with the Amber ff99SB*-ILDN force field to fit the data. In our first application we created synthetic data using the same force field by running five additional $1\,\mu\mathrm{s}$-long simulations. In this case, the only differences between the 'data' and the simulations would arise from insufficient sampling. In our second application, we again created synthetic data, in this case however using the Amber ff15ipq force field (with a modified methyl torsion potential) and the SPC/E$_b$ water model (*Takemura and Kitao, 2012*; *Debiec et al., 2016*; *Hoffmann et al., 2020*). We

performed three 1 μs simulations and used these to generate synthetic NMR relaxation data. This
test enables us to examine how the method behaves where there are real differences between the
potential used in the simulations we use to fit the data, and that give rise to the 'experimental data'.
In contrast to using actual experimental data, however, we here have access to the full underlying
ensembles and dynamics, and thus are able to compare e.g. full spectral density function and ro-
tamer distributions. Finally, we applied ABSURDer to experimental NMR relaxation data, providing
an example of how the approach might be used in practice. The results of these three levels of
complexity are described in the following sections.

Determining block size and assessing convergence

The choice of the block length plays a central role in the calculation of the relaxation rates and
in the possibility to optimize the results via reweighting. For this, two opposing effects need to
be considered (Fig. 2A). On one hand, a long block size allows for a more correct representation
of the correlation functions and thus estimation of the NMR relaxation rates. On the other hand,
it is desirable to increase the variation across the different blocks, as this makes it easier to fit
the relaxation data by reweighting; however this variance decreases with the length of the blocks
(Fig. 2A). Ideally, one would choose a block length that is (i) long enough not to introduce too much
bias compared to calculating correlation functions over the full trajectory and (ii) short enough that
each block is not just a 'converged' representation of the full trajectory. Thus, we chose 10 ns as the
block length that balances these two requirements (Fig. 2A). This choice allows us to calculate the
internal time correlation functions of the methyl C-H bonds up to a maximum lag time of 5 ns, as
also done previously (*Hoffmann et al., 2020*). We note that the chosen block size is also close to
the global tumbling time of the protein ($\tau_r \approx 11$ ns). As NMR spin relaxation experiments are mostly
sensitive to motions on time scales up to approximately τ_r, this explains why shorter block sizes
do not capture the NMR relaxation parameters accurately.

To assess the impact of sampling, we split our three 5 μs-long simulations into 15 1 μs segments
and employed leave-N-out cross-validation to estimate the impact of the amount of sampling for
the estimation of relaxation rates. For each value of N, we calculated the NMR relaxation rates
by averaging over the N segments, and then compared to the average over the remaining $15 - N$
segments by calculating the root mean square deviation (RMSD) between the two sets of rates.
We repeated these calculations for all possible combinations of leaving out N segments, and the
RMSD values were averaged. Finally, we repeated these calculations for all values of N ($N = 1, \ldots, 7$)
(Fig. 2B–D). Even at $N = 1$ we find relative low RMSD values when compared to the spread of
the rates (Fig. 3A–C), and these errors decrease further as additional sampling is included. Thus,
we conclude that using several microseconds of sampling we are able to obtain relatively precise
estimates of the relaxation rates, as would be expected given that they report on ps–ns dynamics
in the protein.

Fitting synthetic data generated with the same force field

We ran five 1 μs-long simulations with the Amber ff99SB*-ILDN force field and calculated NMR relax-
ation parameters from these. In what follows, this data will be referred to as 'NMR' or 'experiments'
to indicate that we treat them as observables from an experiment. We calculated the same observ-
ables as the average over 1500 10 ns blocks coming from the three 5 μs simulations with the same
force field, and compared them to the synthetic experimental data (Fig. 3A–C). As expected the
calculated and experimental values are strongly correlated, in line with the fact that these were
generated from the same underlying physical model. The calculated values of a reduced χ^2 were
25, 10 and 34 for $R(D_z)$, $R(D_y)$ and $R(3D_z^2 - 2)$, respectively.
We then asked whether we could improve the agreement between experiments and simula-
tions by changing the weights of each of the 1500 blocks to reweight the calculated observables.
We thus determined a set of weights that minimize the functional $\mathcal{L}(\mathbf{w}) = \chi^2(\mathbf{w}) - \theta S_{rel}(\mathbf{w})$ at differ-
ent values of θ and plot the resulting χ^2 vs. ϕ_{eff} (Fig. 3A–C; inserts). It is clear that as θ is decreased

Figure 2. Choice of block size and convergence of NMR relaxation rates. (A) Overall (blue) RMSD of NMR relaxation rates and (red) average inverse variance with respect to the full MD data set as a function of the block length. (B-D) Average RMSD of the three NMR relaxation rates obtained by comparing a set of N 1 μs long segments to the rates calculated from the remaining $15 - N$ segments.

Figure 3. ABSURDer applied to synthetic data generated with the same force field. (A – C) Comparison between 'experiment' and simulation for $R(D_z)$, $R(D_y)$ and $R(3D_z^2 - 2)$ both before and after reweighting with ABSURDer. The insets show the behaviour of the respective reduced χ^2 vs. ϕ_{eff} during the reweighting using different values of θ. The chosen $\theta = 300$ (resulting in $\phi_{eff} \approx 0.2$) is shown as a red cross. (D) The spectral density function and $R(D_z)$, $R(D_y)$ and $R(3D_z^2 - 2)$ of THR26-C$^{\gamma 2}$H$^{\gamma 2}$ before and after reweighting with ABSURDer. $J(0)$, $J(\omega)$ and $J(2\omega)$ are shown as black triangles. The errorbars represent the standard error of the mean from averaging over the 1500 10 ns sub-trajectories. (E) The χ_1 angle probability density distribution of THR26 before and after reweighting with ABSURDer.

to put greater weight on fitting the data, the agreement between experiment and simulation can be improved substantially, though at the cost of utilizing only a fraction of the input simulations. For simplicity, we here opted to use a value θ that gives rise to $\phi_{eff} \approx 0.2$, though our conclusions are relatively robust to this choice. At this level of fitting the agreement between experiment and simulations improves substantially (reduced χ^2 approximately 3, 1 and 2 for $R(D_z)$, $R(D_y)$ and $R(3D_z^2 - 2)$, respectively). We note also that $\phi_{eff} \approx 0.2$ represents $3\,\mu s$ of MD simulation, which in itself is sufficient to obtain relatively converged values (Fig. 2B–D).

While the NMR relaxation parameters are the experimental observables, they are calculated from the simulations via the spectral density functions, $J(\omega)$. Spectral density functions are often estimated from experiments using simplified and analytically tractable functions that balance the complexity of the motions and the number of free parameters that can be estimated from limited experiments. Using synthetic data gives us the possibility of comparing the spectral density functions from the simulations, both before and after reweighting, with those that give rise to the experiments. We here exemplify this using a single methyl group from THR26 (Fig. 3D); the code and data to generate plots for all residues are available online (github.com/KULL-Centre/papers/tree/master/2020/AB kummerer-orioli-et-al-2020). The observed NMR relaxation parameters depend on the spectral density at three frequencies ($\omega = 0$, ω_D and ω_{2D}, respectively) and it is thus not surprising that fitting to this data improves agreement at these frequencies. Nevertheless, it is also clear that our reweighting of the trajectories leads to a general improvement between the calculated spectral density function and that which was used to generate the synthetic data. As it is clear from this example (Fig. 3D), and from the overall results on all rates (Fig. 3A–C), it is also evident that discrepancies between experiments and simulations remain. While it is possible to fit the data more closely by decreasing θ this comes with the risk of potential overfitting and diminished trust of the input simulations.

The observed NMR relaxation parameters depend on a complex combination of both overall tumbling and various types of internal motions. In addition to the fast rotation of the methyl groups, this includes fluctuations of the backbone and motions both within and between rotameric states. It has previously been shown that rotamer jumps, in particular when this occurs on a timescale faster than rotational tumbling, can contribute substantially to side chain relaxation (*Best et al., 2005*; *Hu et al., 2005*). We thus calculated the side chain dihedral angles and compared the values before and after reweighting to those from the simulation used to generate the synthetic data. Highlighting again THR26 we see that, due to imperfect sampling of rotamer distributions, even in multi-microsecond simulations, there are clear differences between the distribution of the χ_1 dihedral angle in the simulations used to generate the data compared to that used to fit it (Fig. 3E). Particularly important, however, we also see a clear improvement after the reweighting. Thus, we find that by reweigting a 'dynamic' property (NMR relaxation rates) we also find improvement in a 'static' property such as the distributions of dihedral angles. Similar results were found for a wide range of dihedral angles including also in the backbone (Fig. S1) (see also github.com/KULL-Centre/papers/tree/master/2020/ABSURDer-kummerer-orioli-et-al-2020).

Fitting synthetic data generated with a different force field

The calculations above demonstrate how ABSURDer can be used to fit NMR data using MD simulations, but neglects the challenges arising from imperfect force fields. Thus, we increased the complexity of our comparisons by generating synthetic data with a different force field than the one we used to fit the data. We thus ran three $1\,\mu s$-long simulations with the Amber ff15ipq force field and calculated synthetic NMR relaxation parameters from these. We again used ABSURDer to fit the 1500 $10\,ns$ blocks to this data. In this case, imperfect agreement arises both due to insufficient sampling and differences in the underlying potential, leading to differences both in the dynamics and thermodynamic averages.

This added complexity is evident when we compare 'experimental' and calculated NMR relaxation rates, which show substantial differences and reduced χ^2 values that are about an order

Figure 4. ABSURDer applied to synthetic data generated with different force fields. (A - C) $R(D_z)$, $R(D_y)$ and $R(3D_z^2 - 2)$ from both data sets are compared before and after reweighting with ABSURDer. The insets show the behaviour of the respective reduced χ^2 vs. ϕ_{eff} during the reweighting using different values of θ. The chosen $\theta = 4500$ corresponding to $\phi_{eff} \approx 0.2$ is shown as a red cross. (D) Probability density distribution of the χ_1 and χ_2 angles of ILE9 before and after applying ABSURDer, and the corresponding 'experimental' data.

---

of magnitude greater than in the situation described above (78, 119 and 90 for $R(D_z)$, $R(D_y)$ and
$R(3D_z^2-2)$, respectively) (Fig. 4A–C). We can improve the agreement by applying the reweighting pro-
cedure, but in contrast to the situation described above, it is substantially more difficult to obtain
a good agreement. Thus, when selecting θ to give $\phi_{eff} = 0.2$ we reduce the χ^2 values by about two-
fold, compared to the about 10-fold in the case described above. Nevertheless, it is clear that by
changing the weights of the 1500 blocks we can improve the agreement between experiment and
simulations and remove a number of outliers (Fig. 4A–C). Looking at the different methyl groups,
we do not find that any specific residue gives rise to substantially worse agreement than others.

We again examined the consequences of the reweighting on the dihedral angles distributions,
in this case focusing on ILE9 (Fig. 4D). The NMR relaxation experiments probe both the dynamics
of the C$^\gamma$ and C$^\delta$ in isoleucine residues, and the χ_1 and χ_2 dihedrals may display substantial cou-
plings. We thus calculated the two-dimensional probability densities before and after reweighting
and compared to the distribution from the Amber ff15ipq simulations that we used to generate the
synthetic data. Despite systematic differences remain between the reweighted and 'experimental'
probability densities, one can clearly notice that ABSURDer is able to substantially correct the prob-
ability density around the χ_1 gauche$^+$ state by consequently decreasing the population in the χ_2
gauche$^+$ rotamer, which is populated with low probability in the 'experimental' data. We note that
although it is not unexpected, it is not trivial that a reweighting procedure employing kinetic data
is able to correct static quantities such as equilibrium probability density. As above, we also find
improvements in the backbone dihedral angles (Fig. S2).

Fitting experimental data

287   Having used synthetic data to show how ABSURDer makes it possible to fit MD simulations using
288   NMR relaxation experiments, we now proceed to fit to experimental data on T4L. Before discussing
289   the results, however, some comments are in order. In both cases above where we employed syn-

Figure 5. ABSURDer applied to experimental data. (A - C) $R(D_z)$, $R(D_y)$ and $R(3D_z^2 - 2)$ from both data sets are compared before and after reweighting with ABSURDer. The insets show the behaviour of the respective reduced χ^2 vs. ϕ_{eff} during the reweighting using different values of θ. The chosen $\theta = 1400$ corresponding to $\phi_{eff} \approx 0.2$ is shown as a red cross. (D-G) Distributions of $R(D_z)$, $R(D_y)$ and $R(3D_z^2 - 2)$ over the blocks for ALA134-C$^\beta$H$^\beta$ (D), LEU13-C$^{\delta1}$H$^{\delta1}$ (E), MET102-C$^\epsilon$H$^\epsilon$ (F) and VAL103-C$^{\gamma2}$H$^{\gamma2}$ (G) before and after reweighting with ABSURDer. Vertical lines represent the average relaxation rate for the respective data set.

₂₉₀ thetic data to mimic experimental NMR relaxation rates, we used relaxation rates for all 100 methyl
₂₉₁ groups in T4L. In practice, however, not all of these can be measured accurately e.g. due to over-
₂₉₂ lapping peaks or artifacts from strong ^{13}C–^{13}C coupling in certain residues (*Hoffmann et al., 2018b*).
₂₉₃ Thus, below we use only the 73 methyl groups whose relaxation rates were recently measured
₂₉₄ (*Hoffmann et al., 2018b*). To examine the effect of optimizing against a subset of residues we first
₂₉₅ repeated the reweighting procedure using the synthetic data, but restricting to the set of 73 methyl
₂₉₆ groups, and used the remaining 27 as a test set (Fig. S3). For both sources of synthetic data we
₂₉₇ find that optimizing on the 73 methyl groups lead to improvements in the remaining 27, unless
₂₉₈ aggressive reweighting to low ϕ_{eff} is employed.

₂₉₉ We thus proceeded to use the previously measured data (*Hoffmann et al., 2018b*) recorded at
₃₀₀ $950\,\text{MHz}$, and compared these to the NMR relaxation parameters from the three $5\,\mu\text{s}$ simulations,
₃₀₁ calculated as averages over the 1500 $10\,\text{ns}$ blocks (Fig. 5A–C). We find a reasonable agreement
₃₀₂ (reduced χ^2 values of about 119, 179 and 92 for $R(D_z)$, $R(D_y)$ and $R(3D_z^2 - 2)$, respectively), in line
₃₀₃ with similar calculations previously reported from ten $0.3\,\mu\text{s}$ simulations. In contrast to the results
₃₀₄ described above for the synthetic data we observe, however, some systematic differences with the
₃₀₅ calculated rates on average being 16% lower than experiments.

₃₀₆ We applied ABSURDer to improve agreement with the NMR experiments. When fitting to $\phi_{eff} =$
₃₀₇ 0.2 we are only above to obtain modest improvements of the calculated rates, decreasing the χ^2
₃₀₈ values by on average 30%. Again we do not find the remaining differences to be dominated by a
₃₀₉ single residue type.

₃₁₀ We examined why some relaxation rates can be improved while others appear not to be im-
₃₁₁ proved by ABSURDer. Reweighting approaches such as ABSURDer rely on finding a subensemble
₃₁₂ (or subset of trajectories) that fits the experimental data better than the full ensemble when aver-
₃₁₃ aged uniformly. Thus, the successful application requires that the calculated rates for the different

³¹⁴ blocks can be combined linearly to fit the data. We thus calculated the distribution of the relaxation
³¹⁵ rates over the 1500 blocks for selected methyl groups and compared the averages before and after
³¹⁶ reweighting to the experiments (Fig. 5D–G). For some residues and relaxation rates, exemplified
³¹⁷ by ALA134-$C^\beta H^\beta$ and LEU13-$C^{\delta 1}H^{\delta 1}$ in Fig. 5D–E, we find a relatively broad distribution where the
³¹⁸ relaxation rates calculated for the different blocks overlap with the experimental rates. In these
³¹⁹ cases it is possible to improve agreement by increasing the weights of some blocks and decreas-
³²⁰ ing the weights of others. In several other cases, exemplified by MET102-$C^\epsilon H^\epsilon$ and VAL103-$C^{\gamma 2}H^{\gamma 2}$
³²¹ in Fig. 5F–G, the experimental relaxation rates lie outside the range sampled in our (blocks of) MD
³²² simulations. In such cases, reweighting cannot improve agreement substantially, because no linear
³²³ combination of the blocks can get close to the experiments.

Conclusions

³²⁵ We have here presented ABSURDer, an extension of the ABSURD approach (*Salvi et al., 2016*),
³²⁶ and applied it to NMR relaxation measurements of side chain dynamics. We have described and
³²⁷ validated ABSURDer using synthetic data and applied it to experimental NMR data. As expected, for
³²⁸ the three levels of 'complexity' we obtain different levels of agreement after reweighting between
³²⁹ simulated and 'experimental' data.

³³⁰ When we used the same force field to generate synthetic data as that used to fit the data
³³¹ (ff99SB*-ILDN) we find good agreement even before reweighting, in line with the fact that the cal-
³³² culations of the relaxation parameters are relatively precise even using only a few microseconds
³³³ of simulations (Fig. 2). Nevertheless, we can improve agreement further by reweighting and find,
³³⁴ for example, improvements in dihedral angle distributions when we fit against the relaxation rates.
³³⁵ As expected from the fact that we fit against the relaxation rates, we also find overall improvement
³³⁶ of the spectral density functions after reweighting.

³³⁷ The overall observations were similar when we generated synthetic data using Amber ff15ipq
³³⁸ and fitted using simulations generated with ff99SB*-ILDN. Here, we are still able to achieve a sub-
³³⁹ stantial improvement in the quality of the fit (around a 50% decrease in the overall χ^2). Notably,
³⁴⁰ also in this case the overall agreement between 'experimental' and simulated spectral densities
³⁴¹ and rotamer distributions increases upon reweighting.

³⁴² When we fitted the experimental NMR data to the simulations with ff99SB*-ILDN we obtain a
³⁴³ more modest improvement (an overall 30% decrease in the χ^2). Examining the probability distri-
³⁴⁴ bution of rates over the different blocks and residues, we find that while some of these are broad
³⁴⁵ and highly overlapping with the experimental average, many others are sharp and incompatible
³⁴⁶ with the experimentally determined rates. Clearly, it is difficult to reweight when there is only little
³⁴⁷ overlap between the experimental value and the rates calculated from the different blocks. This
³⁴⁸ problem is exacerbated by the fact that we fit all rates simultaneously leading to a problem akin
³⁴⁹ to the 'curse of dimensionality'.

³⁵⁰ The natural question arises whether it is possible to increase the agreement between simulated
³⁵¹ and experimental NMR relaxation rates further. As described in the Introduction, the main sources
³⁵² of error between simulated and experimental data are: (i) insufficient sampling, (ii) force field in-
³⁵³ accuracies, and (iii) remaining errors in the forward model used to estimate the relaxation rates
³⁵⁴ from the trajectories. In the absence of a measure of ground truth it may, however, be difficult to
³⁵⁵ disentangle these effects.

³⁵⁶ Our analysis of convergence due to sampling (Fig. 2B–D) show that the rates calculated from the
³⁵⁷ 15 μs of simulation are rather precise compared e.g. to the deviation between experimental and
³⁵⁸ calculated rates (Fig. 5). While this suggests that sampling is not the main source of deviation, the
³⁵⁹ situation during reweighting is more complicated. Indeed, the selection of the block size demon-
³⁶⁰ strates one of the complicating factors (Fig. 2A); using a too short block size leads to systematic
³⁶¹ errors while using too long blocks leads to 'local convergence' and thus a narrow distribution of
³⁶² rates across the blocks. While a 10 ns block size strikes a reasonable balance between these oppos-
³⁶³ ing factors it is clear that there in many cases is not sufficient overlap between the distribution of

364 calculated rates and the experimental values (Fig. 5). Clearly, additional sampling to obtain more
365 blocks could alleviate this situation by increasing the number of samples in the tails of these distri-
366 butions, though in many cases the deviations are so large that substantially more sampling would
367 be needed.

368 As reweighting methods such as ABSURDer rely on an overlap between experiment and cal-
369 culated parameters prior to reweighting, they are aided by good agreement before reweighting
370 (*Hummer and Köfinger, 2015*; *Orioli et al., 2020*; *Larsen et al., 2020*). Indeed the extent of reweight-
371 ing needed to obtain good agreement is related to a measure of the error in the force field (*Qian,*
372 *2001*; *Orioli et al., 2020*). Thus, as starting point for our calculations we used force fields that
373 have recently been shown to give improved agreement to side chain relaxation data (*Hoffmann*
374 *et al., 2018a*,b, *2020*). Those studies used quantum-level calculations to modify the barriers for
375 the methyl spinning, and demonstrated improved agreement with experimental NMR relaxation
376 rates. More specifically, it was shown that methyl spinning barriers where generally too high, and
377 that a small decrease (obtained by fitting to coupled cluster quantum chemical calculations of iso-
378 lated dipeptides) lead to increased spinning rates and better agreement with experimental data in
379 both T4L and ubiquitin (*Hoffmann et al., 2018a*,b, *2020*). Other effects might, however, contribute
380 to the imperfect agreement between experiments and simulations. For example, the quantum
381 calculations did not include information about how methyl spinning rates are influenced e.g. by
382 side-chain packing or tunneling effects (*Chatfield and Wong, 2000*; *Chatfield et al., 2004*).

383 A final source of error between experiment and simulation that may also prevent reweighting is
384 the accuracy of the forward model used to calculate the relaxation rates from simulations. Here we
385 have used a recently described approach to calculate deuterium relaxation rates from MD simula-
386 tions (*Hoffmann et al., 2018a*,b). While the approach is highly accurate, there are some sources of
387 errors including for example the treatment of tumbling motions, non-symmetry of partially deuter-
388 ated methyl groups, finite sampling of the fastest dynamics (we save frames every $1\,\mathrm{ps}$) and fitting
389 of the correlation functions to a fixed number of exponentials.

390 In summary, we have presented ABSURDer and applied it to study motions in methyl containing
391 residues in a folded protein. The approach is general and may be extended to other types of
392 residues and nuclei, to other NMR measurements such as measurements of nuclear Overhauser
393 enhancements, and even to data beyond those generated by NMR.

Methods

Sampling

396 The starting structure for our MD simulations was the X-ray structure of the cystein-free T4L SER44GLY
397 mutant (PDB 107L) (*Blaber et al., 1993*) where we changed GLY44 back to a serine. We performed
398 three different sets of simulations: Three $5\,\mathrm{\mu s}$ long ("MD Data" in all three analyses) and five $1\,\mathrm{\mu s}$
399 long ("NMR Data" in the first synthetic data set) simulations, both using the AMBER ff99SB*-ILDN
400 / TIP4P-2005 protein force field (*Lindorff-Larsen et al., 2010*; *Hornak et al., 2006*; *Best and Hum-*
401 *mer, 2009*), and three $1\,\mathrm{\mu s}$ long simulations using the AMBER ff15ipq / SPC/E$_b$ (*Debiec et al., 2016*)
402 force field ("NMR Data" in the second synthetic data set). We applied a modification to the methyl
403 rotation barriers to both force fields as recently described (*Hoffmann et al., 2018a*, *2020*). In the
404 case of the AMBER ff15ipq simulation, we used the Amber input preparation module LEAP from
405 AmberTools17 (*Case et al., 2017*) to set up the system and we afterwards converted the topology
406 to GROMACS file format using ParmED (*Swails et al., 2010*).

407 All MD simulations were carried out with GROMACS v2018.1. We used a periodic truncated
408 dodecahedron of $400\,\mathrm{nm}^3$ volume as a simulation box, keeping 1.2 nm distance from the pro-
409 tein which was centered in the cell. We solvated the system with 12230 TIP4P-2005 (*Abascal and*
410 *Vega, 2005*) water molecules and 12247 SPC/E$_b$water molecules, respectively, retaining crystal wa-
411 ters. To neutralize the system and simulate it in a physiological salt concentration, we also added
412 $150\,\mathrm{mmol\,L^{-1}}$ Na$^+$/Cl$^-$ ions. We minimized the systems with 50000 steps of steepest descent and

equilibrated them for $200\,\mathrm{ps}$ in the *NPT* ensemble, using harmonic position restraints (force con-
stants of $1000\,\mathrm{kJ\,mol^{-1}\,nm^{-2}}$) on the heavy atoms of the protein. Equations of motion were inte-
grated using the leap-frog algorithm. We set a cut-off of $1\,\mathrm{nm}$ for the Van der Waals and Coulomb
interactions and employed PME for long-range electrostatics (*Essmann et al., 1995*) with 4^{th} order
cubic interpolation and $0.16\,\mathrm{nm}$ Fourier grid spacing. We ran the production simulations in the *NPT*
ensemble, using the velocity re-scaling thermostat (*Bussi et al., 2007*) with a reference temperature
of $300\,\mathrm{K}$ and thermostat timescale of $\tau_T = 1\,\mathrm{ps}$, and the Parrinello-Rahman barostat with a reference
pressure of $1\,\mathrm{bar}$, a barostat timescale $\tau_P = 2\,\mathrm{ps}$ and an isothermal compressibility of $4.5 \times 10^{-5}\,\mathrm{bar^{-1}}$.
Finally, we saved the protein coordinates every $1\,\mathrm{ps}$ and, after running the simulations, we removed
the overall tumbling of the protein by fitting to a reference structure.

Spectral Density Mapping

We used a previously described approach (*Hoffmann et al., 2018b*) to calculate NMR relaxation
rates from computed spectral densities. The original work used multiple $300\,\mathrm{ns}$-long simulations,
however, we analysed our simulations in a block-wise fashion, using $10\,\mathrm{ns}$ long, non-overlapping
blocks. This yielded 1500 blocks for the set of 3x $5\,\mathrm{\mu s}$-long simulations, 500 blocks for the set of
5x $1\,\mathrm{\mu s}$-long simulations and 300 blocks for the 3x $1\,\mathrm{\mu s}$ long simulations. We started by calculating
the internal time correlation function (TCF), *i.e.* without global tumbling, of the three C_{methyl}-H^i_{methyl}
($i = 1, 2, 3$) bond vectors up to a maximum lag-time of $5\,\mathrm{ns}$ for all side-chain methyl groups. Next,
we fitted the internal TCFs with six exponential functions and an offset,

$$C_{\mathrm{int,exp}}(t) = \sum_{i=1}^{6} A_i e^{-t/\tau_i} + S^2_{\mathrm{long}}, \tag{1}$$

432   where $\sum_{i=1}^{6} A_i + S^2_{\mathrm{long}} = 1, 0 \le A_i \le 1, 0 \le S^2_{\mathrm{long}} \le 1, 0 \le \tau_1 \le 10ps, 0 \le \tau_2 \le 50ps, 0 \le \tau_3 \le 200ps, 0 \le \tau_4 \le$
433    $500ps, 0 \le \tau_5 \le 1000ps, \tau_6 \ge 0$ and S^2_{long} is the long-time limit order parameter. Since the protein was
434   found to be best represented by an axially symmetric tumbling model (*Hoffmann et al., 2018b*), we
435   introduced methyl-specific global tumbling times $\tau_{R,i}$ by multiplying the internal TCF with a single-
436   exponential thus, yielding the total TCF:

$$C(t) = C_{\mathrm{int,exp}}(t) e^{-t/\tau_{R,i}}. \tag{2}$$

437   In this case, we applied the experimental methyl group-specific tumbling times, which, for the
438   synthetic data sets, we extracted based on analysing backbone N–H relaxation (*Maragakis et al.,*
439   *2008*; *Hoffmann et al., 2018b*). Subsequently, we transformed the total TCF into a spectral density
440   function:

$$J(\omega) = \sum_{i=1}^{6} \frac{A_i \tau_i^{\mathrm{eff}}}{1 + (\omega \tau_i^{\mathrm{eff}})^2} + \frac{S^2_{\mathrm{long}} \tau_{R,i}}{1 + (\omega \tau_{R,i})^2}, \tag{3}$$

441   where $\tau_i^{\mathrm{eff}} = (\tau_i \tau_{R,i})/(\tau_i + \tau_{R,i})$. Finally, we calculated the relaxation rates directly from $J(0)$, $J(\omega_D)$,
442    $J(2\omega_D)$ via

$$R\left(D_z\right) = \frac{1}{32} \left(\frac{qQe^2}{\hbar}\right)^2 \left[J\left(\omega_D\right) + 4J\left(2\omega_D\right)\right], \tag{4}$$

$$R\left(D_y\right) = \frac{1}{32} \left(\frac{qQe^2}{\hbar}\right)^2 \left[9J(0) + 15J\left(\omega_D\right) + 6J\left(2\omega_D\right)\right], \tag{5}$$

$$R\left(3D_z^2 - 2\right) = \frac{1}{32} \left(\frac{qQe^2}{\hbar}\right)^2 \left[3J\left(\omega_D\right)\right], \tag{6}$$

where $\left(qQe^2/\hbar\right)^2$ is the quadrupolar coupling constant of deuterium. We used $145.851\,\mathrm{MHz}$ as the
Larmor frequency ω_D of deuterium for all data sets which corresponds to a Bruker magnetic field
strength of $22.3160\,\mathrm{T}$ that was also used to measure the experimental NMR relaxation rates.
We used this approach for each side-chain methyl group and each of the 1500 blocks to calcu-
late sets of the three relaxation rates, which we employed as a input to the ABSURDer reweighting.

When generating synthetic data, we calculated average rates over the blocks, and we used the
standard error of the mean over the blocks as the errors when calculating χ^2 and for reweighting
with ABSURDer.

453 Reweighting

Our reweighting approach builds upon the previously described ABSURD method (*Salvi et al.,*
2016), where a longer set of trajectories are divided into *blocks*, each of which is long enough to
estimate the relaxation rates of interest. Finally, a weighted average of the rates is performed over
the blocks, where the weights, **w**, are obtained through the optimisation of a functional which de-
termines the agreement between simulated and experimental rates. In particular, the ABSURD
functional is given by

$$\mathcal{R}(\mathbf{w}) = \sum_{r=1}^{N_r} \sum_{n=1}^{M} \left[\frac{R_{r,n}^{\text{exp}} - \sum_{i=1}^{N} w_i R_{r,n}^i}{\sqrt{(\sigma_{r,n}^{\text{exp}})^2 + (\sigma_{r,n}^{\text{calc}})^2}} \right]^2 , \quad \sum_{i=1}^{N} w_i = 1 \tag{7}$$

where r ranges over the N_r types of experimental NMR relaxation rates, M is the overall number of
measured rates and N represents the number of blocks which the trajectory has been cut into, $\sigma_{r,n}^{\text{exp}}$
is the experimental error on the n-th rate of the r-th type, $\sigma_{r,n}^{\text{calc}}$ is the standard error on the simulated
rates, estimated by averaging all the n-th rates of the r-th type over the different blocks. The optimal
set of weights is obtained as the corresponding minimum in weight space, $\mathbf{w}^* = \min_{\mathbf{w}} \mathcal{R}(\mathbf{w})$. Each
weight is associated to a block from the trajectory and it encodes the relevance of each block to
the dynamical ensemble.

The functional in Eq. 7 is unregularised, which sometimes may lead to the risk of overfitting (*Ori-*
oli et al., 2020). In the original implementation of ABSURD, this was avoided by extensive cross-
validation using multiple relaxation rates, measured at multiple magnetic fields. A complementary
approach is to use the MD simulations as a statistical prior and to balance the trust in the data and
the experiments (*Orioli et al., 2020*).

We take this latter approach by introducing an extension to the ABSURD functional (which we
denominate ABSURDer) to make it less prone to overfitting. In particular, we add a Shannon
relative-entropy restraint $S(\mathbf{w})$ (*Kullback and Leibler, 1951*; *Gull and Daniell, 1978*) and the new
functional takes the following form:

$$\mathcal{T}(\mathbf{w}) = \sum_{r=1}^{N_r} \sum_{n=1}^{M} \left[\frac{R_{r,n}^{\text{exp}} - \sum_{i=1}^{N} w_i R_{r,n}^i}{\sqrt{(\sigma_{r,n}^{\text{exp}})^2 + (\sigma_{r,n}^{\text{calc}})^2}} \right]^2 + \theta \sum_{i=1}^{N} w_i \log\left(\frac{w_i}{w_i^0}\right) = \chi^2(\mathbf{w}) - \theta S(\mathbf{w}) \tag{8}$$

where $w_i^0 = 1/N$ represent the initial, uniform weights provided by the prior. The parameter θ, is
used to set the balance between our trust in the prior and in the experimental data. In particular,
for $\theta \to \infty$, an infinite trust is put on the prior and the weights w_i stay at their initial values. In the
opposite limit, $\theta \to 0$, all the trust is put on the experimental data and the weights are allowed to
change freely to accommodate this trust. In this case, the functional $\mathcal{T}(\mathbf{w})$ reduces to that used in
ABSURD. The two limits of θ can be seen also from parameter $\phi_{\text{eff}}(\mathbf{w}) = \exp(S(\mathbf{w}))$, which provides
a measure of the effective fraction of blocks retained for a given set of weights. In particular, for
$\theta \to \infty$ we have $\phi_{\text{eff}}(\mathbf{w}) = 1$, as $w_i \simeq w_i^0$. On the other hand, for $\theta \to 0$ and in the case where the
prior is very different from the experimental data, $\phi_{\text{eff}}(\mathbf{w}) \to 0$.

We minimised functional $\mathcal{T}(\mathbf{w})$ using the limited memory Broyden-Fletcher-Goldfarb-Shanno
algorithm (L-BFGS-B) in its SciPy implementation (*Virtanen et al., 2020*). In each reweighting run
we employed 30 values of θ selected in the range $[0, 8000]$. To accelerate convergence, we started
by optimising the $\theta = 8000$ functional using $\mathbf{w}_0$ as a guess set of weights, then we employed the
obtained optimal weights as guess for the successive minimisation run and so on, iteratively. We
stress that the value of $\mathbf{w}_0$ in Eq. 8 remained unchanged throughout the minimisation, regardless

of the choice of the guess input weights. The ABSURDer code is available under a GNU GPL v3.0 license at github.com/KULL-Centre/ABSURDer and scripts and data to generate the results in this paper are available at github.com/KULL-Centre/papers/tree/master/2020/ABSURDer-kummerer-orioli-et-al-2020.

Acknowledgments

We are grateful to Profs. Lars V. Schäfer and Frans A.A. Mulder for discussions, help and comments on the manuscript. We acknowledge support by a grant from the Lundbeck Foundation to the BRAINSTRUC structural biology initiative, the NordForsk Nordic Neutron Science Programme, the Carlsberg Foundation, and the Novo Nordisk Foundation [grant number NNF15OC0016360].

References

Abascal JL, Vega C. A general purpose model for the condensed phases of water: TIP4P/2005. The Journal of chemical physics. 2005; 123(23):234505.

Andrae R, Schulze-Hartung T, Melchior P. Dos and don'ts of reduced chi-squared. arXiv preprint arXiv:10123754. 2010; .

Bernardi RC, Melo MC, Schulten K. Enhanced sampling techniques in molecular dynamics simulations of biological systems. Biochimica et Biophysica Acta (BBA)-General Subjects. 2015; 1850(5):872–877.

Best RB, Clarke J, Karplus M. What contributions to protein side-chain dynamics are probed by NMR experiments? A molecular dynamics simulation analysis. Journal of molecular biology. 2005; 349(1):185–203.

Best RB, Hummer G. Optimized molecular dynamics force fields applied to the helix- coil transition of polypeptides. The journal of physical chemistry B. 2009; 113(26):9004–9015.

Best RB, Vendruscolo M. Determination of protein structures consistent with NMR order parameters. Journal of the American Chemical Society. 2004; 126(26):8090–8091.

Blaber M, Zhang XJ, Matthews BW. Structural basis of amino acid alpha helix propensity. Science. 1993; 260(5114):1637–1640.

Bonomi M, Heller GT, Camilloni C, Vendruscolo M. Principles of protein structural ensemble determination. Current opinion in structural biology. 2017; 42:106–116.

Bottaro S, Bengtsen T, Lindorff-Larsen K. Integrating molecular simulation and experimental data: a Bayesian/maximum entropy reweighting approach. In: *Structural Bioinformatics* Springer; 2020.p. 219–240.

Bottaro S, Bussi G, Kennedy SD, Turner DH, Lindorff-Larsen K. Conformational ensembles of RNA oligonucleotides from integrating NMR and molecular simulations. Science advances. 2018; 4(5):eaar8521.

Bottaro S, Lindorff-Larsen K. Biophysical experiments and biomolecular simulations: A perfect match? Science. 2018; 361(6400):355–360.

Bowman GR. Accurately modeling nanosecond protein dynamics requires at least microseconds of simulation. Journal of computational chemistry. 2016; 37(6):558–566.

Bremi T, Brüschweiler R, Ernst RR. A protocol for the interpretation of side-chain dynamics based on NMR relaxation: application to phenylalanines in antamanide. Journal of the American Chemical Society. 1997; 119(18):4272–4284.

Brookes DH, Head-Gordon T. Experimental inferential structure determination of ensembles for intrinsically disordered proteins. Journal of the American Chemical Society. 2016; 138(13):4530–4538.

Bussi G, Donadio D, Parrinello M. Canonical sampling through velocity rescaling. The Journal of chemical physics. 2007; 126(1):014101.

Case D, Cerutti D, Cheatham III T, Darden T, Duke R, Giese T, Gohlke H, Goetz A, Greene D, Homeyer N, et al. AMBER 2017. San Francisco: University of California. 2017; .

Case DA. Molecular dynamics and NMR spin relaxation in proteins. Accounts of chemical research. 2002; 35(6):325–331.

536 Cesari A, Reißer S, Bussi G. Using the maximum entropy principle to combine simulations and solution exper-
537 iments. Computation. 2018; 6(1):15.

538 Chatfield DC, Augsten A, D'cunha C. Correlation times and adiabatic barriers for methyl rotation in SNase.
539 Journal of Biomolecular NMR. 2004; 29(3):377–385.

540 Chatfield DC, Wong SE. Methyl motional parameters in crystalline l-alanine: molecular dynamics simulation
541 and NMR. The Journal of Physical Chemistry B. 2000; 104(47):11342–11348.

542 Chen Pc, Shevchuk R, Strnad FM, Lorenz C, Karge L, Gilles R, Stadler AM, Hennig J, Hub JS. Combined small-angle
543 X-ray and neutron scattering restraints in molecular dynamics simulations. Journal of chemical theory and
544 computation. 2019; 15(8):4687–4698.

545 Clore GM, Szabo A, Bax A, Kay LE, Driscoll PC, Gronenborn AM. Deviations from the simple two-parameter
546 model-free approach to the interpretation of nitrogen-15 nuclear magnetic relaxation of proteins. Journal
547 of the American Chemical Society. 1990; 112(12):4989–4991.

548 Cordeiro TN, Chen Pc, De Biasio A, Sibille N, Blanco FJ, Hub JS, Crehuet R, Bernadó P. Disentangling polydis-
549 persity in the PCNA- p15PAF complex, a disordered, transient and multivalent macromolecular assembly.
550 Nucleic acids research. 2017; 45(3):1501–1515.

551 Crehuet R, Buigues PJ, Salvatella X, Lindorff-Larsen K. Bayesian-Maximum-Entropy reweighting of IDP ensem-
552 bles based on NMR chemical shifts. Entropy. 2019; 21(9):898.

553 Debiec KT, Cerutti DS, Baker LR, Gronenborn AM, Case DA, Chong LT. Further along the road less traveled:
554 AMBER ff15ipq, an original protein force field built on a self-consistent physical model. Journal of chemical
555 theory and computation. 2016; 12(8):3926–3947.

556 Essmann U, Perera L, Berkowitz ML, Darden T, Lee H, Pedersen LG. A smooth particle mesh Ewald method.
557 The Journal of chemical physics. 1995; 103(19):8577–8593.

558 Gu Y, Li DW, Brüschweiler R. NMR order parameter determination from long molecular dynamics trajectories
559 for objective comparison with experiment. Journal of chemical theory and computation. 2014; 10(6):2599–
560 2607.

561 Gull SF, Daniell GJ. Image reconstruction from incomplete and noisy data. Nature. 1978; 272(5655):686–690.

562 van Gunsteren WF, Daura X, Hansen N, Mark AE, Oostenbrink C, Riniker S, Smith LJ. Validation of molecular
563 simulation: an overview of issues. Angewandte Chemie International Edition. 2018; 57(4):884–902.

564 Halle B, Wennerström H. Interpretation of magnetic resonance data from water nuclei in heterogeneous
565 systems. The Journal of Chemical Physics. 1981; 75(4):1928–1943.

566 Henriques J, Cragnell C, Skepö M. Molecular dynamics simulations of intrinsically disordered proteins: force
567 field evaluation and comparison with experiment. Journal of chemical theory and computation. 2015;
568 11(7):3420–3431.

569 Hoffmann F, Mulder FA, Schäfer LV. Predicting NMR relaxation of proteins from molecular dynamics simula-
570 tions with accurate methyl rotation barriers. The Journal of Chemical Physics. 2020; 152(8):084102.

571 Hoffmann F, Mulder FA, Schäfer LV. Accurate methyl group dynamics in protein simulations with AMBER force
572 fields. The Journal of Physical Chemistry B. 2018; 122(19):5038–5048.

573 Hoffmann F, Xue M, Schäfer LV, Mulder FA. Narrowing the gap between experimental and computational de-
574 termination of methyl group dynamics in proteins. Physical Chemistry Chemical Physics. 2018; 20(38):24577–
575 24590.

576 Hornak V, Abel R, Okur A, Strockbine B, Roitberg A, Simmerling C. Comparison of multiple Amber force fields
577 and development of improved protein backbone parameters. Proteins: Structure, Function, and Bioinfor-
578 matics. 2006; 65(3):712–725.

579 Hu H, Hermans J, Lee AL. Relating side-chain mobility in proteins to rotameric transitions: insights from molec-
580 ular dynamics simulations and NMR. Journal of biomolecular NMR. 2005; 32(2):151–162.

581 Hummer G, Köfinger J. Bayesian ensemble refinement by replica simulations and reweighting. The Journal of
582 chemical physics. 2015; 143(24):12B634_1.

583 **Köfinger J**, Stelzl LS, Reuter K, Allande C, Reichel K, Hummer G. Efficient ensemble refinement by reweighting.
584 Journal of chemical theory and computation. 2019; 15(5):3390–3401.

585 **Kullback S**, Leibler RA. On information and sufficiency. The annals of mathematical statistics. 1951; 22(1):79–
586 86.

587 **Larsen AH**, Wang Y, Bottaro S, Grudinin S, Arleth L, Lindorff-Larsen K. Combining molecular dynamics simula-
588 tions with small-angle X-ray and neutron scattering data to study multi-domain proteins in solution. PLoS
589 computational biology. 2020; 16(4):e1007870.

590 **Li DW**, Brüschweiler R. NMR-based protein potentials. Angewandte Chemie International Edition. 2010;
591 49(38):6778–6780.

592 **Li DW**, Brüschweiler R. Iterative optimization of molecular mechanics force fields from NMR data of full-length
593 proteins. Journal of chemical theory and computation. 2011; 7(6):1773–1782.

594 **Lindorff-Larsen K**, Best RB, DePristo MA, Dobson CM, Vendruscolo M. Simultaneous determination of protein
595 structure and dynamics. Nature. 2005; 433(7022):128–132.

596 **Lindorff-Larsen K**, Piana S, Palmo K, Maragakis P, Klepeis JL, Dror RO, Shaw DE. Improved side-chain torsion
597 potentials for the Amber ff99SB protein force field. Proteins: Structure, Function, and Bioinformatics. 2010;
598 78(8):1950–1958.

599 **Lipari G**, Szabo A. Model-free approach to the interpretation of nuclear magnetic resonance relaxation
600 in macromolecules. 1. Theory and range of validity. Journal of the American Chemical Society. 1982;
601 104(17):4546–4559.

602 **Lipari G**, Szabo A, Levy RM. Protein dynamics and NMR relaxation: comparison of simulations with experiment.
603 Nature. 1982; 300(5888):197–198.

604 **Maragakis P**, Lindorff-Larsen K, Eastwood MP, Dror RO, Klepeis JL, Arkin IT, Jensen MØ, Xu H, Trbovic N, Friesner
605 RA, et al. Microsecond molecular dynamics simulation shows effect of slow loop dynamics on backbone
606 amide order parameters of proteins. The Journal of Physical Chemistry B. 2008; 112(19):6155–6158.

607 **McCammon JA**, Gelin BR, Karplus M. Dynamics of folded proteins. Nature. 1977; 267(5612):585–590.

608 **Nerenberg PS**, Head-Gordon T. New developments in force fields for biomolecular simulations. Current opin-
609 ion in structural biology. 2018; 49:129–138.

610 **Nodet G**, Salmon L, Ozenne V, Meier S, Jensen MR, Blackledge M. Quantitative description of backbone con-
611 formational sampling of unfolded proteins at amino acid resolution from NMR residual dipolar couplings.
612 Journal of the American Chemical Society. 2009; 131(49):17908–17918.

613 **Norgaard AB**, Ferkinghoff-Borg J, Lindorff-Larsen K. Experimental parameterization of an energy function for
614 the simulation of unfolded proteins. Biophysical journal. 2008; 94(1):182–192.

615 **O'Brien ES**, Wand AJ, Sharp KA. On the ability of molecular dynamics force fields to recapitulate NMR derived
616 protein side chain order parameters. Protein Science. 2016; 25(6):1156–1160.

617 **Orioli S**, Larsen AH, Bottaro S, Lindorff-Larsen K. Chapter Three - How to learn from inconsistencies: Integrating
618 molecular simulations with experimental data. In: Strodel B, Barz B, editors. *Computational Approaches for*
619 *Understanding Dynamical Systems: Protein Folding and Assembly*, vol. 170 of Progress in Molecular Biology and
620 Translational Science Academic Press; 2020.p. 123 – 176. http://www.sciencedirect.com/science/article/pii/
621 S1877117319302121, doi: https://doi.org/10.1016/bs.pmbts.2019.12.006.

622 **Piana S**, Robustelli P, Tan D, Chen S, Shaw DE. Development of a force field for the simulation of single-chain
623 proteins and protein-protein complexes. Journal of Chemical Theory and Computation. 2020; .

624 **Prompers JJ**, Brüschweiler R. General framework for studying the dynamics of folded and nonfolded pro-
625 teins by NMR relaxation spectroscopy and MD simulation. Journal of the American Chemical Society. 2002;
626 124(16):4522–4534.

627 **Qian H**. Relative entropy: Free energy associated with equilibrium fluctuations and nonequilibrium deviations.
628 Physical Review E. 2001; 63(4):042103.

629 **Rauscher S**, Gapsys V, Gajda MJ, Zweckstetter M, de Groot BL, Grubmüller H. Structural ensembles of intrinsi-
630 cally disordered proteins depend strongly on force field: a comparison to experiment. Journal of chemical
631 theory and computation. 2015; 11(11):5513–5524.

632 Robustelli P, Piana S, Shaw DE. Developing a molecular dynamics force field for both folded and disordered
633 protein states. Proceedings of the National Academy of Sciences. 2018; 115(21):E4758–E4766.

634 Salvi N, Abyzov A, Blackledge M. Multi-timescale dynamics in intrinsically disordered proteins from NMR relax-
635 ation and molecular simulation. The journal of physical chemistry letters. 2016; 7(13):2483–2489.

636 Salvi N, Abyzov A, Blackledge M. Solvent-dependent segmental dynamics in intrinsically disordered proteins.
637 Science advances. 2019; 5(6):eaax2348.

638 Showalter SA, Johnson E, Rance M, Brüschweiler R. Toward quantitative interpretation of methyl side-chain
639 dynamics from NMR by molecular dynamics simulations. Journal of the American Chemical Society. 2007;
640 129(46):14146–14147.

641 Skrynnikov NR, Millet O, Kay LE. Deuterium spin probes of side-chain dynamics in proteins. 2. Spectral density
642 mapping and identification of nanosecond time-scale side-chain motions. Journal of the American Chemical
643 Society. 2002; 124(22):6449–6460.

644 Swails J, Hernandez C, Mobley D, Nguyen H, Wang L, Janowski P, ParmEd; 2010.

645 Takemura K, Kitao A. Water model tuning for improved reproduction of rotational diffusion and NMR spectral
646 density. The Journal of Physical Chemistry B. 2012; 116(22):6279–6287.

647 Vasile F, Tiana G. Determination of structural ensembles of flexible molecules in solution from NMR data
648 undergoing spin diffusion. Journal of chemical information and modeling. 2019; 59(6):2973–2979.

649 Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, Burovski E, Peterson P, Weckesser
650 W, Bright J, van der Walt SJ, Brett M, Wilson J, Jarrod Millman K, Mayorov N, Nelson ARJ, Jones E, Kern R, Larson
651 E, Carey C, et al. SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python. Nature Methods.
652 2020; 17:261–272. doi: https://doi.org/10.1038/s41592-019-0686-2.

653 Supporting Material

Supporting Figure 1. Effect of ABSURDer reweighting on backbone dihedral angle distributions. Difference RMSD (ΔRMSD) between distributions for the backbone (A) ϕ and (B) ψ dihedral angles. We calculated the RMSD of the dihedral angle distributions before and after reweighting, in both cases using the distribution used to generate the synthetic data generated by Amber ff99SB*-ILDN as the reference. Thus, large positive values of ΔRMSD indicate highly improved distributions after reweighting whereas negative values indicate distributions worsened by the application of ABSURDer. The bars are colored by residue type and asterisks mark residues highlighted in panels C–F. Examples of residues for which ABSURDer (C,D) substantially improve the dihedral angle distributions and (E,F) worsens them.

Supporting Figure 2. Effect of ABSURDer reweighting on backbone dihedral angle distributions. Difference RMSD (ΔRMSD) between distributions for the backbone (A) ϕ and (B) ψ dihedral angles. We calculated the RMSD of the dihedral angle distributions before and after reweighting, in both cases using the distribution used to generate the synthetic data generated by Amber ff15ipq as the reference. Thus, large positive values of ΔRMSD indicate highly improved distributions after reweighting whereas negative values indicate distributions worsened by the application of ABSURDer. The bars are colored by residue type and asterisks mark residues highlighted in panels C–F. Examples of residues for which ABSURDer (C,D) substantially improve the dihedral angle distributions and (E,F) worsens them.

Supporting Figure 3. Effect of ABSURDer reweighting on NMR relaxation rates not included in the optimisation. We used ABSURDer with synthetic data to fit the relaxation rates for the (blue) 73 methyl groups that have been probed experimentally, and (red) cross-validate across the remaining 27 methyl groups. (A) Synthetic data generated with Amber ff99SB*-ILDN. (B) Synthetic data generated with Amber ff15ipq.