A Practical Alzheimer Disease Classifier via Brain Imaging-Based Deep Learning on 85,721 Samples

Bin Lu[1,2], Hui-Xian Li[1,2], Zhi-Kai Chang[1,2], Le Li[3], Ning-Xuan Chen[1,2], Zhi-Chen Zhu[1,2], Hui-Xia Zhou[1,2], Xue-Ying Li[1,4,5], Yu-Wei Wang[1,2], Shi-Xian Cui[1,4,5], Zhao-Yu Deng[1,2], Zhen Fan[6], Hong Yang[7], Xiao Chen[1,2], Paul M. Thompson[8], Francisco Xavier Castellanos[9,10], Chao-Gan Yan[1,2,11,12*], for the Alzheimer's Disease Neuroimaging Initiative[**]

[1]CAS Key Laboratory of Behavioral Science, Institute of Psychology, Beijing, China; [2]Department of Psychology, University of Chinese Academy of Sciences, Beijing, China; [3]Center for Cognitive Science of Language, Beijing Language and Culture University, Beijing, China; [4]Sino-Danish College, University of Chinese Academy of Science, Beijing, China; [5]Sino-Danish Center for Education and Research, Beijing, China; [6]Department of Neurosurgery, Huashan Hospital, Fudan University, Shanghai Neurosurgical Clinical Center, Shanghai, China; [7]Department of Radiology, The First Affiliated Hospital, College of Medicine, Zhejiang University, Hangzhou, Zhejiang, China; [8]Imaging Genetics Center, Mark & Mary Stevens Institute for Neuroimaging & Informatics, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA; [9]Department of Child and Adolescent Psychiatry, NYU Grossman School of Medicine, New York, NY, USA; [10]Nathan Kline Institute for Psychiatric Research, Orangeburg, NY, USA; [11]International Big-Data Center for Depression Research, Institute of Psychology, Chinese Academy of Sciences, Beijing, China; [12]Magnetic Resonance Imaging Research Center, Institute of Psychology, Chinese Academy of Sciences, Beijing, China.

*Corresponding author:

Chao-Gan Yan, Ph.D.

CAS Key Laboratory of Behavioral Science, Institute of Psychology, Beijing, China

16 Lincui Road, Chaoyang District, Beijing 100101, China

Tel: +86-10-64101582

Fax: +86-10-64101582

E-mail: yancg@psych.ac.cn

**Data used in preparation of this article were obtained from the Alzheimer's Disease

Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). As such, the investigators within

the ADNI contributed to the design and implementation of ADNI and/or provided data but did

not participate in analysis or writing of this report. A complete listing of ADNI investigators can

be found at:

http://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

Abstract

Beyond detecting brain lesions or tumors, comparatively little success has been attained in identifying brain disorders such as Alzheimer's disease (AD), based on magnetic resonance imaging (MRI). Many machine learning algorithms to detect AD have been trained using limited training data, meaning they often generalize poorly when applied to scans from previously unseen populations. Therefore, we built a practical brain MRI-based AD diagnostic classifier using deep learning/transfer learning on a dataset of unprecedented size and diversity. A retrospective MRI dataset pooled from more than 217 sites/scanners constituted one of the largest brain MRI samples to date (85,721 scans from 50,876 participants) between January 2017 and August 2021. Next, a state-of-the-art deep convolutional neural network, Inception-ResNet-V2, was built as a sex classifier with high generalization capability. The sex classifier achieved 94.9% accuracy and served as a base model in transfer learning for the objective diagnosis of AD. After transfer learning, the model fine-tuned for AD classification achieved 90.9% accuracy in leave-sites-out cross-validation on the Alzheimer's Disease Neuroimaging Initiative (ADNI, 6,857 samples) dataset and 94.5%/93.6%/91.1% accuracy for direct tests on three unseen independent datasets (AIBL, 669 samples / MIRIAD, 644 samples / OASIS, 1,123 samples). When this AD classifier was tested on brain images from unseen mild cognitive impairment (MCI) patients, MCI patients who converted to AD were 3 times more likely to be predicted as AD than MCI patients who did not convert (65.2% vs 20.6%). Predicted scores from the AD classifier showed significant correlations with illness severity. In sum, the proposed AD classifier offers a medical-grade marker that has potential to be integrated into AD diagnostic practice.

Keywords

Alzheimer's disease, convolutional neural network, magnetic resonance brain imaging, sex differences, transfer learning

1. Introduction

Magnetic resonance imaging (MRI) is widely used in neuroradiology to detect brain lesions including stroke, vascular disease, and tumor tissue. Still, MRI has been less useful in definitively identifying degenerative diseases including Alzheimer's disease (AD), mainly because signatures of the disease are diffuse within the images and hard to distinguish from normal aging. Machine learning and deep learning methods have been trained on relatively small datasets, but limited training data often leads to poor generalization performance on new datasets not used the train the algorithms. In the current study, we aim to create a practical brain imaging-based AD classifier with high generalization capability via learning/transfer learning on a diverse range of large-scale datasets.

In recently updated AD diagnostic criteria, such as those proposed by International Working Group (IWG-2) Criteria for Alzheimer's Disease Diagnosis and The National Institute on Aging – Alzheimer's Association (NIA-AA) Alzheimer's Diagnostic Framework, markers such as amyloid measures from cerebrospinal fluid (CSF) and amyloid-sensitive positron emission tomography (PET) have been integrated into the diagnosis of AD (1, 2). Diagnostic sensitivity and specificity have been greatly improved by these markers (1, 3). Even so, the invasive nature and lower availability of these markers limits their application in routine clinical settings. Thus, accurate diagnosis of AD and its early stage using a non-invasive and widely available technology is critically important. Structural MRI is a more promising candidate for imaging-based auxiliary diagnosis for AD considering its non-invasive nature and wider availability than PET. In addition, well-developed MRI data preprocessing pipelines make it feasible to integrate MRI markers into automatic end-to-end deep learning algorithms. Deep learning has already been successfully deployed in real-world scenarios such as extreme weather condition prediction (4), aftershock pattern prediction (5) and automatic speech recognition (6). In clinical scenarios, convolutional neural networks (CNN) – a widely-used architecture that is well-suited for image-based deep learning – have been successfully used

for objective diagnosis of retinal diseases (7), skin cancer (8), and for breast cancer

screening (9).

However, prior attempts at MRI-based AD diagnosis have yet to attain clinical utility. A major

challenge for brain MRI-based algorithms, especially if they are trained on limited data, is

their failure to generalize. For example, a brain imaging-based classifier may give precise

predictions for testing samples from a specific hospital from which the training dataset came.

However, performance of the classifier declines dramatically when directly applied to

samples from another unknown hospital (10). One critical reason for performance

discrepancy is that brain imaging data vary depending on scanner characteristics such as

scanner vendor, magnetic field strength, head coil hardware, pulse sequence, applied

gradient fields, reconstruction methods, scanning parameters, voxel size, field of view, etc.

Participants also differ in sex, age, race/ethnicity, and education. Robust methods need to

work well on diverse populations. These variations in the scans – and in the populations

studied – make it hard for a brain imaging-based classifier trained on data from a single site

(or a few sites) to generalize to data from unseen sites/scanners. This has prevented brain

imaging-based classifiers from becoming practically useful in clinical settings. Most brain

MRI-based studies either did not include independent validation (11-13) or did not achieve

satisfactory performance in independent validations (14). In fact, reviews of brain

imaging-based AD classifiers suggest that most machine learning methods have been

trained on samples in the hundreds, with only 2 out of 81 studies (15), 0 out of 16 studies (16),

and 6 out of 114 studies (17) (of those included in recent systematic reviews) including

independent dataset validations, raising doubts about the generalizability of the models.

Another bottleneck in developing a practical brain imaging-based classifier involves the

variety and comprehensiveness of training datasets. Directly training AD models on datasets

that only contain several hundred samples may result in overfitting with poor generalization to

unseen test data (15). The transfer learning framework has been proposed to solve this

problem, by training a model on a certain characteristic for which abundant samples are

available, and fine-tuning it to another characteristic, or for similar tasks, in smaller samples

(18). Published evidence shows that pretrained models can outperform models trained from

scratch in classification accuracy and robustness (19, 20). In medical imaging, transfer

learning has been successfully applied to diagnose retinal disease (7) and skin cancer (8).

Nonetheless, in brain imaging, no study has encompassed the tens of thousands of openly

shared brain images to promote the generalizability of an AD classifier. Thus, in the current

study, we used one of the largest and most diverse samples to date (N = 85,721 from more

than 217 sites/scanners, see Table 1) to pre-train a brain imaging-based classifier with high

generalizability. We chose a sex classifier rather than an age predictor as the base model for

transfer learning, because age prediction error may contain biological meaning (e.g.,

increased predicted age may indicate accelerated aging (21)). Thus, it can be hard to

measure the true performance of an age predictor while participant sex is more stable for

classification. Subsequently, the pre-trained sex classifier was fine-tuned for AD classification

and was validated through leave-sites-out cross-validation and three independent validations.

The goal of the present study was to build a practical AD classifier with high generalizability.

We incorporated three design features to improve the method's clinical utility. First, we trained

and tested the algorithm on a dataset of unprecedented size and diversity (from more than

217 sites/scanners). The variety of training samples was critical for improving model

generalizability. Second, rigorous leave-datasets/sites-out cross-validation and independent

validations were implemented to assure that classifier accuracy would be robust to

site/scanner variability. Third, compared to 2D modules (feature detectors) typically used in

CNNs for natural images, here, fully 3D convolution filters were used to capture more

sophisticated and distributed spatial features for diagnostic classification. We also openly

share our preprocessed data, trained model, code, and have built an online predicting

148 website for anyone interested in testing our classifier.

2. Methods

Data acquisition. We submitted data access applications to nearly all the open-access brain imaging data archives and received permissions from the administrators of 34 datasets. The full dataset list is shown in Table 1. Deidentified data were contributed from datasets collected with approvals from local Institutional Review Boards. The reanalysis of these data was approved by the Institutional Review Board of Institute of Psychology, Chinese Academy of Sciences. All participants had provided written informed consent at their local institution. All 50,876 participants (contributing 85,721 samples) had at least one session with a T1-weighted structural brain image and information on their sex and age. For participants with multiple sessions of structural images, each image was considered an independent sample for data augmentation in training. Importantly, scans from the same person were never split into training and testing sets, as that could artifactually inflate performance.

Table 1: **Datasets used in the present study**

Full Name of Dataset	Number of T1 Scans	T1 Scans after QC	Age (mean±std)	Number of Subjects	Number of Sites	Manufacturers	Field Strength
Adolescent Brain Cognition Development	31176	30222	13.76±10.08	11875	21	SIE/PHI/GE	3T
UK Biobank	20124	19744	63.1±7.46	20124	4	SIE	3T
Alzheimer's Disease Neuroimaging Initiative	16596	16431	74.97±7.4	2546	57	SIE/PHI/GE	3T/1.5T
Open Access Series of Imaging Studies	3150	3099	67.54±20.64	1664	(5)	SIE	3T/1.5T
REST-meta-MDD sample	2380	2363	36.2±15.11	2380	17	SIE/PHI/GE	3T/1.5T
Brain Genomics Superstruct Project	1570	1552	21.54±2.89	1570	2	SIE	3T
Human Connectome Project	1267	1220	/	1267	1	SIE	7T/3T
Autism Brain Imaging Data Exchange	1102	1073	17.09±8.06	1102	17	SIE/PHI/GE	3T/1.5T
Autism Brain Imaging Data Exchange II	1043	1019	15.16±9.39	1043	19	SIE/PHI/GE	3T/1.5T
1000 Functional Connectomes Project	897	864	25.8±10.76	897	33	SIE/PHI/GE	3T/1.5T
ADHD-200 Sample	876	864	12.35±3.28	876	8	SIE/PHI	3T/1.5T
Consortium for Reliability and Reproducibility	714	691	23.45±12.31	715	2	SIE/GE	3T
Cambridge Centre for Ageing Neuroscience (Cam-CAN)	652	523	54.36±18.55	652	1	SIE	3T
Enhanced Nathan Kline Institute - Rockland Sample	646	616	38.63±21.21	646	1	SIE	3T
Southwest University Longitudinal Imaging Multimodal	586	581	20.1±1.3	586	1	SIE	3T
Child Mind Institute Healthy Brain Network	572	506	10.74±3.65	572	3	SIE	3T/1.5T
Establishing Moderators and Biosignatures of Antidepressant	540	523	/	540	4	SIE/PHI/GE	3T
Southwest University Adult Lifespan Dataset	493	483	45.16±17.45	493	1	SIE	3T
Max Planck Institute Leipzig Mind-Brain-Body Dataset	316	291	/	316	1	SIE	3T
Beijing Enhanced Sample	180	176	21.22±1.94	180	1	SIE	3T
Nathan Kline Institute - Rockland Sample	167	151	35.59±20.71	167	1	SIE	3T
The Center for Biomedical Research Excellence	147	137	/	147	1	SIE	3T

The Age-ility Project	110	105	21.87±5.39	110	1	SIE	3T
Parkinson's Disease Datasets	68	65	66.18±7.58	68	2	SIE	3T/1.5T
Power et al., 2012 Neuroimage Sample	63	62	14.25±6.05	63	1 (2)	SIE	3T
NYU Institute for Pediatric Neuroscience	47	45	30.4±8.98	47	1	SIE	3T
Beijing Eyes Open Eyes Closed Sample	46	44	22.54±2.18	46	1	SIE	3T
Multi-Modal MRI Reproducibility Resource	42	42	31.76±9.35	42	1	PHI	3T
Adelstein et al., 2011, PLoS ONE Sample	39	36	29.59±8.38	39	1	SIE	3T
Cleveland CCF	31	29	43.55±11.14	31	1	SIE	3T
Virginia Tech Carilion Research Institute	25	24	26.84±8.17	25	1 (3)	SIE	3T
Beijing Short TR Sample	24	23	23.71±6.74	24	1	SIE	3T
FIND Lab sample	13	12	24.08±3.73	13	1	GE	3T
The Midnight Scan Club dataset	10	10	29.1±3.35	10	1	SIE	3T
	85712	83735		50876	217		

Note: Abbreviation: SIE = Siemens, Phi = Philips, GE = General Electric. Numbers in the bracket indicate the number of scanners.

MRI preprocessing. We did not feed raw data into the classifier for training but used accepted pre-processing pipelines that are known to generate useful features from brain scans. The brain structural data were segmented and normalized to acquire grey matter density (GMD) and grey matter volume (GMV) maps. Specifically, we used the voxel-based morphometry (VBM) analysis module within Data Processing Assistant for Resting-State fMRI (DPARSF) (22), which is based on SPM (23), to segment individual T1-weighted images into grey matter, white matter, and cerebrospinal fluid (CSF). Then, the segmented images were transformed from individual native space to MNI-152 space (a coordinate system created by Montreal Neurological Institute (24)) using the Diffeomorphic Anatomical Registration Through Exponentiated Lie algebra (DARTEL) tool (25). Two voxel-based structural metrics, GMD and GMV were fed into the deep learning classifier as two features for each participant. GMD is the output of the unmodulated tissue segmentation map in MNI space. GMV is calculated by multiplying the voxel value in GMD by the Jacobian determinants derived from the spatial normalization step (modulated) (26). Medical imaging-based classifiers could reach better or similar classification performances using an enhancing preprocessing procedure (27, 28).

Quality control. Poor quality raw structural images would produce distorted GMD and GMV maps during segmentation and normalization. To remove such participants from affecting the training of the classifiers, we excluded participants in each dataset with a spatial correlation exceeding the threshold defined by (mean - 2SD) of the Pearson's correlation between each participant's GMV map and the grand mean GMV template. The grand mean GMV template was generated by randomly selecting 10 participants from each dataset (image quality visually checked for each participant) and averaging the GMV maps of all these 340 (from 34 datasets) participants. After quality control, 83,735 samples were retained for classifier training (Figure S1).

Deep learning: classifier training and testing for sex. As the feature maps of brain MRI

were three-dimensional (3D) rather than two-dimensional (2D), we could not directly use 2D

pretrained models such as models trained based on ImageNet. Meanwhile, few pretrained

3D CNN models based on large-scale datasets, especially brain MRI datasets, exist.

Therefore, we had to pretrain a brain MRI-based model for the further transfer learning

procedure. It could have been more efficient to pretrain the model on another

neurodegenerative disorder such as Parkinson's disease (29, 30). However, we were unable

to locate other neurodegenerative disorder datasets with tens of thousands of samples.

Hence, we chose the sex classification task to pretrain the model because sex is a commonly

available phenotype for any kind of datasets. We trained a 3-dimensional

Inception-ResNet-v2(31) model adopted from its 2-dimensional version in the Keras built-in

application (see Figure 1A for its structure). This is a state-of-the-art pattern recognition

model, which integrates two classical series of CNN models, Inception and ResNet. We

replaced the convolution, pooling, and normalization modules with their 3-dimensional

versions and adjusted the number of layers and convolutional kernels to make them suitable

for 3-dimensional MRI inputs (e.g., GMD and GMV as different input channels). The present

model consists of one stem module, three groups of convolutional modules

(Inception-ResNet-A/B/C) and two reduction modules (Reduction-A/B). The model can take

advantage of convolutional kernels with different shapes and sizes, and can extract features

of different sizes. The model also can mitigate vanishing gradients and exploding gradients

by adding residual modules. We utilized the Keras built-in stochastic gradient descent

optimizer with learning rate = 0.01, Nesterov momentum = 0.9, decay = 0.003 (e.g., learn rate

= learn rate$_0$ × (1 / (1 + decay × batch))). The loss function was set to binary cross-entropy.

The batch size was set to 24 and the training procedure lasted 10 epochs for each fold. To

avoid potential overfitting, we randomly split 600 samples out of the training sample as a

validation sample and set a checking point at the end of every epoch. We saved the model in

which the epoch classifier showed the lowest validation loss. Thereafter, the testing sample

was fed into this model to test the classifier.

Figure 1: **Flow diagram for the Alzheimer disease (AD) transfer learning framework and**

cross-validation procedure

(A) Schema for the 3D Inception-ResNet-V2 model and the transfer learning framework for

the Alzheimer disease classifier. (B) Schematic diagram for the leave-datasets-out 5-fold

cross-validation for the sex classifier. (C) Schematic diagram for the leave-sites-out 5-fold

cross-validation for the AD classifier.

While training the sex classifier, random cross-validation may share participants from the

same sites between training and testing samples, so the model may not generalize well to

datasets from unseen sites due to site information leakage during training. To ensure

generalizability, we used cross-dataset validation. In the testing phase, all the data from a

given dataset would never be seen during the classifier training phase. This also ensured the

data from a given site (and thus a given scanner) were unseen by the classifier during

training (see Figure1B for an illustration). This strict setting can limit classifier performance,

but it makes it feasible to generalize to any participant at any site (scanner). Five-fold

cross-dataset validation was used to assess classifier accuracy. Of note, 3 datasets were

always kept in the training sample due to the massive number of samples: Adolescent Brain

Cognition Development (ABCD) (n = 31,176), UK Biobank (n = 20,124), and the Alzheimer's

Disease Neuroimaging Initiative (ADNI) (n = 16,596). The remaining 31 datasets were

randomly allocated to training and testing samples. The allocating schemas were the solution

that balanced the sample size of 5 folds the best from 10,000 random allocating procedures.

Both healthy normal control and brain-related disorder patient samples in the 34 datasets

were used to train the sex classifier.

Transfer learning: classifier training and testing for AD. After obtaining a highly robust

and accurate brain imaging-based sex classifier as a base model, we used transfer learning

to further fine-tune the AD classifier. Rather than retaining the intact sophisticated structure of

the base model (Inception-ResNet-V2), we only leveraged the pre-trained weights in the stem

module and simplified the upper layers (e.g., replacing Inception-ResNet modules with

ordinary convolutional layers). The retained bottom structure of the model works as a feature

extractor and can take advantage of the massive training of the sex classifier. The pruned

upper structure of the AD model can avoid potential overfitting and promote generalizability

by reducing the number of parameters (10 million parameters for the AD classifier vs. 54

million parameters for the sex classifier). This derived AD classifier was fine-tuned on the

ADNI dataset (2,186 samples from 380 AD patients and 4,671 samples from 698 normal

controls (NCs), 76 ± 7 years, 3,493 samples from women). ADNI was launched in 2003

(Principal Investigator: Michael W. Weiner, MD) to investigate biological markers of the

progression of MCI and early AD (see www.adni-info.org). We used the Keras built-in

stochastic gradient descent optimizer with learning rate = 0.0003, Nesterov momentum = 0.9,

decay = 0.002. The loss function was set to binary cross-entropy. The batch size was set to

24 and the training procedure lasted 10 epochs for each fold. Like the cross-dataset

validation for sex classifier training, five-fold cross-site validation was used to assess

classifier accuracy (see Figure 1C for an illustration). By ensuring that the data from a given

site (and thus a given scanner) were unseen by the classifier during training, this strict

strategy made the classifier generalizable with non-inflated accuracy, thus better simulating

realistic clinical applications than traditional five-fold cross-validation. Other than using

GMD+GMV as the input in transfer learning, we also used GMD, GMV or z-standardized

normalized raw T1-weighted images as the input for the sex/AD classifiers to verify the

influence of input format (Table 2). We also trained an age prediction model instead of the sex

classifier in transfer learning to verify the influence of the base-model. We used the same

structure as the sex classifier, except for adding a fully-connected layer with 128 neurons with

"ELU" activation function before the final layer; we also changed the dropout rate from 0.5 to

0.2 following the parameters in the brain age prediction model reported by Jonsson et al. (21).

Finally, we compared the performance of Inception-ResNet-V2 structure with the

performances of some light-weight structures such as VGG19, DenseNet-201 and

MobileNet-V2. The performances of these structures are listed in Table S1. In sum, the

performance of models with light-weight structures were lower than that of the

Inception-ResNet-V2 model, so that we chose Inception-ResNet-V2 for the model structure.

Furthermore, to test the generalizability of the AD classifier, we directly tested it on three

unseen independent AD samples, i.e., the Australian Imaging, Marker and Lifestyle Flagship

Study of Ageing (AIBL) (32), the Minimal Interval Resonance Imaging in Alzheimer's Disease

cohort (MIRIAD) (33), and the Open Access Series of Imaging Studies (OASIS) (34). We

averaged the sigmoid activation output scores of the 5 AD classifiers in five-fold

cross-validation on ADNI to obtain the final classification for each sample. We used

diagnoses provided by the qualified physicians for the AIBL and MIRIAD datasets as the

sample labels (115 samples from 82 AD patients and 554 samples from 324 NCs in AIBL, 74

± 7 years, 374 samples from women; 409 samples from 46 AD patients and 235 samples

from 23 NCs in MIRIAD, 70 ± 7 years, 358 samples from women). As OASIS did not specify

the criteria for an AD diagnosis, we adopted criteria of MMSE and clinical dementia rating

(CDR) modified from the ADNI-1 protocol manual to define AD and NC samples. Specifically,

criteria for AD are (1) MMSE ≤ 22 and (2) CDR ≥ 1.0, and criteria for NC are (1) MMSE > 26

and (2) CDR = 0. Thus, we tested the model on 137 samples from 34 AD patients and 986

samples from 213 NC participants in the OASIS dataset after quality control, age 75 ± 10

years, 772 samples from women. Of note, the scanning conditions and recruitment criteria of

these independent datasets differed much more than variations among different ADNI sites

(where scanning and recruitment was deliberately coordinated), so we expected the AD

classifier to achieve lower performance. We created heterogeneous distributions by randomly

selecting 50% samples in each independent testing datasets 1,000 times to validate the

stability of the model. The 95% confidence intervals of the classification performance metrics

were produced from the random selection procedure.

We further investigated whether the AD classifier could predict disease progression in people

with mild cognitive impairment (MCI). MCI is a syndrome defined as relative cognitive decline

without symptoms interfering with daily life; even so, more than half of MCI patients progress

to dementia within 5 years (35). The stable MCI (sMCI) samples were defined as "scans from

an individual who was once diagnosed as MCI in any phase of ADNI and has not progressed

to AD by the end of the ADNI follow-up", and the progressive MCI (pMCI) samples were

defined as "scans from a participant who was once diagnosed as MCI in any phase of ADNI

and who has progressed to AD". The scans labeled as "conversion" or "AD" (after conversion)

for pMCI and the last scan for sMCI were excluded in the present study for precision. We

screened imaging records of the MCI patients who converted to AD later in the ADNI 1/2/'GO'

phases, and collected 2,371 images from 243 participants labeled as 'pMCI'. We also

assembled 4,018 samples from 524 participants labeled 'sMCI' without later progression for

contrast. We directly fed all these MCI images into the AD classifier without further fine-tuning,

thus evaluating the performance of the AD classifier on unseen MCI information.

Interpretation of the deep learning classifiers.

To better understand the brain imaging-based deep learning classifier, we calculated

occlusion maps for the classifiers. We repeatedly tested the images in the testing sample

using the model with the highest accuracy within the 5 folds, while successively masking

brain areas (volume = 18mm*18mm*18mm, step = 9mm) of all input images. The accuracy

achieved on "intact" samples by the classifier minus accuracy achieved on "defective"

samples indicated the "importance" of the occluded brain area for the classifier. The occlusion

maps were calculated for both sex and AD classifiers. To investigate the clinical significance

of the output of the AD classifier, we calculated the Spearman's correlation coefficient

between the predicted scores and MMSE scores of AD, NC, and MCI samples. We also used

general linear models (GLM) to verify whether the predicted scores (or MMSE score) showed

a group difference between people with sMCI and pMCI. The age and sex information of MCI

participants was included in this GLM as covariates. We selected the T1-weighted images

from the first visit for each MCI subject and finally collected data from 243 pMCI patients and

524 sMCI patients.

3. Results

3.1. Large-Scale Brain imaging Data

Only brain imaging data with enough size and variety can make deep learning accurate and robust enough to build a practical classifier. We received permissions from the administrators of 34 datasets (85,721 samples of 50,876 participants from more than 217 sites/scanners, see Table 1; some datasets did not require applications). After quality control, all these samples were used to pre-train the stem module to achieve better generalization for further AD classifier training. The T1-weighted images were collected through Magnetization-Prepared Rapid Gradient-Echo Imaging (MPRAGE) or Inversion Recovery Fast Spoiled Gradient Recalled Echo (IR-FSPGR) sequences of 1.5 tesla or 3 tesla MR scanners. The raw acquisition voxel sizes ranged from 0.7mm×0.7mm×0.7mm to 1.3mm×1.3mm×1.2mm. For further fine-tuning of the AD classifier, ADNI, AIBL, MIRIAD, and OASIS were selected to train and test the model.

3.2. Performance of the sex classifier

We trained a 3-dimensional Inception-ResNet-v2 model adapted from its 2-dimensional version in the Keras built-in application (see Figure 1A for structure). As noted in Methods, we did not feed raw data into the classifier for training, but used prior knowledge regarding helpful analytic pipelines. Grey matter density (GMD) and grey matter volume (GMV) maps were treated as different input channels for models. To ensure generalizability, five-fold cross-dataset validation was used to assess classifier accuracy. The five-fold cross-dataset validation accuracies were: 94.8%, 94.0%, 94.8%, 95.7%, and 95.8%. Taken together, accuracy was 94.9% in testing samples when pooling results across the five folds. The area under the curve (AUC) of the receiver operating characteristic (ROC) curve was the classifier performance index at various threshold settings (36, 37). The AUC of the sex classifier reached 0.981 (Figure 2). In short, our model can classify the sex of a participant based on brain structural imaging data from anyone on any scanner with an accuracy of about 95%.

364   Interested   readers   can   test   this   model   on   our   online   prediction   website

365   (http://brainimagenet.org).

Figure 2: Performance of the sex classifier

(A) The receiver operating characteristic curve of the sex classifier. (B) The tensorboard

monitor graph of the sex classifier in the training sample. The curve was smoothed for better

visualization. (C) The tensorboard monitor graph of the sex classifier in the validation sample.

3.3. Performance of the AD classifier

After creating a practical brain imaging-based classifier for sex with high cross-dataset accuracy, we used transfer learning to see if we could classify patients with AD. The AD classifier achieved an accuracy of 90.9% (accuracy = 93.2%, 90.3%, 92.0%, 94.4%, and 86.7% in 5 cross-site folds) in the test samples. The 95% confidence interval of accuracy was [90.2%, 91.5%]. Average sensitivity and specificity were 0.838 and 0.942, respectively. The 95% confidence intervals of sensitivity and specificity were [0.824, 0.854] and [0.935, 0.948], respectively. The ROC AUC reached 0.963 when results from the 5 testing samples were taken together (see Figure 3 and Table 2). The 95% confidence interval of ROC AUC was [0.958, 0.967]. The AD classifier achieved an average accuracy of 91.4% on 3T field strength MR testing samples and achieved an average accuracy of 91.1% on 1.5T MR testing samples. The accuracy in 3T MR testing sample did not differ significantly from that of 1.5T MR testing sample ($p = 0.316$, by permutation test of randomly allocating the testing samples into 1.5T or 3T groups and calculating the accuracy difference between the two groups 100,000 times, Figure S2). In sum, the AUCs of models taking other types of images as the input (e.g., raw T1-weighted images) were slightly lower than that of the GMD+GMV image-based model. The GMD and GMV maps contained a priori knowledge from brain science, which might partly explain the better performance of GMD+GMV derived models.

Figure 3: **Performance of the Alzheimer's disease (AD) classifier**

Left panel shows the training and testing performance of the AD classifier on ADNI sample.

Right panel shows the testing performance of the AD classifier on independent samples. (A)

The receiver operating characteristic curve of the AD classifier. (B) The tensorboard monitor

panel of the AD classifier in the training sample. (C) The tensorboard monitor panel of the AD

classifier in the validation sample. (D) The ROC curve of AD classifier tested on the AIBL

sample. (E) The ROC curve of the AD classifier tested on the MIRIAD sample. (F) The ROC

curve of the AD classifier tested on the OASIS sample. (G) The ROC curve of the AD

classifier tested on the ADNI MCI sample.

To test the generalizability of the AD classifier, we applied it to unseen independent AD

datasets, i.e., AIBL, MIRIAD, and OASIS. The AD classifier achieved 94.5% accuracy in AIBL

with 0.966 AUC (Figure 3D). Sensitivity and specificity were 0.881 and 0.958, respectively.

The AD classifier achieved 93.6% accuracy in MIRIAD with 0.994 AUC (Figure 3E).

Sensitivity and specificity were 0.897 and 1.000, respectively. The AD classifier achieved

91.1% accuracy in OASIS with 0.976 AUC (Figure 3F). Sensitivity and specificity were 0.932

and 0.908, respectively.

Table 2: **Performance of the Alzheimer's disease classifier**

Input Feature	Accuracy				AUC				Sensitivity				Specificity			
	ADNI	AIBL	MIRIAD	OASIS	ADNI	AIBL	MIRIAD	OASIS	ADNI	AIBL	MIRIAD	OASIS	ADNI	AIBL	MIRIAD	OASIS
Sex classifier as base model																
GMV+GMD	0.909	0.945	0.936	0.911	0.963	0.966	0.994	0.976	0.838	0.881	0.897	0.932	0.942	0.958	1.000	0.908
	[0.902, 0.915]	[0.926, 0.965]	[0.917, 0.955]	[0.895, 0.928]	[0.958, 0.967]	[0.945, 0.987]	[0.991, 0.997]	[0.966, 0.985]	[0.824, 0.854]	[0.816, 0.943]	[0.869, 0.928]	[0.889, 0.972]	[0.935, 0.948]	[0.941, 0.974]	[1.000, 1.000]	[0.890, 0.926]
GMD	0.879	0.941	0.944	0.906	0.940	0.952	0.990	0.946	0.762	0.803	0.959	0.793	0.933	0.968	0.920	0.922
	[0.872, 0.887]	[0.923, 0.958]	[0.926, 0.962]	[0.889, 0.924]	[0.935, 0.946]	[0.926, 0.978]	[0.985, 0.995]	[0.927, 0.964]	[0.744, 0.782]	[0.732, 0.875]	[0.939, 0.980]	[0.722, 0.862]	[0.926, 0.941]	[0.953, 0.981]	[0.886, 0.955]	[0.905, 0.940]
GMV	0.903	0.943	0.916	0.862	0.960	0.964	0.991	0.949	0.829	0.857	0.894	0.866	0.937	0.960	0.953	0.861
	[0.895, 0.910]	[0.926, 0.962]	[0.894, 0.936]	[0.841, 0.880]	[0.955, 0.964]	[0.940, 0.986]	[0.986, 0.995]	[0.932, 0.965]	[0.812, 0.844]	[0.783, 0.923]	[0.865, 0.922]	[0.806, 0.924]	[0.930, 0.944]	[0.943, 0.979]	[0.927, 0.983]	[0.840, 0.882]
T1-weighted	0.866	0.947	0.937	0.730	0.928	0.961	0.991	0.940	0.766	0.820	0.935	0.918	0.912	0.971	0.940	0.704
	[0.857, 0.874]	[0.929, 0.965]	[0.917, 0.955]	[0.701, 0.757]	[0.922, 0.934]	[0.939, 0.983]	[0.987, 0.995]	[0.920, 0.963]	[0.748, 0.785]	[0.745, 0.893]	[0.911, 0.959]	[0.871, 0.969]	[0.904, 0.920]	[0.958, 0.985]	[0.911, 0.972]	[0.674, 0.735]
Age predictor as base model																
GMV+GMD	0.901	0.950	0.942	0.892	0.957	0.965	0.994	0.965	0.812	0.881	0.907	0.866	0.943	0.964	1.000	0.896
	[0.894, 0.908]	[0.933, 0.968]	[0.923, 0.958]	[0.875, 0.909]	[0.953, 0.962]	[0.941, 0.988]	[0.991, 0.997]	[0.952, 0.979]	[0.795, 0.829]	[0.815, 0.946]	[0.879, 0.935]	[0.806, 0.928]	[0.936, 0.950]	[0.947, 0.980]	[1.000, 1.000]	[0.877, 0.914]

The ADNI dataset contained 2,186 AD samples and 4,671 NC samples. The AIBL dataset contained 115 AD samples and 554 NC samples. The MIRIAD dataset contained 409 AD samples and 235 NC samples. The OASIS dataset contained 137 AD samples and 986 NC samples. The sample sizes shown here are the numbers of T1-weighted brain MRI scans. Abbreviations: AD = Alzheimer's disease participants, NC = normal control participants, GMD = grey matter density map, GMV = grey matter volume map, T1-weighted = normalized 3D T1-weighted structural image, AUC = area under the curve. The 95% confidence intervals are listed in brackets.

Importantly, although the AD classifier had never "seen" brain imaging data from subjects

with MCI, we directly tested it on the MCI dataset in ADNI to see if it would have the potential

to predict the progression of MCI to AD. We reasoned that even though people with MCI do

not yet have AD, their scans may appear closer to the AD class learned by the deep learning

model. We found that the AD classifier predicted 65.2% of pMCI patients as being in the AD

class but only 20.4% of sMCI patients were predicted as having AD (Figure 3F). If the

percentage of pMCI patients who were predicted as AD was considered as the sensitivity and

the percentage of sMCI patients who were predicted as AD was considered as 1-specificity,

the AUC of the ROC curve of the AD classifier reached 0.82. These results suggest that the

classifier is practical for screening MCI patients to determine risk of progression to AD. In

sum, we believe our AD classifier can provide important insights relevant to computer-aided

diagnosis and prediction of AD, and we have freely provided it on the website

http://brainimagenet.org. Importantly, classification results by the online classifier should be

interpreted with caution, as they are probabilistic and cannot replace diagnosis by licensed

clinicians.

As a supplementary analysis, we also tested transfer learning of the AD classifier using the

intact structure of the base model (Figure S3). The performance of the model was uniformly

somewhat inferior to the optimized AD classifier. The "intact" AD classifier achieved an

average accuracy of 88.4% with 0.938 AUC in the ADNI test samples (Figure S4). Average

sensitivity and specificity were 0.814 and 0.917, respectively. When tested on independent

samples, the AD classifier achieved 91.2% accuracy in AIBL with 0.948 AUC. Sensitivity and

specificity were 0.851 and 0.924, respectively. The AD classifier achieved 93.9% accuracy in

MIRIAD with 0.995 AUC. Sensitivity and specificity were 0.905 and 0.996, respectively

(Figure S5). The AD classifier achieved 86.1% accuracy in OASIS with 0.921 AUC.

Sensitivity and specificity were 0.789 and 0.881, respectively. When tested on MCI samples,

63.2% of pMCI patients were predicted as having AD and 22.1% of sMCI patients were

predicted as having AD by the AD classifier.

3.4. Interpretation of the deep learning classifiers

To better understand the brain imaging-based deep learning classifier, we calculated

occlusion maps for the classifiers. The occlusion map showed that hypothalamus, superior

vermis, pituitary, thalamus, amygdala, putamen, accumbens, hippocampus, and

parahippocampal gyrus played critical roles in predicting sex (Figure 4A). The occlusion map

for the AD classifier highlighted that the hippocampus and parahippocampal gyrus -

especially in the left hemisphere - played unique roles in predicting AD (Figure 4B, Figure S6).

Another visualization technique, Gradient-weighted Class Activation Mapping (Grad-CAM)

(38), also showed relatively high weights of these regions in the input feature maps (Figure

S9).

Figure 4: **Interpretation of the deep learning classifiers with occlusion maps**

Classifier performance dropped considerably when the brain areas rendered in red were

masked out of the model input. (A) Occlusion maps for the sex classifier. Hypothalamus and

pituitary were marked in dashed line and solid line. (B) Occlusion maps for the Alzheimer

disease classifier.

To investigate the clinical significance of the output of the AD classifier, we calculated the

Spearman's correlation coefficient between the scores predicted by the classifier and MMSE

scores in AD, NC, and MCI samples, although the classifier had not been trained with MMSE

scores. This analysis confirmed significant negative correlations between the predicted

scores and MMSE scores for AD (r = -0.37, $p < 1 \times 10^{-55}$), NC (r = -0.11, $p < 1 \times 10^{-11}$), MCI (r

= -0.52, $p < 1 \times 10^{-307}$), and the overall samples (r = -0.64, $p < 1 \times 10^{-307}$) (Figure 5, Figure S7).

As lower MMSE scores indicate more severe cognitive impairment in AD and MCI patients,

we confirmed that the more severe the disease, the higher the classifier's predicted score. In

addition, both the predicted scores and MMSE scores differed significantly between pMCI

and sMCI (predicted scores: t = 13.88, $p < 0.001$, Cohen's d = 1.08; MMSE scores: t = -9.42,

$p < 0.01$, Cohen's d = -0.73, Figure S8). Importantly, the effect sizes of the classifier's

predicted scores were much larger than those for the behavioral measure (MMSE scores).

Figure 5: **Correlations between the output of the Alzheimer's disease (AD) classifier**

and severity of illness

The predicted scores from the AD classifier showed significant negative correlations with the

mini-mental state examination (MMSE) scores of AD, normal control (NC) and mild cognitive

impairment (MCI) samples. (A) Correlation between the predicted scores from the AD

classifier and MMSE scores of AD samples. (B) Correlation between the predicted scores

from the AD classifier and MMSE scores of MCI samples. (C) Correlation between the

predicted scores from the AD classifier and MMSE scores of NC samples. (D) Correlation

between the predicted scores from the AD classifier and the MMSE scores of AD, NC, and

MCI samples, combined.

4. Discussion

Using an unprecedentedly diverse brain imaging sample, we pre-trained a sex classifier with about 95% accuracy which served as a base-model for transfer learning to promote model generalizability. After transfer learning, the model fine-tuned to AD achieved 90.9% accuracy in stringent leave-sites-out cross-validation and 94.2%/93.6%/91.1% accuracy for direct tests on three unseen independent datasets. Predicted scores from the AD classifier correlated significantly with illness severity. The AD classifier also showed the potential to predict the prognosis of MCI patients.

The high accuracy and generalizability of our deep neural network classifiers demonstrate that brain imaging has the practical potential to be auxiliary to the diagnostic process. One of the most prominent advantages of the present protocol is its outstanding generalizability, as validated by leave-sites-out validations and three independent-dataset validations. Performance of the AD classifier remained consistent despite considerable scanner/participant variations across four datasets used in the present study (e.g., ADNI, AIBL, MIRIAD and OASIS). Specifically, accuracies always exceeded 90% and AUCs always exceeded 0.96 in all four datasets. The present model outperformed models in recent studies whose accuracies range from 72.3% to 95% (39) or from 77% to 87% (14) using the same independent datasets (e.g., AIBL, MIRIAD, and OASIS). In addition, the analogous accuracies achieved on 1.5T and 3T MR ADNI imaging data further supported the robustness of the present classifier.

Of note, the output of the deep neural network model is a continuous variable, so the threshold can be adjusted to change sensitivity and specificity for certain purposes. For example, when tested on the AIBL dataset, sensitivity and specificity results were 0.881 and 0.958, respectively, as the default threshold was set at 0.5. However, for screening, the false-negative rate should be minimized even at the cost of higher false-positive rates. If we lower the threshold (e.g., to 0.2), sensitivity can be improved to 0.921 at a cost of decreasing

specificity to 0.885. Thus, in our freely available AD prediction website, users can obtain

continuous outputs and adjust the threshold to suit their specific purposes. Here, the

sensitivity-specificity tradeoff of the AD classifier was consistent across different testing sites,

so that physicians in diverse clinical settings can have consistent expectations for the

classification tendencies of the classifier.

Beyond the feasibility of being integrated into the diagnostic process, the present AD model

also showed potential to predict the progression of MCI patients. First, the present model was

able to quantify key disease milestones by predicting disease progression in MCI patients. To

wit, people with pMCI were 3 times more likely to be classified as AD than sMCI (65.2% vs

20.4%). Recently, a review on predicting progression from MCI noted that about 40% of

studies had methodological issues, such as lack of a test dataset, data-leakage in feature

selection or parameter tuning, and leave-one-out validation performance bias (40). The

present AD classifier was only trained on AD/NC samples and was not fine-tuned using MCI

data, so data leakage was avoided. The estimated true AUC of current published state-of-art

classifiers for predicting progression of MCI is about 0.75 (15, 40). The proposed AD

classifier here outperformed that benchmark (AUC = 0.82). Considering the discouraging

clinical trial failures of AD treatments, early identification of people with MCI with high

potential to progress to AD would help in the evaluation of early treatments (41).

Although deep-learning algorithms have often been referred to as "black boxes" for their poor

interpretability, our subsequent analyses showed that the current MRI-based AD marker was

aligned with pathological findings and clinical insights. For example, AD induced brain

structural changes have been frequently reported by MRI studies. Among all the structural

findings, hippocampal atrophy is the most prominent change and is used in imaging assisted

diagnosis (42, 43). Neurobiological changes in the hippocampus typically precede

progressive neocortical damage and AD symptoms. The convergence of our deep learning

system and human physicians on alterations in hippocampal structure for classifying AD

patients is in line with the crucial role of the hippocampus in AD. On the other hand, its

maximum absolute value was only about 3.1%, which means that even if the most important

brain area was eliminated from input, accuracy only dropped from 90.9% to about 88.8%.

Interestingly, brain atrophy in AD has been frequently reported to be left lateralized (44, 45).

Compared to the un-optimized AD classifier, a slight left hemisphere preference for input

features may help explain the improved performance of the optimized AD classifier.

Rather than indiscriminately imitating the structure of the base model in transfer learning, the

present AD classifier significantly simplified the model before the fine-tuning procedure. To wit,

the performance of the unoptimized AD classifier was far poorer than that of the optimized AD

classifier in accuracy, sensitivity, specificity, and in independent validation performance.

Truncating or pruning models before transfer learning has been found to facilitate the

performance of the transferred models (46, 47). As the sample for training the AD classifier is

considerably smaller than that used to train the sex classifier, the simplified model structure

may have helped to avoid overfitting and improve generalizability.

By precisely predicting sex, the present study provides evidence of sex differences in human

brain. Daphna and colleagues extracted hundreds of VBM features from structural MRI and

concluded that "the so-called male/female brain" does not exist as no individual structural

feature supports a sexually dimorphic view of human brains (48), which was supported by a

recent large-scale review (49). However, human brains may embody sexually dimorphic

features in a multivariate manner. The high accuracy and generalizability of the present brain

image-based sex classifier implies a "brain sex" is kind of recognizable in a

1,981,440-dimension (96*120*86*2) feature space, thus need to be further investigated in the

future. Among those 1,981,440 features, the hypothalamus and pituitary played the most

critical roles in predicting sex. The hypothalamus regulates testosterone secretion through

the hypothalamic-pituitary-gonadal axis and thus plays a critical role in brain masculinization

(50). Men have significantly larger hypothalamus than women relative to cerebrum size (51).

In addition, cerebellum – especially the vermis – strongly contributed to sex classification, in

line with MRI morphology studies of sex differences in cerebellum (52, 53). Taken together,

our machine learning evidence suggests that male/female brain differences do exist, in the

sense that accurate classification is possible.

In the deep learning field, the emergence of ImageNet tremendously accelerated the

evolution of computer vision (54). ImageNet provided large amounts of well-labeled image

data for researchers to pre-train their models. Studies have shown that pre-trained models

can facilitate the performance and robustness of subsequently fine-tuned models (19). The

present study confirms that the "pre-train + fine-tuning" paradigm also works for MRI-based

auxiliary diagnosis. Unfortunately, no such a well-preprocessed dataset exists in the brain

imaging domain. As data organization and preprocessing of MRI data require tremendous

time, manpower and computational load, these constraints impede scientists from other fields

entering brain imaging. Open access to large amounts of preprocessed brain imaging data is

fundamental to facilitate the participation of a broader range of researchers. Beyond building

and sharing a practical brain imaging-based deep learning classifier, we will openly share all

sharable preprocessed data to invite researchers (especially computer scientists) to join the

efforts to create predictive models using brain images (Link_To_Be_Added upon publication;

preprocessed data of some datasets will not be shared as the raw data owners do not allow

sharing of data derivatives). We anticipate that this dataset may boost the clinical utility of

brain imaging as ImageNet has done in computer vision research. We openly share our

models   to   allow   other   researchers   to   deploy   them

(https://github.com/Chaogan-Yan/BrainImageNet). Our code is also openly shared as well,

allowing other researchers to replicate the present results and further develop brain

imaging-based classifiers based on our existing work. Finally, we have also built a

demonstration website for classifying sex and AD (http://brainimagenet.org). Users can

upload raw T1-weighted or preprocessed GMD and GMV data to make predictions of sex or

AD labels in real-time.

Limitations of the current study should be acknowledged. Considering the lower

reproducibility of functional MRI compared to structural MRI, only structural MRI derived

images were used in the present deep learning model. Even so, functional measures of

physiology and activation may further improve the performance of sex and brain disorder

classifiers. In future studies, functional MRI, especially resting-state functional MRI, may

provide additional information for model training. Furthermore, with advances in software

such as FreeSurfer(55), fmriprep(56), and DPABISurf, surface-based algorithms have shown

their superiority when compared with traditional volume-based algorithms (57).

Surface-based algorithms are more time consuming to run in terms of computation load, but

can provide more precise brain registration and reproducibility. Future studies should take

surface-based images as inputs for deep learning models. In addition, the present AD

classification model was built based on labels provided by the ADNI database. Using

post-mortem neuropathological data, the gold standard for AD diagnoses, could further

advance the clinical value of MRI-based markers.

In summary, we pooled MRI data from more than 217 sites/scanners to constitute one of the

largest brain MRI samples to date, with the preprocessed imaging data derivatives openly

shared to the scientific community whenever allowed. The brain imaging-based AD classifier

derived from transfer learning achieved both high rates of accuracy and generalizability,

which were validated by strict cross-sites-validation and independent datasets validation. The

AD classifier was able to predict the progression of MCI patients non-invasively. The present

study demonstrates the feasibility of the transfer learning framework in brain disorder

applications. Future work should examine such a framework to assess psychiatric disorders,

to predict treatment response, and individual differences more broadly.

Declarations

Ethics approval and consent to participate

Deidentified data were contributed from datasets collected with approval by local Institutional

Review Boards. The reanalysis of these data was approved by the Institutional Review Board

of Institute of Psychology, Chinese Academy of Sciences. All participants had provided

written informed consent at their local institutions.

Consent for publication

Not applicable.

Availability of data and materials

The imaging, phenotype and clinical data used for the training, validation and test sets were

obtained by application from the administrators of 34 datasets. The preprocessed brain

imaging data for which raw data owners permit sharing data derivatives will be available on

(Link_To_Be_Added upon publication). The code for training and testing the model are

openly shared at https://github.com/Chaogan-Yan/BrainImageNet. Demonstration website for

classifying sex and AD is available at http://brainimagenet.org.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the Sci-Tech Innovation 2030 - Major Project of Brain Science

and Brain-inspired Intelligence Technology (grant number: 2021ZD0200600), National Key

R&D Program of China (grant number: 2017YFC1309902), the National Natural Science

Foundation of China (grant numbers: 82122035, 81671774, 81630031), the 13th Five-year

Informatization Plan of Chinese Academy of Sciences (grant number: XXH13505), the Key

Research Program of the Chinese Academy of Sciences (grant NO. ZDBS-SSW-JSC006),

Beijing Nova Program of Science and Technology (grant number: Z191100001119104), and

the Scientific Foundation of Institute of Psychology, Chinese Academy of Sciences (grant

number: E2CX4425YZ).

Authors' contributions

C.-G.Y. designed the overall experiment. B.L., H.-X.L., L.L., N.-X.C., Z.-C.Z., H.-X.Z., X.-Y.L.,

Y.-W.W., S.-X.C., Z.-Y.D., Z.F., H.Y. and X.C. applied and preprocessed imaging data.

H.-X.L. and B.L sorted the phenotype information of datasets. B.L. designed the model

architectures and trained the models, B.L., Z.-K.C and C.-G.Y. built the online classifiers.

C.-G.Y. provided technical supports and supervised the project. B.L. and C.-G.Y. wrote the

paper, P.M.T edited the paper and suggested supplementary analysis, F.X.C. edited and

polished the paper. All authors approved the manuscript and had final responsibility for the

decision to submit for publication.

Acknowledgements

Data used in the preparation of this article for training and testing the sex classifier was

obtained from the Adolescent Brain Cognitive Development (ABCD) Study

(https://abcdstudy.org), held in the NIMH Data Archive (NDA). This is a multisite, longitudinal

study designed to recruit more than 10,000 children ages 9-10 and follow them over 10 years

into early adulthood. The ABCD Study is supported by the National Institutes of Health and

additional federal partners under award numbers U01DA041048, U01DA050989,

U01DA051016, U01DA041022, U01DA051018, U01DA051037, U01DA050987,

U01DA041174, U01DA041106, U01DA041117, U01DA041028, U01DA041134,

U01DA050988, U01DA051039, U01DA041156, U01DA041025, U01DA041120,

U01DA051038, U01DA041148, U01DA041093, U01DA041089. A full list of supporters is

available at https://abcdstudy.org/federal-partners.html. A listing of participating sites and a

complete   listing   of   the   study   investigators   can   be   found   at

https://abcdstudy.org/scientists/workgroups/. ABCD consortium investigators designed and

implemented the study and/or provided data but did not necessarily participate in analysis or

writing of this report. This manuscript reflects the views of the authors and may not reflect the

opinions or views of the NIH or ABCD consortium investigators. This research has been

conducted using the UK Biobank Resource. Data collection and sharing for the training and

testing the sex and AD classifiers were funded by the Alzheimer's Disease Neuroimaging

Initiative (ADNI) (National Institutes of Health Grant U01 AG024904) and DOD ADNI

(Department of Defense award number W81XWH-12-2-0012). ADNI is funded by the

National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering,

and through generous contributions from the following: AbbVie; Alzheimer's Association;

Alzheimer's Drug Discovery Foundation; Araclon Biotech; BioClinica, Inc.; Biogen;

Bristol-Myers Squibb Company; CereSpir, Inc.; Cogstate; Eisai Inc.; Elan Pharmaceuticals,

Inc.; Eli Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd and its affiliated company

Genentech, Inc.; Fujirebio; GE Healthcare; IXICO Ltd.; Janssen Alzheimer Immunotherapy

Research   &   Development,   LLC.;   Johnson   &   Johnson   Pharmaceutical   Research   &

Development LLC.; Lumosity; Lundbeck; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.;

NeuroRx Research; Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer

Inc.;   Piramal   Imaging;   Servier;   Takeda   Pharmaceutical   Company;   and   Transition

Therapeutics. The Canadian Institutes of Health Research is providing funds to support ADNI

clinical sites in Canada. Private sector contributions are facilitated by the Foundation for the

National   Institutes   of   Health   (www.fnih.org).   The   grantee   organization   is   the   Northern

California   Institute   for   Research   and   Education,   and   the   study   is   coordinated   by   the

Alzheimer's Therapeutic Research Institute at the University of Southern California. ADNI

data are disseminated by the Laboratory for Neuro Imaging at the University of Southern

California. Data used in the preparation of this article were obtained from the MIRIAD

database. The MIRIAD investigators did not participate in analysis or writing of this report.

The MIRIAD dataset is made available through the support of the UK Alzheimer's Society

(Grant RF116). The original data collection was funded through an unrestricted educational

grant from GlaxoSmithKline (Grant 6GKC).

Authors' information

Chao-Gan YAN, Ph.D.

Dr. Chao-Gan Yan is a professor at the Institute of Psychology, Chinese Academy of

Sciences (IPCAS). He is the Director of the Magnetic Resonance Imaging Research Center,

the Director of the International Big-Data Center for Depression Research, and the Principal

Investigator of The R-fMRI Lab located at IPCAS. His work has been widely cited in the

scientific community (total citations > 16000,

http://scholar.google.com/citations?user=IJQ9B58AAAAJ), achieving an h-index of 38. Six of

his first-author/corresponding author papers ranked as ESI Top 1% Highly Cited Papers, 2 of

which were ESI Top 1‰. He has been ranked as 2019, 2020 and 2021 most cited Chinese

Researchers by Elsevier. He was awarded the 2021 Early Career Investigator Award by The

Organization for Human Brain Mapping (OHBM). Additionally, he currently serves as

associate editor for Neuroimage: Reports and serves on the editorial boards of NeuroImage

and Journal of Neuroscience Methods. His research mainly focuses on resting-state fMRI

(R-fMRI) computational methodology, mechanisms of spontaneous brain activity, and their

applications in depression. He has addressed fundamental methodological issues (e.g.,

impact of head motion (ESI Top 1‰ of highly cited papers), standardization (ESI Top 1% of

highly cited papers) and multiple comparison correction (ESI Top 1% of highly cited papers))

on the study of resting-state functional connectomics. He has also developed data

processing and analysis toolbox for R-fMRI, DPABI (ESI Top 1‰ of highly cited papers) and

DPARSF, which has been cited over 2000 times. He initiated a Chinese consortium for big

data of brain imaging of depression (DIRECT), performed a big data study of depression

neuroimaging (PNAS 2019, ESI Top 1% of highly cited papers), and studied the brain

mechanisms of depression through a longitudinal study of an animal model. He has

published 70+ peer-reviewed articles (30+ as first or corresponding author) in prestigious

journals including PNAS, Molecular Psychiatry, NeuroImage, Human Brain Mapping.

References

1. Dubois B, Feldman HH, Jacova C, Hampel H, Molinuevo JL, Blennow K, et al. Advancing research diagnostic
criteria for Alzheimer's disease: the IWG-2 criteria. The Lancet Neurology. 2014;13(6):614-29.

2. Jack Jr CR, Albert MS, Knopman DS, McKhann GM, Sperling RA, Carrillo MC, et al. Introduction to the
recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic
guidelines for Alzheimer's disease. Alzheimer's & dementia. 2011;7(3):257-62.

3. Rice L, Bisdas S. The diagnostic value of FDG and amyloid PET in Alzheimer's disease-A systematic review.
Eur J Radiol. 2017;94:16-24.

4. Ham Y-G, Kim J-H, Luo J-J. Deep learning for multi-year ENSO forecasts. Nature. 2019;573(7775):568-72.

5. DeVries PMR, Viegas F, Wattenberg M, Meade BJ. Deep learning of aftershock patterns following large
earthquakes. Nature. 2018;560(7720):632-4.

6. Liu W, Wang Z, Liu X, Zeng N, Liu Y, Alsaadi FE. A survey of deep neural network architectures and their
applications. Neurocomputing. 2017;234:11-26.

7. Kermany DS, Goldbaum M, Cai W, Valentim CC, Liang H, Baxter SL, et al. Identifying medical diagnoses and
treatable diseases by image-based deep learning. Cell. 2018;172(5):1122-31.

8. Esteva A, Kuprel B, Novoa RA, Ko J, Swetter SM, Blau HM, et al. Dermatologist-level classification of skin
cancer with deep neural networks. Nature. 2017;542(7639):115-8.

9. McKinney SM, Sieniek M, Godbole V, Godwin J, Antropova N, Ashrafian H, et al. International evaluation
of an AI system for breast cancer screening. Nature. 2020;577(7788):89-94.

10. Cai XL, Xie DJ, Madsen KH, Wang YM, Bogemann SA, Cheung EFC, et al. Generalizability of machine
learning for classification of schizophrenia based on resting-state functional MRI data. Hum Brain Mapp.
2020;41(1):172-84.

11. Suk HI, Lee SW, Shen D, Alzheimer's Disease Neuroimaging I. Hierarchical feature representation and
multimodal fusion with deep learning for AD/MCI diagnosis. NeuroImage. 2014;101:569-82.

12. Bashyam VM, Erus G, Doshi J, Habes M, Nasralah I, Truelove-Hill M, et al. MRI signatures of brain age and
disease over the lifespan based on a deep brain network and 14 468 individuals worldwide. Brain.
2020;143(7):2312-24.

13. Moradi E, Pepe A, Gaser C, Huttunen H, Tohka J, Alzheimer's Disease Neuroimaging I. Machine learning
framework for early MRI-based Alzheimer's conversion prediction in MCI subjects. NeuroImage.

2015;104:398-412.

14.   Qiu S, Joshi PS, Miller MI, Xue C, Zhou X, Karjadi C, et al. Development and validation of an interpretable
deep learning framework for Alzheimer's disease classification. Brain. 2020;143(6):1920-33.

15.   Rathore S, Habes M, Iftikhar MA, Shacklett A, Davatzikos C. A review on neuroimaging-based classification
studies and associated feature extraction methods for Alzheimer's disease and its prodromal stages.
NeuroImage. 2017;155:530-48.

16.   Jo T, Nho K, Saykin AJ. Deep Learning in Alzheimer's Disease: Diagnostic Classification and Prognostic
Prediction Using Neuroimaging Data. Front Aging Neurosci. 2019;11:220.

17.   Ebrahimighahnavieh MA, Luo S, Chiong R. Deep learning to detect Alzheimer's disease from neuroimaging:
A systematic literature review. Comput Methods Programs Biomed. 2020;187:105242.

18.   Yosinski J, Clune J, Bengio Y, Lipson H, editors. How transferable are features in deep neural networks? Adv
Neural Inf Process Syst; 2014.

19.   Hendrycks D, Lee K, Mazeika M. Using pre-training can improve model robustness and uncertainty. arXiv
preprint arXiv:190109960. 2019.

20.   Tajbakhsh N, Shin JY, Gurudu SR, Hurst RT, Kendall CB, Gotway MB, et al. Convolutional neural networks
for medical image analysis: Full training or fine tuning? IEEE Trans Med Imaging. 2016;35(5):1299-312.

21.   Jonsson BA, Bjornsdottir G, Thorgeirsson TE, Ellingsen LM, Walters GB, Gudbjartsson DF, et al. Brain age
prediction using deep learning uncovers associated sequence variants. Nat Commun. 2019;10(1):5409.

22.   Yan CG, Zang YF. DPARSF: A MATLAB Toolbox for "Pipeline" Data Analysis of Resting-State fMRI. Front Syst
Neurosci. 2010;4:13.

23.   Friston KJ, Holmes AP, Worsley KJ, Poline JP, Frith CD, Frackowiak RS. Statistical parametric maps in
functional imaging: a general linear approach. Hum Brain Mapp. 1994;2(4):189-210.

24.   Fonov VS, Evans AC, McKinstry RC, Almli CR, Collins DL. Unbiased nonlinear average age-appropriate brain
templates from birth to adulthood. NeuroImage. 2009;47.

25.   Goto M, Abe O, Aoki S, Hayashi N, Miyati T, Takao H, et al. Diffeomorphic Anatomical Registration Through
Exponentiated Lie Algebra provides reduced effect of scanner for cortex volumetry with atlas-based method in
healthy subjects. Neuroradiology. 2013;55(7):869-75.

26.   Good CD, Johnsrude IS, Ashburner J, Henson RN, Friston KJ, Frackowiak RS. A voxel-based morphometric
study of ageing in 465 normal adult human brains. NeuroImage. 2001;14(1):21-36.

27.   Altan G. DeepOCT: An explainable deep learning architecture to analyze macular edema on OCT images.
Engineering Science and Technology, an International Journal. 2022;34.

28.   Altan SSN, Gokhan. CLAHE based Enhancement to Transfer Learning in COVID-19 Detection.
https://dergiparkorgtr/en/pub/gmbd2022.

29.   Leung KH, Rowe SP, Pomper MG, Du Y. A three-stage, deep learning, ensemble approach for prognosis in
patients with Parkinson's disease. EJNMMI Res. 2021;11(1):52.

30.   Solana-Lavalle G, Rosas-Romero R. Classification of PPMI MRI scans with voxel-based morphometry and
machine learning to assist in the diagnosis of Parkinson's disease. Comput Methods Programs Biomed.
2021;198:105793.

31.   Szegedy C, Ioffe S, Vanhoucke V, Alemi AA, editors. Inception-v4, Inception-ResNet and the Impact of
Residual Connections on Learning. National Conference on Artificial Intelligence; 2016.

32.   Ellis KA, Rowe CC, Villemagne VL, Martins RN, Masters CL, Salvado O, et al. Addressing population aging
and Alzheimer's disease through the Australian Imaging Biomarkers and Lifestyle study: Collaboration with the
Alzheimer's Disease Neuroimaging Initiative. Alzheimer's & dementia. 2010;6(3):291-6.

33.   Malone IB, Cash D, Ridgway GR, MacManus DG, Ourselin S, Fox NC, et al. MIRIAD--Public release of a

multiple time point Alzheimer's MR imaging dataset. NeuroImage. 2013;70:33-6.

34. Marcus DS, Wang TH, Parker J, Csernansky JG, Morris JC, Buckner RL. Open Access Series of Imaging
Studies (OASIS): cross-sectional MRI data in young, middle aged, nondemented, and demented older adults. J
Cogn Neurosci. 2007;19(9):1498-507.

35. Gauthier S, Reisberg B, Zaudig M, Petersen RC, Ritchie K, Broich K, et al. Mild cognitive impairment. The
lancet. 2006;367(9518):1262-70.

36. Altan G. Deep Learning-based Mammogram Classification for Breast Cancer. International Journal of
Intelligent Systems and Applications in Engineering. 2020;8(4):171-6.

37. Altan G, Kutlu Y, Allahverdi N. Deep Learning on Computerized Analysis of Chronic Obstructive Pulmonary
Disease. IEEE J Biomed Health Inform. 2019.

38. Selvaraju RR, Cogswell M, Das A, Vedantam R, Parikh D, Batra D. Grad-CAM: Visual Explanations from
Deep Networks via Gradient-Based Localization. International Journal of Computer Vision. 2019;128(2):336-59.

39. Yee E, Ma D, Popuri K, Wang L, Beg MF, The Alzheimer's Disease Neuroimaging I, et al. Construction of
MRI-Based Alzheimer's Disease Score Based on Efficient 3D Convolutional Neural Network: Comprehensive
Validation on 7,902 Images from a Multi-Center Dataset. J Alzheimers Dis. 2021;79(1):47-58.

40. Ansart M, Epelbaum S, Bassignana G, Bone A, Bottani S, Cattai T, et al. Predicting the progression of mild
cognitive impairment using machine learning: A systematic, quantitative and critical review. Med Image Anal.
2021;67:101848.

41. Selkoe DJ. Preventing Alzheimer's disease. Science. 2012;337(6101):1488-92.

42. Frisoni GB, Fox NC, Jack CR, Jr., Scheltens P, Thompson PM. The clinical use of structural MRI in Alzheimer
disease. Nat Rev Neurol. 2010;6(2):67-77.

43. Abrol A, Bhattarai M, Fedorov A, Du Y, Plis S, Calhoun V, et al. Deep residual learning for neuroimaging: An
application to predict progression to Alzheimer's disease. J Neurosci Methods. 2020;339:108701.

44. Wachinger C, Salat DH, Weiner M, Reuter M, Initiative AsDN. Whole-brain analysis reveals increased
neuroanatomical asymmetries in dementia for hippocampus and amygdala. Brain. 2016;139(12):3253-66.

45. Derflinger S, Sorg C, Gaser C, Myers N, Arsic M, Kurz A, et al. Grey-matter atrophy in Alzheimer's disease is
asymmetric but not lateralized. Journal of Alzheimer's Disease. 2011;25(2):347-57.

46. Liu J, Wang Y, Qiao Y, editors. Sparse deep transfer learning for convolutional neural network. Proceedings
of the Thirty-First AAAI Conference on Artificial Intelligence; 2017.

47. Ke A, Ellsworth W, Banerjee O, Ng AY, Rajpurkar P. CheXtransfer: Performance and Parameter Efficiency of
ImageNet Models for Chest X-Ray Interpretation. arXiv preprint arXiv:210106871. 2021.

48. Joel D, Berman Z, Tavor I, Wexler N, Gaber O, Stein Y, et al. Sex beyond the genitalia: The human brain
mosaic. Proc Natl Acad Sci U S A. 2015;112(50):15468-73.

49. Eliot L, Ahmed A, Khan H, Patel J. Dump the "dimorphism": Comprehensive synthesis of human brain
studies reveals few male-female differences beyond size. Neurosci Biobehav Rev. 2021;125:667-97.

50. Forest MG, Peretti ED, Bertrand J. Hypothalamic-pituitary-gonadal relationships in man from birth to
puberty. Clin Endocrinol (Oxf). 1976;5(5):551-69.

51. Makris N, Swaab DF, Der Kouwe AJWV, Abbs B, Boriel D, Handa RJ, et al. Volumetric parcellation
methodology of the human hypothalamus in neuroimaging: Normative data and sex differences. NeuroImage.
2013;69:1-10.

52. Raz N, Gunningdixon FM, Head D, Williamson A, Acker JD. Age and Sex Differences in the Cerebellum and
the Ventral Pons: A Prospective MR Study of Healthy Adults. Am J Neuroradiol. 2001;22(6):1161-7.

53. Raz N, Dupuis JH, Briggs SD, Mcgavran C, Acker JD. Differential effects of age and sex on the cerebellar
hemispheres and the vermis: a prospective MR study. Am J Neuroradiol. 1998;19(1):65-71.

54.   Deng J, Dong W, Socher R, Li L-J, Li K, Fei-Fei L, editors. Imagenet: A large-scale hierarchical image
database. 2009 IEEE conference on computer vision and pattern recognition; 2009: Ieee.
55.   Fischl B. FreeSurfer. NeuroImage. 2012;62(2):774-81.
56.   Esteban O, Markiewicz CJ, Blair RW, Moodie CA, Isik AI, Erramuzpe A, et al. fMRIPrep: a robust
preprocessing pipeline for functional MRI. Nat Med. 2019;16(1):111-6.
57.   Coalson TS, Van Essen DC, Glasser MF. The impact of traditional neuroimaging methods on the spatial
localization of cortical areas. Proc Natl Acad Sci U S A. 2018;115(27):e6356-e65.
860