Nucleotide substitutions during speciation may explain substitution rate

variation

Thijs Janzen[1*], Folmer Bokma[2], Rampal S. Etienne[1]

[1] Groningen Institute for Evolutionary Life Sciences, University of Groningen, Box

11103, 9700 CC Groningen, The Netherlands

[2] Center for Ecological and Evolutionary Synthesis (CEES) Department of
BioSciences, University of Oslo, PO Box 1066, Blindern, 0316 Oslo, Norway
10

* corresponding author: t.janzen@rug.nl

ABSTRACT

Although molecular mechanisms associated with the generation of mutations are

highly conserved across taxa, there is widespread variation in mutation rates

between evolutionary lineages. When phylogenies are reconstructed based on

nucleotide sequences, such variation is typically accounted for by the

assumption of a relaxed molecular clock, which, however, is just a statistical

distribution of mutation rates without any underlying biological mechanism.

Here, we propose that variation in accumulated mutations may be partly

explained by an elevated mutation rate during speciation. Using simulations, we

show how shifting mutations from branches to speciation events impacts

inference of branching times in phylogenetic reconstruction. Furthermore, the

resulting nucleotide alignments are better described by a relaxed than by a strict

molecular clock. Thus, elevated mutation rates during speciation potentially

explain part of the variation in substitution rates that is observed across the tree

of life.

Keywords: molecular clock, speciation, phylogenetic reconstruction, substitution

rate variation

INTRODUCTION

Phenotypic diversification occurs at a higher rate in some clades than in others

(Simpson 1945; van Valen 1985; Ricklefs 2006; Rabosky et al. 2007; Jansson and

Davies 2008) and similarly, there is substantial variation across evolutionary

lineages in the rate of molecular evolution (King and Wilson 1975), such as that

of nucleotide sequences (Nabholz et al. 2008; Bromham 2011; Dowle et al. 2013;

Sung et al. 2016). As a consequence, studies attempting to reconstruct the

phylogeny of a clade often find that the sequence data do not support the

assumption of a strict molecular clock, i.e. constant substitution rates across

lineages. For such cases, phylogenetic inference software allows one to use a

relaxed molecular clock (Drummond et al. 2006; Lepage et al. 2007), which

assumes that the substitution rate varies between lineages according to a

statistical distribution such as a gamma or lognormal distribution. However, the

relaxed molecular clock thus introduces at least one additional degree of

freedom, namely the variance of the distribution of substitution rates. Moreover,

it is a rather ad-hoc solution with little underlying biological reasoning.

Here, we present a more explicit biological explanation for variation in

substitution rates across the tree of life; we posit that differences in apparent

substitution rates between lineages are due to processes acting exclusively or

predominantly during speciation. Such processes might include founder effects,

bottlenecks, inbreeding, hybridization, selection for an increased mutation rate,

divergent selection and local adaptation, all of which could lead to a short,

temporary surge in the substitution rate. Due to (effectively) random extinction

of lineages, different branches of a reconstructed phylogeny will differ in how

often they experienced such short episodes of accelerated substitution rates,

resulting in differences in apparent substitutions rates along these branches. It is

worthwhile to investigate and model this process, because empirical data

indicate that inferred substitution rates correlate with numbers of branching

events (Pagel et al. 2006; Eo and DeWoody 2010; Lanfear et al. 2010; Duchene

and Bromham 2013; Bromham et al. 2015 but see Goldie et al. 2011), indicating

that such a process might drive observed patterns.

METHODS

We propose a model where substitutions accumulate not only along the

branches of a phylogeny, but also at speciation events including not only the

internal nodes of the phylogeny but also those pruned from the phylogeny by

extinction. We first make the standard assumption that gradual sequence

evolution along a phylogenetic branch can be modeled as a time-homogeneous

Markov process with substitution matrix:

79
$$\mathbf{Q} = \begin{bmatrix} -\mu_{AT} - \mu_{AC} - \mu_{AG} & \mu_{AT} & \mu_{AC} & \mu_{AG} \\ \mu_{TA} & -\mu_{TA} - \mu_{TC} - \mu_{TG} & \mu_{TC} & \mu_{TG} \\ \mu_{CA} & \mu_{CT} & -\mu_{CA} - \mu_{CT} - \mu_{CG} & \mu_{CG} \\ \mu_{GA} & \mu_{GT} & \mu_{GC} & -\mu_{GA} - \mu_{GT} - \mu_{GC} \end{bmatrix}$$

80 where μ_{ij} denotes the mutation rate from nucleotide i to nucleotide j. The

81 transition probabilities of nucleotide substitutions after time t of gradual

82 sequence evolution are then given by the matrix

$$\mathbf{P_a}(t) = \exp(\mathbf{Q}t)$$

where the subscript $\mathbf{a}$ indicates anagenetic change, i.e. gradual accumulation of

substitutions over time. This matrix can be multiplied with an initial probability

vector at time $t = 0$ to yield the probabilities for each of the four nucleotides at

time t.

In addition to gradual sequence evolution over time, we assume that sequences

may change rapidly during speciation. We can thus assume another matrix $\mathbf{P_c}$

(subscript c for "cladogenetic") that describes the nucleotide transition

probabilities during a single speciation event.

The processes that may accelerate sequence evolution during speciation, such as

founder effects, bottlenecks, inbreeding, hybridization, and adaptation to novel

environments, may well result in different kinds of substitutions than those that

take place over time in established species. However, for mathematical

convenience we will here assume that we can write:

$$\mathbf{P}_c = \exp(\mathbf{Q}\tau)$$

where τ is a parameter that measures the effect of substitutions during

speciation. In other words, we assume that nucleotide sequence evolution is only

accelerated during speciation events, but not qualitatively altered. The

acceleration is then measured by parameter τ: a single speciation event causes as

much sequence evolution as τ years of gradual evolution over time. Note that this

interpretation requires that the μ_{ij} used in $\mathbf{P}_c$ must be identical to that used in $\mathbf{P}_a$.

For $\tau = 0$, $\mathbf{P}_c$ becomes the identity matrix, and our model reduces to the standard

model of sequence evolution that only assumes substitutions along phylogenetic

branches.

For simplicity we assume in our simulations that sequence evolution can be

modeled as a Jukes-Cantor process, for which $\mathbf{Q}$ is given by:

$$\mathbf{Q} = \begin{pmatrix} -\dfrac{3\mu}{4} & \dfrac{\mu}{4} & \dfrac{\mu}{4} & \dfrac{\mu}{4} \\ \dfrac{\mu}{4} & -\dfrac{3\mu}{4} & \dfrac{\mu}{4} & \dfrac{\mu}{4} \\ \dfrac{\mu}{4} & \dfrac{\mu}{4} & -\dfrac{3\mu}{4} & \dfrac{\mu}{4} \\ \dfrac{\mu}{4} & \dfrac{\mu}{4} & \dfrac{\mu}{4} & -\dfrac{3\mu}{4} \end{pmatrix}$$

Given a rooted phylogeny and a root sequence (e.g. some arbitrary sequence

assumed to represent the ancestral sequence), several existing software

packages (e.g. the R package phangorn, Schliep 2011; the python module

pyvolve, Sipos *et al.* 2011; the R package phylosim, Spielman & Wilke 2015)

provide algorithms to simulate sequence evolution along the branches of the

phylogeny, by applying the transition matrix sequentially along the phylogenetic

tree. Here, we extend this methodology to also include substitutions

accumulated at the nodes of the phylogeny. We implemented this in the R

package 'nodeSub' (https://github.com/thijsjanzen/nodeSub).

Testing the impact of node substitution models using simulations

To identify the amount of error in phylogenetic inference caused by

assuming a (relaxed) molecular clock when substitutions actually arise (in part)

during speciation, we simulated sequence evolution on known trees and then

reconstructed the phylogeny from the simulated sequences, assuming strict and

relaxed molecular clocks. We simulated sequence evolution with the node

substitution model introduced above, with various degrees of acceleration (τ) at

the nodes of the tree (including $\tau = 0$), and with various extinction rates. We then

compared the resulting trees with the original true tree using a number of

statistics: the gamma statistic (Pybus and Harvey 2000), the beta statistic

(Aldous 2001), the mean branch length (Faith 1992; Clarke and Warwick 2001),

tree height, the normalized Lineages Through Time (nLTT) statistic (Janzen et al.

2015) and the Jenson-Shannon distance metric comparing the Laplacian

spectrum (Lewitus and Morlon 2016).

To correct for any error in phylogenetic inference incurred by processes

independent of the node substitution model chosen (e.g. stochasticity in

substitution accumulation, stochasticity in phylogenetic tree creation), we also

inferred a phylogenetic tree for a *twin* (sensu Bilderbeek, Laudanno & Etienne

2020) alignment, where we simulated an alignment with the exact same number

of substitutions, using the same true tree, but using a standard, strict-clock,

substitution model. Then, for the distribution of trees resulting from this *twin*

alignment, we estimated the same summary statistics. The error introduced by

the node substitution model is then the difference between the error of the node

substitution posterior and the error in the *twin* posterior.

Phylogenetic reconstruction was performed with BEAST2 (Bouckaert et

al. 2019) using the R package *babette* (Bilderbeek and Etienne 2018). BEAST2

inference was performed using default priors with a birth-death prior as tree

prior (or a Yule prior if the extinction rate is zero), and assuming the JC69

nucleotide substitution model and a strict clock model. The BEAST chain was run

for 10 million steps, whilst sampling a tree every 5000 steps. After completion,

the first 10% of the chain was discarded as burn-in.

Obtaining a twin alignment

We generated a twin alignment conditional on a phylogeny, a node substitution

model, and a mutation rate. Because an alignment generated using a node

substitution model has accumulated substitutions at the nodes in addition to

those along the branches, generally the overall number of substitutions

accumulated is higher than an alignment simulated using a branch-only model.

Thus, in order to generate a *twin* alignment that contains the same amount of

information (substitutions) we need to increase the mutation rate. This can be

done by calculating the estimated time spent at the nodes, relative to the time

spent on the branches. This provides a correction factor which we used to make

a first educated guess as to which mutation rate is most likely to generate an

alignment with the branch-only model with the same number of substitutions.

We obtain:

$$\mu_{twin} = \mu\left(1 + \frac{\tau(2N + H)}{\sum t_{branch}}\right) \quad (1)$$

where μ is the mutation rate used in the node substitution model, τ is the time

spent on the node, N is the number of internal nodes in the tree, H is the number

of hidden nodes in the tree and $\sum t_{branch}$ is the total branch length of the tree. On

the assumption that the rates of speciation (b) and extinction (d) have been

constant throughout the tree, the number of hidden nodes H_{branch} on a branch

with endpoints t_0 and t_1 follows a Poisson distribution (Bokma et al. 2012;

Manceau et al. 2020):

$$H_{branch} = 2b(t_0 - t_1) + 2\ln\frac{b\exp\big((b-d)t_1\big) - d}{b\exp\big((b-d)t_0\big) - d} \ (2)$$

We simulated sequences of 10 kb, with μ = 0.001. We used this educated guess as

a starting point to find the mutation rate that yields on average the same number

of mutations as observed in our alignment simulated with the node substitution

model (because the latter is just a single realization, which may be quite different

from the expected number of mutations). Then, using our corrected mutation

rate, we generated twin alignments until we obtained one exactly matching the

number of mutations of the alignment simulated with the node substitution

model.

Simulation settings

We generated birth-death trees with varying degrees of extinction rate d in [0,

0.1, 0.3, 0.5] and a single speciation rate of b = 1. Trees were simulated

conditional on 100 tips.

Varying the time spent on the nodes relative to the crown age

We varied τ in [0, 0.01, 0.05, 0.1, 0.2, 0.4] times the crown age (e.g. when the

crown age of the simulated tree is 3MY, τ = [0, 0.03, 0.15, 0.3, 0.6, 1.2] MY). Again,

for each combination of τ and extinction (d = [0, 0.1, 0.3, 0.5]) we simulated 100

trees and for each tree we generated one node substitution alignment and one

twin alignment.

The impact of tree balance

In unbalanced trees, some terminal branches are connected by many more past

branching events to the root of the tree than other terminal branches. Hence, we

expect that balance of a tree might have a substantial effect on the error in

phylogenetic inference: less balanced trees are expected to have higher error. To

test this, we compared fully balanced (β = 10.0) with extremely unbalanced,

caterpillar trees (β = -2). We did so by simulating the branching times of a birth-

death tree, and assigning these to a fully balanced or fully unbalanced topology.

Thus, the only difference between the trees is the topology. Then, for both the

balanced and unbalanced tree a node substitution alignment was generated, with

the same number of total substitutions, and setting τ as a function of crown age.

As an extra check, we also generated a node substitution alignment for the

original birth-death tree from which the branching times were used. For all three

alignments we inferred a phylogenetic tree as in the other scenarios, and

compared the error in phylogenetic inference. Calculating the Laplacian

Spectrum for caterpillar trees with extreme unbalance returned NA values, and

hence we omitted the Laplacian Spectrum summary statistic in this analysis.

Support for strict and relaxed clock models

To test whether the alignment originating from a process with node

substitutions was better described by a relaxed than by a fixed clock model, we

repeated the analysis, but now inferred the marginal likelihood of the relaxed

clock and strict clock models using the "NS" package for BEAST2, which applies

Nested Sampling to obtain the marginal posterior likelihood for both models

(Russel et al. 2019). We used the function 'bbt_run' from the *babette* package

(Bilderbeek and Etienne 2018) in combination with the function

'create_ns_mcmc' from the *beautier* package (Bilderbeek and Etienne 2018). This

performs a Nested Sampling MCMC run using BEAST2, which runs until

convergence is detected. Then, we converted the obtained marginal likelihoods

to a relative weight for each model (by dividing both marginal likelihoods by

their sum) which allows for comparison of posterior support for each model

across parameter settings and trees.

RESULTS

Summary statistics

We compared summary statistics of trees inferred from alignments using the

node substitution model, with summary statistics of *twin* trees inferred from

alignments with identical information content, but generated without the node

substitution model (e.g. with only substitutions along the branches, and a fixed

clock rate). We find that particularly summary statistics influenced by branching

times are affected (Figure 2, gamma and nLTT statistic). For these summary

statistics, we find an increased difference as the node substitution rate τ

increases. The differences in the value of these summary statistics manifest

themselves across all extinction rates, although the effect tends to be somewhat

smaller for higher extinction rates.

240
Figure 1. Difference in summary statistic values between trees inferred from an
alignment generated with node substitutions (colored boxplots), and *twin* trees
that were inferred from an alignment generated without node substitions (grey
boxplots). Shown are the residual errors, reflecting distributions corrected by
subtracting the median value of the *twin* trees. As a result, the distribution of the
standard substitution model is always centered on 0. We explore τ as a fraction
of crown age (horizontal axis), and the impact of extinction (*d*, columns). The
dotted line indicates zero difference between the two substitution models. The
summary statistics are the beta and gamma statistic, Laplacian spectrum, mean
branch length, nLTT statistic, and crown age.
251

Figure 2. Effect of the node substitution model for phylogenies differing in balance **A**) example plots of a randomly generated birth-death tree (top), a fully balanced tree generated using the same branching times as the birth-death tree (middle) and a very unbalanced tree generated using the same branching times as the birth-death tree (bottom). Shown are trees with 20 tips for illustrative purposes, but results in **B** are from trees with 100 tips. **B)** Difference in summary statistic with the true birth-death tree for phylogenetic trees inferred from alignments generated using the node substitution model on either balanced, unbalanced or random trees. We explore τ as a fraction of crown age (horizontal axis), and the impact of extinction (d, columns). The dotted line indicates zero difference with the true tree. The summary statistics are the beta and gamma statistic, mean branch length, nLTT statistic and tree height.

Tree balance clearly influences the sensitivity of inference to node substitutions

(Figure 2). The inference error is larger for unbalanced trees, again only for the

gamma and the nLTT statistic. Fully balanced trees show slightly less error than

birth-death trees. Overall, all three types of trees show an increased error when

alignments are generated using the node substitution model. Errors are

particularly large for the beta statistic, but that is expected because it measures

topological features of the tree that we modified artificially.

Support for strict and relaxed clock models

We compare the relative support for each model, reflected by the relative weight

of the marginal likelihood. With an increasing amount of time spent at the nodes

τ, the median weight of the relaxed clock model increases for the node

substitution alignment, with generally (across extinction rates) a higher weight

than the strict clock model for values of τ that are equal or larger than 0.1 times

the crown age (Figure 3). For the *twin* alignment, the strict clock model is

preferred, as expected, because this is the generating model.

For low values of τ (smaller than 0.1 times the crown age), we do not find any

effect of the balance of the tree on the marginal likelihood of the relaxed clock

model (Figure 4), in line with our finding above. However, for intermediate

values of τ (0.1 and 0.2), we find that unbalanced trees tend to have a higher

marginal weight for the relaxed clock model. For high values of τ (0.4), we find

that the marginal weight for the relaxed clock model is always higher, regardless

of the balance of the tree.

Figure 3. Marginal likelihood weight of the relaxed and strict clock model for
varying time spent on the nodes (τ), where τ is chosen as fraction of the crown
age. Alignments generated with a node substitution model (top row) are
compared with alignments generated without node substitutions (bottom row).
Per parameter combination, 100 replicate trees were analyzed. Because many
dots are plotted on top of each other, we use solid lines to indicate the best fitting
locally estimated scatterplot smoothing (LOESS), and the 95% Confidence
interval (grey shaded area) of the LOESS curve. As the time spent on the nodes
increases, posterior support for the relaxed clock model increases, but only if the
alignment was generated with a node substitution model.

Figure 4. Marginal likelihood weight of the relaxed clock model for trees of
varying balance, split out across different extinction rates (d = [0, 0.1, 0.3, 0.5])
and time spent on the nodes (τ), where τ is chosen as fraction of the crown age
(e.g. τ = 0.1 reflects a node time of 10% of the crown age). Per parameter
combination, 100 replicate trees were analyzed. Because many dots are plotted
on top of each other, we use solid lines to indicate the best fitting linear
regression and the 95% confidence interval (grey shaded area) of regression.

Box 1

The general framework we have described here, allows for the exploration

of the impact of accumulation of substitutions during speciation events, which

we can vary in relation to the accumulation of substitutions over time between

speciation events. As the accumulation of substitutions during speciation events

becomes more prominent, the impact on the resulting tree becomes more and

more pronounced, in particular the impact on branching time estimates.

The next question is what proportion of substitutions in real data is the

result of processes acting during speciation, and what proportion arises

gradually over time between speciation events. Future work should employ a

likelihood framework able to infer the expected number of substitutions

accumulated during speciation events. The joint likelihood of the phylogenetic

tree and the sequence alignment can be computed using an SSE framework; one

needs to repeatedly apply equations (A1), (A2) and (A3) from Goldberg & Igic

(2012) to each nucleotide from tip to root of the phylogeny, where we assume

that the speciation and extinction rates do not depend on the nucleotide. The

equations then become (with D_{Ni} the probability of the nucleotides at the focal

locus in subclade N being in their observed states when the nucleotide at time t

is equal to i, and E_i the probability of extinction of the clade when the nucleotide

at time t is i, and D_{Ci} the probability just before an observed node in the

phylogeny):

$$\frac{dD_{Ni}}{dt} = -\left(\lambda + \sum_j r_{ij} + \mu\right)D_{Ni}$$

$$+ \sum_j r_{ij}D_{Nj} + \lambda\sum_{j,k}\left(P_{c,ij}D_{Nj}P_{c,ik}E_k + P_{c,ij}E_jP_{c,ik}D_{Nk}\right)$$

$$\frac{dE_i}{dt} = -\left(\lambda + \sum_j r_{ij} + \mu\right)E_i + \sum_j r_{ij}E_j + \mu + \lambda\sum_{j,k}\left(P_{c,ij}E_jP_{c,ik}E_k\right)$$

$$D_{Ci} = \lambda\sum_{j,k}\left(P_{c,ij}D_{Nj}P_{c,ik}D_{Mk} + P_{c,ij}D_{Mj}P_{c,ik}D_{Nk}\right)$$

This likelihood computation can be implemented in software able to

accommodate joint estimation, such as RevBayes (Hohna et al. 2016).

DISCUSSION

We have shown that an increased substitution rate during speciation

events potentially provides a mechanistic explanation of variation in substitution

rates across the branches of phylogenetic trees. We have shown that trees

inferred from alignments generated with this substitution model differ

substantially from trees inferred from alignments generated with a standard

substitution model, especially concerning branching times. Furthermore, we find

that this new substitution model can potentially explain widespread support for

relaxed molecular clocks.

One might expect that a high extinction rate, by elevating numbers of

hidden nodes, leads to a greater impact of node substitutions. It may therefore be

counterintuitive that in our simulation study we did not find such an effect of

higher rates of extinction. However, we conditioned our alignments on having

the same total number of substitutions. Thus, with higher extinction and hence

more hidden nodes, fewer substitutions occur on the observed nodes, while the

number of hidden nodes is proportional to branch length (Eq.2) and can hence

be interpreted as substitutions on the branches. The result is that with higher

extinction, the signature of node substitutions decreases.

The model we propose here takes the simplest form, assuming a JC69

substitution matrix, identical substitution rates, identical substitution matrices

between nodes and branches, and constant birth-death rates over time. These

assumptions were made as a most basic starting point, but can be relaxed in

future analyses, for instance by introducing a different substitution matrix at the

nodes, or by studying the effect of node substitutions on trees that are generated

by density dependent speciation rates (Etienne et al. 2012). By starting with the

most tractable version of the node substitution model, we have provided a first

proof of concept of the potential impact of node substitutions without

overcomplicating matters.

Previous methods have applied rather ad-hoc corrections to account for

differences in substitution rates across different branches in the same

phylogeny, typically referred to as the 'relaxed clock' approach. These methods

provide satisfying statistical solutions to account for variation in substitution

rates, but refrain from providing biological explanations for this observed

phenomenon. The node substitution model we introduce here provides this

explanation: branches that have accumulated a number of 'hidden' branching

events, e.g. speciation events of species that have subsequently gone extinct,

have a higher number of accumulated substitution events during these 'hidden'

speciation events. When we compared the marginal likelihood of the relaxed

clock model versus the strict clock model for alignments generated with the

node substitution model, we found that marginal likelihoods for the relaxed

clock model are much higher. This indicates that our proposed process of

accumulating substitutions during speciation events can generate patterns in the

alignment that are picked up by phylogenetic methods as evidence for a relaxed

clock model, without actually using a relaxed clock model. This does not

necessarily mean that our model provides a perfect explanation, but we do want

to emphasize that our model provides a first potential mechanistic explanation

for the observed patterns.

Our model reflects many of the aspects of the substitution spikes

introduced by Manceau *et al.* (Manceau et al. 2020). In that model, Manceau *et al.*

introduce an alternative solution for punctuated equilibria-like patterns in

molecular evolution through the implementation of spikes of substitution, e.g.

moments in time at which there is an increased substitution rate. They let these

moments occur at speciation events, and also model the probability of such an

event happening at a speciation event (rather than assuming they always do, like

we did here). Furthermore, they provide a first foray into formulating a

likelihood for a node-associated substitution model, but only provide a

likelihood of the alignment, given the tree. Whereas their work modulates the

impact of substitutions at the nodes through modifying the substitution rate, our

work modulates this impact through modifying the time spent at the node. We

think these two models are highly complementary, and we embrace future

endeavors towards combining insights from both of them.

The notion of accelerated evolution during speciation events ties in

closely with the theory of punctuated equilibrium; where Eldridge and Gould

proposed that perhaps evolution is not a gradual process, but rather a process

with distinct bursts of phenotypic and morphological change. (Eldredge and

Gould 1972). Our framework provides a first step towards being able to test this

notion, where information on the estimated fraction of substitutions

accumulated at the nodes can directly inform us about whether the majority of

mutations is accumulated during periods of stasis (e.g. at the branches), or

during speciation (e.g. at the nodes).

With our introduction of the node substitution model, we hope to stimulate

discussion on the biological explanation of variation in substitution rates within

and across phylogenies. Furthermore, we hope to have set a first step in

improving our understanding of this variation, and improving phylogenetic

inference as a whole.

REFERENCES

Aldous D. 2001. Stochastic models and descriptive statistics for phylogenetic

trees, from Yule to today. Stat. Sci. 16:23–34.

Bilderbeek R.J.C., Etienne R.S. 2018. babette⬚: BEAUti 2 , BEAST2 and Tracer for

R. 2018:2034–2040.

Bilderbeek R.J.C., Laudanno G., Etienne R.S. 2020. Quantifying the importance of

an inference model in Bayesian phylogenetics. bioRxiv.

Bokma F., van den Brink V., Stadler T. 2012. Unexpectedly many extinct

hominins. Evolution. 66:2969–2974.

Bouckaert R., Vaughan T.G., Barido-Sottani J., Duchene S., Fourment M.,

Gavryuskina A., Heled J., Jones G., Kuhnert D., De Maio N., Matschiner M., K.

Mendes F., Muller N.F., Ogilvie H.A., du Plessis L., Popinga A., Rambaut A.,

Rasmussen D., Siveroni I., Suchard M.A., Wu C.H., Xie D., Zhang C., Stadler T.,

Drummond A.J. 2019. BEAST 2 .5 ⬚: An advanced software platform for

Bayesian evolutionary analysis. PLoS Comput. Biol. 15:e1006650.

Bromham L. 2011. The genome as a life-history character: Why rate of molecular

evolution varies between mammal species. Philos. Trans. R. Soc. B Biol. Sci.

366:2503–2513.

Bromham L., Hua X., Lanfear R., Cowman P.F. 2015. Exploring the Relationships

between Mutation Rates , Life History , Genome Size , Environment , and

Species Richness in Flowering Plants. 185:507–524.

Clarke K., Warwick R. 2001. A further biodiversity index applicable to species

lists: variation in taxonomic distinctness. Mar. Ecol. Prog. Ser. 216:265–278.

Dowle E.J., Morgan-Richards M., Trewick S.A. 2013. Molecular evolution and the

latitudinal biodiversity gradient. Heredity (Edinb). 110:501–510.

Drummond A.J., Ho S.Y.W., Phillips M.J., Rambaut A. 2006. Relaxed phylogenetics

and dating with confidence. PLoS Biol. 4:e88.

Duchene D., Bromham L. 2013. Rates of molecular evolution and diversification

in plants: Chloroplast substitution rates correlate with species-richness in

the Proteaceae. BMC Evol. Biol. 13:1.

Eldredge N., Gould S.J. 1972. Punctuated equilibria: an alternative to phyletic

gradualism. Models In Paleobiology. Freeman Cooper and Co. p. 82–115.

Eo S.H., DeWoody J.A. 2010. Evolutionary rates of mitochondrial genomes

correspond to diversification rates and to contemporary species richness in

birds and reptiles. Proc. R. Soc. B Biol. Sci. 277:3587–3592.

Etienne R.S., Haegeman B., Stadler T., Aze T., Pearson P.N., Purvis A., Phillimore

A.B. 2012. Diversity-dependence brings molecular phylogenies closer to

agreement with the fossil record. Proc. R. Soc. B Biol. Sci. 279:1300–1309.

Faith D.P. 1992. Conservation evaluation and phylogenetic diversity. Biol.

Conserv. 61:1–10.

Goldie X., Lanfear R., Bromham L. 2011. Diversification and the rate of molecular

evolution: No evidence of a link in mammals. BMC Evol. Biol. 11:286.

Hohna S., Landis M.J., Heath T.A., Boussau B., Lartillot N., Moore B.R.,

Huelsenbeck J.P., Ronquist F. 2016. RevBayes: Bayesian phylogenetic

inference using graphical models and an interactive model-specification

language. Syst. Biol. 65:726–736.

Jansson R., Davies T.J. 2008. Global variation in diversification rates of flowering

plants: Energy vs. climate change. Ecol. Lett. 11:173–183.

Janzen T., Höhna S., Etienne R.S.R.S. 2015. Approximate Bayesian Computation of

diversification rates from molecular phylogenies: introducing a new

efficient summary statistic, the nLTT. Methods Ecol. Evol. 6:566–575.

King M.C., Wilson A.C. 1975. Evolution at two levels in humans and chimpanzees.

Science. 188:107–116.

Lanfear R., Ho S.Y.W., Love D., Bromham L. 2010. Mutation rate is linked to

diversification in birds. Proc. Natl. Acad. Sci. U. S. A. 107:20423–20428.

Lepage T., Bryant D., Philippe H., Lartillot N. 2007. A general comparison of

relaxed molecular clock models. Mol. Biol. Evol. 24:2669–80.

Lewitus E., Morlon H. 2016. Characterizing and comparing phylogenies from

their laplacian spectrum. Syst. Biol. 65:495–507.

Manceau M., Marin J., Morlon H., Lambert A. 2020. Model-based inference of

punctuated molecular evolution. Mol. Biol. Evol.:msaa144.

Nabholz B., Glémin S., Galtier N. 2008. Strong variations of mitochondrial

mutation rate across mammals - The longevity hypothesis. Mol. Biol. Evol.

25:120–130.

Pagel M., Venditti C., Meade A. 2006. Large punctuational contribution of

speciation to evolutionary divergence at the molecular level. Science.

314:119–121.

Pybus O., Harvey P. 2000. Testing macro–evolutionary models using incomplete

molecular phylogenies. Proc. R. Soc. B Biol. Sci. 267:2267–72.

Rabosky D.L., Donnellan S.C., Talaba A.L., Lovette I.J. 2007. Exceptional among-

lineage variation in diversification rates during the radiation of Australia's

most diverse vertebrate clade. Proc. R. Soc. B Biol. Sci. 274:2915–2923.

Ricklefs R.E. 2006. Global variation in the diversification rate of passerine birds.

Ecology. 87:2468–2478.

Russel P.M., Brewer B.J., Klaere S., Bouckaert R.R. 2019. Model Selection and

Parameter Inference in Phylogenetics Using Nested Sampling. Syst. Biol.

68:219–233.

Schliep K.P. 2011. phangorn⬚: phylogenetic analysis in R. 27:592–593.

Simpson G.G. 1945. SECTION OF BIOLOGY: Tempo and Mode in Evolution. Trans.

N. Y. Acad. Sci. 8:45–60.

Sipos B., Massingham T., Jordan G.E., Goldman N. 2011. PhyloSim - Monte Carlo

simulation of sequence evolution in the R statistical computing

environment. .

Spielman S.J., Wilke C.O. 2015. Pyvolve⬚: A Flexible Python Module for

Simulating Sequences along Phylogenies. :1–7.

Sung W., Ackerman M.S., Dillon M.M., Platt T.G., Fuqua C., Cooper V.S., Lynch M.

2016. Evolution of the insertion-deletion mutation rate across the tree of

life. G3 Genes, Genomes, Genet. 6:2583–2591.

van Valen L.M. 1985. Why and how do mammals evolve unusually rapidly. Evol.

Theory. 7:127–132.

508