1

Multigenerational downregulation of insulin/IGF-1 signalling in adulthood improves lineage survival, reproduction, and fitness in *C. elegans*

Elizabeth M.L. Duxbury[1*], Hanne Carlsson[1], Kris Sales[1], Simone Immler[1], Tracey Chapman[1] and Alexei A. Maklakov[1]

[1]School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich, NR4 7TJ, UK. [*]email: E.Duxbury@uea.ac.uk

Short running title: Reduced IIS protects soma and germline

Keywords: ageing, life history evolution, mutation accumulation, germline, senescence

Abstract

Adulthood-only downregulation of insulin/IGF-1 signalling (IIS), an

evolutionarily conserved pathway regulating resource allocation between

somatic maintenance and reproduction, increases lifespan without fecundity

cost in the nematode, *Caenorhabditis elegans*. However, long-term multi-

generational effects of reduced IIS remain unexplored, and it has been

proposed that inter- and trans-generational effects of reduced IIS can have

adverse effects on offspring fitness. To test this hypothesis, we ran a mutation

accumulation (MA) experiment and downregulated IIS in half of the 400 MA

lines by silencing *daf-2* gene expression using RNA interference (RNAi) across

40 generations. Contrary to the prediction, adulthood-only *daf-2* RNAi reduced

extinction of MA lines both under UV-induced and spontaneous mutation

accumulation suggesting net benefit of multi-generational downregulation of

IIS. Fitness of the surviving UV-induced MA lines was higher under *daf-2* RNAi.

Reduced IIS increased inter-generational F1 offspring fitness under UV stress

but had no quantifiable trans-generational effects. Functional *hrde-1* was

required for both the benefits of multi-generational *daf-2* RNAi and for inter-

generational effects. Overall, we did not find any life-history costs of multi-

generational reduction of IIS and the benefits became more apparent under

stress. Our results thus suggest that multi-generational adulthood-only

reduction of IIS via *daf-2* RNAi has net beneficial effect on fitness. We discuss

the challenges these findings pose for the evolutionary theory of ageing,

suggest possible solutions and how future research may address them.

Introduction

Ageing, the physiological deterioration of an organism leading to increased probability of death and decreased reproduction with advancing adult age, is taxonomically ubiquitous but remains incompletely understood[1-4]. While there is broad agreement that ageing evolves because natural selection gradients on traits decline after reproductive maturity[5-9], the proximate causes of late-life deterioration are less clear[1,2,10,11]. The "disposable soma" theory of ageing suggests that ageing evolves due to competitive resource allocation between the germline and the soma[12-14]. It follows that increased investment in somatic maintenance leading to longer lifespan trades-off with traits associated with reproduction, such as the number or size of progeny[15,16]. Despite considerable support for trade-offs between somatic maintenance and reproduction, growing empirical work questions the universality of such resource allocation trade-offs in the evolution of ageing[2,11,17,18]. Several studies have shown that experimentally increased lifespan, often via the downregulation of genes in nutrient-sensing signalling pathways in adulthood, is not detrimental to reproduction[2,18,19,20,21]. These results are in line with the hypothesis that a gradual decline in selection gradients with advancing age after reproductive maturity results in physiological processes that are optimised for successful early life reproduction, but start causing harm in later life, contributing to somatic deterioration[2,22-28].

However, it has been proposed that reduced offspring quality, for example via reduced germline maintenance, can be a hidden cost of lifespan extension via reduced nutrient-sensing signalling[18]. Germline maintenance, the repair and surveillance of genomic and proteomic integrity in germline stem cells and gametes, is energetically expensive[18,29] and germline signalling plays a key role in resource

allocation to somatic maintenance[18,29,30,31]. Germline ablation results in increased

somatic maintenance and lifespan in *Drosophila melanogaster* fruitflies[32] and

Caenorhabditis elegans nematodes[33-37], although lifespan extension in *C. elegans*

requires an intact somatic gonad[33]. It is important to note that the requirement of the

intact somatic gonad for lifespan extension in germline-less worms does not negate

the possibility of a resource allocation trade-off as sometimes implied, but only shows

that germline signalling is required to mediate the effect[18,32,34]. Similarly, recent work

in *Danio rerio* zebrafish suggests that germline ablation increases somatic

maintenance under stress[38]. Furthermore, nutritional stress in *C. elegans* results in

germline reduction and increased lifespan[39]. When soma-to-germline communication

is disrupted, the number of germ cells is unaffected and lifespan extension is

abolished[39]. Taken together, these results suggest that germline maintenance is

costly and can trade-off with life-history traits such as lifespan and offspring

number[18,29]. The corollary is that increased investment in somatic maintenance and

longevity can trade-off with offspring quality by increasing germline mutation load in

offspring.

We tested this hypothesis by focusing on lifespan extension via reduced

insulin/IGF-1 signalling (IIS) in *Caenorhabditis elegans*. IIS is an evolutionarily

conserved pathway that regulates the physiological response of organisms to their

environment[3]. IIS mechanistically links nutrient intake with development, growth,

reproduction and lifespan across diverse taxa[3,31,40]. Reduced IIS, via genetic and

environmental interventions, consistently extends lifespan[31]. Mutations that reduce

the function of *daf-2* gene in *C. elegans*, the homologue of human IGF-1, increase

lifespan and reduce fitness[41]. However, previous work showed that reducing IIS in

adulthood via *daf-2* RNA interference (RNAi) knockdown in a single generation

increases lifespan without a cost to offspring number[19], quality[20] or fitness in variable

environments[28]. It is possible, however, that multi-generational *daf-2* knockdown will

expose a hidden cost to offspring number and/or offspring quality via accumulation of

germline mutations. Here we tested this prediction via adulthood-only *daf-2* RNAi

knockdown for 40 consecutive generations in 400 spontaneous and UV-induced

mutation accumulation (MA) lines. We found that multi-generational adulthood-only

daf-2 knockdown reduced lineage extinction under both spontaneous and UV-

induced MA and increased fitness of remaining lineages under UV-induced MA.

Reduced IIS protected MA lines from infertility, egg hatching failure and

developmental abnormalities and increased offspring fitness under UV stress but we

found no evidence of trans-generational effects of adulthood-only *daf-2* RNAi.

Materials and Methods

Supplementary methods details are included in the Supplementary Material.

Nematode stocks and culture. The nematode (roundworm) *Caenorhabditis elegans*

N2 wild-type (Bristol) and heritable RNAi deficiency 1 (*hrde-1*) mutant strains were

defrosted from stocks acquired from Caenorhabditis Genetics Center (University of

Minnesota, USA, funded by NIH Office of Research Infrastructure Programs, P40

OD010440) and from the lab of Prof. Eric Miska (Gurdon Institute, University of

Cambridge, UK), respectively, and stored at -80°C until use.

Defrosted *C. elegans* strains were reared through two generations prior to set-

up, on NGM agar supplemented with the fungicide nystatin and antibiotics

streptomycin and ampicillin, to prevent infection (each at 100 μg/ml, as standard

recipes, e.g.[42]) and seeded with the antibiotic-resistant *Escherichia coli* bacterial

strain OP50-1 (pUC4K, from J. Ewbank at the Centre d'Immunologie de Marseille-

Luminy, France). We bleached eggs from the grandparents of experimental

individuals, to standardise parental age and remove any infection or temperature

effects from defrosting, prior to experiments.

Reducing IIS via *daf-2* RNAi feeding in adulthood. To downregulate adulthood

expression of the insulin-like sensing signalling receptor homolog, *daf-2*, we fed late-

L4 larvae with *Escherichia coli* bacteria expressing *daf-2* double-stranded RNA

(dsRNA), that decreases mRNA levels of the complementary transcribed *daf-2*

systemically[43,44]. The *daf-2* gene is upstream of and inhibits the action of master

regulator daf-16/FOXO. RNase-III deficient, IPTG-inducible HT115 *E. coli* bacteria

with an empty plasmid vector (L4440) was used as the control (as[19,20,44]). RNAi

clones were acquired from the Vidal feeding library (Source BioScience, created by

M. Vidal lab, Harvard Medical School, USA) and all clones were verified via

sequencing prior to delivery.

Validation of *daf-2* gene expression under RNAi. We quantified the extent of

downregulation of *daf-2* in Day 2 adults following feeding RNAi, using quantitative

reverse transcriptase polymerase chain reaction (qRT-PCR), in a sample of lines

from all eight MA treatments below, that had been frozen at generation 20 (mean n=

6 MA lines per treatment, with mean n=3 individual worms per MA line; Table S1).

Worms for the gene expression assay were maintained under standard conditions

(as the main experiment) and were reared for two generations on empty vector prior

to the assay, to remove freezing effects and age-synchronise. RNAi was applied from

late-L4 stage. Further method details are in included in Supplementary Material.

Ultraviolet wavelength C (UV-C) irradiation. To induce mutations we UV-irradiated

Day 2 adults, with a calibrated ultraviolet-C (UV-C) radiation dose of 46J/m^2

(wavelength 254nm), via 20 second exposure to the UV-C radiation emitted from the

lamp of a Thermo Scientific Heraguard ECO Safety Cabinet (calibration details in

Supplementary Methods). This dose is in the range of previous UV-C irradiation

doses for *C. elegans* adults or eggs[45-48]. Our pilot work showed that this dose

reduced the fecundity of Day 2 adults laying at 20°C by 61% compared with un-

exposed sham controls (data not shown). Non-irradiated control worms received a

sham-irradiation, by being positioned in identical orientation under the UV-C lamp for

20 seconds, while it was switched off.

Mutation accumulation (MA) lines. To determine the effects of reduced adulthood

IIS, via *daf-2* RNAi, on spontaneous and UV-induced mutation accumulation, we

established 800 MA lines, across two genetic backgrounds. The eight experimental

treatments were the full-factorial combinations of genotype (N2 or *hrde-1* mutant),

RNAi treatment (empty vector or *daf-2* RNAi) and irradiation (UV or sham), with 100

MA lines per treatment. We ran each of the eight treatments in parallel, divided into

two time-staggered independent blocks of 50 lines per treatment, for logistical

reasons and to capture any between-block variation. We tracked MA lines until the

majority of UV-irradiated lines were extinct- 40 generations for N2 and 25

generations for *hrde-1* mutants. Every MA generation, *daf-2* RNAi or empty vector

was delivered from the start of adulthood onwards, and all lines developed on empty

vector. More details are provided in the Supplementary Methods.

Each MA line was propagated as a single individual hermaphrodite per

generation, to create successive genetic bottlenecks, allowing *de novo* deleterious

mutations to accumulate in the relative absence of selection[49-51], if they had no effect

on developmental time or viability, and in the absence of mating. MA is a common

approach used in several species, to study the evolutionary genetics of *de novo*

mutation rates[52]. In wild type *C. elegans*, the spontaneous mutation rate in MA lines

is positively correlated with the rate of fitness decay[53].

Effect of 20 generations of spontaneous and UV-induced mutation

accumulation on fitness and life history of surviving MA lines. To test for life

history differences between the N2 wild-type MA lines after 20 generations of MA and

of adulthood *daf-2* RNAi, we assayed age-specific reproduction, egg size, male

production and adult heat shock resistance (n=100 grand-offspring from each of the

four N2 MA treatments). Further details in Supplementary Methods.

Individual age-synchronised, late-L4 hermaphrodites were set up on separate

e.v. plates, labelled with a unique identifier, to blind experimenters to treatment

identity, and thus avoid bias. The two experimenters conducted the life history assay

in two simultaneous time blocks and each block had an identical and equal

representation of individual worms from all four treatments (n=50 per treatment per

block).

For all experiments, we assayed age-specific offspring production (fecundity)

over the first 6 days of adult life (the reproductive window for *C. elegans*

hermaphrodites), by transferring to fresh plates every 24h, to acquire daily

reproduction counts. Plates were scored for viable adult offspring 2.5 days later.

Individual fitness (lambda) was calculated- a measure analogous to the intrinsic rate

of population growth, that is weighted by early reproduction and fast development

[15,20,54,55]. Total lifetime reproduction (lifetime reproductive success) was calculated for

each individual as the sum of age-specific offspring counts across the first 6 days of

adulthood.

To determine if reduced IIS influenced parental resource allocation into eggs,

under MA, we measured the area of a single egg (in mm^2) produced by 75 individuals

per treatment in the generation 20 age-specific reproduction assay, at peak

reproduction (Day 2). Egg size is commonly used as a proxy for parental

investment[20,56]. We photographed eggs, within 2 hours of lay, under 12x

magnification light microscopy (Leica M165C with Lumenera Infinity 2-7C digital

microscope camera).

Transgenerational effects of *daf-2* RNAi on offspring fitness in N2 and *hrde-1*

mutant backgrounds. To determine whether *daf-2* RNAi was transferred

transgenerationally via the germline, we assayed, in a separate experiment, whether

offspring fitness benefits persisted across three generations of offspring (F1, F2 and

F3) from Day 2 parents treated with *daf-2* RNAi or empty vector control throughout

adulthood, in N2 wild-type and *hrde-1* mutant backgrounds. Offspring of each

generation were maintained on empty vector (n=30 individuals per treatment and per

genotype) and assayed for daily reproduction.

Statistical analyses. All analyses were conducted in R version 4.0.0[57]. Fitness and

total lifetime reproduction data were plotted using the R package 'dabestr' (data

analysis using bootstrap-coupled estimation[58]), for visualisation.

Parental survival in the intergenerational (first) experiment was analysed using

a Cox proportional hazards mixed effects model[59] ('coxme'), fitting RNAi, UV and

their interaction as fixed effects and plate as a random effect. Separate analyses

were conducted with matricides either classed as deaths or as censors. The age at

death response variable contained a coding variable to distinguish deaths from

censored individuals (due to accidental losses). Extinction (survival trajectories)

under mutation accumulation was analysed using Cox proportional hazards

regression analysis ('coxph' function in 'survival' package). The generation number at

which each MA line went extinct was the response variable (instead of age at death)

and this contained a coding variable to distinguish extinctions from censors, as

above. To test the effect of experimental block or genotype (either N2 wildtype or

hrde-1 mutant) on extinction under MA, maximal models were fitted with either a

three-way RNAi x UV x block interaction, or a RNAi x UV x genotype interaction, and

step-wise model simplification conducted[60]. Block was included as a fixed effect to

test for repeatability between the two blocks. For all survival analyses, the z-scores

and p-values were reported, with the z-score defined as the log of the hazard ratio

(risk of death, exp(coef)) divided by the standard error of the log hazard ratio.

Age-specific reproduction was analysed using generalised linear mixed effects

models to account for temporal pseudoreplication of repeated measures on the same

individuals across lifetime, with the template model builder package ('glmmTMB' in

R[61,62]). Models fitted with Poisson, Zero-Inflated Poisson, Generalised Poisson and

Zero-Inflated Generalised Poisson error structure were compared using AIC values

of model fit ('AICtab' function in 'bbmle' package), following[62], to account for under-

or over-dispersion and for zero-inflation, when it was found to occur in simulated

residuals generated with the 'DHARMa' package[63]. Age and a quadratic form of age

(age^2) were fitted as fixed effects in both the conditional and zero-inflation model

formula, and significance assessed in each case. The age^2 term controlled for the

curved (non-linear) trajectory of reproduction across age[64]. UV irradiation, RNAi

treatment and their three-way interaction with age or age^2 were fitted as fixed effects,

and experimental block, observer and a unique plate identifier as random effects, the

latter to control for repeated measures of fecundity from the same individuals across

lifetime. Genotype was substituted with UV as a fixed effect to assess the

transgenerational effects of *daf-2* RNAi on N2 versus *hrde-1* mutants. Models that did

not converge were not included in model comparison, and the converging model with

best AIC fit was presented (as[62]).

Fitness (lambda) was calculated using the 'lambda' function in the 'popbio'

package. More details are provided in the Supplementary Material. Fitness, total

reproduction and egg size data were analysed using a generalised linear model

(GLM) with Gaussian error structure ('lm' function in 'stats' package). Locomotion, a

measure of stress resistance to heat shock, was coded as a binary response variable

(normal or abnormal movement)) and analysed using a GLM with binomial error

structure.

Results and Discussion

Parental *daf-2* RNAi in adulthood extends lifespan and improves offspring fitness under UV-induced stress. Adulthood-only *daf-2* RNAi in N2 wildtype *C. elegans* nematodes, significantly extended parental lifespan relative to empty vector (e.v.) controls under benign conditions (no irradiation) and under ultraviolet-C ("UV") irradiation-induced stress (Cox proportional hazards mixed effects model, coxme, with matricides censored, RNAi: z=10.530, p<0.001; UV: z=0.070, p=0.940; RNAi x UV: z= 1.500, p=0.130; matricides classed as dead: RNAi: z=10.400, p<0.001; UV: z=0.250, p=0.800; RNAi x UV: z= 1.310, p=0.190; Fig. 1a).

There was no cost to parental reproduction, neither under benign conditions nor when parents were UV-irradiated (Fig. S1). Reproduction was estimated as age-specific offspring production (fecundity, Generalised Poisson, RNAi x UV x Age: z = -2.546, p=0.0109), as individual fitness (lambda, generalised linear model, GLM, RNAi: t=-0.065, df=1, p=0.948; UV: t=-3.328, df=1, p=0.0104; RNAi x UV: t= 0.159, df=1, p=0.874) and as total lifetime reproduction (LRS) (GLM, RNAi: t=-1.887, df=1, p=0.0606; UV: t=-3.665, df=1, p<0.001; RNAi x UV: t= 0.066, df=1, p=0.948).

However, *daf-2* RNAi significantly increased offspring fitness both under benign conditions and when parents were UV-irradiated (lambda, GLM: RNAi: t=-2.305, df=1, p=0.0222; UV: t=-1.625, df=1, p=0.1059; RNAi x UV: t=- 0.772, df=1, p=0.441; Fig. 1b). This absence of fitness costs for the offspring of UV-irradiated parents, despite the costs of irradiation for parental reproduction, suggests that irradiated parents may have invested in germline repair mechanisms, which improved the fitness of their offspring[18,38,65].

Fig. 1| Adulthood *daf-2* RNAi in UV-irradiated and non-irradiated parents
increases parental lifespan and offspring fitness. a, Parental survival, n=50 per

treatment. **b,** Offspring fitness. UV irradiation status and adulthood RNAi treatment

(either *daf-2* RNAi, 'daf-2', or empty vector control, 'ev') of parents is indicated.

Offspring were maintained on empty vector and were not irradiated. Mean (effect

size) and 95% confidence intervals shown were derived using non-parametric

bootstrap resampling in 'dabestr' R package[58].

The impact of parental *daf-2* RNAi and UV treatment on offspring age-specific

reproduction varied with offspring age as did the number of offspring with zero

fecundity (zero-inflated generalised Poisson, ZIGP, RNAi x age: z=4.864, p<0.001;

UV x age: z=3.678, p<0.001; RNAi x UV: z=-0.175, p=0.861; ZI varied with age: z=-

0.227, p=0.00656 and age^2: z=3.301, p<0.001; Fig. S2). There was no effect of

parental treatment (neither *daf-2* RNAi, UV irradiation nor their interaction) on

offspring total lifetime reproduction (GLM, RNAi: t=-1.141, df=1, p=0.255; UV: t=-

1.626, df=1, p=0.106; RNAi x UV: t= -0.053, df=1, p=0.958; Fig. S2).

Our findings show that under adulthood-only *daf-2* RNAi, parents increase

investment into somatic maintenance resulting in increased lifespan with no survival

or reproductive cost to themselves or their offspring under benign conditions

validating previous work in *C. elegans*[19,20]. Importantly, we reveal that the absence of

a longevity-fecundity trade-off in parents persisted under stressful conditions, when

organisms have to invest into repairing UV-induced damage.

Furthermore, we show that *daf-2* knockdown in adult parents primarily

influenced the timing of reproduction in their offspring. Specifically, it caused a shift to

increased early life reproduction, improving offspring individual fitness, rather than an

increase in total reproduction. Whilst an earlier study found increased total

reproduction in the first generation of offspring from *C. elegans* parents treated with

daf-2 RNAi, across N2 wild-type and two other genetic backgrounds, this was also

accompanied by increased offspring fitness[20], in agreement with our results here.

Under UV-induced mutagenesis, the costs of germline maintenance would be

expected to become more apparent[29,38,66]. However we show that adulthood-only

daf-2 RNAi improves fitness of F1 offspring even when parents are under stress.

Multigenerational daf-2 RNAi in adulthood protects against extinction. We found

that daf-2 RNAi reduced extinction under both UV-induced and spontaneous MA

across 40 generations (Cox proportional hazards regression analysis, coxph, RNAi:

z=2.159, p=0.031; UV: z=11.469, p<0.001; RNAi x UV: z= -0.537, p=0.591; Fig. 2a).

Extinction results did not differ significantly between two independent experimental

blocks (coxph, z=-0.942, p=0.346; Fig. S3). The major causes of MA line extinction

(Fig. 2b) were infertility (failure to lay eggs), or sterility (the production of eggs that

did not hatch), indicative of the underlying germline damage. Infertility and sterility

were likely associated with observed reproductive abnormalities such as a deformed

vulva, abnormal external growth close to the vulva (possible tumour) and cavities in

the reproductive tract in place of embryos or oocytes (Fig. 2c). It is known that vulva-

less C. elegans mutants are unable to lay eggs and can die from internal

hatching[67,68].

It is possible that UV irradiation may also have caused non-genetic damage

(to proteins or other macromolecules) that was not inherited but may also have

contributed to sterility or infertility, and that daf-2 RNAi additionally may have

protected against. Despite this, previous work in C. elegans has shown that UV

325   irradiation most commonly induces base substitutions, which are subject to the

326   nucleotide excision repair pathway[69].

327   The reduction in extinction observed in our MA lines with *daf-2* RNAi shows

328   the benefits of increased investment into germline maintenance that became more

329   pronounced across multiple generations of UV-induced and spontaneous MA.

Fig. 2| Reduced adulthood IIS, via *daf-2* RNAi, protects against N2 wild-type

extinction under mutation accumulation. a, Transgenerational survival in N2

under spontaneous and UV-induced mutation accumulation. Vertical dotted line at

generation 20 indicates timing of life history assay. Sample size of 100 lines per

RNAi strain by irradiation treatment combination. **b,** Causes of extinction of N2 wild-

type MA lines indicate germline damage. **c,** Representative images of germline

damage. Brown scale bar of 0.99 mm for all images.

Multigenerational *daf-2* RNAi increased fitness of surviving MA lines. We next

tested for the effects of *daf-2* RNAi on life history traits of surviving MA lines. We

were interested in how IIS influences potentially detrimental effects of MA that are not

sufficiently severe to cause line extinction. We assayed age-specific reproduction,

egg size, male production and adult heat shock resistance in grand-offspring from the

spontaneous and UV-induced MA lines on *daf-2* RNAi versus control treatments, at

generation 20 of MA, following two generations of rearing under common garden

conditions (no irradiation, on empty vector control) to attenuate direct effects of

irradiation and RNAi. Generation 20 was a point at which there was a pronounced

benefit of *daf-2* RNAi for protection against extinction in the UV-induced MA lines, but

no clear difference in extinction trajectories for the spontaneous (non-irradiated) MA

lines (Fig. 2a).

Adulthood-only *daf-2* RNAi across 20 generations of mutation accumulation

significantly increased individual fitness in the surviving irradiated MA lines, but there

was no effect of *daf-2* RNAi on the fitness of non-irradiated lines (GLM, irradiated,

RNAi: $t=-2.804$, $p=0.006$; non-irradiated, RNAi: $t=0.647$, $p=0.518$; all data, RNAi x

UV: t=-2.886, p=0.004; Fig. 3a), in agreement with the results on line extinction at generation 20 (Fig. 2a).

The increased fitness of irradiated *daf-2* RNAi MA lines was driven by their improved early life fecundity (Day 1 and Day 2 offspring production), relative to irradiated controls (ZIGP, RNAi x age: z=-1.981, p=0.0475; RNAi x age^2: z=3.334, p<0.001; Fig. 3b), an effect absent in the non-irradiated lines (ZIGP, non-UV, RNAi: z=0.05, p=0.963; all data: RNAi x UV x age: z=-2.50, p=0.0125; RNAi x UV x age^2: z=3.58, p<0.001; ZI intercept: z=-18.27, p<0.001). There was no effect on total reproduction (GLM, UV: t=-0.811, p=0.419; non-UV: t=0.875, p=0.383; UV x RNAi: t=-1.138, p=0.256; Fig. S4).

We suggest that the fitness benefits of *daf-2* RNAi for irradiated MA lines were most likely due to genetic differences between the treatments, as the benefits persisted after two generations of common garden rearing and therefore were not due to the direct exposure of offspring to RNAi or UV (neither as adults nor as eggs). This conclusion is reinforced by the finding that parental *daf-2* RNAi effects on offspring fitness do not persist beyond F1 (see below, Fig. 5). Variation in fitness was considerably greater across irradiated than across non-irradiated MA lines (Fig. 3a), suggesting that irradiation-induced *de novo* mutations generated greater genetic and phenotypic variation compared to spontaneous MA. The spontaneous mutation rate in N2 wild type *C. elegans* is estimated as one *de novo* mutation per individual per generation, under standard conditions[69,70]. The close association between extinction trajectories during the first 20 generations of MA (Fig. 2a) and the fitness of the surviving MA lines (Fig. 3a) suggests that a threshold of accumulating deleterious

mutations needs to be crossed before the effect size is sufficiently severe to result in

extinction.

The most parsimonious explanation for the reduced extinction and increased

fitness of *daf-2* RNAi irradiated lines would therefore be due to genetic differences in

the rate of MA. To directly distinguish between genetic and non-genetic effects

associated with MA line extinction, future work could use whole genome sequencing

(WGS) to quantify mutation rates between the MA treatments.

Fig. 3| The effect of *daf-2* RNAi on fitness and age-specific reproduction after

20 generations of MA. a, Individual fitness (lambda) of the grand-offspring of N2

wild-type MA lines at generation 20. Fitness was assayed in a standard common

garden environment, on the empty vector control and no irradiation; following two

generations of rearing under standard conditions, from MA generation 20. Mean and

95% confidence intervals shown. **b,** Age-specific reproduction in the grand-offspring

of N2 MA lines at generation 20.

To determine if increased offspring fitness in the surviving *daf-2* RNAi-treated

MA lines, was associated with greater parental resource allocation into their eggs, we

measured egg size, as a proxy for parental investment (n=75 eggs measured, one

per individual taken from the Generation 20 fitness assay). Previous work has shown

that reduced IIS, either via dietary restriction or via *daf-2* RNAi, increases mean egg

size[20,56].

We found that grandparental *daf-2* RNAi resulted in grand-offspring (F2) that

laid smaller eggs if their grand-parents from the MA lines had been irradiated, but

there was no significant effect on the size of eggs laid by F2 offspring descended

from non-irradiated grandparents treated with *daf-2* RNAi following 20 generations of

MA (GLM, UV lines, RNAi: t=2.362, df=1, p= 0.0195; non-UV lines, RNAi: t=-1.246,

df=1, p=0.215; all data, RNAi x UV: t=2.642, df=1, p=0.00869; RNAi x UV x Block: t=-

0.774, df=1, p=0.440; Fig. 4). This is contrary to the increase in F1 egg size under

reduced parental IIS found in previous work under benign conditions[20,56].

Fig. 4| The effect of *daf-2* RNAi on egg size after 20 generations of spontaneous

or UV-induced MA. Egg size (area, mm^2) was reduced in the grand-offspring of UV-

induced MA lines on *daf-2* RNAi ('daf-2') relative to those from UV-irradiated MA lines

on empty vector control ('ev'). However, this difference in egg size was absent from

the spontaneous ('non-UV') MA lines, following 20 generations of MA. Grand-

offspring were all non-irradiated and kept on ev. Mean and 95% confidence intervals

are shown.

Our results indicate that even though F2 offspring from irradiated *daf-2* RNAi

MA lines laid smaller eggs, this did not seem to impact negatively on the quality

(fitness) of offspring they produced. It is possible that a smaller egg size could be a

phenotypically plastic response to UV radiation, perhaps to improve stress resistance

to irradiation. Alternatively, it could be the result of a trade-off between improved

investment into germline genetic quality and egg size. At present, it is unclear why

grandparental *daf-2* RNAi resulted in smaller eggs in F2.

Exposure to certain environmental stresses (e.g. high temperature[71],

starvation[72], increased mutation rate[73]), increases male production and outcrossing in

C. elegans. However, we found no increase in the proportion of males produced by

the grand-offspring in irradiated MA lines. Only one male developed from the 293

eggs assayed.

Reduced IIS increases the activity of heat shock factor 1 which mediates

lifespan extension[74,75]. Heat shock responses are conserved across diverse taxa and

positively associated with lifespan (as reviewed by[76,77]). To test the stress resistance

of post-reproductive (Day 7) adults, we assayed survival and locomotion under acute

heat shock for 1 hour and 45 minutes at 37°C (following[78]). We found no effect of MA

line treatment (neither UV irradiation nor *daf-2* RNAi) on the survival of untreated F2

offspring following heat shock. Only 4% of the 371 individuals were dead by 24 hours

after heat shock. However F2 offspring from non-irradiated *daf-2* RNAi MA lines

recovered normal locomotion faster following heat shock than F2 from non-irradiated

MA control lines, but this benefit was not seen in descendants from irradiated MA

lines (Binomial GLM, 3h post-heat shock, RNAi: z=-2.274, df=1, p=0.0229; UV: z=-

2.078, df=1, p=0.0377; RNAi x UV: z=2.304, df=1, p= 0.0212; 24h post-heat shock,

UV: z=-0.206, df=1, p=0.837; RNAi: z=-0.045, df=1, p=0.964; RNAi x UV: z=1.347,

df=1, p=0.178; Fig. S5). This is in line with previous work, which found survival after

heat shock to more than double with age, from the first to the fourth day of adulthood

in *C. elegans*[78].

Parental *daf-2* RNAi does not affect offspring fitness transgenerationally.

Transgenerational epigenetic inheritance of RNAi can occur in *C. elegans* via

transmission of small interfering RNAs[43,79]. This transgenerational inheritance of

RNAi can last for several generations[79-81] and requires the germline argonaut protein,

HRDE-1, that is absent in *hrde-1* (heritable RNAi defective- 1) mutants[82]. To

determine whether there was direct transgenerational transfer of *daf-2* RNAi via the

germline and thus for how many generations the effects of parental *daf-2* RNAi could

persist, we assayed the fitness effects of three generations of offspring (F1, F2 and

F3) from *daf-2* RNAi treated versus control parents. Two genetic backgrounds were

used: *C. elegans* N2 wild-types, and *hrde-1* mutants that did not transfer RNAi

transgenerationally.

Parental effects of *daf-2* RNAi on the fitness of descendants, were absent

after the first generation of offspring in the N2 wild-types and absent across all

generations of offspring in the *hrde-1* mutants (GLM, F1, RNAi x genotype: t=-2.160,

df=1, p=0.0329; F2, RNAi x genotype: t=0.372, df=1, p=0.711; RNAi: t=1.090, df=1,

p=0.278; genotype: t=7.044, df=1, p<0.001; F3, RNAi x genotype: t=- 0.973, df=1,

p=0.332; RNAi: t=0.359, df=1, p=0.721; genotype: t=5.302, df=1, p<0.001; all data:

RNAi x generation: F=4.450, df=2, p=0.0124; Fig. 5). Furthermore, there was no

effect of *daf-2* RNAi on age-specific reproduction after the first generation of

offspring, for either N2 or *hrde-1* backgrounds (Fig. S6; Table S2). There was also no

daf-2 RNAi effect on total reproduction for F2 and F3 offspring generations (Fig. S7;

Table S2).

Fig. 5| The effects of *daf-2* RNAi on offspring fitness do not persist beyond the first offspring generation. Fitness of the: **a,** first (F1); **b,** second (F2) and **c,** third (F3) generation of offspring from parents treated with *daf-2* RNAi ('daf-2') or an empty vector control ('ev'), in N2 wild-type and RNAi inheritance deficient *hrde-1* mutant backgrounds. All offspring generations were untreated (kept on ev).

The absence of fitness benefits in the second and third generations of offspring from *daf-2* RNAi parents strongly suggests that the life-history differences between the *daf-2* RNAi and control irradiated MA lines at generation 20 in the common garden experiment were due to genetic differences and not the direct effects of RNAi. These results also imply a lack of transgenerational inheritance of *daf-2* RNAi beyond the first generation of offspring. Interestingly, the heritable RNAi deficiency 1 (*hrde-1*) gene was required for the fitness benefits of parental *daf-2* RNAi in irradiated offspring, as these offspring fitness benefits were absent in the *hrde-1* mutant background. The *hrde-1* mutants had overall lower fitness and total reproduction than N2 wild-type.

hrde-1 is required for *daf-2* RNAi to confer germline protection under MA. *hrde-1* encodes an Argonaut protein that plays a key role in nuclear RNAi, RNAi inheritance and promoting germline immortality[83]. To determine if functional *hrde-1* was necessary for the protective effects of *daf-2* RNAi under spontaneous and UV-induced MA, we ran 400 MA lines in parallel to the N2 MA experiment, using the *C. elegans hrde-1* mutant background and reduced IIS, via *daf-2* RNAi, in half of the MA lines. The heritable RNAi deficiency (*hrde-1* mutant) resulted in the rapid extinction of

irradiated MA lines and the loss of the protective effects of *daf-2* RNAi under UV-

induced MA, across 25 generations (RNAi: z=-4.016, p<0.001; UV: z=12.370,

p<0.001; RNAi x UV: z=-1.758, p=0.079; Fig. 6). In fact, irradiated *daf-2* RNAi lines

went extinct faster than controls, in the *hrde-1* mutant background. This represented

a significant difference between *hrde-1* mutant and N2 wild type backgrounds in the

effect of *daf-2* RNAi (although not UV) on MA line extinction (RNAi x genotype:

z=4.480, p<0.001; genotype: z=-8.083, p<0.001; RNAi: z=-4.351, p<0.001; UV x

genotype: z=-0.487, p=0.626; UV: z=17.148, p<0.001).

The major cause of extinction in irradiated *hrde-1* mutant MA lines was

developmental arrest, followed by infertility (Fig. S8). This strongly suggests that

functional RNAi is required for *daf-2* RNAi protection against extinction, and for

normal development and reproduction under UV-induced MA.

Fig. 6| The effect of *daf-2* RNAi on *hrde-1* mutant multigenerational survival under mutation accumulation. Sample size of 100 MA lines for each RNAi by UV irradiation treatment combination.

We show the important role of *hrde-1* in the *daf-2* RNAi-mediated protection of irradiated offspring under mutation. Our results suggest that the interaction between UV-induced germline damage and a deficiency in transgenerational inheritance of RNAi reverses the protective benefits of *daf-2* RNAi. This supports previous work suggesting that *hrde-1* mutants suffer from progressive sterility under high temperatures implying their increased sensitivity to environmental stresses driven in part by defects in gametogenesis[82].

Multigenerational *daf-2* RNAi in adulthood downregulates *daf-2* expression in MA lines. We confirmed, using qRT-PCR analysis, that feeding nematodes bacteria that expresses double-stranded RNA for *daf-2* significantly downregulated *daf-2* expression. The extent of *daf-2* downregulation at generation 20 varied between the MA treatments (Table S3; Fig. S9). For N2 wild types, *daf-2* expression was downregulated by 23% in UV-induced MA lines and by 12% in spontaneous MA lines (based on mean gene expression fold change, $2^{-\Delta\Delta CT}$). For *hrde-1* mutants under spontaneous MA, *daf-2* expression was more subtly downregulated (by 6% fold change; Table S3; Fig. S9). We were unable to analyse data from UV-induced MA *hrde-1* mutant lines as they did not recover well from freezing (only n=1 MA line for the *daf-2* RNAi treatment; Fig. S9).

The degree of downregulation of *daf-2* expression between MA treatments, interestingly matched the different extent of life history divergence between *daf-2*

RNAi and empty vector treatments for the respective MA treatments at generation

20, in both N2 and *hrde-1* mutant extinction trajectories (Fig. 2a; Fig. 6) and in N2

fitness (Fig. 3a). The relatively modest degree of *daf-2* downregulation in the

spontaneous MA lines for N2, and particularly for *hrde-1* mutants, could be due to an

interaction between RNAi knockdown and genetic background, or a possible

'acclimatisation' to the multigenerational application of *daf-2* RNAi in adulthood, but

the exact reasons would require further investigation.

Technical replicates were highly repeatable for all samples (CV < 2.4% and

CV < 1.5% for 98% of samples). Biological replicates showed more variation in

relative gene expression within each MA line (mean CV= 7%; max. CV =8%), but

within the expected range for *C. elegans* based on previous qRT-PCR analyses for

different genes[84,85].

Conclusion

We found that reduced insulin signalling in adulthood, via *daf-2* RNAi, protects

against extinction under both UV-induced and spontaneous mutation accumulation in

C. elegans. Furthermore, the fitness of the surviving UV-irradiated MA lines was

higher under *daf-2* RNAi. Most extinctions occurred because of infertility, egg

hatching failure and developmental abnormalities suggesting that mutation

accumulation directly contributed to the observed differences between the RNAi

treatments. Future assessment of mutation rates in the MA lines could be useful to

distinguish between genetic and non-genetic effects on extinction. Germline

protection under *daf-2* RNAi requires nuclear argonaut *hrde-1* because fitness of

hrde-1; *daf-2* RNAi worms was reduced both in one-generation and in multi-generation experiments. This is in line with previous work suggesting that *hrde-1* is required for transgenerational inheritance and germline immortality[82,83].

We set out to test whether adulthood-only *daf-2* RNAi, known to extend lifespan without an obvious cost to parental fecundity[19,20], trades-off with detrimental fitness effects when applied across multiple generations. However, we found that multigenerational downregulation of *daf-2* via adulthood-only RNAi has positive effects on the germline and protects mutation accumulation lines from extinction. This was particularly so when germline mutation rate was increased by low-level UV radiation. Our results therefore suggest that while wild-type levels of *daf-2* expression promote development, growth and early-life reproduction, they are costly for late-life fitness and this cost accumulates across generations leading to lineage extinction.

Antagonistic pleiotropy theory of ageing (AP) maintains that genes that increase fitness in early life at the expense of fitness late in life can be overall beneficial for fitness and go to fixation[6]. In line with AP, downregulation of *daf-2* expression during development reduces fitness[19,41], but what is the physiological basis of the putative trade-off? Our findings that adulthood-only downregulation of *daf-2* expression protects both the germline and the soma under benign conditions and under UV-induced stress, argue against the idea that resource allocation underlies this trade-off. However, the results are in line with the hypothesis that selection optimises physiological processes for development and early-life reproduction but fails to optimise the regulation of gene expression associated with these same physiological process later in life[2,10,22-24]. This could be either because selection on insulin signalling in adulthood is too weak in *C. elegans*[86] or because of

constraints that prevent the evolution of age-specific gene expression[87,88]. Future

studies should focus on investigating the relative importance of trade-offs and

constraints in the evolution of ageing across taxa.

Data availability. The data will be available on FigShare upon publication.

Code availability. The codes for analyses will be available on GitHub upon

publication.

Author contributions

E.D., A.A.M. and H.C. conceived and designed the study, with T.C. and S.I. providing

additional advice. E.D., H.C. and K.S. performed the experiments and collected the

data, which E.D. and K.S. analysed. E.D. and A.A.M. wrote the manuscript, with

comments from all authors.

Acknowledgements

The authors thank M. Lind and E. Ivimey-Cook for statistical discussions, and T. van

Baalen, N. Edden and A. Dehem for experimental help. This study was funded by an

ERC Consolidator Grant (GermlineAgeingSoma/724909) to A.A.M.

Competing interests

The authors declare no competing interests.

References

1. Flatt, T. & Partridge, L. Horizons in the evolution of aging. *BMC Biol.* **16**, 93 (2018).
2. Maklakov, A. A. & Chapman, T. Evolution of ageing as a tangle of trade-offs: energy versus function. *Proc. R. Soc. B* **286**, 20191604 (2019).
3. Regan, J. C., Froy, H., Walling, C. A., Moatt, J. P. & Nussey, D. H. Dietary restriction and insulin-like signalling pathways as adaptive plasticity: a synthesis and re-evaluation. *Functional Ecology* **34**, 107-128 (2019).
4. Gaillard, J-M & Lemaître, J-F. An integrative view of senescence in nature. *Funct. Ecol.* **34,** 4–16 (2020).
5. Medawar, P. B. *An Unsolved Problem of Biology* (H. K. Lewis, 1952).
6. Williams, G. C. Pleiotropy, natural selection, and the evolution of senescence. *Evolution* **11**, 398–411 (1957).
7. Hamilton, W. D. The moulding of senescence by natural selection. *J. Theor. Biol.* **12**, 12-45 (1966).
8. Day, T. & Abrams, P. Density dependence, senescence, and Williams' hypothesis. *Trends Ecol. Evol.*, **35**, P300-302 (2020).
9. Caswell, H. & Shyu, E. in *The Evolution of Senescence in the Tree of Life* (eds. R. P. Shefferson, O.R. Jones, & R. Salguero-Gomez) 56-82 (Cambridge University Press, 2017).
10. Gems, D. & Partridge, L. Genetics of longevity in model organisms: debates and paradigm shifts. *Annu. Rev. Physiol.* **75,** 621-644 (2013).
11. Flatt, T. Life-history evolution and the genetics of fitness components in *Drosophila melanogaster*. *Genetics* **214,** 3-48 (2020).
12. Kirkwood, T. B. L. Evolution of ageing. *Nature* **270**, 301–304 (1977).
13. Kirkwood, T. B. L. & Holliday, R. The evolution of ageing and longevity. *Proc. R. Soc. B* **205**, 531–546 (1979).
14. Kirkwood, T. B. L. in *Evolution of Longevity in Animals: A Comparative Approach* (eds. A. D. Woodhead & K. H. Thompson). Immortality of the germ-line versus disposability of the soma 209-218 (Plenum Press, 1987).
15. Stearns, S. C. *The evolution of life histories* (Oxford Univ. Press, 1992).
16. Roff, D. *Life History Evolution* (Sinauer Associates, 2002).
17. Edward, D. A. & Chapman, T. in *Mechanisms of Life History Evolution: The Genetics and Physiology of Life History Traits and Trade-Offs* (eds. T. Flatt & A. Heyland). Mechanisms underlying reproductive trade-offs: Costs of reproduction 137-152 (Oxford Univ, Press, 2012).
18. Maklakov, A. A. & Immler, S. The expensive germline and the evolution of ageing. *Curr. Biol.* **26**, R577-R586 (2016).
19. Dillin, A., Crawford, D. K. & Kenyon, C. Timing requirement for insulin/IGF-1 signalling in *C. elegans*. *Science* **298**, 830-834 (2002).
20. Lind, M. I. L., Ravindran, S., Sekajova, Z., Carlsson, H., Hinas, A. & Maklakov, A. A. Experimentally reduced insulin/IGF-1 signalling in adulthood extends lifespan of parents and improves Darwinian fitness of their offspring. *Evol. Lett.* **4**, 737-744 (2019).

21. Flatt, T. Survival costs of reproduction in *Drosophila. Exp. Gerontol.* **46**, 369-375
(2011).
22. de Magalhaes, J. P. & Church, G. M. Genomes optimize reproduction: aging as a
consequence of the developmental program. *Physiology* **20**, 252-259 (2005).
23. de Magalhaes, J. P. Programmatic features of aging originating in development:
aging mechanisms beyond molecular damage? *FASEB J.* **26,** 4821-4826 (2012).
24. Blagosklonny, M. V. Ageing is not programmed. *Cell Cycle* **12**, 3736-3742 (2013).
25. Ezcurra, M. et al. *C. elegans* eats its own intestine to make yolk leading to
multiple senescent pathologies. *Curr. Biol.* **28**, 2544-2556 (2018).
26. Wang, H. et al. A parthenogenetic quasi-program causes teratoma-like tumors
during aging in wild-type *C. elegans. npj Aging Mech Dis* **4,** 6 (2018).
27. Sala, A. J., Bott, L. C., Brielmann, R. M. & Morimoto, R. I. Embryo integrity
regulates maternal proteostasis and stress resilience. *Genes Dev.* **34**, 1-10
(2020).
28. Carlsson, H. et al. Ageing as "early-life inertia": Disentangling life-history trade-
offs along a lifetime of an individual. *Evol. Lett.*, **5**, 551–564 (2021).
29. Berger, D., Stangberg, J., Grieshop, K., Martinossi-Allibert, I. & Arnqvist, G.
Temperature effects on life-history trade-offs, germline maintenance and mutation
rate under simulated climate warming. *Proc. R. Soc. B* **284,** 20171721 (2017).
30. Labbadia, J. & Morimoto R. I. Repression of the heat shock response is a
programmed event at the onset of reproduction. *Mol. Cell* **59**, 639–650 (2015).
31. Kenyon, C. J. The genetics of ageing. *Nature* **464**, 504-512 (2010).
32. Flatt, T. et al. *Drosophila* germ-line modulation of insulin signalling and lifespan.
PNAS 105, 6368-6373 (2008).
33. Hsin, H. & Kenyon C. Signals from the reproductive system regulate the lifespan
of *C. elegans. Nature* **399**, 362–366 (1999).
34. Arantes-Oliveira, N., Apfield, J., Dillin, A. & Kenyon, C. Regulation of life-span by
germ-line stem cells in *Caenorhabditis elegans. Science* **295**, 502-505 (2002).
35. Shemesh, N., Shai, N. & Ben-Zvi, A. Germline stem cell arrest inhibits the
collapse of somatic proteostasis early in Caenorhabditis elegans adulthood. *Aging
Cell* **12**, 814–822 (2013).
36. Ermolaeva, M. A. et al. DNA damage in germ cells induces immune response
triggering systemic stress resistance. *Nature*, **501**: 416-420 (2013).
37. Antebi, A. Regulation of longevity by the reproductive system. *Exp. Gerontol.* **48**,
596-602 (2013).
38. Chen, H-y., Jolly, C., Bublys, K., Marcu, D. & Immler, S. Trade-off between
somatic and germline repair in a vertebrate supports the expensive germ line
hypothesis. *PNAS* **117**, 8973-8979 (2020).
39. Thondamal, M., Witting, M., Schmitt-Kopplin, P. & Aguilaniu, H. Steroid hormone
signalling links reproduction to lifespan in dietary-restricted *Caenorhabditis
elegans. Nat Commun.* **5,** 4879 (2014).
40. Fontana, L., Partridge, L. & Longo, V. Extending healthy life span- from yeast to
humans. *Science* **328**, 321-326 (2010).
41. Chen, J. et al. A demographic analysis of the fitness cost of extended longevity in
Caenorhabditis elegans. *J. Gerontol A Biol Sci Med Sci.* **62**, 126-135 (2007).

42. Lionaki, E. & Tavernarakis, N. in *Cell senescence* (eds. L. Galluzzi, I. Vitale, O.
Kepp & G.Kroemer) High-throughput and longitudinal analysis of aging and
senescent decline in *Caenorhabditis elegans* 485–500 (Humana Press, 2013).
43. Fire, A. et al. Potent and specific genetic interference by double-stranded RNA in
Caenorhabditis elegans. Nature **391,** 806-811 (1998).
44. Timmons, L., Court, D. L. & Fire, A. 2001. Ingestion of bacterially expressed
dsRNAs can produce specific and potent genetic interference in *Caenorhabditis
elegans. Gene* **263**, 103–112 (2001).
45. Hyun et al. Longevity and resistance to stress correlate with DNA repair capacity
in *Caenorhabditis elegans. Nucleic Acids Res.* **36**, 1380-1389 (2008).
46. Boyd et al. Nucleotide excision repair genes are expressed at low levels and are
not detectably inducible in *Caenorhabditis elegans* somatic tissues, but their
function is required for normal adult life after UVC exposure. *Mutat. Res.* **683**, 57-
67.
47. Stergiou, L., Doukoumetzidis, K., Sendoel, A. & Hengartner, M. O. The nucleotide
excision repair pathway is required for UV-C-induced apoptosis in Caenorhabditis
elegans. *Cell Death Differ.* **14**, 1129-1138 (2007).
48. Meyer et al. Decline of nucleotide excision repair capacity in aging *Caenorhabditis
elegans. Genome Biol.* **8**, R70.
49. Vassilieva, L. L., Hook, A. M. & Lynch, M. The fitness effects of spontaneous
mutations in *Caenorhabditis elegans. Evolution* **54**, 1234-1246 (2000).
50. Keightley, P. D. & Bataillon, T. M. Multigeneration maximum-likelihood analysis
applied to mutation-accumulation experiments in *Caenorhabditis elegans.
Genetics* **154**, 1193-1201.
51. Ajie, B. C., Estes, S., Lynch, M., Phillips, P.C. Behavioral degradation under
mutation accumulation in *Caenorhabditis elegans. Genetics* **170**, 655-660 (2005).
52. Halligan, D. L. & Keightley, P.D. Spontaneous mutation accumulation studies in
evolutionary genetics. *Annu. Rev. Ecol. Evol. Syst.* **40**, 151-172 (2009).
53. Baer, C. F. et al. Comparative evolutionary genetics of spontaneous mutations
affecting fitness in rhabditid nematodes. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 5785-
5790 (2005).
54. Brommer, J. E., Merila, J. & Kokko, H. Reproductive timing and individual fitness.
Ecol. Lett. **5**, 802–810 (2002).
55. Lind, M. I., Zwoinska, M. K., Meurling, S., Carlsson, H. & Maklakov, A. A. 2016.
Sex-specific trade-offs with growth and fitness following lifespan extension by
rapamycin in an outcrossing nematode, *Caenorhabditis remanei. J. Gerontol. Ser.
A Biol. Sci. Med. Sci.* **71**, 882–890 (2016).
56. Hibshman, J. D., Hung, A. & Baugh, L. R. Maternal diet and insulin like signaling
control intergenerational plasticity of progeny size and starvation resistance.
PLoS Genet. **14**, e1007639 (2016).
57. R Core Team. R: A Language and Environment for Statistical Computing (R
Foundation for Statistical Computing, 2020).
58. Ho, J., Tumkaya, T., Aryal, S., Choi, H. & Claridge-Chang, A. Moving beyond p
values: everyday data analysis with estimation plots. *Nat. Methods* **16**, 565-566
(2019).
59. Therneau, T. M. coxme: Mixed Effects Cox Models. R package version 2.2-14
(2019).

60. Bolker, B. M. et al. Generalized linear mixed models: a practical guide for ecology and evolution. *TREE* **24**, 127–135 (2009).

61. Brooks, M. E. et al. glmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. *R Journal* **9**, 378–400 (2017).

62. Brooks, M. E. et al. Statistical modelling of patterns in annual reproductive rates. *Ecology* **100**, e02706 (2019).

63. Hartig, F. DHARMa: residual diagnostics for hierarchical (multi-level/mixed) regression models. R package version 0.2.0. https://cran.r-project.org/web/packages/DHARMa/vignettes/DHARMa.html (2020).

64. Bates, D., Machler, M., Bolker, B. M. & Walker, S. C. Fitting linear mixed-effects models using lme4. *J. Stat. Softw.* **67**, 1-48 (2015).

65. Smelick, C. & Ahmed, S. Achieving immortality in the *C. elegans* germline. *Ageing Res. Rev.* **4**, 67-82 (2005).

66. Sniegowski, P. D., Gerrish, P. J., Johnson, T. & Shaver, A. The evolution of mutation rates: separating causes from consequences. *BioEssays* **22**, 1057–1066 (2000).

67. Ferguson, E. L. & Horvitz, H. R. Identification and characterization of 22 genes that affect the vulval cell lineages of the nematode *Caenorhabditis elegans*. *Genetics* **110**, 17–72 (1985).

68. Sternberg, P. W. & Horvitz, H. R. The combined action of two intercellular signalling pathways specifies three cell fates during vulval induction in *C. elegans*. *Cell* **58**, 679–693 (1989).

69. Meier, B. et al. *C. elegans* whole-genome sequencing reveals mutational signatures related to carcinogens and DNA repair deficiency. *Genome Res.* **24**, 1624-1636 (2014).

70. Denver, D. R. et al. Variation in base-substitution mutation in experimental and natural lineages of *Caenorhabditis nematodes*. *Genome Biol Evol.* **4**, 513-522 (2012).

71. Nigon, V. & Dougherty, E. C. Reproductive patterns and attempts at reciprocal crossing of *Rhabditis elegans* Maupas, 1900, and *Rhabditis briggsae* Dougherty and Nigon, 1949 (Nematoda: Rhabditidae). *J. Exp. Zool.* **112**, 485–503 (1949).

72. Morran, L. T., Cappy, B. J., Anderson, J. L. & Phillips, P. C. Sexual partners for the stressed: facultative outcrossing in the self-fertilizing nematode *Caenorhabditis elegans*. *Evolution* **63**, 1473-1482 (2009).

73. Cutter, A. Mutation and the experimental evolution of outcrossing in *Caenorhabditis elegans*. *J. Evol. Biol.* **18**, 27-34 (2005).

74. Hsu, A. L., Murphy, C. T. & Kenyon, C. Regulation of aging and age-related disease by DAF-16 and heat-shock factor. *Science* **300**, 1142–1145 (2003).

75. Seo, K. et al. Heat shock factor 1 mediates the longevity conferred by inhibition of TOR and insulin/IGF-1 signaling pathways in *C. elegans. Aging Cell* **12**, 1073-1081 (2013).

76. Åkerfelt, M., Morimoto, R. I. & Sistonen, L. Heat shock factors: integrators of cell stress, development and lifespan. *Nat. Rev. Mol. Cell. Biol.* **11**, 545–555 (2010).

77. Muñoz M. J. Longevity and heat stress regulation in *Caenorhabditis elegans*. *Mech. Ageing Dev.* **124**, 43–48 (2003).

78. Zevian S. C. & Yanowitz, J. L. Methodological considerations for heat shock of the nematode *Caenorhabditis elegans*. *Methods* **68**, 450-457 (2014).

79. Posner, R. et al. Neuronal small RNAs control behaviour transgenerationally. *Cell*
177, 1814-1826 (2019).
80. Rechavi, O. et al. Starvation-induced transgenerational inheritance of small RNAs
in *C. elegans. Cell* **158**, 277-287.
81. Ni, J. Z. et al. A transgenerational role of the germline nuclear RNAi pathway in
repressing heat stress-induced transcriptional activation in *C. elegans. Epigenet.*
Chromatin **9**, 3.
82. Buckley, B. et al. A nuclear Argonaute promotes multigenerational epigenetic
inheritance and germline immortality. *Nature* **489,** 447–451 (2012).
83. Spracklin, G. et al. The RNAi inheritance machinery of *Caenorhabditis elegans.*
Genetics **206,** 1403-1416 (2017).
84. Chauve, L. et al. High-throughput quantitative RT-PCR in single and bulk *C.*
elegans samples using nanofluidic technology. *J. Vis. Exp.* **159**, e61132 (2020).
85. Lind, M. I. et al. Cost-free lifespan extension via optimization of gene expression
in adulthood aligns with the developmental theory of ageing. *Proc. Royal Soc. B*
288, 20201728 (2021).
86. Chen, J., Lewis, E. E., Carey, J. R., Caswell, H. & Caswell-Chen, E. P. The
ecology and biodemography of *Caenorhabditis elegans. Exp. Gerontol.* **41,** 1059-
1065 (2006).
87. Bonsall, M. B. Longevity and ageing: appraising the evolutionary consequences
of growing old. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **361,** 119-135 (2006).
88. Cohen, A. A., Coste, C. F. D., Li, X.-Y., Bourg, S. & Pavard, S. Are trade-offs
really the key drivers of ageing and life span? *Funct. Ecol.* **34,** 153-166 (2020).
802

803 **Supplementary Material**

805 **Supplementary Figures and Tables**

Supplementary Figure 1: No effect of *daf-2* RNAi in adulthood on parental reproduction. a, Age-specific reproduction. **b,** Fitness (lambda). **c,** Total reproduction. Mean and 95% confidence intervals shown.

Supplementary Figure 2: The effect of *daf-2* RNAi in UV-irradiated adult parents, on the reproduction of their offspring. a, Age-specific reproduction. **b,** Total reproduction. Parents were UV-irradiated or not and on *daf-2* RNAi ('daf-2') or

816   an empty vector control ('ev'). Parental *daf-2* RNAi affected offspring age-specific

817   reproduction (when parents were irradiated), but not offspring total reproduction.

820
Supplementary Figure 3: Transgenerational survival under mutation

accumulation in wild-type (N2) background, by block. a, Block 1. **b,** Block 2.

Sample size of 50 MA lines per treatment per block.

Supplementary Figure 4: No difference in total reproduction of surviving MA
lines at generation 20 in wild-type (N2) background. Total lifetime reproduction of
MA lines was calculated from the sum of daily reproduction in a standard common
garden environment, on the empty vector control and no irradiation; following two
generations of rearing under standard conditions, from MA generation 20.
Treatments of MA lines indicated.

Supplementary Figure 5: Effect of heat shock on locomotion of Day 7 adults in surviving MA lines at generation 20 in wild-type (N2) background. Locomotion (a measure of heat shock resistance) was recorded in the same individuals at 3 hours and 24 hours after the end of heat shock (37°C for 1 hour 45 minutes). Same individuals as used in the reproduction assay of MA lines from generation 20. Treatments of MA lines indicated (n= 93, 87, 84, 83 individuals per treatment, L-R respectively). The few individuals that died from heat shock were excluded from locomotion analysis.

Supplementary Figure 6: The effects of *daf-2* RNAi on offspring age-specific reproduction do not persist beyond the first offspring generation. Reproduction counts for the first (F1), second (F2) and third (F3) generation of offspring from parents treated with *daf-2* RNAi ('daf') or an empty vector control ('ev'), in N2 wild-type and RNAi inheritance

deficient *hrde-1* mutant backgrounds. All offspring generations were kept on ev. Sample
sizes: N2 (n=30), N2; *daf-2* RNAi (n=30), *hrde-1* (n=29) and *hrde-1*; *daf-2* RNAi (n=30).

Supplementary Figure 7: The effects of *daf-2* RNAi on offspring total lifetime
reproduction do not persist beyond the first offspring generation. Total lifetime
reproduction counts are the sum of age-specific reproduction counts for each individual, in
the first (F1), second (F2) and third (F3) generation of offspring from parents treated with *daf-*
2 RNAi ('daf-2') or an empty vector control ('ev'), in N2 wild-type and RNAi inheritance
deficient *hrde-1* mutant backgrounds. All offspring generations were untreated (kept on ev).

Supplementary Figure 8: a, Causes of extinction of mutation accumulation (MA) lines in *hrde-1* mutant background. b, Representative images showing impacts of germline damage in *hrde-1* mutant MA lines. Brown scale bar of 0.992 mm for all images.

Supplementary Figure 9. Normalised *daf-2* expression following *daf-2* RNAi
treatment versus untreated empty vector (ev) controls, from mutation
accumulation (MA) lines at generation 20. UV treatment indicates the background
of spontaneous (non-UV) or UV-induced (UV) MA from N2 and *hrde-1* mutant
genotypes. RNAi was delivered from late-L4 stage and gene expression quantified in
Day 2 adults using qRT-PCR. Arithmetic mean of biological replicates per MA line
shown as a red diamond with +/- 1 standard error bars. Each colour represents a
different MA line for each treatment combination, from which n=1-6 individual worms
(separate points) were assayed. Normalised *daf-2* expression ($2^{-\Delta CT}$) was calculated
relative to expression of the *actin-3* reference gene[89].

Supplementary Table 1: Sample sizes for gene expression assay. Total number of individual worms assayed for which N=1-6 individual worms were measured per MA line, for the indicated number (N) of MA lines per treatment: N2 or *hrde-1* mutant backgrounds, for UV-induced or spontaneous (non-UV) MA, on *daf-2* RNAi or empty vector (ev) control bacteria.

MA treatment	Total N	N, MA lines
N2 non-UV ev	15	5
N2 non-UV *daf-2* RNAi	16	5
N2 UV ev	18	6
N2 UV *daf-2* RNAi	24	8
hrde-1 mutant non-UV ev	22	10
hrde-1 mutant non-UV *daf-2* RNAi	25	10
hrde-1 mutant UV ev	8	6
hrde-1 mutant UV *daf-2* RNAi	4	1

886 **Supplementary Table 2a:** Full output of zero-inflated generalised Poisson model
887 used to analyse age-specific reproduction data from the first (F1), second (F2) and
888 third (F3) generation of offspring produced by parents treated with *daf-2* RNAi or an
889 empty vector control, in wild-type N2 or *hrde-1* mutant genotypes.

	z	p
F1 generation:		
RNAi x age	2.376	**0.0175***
RNAi x genotype	-1.584	0.113
genotype x age	2.627	**0.00861***
genotype x age^2	-4.257	**<0.001***
F2 generation:		
RNAi	0.137	0.891
RNAi x genotype	0.613	0.540
genotype x age	4.615	**<0.001***
genotype x age^2	-5.309	**<0.001***
F3 generation:		
RNAi	0.656	0.512
RNAi x genotype	-0.853	0.393
genotype x age	5.630	**<0.001***
genotype x age^2	-6.383	**<0.001***

892 **Supplementary Table 2b:** Full output of generalised linear model with Gaussian
893 error structure used to analyse total reproduction data from the first (F1), second (F2)
894 and third (F3) generation of offspring produced by parents from part **a**.

	t	df	p
F1 generation:			
RNAi x genotype	-2.888	1	**0.00464***
F2 generation:			
RNAi	0.286	1	0.775
genotype	3.787	1	**<0.001***
RNAi x genotype	1.309	1	0.193
F3 generation:			
RNAi	0.380	1	0.704
genotype	3.890	1	**<0.001***
RNAi x genotype	-0.746	1	0.457

Supplementary Table 3a: Relative gene expression (ΔCt) in N2 MA lines. The effect of *daf-2* RNAi from the late-L4 stage on the expression of *daf-2* relative to the *actin-3* reference gene (ΔCt from qRT-PCR) at day 2 of adulthood, after 20 generations of MA. RNAi and UV treatment contrasts from linear mixed effect model analysis of variance table.

Factor	Sum Sq	Mean Sq	F	d.f.	p
RNAi treatment	1.238	1.238	6.168	1	**0.001**
UV treatment	0.014	0.014	0.071	1	0.793
RNAi x UV treatment	0.021	0.021	0.107	1	0.747

Supplementary Table 3b: Relative gene expression (ΔCt) in *hrde-1* mutant MA lines. The effect of *daf-2* RNAi from the late-L4 stage on the expression of *daf-2* relative to the *actin-3* reference gene (ΔCt from qRT-PCR) at day 2 of adulthood. RNAi treatment contrast from linear mixed effect model analysis of variance table. Data for non-irradiated *hrde-1* mutant background, after 20 generations of MA.

Factor	Sum Sq	Mean Sq	F	d.f.	p
RNAi treatment	0.012	0.012	0.094	1	0.7618

Supplementary Table 3c. Relative gene expression (ΔCt) in combined analysis of N2 and *hrde-1* mutant MA lines. RNAi, UV and Genotype treatment contrasts and their interactions from linear mixed effect model analysis of variance table.

Factor	Sum Sq	Mean Sq	F	d.f.	p
RNAi treatment	0.414	0.414	2.333	1	0.134
UV treatment	0.007	0.007	0.038	1	0.846
Genotype	0.032	0.032	0.183	1	0.671
Genotype x RNAi	0.074	0.074	0.420	1	0.520
RNAi x UV	0.0002	0.0002	0.001	1	0.972

913 **Supplementary Table 4. Primer sequences**. Forward (fwd) and reverse (rev)
914 sequences are listed in the 5' to 3' direction. Sequences acquired from[84] for *daf-2*
915 and[90] for the *actin-3* reference gene.

Gene	Primer Sequences
daf-2	Fwd: GTGGCGTGAGAATGAAGTGAG
	Rev: GGCTTATCGGCTACAATCGTC
actin-3	Fwd: CCAAGAGAGGTATCCTTACCCTCAA
	Rev: AAGCTCATTGTAGAAGGTGTGATGC

                                    Supplementary Methods

We conducted four experiments to our test our main hypotheses about the effects of

reduced insulin/IGF-1 signalling (IIS) in adulthood, via adult-only *daf-2* RNAi, on the

soma and the germline:

1. Inter-generational effects of reduced IIS in parents under UV-induced stress, on

parental and offspring fitness and reproduction.

2. Effects of reduced IIS on 40 generations of spontaneous and UV-induced mutation

accumulation, in N2 wild-type and RNAi inheritance deficient (*hrde-1*) mutant

backgrounds.

3. Life history and fitness effects of 20 generations of spontaneous and UV-induced

mutation accumulation under reduced adulthood IIS, on the surviving MA lines.

4. Transgenerational effects of *daf-2* RNAi on offspring fitness in N2 and *hrde-1*

mutant backgrounds.

Nematode stocks and culture

All experimental lines were kept at 20°C, 60% relative humidity and in darkness,

consistent with standard *C. elegans* rearing protocol[91]. *C. elegans* is a valuable

model system due to its short life cycle, ease of genetic manipulation and its normal

reproductive state as self-fertilising hermaphrodites. Males were excluded from our

experiments and occur at a very low frequency of approximately 0.3% under benign

lab conditions and in nature[92,93].

During all experiments, worms were kept on 35mm NGM agar plates

(supplemented with 1mM IPTG and 50μg/mL of antibiotic ampicillin, to inhibit the

growth of bacteria other than our antibiotic resistant *E. coli*) and seeded with 0.1mL

of the e.v. control or *daf-2* RNAi bacteria, 24 to 48 hours before use, for *ad libitum*

bacterial growth. Bacterial cultures were prepared prior to the experiments, by

growing in LB supplemented with 50μg/mL ampicillin (as[94]).

Reducing IIS via *daf-2* RNAi feeding in adulthood

945   RNAi treatment was applied from the late-L4 stage, immediately prior to the onset of

946   adult sexual maturity and all individuals developed on empty vector (e.v.) control *E.*

947   *coli* prior to this stage. Normal development, which requires functional *daf-2*, was

948   therefore unaffected[19,31].

Validation of *daf-2* gene expression under RNAi

Worms were collected on day two of adulthood, to assay the downregulation of *daf-2*

in individuals at peak reproduction. To do this we picked individual worms onto

unseeded plates and allowed them to crawl around to remove surface bacteria and

separate day two adults from their eggs[84], and to ensure ingested bacteria was

excreted[95].

Individual cleaned worms were transferred into 10 μl of worm lysis buffer

(containing 1:100 diluted DNase; both from Ambion Power SYBR Green Cells-to-Ct

kit) in the separate domed lids of 0.2 ml PCR tubes, immediately spun down and

flash frozen in liquid nitrogen[84]. To crack the tough nematode cuticle and release

nucleic acids, we performed 10 freeze-thaw cycles, by transferring the PCR tubes

between liquid nitrogen and a 40°C, before homogenising the samples in a thermal

mixer (Eppendorf ThermoMixer C) set at 4°C, for 30 minutes at 1800 rpm[84]. We

confirmed, using Nanodrop spectrometry (Thermo Scientific) that it was possible to

obtain ~30-55 ng of RNA from our single worm samples (as[84,96]).

We reverse transcribed DNase-treated RNA using the Ambion Power SYBR

Green Cells-to-Ct kit, following the manufacturer protocol. We included a no reverse

transcriptase control (NRTC) per RNAi treatment, for which the RT enzyme was

substituted with nuclease-free water. The synthesised cDNA was used undiluted for

PCR and qRT-PCR.

To confirm that any contaminating genomic DNA had been removed, we

performed a standard PCR with a 10 µl reaction and an annealing temperature of

60°C. We ran 5 µl of the PCR reaction on a 1% agarose gel using ethidium bromide

and confirmed both the absence of amplification in the NRTCs verifying the

successful removal of any contaminating gDNA, and also the successful amplification

of cDNA for primer pairs.

The qRT-PCR was performed on an Applied Biosystems 7500 Real-Time PCR

System using the Power SYBR Green Cells-to-Ct kit (Ambion) with the following PCR

cycle: 95°C for 10 minutes, followed by 40 cycles of: 95°C for 15 s and 60°C for one

minute. The total reaction volume was 20 µl of which 4 µl was cDNA. We used

primers specific for the target gene of interest (*daf-2*) and for a reference gene- the

housekeeping gene, *actin-3* (T04C12.4), commonly used for *C. elegans*[90]. Primer

sequences are listed in Table S4. Primers were designed based on MIQUE

guidelines[97] and taken from[84] for *daf-2* and[90] for *actin-3.*

Two qRT-qPCR reactions (technical replicates) were carried out per sample

per primer pair, to check for repeatability and RNAi treatments were split evenly

across the six plates, to control for any minimal plate effects. We also included two

negative template controls (nuclease-free water substituted for cDNA) and one

NRTC per primer pair per plate, to test for any contamination.

To calculate relative gene expression, we determined ΔCt as the difference

between the qRT-PCR cycle thresholds (Ct values) of the target gene of interest (*daf-*

2) and the reference gene, for each sample. The arithmetic mean of the Ct values for

the two technical replicates per gene, per sample was used in ΔCt calculations.

Statistical analyses were performed on ΔCt[84], using a linear mixed effects model with

Gaussian error structure (lme4 package), to determine the effect of RNAi treatment

(*daf-2* RNAi versus empty vector controls), UV treatment and genotype (N2 or *hrde-1*

mutant) and their interaction on relative gene expression. MA line was fitted as a

random effect. The Shapiro-Wilk's normality test confirmed that ΔCt values satisfied

the normality assumption of the linear models (W=0.983, p=0.428).

The coefficient of variation (CV, %) in ΔCt between biological replicates for

each MA line, and in Ct values between technical replicates per gene, per sample,

was calculated as the standard deviation divided by the mean for each

comparison[84,98], to determine biological variation in relative gene expression between

individual worms and repeatability of the qPCR results, respectively.

To quantify fold change in gene expression ($2^{-\Delta\Delta CT}$), we calculated the difference in

the relative levels of mRNA for the target gene of interest compared to the reference

gene (ΔCt) between untreated controls and RNAi treated samples, using mean ΔCt

across all biological replicates per RNAi treatment[89].

Ultraviolet wavelength C (UV-C) irradiation

We calibrated the UV-C lamp of the Thermo Scientific Heraguard ECO Safety

Cabinet to determine the exposure time required to deliver an irradiation dose in the

range of previous work for *C. elegans*[45-48]. First, we measured the UV intensity (also

known as irradiance or fluence rate) of UV-C radiation emitted at a distance of 66cm

from the centre of the lamp to the surface on which our nematodes on plates were

placed for irradiation, as 2.3W/m^2, using a Macam spectroradiometer and verified

with a Sigma Laser Power Energy Meter (Coherent). As the radiation dose is the

product of intensity and exposure time (as approved by the International UV

Association), we calculated a required exposure time of 20 seconds.

The UV lamp was allowed to warm up and stabilise in temperature for 10

minutes, before irradiation of worms, as UV intensity increases with initial increases

in temperature of the UV lamp, but plateaus after at least five minutes[99].

Worms were irradiated on their *E. coli* seeded plates, without lids (plastic

would otherwise absorb UV irradiation) and positioned equidistant from the centre of

the lamp, in batches of maximum 30 plates, to ensure radiation dose did not differ

significantly between plates.

UV irradiation was timed at exactly 24 hours (+/- 30 minutes) after the RNAi

treatment began at the onset of adulthood and prior to peak reproduction, to allow

time for reduced IIS in individuals on *daf-2* RNAi[44,100]. This timing allowed us to

induce germline mutagenesis, as adult somatic tissue in *C. elegans* is post-mitotic

and very resistant to irradiation, whereas germline tissue (eggs, developing oocytes

and germline stem cells) is still actively dividing (undergoing meiosis and mitosis) and

so is more sensitive to UV irradiation[36]. As spermatogenesis is completed during the

late-L4 stage[101], it would not have been directly affected by irradiation.

Nematodes were transferred to new seeding immediately after UV irradiation,

in case the seeding or plate was affected by UV irradiation, and worms were placed

just outside the seeding on the new plate, to minimise contamination with mutated

bacteria (as[103,104]). UV-irradiated worms were allowed 8 hours (+/- 30 minutes)

recovery, to provide sufficient time for expulsion of irradiated embryos, before starting

egg laying[47,104]. Eggs laid were therefore most likely irradiated either as oocytes, or

as germline stem cells[101,104].

1039 Inter-generational effects of UV irradiation and reduced IIS in parents on

1040 parental and offspring fitness and reproduction

Wild-type N2 parents were either UV-irradiated or not as Day 2 adults and

maintained on *daf-2* RNAi or an empty vector control for the whole of adulthood, in

fully factorial design. Offspring were all non-irradiated and maintained on empty

vector throughout life. We assayed the daily offspring production from unmated,

singly-held hermaphrodite parents, for their entire reproductive period (first six days

of adulthood) and also from the first generation of their offspring, by daily transfers to

fresh plates. To assess parental survival, daily mortality checks were made, with

death being defined as no observed movement after gentle prodding. Worms were

grouped as ten worms per plate after the six-day reproductive period, for logistical

reasons and these groups were maintained as independent non-mixing units, for

daily transfers across lifetime so that plate identity could be included as a random

effect in analysis.

Mutation accumulation (MA) design

Each generation, for all treatments, we allowed mid-day 2 to mid-day 3 adults to lay

eggs onto empty vector plates, from which we picked a single late-L4 larva per line to

form the next generation of MA. The late-L4 stage is easily identifiable in *C. elegans*

(vulva cells are visible in the vulva lumen and prior to vulval protrusion at sexual

maturity), which allowed repeatable and consistent age-controlled set-up of each

generation. The duration of the egg lay period was optimised, to account for

differences in developmental timing between treatments. As such, even though the

development time was longer for eggs laid by parents from UV-irradiated MA lines

and for all treatments with increasing MA generation, we limited selection on

development time unevenly between treatments, as the next MA generation was set-

up at an identical late-L4 developmental stage for all treatments. Age of parental egg

lay was alternated between mid-Day 2 and mid-Day 3 every second generation, to

limit selection on parental age at reproduction and so the offspring forming each new

MA generation came from parents during their period of peak reproduction, when

most individuals were reproducing.

We recorded the generation at which extinction occurred and the cause of

extinction (death, failure to produce viable eggs or failure to reach sexually mature

adult). Individuals that were lost, underwent matricide (internal hatching that killed the

parent) prior to the Day 2 egg lay, died from the expulsion of internal tissue, became

infected or desiccated on the plate wall were censored. We photographed a sample

of Day 1 adult worms with visible developmental or reproductive abnormalities under

light microscopy (camera specifications in egg size measurement section below),

including worms with stunted growth, apparent aberrant external growth (tumour),

deformed vulva, or abnormal cavities in the reproductive tract in place of oocytes or

embryos.

Effect of 20 generations of spontaneous and UV-induced mutation accumulation on fitness and life history of surviving MA lines

Prior to the life history assay, individuals taken from MA lines at generation 20 were

reared for two subsequent generations under common garden conditions (no

irradiation, on empty vector control), to reduce direct parental effects from exposure

to RNAi and UV, and hence look for genetic differences between MA treatments. The

life history assay was conducted in the common garden environment.

Age-specific reproduction, fitness and total lifetime reproduction assays

The first 4 days of adult reproduction were used to calculate fitness and to analyse

age-specific reproduction, as day 5 and 6 reproduction was zero for almost all

individuals. Parental fitness and age-specific reproduction were analysed for day 2 to

4 inclusive, for the first intergenerational experiment, as UV-irradiation was

administered at the start of day 2 of adulthood.

Egg size measurement

Only eggs laid at the normal gastrula (approximately 30-cell) stage of development

were photographed, as egg shape and size can vary after 9 hours of ex-utero

development. Egg size (area in mm^2 enclosed within a free-drawn ellipse around the

egg perimeter) was calculated from photos using *Image J Fiji program* v.1.51[105],

whilst blind to treatment identity.

To minimise experimenter error, each egg size measurement was taken twice

and an average taken, and the same experimenter measured all eggs. Egg

measurement was completed across three measurement days, using identical

ImageJ settings and scale calibration. Treatments and blocks were stratified across

measurement days and randomly dispersed within each day, to avoid bias. Egg area

measurements were strongly repeatable between replicate measurements on the

same egg (Spearman's rank correlation coefficient, rho= 0.899), there was no

temporal autocorrelation between measurement days (Kruskal-Wallis rank sum test,

χ^2=299, df=314, p=0.721) and only weak temporal autocorrelation on the first

measurement day (Spearman's rank, rho=0.398).

Male production

We determined the percentage of eggs that developed into adult males (n= 75 eggs

assayed/treatment), one egg per individual Day 2 worm, as above, to determine if the

stress induced by irradiation or mutation accumulation would increase male

production above standard N2 wild-type levels, as seen previously for heat stress

and starvation in *C. elegans*[71,72].

Adult heat shock resistance

To test the stress resistance of post-reproductive (Day 7) adults, we assayed survival

following acute heat shock of 1 hour and 45 minutes at 37°C (following[78]), in the

same individuals used for the Generation 20 age-specific reproduction assay. At 3

and 24 hours post-heat shock we recorded any individuals that had died and

categorised locomotion as normal movement (crawling spontaneously or when gently

prodded), uncoordinated movement of head and tail with no forward or backward

trajectory, or just head movements; following[106,107]. Previous work confirmed that

from 12 hours after heat shock is a reliable time point to score long-term survival in

C. elegans adults[78]. All worms were heat-shocked simultaneously, grouped by

treatment, five worms per plate and the positions of plates in the oven were stratified

by treatment, to control for any positional differences in heat exposure.

Statistical analyses

The 'dabest' Cummings estimation plots display all datapoints from each treatment

as a swarm-plot, with the mean ± standard deviation of treatment as a gapped line

adjacent to the points. The mean difference (effect size) and its 95% confidence

interval (CI) is estimated for pairwise comparisons of treatments via non-parametric

bootstrap resampling (n=5000) and displayed below the main plots, for visualisation.

When the 95% CI vertical bar does not cross the x-axis (mean difference equal to

zero), there is a significant difference between the pairwise treatment comparison.

Individual fitness (lambda) incorporates age-specific reproduction, with emphasis on

early reproduction and fast development. It is therefore an appropriate measure

when timing is important for fitness[54]. Individual fitness was calculated from the life

table of age-specific reproduction, by solving the Euler-Lotka equation using the

lambda function in the popbio package in R (as[20,85]). The life table was constructed

using a common development time of two days without reproduction. Worms

matured and reproduction began during day three (first day of adulthood) when fast-

developing worms would have matured earlier and have had more reproduction on

this day. Since lambda is a rate-sensitive measure, fast development, resulting in

increased early life reproduction, would translate into higher values of lambda.

Supplementary References

89. Schmittgen, T. D. & Livak, K. J. Analyzing real-time PCR data by the comparative C-T method. *Nat. Protoc.* **3**, 1101-1108 (2008).
90. Akay, A. et al. The helicase Aquarius/EMB-4 is required to overcome intronic barriers to allow nuclear RNAi pathways to heritably silence transcription. *Dev. Cell.* **42**, 241-255 (2017).
91. Brenner S. The genetics of Caenorhabditis elegans. *Genetics* **77**, 71-94 (1974).
92. Teotónio, H., Manoel, D. & Phillips, P. C. Genetic variation for outcrossing among *Caenorhabditis elegans* isolates. *Evolution* **60**, 1300-1305 (2006).
93. Barriere, A. & Felix, M.-A. High local genetic diversity and low outcrossing rate in *Caenorhabditis elegans* natural populations. *Curr. Biol.* **15**, 1176-1184 (2005).
94. Hinas, A., Wright, A. J. & Hunter, C. P. SID-5 is an endosome-associated protein required for efficient systemic RNAi in *C. elegans*. *Curr. Biol.* **22**, 1938–1943 (2012).
95. Ghafouri, S. & McGhee, J. D. Bacterial residence time in the intestine of *Caenorhabditis elegans*. *Nematology* **9**, 87-91 (2007).
96. Ly, K., Reid, S. J. & Snell, R. G. Rapid RNA analysis of individual *Caenorhabditis elegans*. *Methodsx* **2**, 59-63 (2015).
97. Bustin, S. A. et al. The MIQE Guidelines: minimum information for publication of quantitative real-time PCR experiments. *Clin. Chem.* **55**, 611-622 (2009).
98. Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acids Res.* **29**, e45 (2001).
99. Lawal, O. et al. Method for the measurement of the output of monochromatic (254nm) low-pressure UV lamps. *IUVA News* **19**, 9-16 (2017).
100. Conte Jr, D., MacNeil, L. T., Walhout, A. J. M. & Mello, C. C. RNA interference in *Caenorhabditis elegans*. *Curr. Protoc. Mol. Biol.* **109**, 1-30 (2015).
101. Kimble, J. E. & White, J. G. On the control of germ cell development in *Caenorhabditis elegans*. *Dev. Biol.* **81**, 208–219 (1981).
102. Stewart, H. I., Rosenbluth, R. E. & Baillie, D. L. Most ultraviolet irradiation induced mutations in the nematode *Caenorhabditis elegans* are chromosomal rearrangements. *Mutat. Res.* **249**, 37-54 (1991).
103. Coohill, T., Marshall, T., Schubert, W. & Nelson G. Ultraviolet mutagenesis of radiation-sensitive (rad) mutants of the nematode *Caenorhabditis elegans*. *Mutat. Res.* **209**, 99-106 (1988).
104. Sakashita, T. et al. Radiation biology of *Caenorhabditis elegans*: germ cell response, aging and behaviour. *J. Radiat. Res.* **51**, 107-121 (2010).

105.Schindelin J. et al. Fiji: an open-source platform for biological-image analysis.
Nat. Methods. **9**, 676-682 (2012).
106.Herndon, L. A. et al. Stochastic and genetic factors influence tissue-specific
decline in ageing *C. elegans. Nature* **419**, 808-814 (2002).
107.Jovic, K. et al. Temporal dynamics of gene expression in heat-stressed
Caenorhabditis elegans. PLoS ONE **12**, e018944 (2017).
1189