What evolutionary processes maintain MHCIIβ diversity within and among

populations of stickleback?

Foen Peng[1], Kimberly M. Ballare[2], S. Hollis Woodard[3], Stijn den Haan[4], Daniel I.

Bolnick[1]

[1] Department of Ecology and Evolutionary Biology, University of Connecticut, Storrs CT,

USA

[2] Department of Ecology and Evolutionary Biology, University of California, Santa Cruz

CA, USA

[3] Department of Entomology, University of California, Riverside CA, USA

[4] Central European University, Budapest, Hungary

Correspondence:

Daniel I. Bolnick

Email: daniel.bolnick@uconn.edu

Abstract

Major Histocompatibility Complex (MHC) genes encode for proteins that recognize

foreign protein antigens to initiate T-cell mediated adaptive immune responses. They are

often the most polymorphic genes in vertebrate genomes. How evolution maintains this

diversity is still an unsettled issue. Three main hypotheses seek to explain the

maintenance of MHC diversity by invoking pathogen-mediated selection: heterozygote

advantage, frequency-dependent selection, and fluctuating selection across landscapes

or through time. Here, we use a large-scale field parasite survey in a stickleback

metapopulation to test predictions derived from each of these hypotheses. We identify

over a thousand MHCIIβ alleles and find that many of them covary positively or

negatively with parasite load, suggesting that these genes contribute to resistance or

susceptibility. However, despite our large sample-size, we find no evidence for the

widely-cited stabilizing selection on MHC heterozygosity, in which individuals with an

intermediate number of MHC alleles have the lowest parasite burden. Nor do we

observe a rare-allele advantage, or widespread fluctuating selection across populations.

In contrast, we find that MHC diversity is best predicted by neutral genome-wide

heterozygosity and between-population genomic divergence, suggesting neutral

processes are important in shaping the pattern of metapopulation MHC diversity. Thus,

although MHCIIβ is highly diverse and relevant to the type and intensity of macroparasite

infection in these populations of stickleback, the main models of MHC evolution still

provide little explanatory power in this system.

Introduction

Genetic variation is the raw material for natural selection to act upon. Hence, there is a long history of evolutionary studies on how genetic variation is maintained in natural populations (Dobzhansky, 1982), either invoking selection or neutral processes. One of the most dramatic cases of genetic variation concerns polymorphism at Major Histocompatibility Complex (MHC) genes, which are commonly the most variable genes in vertebrate genomes (Sommer, 2005). Surprisingly, despite decades of research on MHC evolution, the evolutionary processes sustaining MHC polymorphism remain unclear, with inconsistent and often ambiguous support for competing hypotheses (Radwan, Babik, Kaufman, Lenz, & Winternitz, 2020). Here, we evaluate predictions of multiple competing MHC evolution models (adaptive and neutral), using an exceptionally large dataset of genetic variation and macroparasite infection in a metapopulation of threespine stickleback (*Gasterosteus aculeatus*) inhabiting lakes on Vancouver Island.

MHC genes play a key role in the adaptive immune system of vertebrates. The main function of these genes is to encode for cell surface proteins, which are used to detect foreign molecules. These molecules are then presented to T cells to initiate appropriate immune responses. The MHC gene family consists of two classes of genes, MHC I and MHC II, each of which can be represented by multiple paralog copies. MHC class I genes are expressed in all nucleated cells and typically bind to peptides derived from intracellular molecules, such as virus protein (Jensen, 2007). MHC class II genes (our focus here) are expressed only in antigen presenting cells, such as macrophages, dendritic cells, and B cells. These cells phagocytize extracellular material, digest the proteins, and if the MHC II proteins bind to the resulting fragments, these are presented on the cell surface to T cells, to possibly initiate an adaptive immune response (Jensen,

2007). The second exon of MHC II gene is often used to characterize the variability of

MHC II genes, because it contains the antigen-binding domain (Sommer, 2005).

The maintenance of MHC diversity is often attributed to pathogen-mediated selection,

although intraspecific processes, such as mate choice (Penn & Potts, 1999), are also

potential sources of selection. As reviewed in Spurgin & Richardson (2010), there are

three non-mutually exclusive pathogen-mediated selection mechanisms to explain MHC

diversity: (1). *Heterozygote advantage.* Individuals with more MHC alleles might be able

to recognize a more diverse set of parasite species, thus have fewer parasites. This

advantage may lead to directional selection for ever-increasing diversity, but some

widely-cited studies have presented evidence for intermediate optima (K. Mathias

Wegner, Kalbe, Kurtz, Reusch, & Milinski, 2003). (2). *Negative frequency-dependent*

selection. In a given population, common MHC alleles that protect against parasites

impose selection on those parasites to change their antigens and evade recognition.

Once parasites evolve strategies to evade these common alleles, individuals carrying

the common alleles may be at a selective disadvantage. By contrast, a rare MHC allele

imposes little selection on the parasites it recognizes (because it is rare), and so may

tend to be more protective. For this negative frequency-dependence to work in the long

run, it must on average hold that rare alleles are more protective (and hence more fit)

than common alleles. (3) *Fluctuating selection.* Parasite community and parasite

genotypes may vary in space and time. Given this, different host MHC alleles may be

selectively favored in different locations and at different times. Gene flow between

locations subject to divergent selection can sustain MHC diversity, as can temporally

fluctuating selection.

Although many studies have examined the roles of the above-mentioned mechanisms in

maintaining MHC diversity, the results are often mixed or even contradictory. For

example, heterozygote advantage was supported in one experimental infection study

with inbred mice (Penn, Damjanovich, & Potts, 2002), but the effect was not found in

another study with outbred mice (Ilmonen et al., 2007). Theoretically, negative frequency

dependent selection is more likely to explain the extraordinary diversity in MHC genes

than heterozygote advantage (Borghans, Beltman, & De Boer, 2004). However,

empirically demonstrating negative frequency dependent selection is more challenging,

as the pattern is often also consistent with other mechanisms. For example, in house

sparrows, population-specific MHC alleles were linked to host resistance or susceptibility

to malaria (Bonneaud, Pérez☐Tris, Federici, Chastel, & Sorci, 2006), but this pattern

could be explained either by negative frequency dependent selection or spatially varying

selection. A previous study of stickleback from three lake-stream population pairs (Stutz

& Bolnick, 2017) tried to distinguish between negative frequency dependent selection or

spatially varying selection. The study took advantage of migration between parapatric

populations with different parasite communities, which in principle would allow them to

partition benefits of rarity within populations, from costs of being a (rare) immigrant to a

foreign habitat. Yet, no overall trend was found to support either mechanism. Such

inconsistent or ambiguous results are typical in the MHC evolution literature. Therefore,

a consensus regarding the relative importance of these mechanisms in maintaining MHC

diversity has not been reached (Radwan et al., 2020).

In contrast to these adaptive hypotheses, whether and how MHC diversity is shaped by

neutral processes is less studied. In small and/or isolated populations, MHC diversity is

often heavily influenced by genetic drift, rather than balancing selection (Hedrick, Parker,

& Lee, 2001; Miller & Lambert, 2004; Seddon & Ellegren, 2004). In large

metapopulations, it is unclear what role neutral processes play in shaping MHC diversity.

MHC studies in threespine stickleback have generated some unique insights in the past

(Eizaguirre, Lenz, Kalbe, & Milinski, 2012; Matthews, Harmon, M'Gonigle, Marchinko, &

Schaschl, 2010; Mccairns, Bourget, & Bernatchez, 2011; Milinski et al., 2005; K. Mathias

Wegner et al., 2003). For example, Wegner et al. (2003) conducted a study on 8

stickleback populations to examine the association between MHC diversity and 15

parasite species. They found populations exposed to a wider range of parasites had

higher MHC allelic diversity, though MHC diversity also positively, albeit weakly,

correlated with neutral diversity. Moreover, at the individual level, parasite burden was

minimized in fish with an intermediate, rather than maximal, number of MHCIIβ alleles.

This result supports the theoretical model of stabilizing selection on MHC heterogeneity

(Nowak, Tarczy-Hornoch, & Austyn, 1992). The logic is that low MHC-diversity

individuals do not have the capacity to recognize diverse parasites, but high MHC-

diversity individuals deplete their T cells via negative selection on self-reactive T cells.

The resulting paucity of T cell receptor diversity also limits their ability to recognize

diverse parasites. However, this intermediate-advantage hypothesis, while widely cited,

has not been replicated in many systems and remains controversial (Borghans, Noest, &

De Boer, 2003). Another example of interesting insights from stickleback is the

involvement of MHC alleles in mate recognition. It was found that female stickleback

could choose their mate based on odor to optimize the number of MHC alleles in their

offspring, though the molecular mechanism is unknown (Milinski et al, 2005).

In this study, we set out to test predictions associated with three main models of MHC

polymorphism invoking parasite-mediated selection, using an extensive field survey of

parasites in a metapopulation of threespine stickleback. We did not find evidence for

stabilizing or positive selection for MHC heterozygosity, contrary to Wegner et al. (2003).

Nor did we observe fluctuating selection across populations. Although we found

significant associations between specific MHC alleles and parasite species both within

and across populations, neutral processes in our dataset best explained within- and

between-population MHC diversity in our data set.

Materials and methods

1. Fish sampling and parasite identification

We used unbaited minnow traps to collect adult threespine stickleback from lake, river,

and estuary sites on Vancouver Island in late May and early June 2009. Collections

were approved by the University of Texas IACUC (07-032201) and a Scientific Fish

Collection Permit from the Ministry of the Environment of British Columbia (NA07-

32612). Fish were euthanized in MS-222 and then preserved in formalin, after the caudal

fin of each individual fish was removed and preserved in pure ethanol for genotyping.

Details are provided in Bolnick & Ballare (2020). These samples were used to examine

within- and between-population variation in diet (Bolnick & Ballare, 2020), parasite

community composition (Bolnick, Resetarits, Ballare, Stuart, & Stutz, 2019), and parasite

species richness (Bolnick, Resetarits, Ballare, Stuart, & Stutz, 2020). The parasite

infection data for this study are archived at Dryad Digital Repository:

https://datadryad.org/stash/dataset/doi:10.5061/dryad.gmsbcc2j1. Here, we focus on a

random subset of 26 sampling sites (N = 1437 stickleback) for which we also genotyped

individuals for MHC IIβ. For each stickleback individual, we recorded sex and body

length and then dissected each fish to count and identify macroparasites as described in

(Stutz & Bolnick, 2017).

2. Genotyping

The methods to sequence and genotype MHC IIβ alleles were identical to Stutz & Bolnick (2017). Briefly, genomic DNA was extracted from fin clips using a Promega Wizard 96-well extraction kit. We used PCR to amplify the second exon of MHC IIβ genes in each fish, with primers and PCR cycles as described in Stutz & Bolnick (2017). This exon contains the hypervariable peptide-binding region that binds to possible parasite antigens (Sommer, 2005). Each specimen was barcoded with a unique combination of forward and reverse primer tags for multiplexing. We used Quant-iT PicoGreen kits (Invitrogen P11496) to quantify DNA concentrations of magnetic bead-purified (Agencourt AMPure XP beads) PCR products, then pooled up to 400 samples in equimolar amounts to construct a library. We used Illumina Mi-Seq to sequence these multiplexed amplicon libraries. Then, we used a Stepwise Threshold Clustering (STC) program (Stutz & Bolnick, 2014), implemented in the AmpliSaS web software (URL: http://evobiolab.biol.amu.edu.pl/amplisat/index.php?amplisas; Sebastian, Herdegen, Migalska, & Radwan, 2016) to distinguish real sequence variants from sequencing error or PCR chimeras. The algorithm was originally validated by sequencing cloned amplicon products (Stutz & Bolnick, 2017). The software outputs a table of individual fish (rows) and unique MHC sequences (columns) with read depths.

To efficiently process data in AmpliSaS, we set the upper limit of read depth for each individual as 5000, which was sufficient to retrieve all the possible MHC alleles (Stutz & Bolnick, 2014). Because low sequencing coverage could bias the number of MHC alleles to be identified, individual fishes with coverage lower than 450 were excluded from this study (Fig. S1). After excluding the 160 individuals with low coverage (leaving N = 1277 individuals in the subsequent analyses), there was no longer a significant linear relationship between sequencing coverage and the number of MHC alleles ($t = 1.46$, $p = $

0.14). The number of unique MHC alleles was inferred based on the translated protein

sequences, thus merging distinct exonic sequences that produce identical amino acid

sequences.

Figure S1. A. The relationship between sequencing depth and the number of MHC

alleles. Each dot represents a fish sample. B. The distribution of sequencing depth

among samples. Individuals with sequencing depth < 450 were excluded from analysis.

3. Data analysis

a. Heterozygote advantage

We tested whether MHC diversity is associated with parasite burden at both the

individual and the population level. At the individual level, we used parasite richness

(*i.e.,* the number of parasite species) to measure parasite burden. Given that MHC

genes locate in many genomic loci and often act in a dominant manner in parasite

resistance, we used allelic richness (*i.e.,* the total number of unique MHC alleles found in

an individual) in this study to evaluate the level of MHC heterozygosity. We used a

mixed effect generalized linear model with Poisson distribution to evaluate the impact of

MHC alleles on parasite richness. The fixed effects are the number of MHC alleles and

log body length, which is known to affect parasite richness in stickleback (Bolnick et al.,

2020) and in other fish species (Calhoun, McDevitt-Galles, & Johnson, 2018). We did

not include sex as a main effect because we found no main effect of sex on parasite

richness in a previous analysis of this dataset (Bolnick et al., 2020). Sample site was

treated as a random effect. In addition, we also evaluated the optimizing hypothesis (*i.e.*,

MHC allelic richness is optimized at an intermediate level, driven by the negative

selection pressure to decrease T cell depletion by self-recognition) by adding a quadratic

MHC diversity term into the mixed effect model. We considered the interaction between

site and the linear and quadratic effects of MHC, using a random slope effect across

sites. Statistical support for the model terms was evaluated by AIC.

At the population level, we calculated the mean value of parasite richness and the

average number of MHC alleles (per fish) in each site. We used a linear regression

model to test if the mean value of parasite richness is associated with the average

number of MHC alleles for each site.

b1. Frequency-dependent selection: is there a rare allele advantage?

If MHC alleles were under negative frequency dependent selection, then on average

rare alleles must confer an advantage (in the form of lower parasite infection) compared

to common alleles. In contrast, MHC alleles with high frequency would be less effective

in parasite detection, because parasites would be evolving strategies to avoid detection

by those alleles. We therefore expect a positive relationship between the frequency of an

allele and its overall effect on parasite infection, where a negative effect denotes

protection and a positive effect suggests susceptibility or survivor's bias. Because MHC

II genes scatter throughout multiple loci in fish genome (Kaufman, 2018), it is difficult to

use allele frequency to estimate the abundance of MHC allele in the population. We

used allele prevalence instead, which was calculated as the percentage of individuals

carrying a focal allele in a population. Note that allele prevalence had different properties

from allele frequency; for example, allele prevalence values of all the alleles in a

population do not sum to 1. Similarly, we also calculated parasite prevalence in a

population as the percentage of individuals infected by a focal parasite. Alleles and

parasites that are too rare (<0.05 prevalence) or too common (>0.95 prevalence) do not

provide sufficient variance to estimate effects. For every moderately prevalent MHC -

parasite combination (both prevalence variable ranging from 0.05 to 0.95) in every

sampling site, we used a generalized linear model with negative binomial distribution to

test if the presence/absence of the focal MHC allele influenced the infection intensity of

the focal parasite in that stickleback population. We corrected p values for multiple-

comparison with BH method. The Z value of the regression models indicated the impact

of the particular MHC allele on the infection rate by a particular parasite. We excluded

the models disproportionately influenced by extreme values, *i.e.* the absolute Z value of

a model would change over 0.5 if excluding the largest data point from the model. After

iterating this procedure for all qualifying MHC-parasite combinations, we used another

mixed-effect linear regression model to examine if the estimated effect sizes (Z) of MHC

alleles were influenced by local allele prevalence. In this model, we treated both

sampling site and focal parasite as random effects.

b2. Frequency-dependent selection: are allele effects inconsistent across lakes?

If stickleback and their parasites engage in a Red Queen race style co-evolution, the

efficacy of any single allele will shift through time. For hosts and parasites with limited

dispersal, physically disconnected sites are unlikely to be in the same phase of the arms

race. As a result, a given MHC allele may have different effects on a given parasite from

one site to the next: effective against defense in some places/times, ineffective or

susceptible at others. Alternatively, if the same allele has similar effects on the same

parasite across different lakes (without gene flow), such fluctuating frequency-dependent

selection is unlikely. To test these alternatives, we first identified the moderately

prevalent MHC-parasite combinations that were present in more than one site. For each

qualified combination, we used a generalized linear model with negative binomial

distribution to examine if the infection intensity of the parasite is influenced by the MHC

allele, sampling site, and the interaction between site and MHC allele (this differs from

the GLMs described in b1, which were done separately for each sample site). We

corrected p values for multiple-comparison with BH method. We used the *anova* function

in R to perform an analysis of deviance for each regression model. It reported the

reductions in the residual deviance as each term of the formula was added in turn. We

evaluated whether, across many parasite-allele combinations, more variation was

explained by the focal allele's main effect (implying consistent protection across

populations), or allele × population interactions (inconsistent protection). Parasites

transmitted by birds could spread in a larger spatial scale, so they are less likely to be

engaged in evolutionary arms race. We did a Chi-squared test to test whether the

parasite taxa, which have birds as final hosts, were more likely to be found in the models

with significant main effect than in the models with significant allele x population effect.

c. Fluctuating selection

The parasite communities of stickleback differ significantly among the different sites on

Vancouver Island (Bolnick et al., 2020; Stutz & Bolnick, 2017; Stutz, Schmerer, Coates,

& Bolnick, 2015). This heterogeneity could select for different MHC genotypes at

different sites. If it is true, we expect that the distributions of MHC genotypes and

parasite species would be correlated across the stickleback metapopulation. The

populations that have similar parasite communities should have similar MHC genotype

compositions, and *vice versa*. To test this hypothesis, we first constructed a Euclidian

distance matrix across all the sites for MHC alleles and for parasite species,

respectively. Then we used Mantel test to examine if the two distance matrices were

correlated. A previous analysis (Bolnick et al., 2019) found no significant effect of as-the-

crow-flies (*i.e.,* Euclidean distance) or as-the-fish-swims (*i.e.,* the shortest in-water path)

geographic distance on parasite community structure, so we omit that as a covariate in

this distance analysis.

d. Neutral evolution

We previously generated reduced representation genomic sequence data for the

populations in this study, using ddRADseq following the protocol of Peterson, Weber,

Kay, Fisher, & Hoekstra (2012) modified as described in Stuart et al. (2017). From these

data we estimated genome-wide mean heterozygosity for each individual fish, and from

this population-level mean heterozygosity. We also calculated pairwise Weir-Cockerham

adjusted F_{ST} between each pair of populations.

If MHC polymorphism is strongly affected by bottlenecks or other neutral population

genetic processes, then we may expect that MHC diversity would be positively

correlated with genomic mean heterozygosity. We tested whether the average number

of MHC alleles (per fish) in each population varied as a function of the mean

heterozygosity. We might then also expect that between-population MHC divergence is

primarily a reflection of shared ancestry due to colonization processes or ongoing gene

flow. To test this, we used a Mantel test to evaluate correlations between among-

population MHC distance matrix and the F_{ST} matrix.

Figure 1. The phylogeny and geographic distribution of sampled stickleback populations.

A. A rooted neighbor-joining tree of stickleback populations based on Fst calculated from

SNPs obtained via ddRAD sequencing (Stuart et al. 2017). Site IDs are in parentheses.

Note that Sayward Estuary (SAY), Pye Outlet (PYO), and Campbell River Marsh (CRM)

are estuary habitat, and the rest are freshwater habitat. B. The distribution of sampling

sites on Vancouver Island, Canada overlaid on Google Map. The lower panel is an

enlarged view of the rectangle box in the upper panel. Each site is represented by a

circle. The color of the circles is consistent with the tip label circles in A.

Results

Sequencing results and parasite information

We kept 1277 stickleback from 26 sites for subsequent analyses, after excluding 160

individuals whose genotype calls may have under-represented their actual diversity due

to low read depth (<450 reads each; Figure S1). These 26 sites were from 7 different

watersheds (sample site information is in Table S1). The habitat types included 21 lakes,

two rivers, and three estuaries. Each sample site had an average sample size of about

50 fish, with a range from 18 to 78 (Mean = 49.11, SD=14.7). We identified a total of

1115 unique MHC alleles, 820 (73.54%) of which were private alleles found in only one

population. On average, each fish had seven distinct MHC alleles, with a range from 3-

15 (M=6.94, SD=1.87). 4.6% variation of MHC allelic richness could be explained by

habitat type (ANOVA, habitat term, $p < 2e-16$), 1.7% variation could be explained by

watershed (ANOVA, watershed term, $p = 4.38e-05$), and 23.0% of variation was

explained by sample site (ANOVA, watershed/site term, $p < 2e-16$). Fish from river

(M=8.38, SD=1.98) and estuary (M=7.1, SD=1.81) habitat had higher MHC allelic

diversity than fish from lake habitat (M=6.77, SD=1.82).

In total, we identified 33 parasite taxa (parasite information is in Table S2). On average,

each fish 2.5 different parasite taxa, with a range from 0 to 10 (Mean = 2.53, SD=1.74).

Fish from river (M=1.94, SD=1.6) and estuary (M=1.15, SD=0.76) habitat had lower

parasite burdens than fish from lake habitat (M=2.82, SD=1.74). 11.8% variation in

parasite richness could be explained by habitat type, 4.9% variation could be explained

by watershed (ANOVA, habitat term, $p < 2e-16$), while 28.3% of variation could be

explained by sample site (ANOVA, watershed/site term, $p < 2e-16$). See Bolnick et al.

(2019 and 2020) for further analysis of ecological factors structuring the parasite

metacommunity (diversity, composition, co-occurrence) within and among lakes.

Figure 2. Testing predictions of heterozygote advantage. A. The relationship between

the number of MHC alleles per individual and its residual parasite richness. For

purposes of plotting, the residual parasite richness for each fish was calculated by using

a mixed-effect generalized linear model to control for the random effect of sampling site

and the fixed effect of the log body length (covariates in the statistical analysis we report

in the text). Each dot represents a fish. The quadratic fit using either the full dataset or

each population was represented by thick black line or thin grey lines, respectively. None

of the quadratic fit using data from individual site was significant after correcting for

multiple comparisons. B. The relationship between the average number of MHC alleles

per-fish in a given population and the average per-fish parasite richness in that

population. Each dot represents a population. The grey area represents 95% confidence

interval. Note that x and y axes are reversed from panel A to B. In A we are specifically

testing whether individual genotype affects their parasite burden, as the reverse direction

of causation is not plausible. In B we are considering the hypothesis that existing

parasite diversity is the driver of evolution of MHC diversity.

a. Heterozygote advantage

We found no evidence for an association between stickleback allelic diversity and

parasite diversity, at either the scale of individual fish or among populations. At the

individual level (Fig. 2A), the best mixed effect generalized linear regression model was

the base model (AIC: 4226.02) that only included the fixed effect of log fish body length

(coefficient = 1.24, z = 8.78, p < 2e-16) and the random intercept of sampling site

(standard deviation: 0.47) conveying that populations differ in parasite diversity (as

previously shown, (Bolnick et al., 2020)). Adding the fixed effect of MHC alleles

(coefficient = 0.01, z = 1.22, p = 0.22) into the base model slightly decreased model fit

(AIC: 4226.53). Adding the quadratic MHC fixed effect further decreased model fit (AIC:

4239.6). No additional explanatory power was given by site by MHC diversity

interactions (linear or quadratic), so we have no support for the possibility that MHC

diversity is under selection in some populations but not others (a summary of all the

models described in this section is available in Table S3). This result suggested that

MHC diversity did not have a significant impact on parasite richness.

In the model just described, the random effect of sampling site explained a large portion

of the total variance of parasite richness, as noted previously (Bolnick et al., 2020).

Populations ranged from as few as an average of 0.5 macroparasite taxa per individual,

to nearly 5 taxa per individual (Fig. 2B). We also observed significant variation in MHC

allelic diversity between populations, ranging from as low as an average of 5.31 alleles

per fish, to as high as 9.71 alleles per fish (Fig. 2B). Among populations, average MHC

diversity was unrelated to average parasite richness (coefficient = 0.03, F = 0.02, df =

24, p = 0.89, Fig. 2B).

Figure 3. Testing predictions of frequency-dependent selection: rare allele advantage. A.

The Z values of all MHC-parasite regressions in one sample site (Lawson Lake), as an

example. Each model used data from one population and regressed the infection

intensity of a focal parasite against the presence/absence of a focal MHC allele. Each

row in the grid represents a MHC allele, and each column represents a parasite taxa.

The color filling represents the effect size. The darker the blue color indicates a stronger

positive association (having the allele confers higher infection load). The darker the

orange color indicates a stronger negative association (e.g., resistance). The MHC-

parasite regressions heavily biased by extreme values are filled as grey and excluded

from further analysis. Two examples (highlighted in black boxes in A) are plotted in B. B.

The infection intensity of parasite Unionidae is negatively and positively influenced by

the presence/absence of MHC allele prot_577 and prot_673, respectively. Note the y

axis of B is on log scale. C. The relationship between the allele prevalence of an MHC

allele and its effect size (Z value) on the infection intensity of certain parasite. Each dot

represents a MHC-parasite combination in a particular population. The solid line depicts

the result of linear regression using all the data. The grey area surrounding the

regression line indicates 95% confidence interval.

b1. Frequency-dependent selection: is there a rare allele advantage

Although we found many MHC-parasite associations, we found no tendency for alleles to

be protective when rare, or vulnerable to infection when common. That is, the Z value of

a given allele's effect on a given parasite in a given site was independent of that alleles'

prevalence in that focal site. Across all sites, there were a total of 5623 MHC-parasite

combinations that were moderately frequent and so included in the analysis. After

excluding the models heavily influenced by outlier values with high leverage, we

obtained 4130 linear regression models (see Fig. 3A for an example of this result from

one population). 45 models had significant negative MHC-parasite associations and 45

models had significant positive MHC-parasite associations (see Fig. 3B for an example).

Some MHC alleles, and parasites, are repeated across multiple regression models from

different populations. None of the models were significant after we corrected for multiple-

comparison with BH method. A mixed effect linear model showed that the impact of

allele prevalence on Z values was not significant (coefficient = 0.09, t = 1.14, p = 0.27)

(Fig. 3C). Therefore, this result did not provide evidence for rare allele advantage of

MHC alleles on parasite infection intensities.

Figure 4. Testing predictions of frequency-dependent selection: inconsistent allele

effects across sites. A. The percentage of deviation residuals explained by the MHC

term and the MHC × sample site interaction term in the regression models, which

regressed the infection load of certain parasite against a particular MHC allele, sample

site, and the MHC × sample site interaction. Each dot represents a MHC-parasite

combination. The dotted line is a line of equal effect size (slope of 1, intercept of 0). The

models with either term significant are plotted with black dots, and all the non-significant

models are in grey. Two significant examples were highlighted in A, and plotted in B and

C. B. The effect of MHC allele prot_1215 on the infection load of parasite Bunoderina in

Mud lake and Little Mud lake. C. The effect of MHC allele prot_110 on the infection load

of parasite Nematode spp4 in McCreight lake and McCreight stream. Note that the y

axes of B and C are on log scale.

b2. Frequency-dependent selection: are allele effects inconsistent across lakes?

We found some cases where the allele effects are inconsistent across lakes (significant

MHC allele × site interaction effects), suggesting there were potentially Red Queen style

co-evolutionary arms race between stickleback population and parasites. We obtained

788 generalized linear models for each moderately prevalent MHC-parasite

combinations which were present in more than one site. 53 models had a significant

interaction term but an insignificant MHC term (11 of them involved parasites with birds

as final hosts), suggesting inconsistent allelic effects across sites. For example, the

allele prot_110 confers susceptibility (higher infection) in McCreight Stream, but has no

effect in McCreight Lake (Fig. 4C). Although none of the models were significant after

correction for multiple-comparison with BH method, the number of significant models

was greater than null expectation with a significance level of 0.05 (792 * 0.05 = 39.6), so

some of the model results were truly significant despite the issue of multiple comparison.

66 models had a significant MHC term but an insignificant interaction term (13 of them

involved parasites with birds as final hosts) (Fig. 4B), suggesting that those alleles had

consistent effects on the focal parasite across sites, and thus did not fit localized Red

Queen race dynamics. Parasites with birds as final hosts were not more likely to be

found in the models with significant MHC term than in the models with significant

interaction term (Chi-squared test, X^2=0.004, p=0.95).

Figure 5. Testing predictions of spatially fluctuating selection. The parasite distance

between populations is not detectably correlated with the MHC distance between

populations. Each dot represents a pair of populations. The ellipse represents 95%

confidence interval prediction.

c. Fluctuating selection

Despite spatial heterogeneity in the distribution of parasite species (Bolnick et al., 2019),

we did not detect parasite-mediated spatially fluctuating selection. The Mantel test

between the distance matrix of parasite community and the distance matrix of MHC

alleles across all populations suggested that the two matrices were not correlated (r =

0.089, p=0.22, Fig. 5). Thus, although MHC alleles and parasite communities both differ

strongly between lakes, there is no detectable relationship between them.

Figure 6. Testing predictions of neutral hypotheses. A. The average number of MHC

alleles per population is positively associated with population mean heterozygosity. Each

dot represents a population. The solid line represents linear regression prediction. Grey

area indicates 95% confidence interval. B. The relationship between MHC distance

between populations and Fst. Each dot represents a pair of populations. The ellipse

indicates 95% confidence interval prediction.

d. Neutral evolution

We found that MHC diversity was significantly associated with genome-wide genetic

diversity that should mostly reflect neutral processes. First, the average MHC allelic

richness covaried positively with population mean heterozygosity (coefficient=25.4,

F=6.57, df = 24, p = 0.02, Fig. 6A). Second, there was a strong positive correlation

between MHC divergence between populations, and genome-wide divergence (Mantel

test: r = 0.51, p=0.002, Fig. 6B). That is, more closely related populations tended to be

more similar at their MHC loci as well.

Discussion

This study represents an effort to comprehensively evaluate multiple major hypotheses

concerning the maintenance of MHC diversity, using observational data from 26

stickleback populations. Despite the large scale of sampling, we did not find strong

support for any of the three popular parasite-mediated selection hypotheses, namely

heterozygote advantage, frequency-dependent selection, and fluctuating selection. In

contrast, neutral processes seem to best explain MHC diversity in this system, both in

terms of allelic richness and between-population divergence.

Consistent with previous studies in other vertebrates (Kaufman, 2018), MHC IIβ genes

are highly polymorphic in the surveyed stickleback metapopulation. We found 1115

unique MHC IIβ alleles from 1277 individuals, with a majority of alleles only present in

one population. However, although on average fish from lake habitats are infected with

more diverse parasite species than the ones from river and estuary habitats (Bolnick et

al., 2020), lake stickleback have less MHC allelic diversity than others. The extremely

high level of polymorphism in MHC genes is often attributed to parasite-mediated

selection; however, the discrepancy between parasite richness and MHC allelic diversity

among habitat types suggests that parasite-mediated selection alone is unlikely to

explain the high level of MHC polymorphism in the studied stickleback populations. This

finding contradicts previous results from studies of stickleback inhabiting different

habitats in Germany (K. M. Wegner, Reusch, & Kalbe, 2003), where lake fish had higher

parasite diversity (as in this study), and higher MHC diversity (unlike this study).

The above finding is further supported through testing specific predictions derived from

heterozygote advantage model. The heterozygote advantage hypothesis predicts that

the number of MHC alleles in the host genome would be *maximized* to recognize a wide

range of parasites. But previous studies (K. Mathias Wegner et al., 2003) in stickleback

found support for the theoretical model that the number of MHC alleles should be

optimized at an intermediate level, rather than *maximized*. The argument is that too

many MHC alleles would reduce the T cell portfolio due to negative selection of self-

recognition T cell receptors. In this study, we applied similar analytical approaches as

Wegner et al. (2003) to a larger sample size and many more populations. Contrary to

those previous results, we did not find support for either the *maximizing* or *optimizing*

hypothesis.

Negative frequency-dependent selection suggests that the frequency of MHC alleles

might experience a cyclical pattern through time, with different low frequency alleles

playing the role of conferring resistance at different time points. For negative frequency

dependence to be effective, on average, it must be true that currently rare alleles exhibit

an advantage over common ones. In this study, we found that allele prevalence in

general is not associated with whether an allele is effective in parasite resistance. We

further examined whether alleles have consistent effects across sites. We found

inconsistent allele effects in some cases, but we also found consistent effects in a larger

fraction of cases. So negative frequency-dependent selection might be able to explain a

limited number of allele-parasite dynamics in our study. But, in most cases MHC alleles

confer equivalent protection in multiple isolated populations, inconsistent with cyclic

changes due to coevolution. Ideally, to demonstrate negative frequency-dependent

selection, one would need to carry out a longitudinal study that samples parasite

genotypes and host genotypes through time. However, this type of study is very

challenging to implement in a field setting, especially for annually reproducing animals.

Finally, fluctuating selection, either temporally or spatially, is frequently invoked to

explain the maintenance of genetic diversity across a landscape (Charbonnel &

Pemberton, 2005; Schemske & Bierzychudek, 2007). Our sampling sites on Vancouver

Island differ in many physical parameters, such as water area, depth, temperature, flow

speed, *etc*. It is also documented that parasite communities differ significantly among

different sites (Bolnick et al., 2020). This spatial heterogeneity in parasite communities is

due to abiotic and biotic variables (e.g., lake size, fish diet; Bolnick et al. (2019)), and

could generate spatially fluctuating selection that maintains MHC diversity. However, we

did not find significant correlation between the parasite distance matrix and the MHC

distance matrix, suggesting that the spatial heterogeneity in parasite community could

not explain the observed spatial variation in MHC diversity.

Interestingly, we found that neutral processes could contribute to MHC diversity in two

different analyses: (a). populations with higher genomic heterozygosity also have higher

MHC allelic richness, and (b). less divergent populations (low genomic Fst) have similar

MHC genotypes and frequencies. Similar to analysis (a), a previous study in stickleback

(K. M. Wegner et al., 2003) also found a positive, albeit weak, association between

neutral genetic diversity and population MHC allelic diversity. The role of neutral

processes in the maintenance of MHC diversity has often been studied in the context of

conservation, which is usually concerned with small and isolated populations (Miller &

Lambert, 2004; Seddon & Ellegren, 2004). Our result suggested that at larger scales,

neutral processes are also important factors in explaining MHC diversity within- and

between-population, even when there is no specific evidence for past bottlenecks.

Parasite-mediated selection is believed to have an important role in maintaining MHC

polymorphism. Consistent with this, we found some associations between certain

parasites and individual MHC allele. However, we did not find strong support for

parasite-mediated selection hypotheses in the scale of metapopulations. There are

several possible explanations. First, in this study, we examined the three hypotheses

separately, but in reality, the mechanisms are not mutually exclusive. For example, low

frequency alleles are more likely to be in a heterozygous state in a population.

Therefore, if an allele were effective in resistance at low frequency, it would be

consistent with both negative frequency-dependent selection and heterozygote

advantage. Furthermore, if these mechanisms work in combination in different MHC-

parasite pairs or in different populations, we may not detect a clear signal of each

mechanism when combining data from multiple MHC-parasite pairs and many

populations. Second, parasites are not the only selective agents that act upon MHC loci.

MHC genes are also involved in interactions with other species, such as symbionts

comprising the gut microbiome (Bolnick et al., 2014), and within species interactions,

such as mate choice (Milinski et al., 2005). It is unknown which MHC genes are relevant

to which selective forces. If other selective forces are also at work, the influences of

parasite-mediated selection on MHC diversity could be obscured. Third, we only

measured parasite richness and MHC allelic richness in this study to represent parasite

diversity and MHC diversity, respectively, so the detailed differences among individual

parasite and individual MHC alleles were not considered. In the future, it is worthwhile to

test the parasite-mediated selection hypotheses with diversity measurements that take

into account of nuances such as the phylogenetic relationship of parasite taxa and the

sequence divergence between distinct MHC alleles. Finally, our survey does not span

multiple seasons, so we cannot evaluate how MHC variation is shaped across time by

temporally fluctuating selection.

To summarize, we used a large-scale field survey to evaluate the three popular parasite-

mediated selection hypotheses on MHC diversity, namely, heterozygote advantage,

negative frequency-dependent selection, and fluctuating selection. We found that neutral

processes best explain MHC diversity (allelic richness and population divergence),

instead of those parasite-mediated selection mechanisms. Because MHC diversity can

be influenced by many selective forces, and the outcome of this selection could be

further shaped by spatial and temporal heterogeneity, our study suggests it may not be

possible to parse out how each selective force influences MHC diversity at large scales

directly from observational data. We propose that it is worthwhile to instead investigate

how each selective force acts upon MHC diversity with experimental approaches (*e.g.*

controlling other selective forces) at smaller scales. Combining insights from these

small-scale controlled studies in a step-wise manner can be a fertile future direction to

understand the complex process of MHC evolution and diversification.

Acknowledgment

This work was supported by the David and Lucille Packard Fellowship for Science and

Engineering (DIB), the Howard Hughes Medical Institute (Early Career Scientist award to

DIB), and by NIH NIAID grant 1R01AI123659-01A1. Field sampling in 2009 was made

possible by assistance from Todasporn Rodbumrung, Chris Harrison, Will Shim, Yuexin

Jiang, Will Stutz. Will Shim provided parasite information in Table S2.

Author Contribution

DIB conceived and designed this project with Will Stutz. DIB and WS performed the field

survey. KB carried out the dissections. SHW performed the MHC sequencing, SDH

performed genotype scoring. FP performed data analysis and wrote the manuscript with

input from DIB. All authors contributed in the revision of the manuscript.

References

Bolnick, D. I., & Ballare, K. M. (2020). Resource diversity promotes among-individual diet variation, but not genomic diversity, in lake stickleback. *Ecology Letters*, *23*(3), 495–505. doi: 10.1111/ele.13448 (Original work published)

Bolnick, D. I., Resetarits, E. J., Ballare, K., Stuart, Y. E., & Stutz, W. E. (2019). Scale-dependent effects of geography, host ecology, and host genetics, on species composition and co-occurrence in a stickleback parasite metacommunity. *BioRxiv*, 672410. doi: 10.1101/672410 (Original work published)

Bolnick, D. I., Resetarits, E. J., Ballare, K., Stuart, Y. E., & Stutz, W. E. (2020). Host patch traits have scale-dependent effects on diversity in a stickleback parasite metacommunity. *Ecography*, *43*(7), 990–1002. doi: 10.1111/ecog.04994 (Original work published)

Bolnick, D. I., Snowberg, L. K., Caporaso, J. G., Lauber, C., Knight, R., & Stutz, W. E. (2014). Major Histocompatibility Complex class IIb polymorphism influences gut microbiota composition and diversity. *Molecular Ecology*, *23*(19), 4831–4845. doi: 10.1111/mec.12846 (Original work published)

Bonneaud, C., Pérez☐Tris, J., Federici, P., Chastel, O., & Sorci, G. (2006). Major Histocompatibility Alleles Associated with Local Resistance to Malaria in a Passerine. *Evolution*, *60*(2), 383–389. doi: 10.1111/j.0014-3820.2006.tb01114.x (Original work published)

Borghans, J. A. M., Beltman, J. B., & De Boer, R. J. (2004). MHC polymorphism under host-pathogen coevolution. *Immunogenetics*, *55*(11), 732–739. doi: 10.1007/s00251-003-0630-5 (Original work published)

Borghans, J. A. M., Noest, A. J., & De Boer, R. J. (2003). Thymic selection does not limit the individual MHC diversity. *European Journal of Immunology*, *33*(12), 3353–3358. doi: 10.1002/eji.200324365 (Original work published)

Calhoun, D. M., McDevitt-Galles, T., & Johnson, P. T. J. (2018). Parasites of invasive

freshwater fishes and the factors affecting their richness. *Freshwater Science*,

37(1), 134–146. doi: 10.1086/696566 (Original work published)

Charbonnel, N., & Pemberton, J. (2005). A long-term genetic survey of an ungulate

population reveals balancing selection acting on MHC through spatial and

temporal fluctuations in selection. *Heredity*, *95*(5), 377–388. doi:

10.1038/sj.hdy.6800735 (Original work published)

Dobzhansky, T. (1982). *Genetics and the Origin of Species* (Vol. 11). Columbia

university press. (Original work published)

Eizaguirre, C., Lenz, T. L., Kalbe, M., & Milinski, M. (2012). Rapid and adaptive evolution

of MHC genes under parasite selection in experimental vertebrate populations.

Nature Communications, *3*(1), 621. doi: 10.1038/ncomms1632 (Original work

published)

Hedrick, P. W., Parker, K. M., & Lee, R. N. (2001). Using microsatellite and MHC

variation to identify species, ESUs, and MUs in the endangered Sonoran

topminnow. *Molecular Ecology*, *10*(6), 1399–1412. doi: 10.1046/j.1365-

294X.2001.01289.x (Original work published)

Ilmonen, P., Penn, D. J., Damjanovich, K., Morrison, L., Ghotbi, L., & Potts, W. K.

(2007). Major histocompatibility complex heterozygosity reduces fitness in

experimentally infected mice. *Genetics*, *176*(4), 2501–2508. doi:

10.1534/genetics.107.074815 (Original work published)

Jensen, P. E. (2007). Recent advances in antigen processing and presentation. *Nature*

Immunology, *8*(10), 1041–1048. doi: 10.1038/ni1516 (Original work published)

Kaufman, J. (2018). Unfinished Business: Evolution of the MHC and the Adaptive

Immune System of Jawed Vertebrates. *Annual Review of Immunology*, *36*, 383–

409. doi: 10.1146/annurev-immunol-051116-052450 (Original work published)

Matthews, B., Harmon, L. J., M'Gonigle, L., Marchinko, K. B., & Schaschl, H. (2010).

Sympatric and Allopatric Divergence of MHC Genes in Threespine Stickleback.

PLOS ONE, *5*(6), e10948. doi: 10.1371/journal.pone.0010948 (Original work

published)

Mccairns, R. J. S., Bourget, S., & Bernatchez, L. (2011). Putative causes and

consequences of MHC variation within and between locally adapted stickleback

demes. *Molecular Ecology*, *20*(3), 486–502. doi: 10.1111/j.1365-

294X.2010.04950.x (Original work published)

Milinski, M., Griffiths, S., Wegner, K. M., Reusch, T. B. H., Haas-Assenbaum, A., &

Boehm, T. (2005). Mate choice decisions of stickleback females predictably

modified by MHC peptide ligands. *Proceedings of the National Academy of*

Sciences, *102*(12), 4414–4418. doi: 10.1073/pnas.0408264102 (Original work

published)

Miller, H. C., & Lambert, D. M. (2004). Genetic drift outweighs balancing selection in

shaping post-bottleneck major histocompatibility complex variation in New

Zealand robins (Petroicidae). *Molecular Ecology*, *13*(12), 3709–3721. doi:

10.1111/j.1365-294X.2004.02368.x (Original work published)

Nowak, M. A., Tarczy-Hornoch, K., & Austyn, J. M. (1992). The optimal number of major

histocompatibility complex molecules in an individual. *Proceedings of the*

National Academy of Sciences of the United States of America, *89*(22), 10896–

10899. doi: 10.1073/pnas.89.22.10896 (Original work published)

Penn, D. J., Damjanovich, K., & Potts, W. K. (2002). MHC heterozygosity confers a

selective advantage against multiple-strain infections. *Proceedings of the*

National Academy of Sciences, *99*(17), 11260–11264. doi:

10.1073/pnas.162006499 (Original work published)

Penn, D. J., & Potts, W. K. (1999). The Evolution of Mating Preferences and Major

Histocompatibility Complex Genes. *The American Naturalist*, *153*(2), 145–164.

doi: 10.1086/303166 (Original work published)

Peterson, B. K., Weber, J. N., Kay, E. H., Fisher, H. S., & Hoekstra, H. E. (2012). Double

Digest RADseq: An Inexpensive Method for De Novo SNP Discovery and

Genotyping in Model and Non-Model Species. *PLOS ONE*, *7*(5), e37135. doi:

10.1371/journal.pone.0037135 (Original work published)

Radwan, J., Babik, W., Kaufman, J., Lenz, T. L., & Winternitz, J. (2020). Advances in the

Evolutionary Understanding of MHC Polymorphism. *Trends in Genetics*, *0*(0).

doi: 10.1016/j.tig.2020.01.008 (Original work published)

Schemske, D. W., & Bierzychudek, P. (2007). Spatial Differentiation for Flower Color in

the Desert Annual Linanthus Parryae: Was Wright Right? *Evolution*, *61*(11),

2528–2543. doi: 10.1111/j.1558-5646.2007.00219.x (Original work published)

Sebastian, A., Herdegen, M., Migalska, M., & Radwan, J. (2016). amplisas: a web server

for multilocus genotyping using next-generation amplicon sequencing data.

Molecular Ecology Resources, *16*(2), 498–510. doi: 10.1111/1755-0998.12453

(Original work published)

Seddon, J. M., & Ellegren, H. (2004). A temporal analysis shows major histocompatibility

complex loci in the Scandinavian wolf population are consistent with neutral

evolution. *Proceedings of the Royal Society of London. Series B: Biological*

Sciences, *271*(1554), 2283–2291. doi: 10.1098/rspb.2004.2869 (Original work

published)

Sommer, S. (2005). The importance of immune gene variability (MHC) in evolutionary

ecology and conservation. *Frontiers in Zoology*, *2*, 16. doi: 10.1186/1742-9994-2-

16 (Original work published)

Stuart, Y. E., Veen, T., Weber, J. N., Hanson, D., Ravinet, M., Lohman, B. K., … Bolnick,

D. I. (2017). Contrasting effects of environment and genetics generate a

continuum of parallel evolution. *Nature Ecology & Evolution*, *1*(6), 1–7. doi:

10.1038/s41559-017-0158 (Original work published)

Stutz, W. E., & Bolnick, D. I. (2014). Stepwise Threshold Clustering: A New Method for

Genotyping MHC Loci Using Next-Generation Sequencing Technology. *PLOS*

ONE, *9*(7), e100587. doi: 10.1371/journal.pone.0100587 (Original work

published)

Stutz, W. E., & Bolnick, D. I. (2017). Natural selection on MHC IIβ in parapatric lake and

stream stickleback: Balancing, divergent, both or neither? *Molecular Ecology*,

26(18), 4772–4786. doi: 10.1111/mec.14158 (Original work published)

Stutz, W. E., Schmerer, M., Coates, J. L., & Bolnick, D. I. (2015). Among-lake reciprocal

transplants induce convergent expression of immune genes in threespine

stickleback. *Molecular Ecology*, *24*(18), 4629–4646. doi: 10.1111/mec.13295

(Original work published)

Wegner, K. M., Reusch, T. B. H., & Kalbe, M. (2003). Multiple parasites are driving major

histocompatibility complex polymorphism in the wild. *Journal of Evolutionary*

Biology, *16*(2), 224–232. doi: 10.1046/j.1420-9101.2003.00519.x (Original work

published)

Wegner, K. Mathias, Kalbe, M., Kurtz, J., Reusch, T. B. H., & Milinski, M. (2003).

Parasite Selection for Immunogenetic Optimality. *Science*, *301*(5638), 1343–

1343. doi: 10.1126/science.1088293 (Original work published)

747