Tau, XMAP215/Msps and Eb1 co-operate interdependently to regulate microtubule polymerisation and bundle formation in axons

Ines Hahn[#, 1], Andre Voelzmann[1], Jill Parkin[1], Judith Fuelle[1], Paula G Slater[2], Laura A Lowery[3], Natalia Sanchez-Soriano[#‡4] & Andreas Prokop[#‡1]

1) The University of Manchester, Manchester Academic Health Science Centre, Faculty of Biology, Medicine and Health, School of Biology, Manchester, UK

2) Department of Biology, Boston College, Chestnut Hill, MA, USA

3) Department of Medicine, Boston University Medical Center, Boston, MA, USA

4) Department of Molecular Physiology & Cell Signalling, Institute of Systems, Molecular & Integrative Biology, University of Liverpool, Liverpool, United Kingdom

Running title**:** Tau, XMAP215 and Eb1, a functional trio for microtubule polymerisation and organisation

Key words: *Drosophila*, neurodegeneration, axons, cytoskeleton, microtubules

[#]   authors for correspondence:

The University of Manchester

Faculty of Life Sciences

Oxford Road

Manchester M13 9PT

Tel: +44-(0)161-27-51556

Fax: +44-(0)161-27-51505

Ines.Hahn@manchester.ac.uk

N.Sanchez-Soriano@liverpool.ac.uk

Andreas.Prokop@manchester.ac.uk

[‡]   joined senior authors

Summary statement:

Eb1, XMAP215 and tau co-operate interdependently in axons to promote the polymerisation of microtubules and their organisation into the parallel bundles required for axonal transport.

Abstract

The formation and maintenance of microtubules requires their polymerisation, but little is known about how this polymerisation is regulated in cells. Focussing on the essential microtubule bundles in axons of *Drosophila* and *Xenopus* neurons, we show that the plus-end scaffold Eb1, the polymerase XMAP215/Msps and the lattice-binder Tau co-operate interdependently to promote microtubule polymerisation and bundle organisation during axon development and

maintenance. Eb1 and XMAP215/Msps promote each other's localisation at polymerising microtubule plus-ends. Tau outcompetes Eb1-binding along microtubule lattices, thus preventing depletion of Eb1 tip pools. The three factors genetically interact and show shared mutant phenotypes: reductions in axon growth, comet size, comet number and comet velocity, as well as prominent deterioration of parallel microtubule bundles into disorganised curled conformations. This microtubule curling is caused by Eb1 plus-end depletion which impairs spectraplakin-mediated guidance of extending microtubules into parallel bundles. Our demonstration that Eb1, XMAP215/Msps and Tau co-operate during the regulation of microtubule polymerisation and bundle organisation, offers new conceptual explanations for developmental and degenerative axon pathologies and how to treat them.

Introduction

Axons are the enormously long cable-like cellular processes of neurons that wire nervous systems. In humans, axons of ≤15μm diameter can be up to two meters long (Prokop, 2020; Prokop, 2021). They are constantly exposed to mechanical challenges, yet have to survive for up to a century; we lose ~40% of axons towards high age and far more in neurodegenerative diseases (Adalbert & Coleman, 2012; Calkins, 2013; Marner *et al*, 2003).

Their growth and maintenance absolutely require parallel bundles of microtubules (MTs) that run all along axons, providing the highways for life-sustaining transport and driving morphogenetic events. Consequently, bundle decay through MT loss or curling is a common feature in axon pathologies (summarised in (Hahn *et al*, 2019; Prokop, 2021). Key roles must be played by MT polymerisation, which is not only essential for the *de novo* formation of MT bundles occurring during axon growth in development, plasticity or regeneration, but also to repair damaged and replace senescent MTs during long-term maintenance (Gasic & Mitchison, 2019; Schaedel *et al*, 2015; Voelzmann *et al*, 2016). However, the molecular mechanisms regulating MT polymerisation in axons are surprisingly little understood.

MT polymerisation is primarily understood *in vitro*, where MTs can undergo polymerisation in the presence of nucleation seeds and tubulin heterodimers; the addition of catalytic factors such as CLASPs, stathmins, tau, Eb proteins or XMAP215 can enhance and refine the process (Aher *et al*, 2020; Al-Bassam *et al*, 2010; Brouhard *et al*, 2008; Drechsel *et al*, 1992; Li *et al*, 2012; Manna *et al*, 2009; Zanic *et al*, 2013). However, we do not know whether mechanisms observed in reconstitution assays are biologically relevant in the context of axons (Voelzmann *et al.*, 2016), especially when considering that none of the above-mentioned factors has genetic links to human neurological disorders on OMIM® (Online Mendelian Inheritance in Man), except Tau/MAPT which executes numerous disease-relevant functions also when detached from MTs (Morris *et al*, 2011).

To identify relevant factors regulating axonal MT polymerisation, we use *Drosophila* primary neurons as one consistent model, which is amenable to combinatorial genetics as a powerful strategy to decipher complex regulatory networks (Prokop *et al*, 2013). Our previous loss-of-function studies of 9 MT plus-end-associating factors in these *Drosophila* neurons (CLASP, CLIP190, dynein heavy chain, APC, p150[Glued], Eb1, Short stop/Shot, doublecortin, Lis1) have taken axon length as a crude proxy readout for net polymerisation, mostly revealing relatively mild axon length phenotypes, with the exception of Eb1 and Shot which cause severe axon shortening {Beaven, 2015 #6487; Sánchez-Soriano, 2009 #7139; Alves-Silva, 2012 #4930; A.V., unpublished data}.

Here we have incorporated more candidate factors and readouts to take these analyses to the

next level. We show that three factors, Eb1, XMAP215/Msps and Tau, share a unique
combination of mutant phenotypes in culture and *in vivo*, including reduced axonal MT
polymerisation in frog and fly neurons. Our data reveal that the three factors co-operate. Eb1
and XMAP215/Msps act interdependently at MT plus-ends, whereas Tau acts through a novel
mechanism: it outcompetes Eb1-binding along MT lattices, thus preventing the depletion of Eb1
pools at polymerising MT plus-ends. By upholding these Eb1 pools, the functional trio also
promotes the bundle conformation of axonal MTs through a guidance mechanism mediated by
the spectraplakin Shot. Our work uniquely integrates molecular mechanisms into understanding
of MT regulation that is biologically relevant for axon growth, maintenance and disease.

Methods

<u>Fly stocks</u>
Loss-of-function mutant stocks used in this study were the deficiencies *Df(3R)Antp17* (*tub^def^*;
removing both *αtub84B* and *αtub84D*; (Duncan & Kaufman, 1975; Jenkins *et al*, 2017),
Df(2L)Exel6015 (*stai^Df^*; (Duncan *et al*, 2013), *Df(3L)BSC553* (*clasp^Df^*; Bloomington stock
#25116; Beaven *et al*, 2015, *Df(3R)tauMR22* (*tau^Df^*; Doerflinger *et al*, 2003) and the loss-of-
function mutant alleles *α-tub84B^KO^* (an engineered null-allele; Jenkins *et al.*, 2017),
*chromosome bows^2^* (*clasp^2^*, an amorph allele; Inoue *et al*, 2000), *Eb1^04524^*, *Eb1^5^* (strong loss-of-
function mutant alleles; Elliott *et al*, 2005), *futsch^P158^* (*MAP1B^-^*; a deficiency uncovering the
futsch locus; Hummel *et al*, 2000), *msps^A^* (a small deletion causing a premature stop after 370
amino acids; gift from H. Ohkura, unpublished), *msps^146^* (Brittle & Ohkura, 2005), *sentin^ΔB^* (*short
spindle2^ΔB^*, *ssp2^ΔB^*; Gluszek *et al*, 2015), *tacc^1^* (*dTACC^1^*; Gergely *et al*, 2000), *shot^3^* (the
strongest available allele of *short stop*; Kolodziej *et al*, 1995; Sánchez-Soriano *et al*, 2009),
*stai^KO^* (Yang *et al*, 2012), *tau^KO^* (a null allele; Burnouf *et al*, 2016). Gal4 driver lines used were
elav-Gal4 (1^st^ and 3^rd^ chromosomal, both expressing pan-neuronally at all stages; Luo *et al*,
1994), *GMR31F10-Gal4* (Bloomington #49685; expressing in T1 medulla neurons; Qu *et al*,
2019). Lines for targeted gene expression were *UAS-Eb1-GFP* and *UAS-shot-Ctail-GFP* (Alves-
Silva *et al*, 2012), *UAS-shot^ΔABD^-GFP* (Qu, 2015), *UAS-shot^3MTLS^-GFP* (Alves-Silva *et al.*, 2012),
UAS-dtau-GFP (Doerflinger *et al.*, 2003), *UAS-GFP-α-tubulin84B* (Grieder *et al*, 2000) and
further lines generated here (see below).

<u>*Drosophila* primary cell culture</u>
Drosophila primary neuron cultures were done as described previously (Prokop *et al*, 2012; Qu
et al, 2017). Stage 11 embryos were treated for 90 s with bleach to remove the chorion, sterilized
for ~30 s in 70% ethanol, washed in sterile Schneider's medium containing 20% fetal calf serum
(Schneider's/FCS; Gibco), and eventually homogenized with micro-pestles in 1.5 centrifuge
tubes containing 21 embryos per 100 μl dispersion medium (Prokop *et al.*, 2012) and left to
incubated for 4 min at 37°C. Dispersion was stopped with 200 μl Schneider's/FCS, cells were
spun down for 4 mins at 650 g, supernatant was removed and cells re-suspended in 90 μl of
Schneider's/FCS, and 30 μl drops were placed in culture chambers and covered with cover
slips. Cells were allowed to adhere to cover slips for 90-120 min either directly on glass or on
cover slips coated with a 5 μg/ml solution of concanavalin A, and then grown as a hanging drop
culture at 26°C as indicated.

To eliminate a potential maternal rescue of mutants (i.e. reduction of the mutant phenotype due
to normal gene product deposition from the wild-type gene copy of the heterozygous mothers

in oocytes (Prokop, 2013), we used a pre-culture strategy (Prokop *et al.*, 2012; Sánchez-
Soriano *et al*, 2010) where cells were incubated in a tube for 5-7 days before they were plated
on coverslips.

For cultures from larval brains, L3 brains (2-3 per cover slip) were dissected in PBS, transferred
into Schneider's/FCS medium, washed three times with medium and then proceed with
homogenisation and dispersion as explained above

Transfection of *Drosophila* primary neurons was executed as described previously (Qu *et al.*,
2019). In brief, 70-75 embryos per 100 μl dispersion medium were used. After the washing step
and centrifugation, cells were re-suspended in 100 μl transfection medium [final media
containing 0.1-0.5 μg DNA and 2 μl Lipofecatmine 2000 (L2000, Invitrogen), incubation
following manufacturer's protocols (Thermo Fisher, Invitrogen) and kept for 24 hrs at 26°C. Cells
were then treated again with dispersion medium, re-suspended in culture medium and plated
out as described above.

<u>*Xenopus* primary neuron experiments</u>

Xenopus primary neuron cultures were obtained from embryonic neural tube. Eggs collected
from female X. laevis frogs were fertilised in vitro, dejellied and cultured following standard
methods (Sive *et al*, 2010). Embryos were grown to stage 22–24 (Nieuwkoop & J., 1994), and
neural tubes were dissected as described (Lowery *et al*, 2012). Three neural tubes were
transferred to an Eppendorf tube containing 150 uL CMF-MMR (0.1 M NaCl, 2.0 mM KCl, 1.0
mM EDTA, 5.0 mM HEPES, pH 7.4), 10 min later centrifuged at 1000 g for 5 min, and 150 uL
of Steinberg's solution (58 mM NaCl, 0.67 mM KCl, 0.44 mM $Ca(NO_3)_2$, 1.3 mM $MgSO_4$, 4.6
mM Tris, pH 7.8) was added to the supernatant to follow with the tissue dissociation using a fire
polished glass Pasteur pipet. Cells were seeded in 100 ug/mL Poly-L-lysine and 10 μg/ml
laminin pre-treated 60mm plates, and after 2 hr the media was replaced by plating culture media
(50% Ringer's, 49% L-15 media, 1% Fetal Bovine Serum, 25 ng/μl NT3 and BDNF, plus 50
μg/ml penicillin/streptomycin and gentamycin, pH 7.4 and filter sterilized) and kept for 24 hr
before imaging.

All experiments were approved by the Boston College Institutional Animal Care and Use
Committee and performed according to national regulatory standards.
The embryos were injected four times in dorsal blastomeres at two-to-four cell stage with 6 ng
of the validated XMAP215 morpholino (MO; Lowery *et al*, 2013), 10 ng of the validated Tau MO
(Liu *et al*, 2015), and/or 5 ng of a newly designed splice site MO for EB3
(3´CTCCCAATTGTCACCTACTTTGTCG5´; for verification see Fig3-S1), in order to obtain a
50% knockdown of each.

To assess EB1 comet dynamics and comet amounts, 300 pg of MACF43-Ctail::GFP, an Eb
protein-binding 43-residue fragment derived from the C-terminal regions of hMACF2 (human
microtubule actin crosslinking factor 2; Fig.3F-F'''; Honnappa *et al*, 2009; Slater *et al*, 2019),
was co-injected with the MO.

<u>Immunohistochemistry</u>

Primary fly neurons were fixed in 4% paraformaldehyde (PFA; in 0.05 M phosphate buffer, pH
7–7.2) for 30 min at room temperature (RT). For anti-Eb1 and anti-GTP-tubulin staining, cells
were fixed for 10 mins at -20°C in +TIP fix (90% methanol, 3% formaldehyde, 5 mM sodium
carbonate, pH 9; stored at -80°C and added to the cells) (Rogers *et al*, 2002), then washed in
PBT (PBS with 0.3% TritonX). Antibody staining and washes were performed with PBT. Staining

reagents: anti-tubulin (clone DM1A, mouse, 1:1000, Sigma; alternatively, clone YL1/2, rat,
1:500, Millipore Bioscience Research Reagents); anti-DmEb1 (gift from H. Ohkura; rabbit,
1:2000; (Elliott *et al.*, 2005); anti-GTP-tubulin (hMB11; human, 1:200; AdipoGen; Dimitrov *et al*,
2008); anti-Shot (1:200, guinea pig; Strumpf & Volk, 1998); anti-Elav (Elav-7E8A10; rat, 1:1000;
Developmental Studies Hybridoma Bank, The University of Iowa, IA, USA) (O'Neill *et al*, 1994);
anti-GFP (ab290, Abcam, 1:500); Cy3-conjugated anti-HRP (goat, 1:100, Jackson
ImmunoResearch); F-actin was stained with phalloidin conjugated with TRITC/Alexa647, FITC
or Atto647N (1:100 or 1:500; Invitrogen and Sigma). Specimens were embedded in ProLong
Gold Antifade Mountant (ThermoFisher Scientific).

Western blot analysis of *Xenopus* embryos

For protein extraction, 10 embryos were transferred to a centrifuge tube with 800 µl lysis buffer
(50 mM Tris pH 7.5, 5% glycerol, 0.2% IGEPAL/NP-40, 1 mM EDTA, 1.5 mM $MgCl_2$, 125 mM
NaCl, 25 mM NaF, 1 mM Na_3VO_4), homogenised with a sterile pestle and, after 10 min,
centrifuged at 13000 rpm for 15-20 min. The supernant was collected and the protein
concentration determined with the Micro BCA^{TM} Protein Assay Kit (Thermo Fisher Scientific).
80 µg protein were loaded into a 10 % SDS gel and stained with anti-Tau (clone Tau46, T9450,
mouse, 1:1000, Sigma-Aldrich).

Dissection of adult heads

For *in vivo* studies, brain dissections were performed in Dulbecco's PBS (Sigma, RNBF2227)
after briefly sedating them on ice. Dissected brains with their laminas and eyes attached were
placed into a drop of Dulbecco's PBS on MatTek glass bottom dishes (P35G1.5-14C),
covered by coverslips and immediately imaged with a 3i Marianas Spinning Disk Confocal
Microscope.

Microscopy and data analysis

Standard imaging was performed with AxioCam 506 monochrome (Carl Zeiss Ltd.) or
MatrixVision mvBlueFox3-M2 2124G digital cameras mounted on BX50WI or BX51 Olympus
compound fluorescent microscopes. For the analysis of *Drosophila* and *Xenopus* primary
neurons, we used the following parameters:

Axon length was measured from cell body to growth cone tip using the segmented line tool of
ImageJ (Alves-Silva *et al.*, 2012; Sánchez-Soriano *et al.*, 2010).

Degree of disorganised MT curling in axons was measured as "MT disorganisation index" (MDI)
described previously (Qu *et al.*, 2019; Qu *et al.*, 2017); in short: the area of disorganised curling
was measured with the freehand selection in ImageJ; this value was then divided by axon length
(see above) multiplied by 0.5 µm (typical axon diameter, thus approximating the expected area
of the axon if it were properly bundled).
Eb1 comet amounts were approximated by using product of comet mean intensity and length.
For this, a line was drawn through each comet (using the segmented line tool in FIJI) and length
as well as mean staining intensity (of Eb1 or GTP-tub in fixed *Drosophila* and MACF43::GFP in
a movie still in *Xenopus* neurons) was determined.

To measure MT curling in the optic lobe of adult flies, *GMR31F10-Gal4* (Bloomington #49685)
was used to express *UAS-α-tubulin84B-GFP* (Grieder *et al.*, 2000) in a subset of lamina axons

which projects within well-ordered medulla columns (Prokop & Meinertzhagen, 2006). Flies
were left to age for 26-27 days (about half their life expectancy) and then their brains were
dissected out, mounted in Mattek dishes and imaged using a 3i spinning disk confocal system
at the ITM Biomedecial imaging facility at the University of Liverpool. A section of the medulla
columns comprising the 4 most proximal axonal terminals was used to quantify the number of
swellings and regions with disorganised curled MTs.

To measure MT polymerisation dynamics, movies were collected on an Andor Dragonfly200
spinning disk upright confocal microscope (with a Leica DM6 FS microscope frame) and using
a *100x/1.40 UPlan SAPO (Oil)* objective. Samples where excited using 488nm (100%) and
561nm (100%) diode lasers via Leica GFP and RFP filters respectively. Images where collected
using a Zyla 4.2 Plus sCMOS camera with a camera gain of 1x. The incubation temperature
was set to 26°C. Time lapse movies were constructed from images taken every 1 s for 1 mins.
To measure comet velocity and lifetime, a line was drawn that followed the axon using the
segmented line tool in ImageJ. A kymograph was then constructed from average intensity in
FIJI using the KymoResliceWide macro (Cell Biology group, Utrecht University) and events
scored via the Velocity Measurement Tool Macro (Volker Baecker, INSERM, Montpellier, RIO
Imaging; J. Rietdorf, FMI Basel; A. Seitz, EMBL Heidelberg). For each condition at least 15 cells
were analysed in ≥2 independent repeats.

Time lapse imaging for *Xenopus* primary cultures was performed with a CSU-X1M 5000
spinning-disk confocal (Yokogawa, Tokyo, Japan) on a Zeiss Axio Observer inverted motorized
microscope with a Zeiss 63× Plan Apo 1.4 numerical aperture lens (Zeiss, Thornwood, NY).
Images were acquired with an ORCA R2 charge-coupled device camera (Hamamatsu,
Hamamatsu, Japan) controlled with Zen software. Time lapse movies were constructed from
images taken every 2 s for 1 min. The MACF43 comets' velocities and lifetime were analysed
with plusTipTracker software. The same parameters were used for all movies: maximum gap
length, eight frames; minimum track length, three frames; search radius range, 5–12 pixels;
maximum forward angle, 50°; maximum backward angle, 10°; maximum shrinkage factor, 0.8;
fluctuation radius, 2.5 pixels; and time interval 2 s.

Images were derived from at least 3 independent experimental repeats performed on different
days, for each of which at least 2 independent culture wells were analysed by taking a minimum
of 20 images per slide. For statistical analyses, Kruskal–Wallis one-way ANOVA with *post hoc*
Dunn's test or Mann–Whitney Rank Sum Tests (indicated as P_{MW}) were used to compare
groups, r and p-value for correlation were determined via non-parametric Spearman correlation
analysis (tests showed that data are not distributed normally). All raw data of our analyses are
provided as supplementary Excel files T1-6.

<u>Molecular biology</u>

To generate the *UAS-msps^{FL}-GFP* (aa1-2050) and *UAS-msps^{ΔCterm}* (aa1-1322) constructs,
eGFP was PCR-amplified from pcDNA3-EGFP and *msps* sequences from cDNA clone
LP04448 (DGRC; FBcl0189229) using the following primers:

msps^{FL} and *msps^{ΔCterm}* fw:
GAATAGGGAATTGGGAATTCGTTAGGCGCGCCAACATGGCCGAGGACACAGAGTAC

msps^{FL} rev:
CAAGAAAGAGAATCATGCCCAAGGGCCCGGTAGCGGCAGCGGTAGCGTGAGCAAGGGC
GAG

msps^{ΔCterm} rev:
GATGGAGGGTCTAAAATCGCATATGGGTAGCGGCAGCGGTAGCGTGAGCAAGGGCGAG
GAG

eGFP fw:
GAGAATCATGCCCAAGGGCCCGGTAGCGGCAGCGGTAGCGTGAGCAAGGGCGAGGAG
CTG

eGFP rev:
CTCTCGGCATGGACGAGCTGTACAAGTAGGCGGCCGCCTCGAGGGTACCTCTAGAG

The *msps* and *eGFP* sequences were introduced into *pUAST-attB* via Gibson assembly
(ThermoFisher) using *EcoRI* and *XhoI*. To generate transgenic fly lines, *pUAST-attB*
constructs were integrated into *PBac{yellow[+]-attP-9A}VK00024* (Bloomington line #9742) via
PhiC31-mediated recombination (outsourced to Bestgene Inc).

Results

<u>Eb1, Msps/XMAP215 and Tau share the same combination of loss-of-function phenotypes in</u>
<u>axons</u>

To identify factors relevant for axonal MT polymerisation, we performed a detailed loss-of-
function study with a selection of candidate factors suggested by the literature (Voelzmann *et*
al., 2016). We included the MT plus-end-associating factors Eb1, Shot, CLASP/Chb
(Chromosome bows) and the XMAP215 homologue Msps. As MT shaft-binding candidates, we
chose Tau and Map1b/Futsch. To explore the impact of tubulin availability on polymerisation,
we used α1-tubulin/αtub84B (the predominant α-tubulin expressed in the fly nervous system;
FlyAtlas 2, University of Glasgow, UK; A.V., unpublished) and Stathmin (a promoter of tubulin
pools; (Duncan *et al.*, 2013; Manna *et al*, 2007); J.F. and A.V., unpublished).

We analysed primary neurons derived from animals carrying loss-of-function mutations of the
respective genes. To exclude that phenotypes are masked by maternal contribution (which is
wild-type gene product deposited in the eggs by the heterozygous mothers), we used two
strategies (Prokop *et al.*, 2012; Sánchez-Soriano *et al.*, 2010): where possible, we analysed late
larval brain-derived primary neurons at 18 HIV (from now on referred to as larval neurons); if
mutants did not reach larval stages, we used embryo-derived neurons that were kept for 5-7
days in pre-culture to deplete maternal product and then plated and grown for 12 hrs (pre-
cultured neurons). In all cases, primary neurons were immuno-stained either for endogenous
tubulin to assess axon length, or for endogenous Eb1 protein to gain a first insight into the
polymerisation state of axonal MTs.

Except for Shot and Futsch, loss of all other factors displayed a significant reduction in the
number of Eb1-positive plus-end comets (Figs. 1A-D, I and 1-S1A). In addition, we measured
the mean intensities and mean lengths of Eb1 comets and used multiplication of these two
parameters to approximate Eb1 amounts at MT plus-ends. The strong hypomorphic *Eb1^{04524}*
mutant allele is known to display severe, but not complete reduction of protein levels (Elliott *et*
al., 2005); accordingly, Eb1 amounts at MT plus-ends were severely, but not completely
diminished in neurons mutant for this allele (Figs. 1D,J and 1-S1B). Out of the remaining seven
candidate factors, only two further genes showed the same qualitative mutant phenotype:
msps^{A/A} showed a reduction almost as strong as *Eb1^{04524/04524}*, and *tau^{KO/KO}* mutant neurons
showed a milder but reliable Eb1 depletion (Fig. 1B,C, J). In all cases, the drop in Eb1 amounts
was to almost equal parts due to reductions in comet length and intensity (Fig. 1-S2A,B); across

all genotypes tested. They correlated well with reduced comet velocities and lifetimes when
assessed in live imaging experiments (Fig. 1K,L and 1-S2C-E; using Shot-Ctail::GFP as readout
for plus-end dynamics; Alves-Silva *et al.*, 2012). Our data demonstrate therefore that
comparable comet length/velocity correlations made *in vitro* (Roostalu *et al*, 2020) are relevant
in the cellular context of axons.

To assess whether reduced comet velocities or numbers correlate with impaired axon growth,
we performed tubulin staining and measured axon length. Out of the eight factors, all but
Futsch/MAP1B showed a decrease in axon length ranging between 10 and 43% when
compared to parallel control cultures with wild-type neurons (Figs. 1E-H, M and 1-S1C). In these
experiments, Tau-, Msps-, Shot- or Eb1-deficient axons showed areas of prominent
disintegration of axonal MT bundles where MTs displayed intertwined, criss-crossing curls
which can be quantified via the MT disorganisation index (MDI; see methods; white arrowheads
in 1F-H,N and 1-S1D). This MT curling phenotype was known for Shot and Eb1 (Alves-Silva *et*
al., 2012), but unexpected for Tau and Msps.

Taken together, out of eight candidate factors, loss of Futsch/MAP1B was the only condition
showing no obvious defects. In contrast, loss of Eb1, Msps and Tau stood out by showing the
same phenotypic pattern: reductions in comet numbers, comet velocities, Eb1 amounts and
axon lengths, as well as a MT curling phenotype shared with loss of Shot. These five
phenotypes appeared consistently milder in *tau* than *Eb1* or *msps* mutant neurons. Our
observations raised the question as to why these three factors display such a striking pattern of
collective phenotypes.

<u>Eb1, Msps and Tau genetically interact during axonal MT regulation both in culture and *in vivo*</u>

To investigate potential functional relationship between Eb1, Msps and Tau, we performed
genetic interaction studies. We used heterozygous conditions (i.e. one mutant and one normal
copy) of genes to reduce their respective protein levels. If reduced protein levels of different
genes combined in the same neurons cause a phenotype, this suggests that they are likely to
function in a common pathway. For our studies, we used larval neurons and five different
readouts: axon length (Fig. 2A), MT curling (Fig. 2A'), Eb1 amounts at MT plus-ends (Fig. 2A''),
comet numbers (Fig. 2-S1A) and comet dynamics (Fig. 2-S1B).

To assess the baseline, we analysed single-heterozygous mutant neurons ($Eb1^{04524/+}$, $msps^{A/+}$
or $tau^{KO/+}$), none of which displayed any phenotypes. Certain double-combinations of these
heterozygous conditions in the same neurons ($Eb1^{04524/+}$ $msps^{A/+}$ and $tau^{KO/+}$ $msps^{A/+}$) displayed
a mild but significant reduction in Eb1 amounts at MT plus-ends and a trend towards stronger
MT curling. Strong enhancement of the phenotypes for all five readouts was observed in
$Eb1^{04524/+}$ $msps^{A/+}$ $tau^{KO/+}$ triple-heterozygous neurons, suggesting that the three factors are
functionally linked when regulating MT plus-ends, axon growth and MT organisation (Fig. 2A'-
A'' and 2-S1A,B).

The triple-heterozygous condition had a similar effect in fly brains *in vivo* when using axons of
T1 medulla neurons in the adult optic lobe as readouts. In control flies analysed at 26-27 days
after eclosure, axons of T1 neurons contained prominent MT bundles labelled by α-tubulin::GFP
(Fig.3A; Qu *et al.*, 2019). In contrast, T1 axons of triple-heterozygous mutant flies aged in
parallel, showed a strong increase in areas where MTs become unbundled and twisted and
axons displayed prominent swellings often containing curled MTs (Fig.3A-D). These data
strongly suggest that the co-operating network of Eb1, Msps and Tau is relevant *in vivo*.

<u>Eb1, Msps and Tau interaction is evolutionarily conserved in *Xenopus* neurons</u>

To assess potential evolutionary conservation, we evaluated whether Eb1, Msps and Tau
interact also in frog primary neurons. In the frog *Xenopus laevis*, *mapt* (*microtubule associated*
protein tau) is the only *tau* homologue, *XMAP215* (*ckap5/cytoskeleton associated protein 5*) the
only *msps* homologue, and *EB3* (*mapre3/microtubule associated protein RP/EB family member*
3) is the only one of three *Eb1* homologues that is prominently expressed in the nervous system
(Bowes *et al*, 2009; Karimi *et al*, 2017). We used morpholinos against these three genes. Similar
to our strategy in *Drosophila*, we analysed MTs by staining for endogenous tubulin (Fig.3E-E'''),
and measured Eb3 comet amounts in live movies using the Eb protein-binding peptide
MACF43-Ctail::GFP as readout (comparable to Shot-Ctail::GFP used in Fig.1K,L; see Methods;
Honnappa *et al.*, 2009; Slater *et al.*, 2019).

To approximate heterozygous mutant conditions used in our *Drosophila* experiments, we
adjusted morpholino concentrations to levels that achieved knock-down of each of the three
genes to ~50% (Fig. 3-S1A-B and Lowery *et al.*, 2013; Slater *et al.*, 2019). Individual knock-
downs to approximately 50% did not cause prominent decreases in MACF43::GFP comet
amounts or increases in MT curling; but when knock-down of all three factors was combined in
the same neurons, we found a reduction in MACF43::GFP comet amounts to 60% and a 4.8
fold increase in MT bundle disintegration and curling (Figs. 3E''',F''',G,H and 3-S1C,D).

Together, the three genes appear to functionally interact also in *Xenopus*, suggesting
evolutionary conservation of the underpinning mechanisms.

<u>Eb1, Msps and Tau form an interdependent functional network with key roles played by Eb1</u>

To investigate the underlying mechanisms, we next asked whether the functions of the three
factors are hierarchically and/or interdependently organised, or whether they regulate the
assessed MT properties through independent parallel mechanisms. To distinguish between
these possibilities, we combined homozygous mutant conditions of the three mutant alleles in
the same neurons (*Eb1$^{04524/04524}$ msps$^{A/A}$ tau$^{KO/KO}$*) and asked whether this condition enhances
phenotypes over single-homozygous conditions (suggesting parallel mechanisms), or whether
they show no further increases (suggesting a hierarchical organisation) (Avery & Wasserman,
1992).

Since the homozygous mutant conditions do not all survive into the late larval stage, we used
embryonic pre-cultured neurons for these experiments. As reference we used *Eb1$^{04524/04524}$*,
msps$^{A/A}$ and *tau$^{KO/KO}$* single mutant neurons; since *Eb1^{04524}* is a strong but not a total loss-of-
function allele (Elliott *et al.*, 2005), we added neurons homozygous for *Df(2R)Exel6050* (from
now referred to as *Eb1^{Df}*), a deficiency uncovering the entire *Eb1* locus (see Methods). The
order of phenotypic strengths in these pre-cultured homozygous mutant neurons consistently
was *Eb1$^{Df/Df}$* > *Eb1$^{04524/04524}$* ≥ *msps$^{1/1}$* > *tau$^{KO/KO}$* for all parameters assessed (Figs.2, 2-S1; see
Discussion). The strong phenotypes of the *Eb1$^{Df/Df}$* allele provided therefore a new reference
bar, displaying values that were not excelled or even reached by any combination of
homozygous mutants, including the triple-homozygous mutant neurons (Fig. 2B-B'').

These results are consistent with a model where the three proteins co-operate interdependently,
through a modality in which Tau has the weakest contribution and Eb1 is the most essential
factor. This is also suggested when extracting data from a range of different genetic conditions
analysed throughout this study and plotting their Eb1 amounts at MT plus-ends against the

degree of MT curling in each condition (Fig.2-S2A). This plot revealed a highly significant
inverse correlation between Eb1 comet amounts and MT disorganisation (Figs. 2C and 2-S2B),
suggesting that Eb1 is the key factor within the functional trio linking out to MT bundle promotion.

<u>Eb1 and Msps depend on each other for MT plus-end localisation</u>

We next investigated the mechanistic links between the three proteins, starting with Msps and
Eb1. When co-expressing Msps::GFP together with Eb1::RFP in *Drosophila* embryonic primary
neurons, both proteins prominently localised at the same MT plus-ends, with Msps::GFP
localising slightly distal to Eb1 (Fig. 4A-A'', Movie M1). Their localisation at the same MT plus-
ends was even clearer in kymographs of live movies where both proteins remained closely
associated during comet dynamics (Fig.4B,B'). These findings agree with localisation data for
Eb proteins and XMAP215 *in vitro* (Maurer *et al*, 2014), suggesting that the same mechanisms
apply in the cellular context of axons and might explain the linked phenotypes we observed for
the two factors. This was further supported by their co-dependence for achieving prominent
plus-end localisation:

We already reported that loss of Msps causes severe Eb1 depletion and comet shortening at
MT plus-ends (Fig1. 1B, D and 1-S1B). This is likely due to the role of Msps as a tubulin
polymerase (Brouhard *et al*, 2008) which helps to sustain a prominent GTP-tubulin cap to which
EB1 can bind (Zanic *et al*., 2013). To test this, we stained *msps^1/1* mutant neurons for GTP-
tubulin, and found that GTP caps displayed reduced lengths comparable to the shortened Eb1
comets (Figs. 1J *vs.* 5-S2C). This result, together with our finding of reduced comet velocity in
msps^1/1 mutant neurons (Fig. 1K), is consistent with the hypothesis that Msps-dependent
polymerisation increases Eb1 amounts at MT plus-ends (see Discussion).

Vice versa, we expressed Msps::GFP in *Eb1^04524/04524* mutant neurons, which likewise revealed
severe depletion of Msps at growing MT plus-ends when compared to controls (Fig. 4C,C';
Movies M2 and M3). These results suggest that Msps/XMAP215 can bind MT plus-ends
independently, but that its binding is enhanced if Eb1 is present (see also Maurer *et al*., 2014;
Zanic *et al*., 2013). One mechanism through which Eb1 might recruit Msps/XMAP215 at MT
plus-ends is through adaptors such as SLAIN in vertebrates or TACC (Transforming acidic
coiled-coil protein) or Sentin in non-neuronal *Drosophila* cells (Brouhard *et al*., 2008; Lee *et al*,
2001; Li *et al*, 2011; Li *et al*., 2012; Lowery *et al*., 2013; Tang *et al*, 2020; van der Vaart *et al*,
2012). However, our functional studies using homozygous mutant neurons carrying *tacc^1* and
sentin^AB loss-of-function mutant alleles failed to display axon shortening or MT curling, thus
arguing against a prominent role of these factors during Eb1-dependent Msps recruitment
(details in Fig.4-S1A,B). To further substantiate these findings, we generated a *msps^ΔCterm*-*GFP*
construct which lacks the C-terminal domain essential for the interaction with adaptors (Fig.4D;
Fox *et al*, 2014; Mortuza *et al*, 2014). When *msps^ΔCterm*-*GFP* or *msps^FL*-*GFP* full length controls
were transfected into *msps^A/146* mutant neurons, we found that both protein variants were
similarly able to improve axon length and MT curling defects (Fig.4D,D'), thus likewise arguing
against the requirement of adaptors to recruit Msps to MT plus-ends in fly neurons.

In conclusion, our data clearly demonstrate that Eb1 and Msps require each other to achieve
prominent MT plus-end localisation in axons: Msps likely maintains Eb1 at MT plus-ends
through promoting GTP-cap formation, as is supported by our findings with GTP-tubulin
staining. *Vice versa,* Eb1-mediated recruitment of Msps might involve roles of Eb1 in the
structural maturation of MT plus-ends (see Discussion; Maurer *et al*., 2014; Zanic *et al*., 2013).

Tau promotes Eb1 pools at MT plus-ends by outcompeting it from lattice binding

Similar to Msps deficiency, we found that also loss of Tau leads to a reduction of Eb1 comet
sizes in both *Drosophila* (Fig.1C,J and 1-S2A,B) and *Xenopus* neurons (Fig.3-S1G). In wild-
type fly neurons, Tau localises along MT lattices and appears not to extend into the Eb1 comet
at the MT plus-end (Fig.5A'A''). This distribution is consistent with reports that Tau and Eb1 do
not overlap at MT plus-ends due to their respective higher affinities for GDP- and GTP-tubulin
(Castle *et al*, 2020; Duan *et al*, 2017), suggesting that Tau promotes Eb1 plus-end localisation
through indirect mechanisms not involving their physical interaction.

We noticed that the reduction of Eb1 comet sizes in Tau-deficient neurons is accompanied by
a 20% increase in the intensity of Eb1 along MT lattices (Fig.5C,E for 6 day pre-culture, Fig.5-
S2A for embryonic cultures). This effect is specific for *tau* and not observed in *msps$^{A/A}$* mutant
neurons (Fig. 5-S2A,B). Tau has previously been shown to protect MTs against Katanin-induced
damage (Qiang *et al*, 2006). Increased MT repair upon loss of Tau could therefore promote the
incorporation of GTP-tubulin along the lattice which would, in turn, recruit Eb1 (Vemu *et al*,
2018). However, MT lattices in Tau-deficient neurons did not show obvious increases in GTP-
tubulin (Fig. 5-S1B, D), arguing against the repair hypothesis. In the same specimens, GTP-
tubulin amounts at MT plus-ends were clearly reduced, consistent with the observed Eb1 comet
depletion in Tau-deficient neurons (Figs. 1J *vs*. 5-S1A-C).

We observed that over-expression of Tau in wild-type neurons decreased Eb1 levels along MT
lattices even further (Fig. 5-S1E), and similar observations were reported *in vitro* with
mammalian versions of the two proteins (Ramirez-Rios *et al*, 2016). We hypothesised therefore
that Tau may competitively prevent lattice-binding of Eb1, as similarly observed for other MAPs
(Qiang *et al*, 2018): if Tau is absent, MT lattices would therefore turn into a sink for Eb1 and
sequester Eb1 pools away from MT plus-ends. Such a mechanism could be particularly relevant
in narrow axons which display a high relative density of MTs (Prokop, 2020). In support of our
hypothesis, we found that *tau$^{KO/KO}$* mutant neurons overexpressing Eb1::GFP showed even
greater Eb1 intensity along MT lattices, but also replenished Eb1 amounts at MT plus-ends (Fig.
5D-F). This treatment was sufficient to suppress Tau-deficient phenotypes, as reflected in the
recovery of Eb1 comet dynamics (improved velocity and lifetime) and strongly reduced MT
curling ('Eb1-GFP' in Fig. 5G-I).

These effects were specific for Tau and Eb1: Firstly, no rescue of *tau$^{KO/KO}$* mutant phenotypes
was observed when expressing Shot-Ctail::GFP (as used in Fig.1K,L) to track MT plus-ends
without altering Eb1 levels ('Ctail-GFP' in Fig.5G,H). Secondly, Eb1::GFP could not restore MT
plus-end dynamics in *msps$^{A/A}$* mutant neurons (Fig. 5-S2D,E), a finding that provides important
insights also into the hierarchical relationships between Eb1, Msps and Tau (see Discussion).

In conclusion, we propose that Tau contributes to MT polymerisation dynamics and MT
organisation indirectly, through preventing that Eb1 is sequestered away from MT plus-ends.
The fact that Eb1::GFP can rescue *tau* mutant phenotypes including their MT curling, further
highlights Eb1's crucial role in MT bundle promotion (compare Fig.2C, Fig.2-S2) and may
suggest it even as a potential therapeutic target (see Discussion).

An Eb1- and spectraplakin-dependent guidance mechanism explains roles of the functional trio
in MT bundle organisation

Since Eb1 amounts at MT plus-ends inversely correlate with MT curling (Figs.2C and 2-S2),
bundle deterioration in either Eb1-, Msps- or Tau-deficient neurons might be a consequence of
their negative impacts on Eb1 localisation. Eb1 at polymerising MT plus-ends has been
proposed to recruit the C-terminus of Shot which simultaneously binds cortical F-actin via its N-
terminus (Fig.6A). Through this cross-linking activity, Shot has been proposed to guide the
extension of MTs along the axonal surface into parallel bundles (Fig.6F); accordingly, depletion
of either Shot or Eb1 causes MT curling; Alves-Silva *et al.*, 2012; Sánchez-Soriano *et al.*, 2009;
Voelzmann *et al*, 2017); Figs.1D, 6E).

In support of this guidance hypothesis, we found that severe MT curling observed in $Eb1^{+/-}$
$msps^{+/-}$ $tau^{+/-}$ triple-heterozygous mutant neurons (which have strongly reduced Eb1 comet
amounts; Fig.2A'') was significantly improved from 7.3-fold (with GFP-expression) to 1.4-fold,
when over-expressing full length Shot-FL::GFP (Fig.6B-C; both compared to wild-type controls).
This rescue could also be mediated through Eb1-independent roles of Shot in MT stabilisation
mediated by the C-terminal Gas2-related domain (Fig.6A; Alves-Silva *et al.*, 2012). We therefore
repeated the rescue experiments with two different Shot variants that maintain MT-stabilising
activity whilst specifically abolishing actin-Eb1 cross-linkage (Fig.6A,F): (1) Shot$^{\Delta ABD}$::GFP lacks
the N-terminal calponin homology domain required for interaction with the actin cortex; (2)
Shot^{3MtLS*}::GFP carries mutations in the C-terminal SxIP motifs required for Eb1 binding. Both
these Shot variants failed to rescue MT curling in $Eb1^{04524/+}$ $msps^{1/+}$ $tau^{KO/+}$ triple-heterozygous
mutant neurons (Fig.6B,C), hence lending further support to the guidance hypothesis based on
Eb1-Shot interaction. This conclusion is further substantiated by the finding that MT curling
observed in $Eb1^{04524/04524}$ mutant neurons is not enhanced in $Eb1^{04524/04524}$ $shot^{3/3}$ double-mutant
neurons, neither in normal embryonic nor in embryonic pre-cultured neurons (Figs. 6D, E).

We conclude that co-operative promotion of MT polymerisation through Eb1, Msps and Tau
appears to perform its additional function in axon bundle organisation through Shot-mediated
MT guidance downstream of Eb1. In this way, the three proteins uphold the numbers/mass as
well as the bundled organisation of MTs as two key features underpinning the formation and
long-term maintenance of axons.

Discussion

New understanding of the role and regulation of MT polymerisation and guidance in axons

Understanding the machinery of MT polymerisation is of utmost importance in axons where MTs
form loose bundles of enormous lengths which are essential for axonal morphogenesis and
serve as life-sustaining transport highways (Prokop, 2020). These bundles must be maintained
in functional state for up to a century in humans (Hahn *et al.*, 2019). To achieve this, MT
polymerisation is required to generate MTs *de novo*, repair or replace them. The underpinning
machinery is expected to be complex (Voelzmann *et al.*, 2016), but deciphering the involved
mechanisms will pay off by delivering new strategies for tackling developmental and
degenerative axon pathologies (Prokop, 2021).

Here we made important advances to this end. Having screened through 13 candidates (this
work and Beaven *et al.*, 2015), we found the three factors Eb1, Msps and Tau to stand out by
expressing the same combination of phenotypes, and by displaying functional interaction in both
Drosophila and *Xenopus* neurons. We found that their functions are not only important to
maintain MT polymerisation, but also to align MTs into parallel arrangements, thus contributing
in two ways to MT bundle formation and maintenance, both in culture and *in vivo*. The observed

impact on MT organisation is also consistent with roles of XMAP215 during MT guidance in
growth cones of frog neurons (Slater *et al.*, 2019).

Our data reveal that various mechanisms observed *in vitro* or in non-neuronal contexts, apply
in the biological context of axons, which was unpredictable for two reasons: Firstly, of the three
proteins only the human tau homologue has OMIM®-listed links to human axonopathies, and
these do not necessarily relate to MT polymerisation (Morris *et al.*, 2011). Secondly, axons and
non-neuronal cells can display significant mechanistic deviations as shown for CLIP170/190
(Beaven *et al.*, 2015) and for the MT localisation of Msps which is facilitated by Sentin or dTACC
in non-neuronal cells, dendrites and *in vitro* (Lee *et al.*, 2001; Li *et al.*, 2011; Li *et al.*, 2012; Tang
et al., 2020) but seemingly not in axons (Fig.4S1). However, other mechanisms we observed in
axons matched previous reports: (1) the complementary binding preferences of Eb1 and Tau
for GTP-/GDP-tubulin (Castle *et al.*, 2020; Zanic *et al*, 2009); (2) the mutual enhancement of
Eb1 and XMAP215/Msps (Kronja *et al*, 2009; Li *et al.*, 2012; Zanic *et al.*, 2013); (3) the
correlation of GTP cap size with comet velocity (Roostalu *et al.*, 2020). Furthermore, we
observed that depletion of α1-tubulin in neurons mutant for *atub84B* or *stathmin* (a promoter of
tubulin availability; (Duncan *et al.*, 2013; Manna *et al.*, 2007); J.F. and A.V., unpublished) affects
comet numbers but not comet dynamics (Fig.1-S1); this is consistent with observations that MT
nucleation *in vitro* is far more sensitive to tubulin levels than polymerisation (Consolati *et al*,
2020).

<u>Eb1 and XMAP215/Msps are core factors promoting MT polymerisation and guidance</u>

Apart from demonstrating the relevance of various molecular mechanisms in the context of
axonal MT regulation, our work provides key insights as to how they integrate into one
consistent mechanistic model of biological function (summarised in Fig.7).

The TOG-domain protein XMAP215/Msps is relevant for neuronal morphogenesis in fly and
Xenopus (Lowery *et al.*, 2013; Tang *et al.*, 2020), likely through its expected function as a MT
polymerase (Al-Bassam & Chang, 2011; Brouhard *et al.*, 2008; Fox *et al.*, 2014; Howard &
Hyman, 2009; Li *et al.*, 2012; Zanic *et al.*, 2013). In contrast, *Drosophila* and vertebrate Eb
proteins are only moderate promoters of MT polymerisation *in vitro* (Li *et al.*, 2012; Ramirez-
Rios *et al.*, 2016; Zanic *et al.*, 2013) and references within), but rather act as scaffolds
(Akhmanova & Steinmetz, 2015). Conserved binding partners of Eb proteins are the
spectraplakins which can guide extending MT plus-ends along actin networks in axons and non-
neuronal cells (Alves-Silva *et al.*, 2012; Voelzmann *et al.*, 2017; Wu *et al*, 2008).

We propose therefore (see Fig.7A) that Eb1 is the key mediator of MT guidance into bundles
(as supported by data throughout this work; Figs.2C, 2-S2, 6), and Msps the key promoter of
MT polymerisation. To execute these functions, both proteins depend on each other: MT plus-
end localisation of Msps is reduced upon loss of Eb1 (Fig.7B) and *vice versa* (Fig.7C). This
mutual dependency is unlikely to involve their physical interaction, since MT plus-end
localisation of Eb1 is known to occur tens of nanometres behind XMAP215 (Maurer *et al.*, 2014;
Zanic *et al.*, 2013), as seems to be the case also for axonal MTs (Fig.4A). Furthermore, our
data do not support an obvious role of the Sentin or dTACC adaptors in mediating Eb1-
XMAP215 interactions (Figs.4D-D'' and 4-S1).

Potential indirect mechanisms explaining this co-dependency are provided by the promotion of
MT polymerisation through XMAP215/Msps which maintains a prominent GTP-cap (Fig.5-S1)
that, in turn, mediates Eb1 binding (see also (Maurer *et al.*, 2014; Zanic *et al.*, 2013); Fig.7C).

Restricted GTP-cap formation as a limiting factor for Eb1 binding would also explain why Eb1
over-expression fails to rescue Msps-deficient phenotypes (Fig.5-S2D-F). *Vice versa*, Eb
proteins promote lateral protofilament contacts which could assist in sheet formation at the very
plus tip (Fig.7B), thus facilitating the binding of XMAP215/Msps (Maurer *et al.*, 2014; Zanic *et*
al., 2013).

<u>Tau contributes through an indirect mechanism of competitive binding to MT lattices</u>

Tau and Map1b/Futsch are known to bind along MT lattices, to promote MT polymerisation *in*
vitro, and to enhance axon growth in mouse and fly neurons through mechanisms that remain
unclear (Brandt & Lee, 1993; Caceres & Kosik, 1990; Cleveland *et al*, 1977; DiTella *et al*, 1996;
Drechsel *et al.*, 1992; Hummel *et al.*, 2000; Kadavath *et al*, 2015; Kiris *et al*, 2010; Levy *et al*,
2005; Liu *et al.*, 2015; Panda *et al*, 1995; Ramirez-Rios *et al.*, 2016; Takei *et al*, 2000; Tymanskyj
et al, 2012; Villarroel-Campos & Gonzalez-Billault, 2014).

In our cellular model, loss of Map1b/Futsch has no obvious effects, whereas Tau shares all
assessed loss-of-function mutant phenotypes with Msps and Eb1, though with weaker
expression. Of these phenotypes, reduced MT plus-end localisation of Eb proteins upon loss of
Tau function (Figs.1C,J) was likewise reported for frog neurons (Fig.3-S1G-I), N1E-115 mouse
neuroblastoma cells and primary mouse cortical neurons (Sayas *et al*, 2015).

One proposed mechanism involves direct interaction where Tau recruits Eb proteins at MT plus
ends (Sayas *et al.*, 2015), consistent with other reports that Tau can bind Eb1 (Buey *et al*, 2011;
Duan *et al.*, 2017). However, further reports argue against overlap of Tau and Eb1 and rather
show that Eb1 and Tau have complementary preferences for GTP- and GDP-tubulin,
respectively (Castle *et al.*, 2020; Duan *et al.*, 2017; Zanic *et al.*, 2009); this is also consistent
with our data (Fig.5A). Furthermore, a recruitment model is put in question by our finding that
Eb1 lattice localisation increases rather than decreases upon loss of Tau (Fig.5E), as similarly
observed for mammalian tau and Eb1 *in vitro* (Ramirez-Rios *et al.*, 2016).

Therefore, we propose a different mechanism based on competitive binding where Tau's
preferred binding to GDP-tubulin along the lattice prevents Eb1 localisation (Fig.5), comparable
to Tau's role in preventing MAP6 from binding in certain regions of the MT lattice (Baas & Qiang,
2019; Qiang *et al.*, 2018). Given the high density of MTs in the narrow space of axons, lattice
binding could generate a sink large enough to reduce Eb1 levels at MT plus-ends, and our
rescue experiments with Eb1::GFP overexpression strongly support this notion (Fig.5). In this
way, loss of Tau generates a condition comparable to a modest Eb1 loss-of-function mutant
phenotype, thus explaining why Tau shares its repertoire of loss-of-function phenotypes with
Msps and Eb1, but with more modest presentation. This competition mechanism might apply in
axons with high MT density, for example explain the reduction in axonal MT numbers upon Tau
deficiency in *C. elegans* (Krieg *et al*, 2017); it might be less relevant in larger diameter axons of
vertebrates where MT densities are low (Prokop, 2020).

<u>Main conclusions and future perspectives</u>

Here we have used a standardised *Drosophila* neuron system amenable to combinatorial
genetics to gain understanding of MT regulation at the cellular level in axons. We propose a
consistent mechanistic model that can integrate all our data, mechanisms reported in the
literature, and our previous mechanistic model explaining Eb1/Shot-mediated MT guidance

(Alves-Silva *et al.*, 2012; Qu *et al.*, 2019). This understanding offers new opportunities to investigate the mechanisms behind other important observations.

For example, the presence of an axonal sleeve of cortical actin/spectrin networks was shown to be important to maintain MT polymerisation, likely relevant in certain axonopathies (Prokop, 2021; Qu *et al.*, 2017); the underlying mechanisms are now far easier to dissect. As another example, we found that loss of either Eb1, XMAP215/Msps or Tau all caused a reduction in comet numbers, consistent with reports of nucleation-promoting roles of XMAP215 (Flor-Parra *et al*, 2018; Roostalu *et al*, 2015; Thawani *et al*, 2018; Wieczorek *et al*, 2015). This might offer opportunities to investigate how MT numbers can be regulated in reproducible, neuron/axon-specific ways, thus addressing a fundamental aspect of axon morphology (Prokop, 2020).

By gradually assembling molecular mechanisms into regulatory networks that can explain axonal MT regulation at the cellular level, i.e. the level at which diseases become manifest, our studies come closer to explaining axonal pathologies which can then form the basis for the development of remedial strategies (Hahn *et al.*, 2019).

Acknowledgements

This work was made possible through support by the BBSRC to A.P (BB/I002448/1, BB/P020151/1, BB/L000717/1, BB/M007553/1) to N.S.S. (BB/M007456/1, BB/R018960/1), by the Leverhulme Trust to I.H. (ECF-2017-247), by the German Research Council (DFG) to A.V. (VO 2071/1-1), by NIH to L.A.L (R01 MH109651), and a postdoctoral fellowship from CONICYT to P.G.S. The Manchester Bioimaging Facility microscopes used in this study were purchased with grants from the BBSRC, The Wellcome Trust and The University of Manchester Strategic Fund. The Fly Facility has been supported by funds from The University of Manchester and the Wellcome Trust (087742/Z/08/Z). We thank Hiro Ohkura for kindly providing DmEb1 antibody and unpublished mutant alleles of *msps*. Stocks obtained from the Bloomington *Drosophila* Stock Center (NIH P40OD018537) were used in this study.

References

Adalbert R, Coleman MP (2012) Axon pathology in age-related neurodegenerative disorders. *Neuropathol Appl Neurobiol* 39: 90–108

Aher A, Rai D, Schaedel L, Gaillard J, John K, Liu Q, Altelaar M, Blanchoin L, Thery M, Akhmanova A (2020) CLASP mediates microtubule repair by restricting lattice damage and regulating tubulin incorporation. *Current Biology*

Akhmanova A, Steinmetz MO (2015) Control of microtubule organization and dynamics: two ends in the limelight. *Nat Rev Mol Cell Biol* 16: 711–726

Al-Bassam J, Chang F (2011) Regulation of microtubule dynamics by TOG-domain proteins XMAP215/Dis1 and CLASP. *Trends Cell Biol* 21: 604-614

Al-Bassam J, Kim H, Brouhard G, van Oijen A, Harrison SC, Chang F (2010) CLASP promotes microtubule rescue by recruiting tubulin dimers to the microtubule. *Dev Cell* 19: 245-258

Alves-Silva J, Sánchez-Soriano N, Beaven R, Klein M, Parkin J, Millard T, Bellen H, Venken KJT, Ballestrem C, Kammerer RA *et al* (2012) Spectraplakins promote microtubule-mediated axonal growth by functioning as structural microtubule-associated proteins and EB1-dependent +TIPs (Tip Interacting Proteins). *J Neurosci* 32: 9143-9158

Avery L, Wasserman S (1992) Ordering gene function: the interpretation of epistasis in
regulatory hierarchies. *Trends in genetics : TIG* 8: 312-316

Baas PW, Qiang L (2019) Tau: it's not what you think. *Trends Cell Biol* 29: 452-461

Beaven R, Dzhindzhev NS, Qu Y, Hahn I, Dajas-Bailador F, Ohkura H, Prokop A (2015)
Drosophila CLIP-190 and mammalian CLIP-170 display reduced microtubule plus end
association in the nervous system. *Mol Biol Cell* 26: 1491-1508

Bowes JB, Snyder KA, Segerdell E, Jarabek CJ, Azam K, Zorn AM, Vize PD (2009) Xenbase:
gene expression and improved integration. *Nucleic Acids Research* 38: D607-D612

Brandt R, Lee G (1993) The balance between tau protein's microtubule growth and nucleation
activities: implications for the formation of axonal microtubules. *J Neurochem* 61: 997-1005

Brittle AL, Ohkura H (2005) Mini spindles, the XMAP215 homologue, suppresses pausing of
interphase microtubules in *Drosophila. EMBO J* 24: 1387-1396

Brouhard GJ, Stear JH, Noetzel TL, Al-Bassam J, Kinoshita K, Harrison SC, Howard J, Hyman
AA (2008) XMAP215 is a processive microtubule polymerase. *Cell* 132: 79-88

Buey RM, Mohan R, Leslie K, Walzthoeni T, Missimer JH, Menzel A, Bjelic S, Bargsten K,
Grigoriev I, Smal I *et al* (2011) Insights into EB1 structure and the role of its C-terminal domain
for discriminating microtubule tips from the lattice. *Mol Biol Cell* 22: 2912-2923

Burnouf S, Gronke S, Augustin H, Dols J, Gorsky MK, Werner J, Kerr F, Alic N, Martinez P,
Partridge L (2016) Deletion of endogenous Tau proteins is not detrimental in *Drosophila. Sci*
Rep 6: 23102

Caceres A, Kosik KS (1990) Inhibition of neurite polarity by tau antisense oligonucleotides in
primary cerebellar neurons. *Nature* 343: 461-463

Calkins DJ (2013) Age-Related Changes in the Visual Pathways: Blame It on the AxonAge-
Related Changes in the Visual Pathways. *Invest Ophthalmol Vis Sci* 54: ORSF 37-41

Castle BT, McKibben KM, Rhoades E, Odde DJ (2020) Tau avoids the GTP cap at growing
microtubule plus ends. *bioRxiv*: 2019.2012.2031.891234

Cleveland DW, Hwo SY, Kirschner MW (1977) Purification of tau, a microtubule-associated
protein that induces assembly of microtubules from purified tubulin. *J Mol Biol* 116: 207-225

Consolati T, Locke J, Roostalu J, Chen ZA, Gannon J, Asthana J, Lim WM, Martino F, Cvetkovic
MA, Rappsilber J *et al* (2020) Microtubule Nucleation Properties of Single Human γTuRCs
Explained by Their Cryo-EM Structure. *Developmental Cell* 53: 603-617.e608

Dimitrov A, Quesnoit MI, Moutel S, Cantaloube I, Po?s C, Perez F (2008) Detection of GTP-
tubulin conformation *in vivo* reveals a role for GTP remnants in microtubule rescues. *Science*
322: 1353-1356

DiTella MC, Feiguin F, Carri N, Kosik KS, Caceres A (1996) MAP-1B/TAU functional
redundancy during laminin-enhanced axonal growth. *J Cell Sci* 109 (Pt 2): 467-477

Doerflinger H, Benton R, Shulman JM, St Johnston D (2003) The role of PAR-1 in regulating
the polarised microtubule cytoskeleton in the Drosophila follicular epithelium. *Development*
130: 3965-3975

Drechsel DN, Hyman AA, Cobb MH, Kirschner MW (1992) Modulation of the dynamic instability
of tubulin assembly by the microtubule-associated protein tau. *Mol Biol Cell* 3: 1141-1154

Duan AR, Jonasson EM, Alberico EO, Li C, Scripture JP, Miller RA, Alber MS, Goodson HV
(2017) Interactions between tau and different conformations of tubulin: implications for tau
function and mechanism. *J Mol Biol* 429: 1424-1438

Duncan IW, Kaufman TC (1975) Cytogenic analysis of chromosome 3 in *Drosophila*
melanogaster: mapping of the proximal portion of the right arm. *Genetics* 80: 733-752

Duncan JE, Lytle NK, Zuniga A, Goldstein LS (2013) The microtubule regulatory protein
Stathmin is required to maintain the integrity of axonal microtubules in *Drosophila*. *PLoS One*
8: e68324

Elliott SL, Cullen CF, Wrobel N, Kernan MJ, Ohkura H (2005) EB1 is essential during *Drosophila*
development and plays a crucial role in the integrity of chordotonal mechanosensory organs.
Mol Biol Cell 16: 891-901

Flor-Parra I, Iglesias-Romero AB, Chang F (2018) The XMAP215 ortholog Alp14 promotes
microtubule nucleation in fission yeast. *Curr Biol* 28: 1681-1691.e1684

Fox JC, Howard AE, Currie JD, Rogers SL, Slep KC (2014) The XMAP215 family drives
microtubule polymerization using a structurally diverse TOG array. *Mol Biol Cell* 25: 2375-
2392

Gasic I, Mitchison TJ (2019) Autoregulation and repair in microtubule homeostasis. *Curr Opin*
Cell Biol 56: 80-87

Gergely F, Kidd D, Jeffers K, Wakefield JG, Raff JW (2000) D-TACC: a novel centrosomal
protein required for normal spindle function in the early Drosophila embryo. *EMBO J* 19: 241-
252

Gluszek AA, Cullen CF, Li W, Battaglia RA, Radford SJ, Costa MF, McKim KS, Goshima G,
Ohkura H (2015) The microtubule catastrophe promoter Sentin delays stable kinetochore-
microtubule attachment in oocytes. *J Cell Biol* 211: 1113-1120

Grieder NC, de Cuevas M, Spradling AC (2000) The fusome organizes the microtubule network
during oocyte differentiation in Drosophila. *Development* 127: 4253-4264

Hahn I, Voelzmann A, Liew Y-T, Costa-Gomes B, Prokop A (2019) The model of local axon
homeostasis - explaining the role and regulation of microtubule bundles in axon maintenance
and pathology *Neural Dev* 14: 10.1186/s13064-13019-10134-13060

Honnappa S, Gouveia SM, Weisbrich A, Damberger FF, Bhavesh NS, Jawhari H, Grigoriev I,
van Rijssel FJ, Buey RM, Lawera A *et al* (2009) An EB1-binding motif acts as a microtubule
tip localization signal. *Cell* 138: 366-376

Howard J, Hyman AA (2009) Growth, fluctuation and switching at microtubule plus ends. *Nat*
Rev Mol Cell Biol 10: 569-574

Hummel T, Krukkert K, Roos J, Davis G, Klämbt C (2000) Drosophila Futsch/22C10 is a
MAP1B-like protein required for dendritic and axonal development. *Neuron* 26: 357-370

Inoue YH, do Carmo Avides M, Shiraki M, Deak P, Yamaguchi M, Nishimoto Y, Matsukage A,
Glover DM (2000) Orbit, a novel microtubule-associated protein essential for mitosis in
Drosophila melanogaster. *J Cell Biol* 149: 153-166

Jenkins BV, Saunders HAJ, Record HL, Johnson-Schlitz DM, Wildonger J (2017) Effects of
mutating alpha-tubulin lysine 40 on sensory dendrite development. *J Cell Sci* 130: 4120-4131

Kadavath H, Hofele RV, Biernat J, Kumar S, Tepper K, Urlaub H, Mandelkow E, Zweckstetter
M (2015) Tau stabilizes microtubules by binding at the interface between tubulin heterodimers.
Proc Natl Acad Sci 112: 7501-7506

Karimi K, Fortriede JD, Lotay VS, Burns KA, Wang DZ, Fisher ME, Pells TJ, James-Zorn C,
Wang Y, Ponferrada V G *et al* (2017) Xenbase: a genomic, epigenomic and transcriptomic
model organism database. *Nucleic Acids Res* 46: D861-D868

Kiris E, Ventimiglia D, Feinstein SC (2010) Quantitative analysis of MAP-mediated regulation of
microtubule dynamic instability in vitro focus on Tau. *Methods Cell Biol* 95: 481-503

Kolodziej PA, Jan LY, Jan YN (1995) Mutations that affect the length, fasciculation, or ventral
orientation of specific sensory axons in the Drosophila embryo. *Neuron* 15: 273-286

Krieg M, Stühmer J, Cueva JG, Fetter R, Spilker K, Cremers D, Shen K, Dunn AR, Goodman
MB (2017) Genetic defects in β-spectrin and tau sensitize C. elegans axons to movement-
induced damage via torque-tension coupling. *Elife* 6: e20172

Kronja I, Kruljac-Letunic A, Caudron-Herger M, Bieling P, Karsenti E (2009) XMAP215-EB1
interaction is required for proper spindle assembly and chromosome segregation in *Xenopus*
egg extract. *Mol Biol Cell* 20: 2684-2696

Lee MJ, Gergely F, Jeffers K, Peak-Chew SY, Raff JW (2001) Msps/XMAP215 interacts with
the centrosomal protein D-TACC to regulate microtubule behaviour. *Nat Cell Biol* 3: 643-649

Levy SF, Leboeuf AC, Massie MR, Jordan MA, Wilson L, Feinstein SC (2005) Three- and four-
repeat tau regulate the dynamic instability of two distinct microtubule subpopulations in
qualitatively different manners. Implications for neurodegeneration. *J Biol Chem* 280: 13520-
13528

Li W, Miki T, Watanabe T, Kakeno M, Sugiyama I, Kaibuchi K, Goshima G (2011) EB1 promotes
microtubule dynamics by recruiting Sentin in *Drosophila* cells. *J Cell Biol* 193: 973-983

Li W, Moriwaki T, Tani T, Watanabe T, Kaibuchi K, Goshima G (2012) Reconstitution of dynamic
microtubules with *Drosophila* XMAP215, EB1, and Sentin. *J Cell Biol* 199: 849-862

Liu Y, Wang C, Destin G, Szaro BG (2015) Microtubule-associated protein tau promotes
neuronal class II beta-tubulin microtubule formation and axon elongation in embryonic
Xenopus laevis. Eur J Neurosci 41: 1263-1275

Lowery LA, Faris AER, Stout A, Van Vactor D (2012) Neural explant cultures from *Xenopus*
laevis. JoVE: e4232

Lowery LA, Stout A, Faris AE, Ding L, Baird MA, Davidson MW, Danuser G, Van Vactor D
(2013) Growth cone-specific functions of XMAP215 in restricting microtubule dynamics and
promoting axonal outgrowth. *Neural Dev* 8: 22

Luo L, Liao YJ, Jan LY, Jan YN (1994) Distinct morphogenetic functions of similar small
GTPases: Drosophila Drac1 is involved in axonal outgrowth and myoblast fusion. *Genes Dev*
8: 1787-1802

Manna T, Grenningloh G, Miller HP, Wilson L (2007) Stathmin family protein SCG10
differentially regulates the plus and minus end dynamics of microtubules at steady state *in*
vitro: implications for its role in neurite outgrowth. *Biochemistry* 46: 3543-3552

Manna T, Thrower DA, Honnappa S, Steinmetz MO, Wilson L (2009) Regulation of microtubule
dynamic instability in vitro by differentially phosphorylated stathmin. *J Biol Chem* 284: 15640-
15649

Marner L, Nyengaard JR, Tang Y, Pakkenberg B (2003) Marked loss of myelinated nerve fibers
in the human brain with age. *J Comp Neurol* 462: 144-152

Maurer SP, Cade NI, Bohner G, Gustafsson N, Boutant E, Surrey T (2014) EB1 accelerates two
conformational transitions important for microtubule maturation and dynamics. *Curr Biol* 24:
372-384

Morris M, Maeda S, Vossel K, Mucke L (2011) The many faces of tau. *Neuron* 70: 410-426

Mortuza GB, Cavazza T, Garcia-Mayoral MF, Hermida D, Peset I, Pedrero JG, Merino N, Blanco
FJ, Lyngsø J, Bruix M *et al* (2014) XTACC3–XMAP215 association reveals an asymmetric
interaction promoting microtubule elongation. *Nature Communications* 5: 5072

Nieuwkoop, J. F (1994) *Normal table of Xenopus laevis (Daudin): A systematical and*
chronological survey of the development from the fertilized egg till the end of metamorphosis.
Garland Pub. Inc. , New York

O'Neill EM, Rebay I, Tjian R, Rubin GM (1994) The activities of two Ets-related transcription
factors required for Drosophila eye development are modulated by the Ras/MAPK pathway.

Cell 78: 137-147.

Panda D, Goode BL, Feinstein SC, Wilson L (1995) Kinetic stabilization of microtubule
dynamics at steady state by tau and microtubule-binding domains of tau. *Biochemistry* 34:
11117-11127

Prokop A (2013) A rough guide to Drosophila mating schemes. *figshare*:
dx.doi.org/10.6084/m6089.figshare.106631

Prokop A (2020) Cytoskeletal organization of axons in vertebrates and invertebrates. *J Cell Biol*
219: e201912081

Prokop A (2021) A common theme for axonopathies? The dependency cycle of local axon
homeostasis. *Cytoskeleton* 78: 52–63

Prokop A, Beaven R, Qu Y, Sánchez-Soriano N (2013) Using fly genetics to dissect the
cytoskeletal machinery of neurons during axonal growth and maintenance. *J Cell Sci* 126:
2331-2341

Prokop A, Küppers-Munther B, Sánchez-Soriano N (2012) Using primary neuron cultures of
Drosophila to analyse neuronal circuit formation and function. *The making and un-making of*
neuronal circuits in Drosophila 69: 225-247

Prokop A, Meinertzhagen IA (2006) Development and structure of synaptic contacts in
Drosophila. *Semin Cell Dev Biol* 17: 20-30

Qiang L, Sun X, Austin TO, Muralidharan H, Jean DC, Liu M, Yu W, Baas PW (2018) Tau does
not stabilize axonal microtubules but rather enables them to have long labile domains. *Curr*
Biol 28: 2181-2189 e2184

Qiang L, Yu W, Andreadis A, Luo M, Baas PW (2006) Tau protects microtubules in the axon
from severing by katanin. *J Neurosci* 26: 3120-3129

Qu Y (2015) Novel concepts of microtubule regulation during axon growth and maintenance.
Faculty of Life Sciences PhD: 205

Qu Y, Hahn I, Lees M, Parkin J, Voelzmann A, Dorey K, Rathbone A, Friel C, Allan V, Okenve
Ramos P *et al* (2019) Efa6 protects axons and regulates their growth and branching by
inhibiting microtubule polymerisation at the cortex. *eLife* 8: e50319

Qu Y, Hahn I, Webb SED, Pearce SP, Prokop A (2017) Periodic actin structures in neuronal
axons are required to maintain microtubules. *Mol Biol Cell* 28 296-308

Ramirez-Rios S, Denarier E, Prezel E, Vinit A, Stoppin-Mellet V, Devred F, Barbier P, Peyrot V,
Sayas CL, Avila J *et al* (2016) Tau antagonizes end-binding protein tracking at microtubule
ends through a phosphorylation-dependent mechanism. *Mol Biol Cell* 27: 2924-2934

Rogers SL, Rogers GC, Sharp DJ, Vale RD (2002) Drosophila EB1 is important for proper
assembly, dynamics, and positioning of the mitotic spindle. *J Cell Biol* 158: 873-884

Roostalu J, Cade NI, Surrey T (2015) Complementary activities of TPX2 and chTOG constitute
an efficient importin-regulated microtubule nucleation module [corrigendum: same issue,
p.1512]. *Nat Cell Biol* 17: 1422-1434

Roostalu J, Thomas C, Cade NI, Kunzelmann S, Taylor IA, Surrey T (2020) The speed of GTP
hydrolysis determines GTP cap size and controls microtubule stability. *eLife* 9

Sánchez-Soriano N, Gonçalves-Pimentel C, Beaven R, Haessler U, Ofner L, Ballestrem C,
Prokop A (2010) Drosophila growth cones: a genetically tractable platform for the analysis of
axonal growth dynamics. *Dev Neurobiol* 70: 58-71

Sánchez-Soriano N, Travis M, Dajas-Bailador F, Goncalves-Pimentel C, Whitmarsh AJ, Prokop
A (2009) Mouse ACF7 and Drosophila Short stop modulate filopodia formation and
microtubule organisation during neuronal growth. *J Cell Sci* 122: 2534-2542

Sayas CL, Tortosa E, Bollati F, Ramirez-Rios S, Arnal I, Avila J (2015) Tau regulates the
localization and function of End-binding proteins 1 and 3 in developing neuronal cells. *J*
Neurochem 133: 653-667

Schaedel L, John K, Gaillard J, Nachury MV, Blanchoin L, Thery M (2015) Microtubules self-
repair in response to mechanical stress. *Nat Mater* 14: 1156-1163

Sive HL, Grainger RM, Harland RM (2010) Microinjection of Xenopus oocytes. *Cold Spring Harb*
Protoc 2010: pdb prot5536

Slater PG, Cammarata GM, Samuelson AG, Magee A, Hu Y, Lowery LA (2019) XMAP215
promotes microtubule-F-actin interactions to regulate growth cone microtubules during axon
guidance. *J Cell Sci*: jcs.224311

Strumpf D, Volk T (1998) Kakapo, a novel *Drosophila* protein, is essential for the restricted
localization of the neuregulin-like factor, Vein, at the muscle-tendon junctional site. *J Cell Biol*
143: 1259-1270

Takei Y, Teng J, Harada A, Hirokawa N (2000) Defects in axonal elongation and neuronal
migration in mice with disrupted tau and map1b genes. *J Cell Biol* 150: 989-1000

Tang Q, Rui M, Bu S, Wang Y, Chew LY, Yu F (2020) A microtubule polymerase is required for
microtubule orientation and dendrite pruning in *Drosophila*. *EMBO J* 39: e103549

Thawani A, Kadzik RS, Petry S (2018) XMAP215 is a microtubule nucleation factor that
functions synergistically with the γ-tubulin ring complex. *Nature Cell Biology* 20: 575-585

Tymanskyj SR, Scales TM, Gordon-Weeks PR (2012) MAP1B enhances microtubule assembly
rates and axon extension rates in developing neurons. *Mol Cell Neurosci* 49: 110-119

van der Vaart B, Franker MA, Kuijpers M, Hua S, Bouchet BP, Jiang K, Grigoriev I, Hoogenraad
CC, Akhmanova A (2012) Microtubule Plus-End Tracking Proteins SLAIN1/2 and ch-TOG
Promote Axonal Development. *J Neurosci* 32: 14722-14728

Vemu A, Szczesna E, Zehr EA, Spector JO, Grigorieff N, Deaconescu AM, Roll-Mecak A (2018)
Severing enzymes amplify microtubule arrays through lattice GTP-tubulin incorporation.
Science 361: eaau1504

Villarroel-Campos D, Gonzalez-Billault C (2014) The MAP1B case: an old MAP that is new
again. *Dev Neurobiol* 74: 953-971

Voelzmann A, Hahn I, Pearce S, Sánchez-Soriano NP, Prokop A (2016) A conceptual view at
microtubule plus end dynamics in neuronal axons. *Brain Res Bulletin* 126: 226-237

Voelzmann A, Liew Y-T, Qu Y, Hahn I, Melero C, Sánchez-Soriano N, Prokop A (2017)
Drosophila Short stop as a paradigm for the role and regulation of spectraplakins. *Semin Cell*
Dev Biol 69: 40-57

Wieczorek M, Bechstedt S, Chaaban S, Brouhard GJ (2015) Microtubule-associated proteins
control the kinetics of microtubule nucleation. *Nat Cell Biol* 17: 907-916

Wu X, Kodama A, Fuchs E (2008) ACF7 regulates cytoskeletal-focal adhesion dynamics and
migration and has ATPase activity. *Cell* 135: 137–148

Yang N, Inaki M, Cliffe A, Rørth P (2012) Microtubules and Lis-1/NudE/dynein regulate invasive
cell-on-cell migration in Drosophila. *PLoS One* 7: e40632

Zanic M, Stear JH, Hyman AA, Howard J (2009) EB1 recognizes the nucleotide state of tubulin
in the microtubule lattice. *PloS one* 4: e7585-e7585

Zanic M, Widlund PO, Hyman AA, Howard J (2013) Synergy between XMAP215 and EB1
increases microtubule growth rates to physiological levels. *Nat Cell Biol* 15: 688–693

Figures

Fig. 1. Eb1, Msps and Tau share the same combination of axonal loss-of-function phenotypes in *Drosophila* primary neurons. **A-H)** Images of representative examples of embryonic primary neurons pre-cultured to 6 days (to deplete maternal gene product; see methods) and either immuno-stained for Eb1 (top) or for tubulin (bottom); neurons were either wild-type controls (ctrl) or carried the mutant alleles *msps^1*, *tau^{KO}* or *Eb1^{04524}* in homozygosis (from left to right); asterisks indicate cell bodies, black arrow heads the axon tips, white arrow heads point at areas of MT curling, dashed squares in A-D are shown as 3.5-fold magnified close-ups below each image with black arrows pointing at Eb1 comets; the axonal outline in D is indicated by a dotted line; scale bar in A represents 15 μm in all images. **I-N)** Quantification of different parameters (as indicated above each graph) obtained from pre-cultured embryonic primary neurons with the same genotypes as shown in A-H. Data were normalised to parallel controls (dashed horizontal line) and are shown as median ± 95% confidence interval (I-M) or mean ± SEM (N); data points in each plot, taken from at least two experimental repeats consisting of 3 replicates each; large open circles in graphs indicate median/mean of independent biological repeats. P-values obtained with Kruskall-Wallis ANOVA test for the different genotypes are indicated in each graph. For raw data see Tab. T1.

Fig. 1-S1. A candidate screen of axonal loss-of-function phenotypes in primary neurons. Graphs show extended data sets for four of the parameters displayed in Fig. 1 (indicated above each graph). Data points/bars representing mutant conditions for different genes are consistently colour-coded in all graphs, and conditions used are indicated below (6HIV, cultured from embryos for 6hrs; 6d/7d pre, cultured from embryos for 12hrs following 6 or 7 days pre-culture; L3, cultured from late larval CNS for 18hrs). Allele names are given as superscript: absence of slash indicates homozygous, presence of slash hetero-allelic conditions. Data were normalised to parallel controls (dashed horizontal line) and are shown as median ± 95% confidence interval (B,C) or mean ± SEM (A,D); data points from at least two experimental repeats consisting of 3 replicates each are shown, large open circles in graphs indicate median/mean of independent biological repeats. P-values obtained with Kruskall-Wallis ANOVA test above data points/bars. For raw data see Tab. T1-S1.

Fig. 1-S2. Correlation of different Eb1 comet properties. **A,B)** Eb1 amount at comets is calculated as the product of comet length (A) and the fluorescent mean intensity of Eb1 comets (B), which are both affected to similar degrees by homozygous condition of *msps^A*, *tau^{KO}* and *Eb1^{04524}* in embryo-derived neurons cultured for 12hrs following 6 day pre-culture (6d pre); data were normalised to controls (dashed horizontal line) and are shown as scatter dot plot with median ± 95% confidence interval of at least three experimental repeats, large open circles in graphs indicate median/mean of independent biological repeats. P-values listed above each plot were obtained with Kruskall-Wallis ANOVA tests. **C)** The table lists data for Eb1 amounts (fixed neurons; compare Fig.1E-H, L) or for comet velocity/lifetime (live imaging; compare Fig.1M,N), all obtained from pre-cultured embryonic primary neurons carrying the same combinations of mutant alleles in homozygosis (indicated on the left; used alleles: *tub84B^{def}*, *msps^A*, *tau$^{KO/Df}$*, *eb1^{04524}*, *stai^{KO}*). **D,E)** Plotting comet velocity or lifetime against Eb1 amounts from different genetic conditions shows good correlation (r and p-value determined via non-

parametric Spearman correlation analysis). For raw data see Tab. T1-S2.

Fig. 2. *Eb1, tau* and *msps* interact genetically. **A-B''**) Axon length, MT curling and Eb1 amount
(as indicated on the right), for primary neurons displaying heterozygous (A-A'', larval cultures)
and homozygous (B-B', embryonic 6d pre-cultures') mutant conditions, alone or in combination.
Data were normalised to parallel controls (dashed horizontal lines) and are shown as scatter
dot plots with median ± 95% confidence interval (A, A'',B, B'') or bar chart with mean ± SEM (A',
B') of at least two independent repeats with 3 replicates each; large open circles in graphs
indicate median/mean of independent biological repeats. P-values above data points/bars were
obtained with Kruskall-Wallis ANOVA tests; used alleles: $msps^A$, tau^{KO}, $Eb1^{04524}$. **C**) The graph
compares Eb1 amounts at MT plus-ends with MT disorganisation (MDI) for a range of genetic
conditions used in this work: green dots show data from pre-cultured embryonic neurons (B' vs.
B''), purple dots show comparable data obtained from larval primary neurons (A' vs. A''); in
addition, green/purple dots contain data from Figs. 1J and 1-S1B,D; black dots show similar
data obtained from primary *Xenopus* neurons (Fig.G,H); r and p-value determined via non-
parametric Spearman correlation analysis; see further detail of these correlations in Fig.2-S2.
For raw data see Tab. T2.

Fig. 2-S1. Comet numbers and dynamics in (trans-) heterozygous conditions **A**) Eb1 comet
numbers in fixed primary neurons cultured for 12hrs following 5 day pre-culture. **B**) Comet
velocity and lifetime obtained from live analyses of primary neurons cultured for 18hrs from
CNSs of late larvae carrying single-, double- or triple-heterozygous conditions (complementing
data in Fig. 2A). In all graphs, data were normalised to parallel controls (dashed horizontal lines)
and are shown as scatter dot plots with median ± 95% confidence interval (B) or bar chart with
mean ± SEM (A) of at least two experimental repeats; large open circles in graphs indicate
median/mean of independent biological repeats. P-values obtained with Kruskal-Wallis ANOVA
test are given above data points/bars; used alleles: $msps^A$, tau^{KO}, $Eb1^{04524}$. For raw data see
Tab. T2-S1.

Fig. 2-S2. Increased Eb1 amounts correlate with decreased MT curling. The data and graph
show further details behind the correlations displayed in Fig.2C. **A**) The table provides
descriptions, data and references for the graph shown in B: coloured numbers (1st column)
correspond to numbers of data points in B; the different allelic combinations and culture
conditions used (2nd column) comprise embryonic neurons cultures for 6 hrs (6HIV), embryonic
neurons cultured for 12 hrs following 6 day preculture (6d pre), neurons cultured from larval
CNSs for 18hrs (L3); respective data for Eb1 amounts (3rd column) and MT curling (4th column)
were obtained from different sets of experiments throughout this work (5th column lists the
figures from where these data originate). **B**) Correlation plot of the data shown in A, with
numbers and colours of data points corresponding to the 1st column; r and p-value determined
via non-parametric Spearman correlation analysis. For raw data see Tab. T2-S2.

Fig. 3. Eb1, Msps and Tau functionally interact in the fly brain and in frog primary neurons. **A,B**)
Medulla region of adult brains at 26-27 days after eclosure, all carrying the *GMR31F10-Gal4*
driver and *UAS-GFP-α-tubulin84B* (*GMR-tub*) which together stain MTs in a subset of lamina

neuron axons that terminate in the medulla; the further genetic background is either wild-type
(A) or triple-heterozygous (*Eb1$^{04524/+}$msps$^{A/+}$ tau$^{KO/KO}$*; B); white/black arrows indicate axonal
swellings without/with MT curling; rectangles outlined by red dashed lines are shown as 2.5 fold
magnified insets where white arrow heads point at disorganised curling MTs. **C,D**) Quantitative
analyses of specimens shown in A and B: relative number of total swellings per axon (C) and
of swellings with MT curling per axon (D); bars show mean ± SEM; P values from Kruskal–
Wallis one-way tests are given above each column, merged sample numbers (i.e. individual
axon bundles) from at least two experimental repeats at the bottom of each bar. **E-E''''**) Primary
Xenopus neurons stained for tubulin (tub): asterisks indicate cell bodies, white arrows indicate
unbundled MTs, white arrowheads unbundled areas with MT curling. **F-F'''**) *Xenopus* neurons
labelled with MACF43::GFP (MACF43): black arrows point at comets (visible as black dots). In
E-F''', black-stippled squares in overview images are shown as 2.5 fold magnified close-ups
below; ↓ behind gene symbols indicates 50% knock-down thus approximating heterozygous
conditions. **G,H**) Quantification of specimens shown in E-F'''' with respect to MT curling (G) and
comet amount of MACF43::GFP (H); data were normalised to parallel controls (dashed
horizontal lines) and are shown as mean ± SEM (G) or median ± 95% confidence interval (H);
merged sample numbers from at least two experimental repeats are shown at the bottom, P-
values obtained with Kruskall-Wallis ANOVA tests above data points/bars. The scale bar in A
represents 15 µm in A,B and 20 µm in E-F''''. For raw data see Tab. T3.

1003

Fig. 3-S1. Support data for *Xenopus* experiments. **A-B'**) A RT-PCR DNA gel (A) and Western
blot (B) and their quantifications (A', B') show the degrees of EB3/Tau knock-downs upon
application of different morpholino concentrations (indicated on top in blots and at the bottom in
graphs); ODC1 and ß-actin are used as loading controls; data are normalised to no-morpholino
controls from two experimental repeats (dashed lines). 50% knock-down of XMAP215 was
achieved by injecting 6 ng of the validated XMAP215 MO as described previously (Lowery *et*
al., 2013; Slater *et al.*, 2019). **C-D**) Different properties of MACF43::GFP comets (as indicated
upon graphs) obtained from *Xenopus* primary neurons, either upon 50% ; data were normalised
to parallel controls (dashed horizontal lines) and are shown as median ± 95% confidence
interval; merged sample numbers from at least two experimental repeats are shown at the
bottom, P-values obtained with Kruskall-Wallis ANOVA and Dunn's posthoc tests above data
points. For raw data see Tab. T3-S1.

1016

Fig. 4. Eb1 and Msps depend on each other for MT plus-end localisation. **A-A''**) Primary
neurons at 6HIV co-expressing Eb1::mCherry (magenta, Eb1) and Msps^{FL}::GFP (green, Msps)
and imaged live; asterisks indicate somata, scale bar represents 10µm, dashed boxes indicate
the positions of the 3.5-fold magnified close-ups shown at the bottom with arrowheads pointing
at the position of Msps::GFP accumulation (same in C,C'). **B,B'**) Kymograph of live movies (as
in A-A'') with the dashed line on the left representing a straightened version of the dashed lines
shown in A' and A'' (i.e. the length of the axon; proximal at the top) and the x-axis indicating
time; arrowheads point at trajectories of Msps and Eb1 which are almost identical. **C**) Primary
neurons expressing Msps::GFP and imaged live, either displaying wild-type background (ctrl)
or being homozygous mutant for *Eb1^{04524}* (*Eb1$^{-/-}$*); white arrowheads point at Msps::GFP comets
which are much smaller in the mutant neurons. **D**) Schematic representations of Msps^{FL}::GFP
and Msps$^{\Delta CTD}$::GFP. **D',D''**) Graphs displaying axon length and MT curling (as indicated) for pre-
cultured embryonic primary neurons expressing GFP or Msps::GFP constructs via the *elav-*

Gal4 driver, either in wild-type or *msps*$^{A/1}$ mutant background; data were normalised to parallel
controls (dashed horizontal lines) and are shown as median ± 95% confidence interval (D') or
mean ± SEM (D'') from at least two experimental repeats; large open circles in graphs indicate
median/mean of independent biological repeats. P-values obtained with Kruskall-Wallis ANOVA
tests are shown above data points/bars. **E**) Model view of the results shown here and in Fig. 1;
note that yellow circles represent GTP-tubulin which mediates the binding of Eb1; for further
explanations see main text and Discussion. For raw data see Tab. T4.

1037

Fig. 4-S1. Loss of TACC or Sentin does not cause obvious axonal phenotypes. Axon length (**A**)
and MT curling (**B**) embryonic 6d pre-cultured neurons which were either wild-type (wt) or
homozygous mutant for *sentin* or *dTACC* (as indicated); data were normalised to parallel
controls (dashed horizontal lines) and are shown as scatter dot plots with median ± 95%
confidence interval (A) or mean ± SEM (B) from at least two experimental repeats; large open
circles in graphs indicate median/mean of independent biological repeats. P-values obtained
with Kruskal-Wallis ANOVA tests are shown above data points/bars. For raw data see Tab. T4-
S1.

1046

Fig. 5. Tau promotes Eb1 pools at MT plus-ends by outcompeting Eb1's association with the
MT lattice. **A-A''**) Example of an embryonic neuron at 6 HIV imaged live with curling MTs to
illustrate Tau binding (green) along the MT lattice, separated from Eb1 comets (magenta);
asterisks indicate the soma, the scale bar represents 10µm, dashed boxes indicate the positions
of the 4-fold magnified close-ups shown at the bottom, white arrowheads point at Eb1 comets
(same in B-D). **B-D**) Primary neurons at 6 HIV stained for Eb1 which are either wild-type (B) or
mutant for *tau*$^{KO/Df}$ (C) or expressing Eb1::GFP driven by *elav-Gal4* in *tau*$^{KO/Df}$ mutant
background (D); white arrowheads indicate Eb1 comets, red arrowheads Eb1 lattice
localisation. **E-I**) Different parameters (as indicated) of control (ctrl) or *tau*$^{KO/Df}$ (*tau*$^{-/-}$) mutant
neurons at 6 HIV without/with *elav-Gal4*-driven expression of Eb1::GFP or Shot-Ctail::GFP (as
indicated); data were normalised to parallel controls (dashed horizontal lines) and are shown as
scatter dot plots with mean ± SEM (I) or median ± 95% confidence interval (E-H) from at least
two independent repeats with 3 experimental replicates; large open circles in graphs indicate
median/mean of independent biological repeats. P-values obtained with Kruskal-Wallis ANOVA
and Dunn's posthoc test as shown above data points/bars. **J**) Model view of the results shown
here; note that yellow circles represent GTP-tubulin which provides higher affinity for Eb1
binding; for further explanations see main text and Discussion. For raw data see Tab. T5.

1064

Fig. 5-S1 In Tau-deficient axons, GTP-tubulin is reduced at MT plus-ends but unchanged at
lattices. **A-B''**) Fixed primary neurons stained for Eb1 (magenta) and GTP-tubulin (green; the
scale bar represents 10µm); asterisks indicate somata, dashed boxes the positions of the 4-fold
magnified close-ups shown at the bottom, arrowheads point at Eb1::GFP comets and GTP
caps. **C,D**) Graphs showing staining intensity of GTP-tubulin at MT plus-ends (C) and along the
MT lattice (D) of embryonic neuron at 6 HIV. **E**) Graphs showing staining intensity of Eb1 along
the MT lattice of neurons without/with *elav-Gal4*-driven expression of *Drosophila* Tau (*dtau*).
Overexpression of *dtau* leads to a reduction of Eb1 at the MT shaft; data were normalised to
parallel controls (dashed horizontal lines) and are shown as scatter dot plots with median ± 95%
confidence interval from at least two independent repeats with 3 experimental replicates; large

open circles in graphs indicate median/mean of independent biological repeats. P-values
obtained with Kruskall-Wallis ANOVA test above data points/bars. For raw data see Tab. T5-
S1.

1078

Fig. 5-S2 Details of Eb1-Msps cross-regulation. **A,B**) Eb1 lattice localisation is unaffected by
loss of Msps in primary neurons at 6HIV (A) or at ~12HIV following 6 day pre-culture (B). **C**)
Staining intensity of GTP-tubulin at MT plus-ends is reduced in *msps$^{A/A}$* and *Eb1$^{04524/04524}$* mutant
embryonic neurons at ~12 HIV following 6 day pre-culture. **D-F**) Expressing Eb1::GFP via *elav-*
Gal4 does not rescue comet velocities (D), comet lifetime (E) and MT curling (F) in primary
neurons of *msps$^{A/146}$* mutants (cultured12 HIV following 6 day pre-culture); data were normalised
to parallel controls (dashed horizontal lines) and are shown as scatter dot plots with mean ±
SEM (F) or median ± 95% confidence interval (A-E) from at least two experimental repeats;
large open circles in graphs indicate median/mean of independent biological repeats. P-values
obtained with Kruskal-Wallis ANOVA tests and Dunn's posthoc analyses are shown above data
points/bars. For raw data see Tab. T5-S2

1090

Fig. 6. Shot-mediated guidance mechanistically links Eb1 at MT plus-ends to bundle
organisation. **A**) Schematic representation of Shot constructs (CH, actin-binding calponin-
homology domains; EF, EF-hand motifs; GRD, MT-binding Gas2-related domain; Ctail,
unstructured MT-binding domain containing Eb1-binding SxIP motifs in blue); in Shot-
3MtLS*::GFP the SxIP motifs are mutated. **B**) Fixed primary neurons at 18HIV obtained from
late larval CNSs stained for GFP (green) and tubulin (magenta), which are either wild-type (top)
or *Eb1$^{04524/+}$msps$^{A/+}$ tau$^{KO/+}$* triple-heterozygous (indicated on right) and express GFP or either
of the constructs shown in D; scale bar 10µm. **C**) Quantification of MT curling of neurons as
shown in B. **D,E**) MT curling in *shot$^{3/3}$ Eb1$^{04524/04524}$* double-mutant neurons is not enhanced over
single mutant conditions assessed in fixed embryonic primary neurons at 6HIV (D) or 12HIV
following 6 day pre-culture (E). In all graphs data were normalised to parallel controls (dashed
horizontal lines) and are shown as mean ± SEM from at least two independent repeats with 3
experimental replicates each; large open circles in graphs indicate median/mean of independent
biological repeats. P-values obtained with Kruskall-Wallis ANOVA tests are shown above bars.
F,F') Model derived from previous work (Alves-Silva *et al.*, 2012), proposing that the
spectraplakin Shot cross-links Eb1 at MT plus-ends with cortical F-actin, thus guiding MT
extension in parallel to the axonal surface; yellow dots represent GTP-tubulins providing high
affinity sites for Eb1-binding.

1109

Fig. 7. A mechanistic model consistent with all reported data. **A**) In wild-type (WT) neurons, the
three factors bind (dark grey arrows) to MTs at different locations: Msps binds to the very tip of
MT plus-ends, Eb1 forms plus-end comets (curly bracket) by binding with higher affinity to GTP-
tubulin (GTP-cap, light beige) but lagging behind the front, and Tau localises along the MT
lattice primarily composed of GDP-tubulin (green). Tau outcompetes (red T-bar) low-affinity
binding of Eb1 along the lattice, thus maintaining more Eb1 at MT plus-ends. Msps enhances
(orange arrow) MT polymerisation (brown), thus sustaining a prominent GTP-cap that Eb1 can
bind to. Eb1 stabilises MT plus-ends (green arrow) by promoting sheet formation of
protofilaments at the plus-end (short Y-shaped tip), thus improving conditions for Msps binding.
Eb1 also recruits the C-terminus of Shot which binds cortical actin via its N-terminus, thus

establishing cross-linkage that can guide (blue arrow) extending MT plus-ends into parallel
bundles. **B-E**) Illustrations explaining the changes triggered by the loss of the different factors
(stippled X); any affected processes are shown as stippled lines with reduced thickness and
reduced font sizes of accompanying texts. **B**) Upon loss of Eb1, the plus-end sheet structure is
weakened (larger Y shape), thus reducing Msps binding and, in turn, reducing polymerisation;
Shot detaches, thus abolishing guidance. **C**) Loss of Msps abolishes enhanced polymerisation
so that the GTP-cap shrinks and less Eb1 binds, thus also weakening Shot binding and MT
guidance. **D**) Upon loss of Tau, more Eb1 can bind to the MT lattice, thus reducing the amounts
available for plus-end association; this causes a modest Eb1 depletion phenotypes with
consequences similar to B, but less pronounced. **E**) Upon loss of Shot, the localisations and
functions of the other three proteins are unaffected, but MT guidance is abolished.

1131

1132 **Movie M1**. Msps::GFP and Eb1::RFP jointly track MT plus-ends. Live movie of a wild-type
1133 neuron co-expressing Msps::GFP and Eb1::RFP; for stills see Fig.4A-A". As indicated, single
1134 channels are shown on the left and middle, and the combined movie on the right. The movie
1135 was acquired at 0.5 frames per second and plays at 0.5 s per frame. The scale bar indicates 10
1136 µm.

1137

1138 **Movie M2**. Msps plus-end localisation in wild-type neurons. Live movie of a wild-type neuron
1139 expressing Msps::GFP; for a still see Fig.4C. The movie was acquired at 1 frame per second
1140 and plays at 0.2 s per frame. The scale bar indicates 10 µm.

1141

1142 **Movie M3**. Msps plus-end localisation is impaired in the absence of Eb1. Live movie of an
1143 *Eb1$^{04524/04524}$* mutant neuron expressing Msps::GFP; for a still see Fig.4C'. The movie was
1144 acquired at 1 frames per second and plays at 0.2 s per frame. The scale bar indicates 10 µm.

Hahn et al. Fig. 1

Hahn et al. Fig. 1-S1

C

	Eb1 amount	comet velocity	comet lifetime
ctrl	1.0000	1.0000	1.0000
tau+/-	1.1130	1.1010	1.0360
msps+/-	1.0430	0.9792	0.9411
eb1+/-	0.9028	1.1430	1.0360
eb1+/- msps+/-	0.7093	0.8496	0.9097
eb1+/- msps+/- tau+/-	0.3989	0.7602	0.7879
msps-/-	0.3796	0.4676	0.6905
tau-/-	0.8918	0.8234	0.7246
tubdef/def	1.1020	1.0720	1.0160
stai-/-	1.0450	0.9107	1.0490

pre-cultures

Hahn et al. Fig. 1-S2

Hahn et al. Fig. 2

A comet numbers (embryonic 6D pre)

B comet dynamics (larval)

comet velocity

comet lifetime

genetic interactions

Hahn et al. Fig. 2-S1

A

		comet Eb1 amount	MDI	Figure
1	6d pre msps$^{1/1}$	0.3796	2.338	Fig. 1
2	6d pre tau$^{KO/KO}$	0.8918	1.388	Fig. 1
3	6d pre Eb1$^{04524/04524}$	0.282	2.191	Fig. 1
4	6d pre msps$^{1/1}$ tau$^{KO/KO}$	0.4592	1.786	Fig. 2
5	6d pre Eb1$^{04524/04524}$ tau$^{KO/KO}$	0.3298	2.063	Fig. 2
6	6d pre Eb1$^{04524/04524}$ msps$^{1/1}$	0.3421	2.109	Fig. 2
7	6d pre Eb1$^{04524/04524}$ msps$^{1/1}$ tau$^{KO/KO}$	0.2077	4.363	Fig. 2
8	7d pre Eb1^{Def}	0	5.814	Fig. 2
9	6d pre tub^{def}	1.102	0.6603	Fig. 1-S1
10	6d pre clasp$^{2/Df}$	1.134	0.4888	Fig. 1-S1
11	6d pre MAP1B^{-}	1.162	0.8491	Fig. 1-S1
12	6HIV Eb1^{04524}	0.3832	2.697	Fig. 1-S1
13	6HIV msps^{A}	0.4884	1.622	Fig. 1-S1
14	6HIV tau^{KO}	0.5217	1.815	Fig. 1-S1
15	6HIV tub^{Df}	1.09	0.43	Fig. 1-S1
16	6HIV stai^{KO1}	1.04	0.69	Fig. 1-S1
17	L3 msps$^{1/+}$	1.043	1.176	Fig. 2
18	L3 tau$^{KO/+}$	1.114	1.11	Fig. 2
19	L3 Eb1$^{04524/+}$	0.9031	1.397	Fig. 2
20	L3 msps$^{1/+}$ tau$^{KO/+}$	0.7372	1.932	Fig. 2
21	L3 tau$^{KO/+}$ Eb1$^{04524/+}$	0.8818	1.243	Fig. 2
22	L3 msps$^{1/+}$ Eb1$^{04524/+}$	0.7095	2.609	Fig. 2
23	L3 msps$^{1/+}$ tau$^{KO/+}$ Eb1$^{04524/+}$	0.399	6.655	Fig. 2
24	L3 elavG4; tau$^{KO/def}$	0.7631	2.758	Fig. 5
25	L3 elav>EB1-GFP, tauKO/def	1.067	1.34	Fig. 5
26	L3 stai$^{Df/KO3}$	1.045	1.197	Fig. 1-S1
27	Xen 50% Tau KD	0.9358	1.411	Fig. 3
28	Xen 50% XMAP215 KD	0.8873	1.892	Fig. 3
29	Xen 50% EB3 KD	0.9271	2.002	Fig. 3
30	Xen 50% XMAP215/Tau/EB3 KD	0.5739	3.822	Fig. 3

B

Hahn et al. Fig. 2-S2

A *GMR-tub*

B *GMR-tub Eb1⁺/⁻ , msps⁺/⁻ , tau⁺/⁻*

C # of swellings/ axon bundle ↑

swelling/axon bundle

p <0.0001

ctrl   *Eb1⁺/⁻ msps⁺/⁻ tau⁺/⁻*

381   285

D # of disor. MT areas/ axon bundle ◁

disorg. MT areas/ axon bundle

<0.0001

ctrl   *Eb1⁺/⁻ msps⁺/⁻ tau⁺/⁻*

382   289

E E′ E″ E‴ E‴′   tub

F F′ F″ F‴ F‴′   MACF43

control *Tau↓* *XMAP↓* *EB3↓* *Tau↓ XMAP↓ EB3↓*   ↓50% KD

G MT disorganisation

MDI

0.0002

0.99   0.9   0.08   <0.001

Control Tau↓ XMAP215↓ EB3↓ EB3↓XMAP215↓Tau↓

64   16   24   53   54

H comet MACF43 amount

comet MACF43 amounts

<0.001

0.99   0.09   0.99   <0.001

Control Tau↓ XMAP215↓ EB3↓ EB3↓XMAP215↓Tau↓

95   40   40   47   41

Hahn et al. Fig. 3

Hahn et al. Fig. 3-S1

Hahn et al. Fig. 4

A   axon length

B   MT disorganisation

embryo pre-cultures

Hahn et al. Fig. 4-S1

A Tau Eb1 **A'** Tau **A"** Eb1

B Eb1 **C** **D**

control | $tau^{-/-}$ | $tau^{-/-}$ Eb1GFP

E Eb1 intensity along MT lattice

rel. Eb1 intensity

<0.0001   <0.0001

ctrl $tau^{-/-}$ ctrl $tau^{-/-}$

Eb1-GFP

embryo cultures

F Comet Eb1 amount

rel. Eb1 amount

<0.0001   0.75

<0.0001

ctrl $tau^{-/-}$ $tau^{-/-}$

Eb1-GFP

G Comet velocity

comet velocity [μm/s]

0.33

0.91   <0.0001

ctrl $tau^{-/-}$ ctrl $tau^{-/-}$

Eb1-GFP   *Ctail-GFP*

H Comet lifetime

comet lifetime [s]

0.71

0.67   <0.0001

ctrl $tau^{-/-}$ ctrl $tau^{-/-}$

Eb1-GFP   *Ctail-GFP*

I MT disorganisation

MDI

<0.0001   0.35

<0.0001

ctrl $tau^{-/-}$ $tau^{-/-}$

Eb1-GFP

J

Tau   Eb1

wt   MT polym.

$tau^{-/-}$

$tau^{-/-}$ EB1-GFP

Hahn et al. Fig. 5

A ctrl **A′** **A″**

Eb1 GTPtub   GTPtub   Eb1

B *tau* **B′** **B″**

Eb1 GTPtub   GTPtub   Eb1

C GTP cap amount (embryo)

rel. GTPtub amount

<0.0001

wt   tau⁻/⁻

D GTP-tub intensity along MT lattice (embryo)

rel. GTPtub intensity

0.007

wt   tau⁻/⁻

E Eb1 intensity along MT lattice (embryo)

rel. Eb1 intensity

<0.0001

ctrl   elav>dtau

Hahn et al. Fig. 5-S1

Hahn et al. Fig. 5-S2

Hahn et al. Fig. 6

Hahn et al. Fig.7