Tau, XMAP215/Msps and Eb1 jointly regulate microtubule polymerisation and bundle
formation in axons

Ines Hahn[#, 1], Andre Voelzmann[1], Jill Parkin[1], Judith Fuelle[1], Paula G Slater[2], Laura A Lowery[3],
Natalia Sanchez-Soriano[#‡4] & Andreas Prokop[#‡1]

1) The University of Manchester, Manchester Academic Health Science Centre, Faculty of
Biology, Medicine and Health, School of Biology, Manchester, UK

2) Department of Biology, Boston College, Chestnut Hill, MA, USA

3) Department of Medicine, Boston University Medical Center, Boston, MA, USA

4) Department of Molecular Physiology & Cell Signalling, Institute of Systems, Molecular &
Integrative Biology, University of Liverpool, Liverpool, United Kingdom

Running title**:** Tau, XMAP215 and Eb1, a functional trio for microtubule polymerisation and
organisation

Key words: *Drosophila*, neurodegeneration, axons, cytoskeleton, microtubules

authors for correspondence:

The University of Manchester

Faculty of Life Sciences

Oxford Road

Manchester M13 9PT

Tel: +44-(0)161-27-51556

Fax: +44-(0)161-27-51505

Ines.Hahn@manchester.ac.uk

N.Sanchez-Soriano@liverpool.ac.uk

Andreas.Prokop@manchester.ac.uk

‡ joined senior authors

Summary statement:

Eb1, XMAP215 and tau operate as a functional unit in axons to promote the polymerisation of
microtubules and their organisation into the parallel bundles essential for axonal transport.

Abstract

Axons are the enormously long cable-like cellular processes of neurons that wire nervous
systems and have to survive for up to a century in humans. We lose ~40% of axons towards
high age and far more in neurodegenerative diseases. Sustaining axons requires axonal
transport and dynamic morphogenetic changes, both crucially dependent on bundles of
microtubules that run all along axons. How polymerisation is regulated to form, repair and

replace microtubules in these bundles during axon development and maintenance is virtually
unknown. Here, we show in axons of *Drosophila* and *Xenopus* neurons alike that Eb1,
XMAP215/Msps and Tau are key players which operate as one functional unit to promote
microtubule polymerisation. Eb1 and XMAP215/Msps are interdependent core factors at the
microtubule tip, whereas Tau outcompetes Eb1 binding at microtubule lattices, thus preventing
its pool depletion at polymerising plus ends. In agreement with their closely interwoven
functions, the three factors show the same combination of axonal loss-of-function mutant
phenotypes including: (1) reduced microtubule polymerisation dynamics and shorter axon
growth, indicating their importance for upholding microtubule mass in axons; (2) prominent
deterioration of parallel microtubule bundles into disorganised curled conformations, indicating
their key roles in promoting essential axonal architecture. We show the latter to occur through
Eb1- and spectraplakin-dependent guidance of extending microtubules. We conclude that Eb1,
XMAP215/Msps and Tau jointly promote microtubule polymerisation, important to regulate the
quantity and bundled organisation of microtubules and offering new ways to think about
developmental and degenerative axon pathologies and how to treat them.

Introduction

Axons are the enormously long cable-like cellular processes of neurons that wire nervous
systems. In humans, axons of ≤15μm diameter can be up to a meter long (Prokop, 2020). They
are constantly exposed to mechanical challenges, yet have to survive for up to a century; we
lose ~40% of axons towards high age and far more in neurodegenerative diseases (Adalbert &
Coleman, 2012, Calkins, 2013, Marner, Nyengaard et al., 2003).

Their growth and maintenance require parallel bundles of microtubules (MTs) that run all along
axons, providing the highways for life-sustaining transport and driving morphogenetic events.
Consequently, bundle decay through MT mass decrease and/or MT disorganisation is a
common feature in axon pathologies (summarised in (Hahn, Voelzmann et al., 2019). Key roles
must be played by MT polymerisation, which is not only essential for the *de novo* formation of
MT bundles during axon growth occuring in development, plasticity or regeneration, but also to
repair and replace senescent MTs during long-term maintenance (Gasic & Mitchison, 2019,
Schaedel, John et al., 2015, Voelzmann, Hahn et al., 2016). However, the molecular
mechanisms driving MT polymerisation in axons are surprisingly little understood.

In vitro, MTs undergo polymerisation in the absence of enzymatic catalysis, but the addition of
factors such as CLASPs, stathmins, tau, Eb proteins or XMAP215 can enhance and refine the
process (Aher, Rai et al., 2020, Al-Bassam, Kim et al., 2010, Brouhard, Stear et al., 2008,
Drechsel, Hyman et al., 1992, Li, Moriwaki et al., 2012, Manna, Thrower et al., 2009, Zanic,
Widlund et al., 2013). Accordingly, many candidate regulators have been proposed in the
literature to regulate MT polymerisation in axons, comprising factors that are MT plus end-
associating, MT shaft-binding or involved in tubulin provision (summarised in (Voelzmann et al.,
2016). We would therefore expect that MT polymerisation in axons is regulated through
functional networks of proteins, but we are far from understanding what the key players are and
how they are regulated by other proteins in these networks.

To gain such knowledge, we have chosen *Drosophila* primary neurons as one consistent model
amenable to combinatorial genetics as a powerful strategy to decipher complex regulatory
networks (Prokop, Beaven et al., 2013). Our previous loss-of-function studies of 9 MT plus end-
associating factors in these *Drosophila* neurons (CLASP, CLIP190, dynein heavy chain, APC,
p150[Glued], Eb1, Short stop/Shot, doublecortin, Lis1) have taken axon length as a crude proxy

readout for net polymerisation, mostly revealing relatively mild axon length phenotypes, with the
exception of Eb1 and Shot which cause severe axon shortening {Beaven, 2015 #6487;
Sánchez-Soriano, 2009 #7139; Alves-Silva, 2012 #4930; A.V., unpublished data}.

Here we have taken these analyses to the next level. We show that three factors, Eb1,
XMAP215/Msps and Tau, share a unique combination of mutant phenotypes, including reduced
axonal MT polymerisation in frog and fly neurons in culture and *in vivo*. Our data reveal that the
three factors operate as a functional unit: whereas Eb1 and XMAP215/Msps act
interdependently at MT plus ends, Tau outcompetes Eb1 binding at MT lattices, thus preventing
its pool depletion at polymerising MT plus ends. By upholding MT polymerisation, this functional
trio also promotes the bundle conformation of axonal MTs through a guidance mechanism
mediated by the spectraplakin Shot.

Methods

Fly stocks

Loss-of-function mutant stocks used in this study were the deficiencies *Df(3R)Antp17* (*tub^{def}*;
removing both *αtub84B* and *αtub84D*; Duncan & Kaufman, 1975, Jenkins, Saunders et al.,
2017), *Df(2L)Exel6015* (*stai^{Df}*; Duncan, Lytle et al., 2013), *Df(3L)BSC553* (*clasp^{Df}*; Bloomington
stock #25116; Beaven, Dzhindzhev et al., 2015), *Df(3R)tauMR22* (*tau^{Df}*; Doerflinger, Benton et
al., 2003) and the loss-of-function mutant alleles *α-tub84B^{KO}* (an engineered null-allele; Jenkins
et al., 2017), *chromosome bows^2* (*clasp^2*, an amorph allele; Inoue, do Carmo Avides et al.,
2000), *Eb1^{04524}*, *Eb1^5* (strong loss-of-function mutant alleles; Elliott, Cullen et al., 2005),
futsch^{P158} (*MAP1B*; a deficiency uncovering the *futsch* locus; Hummel, Krukkert et al., 2000),
msps^A (a small deletion causing a premature stop after 370 amino acids; gift from H. Ohkura,
unpublished), *msps^{146}* (Brittle & Ohkura, 2005), *sentin^{ΔB}* (*short spindle2^{ΔB}*, *ssp2^{ΔB}*; Gluszek,
Cullen et al., 2015), *tacc^1* (*dTACC^1*; Gergely, Kidd et al., 2000), *shot^3* (the strongest available
allele of *short stop*; Kolodziej, Jan et al., 1995, Sánchez-Soriano, Travis et al., 2009), *stai^{KO}*
(Yang, Inaki et al., 2012), *tau^{KO}* (a null allele; Burnouf, Gronke et al., 2016). Gal4 driver lines
used were *elav-Gal4* (1st and 3rd chromosomal, both expressing pan-neuronally at all stages;
(Luo, Liao et al., 1994), *GMR31F10-Gal4* (Bloomington #49685; expressing in T1 medulla
neurons; (Qu, Hahn et al., 2019). Lines for targeted gene expression were *UAS-Eb1-GFP* and
UAS-shot-Ctail-GFP (Alves-Silva, Sánchez-Soriano et al., 2012), *UAS-shot^{ΔABD}-GFP* (Qu,
2015), *UAS-shot^{3MTLS}-GFP* (Alves-Silva et al., 2012), *UAS-dtau-GFP* (Doerflinger et al., 2003),
UAS-GFP-α-tubulin84B (Grieder, de Cuevas et al., 2000) and further lines generated here (see
below).

Drosophila primary cell culture

Drosophila primary neuron cultures were done as described previously (Prokop, Küppers-
Munther et al., 2012, Qu, Hahn et al., 2017). Stage 11 embryos were treated for 90 s with bleach
to remove the chorion, sterilized for ~30 s in 70% ethanol, washed in sterile Schneider's medium
containing 20% fetal calf serum (Schneider's/FCS; Gibco), and eventually homogenized with
micro-pestles in 1.5 centrifuge tubes containing 21 embryos per 100 µl dispersion medium
(Prokop et al., 2012) and left to incubated for 4 min at 37°C. Dispersion was stopped with 200
µl Schneider's/FCS, cells were spun down for 4 mins at 650 g, supernatant was removed and
cells re-suspended in 90 µl of Schneider's/FCS, and 30 µl drops were placed in culture
chambers and covered with cover slips. Cells were allowed to adhere to cover slips for 90-120

min either directly on glass or on cover slips coated with a 5 µg/ml solution of concanavalin A,
and then grown as a hanging drop culture at 26°C as indicated.

To eliminate a potential maternal rescue of mutants (i.e. reduction of the mutant phenotype due
to normal gene product deposition from the wildtype gene copy of the heterozygous mothers in
oocytes (Prokop, 2013), we used a pre-culture strategy (Prokop et al., 2012, Sánchez-Soriano,
Gonçalves-Pimentel et al., 2010) where cells were incubated in a tube for 7 days before they
were plated on coverslips.

For cultures from larval brains, L3 brains (2-3 per cover slip) were dissected in PBS, transferred
into Schneider's/FCS medium, washed three times with medium and then proceed with
homogenisation and dispersion as explained above

Transfection of *Drosophila* primary neurons was executed as described previously (Qu et al.,
2019). In brief, 70-75 embryos per 100 µl dispersion medium were used. After the washing step
and centrifugation, cells were re-suspended in 100 µl transfection medium [final media
containing 0.1-0.5 µg DNA and 2 µl Lipofecatmine 2000 (L2000, Invitrogen), incubation
following manufacturer's protocols (Thermo Fisher, Invitrogen) and kept for 24 hrs at 26°C. Cells
were then treated again with dispersion medium, re-suspended in culture medium and plated
out as described above.

<u>*Xenopus* primary neuron experiments</u>

Xenopus primary neuron cultures were obtained from embryonic neural tube. Eggs collected
from female X. laevis frogs were fertilised in vitro, dejellied and cultured following standard
methods (Sive, Grainger et al., 2010). Embryos were grown to stage 22–24 (Nieuwkoop & J.,
1994), and neural tubes were dissected as described (Lowery, Faris et al., 2012). Three neural
tubes were transferred to an Eppendorf tube containing 150 uL CMF-MMR (0.1 M NaCl, 2.0
mM KCl, 1.0 mM EDTA, 5.0 mM HEPES, pH 7.4), 10 min later centrifuged at 1000 g for 5 min,
and 150 uL of Steinberg's solution (58 mM NaCl, 0.67 mM KCl, 0.44 mM $Ca(NO_3)_2$, 1.3 mM
$MgSO_4$, 4.6 mM Tris, pH 7.8) was added to the supernatant to follow with the tissue dissociation
using a fire polished glass Pasteur pipet. Cells were seeded in 100 ug/mL Poly-L-lysine and 10
µg/ml laminin pre-treated 60mm plates, and after 2 hr the media was replaced by plating culture
media (50% Ringer's, 49% L-15 media, 1% Fetal Bovine Serum, 25 ng/µl NT3 and BDNF, plus
50 µg/ml penicillin/streptomycin and gentamycin, pH 7.4 and filter sterilized) and kept for 24 hr
before imaging.

All experiments were approved by the Boston College Institutional Animal Care and Use
Committee and performed according to national regulatory standards.
The embryos were injected four times in dorsal blastomeres at two-to-four cell stage with 6 ng
of the validated XMAP215 morpholino (MO; Lowery, Stout et al., 2013), 10 ng of the validated
Tau MO (Liu, Wang et al., 2015), and/or 5 ng of a newly designed splice site MO for EB3
(3´CTCCCAATTGTCACCTACTTTGTCG5´; for verification see Fig3-S1), in order to obtain a
50% knockdown of each.

To assess EB1 comet dynamics and comet amounts, 300 pg of MACF43-Ctail::GFP, an Eb
protein-binding 43-residue fragment derived from the C-terminal regions of hMACF2 (human
microtubule actin crosslinking factor 2; Fig.3F-F'''; Honnappa, Gouveia et al., 2009, Slater,
Cammarata et al., 2019), was co-injected with the MO.

<u>Immunohistochemistry</u>

Primary fly neurons were fixed in 4% paraformaldehyde (PFA; in 0.05 M phosphate buffer, pH
7–7.2) for 30 min at room temperature (RT). For anti-Eb1 and anti-GTP-tubulin staining, cells
were fixed for 10 mins at -20°C in +TIP fix (90% methanol, 3% formaldehyde, 5 mM sodium
carbonate, pH 9; stored at -80°C and added to the cells) (Rogers, Rogers et al., 2002), then
washed in PBT (PBS with 0.3% TritonX). Antibody staining and washes were performed with
PBT. Staining reagents: anti-tubulin (clone DM1A, mouse, 1:1000, Sigma; alternatively, clone
YL1/2, rat, 1:500, Millipore Bioscience Research Reagents); anti-DmEb1 (gift from H. Ohkura;
rabbit, 1:2000; Elliott et al., 2005); anti-GTP-tubulin (hMB11; human, 1:200; AdipoGen; Dimitrov,
Quesnoit et al., 2008); anti-Shot (1:200, guinea pig; Strumpf & Volk, 1998); anti-Elav (Elav-
7E8A10; rat, 1:1000; Developmental Studies Hybridoma Bank, The University of Iowa, IA, USA;
O'Neill, Rebay et al., 1994); anti-GFP (ab290, Abcam, 1:500); Cy3-conjugated anti-HRP (goat,
1:100, Jackson ImmunoResearch); F-actin was stained with phalloidin conjugated with
TRITC/Alexa647, FITC or Atto647N (1:100 or 1:500; Invitrogen and Sigma). Specimens were
embedded in ProLong Gold Antifade Mountant (ThermoFisher Scientific).

Western blot analysis of *Xenopus* embryos

For protein extraction, 10 embryos were transferred to a centrifuge tube with 800 µl lysis buffer
(50 mM Tris pH 7.5, 5% glycerol, 0.2% IGEPAL/NP-40, 1 mM EDTA, 1.5 mM $MgCl_2$, 125 mM
NaCl, 25 mM NaF, 1 mM Na_3VO_4), homogenised with a sterile pestle and, after 10 min,
centrifuged at 13000 rpm for 15-20 min. The supernant was collected and the protein
concentration determined with the Micro BCA^{TM} Protein Assay Kit (Thermo Fisher Scientific).
80 µg protein were loaded into a 10 % SDS gel and stained with anti-Tau (clone Tau46, T9450,
mouse, 1:1000, Sigma-Aldrich).

Dissection of adult heads

198   For *in vivo* studies, brain dissections were performed in Dulbecco's PBS (Sigma, RNBF2227)
199   after briefly sedating them on ice. Dissected brains with their laminas and eyes attached were
200   placed into a drop of Dulbecco's PBS on MatTek glass bottom dishes (P35G1.5-14C), covered
201   by coverslips and immediately imaged with a 3i Marianas Spinning Disk Confocal Microscope.

Microscopy and data analysis

Standard imaging was performed with AxioCam 506 monochrome (Carl Zeiss Ltd.) or
MatrixVision mvBlueFox3-M2 2124G digital cameras mounted on BX50WI or BX51 Olympus
compound fluorescent microscopes. For the analysis of *Drosophila* and *Xenopus* primary
neurons, we used the following parameters:

Axon length was measured from cell body to growth cone tip using the segmented line tool of
ImageJ (Alves-Silva et al., 2012, Sánchez-Soriano et al., 2010).

Degree of MT disorganisation in axons was measured as "MT disorganisation index" (MDI)
described previously (Qu et al., 2019, Qu et al., 2017); in short: the area of disorganisation was
measured with the freehand selection in ImageJ; this value was then divided by axon length
(see above) multiplied by 0.5 µm (typical axon diameter, thus approximating the expected area
of the axon if it were not disorganised).

Eb1 comet amounts were approximated by using product of comet mean intensity and length.
For this, a line was drawn through each comet (using the segmented line tool in FIJI) and length

as well as mean staining intensity (of Eb1 or GTP-tub in fixed *Drosophila* and MACF43::GFP in
a movie still in *Xenopus* neurons) was determined.

To measure MT disorganisation in the optic lobe of adult flies, *GMR31F10-Gal4* (Bloomington
#49685) was used to express *UAS-α-tubulin84B-GFP* (Grieder et al., 2000) in a subset of
lamina axons which projects within well-ordered medulla columns (Prokop & Meinertzhagen,
2006). Flies were left to age for 26-27 days (about half their life expectancy) and then their
brains were dissected out, mounted in Mattek dishes and imaged using a 3i spinning disk
confocal system at the ITM Biomedecial imaging facility at the University of Liverpool. A section
of the medulla columns comprising the 4 most proximal axonal terminals was used to quantify
the number of swellings and regions with disorganised MTs.

To measure MT polymerisation dynamics, movies were collected on an Andor Dragonfly200
spinning disk upright confocal microscope (with a Leica DM6 FS microscope frame) and using
a *100x/1.40 UPlan SAPO (Oil)* objective. Samples where excited using 488nm (100%) and
561nm (100%) diode lasers via Leica GFP and RFP filters respectively. Images where collected
using a Zyla 4.2 Plus sCMOS camera with a camera gain of 1x. The incubation temperature
was set to 26°C. Time lapse movies were constructed from images taken every 1 s for 1 mins.
To measure comet velocity and lifetime, a line was drawn that followed the axon using the
segmented line tool in ImageJ. A kymograph was then constructed from average intensity in
FIJI using the KymoResliceWide macro (Cell Biology group, Utrecht University) and events
scored via the Velocity Measurement Tool Macro (Volker Baecker, INSERM, Montpellier, RIO
Imaging; J. Rietdorf, FMI Basel; A. Seitz, EMBL Heidelberg). For each condition at least 15 cells
were analysed in ≥2 independent repeats.

Time lapse imaging for *Xenopus* primary cultures was performed with a CSU-X1M 5000
spinning-disk confocal (Yokogawa, Tokyo, Japan) on a Zeiss Axio Observer inverted motorized
microscope with a Zeiss 63× Plan Apo 1.4 numerical aperture lens (Zeiss, Thornwood, NY).
Images were acquired with an ORCA R2 charge-coupled device camera (Hamamatsu,
Hamamatsu, Japan) controlled with Zen software. Time lapse movies were constructed from
images taken every 2 s for 1 min. The MACF43 comets' velocities and lifetime were analysed
with plusTipTracker software. The same parameters were used for all movies: maximum gap
length, eight frames; minimum track length, three frames; search radius range, 5–12 pixels;
maximum forward angle, 50°; maximum backward angle, 10°; maximum shrinkage factor, 0.8;
fluctuation radius, 2.5 pixels; and time interval 2 s.

Images were derived from at least 2 independent experimental repeats performed on different
days, for each of which at least 3 independent culture wells were analysed by taking a minimum
of 20 images per slide. For statistical analyses, Kruskal–Wallis one-way ANOVA with *post hoc*
Dunn's test or Mann–Whitney Rank Sum Tests (indicated as P_{MW}) were used to compare
groups, r and p-value for correlation were determined via non-parametric Spearman correlation
analysis. All raw data of our analyses are provided as supplementary Excel files T1-6.

<u>Molecular biology</u>

To generate the *UAS-msps^{FL}-GFP* (aa1-2050) and *UAS-msps$^{\Delta Cterm}$* (aa1-1322) constructs,
eGFP was PCR-amplified from pcDNA3-EGFP and *msps* sequences from cDNA clone
LP04448 (DGRC; FBcl0189229) using the following primers:

msps^{FL}   and   *msps$^{\Delta Cterm}$*   fw:
GAATAGGGAATTGGGAATTCGTTAGGCGCGCCAACATGGCCGAGGACACAGAGTAC

msps^{FL}   rev:
CAAGAAAGAGAATCATGCCCAAGGGCCCGGTAGCGGCAGCGGTAGCGTGAGCAAGGGC
GAG

msps^{ΔCterm}   rev:
GATGGAGGGTCTAAAATCGCATATGGGTAGCGGCAGCGGTAGCGTGAGCAAGGGCGAG
GAG

eGFP   fw:
GAGAATCATGCCCAAGGGCCCGGTAGCGGCAGCGGTAGCGTGAGCAAGGGCGAGGAG
CTG

eGFP   rev:
CTCTCGGCATGGACGAGCTGTACAAGTAGGCGGCCGCCTCGAGGGTACCTCTAGAG

The *msps* and *eGFP* sequences were introduced into *pUAST-attB* via Gibson assembly
(ThermoFisher) using *EcoRI* and *XhoI*. To generate transgenic fly lines, *pUAST-attB* constructs
were integrated into *PBac{yellow[+]-attP-9A}VK00024* (Bloomington line #9742) via PhiC31-
mediated recombination (outsourced to Bestgene Inc).

Results

<u>Eb1, Msps/XMAP215 and Tau share the same combination of loss-of-function phenotypes in</u>
<u>axons</u>

Many candidate regulators have been proposed to regulate MT polymerisation in axons,
comprising factors that are MT plus end-associating, MT shaft-binding or involved in tubulin
provision (Voelzmann et al., 2016). To reveal the core regulatory machinery necessary for
axonal MT polymerisation, we performed a detailed loss-of-function study with a set of
candidate factors using MT polymerisation dynamics and axon length as readouts. Of the pool
of MT plus end-associating factors, we included Eb1, Shot, CLASP/Chb (Chromosome bows)
and the XMAP215 homologue Msps; as MT shaft-binding candidates, we chose Tau and
Map1b/Futsch; to explore the impact of tubulin availability on polymerisation, we used α1-
tubulin/αtub84B, the predominant α-tubulin expressed in the fly nervous system (FlyAtlas 2,
University of Glasgow, UK; A.V., unpublished) and the tubulin regulator Stathmin (see
Discussion for more details on these factors).

To study these candidates, we analysed embryo-derived primary neurons carrying loss-of-
function mutations for the respective genes. To exclude that phenotypes are masked by
maternal contribution (i.e. healthy gene product deposited in the eggs by the heterozygous
mothers), we used additional strategies (Prokop et al., 2012, Sánchez-Soriano et al., 2010):
either we analysed late larval brain-derived primary neurons at 18 HIV (from now on referred to
as larval neurons); if mutants did not reach larval stages, we used embryo-derived neurons that
were kept for 5-7 days in pre-culture to deplete maternal product and then plated and grown for
12 hrs (pre-cultured neurons). In all cases, primary neurons were immuno-stained either for
endogenous tubulin to assess axon length, or for endogenous Eb1 protein to gain a first insight
into the polymerisation state of axonal MTs.

Eb1 staining revealed that loss of all factors, except Shot and Futsch, displayed a significant
reduction in the number of plus end comets (Figs. 1A-D, I and 1-S1A). In addition, we measured
the mean intensities and mean lengths of Eb1 comets and used the product of these two

parameters to approximate Eb1 amounts at MT plus ends. The strong hypomorphic $Eb1^{04524}$
mutant allele is known to display severe, but not complete reduction of protein levels (Elliott et
al., 2005); accordingly, Eb1 amounts at MT plus ends were severely, but not completely
diminished in neurons mutant for this allele (Figs. 1D,J and 1-S1B). Out of the remaining seven
candidate factors, only two further genes showed the same qualitative mutant phenotype:
$msps^{A/A}$ showed a reduction almost as strong as $Eb1^{04524}$, and $tau^{KO/KO}$ mutant neurons showed
a milder but reliable Eb1 depletion (Fig. 1B,C, J). In all cases, the drop in Eb1 amounts was to
almost equal parts due to reductions in comet length and in intensity (Fig. 1-S2A,B) and, across
all genotypes tested, they correlated well with reduced comet velocities and lifetimes when
assessed in live imaging experiments (using the C-terminal Shot domain Shot-Ctail::GFP as
readout for plus end dynamics; Fig. 1K,L and 1-S2C-E). Our data demonstrate therefore that
similar comet length/velocity correlations made in *in vitro* (Roostalu, Thomas et al., 2020) are
relevant in cellular contexts.

To assess whether observed reductions in the number or dynamics of MT plus end comets
correlate with impaired axon growth, we performed tubulin staining and measured axon length.
Out of the eight factors, all but Futsch/MAP1B showed a decrease in axon length ranging
between 10 and 43% when compared to parallel control cultures with wild-type neurons (Figs.
1E-H, M and 1-S1C). In these experiments, a fraction of genotypes showed prominent MT
bundle disintegration where MTs are disorganised and display intertwined, criss-crossing curls
(white arrowheads in Fig. 1F-H). When quantifying these phenotypes using the MT
disorganisation index (MDI; see methods), four of eight candidates showed significant MT
disorganisation including loss of Tau, Msps, Shot and Eb1 (Figs. 1F-H,N and 1-S1D), of which
the latter two confirm previous reports (Alves-Silva et al., 2012).

Therefore, out of eight candidate factors assessed with four phenotypic readouts (drop in comet
numbers, reduced Eb1 amounts/dynamics, shorter axons, MT disorganisation), Futsch
deficiency was the only condition showing no obvious defects. In contrast, loss of Eb1, Msps
and Tau stood out by displaying all four phenotypes, of which Tau consistently has a milder
effect. We therefore aimed to understand why these three factors share such a characteristic
combination of phenotypes.

<u>Neuronal Eb1, Msps and Tau are functionally linked and required *in vivo*</u>

Next we assessed whether Eb1, Msps and Tau are functionally related. For this, we performed
genetic interaction studies where heterozygous conditions (i.e. one mutant and one normal
copy) of genes are used to reduce their respective protein levels: if reduced protein levels of
different genes combined in the same neurons cause a phenotype, this suggests that they
function in a common pathway. For our studies, we used larval neurons and five different
readouts: axon length (Fig. 2A), MT disorganisation (Fig. 2A'), Eb1 amounts at MT plus ends
(Fig. 2A''), comet numbers (Fig. 2-S1A) and comet dynamics (Fig. 2-S1B).

To assess the baseline, we analysed single-heterozygous mutant neurons ($Eb1^{04524/+}$, $msps^{A/+}$
or $tau^{KO/+}$), none of which displayed any phenotypes (Figs. 2A-A'' and 2-S1A,B). However, when
bringing heterozygous conditions of these genes together in the same neurons, certain genetic
combinations displayed significant phenotypes (Fig. 2A'-A'' and 2-S1A,B): $Eb1^{04524/+}$ $msps^{A/+}$
and $tau^{KO/+}$ $msps^{A/+}$ double-heterozygous neurons displayed only a mild but significant reduction
in Eb1 amounts at MT plus ends and a trend towards stronger MT disorganisation; the
phenotypes were further enhanced for all five readouts in $Eb1^{04524/+}$ $msps^{A/+}$ $tau^{KO/+}$ triple-
heterozygous neurons. These data suggest that the three factors are functionally linked when

regulating MT plus ends, axon growth and MT organisation.

The triple-heterozygous condition had a similar effect in fly brains *in vivo*. We could show this
using the T1 subset of medulla neurons in the adult optic lobe which contain prominent axonal
MTs, arranged into tight bundles that can be visualised with α-tubulin::GFP labelling (Qu et al.,
2019). We found that the axons of T1 medulla neurons from triple-heterozygous mutant adult
flies show a strong increase in areas of MT bundle disorganisation compared to controls, with
MTs becoming unbundled and twisted and axons displaying prominent swellings (Fig.3A-D).
These data strongly suggest that the functional network of Eb1, Msps and Tau is relevant *in*
vivo.

<u>The functional unit of Eb1, Msps and Tau is evolutionarily conserved in *Xenopus* neurons</u>

In order to evaluate whether the functional network of Eb1, Msps and Tau might be conserved
evolutionarily, we used frog primary neurons. In the frog *Xenopus laevis*, there is only one *tau*
gene (*mapt/microtubule associated protein tau*), *XMAP215* is the only *msps* homologue
(*ckap5/cytoskeleton associated protein 5*), and *EB3* (*mapre3/microtubule associated protein*
RP/EB family member 3) is the only one of three *Eb1* homologues that is prominently expressed
in the nervous system (Bowes, Snyder et al., 2009, Karimi, Fortriede et al., 2017). We used
morpholinos against these three genes. Similar to our strategy in *Drosophila*, we analysed MTs
by staining for endogenous tubulin (Fig.3E-E'''), and measured Eb3 comet amounts in live
movies using the Eb protein-binding peptide MACF43-Ctail::GFP as readout (see Methods;
Honnappa et al., 2009, Slater et al., 2019).

To approximate heterozygous mutant conditions used in our *Drosophila* experiments, we
adjusted morpholino concentrations to levels that achieved knock-down of each of the three
genes to ~50% (Fig. 3-S1A-B and (Lowery et al., 2013, Slater et al., 2019). Individual knock-
downs to approximately 50% did not cause prominent decreases in MACF43::GFP comet
amounts or increases in MT disorganisation; but when knock-down of all three factors was
combined in the same neurons, we found a reduction in MACF43::GFP comet amounts to 60%
and a 4.8 fold increase in MT disorganisation (Figs. 3E''',F''',G,H and 3-S1C,D).

Together, these results suggest that there is functional interaction between the three genes also
in *Xenopus*, and that their operation as a functional unit is evolutionarily conserved. This raises
the question as to what mechanisms can explain the joined function of these proteins.

<u>Eb1, Msps and Tau operate as one functional unit with key roles played by Eb1</u>

We next asked whether the three factors are hierarchically and/or interdependently organised
into a functional unit, or whether they regulate the assessed MT properties through independent
parallel mechanisms. To distinguish between these possibilities (Avery & Wasserman, 1992),
we combined homozygous mutant conditions of the three mutant alleles in the same neurons
(*Eb1$^{04524/04524}$ msps$^{A/A}$ tau$^{KO/KO}$*) and asked whether this condition enhances phenotypes over
single-homozygous conditions (indicating parallel mechanisms), or whether they show no
further increases (reflecting a functional unit).

Since the combined triple-homozygous condition does not survive into the late larval stage, we
used embryonic pre-cultured neurons for these experiments. As reference we used
Eb1$^{04524/04524}$, *msps$^{A/A}$* and *tau$^{KO/KO}$* single mutant neurons; since *Eb1^{04524}* is a strong but not a
total loss-of-function allele (Elliott et al., 2005), we added neurons homozygous for

Df(2R)Exel6050 (*Eb1^Df*), a deficiency uncovering the entire *Eb1* locus (see Methods). Of the
four conditions, *Eb1^{Df/Df}* neurons showed the strongest phenotypes, thus setting the reference
bar for the triple-homozygous mutant neurons; we found that having the mutations combined
did not enhance the phenotypes beyond that of *Eb1^Df* (Fig. 2B-B'').

These results are consistent with a model where the three proteins act as a functional unit.
Within this unit, the three factors show the same qualitative loss-of-function phenotypes.
However, there is a clear order of severity with *Eb1^{Df/Df}* > *msps^{1/1}* > *tau^{KO/KO}* across all phenotypes
assessed (Figs.2, 2-S1; see Discussion), suggesting that Eb1 is of particular importance within
the functional unit. This is also highlighted when plotting Eb1 amounts against MT
disorganisation from a range of different genetic conditions analysed throughout this study
(Fig.2-S2A). This plot revealed that there is a highly significant inverse correlation between Eb1
comet amounts and MT disorganisation (Figs. 2C and 2-S2), suggesting that Eb1 is the key
factor within the functional unit that mediates to MT bundle promotion.

<u>Eb1 and Msps depend on each other for MT plus end localisation</u>

To understand the functional unit, we next focussed on the mechanistic links between Msps
and Eb1. First we started with co-localisation studies by co-expressing Msps::GFP together with
Eb1::RFP in *Drosophila* embryonic primary neurons. In these experiments, both proteins
prominently localised at the same MT plus ends, with Msps::GFP localising slightly distal to Eb1
(Fig. 4A-A'', Movie M1), as was similarly reported *in vitro* (Maurer, Cade et al., 2014). Their
localisation at the same MT plus ends was even clearer in kymographs of live movies where
both proteins remained closely associated during comet dynamics (Fig.4B,B').

In previous sections we reported that loss of Msps causes severe Eb1 depletion at MT plus
ends (Fig1. 1B, D and 1-S1B), suggesting a requirement of Msps for Eb1's MT plus-end
localisation. We hypothesised that the role of Msps as a tubulin polymerase (Brouhard et al.,
2008) helps to sustain a prominent GTP-tubulin cap as important prerequisite for EB1 binding
(Zanic et al., 2013). In support of this hypothesis, immunostaining for GTP-tubulin in *msps^1*
mutant neurons revealed length reductions of GTP caps that were as severe as the reduction
observed for Eb1 comets (Figs. 1J *vs.* 5-S2C). This result together with our finding of reduced
comet velocity in *msps^1* mutant neurons (Fig. 1K) are consistent with our hypothesis that Msps-
dependent polymerisation increases Eb1 amounts at MT plus ends (see Discussion).

Next we investigated whether Eb1 may influence Msps function. We expressed Msps::GFP in
Eb1^{04524} mutant neurons and found a severe depletion of Msps at growing MT ends when
compared to controls (Fig. 4C, C'; Movies M2 and M3), suggesting that Msps is no longer
efficiently recruited to the plus ends of polymerising MTs. Mechanistically, it was shown *in vitro*
and other cell systems that Msps/XMAP215 can bind MT plus ends independently, but that Eb
proteins can enhance its binding (Maurer et al., 2014, Zanic et al., 2013). Such enhanced Eb
protein-dependent recruitment of Msps/XMAP215 can be mediated by adaptors such as SLAIN
in vertebrates or TACC (Transforming acidic coiled-coil protein) or Sentin in non-neuronal
Drosophila cells (Brouhard et al., 2008, Lee, Gergely et al., 2001, Li, Miki et al., 2011, Li et al.,
2012, Lowery et al., 2013, Tang, Rui et al., 2020, van der Vaart, Franker et al., 2012). To test
potential roles of candidate adaptors in fly neurons, we performed functional studies with *tacc^1*
and *sentin^{ΔB}* loss-of-function mutant alleles. We found that primary neurons carrying these
mutant alleles failed to display axon shortening or MT disorganisation, thus arguing against a
prominent role of these factors to Eb1-dependent Msps recruitment (details in Fig.4-S1A,B). As

a complementary approach, we generated a *msps$^{\Delta Cterm}$-GFP* construct which lacks the C-
terminal domain essential for the interaction with adaptors (Fig.4D; Fox, Howard et al., 2014,
Mortuza, Cavazza et al., 2014). When *msps$^{\Delta Cterm}$-GFP* or *msps^{FL}-GFP* length controls were
transfected into *msps$^{A/146}$* mutant neurons, we found that both protein variants were similarly
able to improve axon length and MT disorganisation defects, further arguing against the
requirement of adaptors (Fig.4D,D') in fly neurons.

In conclusion, our data suggest a scenario where Eb1 and Msps require each other to achieve
prominent MT plus-end localisation. Msps likely maintains Eb1 at MT plus ends through
promoting GTP-cap formation. *Vice versa,* Eb1 facilitates Msps recruitment through a
mechanism that might involve structural maturation of MT plus ends (see Discussion; Maurer
et al., 2014, Zanic et al., 2013).

<u>Tau promotes Eb1 pools at MT plus ends by outcompeting it from lattice binding</u>

Similar to Msps, we found that also loss of Tau leads to a reduction of Eb1 comet sizes in both
Drosophila (Fig.1C,J and 1-S2A,B) and *Xenopus* neurons (Fig.3-S1G-I). Immuno-histochemical
studies of wild-type fly neurons revealed Tau localisation along MT lattices which does not
extend into the Eb1 comet at the MT plus end (Fig.5A'A''). This distribution is consistent with
reports that Tau has a higher affinity for GDP-tubulin (Castle, McKibben et al., 2020, Duan,
Jonasson et al., 2017) and does suggest that Tau promotes Eb1 plus end localisation through
an indirect mechanisms which does not involve co-localisation.

We noticed that the reduction of Eb1 comet sizes in Tau-deficient neurons is accompanied by
a 20% increase in the intensity of Eb1 all along MT lattices (Fig.5C,E for 6 day pre-culture, Fig.5-
S2A for 6HIV). This effect is specific to *tau* and not observed in *msps$^{A/A}$* mutant neurons (Fig.
5-S2A,B). Tau has previously been shown to protect MTs against Katanin-induced damage
(Qiang, Yu et al., 2006). Therefore, we reasoned that the increase of Eb1 on MT lattices upon
loss of Tau could be a consequence of MT repair mechanisms leading to an increase in GTP-
tubulin along the lattice which could recruit Eb1 (Vemu, Szczesna et al., 2018). However, in
spite of increased Eb1 binding, MT lattices in Tau-deficient neurons did not show obvious
increases in GTP-tubulin (Fig. 5-S1B, D), arguing against the repair hypothesis. In the same
specimens, GTP-tubulin amounts at MT plus ends were clearly reduced, thus mirroring Eb1
comet depletion in Tau deficient neurons (Figs. 1J *vs*. 5-S1A-C) and suggesting that GTP-
tubulin staining is sufficiently sensitive to make quantitative statements in our model.

We next hypothesised that Tau may competitively prevent Eb1 from binding the MT lattice, as
similarly observed for other MAPs (Qiang, Sun et al., 2018). In support of this notion, we found
that low levels of Eb1 localisation along MT lattices observed in wild-type neurons are further
decreased when Tau is over-expressed (Fig. 5-S1E). A similar phenomenon has been observed
in vitro with mammalian versions of the two proteins (Ramirez-Rios, Denarier et al., 2016)
suggesting that this mechanism is evolutionarily conserved.

We reasoned therefore that MT lattices may turn into a sink for Eb1 if Tau is absent. Given the
high density of MTs in the narrow axons, this aberrant binding could sequester Eb1 pools away
from MT plus ends. To test this possibility, we expressed Eb1::GFP in *tau$^{KO/KO}$* mutant neurons
and stained them with anti-Eb1 antibodies. In support of our hypothesis, these neurons showed
a substantial further increase of Eb1 along MT lattices, but also replenished Eb1 amounts at
MT plus ends (Fig. 5D-F). This treatment was sufficient to suppress Tau-deficient phenotypes,
as reflected in the recovery of Eb1 comet dynamics (improved velocity and lifetime) and strongly

reduced MT disorganisation ('Eb1-GFP' in Fig. 5G-I). The latter finding further supports a role
of Eb1 in promoting the bundled conformation of axonal MTs (Fig.2C, Fig.2-S2; see Discussion).

Specificity of this mechanism for Tau is demonstrated by parallel experiments expressing
Eb1::GFP in $msps^{A/A}$ mutant neurons which failed to restore their MT plus end dynamics (Fig.
5-S2D,E). This result also provides further important information about the hierarchical
relationships between Eb1, Msps and Tau within the functional unit (see Discussion). As a
further control, we used Shot-Ctail::GFP (comparable to MACF43::GFP used in *Xenopus*; Figs.
3F and 6A), thus tracking MT plus ends without altering Eb1 levels. When expressing it in
$tau^{KO/KO}$ mutant neurons, no rescue of MT plus end comet velocity and lifetime was observed
('Ctail-GFP' in Fig.5G,H).

In conclusion, we propose that Tau contributes to MT polymerisation dynamics and MT
organisation in an indirect way through preventing that Eb1 is sequestered away from MT plus
ends. This function of Tau might be particularly important in axons where relative MT densities
are high (Prokop, 2020; see Discussion).

<u>An Eb1- and spectraplakin-dependent guidance mechanisms explains roles of the functional</u>
<u>unit in MT bundle organisation</u>

As explained before, Eb1 amounts at MT plus ends inversely correlate with MT disorganisation
(Figs.2C and 2-S2). Observed MT disorganisation in either Eb1-, Msps- or Tau-deficient
neurons might therefore be a consequence of Eb1 loss from plus ends. We hypothesised that
this phenomenon can be explained through a mechanism involving the *Drosophila* spectraplakin
Shot. Shot has been proposed to link Eb1 at polymerising MT plus ends to cortical F-actin, in
this way guiding the extension of MTs along the axonal surface into parallel bundles.
Accordingly, depletion of either Shot or Eb1 causes MT disorganisation (Alves-Silva et al., 2012,
Voelzmann, Liew et al., 2017; Fig.6F).

In support of this guidance hypothesis, we found that severe MT disorganisation observed in
$Eb1^{+/-}$ $msps^{+/-}$ $tau^{+/-}$ triple-heterozygous mutant neurons (which have reduced Eb1 comet
amounts; Fig.2A'') was significantly improved from 7.3-fold (with GFP-expression) to 1.4-fold
when over-expressing full length Shot-FL::GFP (Fig.6B-C; both compared to wild-type controls).

To exclude that this rescue is mediated through different Shot-dependent mechanisms, in
particular its role in MT stabilisation (Voelzmann et al., 2017), we repeated these experiments
with Shot versions that maintain MT-stabilising activity but affect actin-Eb1 cross-linkage in two
specific ways: (1) Shot$^{\Delta ABD}$::GFP lacks the N-terminal calponin homology domain required for
interaction with the actin cortex; (2) Shot^{3MTLS}::GFP carries mutations in the C-terminal SxIP
motifs required for the specific binding to Eb1 (Fig. 6A and F; see also Fig. 7A). Both Shot
variants failed to rescue MT disorganisation in $Eb1^{04524/+}$ $msps^{1/+}$ $tau^{KO/+}$ triple-heterozygous
mutant neurons (Fig.6B,C), consistent with a model where where Shot guides MTs downstream
of Eb1 through cross-linking it to cortical F-actin. This conclusion is further supported by the
finding that MT disorganisation observed in $Eb1^{04524/04524}$ mutant neurons is not further
enhanced in $Eb1^{04524/04524}$ $shot^{3/3}$ double-mutant neurons (Figs. 6D, E).

We conclude that the polymerisation-promoting functional unit of Eb1, Msps and Tau appears
to act through Shot-mediated MT guidance downstream of Eb1 to perform its function in axonal
bundle organisation, as a key mechanism contributing to the formation and long-term
maintenance of axons.

Discussion

New understanding of the role and regulation of MT polymerisation and guidance in axons

Understanding the machinery of MT polymerisation is of utmost importance in axons where MTs form loose bundles of enormous lengths which are essential for axonal morphogenesis and serve as life-sustaining transport highways (Prokop, 2020). They have to be maintained in functional state for up to a century in humans (Hahn et al., 2019). For this, MT polymerisation is required to generate new MTs, or to repair or replace them. The underpinning machinery is expected to be complex (Voelzmann et al., 2016), but its understanding will deliver new strategies for tackling developmental and degenerative axon pathologies.

Here we made important advances to this end. Having screened through 13 candidates (here and (Beaven et al., 2015), we found the three factors Eb1, Msps and Tau to stand out by expressing the same combination of phenotypes, and displaying strong functional interaction in *Drosophila* that is conserved in *Xenopus* neurons. We found that this machinery is not only important to maintain MT mass by driving polymerisation, but also for arranging MTs into bundles, thus highlighting MT polymerisation as a doubly important process for axon formation and maintenance.

Through carrying out this work in fly, we were able to demonstrate that the functional collaboration of the three factors is relevant *in vivo* and gain an understanding of the underlying mechanisms, both with respect to promoting MT polymerisation and their arrangement into parallel bundles (see details in Fig.7 and below).

Our mechanistic models are strongly supported by published data obtained through *in vitro* studies of MTs that match extremely well with virtually all our findings. Key examples are: (1) the complementary binding preferences of EB1 and Tau for GTP-/GDP-tubulin (Castle et al., 2020, Zanic, Stear et al., 2009); (2) the mutual enhancement of Eb1 and XMAP215/Msps originally suggested in *Xenopus* extracts and then demonstrated in reconstitution essays with purified proteins from fly and vertebrates (Kronja, Kruljac-Letunic et al., 2009, Li et al., 2012, Zanic et al., 2013); (3) the correlation of GTP cap size with comet velocity (Roostalu et al., 2020); (4) the reduction of comet numbers but not dynamics (Fig.1-S1) upon depletion of α1-tubulin or Stathmin (a prominent regulator of tubulin availability; Duncan et al., 2013, Manna, Grenningloh et al., 2007) which are consistent with *in vitro* observations that MT nucleation is far more sensitive to tubulin levels than polymerisation (Consolati, Locke et al., 2020). This list of similarities is in stark support of our model and a clear indication that many *in vitro* observation seem to apply in cellular contexts.

In the following we will discuss a mechanistic model (summarised in Fig.7) able to coherently integrate our experimental findings in neurons with current knowledge obtained from *in vitro* studies.

Eb1 and XMAP215/Msps are core factors promoting MT polymerisation and guidance

At the core of this model lie Eb1 and XMap215/Msps. Of these, vertebrate XMAP215 and fly Msps are both known to be relevant for neuronal morphogenesis in fly and *Xenopus* (Lowery et al., 2013, Tang et al., 2020). Work in non-neuronal cells or in vitro have shown that Msps and XMAP215 both strongly enhance MT polymerisation, consistent with expected functions of this TOG-domain protein as a polymerase (Al-Bassam & Chang, 2011, Brouhard et al., 2008, Fox

et al., 2014, Howard & Hyman, 2009, Li et al., 2012, Zanic et al., 2013).

Eb1 is a known scaffold (Akhmanova & Steinmetz, 2015) and, accordingly, both *Drosophila* and
vertebrate Eb1 seem to be only moderate promoters of MT polymerisation *in vitro* (Li et al.,
2012, Ramirez-Rios et al., 2016, Zanic et al., 2013) and references within). Conserved binding
partners of Eb proteins are the spectraplakins that can guide extending MT plus ends by cross-
linking them to actin, relevant for MT guidance in axons and non-neuronal cells (Alves-Silva et
al., 2012, Voelzmann et al., 2017, Wu, Kodama et al., 2008). Although spectraplakins might not
be the only interactors involved in Eb1-dependent axonal MT guidance (Hahn et al., 2019), their
contribution is prominent and our data clearly support this role of Shot (Fig.7E,E').

Taken together, it seems therefore reasonable to assume that, within the functional unit, Msps
is the key promoter of polymerisation, and Eb1 the key mediator of MT guidance into bundles
(Fig.7A,A'). However, to perform these functions, both factors have to be enriched in sufficient
amounts at MT plus ends. For this, both proteins depend on each other: taking out Eb1 detaches
not only Shot thus abolishing guidance, but also negatively impacts Msps localisation hence
polymerisation (Fig.7B,B'); the same is true in reverse when removing Msps (Fig.7C,C') and
may explain why loss of XMAP215 was reported to affect MT guidance in growth cones of frog
neurons (Slater et al., 2019).

This mutual dependency is unlikely to involve their physical interaction, since MT plus end
localisation of Eb1 is known to occur tens of nanometres behind XMAP215 (Maurer et al., 2014,
Zanic et al., 2013). Furthermore, as detailed in the Results part, our data do not support an
obvious role of adaptors in mediating Eb1- XMAP215 interactions (Figs.4D-D'' and 4-S1). This
deviates from other cellular contexts in which EB1 adaptors are required for Msps/XMAP215
MT plus end localisation (Lee et al., 2001, Li et al., 2011, Li et al., 2012, Nwagbara, Faris et al.,
2014, Tang et al., 2020, van der Vaart et al., 2012), potentially reflecting context-specific
variations in MT regulation; this has similarly been reported for the plus end-associating factor
Clip170/190 which fails to form comets in axons of fly and mouse neurons (Beaven et al., 2015).

Instead, indirect mechanisms for mutual dependency of Eb1 and XMAP215/Msps were
proposed in the context of *in vitro* studies: through promoting MT polymerisation, XMAP215
maintains a prominent GTP-cap thus enabling the binding of substantial amounts of Eb1
molecules (Maurer et al., 2014, Zanic et al., 2013). Restricted GTP-cap formation as a limiting
factor for Eb1 binding would also explain why Eb1 over-expression fails to rescue Msps-
deficient phenotypes (Fig.5-S2D-F). *Vice versa*, it was suggested that Eb1's ability to promote
lateral protofilament contacts could assist in sheet formation at the very plus tip, thus facilitating
the binding of XMAP215/Msps (Maurer et al., 2014, Zanic et al., 2013). We believe this
mechanism to be the most likely to explain our findings.

<u>Tau contributes through an indirect mechanism of competitive binding to MT lattices</u>

Tau and Map1b/Futsch are known to promote MT polymerisation *in vitro* and axon growth in
mouse and fly neurons through mechanisms that remain unclear (Brandt & Lee, 1993, Caceres
& Kosik, 1990, Cleveland, Hwo et al., 1977, DiTella, Feiguin et al., 1996, Drechsel et al., 1992,
Hummel et al., 2000, Kadavath, Hofele et al., 2015, Kiris, Ventimiglia et al., 2010, Levy, Leboeuf
et al., 2005, Liu et al., 2015, Panda, Goode et al., 1995, Ramirez-Rios et al., 2016, Takei, Teng
et al., 2000, Tymanskyj, Scales et al., 2012, Villarroel-Campos & Gonzalez-Billault, 2014).

In our cellular model, loss of Map1b/Futsch has no obvious effects, whereas Tau shares a
surprising number of loss-of-function mutant phenotypes with those of Msps and Eb1. At least

one of these, loss of Eb proteins from MT plus ends, appears an evolutionarily well conserved
phenomenon observed in fly neurons (Figs.1C,J), frog neurons (Fig.3-S1G-I), N1E-115 mouse
neuroblastoma cells and primary mouse cortical neurons (Sayas, Tortosa et al., 2015). Sayas
and co-workers originally reported also tau-mediated recruitment of Eb proteins to the MT lattice
(analogous to MAP2-mediated Eb3 recruitment to MTs in dendrites; Kapitein, Yau et al., 2011).
However, their subsequent *in vitro* work suggested that Tau outcompetes Eb1 at the lattice and
Eb proteins outcompete Tau at MT plus ends (Ramirez-Rios et al., 2016), potentially involving
sequestration through direct interaction in the cytoplasm - as would be consistent with other
reports that Tau can bind Eb1 (Buey, Mohan et al., 2011, Duan et al., 2017).

We do not argue against such interaction and that it can potentially contribute to Eb protein loss
from MT plus ends when tau is over-expressed (Ramirez-Rios et al., 2016); but it cannot explain
why Eb1 comets are reduced when Tau is absent. Instead, the mechanism we propose is
based on the known complementary binding preferences of Eb1 and Tau for GTP-/GDP-tubulin
(Castle et al., 2020, Duan et al., 2017, Zanic et al., 2009). Thus, we find that Tau binds
preferentially along MT shafts but not at plus ends, whereas Eb1 binds in complementary
fashion and displays elevated shaft binding when Tau is absent (Fig.5C and 7D'). Such
competitive binding behaviour is reminiscent of Tau's role in preventing MAP6 from binding in
certain regions of the MT lattice (Baas & Qiang, 2019, Qiang et al., 2018); it is in agreement
with the idea of a MAP code where similar MAP competition is assumed to regulate region-
specific axonal transport (Monroy, Tan et al., 2020).

Given the high density of MTs in the narrow space of axons, we propose that lattice binding can
generate a sink large enough to reduce Eb1 levels at MT plus ends, and our rescue experiments
with Eb1::GFP overexpression strongly support this notion. In this way, loss of Tau generates
a condition comparable to a modest Eb1 loss-of-function mutant phenotype, thus explaining
why Tau shares its repertoire of loss-of-function phenotypes with Msps and Eb1 but with lower
penetrance.

We propose that similar mechanisms might operate in smaller diameter axons in vertebrates,
such as in parallel fibres of the cerebellum where Tau has demonstrated structural roles
(Harada, Oguchi et al., 1994); they might explain the aforementioned findings of small Eb protein
comets in cultured vertebrate neurons, as well as the reduction in MT numbers observed upon
loss of Tau in *C. elegans* (Krieg, Stühmer et al., 2017). However, in larger diameter axons of
vertebrates where MT densities are low (Prokop, 2020), the Eb1 depletion effect might be far
less noticeable.

<u>Main conclusions and future perspectives</u>

Here we have gained new understanding of MT polymerisation regulation in axons and have
been able to propose a solid and consistent mechanistic model. This model aligns well with
known *in vitro* data and our previous mechanistic models explaining Eb1/Shot-mediated MT
guidance (Alves-Silva et al., 2012) and complementary functions of Shot and the cortical
collapse factor Efa6 (Qu et al., 2019).

Developing such refined mechanistic models of increasing complexity is made possible through
using one standardised *Drosophila* neuron system amenable to combinatorial genetic
approaches - with enormous capacity to extend our understanding even further to other factors
known to regulate MT polymerisation *in vitro* (Zanic et al., 2013). Gaining an understanding of
the machinery regulating axonal MT polymerisation opens up new ways to investigate the

mechanisms behind other important observations, such as the finding that cortical actin has a
regulatory impact on MT polymerisation in axons (Qu et al., 2017).

As another example, we found that loss of either Eb1, XMAP215/Msps or Tau all caused a
reduction in comet numbers, potentially reflecting changes in MT nucleation activity, consistent
with reports of nucleation-promoting roles of XMAP215 (Flor-Parra, Iglesias-Romero et al.,
2018, Roostalu, Cade et al., 2015, Thawani, Kadzik et al., 2018, Wieczorek, Bechstedt et al.,
2015). Extending work from MT polymerisation to nucleation is possible in the fly system and
would have the potential to deliver explanations for how numbers of MTs can be regulated in
reproducible, neuron-specific ways, thus addressing a fundamental aspect of axon morphology
(Prokop, 2020).

By gradually assembling molecular mechanisms into regulatory networks that can explain
axonal MT regulation at the cellular level, our studies come closer to explaining axonal
pathologies which can then form the basis for the development of remedial strategies (Hahn et
al., 2019).

Acknowledgements

This work was made possible through support by the BBSRC to A.P (BB/I002448/1,
BB/P020151/1, BB/L000717/1, BB/M007553/1) to N.S.S. (BB/M007456/1, BB/R018960/1), by
the Leverhulme Trust to I.H. (ECF-2017-247), by the German Research Council (DFG) to A.V.
(VO 2071/1-1), by NIH to L.A.L (R01 MH109651), and a postdoctoral fellowship from CONICYT
to P.G.S. The Manchester Bioimaging Facility microscopes used in this study were purchased
with grants from the BBSRC, The Wellcome Trust and The University of Manchester Strategic
Fund. The Fly Facility has been supported by funds from The University of Manchester and the
Wellcome Trust (087742/Z/08/Z). We thank Hiro Ohkura for kindly providing DmEb1 antibody
and unpublished mutant alleles of *msps*. Stocks obtained from the Bloomington *Drosophila*
Stock Center (NIH P40OD018537) were used in this study.

References

Adalbert R, Coleman MP (2012) Axon pathology in age-related neurodegenerative disorders.
Neuropathol Appl Neurobiol 39: 90–108

Aher A, Rai D, Schaedel L, Gaillard J, John K, Liu Q, Altelaar M, Blanchoin L, Thery M,
Akhmanova A (2020) CLASP mediates microtubule repair by restricting lattice damage and
regulating tubulin incorporation. *Current Biology*

Akhmanova A, Steinmetz MO (2015) Control of microtubule organization and dynamics: two
ends in the limelight. *Nat Rev Mol Cell Biol* 16: 711–726

Al-Bassam J, Chang F (2011) Regulation of microtubule dynamics by TOG-domain proteins
XMAP215/Dis1 and CLASP. *Trends Cell Biol* 21: 604-14

Al-Bassam J, Kim H, Brouhard G, van Oijen A, Harrison SC, Chang F (2010) CLASP promotes
microtubule rescue by recruiting tubulin dimers to the microtubule. *Dev Cell* 19: 245-58

Alves-Silva J, Sánchez-Soriano N, Beaven R, Klein M, Parkin J, Millard T, Bellen H, Venken
KJT, Ballestrem C, Kammerer RA, Prokop A (2012) Spectraplakins promote microtubule-
mediated axonal growth by functioning as structural microtubule-associated proteins and
EB1-dependent +TIPs (Tip Interacting Proteins). *J Neurosci* 32: 9143-58

Avery L, Wasserman S (1992) Ordering gene function: the interpretation of epistasis in
regulatory hierarchies. *Trends Genet* 8: 312-316

Baas PW, Qiang L (2019) Tau: it's not what you think. *Trends Cell Biol* 29: 452-461

Beaven R, Dzhindzhev NS, Qu Y, Hahn I, Dajas-Bailador F, Ohkura H, Prokop A (2015)
Drosophila CLIP-190 and mammalian CLIP-170 display reduced microtubule plus end
association in the nervous system. *Mol Biol Cell* 26: 1491-508

Bowes JB, Snyder KA, Segerdell E, Jarabek CJ, Azam K, Zorn AM, Vize PD (2009) Xenbase:
gene expression and improved integration. *Nucleic Acids Research* 38: D607-D612

Brandt R, Lee G (1993) The balance between tau protein's microtubule growth and nucleation
activities: implications for the formation of axonal microtubules. *J Neurochem* 61: 997-1005

Brittle AL, Ohkura H (2005) Mini spindles, the XMAP215 homologue, suppresses pausing of
interphase microtubules in *Drosophila*. *EMBO J* 24: 1387-96

Brouhard GJ, Stear JH, Noetzel TL, Al-Bassam J, Kinoshita K, Harrison SC, Howard J, Hyman
AA (2008) XMAP215 is a processive microtubule polymerase. *Cell* 132: 79-88

Buey RM, Mohan R, Leslie K, Walzthoeni T, Missimer JH, Menzel A, Bjelic S, Bargsten K,
Grigoriev I, Smal I, Meijering E, Aebersold R, Akhmanova A, Steinmetz MO (2011) Insights
into EB1 structure and the role of its C-terminal domain for discriminating microtubule tips
from the lattice. *Mol Biol Cell* 22: 2912-23

Burnouf S, Gronke S, Augustin H, Dols J, Gorsky MK, Werner J, Kerr F, Alic N, Martinez P,
Partridge L (2016) Deletion of endogenous Tau proteins is not detrimental in *Drosophila*. *Sci*
Rep 6: 23102

Caceres A, Kosik KS (1990) Inhibition of neurite polarity by tau antisense oligonucleotides in
primary cerebellar neurons. *Nature* 343: 461-3

Calkins DJ (2013) Age-Related Changes in the Visual Pathways: Blame It on the AxonAge-
Related Changes in the Visual Pathways. *Invest Ophthalmol Vis Sci* 54: ORSF 37-41

Castle BT, McKibben KM, Rhoades E, Odde DJ (2020) Tau avoids the GTP cap at growing
microtubule plus ends. *bioRxiv*: 2019.12.31.891234

Cleveland DW, Hwo SY, Kirschner MW (1977) Purification of tau, a microtubule-associated
protein that induces assembly of microtubules from purified tubulin. *J Mol Biol* 116: 207-25

Consolati T, Locke J, Roostalu J, Chen ZA, Gannon J, Asthana J, Lim WM, Martino F, Cvetkovic
MA, Rappsilber J, Costa A, Surrey T (2020) Microtubule Nucleation Properties of Single
Human γTuRCs Explained by Their Cryo-EM Structure. *Developmental Cell* 53: 603-617.e8

Dimitrov A, Quesnoit MI, Moutel S, Cantaloube I, Po?s C, Perez F (2008) Detection of GTP-
tubulin conformation *in vivo* reveals a role for GTP remnants in microtubule rescues. *Science*
322: 1353-1356

DiTella MC, Feiguin F, Carri N, Kosik KS, Caceres A (1996) MAP-1B/TAU functional
redundancy during laminin-enhanced axonal growth. *J Cell Sci* 109 (Pt 2): 467-77

Doerflinger H, Benton R, Shulman JM, St Johnston D (2003) The role of PAR-1 in regulating
the polarised microtubule cytoskeleton in the Drosophila follicular epithelium. *Development*
130: 3965-75

Drechsel DN, Hyman AA, Cobb MH, Kirschner MW (1992) Modulation of the dynamic instability
of tubulin assembly by the microtubule-associated protein tau. *Mol Biol Cell* 3: 1141-54

Duan AR, Jonasson EM, Alberico EO, Li C, Scripture JP, Miller RA, Alber MS, Goodson HV
(2017) Interactions between tau and different conformations of tubulin: implications for tau
function and mechanism. *J Mol Biol* 429: 1424-1438

Duncan IW, Kaufman TC (1975) Cytogenic analysis of chromosome 3 in *Drosophila*
melanogaster: mapping of the proximal portion of the right arm. *Genetics* 80: 733-52

Duncan JE, Lytle NK, Zuniga A, Goldstein LS (2013) The microtubule regulatory protein
Stathmin is required to maintain the integrity of axonal microtubules in *Drosophila*. *PLoS One*
8: e68324

Elliott SL, Cullen CF, Wrobel N, Kernan MJ, Ohkura H (2005) EB1 is essential during *Drosophila*
development and plays a crucial role in the integrity of chordotonal mechanosensory organs.
Mol Biol Cell 16: 891-901

Flor-Parra I, Iglesias-Romero AB, Chang F (2018) The XMAP215 ortholog Alp14 promotes
microtubule nucleation in fission yeast. *Curr Biol* 28: 1681-1691.e4

Fox JC, Howard AE, Currie JD, Rogers SL, Slep KC (2014) The XMAP215 family drives
microtubule polymerization using a structurally diverse TOG array. *Mol Biol Cell* 25: 2375-92

Gasic I, Mitchison TJ (2019) Autoregulation and repair in microtubule homeostasis. *Curr Opin*
Cell Biol 56: 80-87

Gergely F, Kidd D, Jeffers K, Wakefield JG, Raff JW (2000) D-TACC: a novel centrosomal
protein required for normal spindle function in the early Drosophila embryo. *EMBO J* 19: 241-
52

Gluszek AA, Cullen CF, Li W, Battaglia RA, Radford SJ, Costa MF, McKim KS, Goshima G,
Ohkura H (2015) The microtubule catastrophe promoter Sentin delays stable kinetochore-
microtubule attachment in oocytes. *J Cell Biol* 211: 1113-20

Grieder NC, de Cuevas M, Spradling AC (2000) The fusome organizes the microtubule network
during oocyte differentiation in Drosophila. *Development* 127: 4253-4264

Hahn I, Voelzmann A, Liew Y-T, Costa-Gomes B, Prokop A (2019) The model of local axon
homeostasis - explaining the role and regulation of microtubule bundles in axon maintenance
and pathology *Neural Dev* 14: 10.1186/s13064-019-0134-0

Harada A, Oguchi K, Okabe S, Kuno J, Terada S, Ohshima T, Sato-Yoshitake R, Takei Y, Noda
T, Hirokawa N (1994) Altered microtubule organization in small-calibre axons of mice lacking
tau protein. *Nature* 369: 488-91

Honnappa S, Gouveia SM, Weisbrich A, Damberger FF, Bhavesh NS, Jawhari H, Grigoriev I,
van Rijssel FJ, Buey RM, Lawera A, Jelesarov I, Winkler FK, Wuthrich K, Akhmanova A,
Steinmetz MO (2009) An EB1-binding motif acts as a microtubule tip localization signal. *Cell*
138: 366-76

Howard J, Hyman AA (2009) Growth, fluctuation and switching at microtubule plus ends. *Nat*
Rev Mol Cell Biol 10: 569-74

Hummel T, Krukkert K, Roos J, Davis G, Klämbt C (2000) Drosophila Futsch/22C10 is a
MAP1B-like protein required for dendritic and axonal development. *Neuron* 26: 357-370

Inoue YH, do Carmo Avides M, Shiraki M, Deak P, Yamaguchi M, Nishimoto Y, Matsukage A,
Glover DM (2000) Orbit, a novel microtubule-associated protein essential for mitosis in
Drosophila melanogaster. *J Cell Biol* 149: 153-66

Jenkins BV, Saunders HAJ, Record HL, Johnson-Schlitz DM, Wildonger J (2017) Effects of
mutating alpha-tubulin lysine 40 on sensory dendrite development. *J Cell Sci* 130: 4120-4131

Kadavath H, Hofele RV, Biernat J, Kumar S, Tepper K, Urlaub H, Mandelkow E, Zweckstetter
M (2015) Tau stabilizes microtubules by binding at the interface between tubulin
heterodimers. *Proc Natl Acad Sci* 112: 7501-7506

Kapitein LC, Yau KW, Gouveia SM, van der Zwan WA, Wulf PS, Keijzer N, Demmers J,
Jaworski J, Akhmanova A, Hoogenraad CC (2011) NMDA receptor activation suppresses
microtubule growth and spine entry. *J Neurosci* 31: 8194-209

Karimi K, Fortriede JD, Lotay VS, Burns KA, Wang DZ, Fisher ME, Pells TJ, James-Zorn C,
Wang Y, Ponferrada V G, Chu S, Chaturvedi P, Zorn AM, Vize PD (2017) Xenbase: a
genomic, epigenomic and transcriptomic model organism database. *Nucleic Acids Res* 46:
D861-D868

Kiris E, Ventimiglia D, Feinstein SC (2010) Quantitative analysis of MAP-mediated regulation of
microtubule dynamic instability in vitro focus on Tau. *Methods Cell Biol* 95: 481-503

Kolodziej PA, Jan LY, Jan YN (1995) Mutations that affect the length, fasciculation, or ventral
orientation of specific sensory axons in the Drosophila embryo. *Neuron* 15: 273-286

Krieg M, Stühmer J, Cueva JG, Fetter R, Spilker K, Cremers D, Shen K, Dunn AR, Goodman
MB (2017) Genetic defects in β-spectrin and tau sensitize C. elegans axons to movement-
induced damage via torque-tension coupling. *Elife* 6: e20172

Kronja I, Kruljac-Letunic A, Caudron-Herger M, Bieling P, Karsenti E (2009) XMAP215-EB1
interaction is required for proper spindle assembly and chromosome segregation in *Xenopus*
egg extract. *Mol Biol Cell* 20: 2684-96

Lee MJ, Gergely F, Jeffers K, Peak-Chew SY, Raff JW (2001) Msps/XMAP215 interacts with
the centrosomal protein D-TACC to regulate microtubule behaviour. *Nat Cell Biol* 3: 643-9

Levy SF, Leboeuf AC, Massie MR, Jordan MA, Wilson L, Feinstein SC (2005) Three- and four-
repeat tau regulate the dynamic instability of two distinct microtubule subpopulations in
qualitatively different manners. Implications for neurodegeneration. *J Biol Chem* 280: 13520-
8

Li W, Miki T, Watanabe T, Kakeno M, Sugiyama I, Kaibuchi K, Goshima G (2011) EB1 promotes
microtubule dynamics by recruiting Sentin in Drosophila cells. *J Cell Biol* 193: 973-83

Li W, Moriwaki T, Tani T, Watanabe T, Kaibuchi K, Goshima G (2012) Reconstitution of dynamic
microtubules with *Drosophila* XMAP215, EB1, and Sentin. *J Cell Biol* 199: 849-62

Liu Y, Wang C, Destin G, Szaro BG (2015) Microtubule-associated protein tau promotes
neuronal class II beta-tubulin microtubule formation and axon elongation in embryonic
Xenopus laevis. Eur J Neurosci 41: 1263-75

Lowery LA, Faris AER, Stout A, Van Vactor D (2012) Neural explant cultures from *Xenopus*
laevis. JoVE: e4232

Lowery LA, Stout A, Faris AE, Ding L, Baird MA, Davidson MW, Danuser G, Van Vactor D
(2013) Growth cone-specific functions of XMAP215 in restricting microtubule dynamics and
promoting axonal outgrowth. *Neural Dev* 8: 22

Luo L, Liao YJ, Jan LY, Jan YN (1994) Distinct morphogenetic functions of similar small
GTPases: Drosophila Drac1 is involved in axonal outgrowth and myoblast fusion. *Genes Dev*
8: 1787-1802

Manna T, Grenningloh G, Miller HP, Wilson L (2007) Stathmin family protein SCG10
differentially regulates the plus and minus end dynamics of microtubules at steady state *in*
vitro: implications for its role in neurite outgrowth. *Biochemistry* 46: 3543-52

Manna T, Thrower DA, Honnappa S, Steinmetz MO, Wilson L (2009) Regulation of microtubule
dynamic instability in vitro by differentially phosphorylated stathmin. *J Biol Chem* 284: 15640-
9

Marner L, Nyengaard JR, Tang Y, Pakkenberg B (2003) Marked loss of myelinated nerve fibers
in the human brain with age. *J Comp Neurol* 462: 144-52

Maurer SP, Cade NI, Bohner G, Gustafsson N, Boutant E, Surrey T (2014) EB1 accelerates two
conformational transitions important for microtubule maturation and dynamics. *Curr Biol* 24:
372-84

Monroy BY, Tan TC, Oclaman JM, Han JS, Simó S, Niwa S, Nowakowski DW, McKenney RJ,
Ori-McKenney KM (2020) A combinatorial MAP code dictates polarized microtubule
transport. *Dev Cell* 53

Mortuza GB, Cavazza T, Garcia-Mayoral MF, Hermida D, Peset I, Pedrero JG, Merino N, Blanco
FJ, Lyngsø J, Bruix M, Pedersen JS, Vernos I, Montoya G (2014) XTACC3–XMAP215
association reveals an asymmetric interaction promoting microtubule elongation. *Nature*
Communications 5: 5072

Nieuwkoop, J. F (1994) *Normal table of Xenopus laevis (Daudin): A systematical and*
chronological survey of the development from the fertilized egg till the end of metamorphosis.
Garland Pub. Inc. , New York

Nwagbara BU, Faris AE, Bearce EA, Erdogan B, Ebbert PT, Evans MF, Rutherford EL,
Enzenbacher TB, Lowery LA (2014) TACC3 is a microtubule plus-end tracking protein that

promotes axon elongation and also regulates microtubule plus-end dynamics in multiple
embryonic cell types. *Mol Biol Cell*

O'Neill EM, Rebay I, Tjian R, Rubin GM (1994) The activities of two Ets-related transcription
factors required for Drosophila eye development are modulated by the Ras/MAPK pathway.
Cell 78: 137-47.

Panda D, Goode BL, Feinstein SC, Wilson L (1995) Kinetic stabilization of microtubule
dynamics at steady state by tau and microtubule-binding domains of tau. *Biochemistry* 34:
11117-27

Prokop A (2013) A rough guide to Drosophila mating schemes. *figshare*:
dx.doi.org/10.6084/m9.figshare.106631

Prokop A (2020) Cytoskeletal organization of axons in vertebrates and invertebrates. *J Cell Biol*
219: e201912081

Prokop A, Beaven R, Qu Y, Sánchez-Soriano N (2013) Using fly genetics to dissect the
cytoskeletal machinery of neurons during axonal growth and maintenance. *J Cell Sci* 126:
2331-41

Prokop A, Küppers-Munther B, Sánchez-Soriano N (2012) Using primary neuron cultures of
Drosophila to analyse neuronal circuit formation and function. *The making and un-making of*
neuronal circuits in Drosophila 69: 225-47

Prokop A, Meinertzhagen IA (2006) Development and structure of synaptic contacts in
Drosophila. *Semin Cell Dev Biol* 17: 20-30

Qiang L, Sun X, Austin TO, Muralidharan H, Jean DC, Liu M, Yu W, Baas PW (2018) Tau does
not stabilize axonal microtubules but rather enables them to have long labile domains. *Curr*
Biol 28: 2181-2189 e4

Qiang L, Yu W, Andreadis A, Luo M, Baas PW (2006) Tau protects microtubules in the axon
from severing by katanin. *J Neurosci* 26: 3120-9

Qu Y (2015) Novel concepts of microtubule regulation during axon growth and maintenance.
Faculty of Life Sciences PhD: 205

Qu Y, Hahn I, Lees M, Parkin J, Voelzmann A, Dorey K, Rathbone A, Friel C, Allan V, Okenve
Ramos P, Sánchez-Soriano N, Prokop A (2019) Efa6 protects axons and regulates their
growth and branching by inhibiting microtubule polymerisation at the cortex. *eLife* 8: e50319

Qu Y, Hahn I, Webb SED, Pearce SP, Prokop A (2017) Periodic actin structures in neuronal
axons are required to maintain microtubules. *Mol Biol Cell* 28 296-308

Ramirez-Rios S, Denarier E, Prezel E, Vinit A, Stoppin-Mellet V, Devred F, Barbier P, Peyrot V,
Sayas CL, Avila J, Peris L, Andrieux A, Serre L, Fourest-Lieuvin A, Arnal I (2016) Tau
antagonizes end-binding protein tracking at microtubule ends through a phosphorylation-
dependent mechanism. *Mol Biol Cell* 27: 2924-2934

Rogers SL, Rogers GC, Sharp DJ, Vale RD (2002) Drosophila EB1 is important for proper
assembly, dynamics, and positioning of the mitotic spindle. *J Cell Biol* 158: 873-84

Roostalu J, Cade NI, Surrey T (2015) Complementary activities of TPX2 and chTOG constitute
an efficient importin-regulated microtubule nucleation module [corrigendum: same issue,
p.1512]. *Nat Cell Biol* 17: 1422-34

Roostalu J, Thomas C, Cade NI, Kunzelmann S, Taylor IA, Surrey T (2020) The speed of GTP
hydrolysis determines GTP cap size and controls microtubule stability. *eLife* 9

Sánchez-Soriano N, Gonçalves-Pimentel C, Beaven R, Haessler U, Ofner L, Ballestrem C,
Prokop A (2010) Drosophila growth cones: a genetically tractable platform for the analysis of
axonal growth dynamics. *Dev Neurobiol* 70: 58-71

Sánchez-Soriano N, Travis M, Dajas-Bailador F, Goncalves-Pimentel C, Whitmarsh AJ, Prokop
A (2009) Mouse ACF7 and Drosophila Short stop modulate filopodia formation and
microtubule organisation during neuronal growth. *J Cell Sci* 122: 2534-42

Sayas CL, Tortosa E, Bollati F, Ramirez-Rios S, Arnal I, Avila J (2015) Tau regulates the
localization and function of End-binding proteins 1 and 3 in developing neuronal cells. *J*
Neurochem 133: 653-67

Schaedel L, John K, Gaillard J, Nachury MV, Blanchoin L, Thery M (2015) Microtubules self-
repair in response to mechanical stress. *Nat Mater* 14: 1156-63

Sive HL, Grainger RM, Harland RM (2010) Microinjection of Xenopus oocytes. *Cold Spring Harb*
Protoc 2010: pdb prot5536

Slater PG, Cammarata GM, Samuelson AG, Magee A, Hu Y, Lowery LA (2019) XMAP215
promotes microtubule-F-actin interactions to regulate growth cone microtubules during axon
guidance. *J Cell Sci*: jcs.224311

Strumpf D, Volk T (1998) Kakapo, a novel *Drosophila* protein, is essential for the restricted
localization of the neuregulin-like factor, Vein, at the muscle-tendon junctional site. *J Cell Biol*
143: 1259-1270

Takei Y, Teng J, Harada A, Hirokawa N (2000) Defects in axonal elongation and neuronal
migration in mice with disrupted tau and map1b genes. *J Cell Biol* 150: 989-1000

Tang Q, Rui M, Bu S, Wang Y, Chew LY, Yu F (2020) A microtubule polymerase is required for
microtubule orientation and dendrite pruning in Drosophila. *EMBO J* 39: e103549

Thawani A, Kadzik RS, Petry S (2018) XMAP215 is a microtubule nucleation factor that
functions synergistically with the γ-tubulin ring complex. *Nature Cell Biology* 20: 575-585

Tymanskyj SR, Scales TM, Gordon-Weeks PR (2012) MAP1B enhances microtubule assembly
rates and axon extension rates in developing neurons. *Mol Cell Neurosci* 49: 110-19

van der Vaart B, Franker MA, Kuijpers M, Hua S, Bouchet BP, Jiang K, Grigoriev I, Hoogenraad
CC, Akhmanova A (2012) Microtubule Plus-End Tracking Proteins SLAIN1/2 and ch-TOG
Promote Axonal Development. *J Neurosci* 32: 14722-8

Vemu A, Szczesna E, Zehr EA, Spector JO, Grigorieff N, Deaconescu AM, Roll-Mecak A (2018)
Severing enzymes amplify microtubule arrays through lattice GTP-tubulin incorporation.
Science 361: eaau1504

Villarroel-Campos D, Gonzalez-Billault C (2014) The MAP1B case: an old MAP that is new
again. *Dev Neurobiol* 74: 953-71

Voelzmann A, Hahn I, Pearce S, Sánchez-Soriano NP, Prokop A (2016) A conceptual view at
microtubule plus end dynamics in neuronal axons. *Brain Res Bulletin* 126: 226-37

Voelzmann A, Liew Y-T, Qu Y, Hahn I, Melero C, Sánchez-Soriano N, Prokop A (2017)
Drosophila Short stop as a paradigm for the role and regulation of spectraplakins. *Sem Cell*
Dev Biol 69: 40-57

Wieczorek M, Bechstedt S, Chaaban S, Brouhard GJ (2015) Microtubule-associated proteins
control the kinetics of microtubule nucleation. *Nat Cell Biol* 17: 907-16

Wu X, Kodama A, Fuchs E (2008) ACF7 regulates cytoskeletal-focal adhesion dynamics and
migration and has ATPase activity. *Cell* 135: 137–48

Yang N, Inaki M, Cliffe A, Rørth P (2012) Microtubules and Lis-1/NudE/dynein regulate invasive
cell-on-cell migration in Drosophila. *PLoS One* 7: e40632

Zanic M, Stear JH, Hyman AA, Howard J (2009) EB1 recognizes the nucleotide state of tubulin
in the microtubule lattice. *PloS one* 4: e7585-e7585

Zanic M, Widlund PO, Hyman AA, Howard J (2013) Synergy between XMAP215 and EB1
increases microtubule growth rates to physiological levels. *Nat Cell Biol* 15: 688–693

Figures

Fig. 1. Eb1, Msps and Tau share the same combination of axonal loss-of-function phenotypes *Drosophila* in primary neurons. **A-H)** Images of representative examples of pre-cultured embryonic primary neurons either immuno-stained for Eb1 (top) or for tubulin (bottom); neurons were either wild-type controls (ctrl) or carried the mutant alleles *msps¹*, *tau^{KO}* or *Eb1^{04524}* in homozygosis (from left to right); asterisks indicate cell bodies, black arrow heads the axon tips, white arrow heads point at areas of MT disorganisation, dashed squares in A-D are shown as 3.5-fold magnified close-ups below each image with black rrows pointing at Eb1 comets; the axonal outline in D is indicated by a dotted line; scale bar in A represents 15 µm in all images. **I-N)** Quantification of different parameters (as indicated above each graph) obtained from pre-cultured embryonic primary neurons with the same genotypes as shown in A-H. Sample numbers, and P-values obtained with Kruskall-Wallis ANOVA test for the different genotypes are indicated in each graph. Error bars represent median ± 95% confidence interval (I - M) or mean ± SEM (N). For raw data see Tab. T1.

Fig. 1-S1. A candidate screen of axonal loss-of-function phenotypes in primary neurons. Graphs show extended data sets for four of the parameters displayed in Fig. 1 (indicated above each graph). Data points/bars representing mutant conditions for different genes are consistently colour-coded in all graphs, and conditions used are indicated below (6HIV, cultured from embryos for 6hrs; 6dpre, cultured from embryos for 12hrs following 6 day pre-culture; L3, cultured from late larval CNS for 18hrs). Allele names are given as superscript: no slash indicates homozygous, a present slash hetero-allelic conditions. Data were normalised to parallel controls (dashed horizontal line) and are shown as median ± 95% confidence interval (B,C) or mean ± SEM (A,D); merged sample numbers from at least two experimental repeats consisting of 3 samples each are shown at the bottom, P-values obtained with Kruskall-Wallis ANOVA test above data points/bars. For raw data see Tab. T1-S1.

Fig. 1-S2. Correlation of different Eb1 comet properties. **A,B)** Eb1 amount at comets is calculated as the product of comet length (A) and the fluorescent mean intensity of Eb1 comets (B), which are both to similar degrees affected by homozygous condition of *msps^A*, *tau^{KO}* and *Eb1^{04524}* in embryo derived neurons cultured for 12hrs following 6 day pre-culture (6dpre) ; data were normalised to controls (dashed horizontal line) and are shown as median ± 95% confidence interval; merged sample numbers from at least three experimental repeats are shown at the bottom, P-values obtained with Kruskall-Wallis ANOVA test above data points. **C)** Table lists data for Eb1 amounts (fixed neurons; compare Fig.1E-H, L) or for comet velocity/lifetime (live imaging; compare Fig.1M,N), all obtained from pre-cultured embryonic primary neurons carrying the same combinations of mutant alleles (indicated on the left; used alleles: *tub84B^{def}*, *msps^A*, *tau^{KO/Df}*, *eb1^{04524}*, *stai^{KO}*). **D,E)** Plotting comet velocity or lifetime against Eb1 amounts from different genetic conditions shows a fairly good correlation (r and p-value determined via non-parametric Spearman correlation analysis). For raw data see Tab. T1-S2.

Fig. 2. *Eb1, tau* and *msps* interact genetically. **A-B")** Axon length, MT disorganisation and Eb1

amount (as indicated on the right), for primary neurons displaying heterozygous (A-A'', larval
cultures) and homozygous (B-B', embryonic 6d pre-cultures') mutant conditions, alone or in
combination. Data were normalised to parallel controls (dashed horizontal lines) and are shown
as median ± 95% confidence interval (A, A'',B, B'') or mean ± SEM (A', B'); merged sample
numbers from at least two independent repeats of 3 parallel setups are shown at the bottom, P-
values obtained with Kruskall-Wallis ANOVA test above data points/bars; used alleles: $msps^A$,
tau^{KO}, $Eb1^{04524}$. **C**) Green dots represent data for MT disorganisation (from A',B') plotted against
Eb1 amounts (from B', B'' as well as Fig.1J,L, Fig.1-S1B,D), purple dots show comparable data
obtained from larval primary neurons (from Fig.A', A', Fig.1-S1B,D), and black dots similar data
obtained from primary *Xenopus* neurons (Fig.G,H); r and p-value determined via non-parametric
Spearman correlation analysis; see further detail of these correlations in Fig.2-S2. For raw data
see Tab. T2.

1009

Fig. 2-S1. Genetic interactions and heterozygous combinations. Graphs show data sets
extending on data displayed in Fig. 2 for heterozygous mutant conditions. **A**) Eb1 comet
numbers in fixed primary neurons cultured for 12hrs following 5 day pre-culture. **B**) Comet
velocity and lifetime obtained from live analyses of primary neurons cultured for 18hrs from late
larval CNSs. In all graphs, data were normalised to parallel controls (dashed horizontal lines)
and are shown as median ± 95% confidence interval (B) or mean ± SEM (A); merged sample
numbers from at least two experimental repeats are shown at the bottom, P-values obtained
with Kruskal-Wallis ANOVA test above data points/bars; used alleles: $msps^A$, tau^{KO}, $Eb1^{04524}$.
For raw data see Tab. T2-S1.

1019

Fig. 2-S2. Increasing Eb1 amounts correlate with decreasing MT disorganisation. The data and
graph show further details behind the correlations displayed in Fig.2C. **A**) The table shows for
different allelic combinations and culture conditions (indicated in 2nd column; 6HIV, cultured from
embryos for 6hrs; 6d pre, cultured from embryos for 12hrs following 6 day pre-culture; L3,
cultured from late larval CNS for 18hrs) the respective Eb1 amounts (3rd column) and MT
disorganisation (4th column), as obtained from different sets of experiments throughout this work
(5th column lists the figures from where these data originate); numbers in the 1st column
correspond to numbers of data points in the graph in B. **B**) Correlation plot of the data shown in
A, with numbers and colours of data points corresponding to the 1st column; r and p-value
determined via non-parametric Spearman correlation analysis. For raw data see Tab. T2-S2.

1030

Fig. 3. Eb1, Msps and Tau functionally interact in the fly brain and in frog primary neurons. **A,B**)
Medulla region of adult brains at 26-27 days after eclosure, all carrying the *GMR31F10-Gal4*
driver and *UAS-GFP-α-tubulin84B* (*GMR-tub*) which together stain MTs in a subset of lamina
neuron axons that terminate in the medulla; the further genetic background is either wild-type
(A) or triple-heterozygous ($Eb1^{04524/+}$ $msps^{A/+}$ $tau^{KO/KO}$; B); white/black arrows indicate axonal
swellings without/with MT disorganisation; rectangles outlined by red dashed lines are shown
as 2.5 fold magnified insets where white arrow heads point at disorganised MTs. **C,D**)
Quantitative analyses of specimens shown in A and B with respect to the number of axonal
swelling (C) and swellings with MT disorganisation (D); bars show mean ± SEM; P values from
Kruskal–Wallis one-way tests are given above each column, merged sample numbers (i.e.
individual axon bundles) from at least two experimental repeats at the bottom of each bar. **E-**

F'''') Primary *Xenopus* neurons stained either for tubulin (tub; top; white arrows indicating unbundled MTs, white arrowheads unbundled areas with MT disorganisation), black arrows comets labelled with MACF43::GFP (MACF43; visible as black spots); blacks dashed squares in E-E''' and F-F'''' shown as 2.5 fold magnified close-ups below; ↓ behind gene symbols indicates 50% knock-down thus approximating heterozygous conditions. **G,H)** Quantification of specimens shown in E-F'''' with respect to MT disorganisation (G) and comet MACF43::GFP amount (H); data were normalised to parallel controls (dashed horizontal lines) and are shown as mean ± SEM (G) or median ± 95% confidence interval (H); merged sample numbers from at least two experimental repeats are shown at the bottom, P-values obtained with Kruskall-Wallis ANOVA test above data points/bars. The scale bar in A represents 15 µm in A,B and 20 µm in E-F''''. For raw data see Tab. T3.

Fig. 3-S1. Support data for *Xenopus* experiments. **A-B')** A RT-PCR DNA gel/Western blot and their quantifications showing the degrees of EB3/Tau knock-downs upon application of different morpholino concentrations (indicated on top in blots and at the bottom in graphs), using ODC1 and ß-actin as loading controls; data are normalised to no-morpholino controls from two experimental repeats (dashed lines). 50% knock-down of XMAP215 was achieved by injecting 6 ng of the validated XMAP215 MO as described previously (Lowery et al., 2013, Slater et al., 2019). **C-G)** Different properties of MACF43::GFP comets (as indicated upon graphs) obtained from *Xenopus* primary neurons, either upon 50% (C,D) or 70% (E-G) knock-down of respective genes (indicated as 50/70%KD); data were normalised to parallel controls (dashed horizontal lines) and are shown as median ± 95% confidence interval; merged sample numbers from at least two experimental repeats are shown at the bottom, P-values obtained with Kruskall-Wallis ANOVA and Dunn's posthoc test above data points. For raw data see Tab. T3-S1.

Fig. 4. Eb1 and Msps depend on each other for MT plus end localisation. **A-A")** Primary neurons at 6HIV co-expressing Eb1::mCherry (magenta, Eb1) and Msps^{FL}::GFP (green, Msps) and imaged live; asterisks indicate somata, scale bar represents 10µm, dashed boxes indicate the positions of the 3.5-fold magnified close-ups shown at the bottom with arrowheads pointing at the position of Msps::GFP accumulation (same in C,C'). **B,B')** Kymograph of live movies (as in A-A") with the dashed line on the left representing the dashed lines shown in A' and A" (i.e. the length of the axon; proximal at the top) and the x-axis indicating time; arrowheads point at trajectories of Msps and Eb1 which are almost identical. **C)** Primary neurons expressing Msps::GFP and imaged live, either displaying wild-type background (ctrl) or being homozygous mutant for *Eb1^{04524}* (*Eb1$^{-/-}$*); white arrowheads point at Msps::GFP comets which are much smaller in the mutant neurons. **D)** Schematic representations of Msps^{FL}::GFP and Msps$^{\Delta CTD}$::GFP. **D',D")** Graphs displaying axon length and MT disorganisation (as indicated) for pre-cultured embryonic primary neurons expressing GFP or Msps::GFP constructs via the *elav-Gal4* driver, either in wild-type or *msps$^{A/1}$* mutant background; data were normalised to parallel controls (dashed horizontal lines) and are shown as median ± 95% confidence interval (D') or mean ± SEM (D"); merged sample numbers from at least two experimental repeats are shown at the bottom, P-values obtained with Kruskall-Wallis ANOVA test above data points/bars. **E)** Model view of the results shown here and in Fig. 1; for explanations see main text and Discussion. For raw data see Tab. T4.

Fig. 4-S1. Loss of Tacc or Sentin does not cause obvious axonal phenotypes. Axon length (**A**) and MT disorganisation (**B**) for primary neurons at 6HIV which were either wild-type (wt) or homozygous mutant for *sentin* or *dTACC* (as indicated); data were normalised to parallel controls (dashed horizontal lines) and are shown as median ± 95% confidence interval (A) or mean ± SEM (B); merged sample numbers from at least two experimental repeats are shown at the bottom, P-values obtained with Kruskal-Wallis ANOVA test above data points/bars. For raw data see Tab. T4-S1.

Fig. 5. Tau promotes Eb1 pools at MT plus ends by outcompeting its association with the MT lattice. **A-A"**) Example of neuron imaged live with disorganised MTs to illustrate Tau binding (green) along the MT lattice, separated from Eb1 comets (magenta); asterisks indicate somata, the scale bar represents 10µm, dashed boxes indicate the positions of the 4fold magnified close-ups shown at the bottom, white arrowheads point at Eb1 comets (same in B-D). **B-D**) Primary neurons at 6HIV stained for Eb1 which are either wild-type (B), *tau^{KO/Df}* mutant (C) or *tau^{KO/Df}* mutant plus expressing Eb1::GFP driven by *elav-Gal4* (D); white/red arrowheads indicate Eb1 comets/lattice localisation. **E-I**) Different parameters (as indicated) of control (ctrl) or *tau^{KO/Df}* (*tau^{-/-}*) mutant neurons without/with *elav-Gal4*-driven expression of Eb1::GFP or Shot-Ctail::GFP (as indicated); data were normalised to parallel controls (dashed horizontal lines) and are shown as mean ± SEM (I) or median ± 95% confidence interval (E-H); merged sample numbers from at least two independent repeats with 3 experimental setups each are shown at the bottom, P-values obtained with Kruskal-Wallis ANOVA and Dunn's posthoc test above data points/bars. **J**) Model view of the results shown here; for explanations see main text and Discussion. For raw data see Tab. T5.

Fig. 5-S1 Upon tau deficiency, GTP-tubulin is reduced at MT plus ends but unchanged at lattices. **A-B"**) Fixed primary neurons stained for Eb1 (magenta) and GTP-tubulin (green); asterisks indicate somata, scale bar represents 10µm, dashed boxes indicate the positions of the 4fold magnified close-ups shown at the bottom, arrowheads point at Eb1::GFP comets and GTP caps. **C,D**) Graphs showing staining intensity of GTP-tubulin at MT plus ends (C) and along the MT lattice (D), and that *msps^{A/146}* mutant phenotypes were not rescued by Eb1::GFP expression (C,D). **E**) Graphs showing staining intensity of Eb1 along the MT lattice of neurons without/with *elav-Gal4*-driven expression of *dtau*. Overexpression of *dtau* leads to a reduction of Eb1 at the MT shaft; data were normalised to parallel controls (dashed horizontal lines) and are shown as median ± 95% confidence interval; merged sample numbers from at least two independent repeats with 3 experimental setups each are shown at the bottom, P-values obtained with Kruskall-Wallis ANOVA test above data points/bars. For raw data see Tab. T5-S1.

Fig. 5-S2 Eb1 overexpression does not rescue *msps* mutant phenotypes. Graphs show that Eb1 lattice localisation is unaffected by loss of Msps in primary neurons at 6HIV (A) or at ~12HIV following 6 day preculture (B). Staining intensity of GTP-tubulin at MT plus ends is reduced in *msps^{A/A}* and *Eb1^{04524/04524}* mutant at 6dpre (C). Expressing Eb1::GFP via *elav-Gal4* does not rescue comet velocities (D), comet lifetime € and MT disorganisation (F) in primary neurons of *msps^{A/146}* mutants (cultured12HIV following 6 day preculture); data were normalised to parallel controls (dashed horizontal lines) and are shown as mean ± SEM (F) or median ± 95% confidence interval (A-E); merged sample numbers from at least two experimental repeats are shown at the bottom, P-values obtained with Kruskal-Wallis ANOVA test with Dunn's posthoc analysis above data points/bars. For raw data see Tab. T5-S2

Fig. 6. Shot-mediated guidance as a mechanism linking Eb1 at MT plus ends to bundle organisation. **A**) Schematic representation of Shot constructs (CH, actin-binding calponin-homology domains; EF, EF-hand motifs; GRD, MT-binding Gas2-related domain; Ctail, unstructured MT-binding domain containing Eb1-binding SxIP motifs in blue); in Shot-3MTLS*::GFP the SxIP motifs are mutated. **B**) Fixed primary larval neurons at 18HIV obtained from late larval CNS stained for GFP (green) and tubulin (magenta), which are either wild-type (top) or $Eb1^{04524/+}msps^{A/+}tau^{KO/+}$ triple-heterozygous (indicated on right) expressing GFP or either of the constructs shown in D; scale bar 10µm. **C**) Quantification of MT disorganisation of neurons as shown in B. **D,E**) MT disorganisation in $shot^{3/3}$ $Eb1^{04524/04524}$ double-mutant neurons is not enhanced over single mutant conditions assessed in fixed primary neurons at 6HIV (D) or 12HIV following 6 day pre-culture (E). In all graphs data were normalised to parallel controls (dashed horizontal lines) and are shown as mean ± SEM; merged sample numbers from at least two independent repeats with 3 experimental setups each are shown at the bottom, P-values obtained with Kruskall-Wallis ANOVA test above bars. **F,F'**) Model derived from previous work (Alves-Silva et al., 2012), proposing that the spectraplakin Shot cross-links Eb1 at MT plus ends with cortical F-actin, thus guiding MT extension in parallel to the axonal surface.

Fig. 7. Model summarising the common functional pathway of Eb1, Msps and Tau. Images on the left show functional diagrams and on the right cartoon models for wild-type and four mutant conditions as indicated. **A,A'**) In wild-type (WT) neurons, the three factors bind independently to MTs: Msps to the very tip of plus ends, Eb1 to the GTP-cap (GTP-tubulin shown with yellow dots) but lagging behind the front, and Tau along the lattice. Tau promotes Eb1 pools at MT plus ends (green arrow in A) by outcompeting its binding along the lattice (green T-bar in A'). Eb1 and Msps mutually promote each other's localisation (stippled black and red arrows in A, A'): Msps is the main promoter of polymerisation (brown arrow), thus sustaining a prominent GTP-tubulin cap (pale yellow box in A) for Eb1; Eb1 potentially promotes sheet formation of protofilaments at the plus tip (red arrow), thus helping Msps to bind; Eb1 links to Shot (orange double-arrow) which provides MT plus end guidance (blue dashed arrow). **B,B'**) Upon loss of Eb1, the plus end sheet structure is weakened, thus weakening Msps binding and, in turn, reducing polymerisation (thinner/shorter arrows, smaller font); Shot detaches, thus abolishing guidance (curved off-track extension arrow). **C,C'**) Upon loss of Msps, its catalysis of polymerisation is abolished, the GTP-tubulin cap shrinks, less Eb1 binds, thus strongly weakening Shot binding and guidance. **D,D'**) Upon loss of Tau, Eb1 is recruited to the MT lattice, thus reducing its plus end amounts; reduced Eb1 negatively impacts on Shot and Msps causing the milder guidance and polymerisation effects we observed. **E,E'**) Upon loss of Shot, the localisation of the other three proteins is unaffected, but guidance is abolished.

Movie M1. Msps::GFP and Eb1::RFP jointly track MT plus ends. Live movie of a wild-type neuron co-expressing Msps::GFP and Eb1::RFP; for stills see Fig.4A-B'. As indicated, single channels are shown on the left and middle, and the combined movie on the right. The movie was acquired at 0.5 frames per second, and play at 0.5 s per frame. The scale bar indicates 10 µm.

Movie M2. Msps plus end localisation in wild-type neurons. Live movie of a wild-type neuron expressing Msps::GFP; for stills see Fig.4C. The movie was acquired at 1 frame per second, and plays at 0.2 s per frame. The scale bar indicates 10 µm.

1183 **Movie M3**. Msps plus end localisation is impaired in the absence of Eb1. Live movie of an
1184 *Eb1$^{04524/04524}$* mutant neuron expressing Msps::GFP; for stills see Fig.4C'. The movie was
1185 acquired at 1 frames per second, and plays at 0.2 s per frame. The scale bar indicates 10 μm.

Hahn et al. Fig. 1

Hahn et al. Fig. 1-S1

A **comet length**

B **comet intensity**

C

	Eb1 amount	comet velocity	comet lifetime
ctrl	1.0000	1.0000	1.0000
tau+/-	1.1130	1.1010	1.0360
msps+/-	1.0430	0.9792	0.9411
eb1+/-	0.9028	1.1430	1.0360
eb1+/- msps+/-	0.7093	0.8496	0.9097
eb1+/- msps+/- tau+/-	0.3989	0.7602	0.7879
msps-/-	0.3796	0.4676	0.6905
tau-/-	0.8918	0.8234	0.7246
tubdef/def	1.1020	1.0720	1.0160
stai-/-	1.0450	0.9107	1.0490

D

p = 0.0028
r = 0.8320

E

p = 0.008
r = 0.7786

Hahn et al. Fig. 1-S2

Hahn et al. Fig. 2

Hahn et al. Fig. 2-S1

A

		comet Eb1 amount	MDI	Figure
1	6d pre *msps*[1/1]	0.3796	2.338	Fig. XXX1
2	6d pre *tau*[KO/KO]	0.8918	1.388	Fig. XXX1
3	6d pre *Eb1*[04524/04524]	0.282	2.191	Fig. XXX1
4	6d pre *msps*[1/1] *tau*[KO/KO]	0.4592	1.786	Fig. XXX2
5	6d pre *Eb1*[04524/04524] *tau*[KO/KO]	0.3298	2.063	Fig. XXX2
6	6d pre *Eb1*[04524/04524] *msps*[1/1]	0.3421	2.109	Fig. XXX2
7	6d pre *Eb1*[04524/04524] *msps*[1/1] *tau*[KO/KO]	0.2077	4.363	Fig. XXX2
8	7d pre *Eb1*[Def]	0	5.814	Fig. XXX2
9	6d pre *tub*[def]	1.102	0.6603	Fig. XXX1-S1
10	6d pre *clasp*[2/Df]	1.134	0.4888	Fig. XXX1-S1
11	6d pre *MAP1B*[-]	1.162	0.8491	Fig. XXX1-S1
12	6HIV *Eb1*[04524]	0.3832	2.697	Fig. XXX1-S1
13	6HIV *msps*[A]	0.4884	1.622	Fig. XXX1-S1
14	6HIV *tau*[KO]	0.5217	1.815	Fig. XXX1-S1
15	6HIV *tub*[Df]	1.09	0.43	Fig. XXX1-S1
16	6HIV *stai*[KO1]	1.04	0.69	Fig. XXX1-S1
17	L3 *msps*[1/+]	1.043	1.176	Fig. XXX2
18	L3 *tau*[KO/+]	1.114	1.11	Fig. XXX2
19	L3 *Eb1*[04524/+]	0.9031	1.397	Fig. XXX2
20	L3 *msps*[1/+] *tau*[KO/+]	0.7372	1.932	Fig. XXX2
21	L3 *tau*[KO/+] *Eb1*[04524/+]	0.8818	1.243	Fig. XXX2
22	L3 *msps*[1/+] *Eb1*[04524/+]	0.7095	2.609	Fig. XXX2
23	L3 *msps*[1/+] *tau*[KO/+] *Eb1*[04524/+]	0.399	6.655	Fig. XXX2
24	L3 elavG4; *tau*[KO/def]	0.7631	2.758	Fig. XXX5
25	L3 elav>EB1-GFP, tauKO/def	1.067	1.34	Fig. XXX5
26	L3 *stai*[Df/KO3]	1.045	1.197	Fig. XXX1-S1
27	Xen 50% Tau KD	0.9358	1.411	Fig. XXX3
28	Xen 50% XMAP215 KD	0.8873	1.892	Fig. XXX3
29	Xen 50% EB3 KD	0.9271	2.002	Fig. XXX3
30	Xen 50% XMAP215/Tau/EB3 KD	0.5739	3.822	Fig. XXX3

B

Hahn et al. Fig. 2-S2

A *GMR-tub*

B *GMR-tub Eb1⁺ᐟ⁻, msps⁺ᐟ⁻, tau⁺ᐟ⁻*

C # of swellings/ axon bundle ↑

swelling/axon bundle

<0.0001

ctrl   *Eb1⁺ᐟ⁻ msps⁺ᐟ⁻ tau⁺ᐟ⁻*

381   285

D # of disor. MT areas/ axon bundle ◁

disorg. MT areas/ axon bundle

<0.0001

ctrl   *Eb1⁺ᐟ⁻ msps⁺ᐟ⁻ tau⁺ᐟ⁻*

382   289

E | **E'** | **E''** | **E'''** | **E''''**

tub

F | **F'** | **F''** | **F'''** | **F''''**

MACF43

control | *Tau↓* | *XMAP↓* | *EB3↓* | *Tau↓ XMAP↓ EB3↓*

↓50% KD

G MT disorganisation

MDI

0.0002

0.99 0.9 0.08 <0.001

Control Tau↓ XMAP215↓ EB3↓ EB3↓XMAP215↓Tau↓

64 16 24 53 54

H comet MACF43 amount

comet MACF43 amounts

<0.001

0.99 0.09 0.99 <0.001

Control Tau↓ XMAP215↓ EB3↓ EB3↓XMAP215↓Tau↓

95 40 40 47 41

Hahn et al. Fig. 3

Hahn et al. Fig. 3-S1

Hahn et al. Fig. 4

Hahn et al. Fig. 4-S1

Hahn et al. Fig. 5

Hahn et al. Fig. 5-S1

Hahn et al. Fig. 5-S2

Hahn et al. Fig. 6

Hahn et al. Fig.7